

Many body localization proximity effects in platforms of coupled spins and bosons

J. Marino¹ and R.M. Nandkishore¹

¹*Department of Physics and Center for Theory of Quantum Matter,
University of Colorado Boulder, Boulder, Colorado 80309, USA*

We discuss the onset of many body localization in a one-dimensional system composed of a XXZ quantum spin chain and a Bose-Hubbard model linearly coupled together. We consider two complementary setups depending whether spatial disorder is initially imprinted on spins or on bosons; in both cases, we explore the conditions for the disordered portion of the system to localize by proximity of the other clean half. Assuming that the dynamics of one of the two parts develops on shorter time scales than the other, we can adiabatically eliminate the fast degrees of freedom, and derive an effective hamiltonian for the system's remainder using projection operator techniques. Performing a locator expansion on the strength of the many-body interaction term or on the hopping amplitude of the effective hamiltonian thus derived, we present results on the stability of the many-body localized phases induced by proximity effect. We also briefly comment on the feasibility of the proposed model through modern quantum optics architectures, with the long-term perspective to realize experimentally, in composite open systems, Anderson or many-body localization proximity effects.

I. INTRODUCTION

Inhibition of energy transport induced by the presence of spatial disorder has its origins at the middle of last century [1, 2], and its phenomenology has been confined to free systems till the last decade. More recently, a number of works ranging from applications of perturbation theory [3–5] to numerical simulations [6–8] have established that a localized phase exhibiting absence of diffusion and equilibration on long time scales, can survive the presence of many body interactions. For instance, a one dimensional disordered quantum system can show a many-body localized phase [9–11] even at strong interactions or at high energy densities, failing to act as its own bath. The existence of a many-body localized phase has been recently proven rigorously for one-dimensional systems on a lattice with short-range interactions [12], and explored in experiments with cold gases [13–15]. This novel phase of matter not only displays breakdown of ergodicity as salient feature, but it also shows a rich phenomenology, including connections to integrability [16–22], unusual response properties [23, 24], a rich pattern of quantum entanglement [25–29], and new types of order that cannot arise in equilibrium [30–32]. This many body localization (MBL) is accordingly drawing considerable interest.

While much effort has been devoted to the question of when an isolated quantum system can be localized [33–43], in reality, any system is unavoidably coupled to an environment, and understanding the interplay of a many-body localized phase communicating with a thermal bath is of paramount importance, both for the exploration of the phenomenon in experiments, as well as a to understand its robustness to ergodic perturbations. The naive expectation that the bath can provide sufficient energy and phase-space to facilitate the hopping in an otherwise localized system, has been confirmed by a series of theoretical studies [41, 44–52] and by a recent experiment [53]. These works have however shown that the interplay of localization and dissipation can leave signatures on the

evolution of observables of interest at intermediate times, before thermalization establishes eventually.

These findings suggest an inherent fragility of disorder-induced localized phases of matter to the coupling to the environment. However, there exist other captivating scenarios when open systems supporting a localized phase are not necessarily doomed to a restoration of thermodynamic equilibrium; in addition to dissipation engineering via non-local Lindbladians capable to drive a system into a state with desired localization properties, the possibility of a *many body localization proximity effect* represents a promising direction. The phenomenon has been originally discussed in a toy model of two interacting systems composed of different elementary degrees of freedom [36]: when a disordered system is coupled to a clean one with comparable size acting virtually as a bath, the former can in turn, for certain choice of interaction and disorder strengths, induce localization on the latter. Starting from the analysis in Ref. [36], we aim at presenting in this work a realistic setup (see for some experimental proposals Sec. V) where a many body proximity effect can occur in a quantum spin chain coupled to interacting bosons hopping on a lattice (Sec. II). Our analysis makes use of the Lindblad formalism, which has not been previously applied to the MBL proximity effect. We will consider both disorder imprinted on the spin (Sec. III) or bosonic sector (Sec. IV), and establish the conditions for the robustness of a many-body localized phase, induced by the proximity of the disordered system into the clean one. We provide a series of estimates for the borders of the localization/delocalization transition in various parameter regimes, basing our study on the combination of adiabatic elimination techniques and locator expansion, which can serve as guideline for subsequent numerical and analytical developments. We also discuss the more natural occurrence of the clean, ergodic system acting as a bath, which delocalizes the disordered one. In view of proposing viable platforms to observe these effects, we also suggest at the end of the paper (Sec. V) some po-

tentially interesting quantum optics platforms where the phases discussed in this work might be experimentally explored in the future.

II. THE MODEL

We consider a one dimensional model composed of a XXZ quantum spin chain coupled linearly to a Bose-Hubbard model, as given by the hamiltonian (see also Fig. 1)

$$H = -J \sum_{\langle i,j \rangle} b_i^\dagger b_j + \frac{U}{2} \sum_i n_i(n_i - 1) - \sum_i \omega_i b_i^\dagger b_i + H_{int} + \alpha \sum_{\langle i,j \rangle} \sigma_i^+ \sigma_j^- + \lambda \sum_i \sigma_i^z \sigma_{i+1}^z + \sum_i h_i \sigma_i^z, \quad (1)$$

The term H_{int} denotes a spin-boson, dipolar-like (see Sec. V for the choice of terminology in connection with potential experimental realisations), interconversion term

$$H_{int} = g \sum_i (\sigma_i^+ b_i + h.c.), \quad (2)$$

while $n_i = b_i^\dagger b_i$ labels bosonic occupation numbers, and σ_i^α ($\alpha = x, y, z$) are Pauli matrices describing spins-1/2 on sites $i = 1, \dots, N$; the sum $\sum_{\langle i,j \rangle} \dots$ is intended over next-nearest-neighbours. The transverse field h_i and the chemical potential ω_i are taken site-dependent, since they will alternatively host spatial disorder.

A Jordan-Wigner transformation renders the XXZ quantum spin chain an interacting fermionic model, which, in presence of disorder, constitutes a paradigm for the many body localization transition [6, 7]. We therefore first imprint disorder on the transverse fields, h_i , drawn accordingly from a uniform probability distribution of width, W , and consider the emergence of a many-body localization transition in the bosonic chain by proximity of the disordered spin model, Sec. III. Alternatively, we consider on-site disorder (uniformly drawn from a distribution of width Ω) on the chemical potential of bosons, ω_i (see Sec. IV), and study the impact of the latter on the quantum spin chain. A schematic representation of the setups considered is provided in Fig. 1.

In order to estimate the border of the many-body localized phase, we will always consider a disorder sufficiently strong to push the disordered sector of the system (spins or bosons) deep into the localized phase and evaluate its impact on the latter portion, assumed clean. A fully many-body-localized system (i.e. with all the many-body eigenstates localized), renders more transparent the analysis, since the effective hamiltonian can be approximated by a sum of several single particle terms, with the range of many-body interactions falling exponentially with a short correlation length if disorder is sizeable. Under this hypothesis the conserved, l -bits of the many-body




FIG. 1. We consider an interacting quantum spin chain coupled to a Bose-Hubbard model by a linear, dipole-like interaction term, $H_{int} = g \sum_i (\sigma_i^+ b_i + h.c.)$, see hamiltonian (1). When the typical energy scale ruling the dynamics of the bosons (or of the spins) is larger than the coupling g , one can adiabatically eliminate the bosonic (spin) sector, and derive an effective hamiltonian for the remainder of the system, *viz.* spins (or bosons). We study the onset of a many body localized phase in these effective hamiltonians derived in adiabatic elimination, imprinting first disorder on the spin sector (S) in Sec. III, and later in Sec. IV on the bosonic (B) one.

localized phase [12, 18, 54] are close to original bare degrees of freedom employed in hamiltonian (1), which can be therefore employed in the subsequent analysis without qualitatively affecting the generality of the conclusions.

III. DISORDER IN THE SPIN SECTOR

A. MBL proximity effect for bosons

The first case considered in our analysis consists of spins subject to disordered fields, $h_i \in [-W/2, W/2]$, along the \hat{z} -direction, and linearly coupled to bosons as described by the hamiltonian

$$H = -J \sum_{\langle i,j \rangle} b_i^\dagger b_j + \frac{U}{2} \sum_i n_i(n_i - 1) - \omega_0 \sum_i b_i^\dagger b_i + H_{int} + \sum_i h_i \sigma_i^z. \quad (3)$$

Compared to (1), we have assumed a constant chemical potential $\omega_i = \omega_0, \forall i = 1, \dots, N$. The hamiltonian (3) models the transfer of disorder from a bath of free spins into a clean system of bosons via the linear term H_{int} ; as we commented in Sec. II, spins precessing in strong ($W \gg \lambda, \alpha$), disordered transverse fields h_i , can be interpreted as resulting from a disordered XXZ spin chain, deep in the many-body localized phase. In the $W \rightarrow \infty$ limit all other terms in the spin Hamiltonian can be neglected; at large but finite W , the spin Hamiltonian can be written in terms of emergent local integrals of motion or ‘ l -bits,’ which are approximately equal to the σ_i^z , with some dressing by higher spin terms [12, 18, 54].

Within the energy scale separation $g \ll W$, i.e. assuming that the spin bath has faster dynamics than the

bosonic system ($\omega \ll W$), we can integrate out the former and find an effective dynamics for the bosons, using projection operator techniques [55]. In particular, denoting with $\rho_b(t)$ the reduced density matrix for the bosons at time t , we can write an effective master equation, where spin dynamics has been averaged out [55, 56], as a result of second order perturbation theory in the coupling g ,

$$\dot{\rho}_b(t) = \mathcal{L}_b \rho_b(t) - \int_0^\infty d\tau \text{Tr}_s \{ [H_{int}, e^{\tau \mathcal{L}_s} ([H_{int}, \rho_b(t) \otimes \rho_s^0])] \}. \quad (4)$$

In Eq. (4), the density matrix ρ_s^0 encodes the state in which the spins are frozen as a result of the above mentioned energy scale separation, and \mathcal{L}_s is the Liouvillian evolver of the spin sub-system; finally, $\text{Tr}_s \{ \dots \}$ denotes the trace on the spin's Hilbert space.

Eq. (4) holds in the generic case of dynamics with co-existent hamiltonian and Lindbladian processes [55]. Although for large part of the paper, we will consider purely hamiltonian evolutions, in Sec. V we study the impact of dissipation on the spin sector, where the use of Eq. (4) is convenient to perform adiabatic elimination.

In order to apply (4), we first assume that the bosonic gas is in the atomic limit ($J = 0$) and subsequently (see discussion after Eq. (6)) we perform a locator expansion in the hopping J , in order to find the border of the many body localization transition as a function of the couplings appearing in Eq. (3). The effective hamiltonian, H'_b , for the bosonic system is easily identified writing the right hand side of (4) as the generator of a unitary dynamics,

$$\dot{\rho}_b(t) = -i[H'_b, \rho_b(t)]. \quad (5)$$

The trace over the spin degrees of freedom is performed assuming that the spin system is in an eigenstate of the disordered spin hamiltonian, $h_s = \sum_i h_i \sigma_i^z$. The computations on the right hand side of Eq. (4) can be easily carried (see for instance [56, 57] and the Appendix), as they require to freely evolve the spin operators in H_{int} with $e^{\tau \mathcal{L}_s}$ (in this case, we have simply $\mathcal{L}_s = ih_s$), and to evaluate the time integral in the variable τ . Since spin states are localized in real space (as consequence of having assumed the XXZ spin chain deep in the localized phase), the new effective disordered bosonic term in H'_b will be local in space as well: indeed, the matrix element $\sum_{i,j} \langle \sigma_i^- \sigma_j^+ \rangle b_i^\dagger b_j$, occurring in the evaluation of Eq. (4), will have local spatial support, $\langle \sigma_i^- \sigma_j^+ \rangle \sim \delta_{i,j}$, because strong disorder rules out spin states, ρ_s^0 , delocalized in real space. Eq. (4), therefore, yields the effective bosonic hamiltonian

$$H'_b = -J \sum_{\langle i,j \rangle} b_i^\dagger b_j + \frac{U}{2} \sum_i n_i(n_i - 1) - \sum_i \left(\omega_0 + \frac{g^2}{s_i h_i} \right) n_i, \quad (6)$$

where the disordered dressing of the bosonic frequency ω_0 is the main effect of the adiabatic elimination of the spins. The term s_i in (6) is reminiscent of the state of the

spin bath, ρ_s^0 , and it can acquire only the values $s_i = \pm 1$, depending whether in ρ_s^0 the spin on the lattice site i , was respectively oriented up or down along the \hat{z} direction; however, since h_i is drawn from a distribution with support in $[-W/2, W/2]$ and centred around zero mean, the sign of s_i cannot qualitatively alter the results of the following analysis. The impact of the hopping term $\propto J$, on the otherwise trivially localized boson hamiltonian (Eq. (6) with $J = 0$), will now be discussed in locator expansion. As a side remark, we notice that if the dynamics of spins is itself dissipative, terms in the Lindblad form will appear in the effective Liouvillian for the bosons as a result of the calculations contained in Eq. (4); see for instance Sec. IV C.

As anticipated, we now consider the effect of the hopping J in locator expansion. A regime of interest is the one where ω_0 is negligible compared to the disordered correction $\propto g^2$ arising from adiabatic elimination of spins, $\omega_0 \ll g^2/W$. At $J = 0$, hamiltonian (6) is trivially localized because in the atomic limit; switching the hopping, locator expansion [33, 36] predicts a critical value of J , at which transport is restored in the system

$$J \gtrsim J_c \simeq \frac{g^2}{W} \exp\left(-\frac{1}{2} \Sigma_T \xi\right). \quad (7)$$

In the above expression ξ is the localization length of bosons and Σ_T the entropy density at fixed temperature T (we recall that in the estimate (7) and in the others following in the next sections, the localization length ξ is measured in units of the lattice spacing of (1), and therefore adimensional). Formula (7) is derived (see Ref. [36]) requiring that the typical matrix element, $J \exp(-\frac{1}{2} \Sigma_T \xi)$, among states connected by the bosonic hopping term is larger than their off-shell energy level spacing, $(g^2/W) \exp(-\Sigma_T \xi)$, indicating the breaking of the locator expansion, and the onset of a delocalized ergodic phase in the system. However, as long as $J \lesssim J_c$, the bosonic system described by (6) will be in a localized phase induced by the coupling to the original strongly disordered spin model and representing accordingly a typical instance of *many-body localization proximity effect*.

B. Delocalization of the disordered spin bath

We now consider a regime of energy scales separation complementary to the one discussed above. Specifically, we set to zero the spin hopping term ($\alpha = 0$ in Eq. (1)), and discuss the generation of a next-neighbour spin-flipping term when the clean bosonic system is traced out. The impact of the transverse spin-spin interaction term $\propto \lambda$ will be analysed in locator expansion. When $g \ll \omega_0$ in hamiltonian (3), we can adiabatically eliminate the bosons, if they evolve faster compared to spins, $\omega_0 \gg W$, and derive an effective hamiltonian for the latter using once again the formula (4) where the role of bosons and spins are now interchanged with respect

to Sec. III A. Since the bosons are clean from disorder, we can now assume that they are frozen in a delocalized state in real space, contrary to the 'l-bits' of Sec. III A; this carries the consequence that the effective spin hamiltonian can inherit an hopping term $\propto g^2 \sum_{\langle i,j \rangle} \sigma_i^+ \sigma_j^-$ after bosons are traced away, because the matrix element $\langle b_i b_j^\dagger \rangle$ can have non-local support on the lattice.

The effective hamiltonian for the spins degrees of freedom reads then,

$$H'_s = \sum_i h_i \sigma_i^z + \sum_i \frac{g^2}{\omega_0 + U(1 + 2\bar{n}_i)} (\sigma_i^+ \sigma_{i+1}^- + h.c.). \quad (8)$$

For instance, in order to generate a hopping to next neighbouring sites as in Eq. (8), is sufficient to trace out a bosonic state, $|\psi\rangle_b$, written as a superposition $|\psi\rangle_b = c_1 |\{01\}_i\rangle + c_2 |\{10\}_i\rangle$ (with c_1 and c_2 arbitrary constants), where the shorthand $|\{01\}_i\rangle$ stands for a state with sites alternatively occupied and unoccupied by bosons, $|010101\dots01\rangle$, and the state $|\{10\}_i\rangle = |101010\dots10\rangle$ is just obtained by a unit shift of the lattice site position in $|\{01\}_i\rangle$. The number \bar{n}_i in (8) is the expectation value of the conserved, local bosonic number \hat{n}_i (we recall that we considering the atomic limit $J = 0$ for bosons).

A one dimensional, non-interacting disordered system is an Anderson insulator at any disorder strength; in order to study the potential onset of delocalization in this system via many body spin-spin interactions, we treat in locator expansion the term

$$V = \lambda \sum_i \sigma_i^z \sigma_{i+1}^z \quad (9)$$

in the hamiltonian (8) (therefore now H'_s has been dressed by V), having in mind the structure of the XXZ spin chain in (1). Since V is a quartic fermionic interaction after Jordan-Wigner fermionization, we can estimate the magnitude of the typical term of the locator series [33, 36] connecting two localized states $|\alpha\rangle$ and $|\beta\rangle$ of the spin hamiltonian (8) with energies respectively E_α and E_β ,

$$\frac{\langle \alpha | V | \beta \rangle}{E_\alpha - E_\beta} \sim \frac{\lambda \xi^{-d}}{J' \xi^{-4d}} \quad (10)$$

where $d = 1$ in our case and $J' \sim g^2/(\omega_0 + U)$ is the effectively generated hopping in hamiltonian (8). The estimate for the matrix element of the interaction $\langle \alpha | V | \beta \rangle$ is derived [33] in the basis of the localized wave functions of the spins, and requires to take properly into account the normalization of the localized states, $\propto \xi^{-d/2}$, while the energy level spacing in the denominator, $\Delta E = E_\alpha - E_\beta$, is proportional to ξ^{-4d} , since the interaction involves four localized fermions, acting each one on a region of order $\sim \xi^{-d}$. Since the model is one dimensional, locator expansion for the free model (8) predicts a localization length smaller than lattice spacing and scaling as the logarithm of the ratio between disorder and hopping

strengths, $\xi \propto 1/\log(W/J')$, since the spin chain (8) will be in a strongly localized regime for the energy scale separation adopted. Therefore, the threshold in the coupling λ for delocalizing the disordered spin bath is, using Eq. (10),

$$\lambda \gtrsim \lambda_c \simeq J' \log^3(W/J'), \quad (11)$$

since as far as $\lambda \lesssim \lambda_c$ and the locator expansion converges, the eigenstates of the interacting problem would remain close to those of the Anderson insulator (8) present at $\lambda = 0$.

In a similar fashion, if we start with bosons in a hopping dominated regime ($U = 0$, $t \neq 0$) and with spins in a strongly localized phase, the adiabatic elimination of the latter will imprint on-site disorder on the former which will immediately localize the bosons at any disorder strength (once again, the system is one dimensional). If one treats on-site bosonic interactions in locator expansion, one can easily show, following the same procedure outlined above, that the spin-induced, localized phase of bosons melts as interaction's strength, U , reaches values $\propto t \log^3(g^2/(Wt))$, where we have employed again strong localization theory.

IV. DISORDER IN THE BOSONIC SECTOR

A. MBL proximity effect for spins

The setup of the second part of this work complements the one of the previous Section. Disorder is now imprinted on bosons, specifically, the chemical potentials ω_i are drawn from a uniform distribution of width Ω and with zero mean value; complementing Sec. III A, we aim at inspecting the impact of the disorder transfer from bosons to the quantum spin chain hamiltonian with uniform transverse fields, $h_i = h_0$, cfr. Eq. (1),

$$H = \sum_i h_0 \sigma_i^z + \lambda \sum_i \sigma_i^z \sigma_{i+1}^z + \alpha \sum_{\langle i,j \rangle} \sigma_i^+ \sigma_j^- + H_{int} + \sum_i \omega_i b_i^\dagger b_i. \quad (12)$$

Since we now wish to adiabatically eliminate the bosons, we adopt the energy scales separation $g \ll \Omega$ and $h_0 \ll \Omega$, while hopping will be subsequently switched on in locator expansion. The choice of a diagonal bosonic hamiltonian can, once again, be considered the extreme limit of a strongly disordered, interacting bosonic system, deep in its many-body localized phase (see the discussion in Sec. II and Refs. [12, 18, 54]). In the spin 'atomic' limit $\alpha \rightarrow 0$ (σ_i^z is conserved at each lattice site), the calculations necessary to adiabatically eliminate bosons using Eq. (4), can be straightforwardly performed. Assuming the bosons frozen into a localized state, we find that they contribute with a disordered shift to the constant trans-

verse field h_0 , in the effective spin hamiltonian

$$H'_s = \sum_i \left(h_0 + \frac{g^2}{\omega_i} \right) \sigma_i^z + \lambda \sum_i \sigma_i^z \sigma_{i+1}^z + \alpha \sum_{\langle i,j \rangle} \sigma_i^+ \sigma_j^-. \quad (13)$$

Considering in this case the hopping as a perturbation $V = \alpha \sum_{\langle i,j \rangle} \sigma_i^+ \sigma_j^-$, capable to connect states which otherwise would be localized, we can estimate the magnitude of the characteristic term of the locator expansion

$$\frac{\langle \alpha | V | \beta \rangle}{E_\alpha - E_\beta} \sim \frac{\alpha}{g^2 / \Omega \exp(-s_T \xi)}, \quad (14)$$

using an argument analogous to the one employed for (7). We can then conclude that the series would converge for

$$\alpha \lesssim g^2 / \Omega \exp(-\Sigma_T \xi), \quad (15)$$

and the system's eigenstates would remain close to the localized ones (as in Eq. (7), Σ_T is the entropy density at temperature T). It is straightforward to notice the analogies between the hamiltonians (6) and (13), and their respective results.

B. Delocalization of the disordered bosonic bath

In the complementary limit of spin dynamics occurring on faster time scales than the bosonic one, $g \ll h_0$, $\Omega \ll h_0$, we find that the impact of the clean, spin bath is to restore ergodicity in the disordered boson hamiltonian. As in Sec. III B, we assume that there is no bare bosonic hopping in (1), and integrate out the spin sector frozen in a delocalized state, deriving an effective tunnelling for bosons. Indeed, the effective bosonic hamiltonian reads in this case

$$H'_b = \sum_i \omega_i n_i - \sum_{\langle i,j \rangle} \frac{g^2}{h_0} (b_i^\dagger b_j + h.c.), \quad (16)$$

which again is a model of disordered free particles in one dimension, localized at any disorder strength Ω . A locator expansion using the on-site many-body bosonic interaction term as perturbation

$$V = U \sum_i n_i (n_i - 1), \quad (17)$$

leads to the estimate

$$U_c \sim \frac{g^2}{h_0} \log^3 \left(\frac{\Omega h_0}{g^2} \right), \quad (18)$$

for the magnitude of the critical interaction strength for delocalization, where in Eq. (18) we have employed the expression of ξ for a strongly disordered system one dimension, $\xi \propto \log \left(\frac{\Omega h_0}{g^2} \right)$, replacing the effective hopping $\propto g^2/h_0$ (compare with Sec. III B).

Analogue results to those discussed at the end of Sec. III B, are found if we consider the bosons in the localized phase coupled to a XX-quantum spin chain ($\lambda = 0$), and treating the spin-spin interaction along the \hat{z} direction in locator expansion. As expected, once again, the latter gets in turn localized by the former: eliminating the bosons imprints on-site disorder on the spins, which gets localized until the interaction strength λ ramps up to a threshold which would restore ergodicity in the system.

C. Coupling to thermal spins and dissipation

The generality of Eq. (4) allows us to adiabatically eliminate systems with both coherent and dissipative dynamics. This offers the opportunity to briefly comment on the fate of the Bose Hubbard model in (3) with on-site disorder of strength $\propto \Omega$ (as the one considered through this section), in contact with a system of spins at equilibrium with a thermal bath at temperature T . This can be, for instance, achieved adding to the simple spin hamiltonian in (3), on-site Lindblad operators

$$L_i = \sigma_i^+, \quad L_i = \sigma_i^-, \quad (19)$$

respectively with rates $\gamma_+ = \gamma n_T$ and $\gamma_- = \gamma(n_T + 1)$, where n_T is the spin thermal distribution at frequency h_0 and temperature T . We assume the scale separation h_0 , $\gamma_\pm \gg g$, and consider the spins frozen in their thermal state at temperature T . The adiabatic elimination of the spin sector adds an effective dissipation to the bosons (see for instance [57]), described by the dephasing jump operator, $\mathcal{L}_i = b_i^\dagger b_i$, and occurring at rate $\Gamma = \tilde{\gamma} g^2 / (\tilde{\gamma}^2 + h_0^2)$ where $\tilde{\gamma} = \gamma / \tanh(h_0/2T)$; the contribution to the bosonic hamiltonian is inconsequential, since the thermal state $\rho_T \sim e^{-\sum_i h_0 \sigma_i^z / T}$ is factorised into a product of local density matrices at each site, yielding a constant energy level shift to the disordered chemical potential, ω_i .

Unless $h_0 \gg \tilde{\gamma}$, which would render the dissipation rate Γ small, we arrive at the natural conclusion that, on time scales of the order $\sim 1/\Gamma$, the localized phase melts under the effect of the dephasing operators \mathcal{L}_i , without need to invoke a delocalization transition mediated by coherent many body interactions.

V. PERSPECTIVES

A. Experimental implementations

The model (1) discussed in this work originates from the coupling of two paradigmatic hamiltonians of condensed matter physics. In this Section, we briefly outline possible realizations of the many-body localization proximity effects presented here, from the perspective of quantum many body simulation using quantum optics platforms.

A first simple experimental implementation of our set-up consists in considering circuit QED-arrays [58], modelling a Jaynes-Cummings lattice with nearest neighbour photon hopping. The latter consists of a one-dimensional array of micro-wave resonators supporting photonic excitations at a given frequency set by the cavity modes; a linear dipole interaction of the type discussed in this work, H_{int} , converts bosons into excitations of superconducting qubits (spins one-half), each of them coupled to one of the resonators. Tunnelling among nearby photonic cavities implements a kinetic hopping term of the Hubbard type, leading to a competition, even in the clean system, among delocalization of photons, and the trapping of the associated polaritons, arising from effective on-site interactions [58]. In turn, the phase diagrams of the Jaynes-Cummings and of the Bose-Hubbard model bear several similarities [58], and it would be of interest to explore, in the laboratory, the impact of disorder present in individual cavity frequencies [59], using as guidelines the prediction of this work. Indeed, the increasing scalability of circuit QED lattice experiments with the consequent possibility to observe phase transitions [60], make in principle these hybrid qubit-photon architectures promising grounds to realise many-body localization proximity effects.

A more complex, yet closer, platform to implement the ideas presented in this work, are bosonic Hubbard models of photons on a lattice [61–63] (equipped with effective two-body photon-photon interactions arising from Kerr non-linearities), coupled to superconducting qubits with a photon-spin dipole interaction term; varying the photonic (or spin) modes from cavity to cavity (or qubit), one can simulate disorder in this system. Furthermore, when qubits are placed next to each other, electrostatic or magnetic interactions among them become relevant, mimicking the spin-spin interaction terms in our starting hamiltonian (1).

Another possibility to start with intrinsic spin interactions, as a term of the type $\sim \sum_i \sigma_i^z \sigma_{i+1}^z$, are Rydberg-dressed spin lattices coupled to interacting photons [64]. However, off-diagonal spin couplings terms, $\propto \sum_i \sigma_i^+ \sigma_{i+1}^-$, would be effectively generated only at higher orders in perturbation theory, changing some of the estimates contained in the previous Sections.

B. Future directions

Our results concerning the onset of many-body localization proximity effects in physically feasible platforms, are based on a combination of locator expansion and adiabatic elimination methods, and constitute a promising first route towards a systematic study of many-body localization in realistic systems made up of different species. As a first direction, we foresee a study of Anderson localization in a disordered Jaynes-Cummings lattice (see also discussion at the beginning of Sec. V A) using methods suited for one dimensional disordered systems, such as strong disorder renormalization group. At the next level of complexity, there is the challenging possibility to derive the effective strong disorder hamiltonian for the many-body spin-boson model in (1) using flow equation methods [65], which have already proven successful [66] in the description of the paradigmatic many-body localization transition of the XXZ quantum spin chain. Another promising avenue consists in going beyond our perturbative estimates with a real-space renormalisation group procedure [67, 68]; for instance Ref. [69] adopts, as in our work, an energy scale separation to derive as well an effective hamiltonian, and extract from the latter a flow of the probability distribution of the random couplings of a disordered, interacting quantum spin chain. This approach might lead to clearer characterisation of the many-body localized phases induced by proximity effect, which up to now has been characterised only through locator expansion techniques.

Finally, an agenda for future studies of the many-body localization transition in quantum optics, would require a systematic inclusion of dissipation since experimental implementations would be plagued at long times by photon losses or spin decoherence/dephasing terms (see for instance Sec. IV C), which would eventually spoil the features of the localized phase.

Acknowledgements– We thank I. Carusotto, R. Lewis-Swan and A. M. Rey for discussions on related topics. This material is based upon work supported by the Air Force Office of Scientific Research under award number FA9550-17-1-0183.

-
- [1] P. W. Anderson, “Absence of diffusion in certain random lattices,” *Phys. Rev.* **109**, 1492–1505 (1958).
 [2] E. Abrahams, P. W. Anderson, D. C. Licciardello, and T. V. Ramakrishnan, “Scaling theory of localization: Absence of quantum diffusion in two dimensions,” *Phys. Rev. Lett.* **42**, 673–676 (1979).
 [3] Boris L. Altshuler, Yuval Gefen, Alex Kamenev, and Leonid S. Levitov, “Quasiparticle lifetime in a finite system: A non-perturbative approach,” *Phys. Rev. Lett.* **78**,

- 2803–2806 (1997).
 [4] D.M. Basko, I.L. Aleiner, and B.L. Altshuler, “Metal-insulator transition in a weakly interacting many-electron system with localized single-particle states,” *Annals of Physics* **321**, 1126 – 1205 (2006).
 [5] I. V. Gornyi, A. D. Mirlin, and D. G. Polyakov, “Electron transport in a disordered luttinger liquid,” *Phys. Rev. B* **75**, 085421 (2007).
 [6] Arijeet Pal and David A. Huse, “Many-body localization

- phase transition,” *Phys. Rev. B* **82**, 174411 (2010).
- [7] M. Znidaric, T. Prosen, and P. Prelovsek, “Many-body localization in the heisenberg xxx magnet in a random field,” *Phys. Rev. B* **77**, 064426 (2008).
- [8] Vadim Oganesyan and David A. Huse, “Localization of interacting fermions at high temperature,” *Phys. Rev. B* **75**, 155111 (2007).
- [9] Rahul Nandkishore and David A. Huse, “Many-body localization and thermalization in quantum statistical mechanics,” *Annual Review of Condensed Matter Physics* **6**, 15–38 (2015).
- [10] Ronen Vosk, David A. Huse, and Ehud Altman, “Theory of the many-body localization transition in one-dimensional systems,” *Phys. Rev. X* **5**, 031032 (2015).
- [11] D. A. Abanin and Z. Papić, “Recent progress in many-body localization,” *ArXiv e-prints* (2017), arXiv:1705.09103 [cond-mat.dis-nn].
- [12] John Z. Imbrie, “Diagonalization and many-body localization for a disordered quantum spin chain,” *Phys. Rev. Lett.* **117**, 027201 (2016).
- [13] M. Schreiber, S. S. Hodgman, P. Bordia, H. P. Lüschen, M. H. Fischer, R. Vosk, E. Altman, U. Schneider, and I. Bloch, “Observation of many-body localization of interacting fermions in a quasi-random optical lattice,” *Science* **349**, 842–845 (2015).
- [14] Pranjali Bordia, Henrik P. Lüschen, Sean S. Hodgman, Michael Schreiber, Immanuel Bloch, and Ulrich Schneider, “Coupling identical one-dimensional many-body localized systems,” *Phys. Rev. Lett.* **116**, 140401 (2016).
- [15] J. Choi, S. Hild, J. Zeiher, P. Schauss, A. Rubio-Abadal, T. Yefsah, V. Khemani, D. A. Huse, I. Bloch, and C. Gross, “Exploring the many-body localization transition in two dimensions,” *Science* **352**, 1547–1552 (2016).
- [16] Jens H. Bardarson, Frank Pollmann, and Joel E. Moore, “Unbounded growth of entanglement in models of many-body localization,” *Phys. Rev. Lett.* **109**, 017202 (2012).
- [17] Maksym Serbyn, Z. Papić, and Dmitry A. Abanin, “Local conservation laws and the structure of the many-body localized states,” *Phys. Rev. Lett.* **111**, 127201 (2013).
- [18] David A. Huse, Rahul Nandkishore, and Vadim Oganesyan, “Phenomenology of fully many-body-localized systems,” *Phys. Rev. B* **90**, 174202 (2014).
- [19] V. Ros, M. Müller, and A. Scardicchio, “Integrals of motion in the many-body localized phase,” *Nuclear Physics B* **891**, 420 – 465 (2015).
- [20] A. Chandran, A. Pal, C. R. Laumann, and A. Scardicchio, *ArXiv e-prints*: 1605.00655.
- [21] Scott D. Geraedts, R. N. Bhatt, and Rahul Nandkishore, “Emergent local integrals of motion without a complete set of localized eigenstates,” *Phys. Rev. B* **95**, 064204 (2017).
- [22] S. A. Parameswaran and S. Gopalakrishnan, “Non-Fermi glasses: fractionalizing electrons at finite energy density,” *ArXiv e-prints* (2016), arXiv:1608.00981 [cond-mat.dis-nn].
- [23] Vedika Khemani, Rahul Nandkishore, and S. L. Sondhi, “Nonlocal adiabatic response of a localized system to local manipulations,” *Nat Phys* **11**, 560–565 (2015).
- [24] Sarang Gopalakrishnan, Markus Müller, Vedika Khemani, Michael Knap, Eugene Demler, and David A. Huse, “Low-frequency conductivity in many-body localized systems,” *Phys. Rev. B* **92**, 104202 (2015).
- [25] Jens H. Bardarson, Frank Pollmann, and Joel E. Moore, “Unbounded growth of entanglement in models of many-body localization,” *Phys. Rev. Lett.* **109**, 017202 (2012).
- [26] Scott D. Geraedts, Rahul Nandkishore, and Nicolas Regnault, “Many-body localization and thermalization: Insights from the entanglement spectrum,” *Phys. Rev. B* **93**, 174202 (2016).
- [27] Vedika Khemani, S. P. Lim, D. N. Sheng, and David A. Huse, “Critical properties of the many-body localization transition,” *Phys. Rev. X* **7**, 021013 (2017).
- [28] Z.-C. Yang, A. Hama, S. M. Giampaolo, E. R. Mucciolo, and C. Chamon, “Entanglement Complexity in Quantum Many-Body Dynamics, Thermalization and Localization,” *ArXiv e-prints* (2017), arXiv:1703.03420 [cond-mat.str-el].
- [29] S D Geraedts, N Regnault, and R M Nandkishore, “Characterizing the many-body localization transition using the entanglement spectrum,” *New Journal of Physics* **19**, 113021 (2017).
- [30] David A. Huse, Rahul Nandkishore, Vadim Oganesyan, Arijeet Pal, and S. L. Sondhi, “Localization-protected quantum order,” *Phys. Rev. B* **88**, 014206 (2013).
- [31] David Pekker, Gil Refael, Ehud Altman, Eugene Demler, and Vadim Oganesyan, “Hilbert-glass transition: New universality of temperature-tuned many-body dynamical quantum criticality,” *Phys. Rev. X* **4**, 011052 (2014).
- [32] Ronen Vosk and Ehud Altman, “Dynamical quantum phase transitions in random spin chains,” *Phys. Rev. Lett.* **112**, 217204 (2014).
- [33] R. Nandkishore and A. C. Potter, “Marginal anderson localization and many-body delocalization,” *Phys. Rev. B* **90**, 195115 (2014).
- [34] Rahul Nandkishore, “Many-body localization and delocalization in the two-dimensional continuum,” *Phys. Rev. B* **90**, 184204 (2014).
- [35] I. V. Gornyi, A. D. Mirlin, M. Müller, and D. G. Polyakov, “Absence of many-body localization in a continuum,” *ArXiv e-prints* (2016), arXiv:1611.05895 [cond-mat.dis-nn].
- [36] Rahul Nandkishore, “Many-body localization proximity effect,” *Phys. Rev. B* **92**, 245141 (2015).
- [37] Andrew C. Potter and Romain Vasseur, “Symmetry constraints on many-body localization,” *Phys. Rev. B* **94**, 224206 (2016).
- [38] Wojciech De Roeck and François Huveneers, “Asymptotic quantum many-body localization from thermal disorder,” *Communications in Mathematical Physics* **332**, 1017–1082 (2014).
- [39] Wojciech De Roeck, Francois Huveneers, Markus Müller, and Mauro Schiulaz, “Absence of many-body mobility edges,” *Phys. Rev. B* **93**, 014203 (2016).
- [40] Rahul Nandkishore and Sarang Gopalakrishnan, “Many body localized systems weakly coupled to baths,” *Annalen der Physik*, 1521-3889 (2016).
- [41] W. De Roeck and F. Huveneers, “Stability and instability towards delocalization in MBL systems,” *ArXiv e-prints* (2016), arXiv:1608.01815 [cond-mat.dis-nn].
- [42] S. A. Parameswaran and S. Gopalakrishnan, “Spin-catalyzed hopping conductivity in disordered strongly interacting quantum wires,” *Phys. Rev. B* **95**, 024201 (2017).
- [43] Rahul M. Nandkishore and S. L. Sondhi, “Many-body localization with long-range interactions,” *Phys. Rev. X* **7**, 041021 (2017).
- [44] Rahul Nandkishore, Sarang Gopalakrishnan, and David A. Huse, “Spectral features of a many-body-

- localized system weakly coupled to a bath,” Phys. Rev. B **90**, 064203 (2014).
- [45] Sarang Gopalakrishnan and Rahul Nandkishore, “Mean-field theory of nearly many-body localized metals,” Phys. Rev. B **90**, 224203 (2014).
- [46] Rahul Nandkishore and Sarang Gopalakrishnan, “Many body localized systems weakly coupled to baths,” Annalen der Physik , 1521–3889 (2016).
- [47] Sumilan Banerjee and Ehud Altman, “Variable-range hopping through marginally localized phonons,” Phys. Rev. Lett. **116**, 116601 (2016).
- [48] Mark H Fischer, Mykola Maksymenko, and Ehud Altman, “Dynamics of a many-body-localized system coupled to a bath,” Phys. Rev. Lett. **116**, 160401 (2016).
- [49] Emanuele Levi, Markus Heyl, Igor Lesanovsky, and Juan P. Garrahan, “Robustness of many-body localization in the presence of dissipation,” Phys. Rev. Lett. **116**, 237203 (2016).
- [50] M. Medvedyeva, T. Prosen, and M. Znidaric, “Influence of dephasing on many-body localization,” Phys. Rev. B **93**, 094205 (2016).
- [51] D. J. Luitz, F. Huveneers, and W. De Roeck, “How a small quantum bath can thermalize long localized chains,” Phys. Rev. Lett. **119**, 150602 (2017).
- [52] P. Ponte, C. Laumann, D. Huse, and A. Chandran, “Thermal inclusions: how one spin can destroy a many-body localized phase,” Philosophical Transactions A (2017).
- [53] H. P. Lüschen, P. Bordia, S. S. Hodgman, M. Schreiber, S. Sarkar, A. J. Daley, M. H. Fischer, E. Altman, I. Bloch, and U. Schneider, “Signatures of many-body localization in a controlled open quantum system,” Phys. Rev. X **7**, 011034 (2017).
- [54] J. Z Imbrie, V. Ros, and A. Scardicchio, arXiv:cond-mat 1609.08076 (2016).
- [55] H-P Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, 2007).
- [56] D. Marcos, A. Tomadin, S. Diehl, and P. Rabl, “Photon condensation in circuit qed by engineered dissipation,” New J. Phys. **14**, 055005 (2012).
- [57] G. S. Agarwal, R. R. Puri, and R. P. Singh, “Atomic schrödinger cat states,” Phys. Rev. A **56**, 2249–2254 (1997).
- [58] A. Houck, H. E. Tureci, and J. Koch, “On-chip quantum simulation with superconducting circuits,” Nature Physics **8**, 292 – 299 (2012).
- [59] D. L. Underwood, W. E. Shanks, J. Koch, and A. A. Houck, “Low-disorder microwave cavity lattices for quantum simulation with photons,” Phys. Rev. A **86**, 023837 (2012).
- [60] M. Fitzpatrick, N. M. Sundaresan, Andy C. Y. Li, J. Koch, and A. A. Houck, “Observation of a dissipative phase transition in a one-dimensional circuit qed lattice,” Phys. Rev. X **7**, 011016 (2017).
- [61] M. Hartmann, F. Brandao, and M. Plenio, “Quantum many-body phenomena in coupled cavity arrays,” Laser Photonics Rev. **2**, 527.
- [62] A. Tomadin and R. Fazio, “Many-body phenomena in qed-cavity arrays,” J. Opt. Soc. Am. B, **27**, A130 (2010).
- [63] D. E Chang, A. Sorensen, E. Demler, and M. Lukin, “A single-photon transistor using nanoscale surface plasmons,” Nature Physics, **3**, 807-812 (2007).
- [64] J. Zeiher and *et al.*, “Many-body interferometry of a rydberg-dressed spin lattice,” Nature Physics, **12**, 1095-1099 (2016).
- [65] David Pekker, Bryan K. Clark, Vadim Oganesyan, and Gil Refael, “Fixed points of wegner-wilson flows and many-body localization,” Phys. Rev. Lett. **119**, 075701 (2017).
- [66] S. Thomson and M. Schiro, arXiv:cond-mat , 1707.06981 (2017).
- [67] Chandan Dasgupta and Shang-keng Ma, “Low-temperature properties of the random heisenberg antiferromagnetic chain,” Phys. Rev. B **22**, 1305–1319 (1980).
- [68] Daniel S. Fisher, “Random antiferromagnetic quantum spin chains,” Phys. Rev. B **50**, 3799–3821 (1994).
- [69] Ronen Vosk and Ehud Altman, “Many-body localization in one dimension as a dynamical renormalization group fixed point,” Phys. Rev. Lett. **110**, 067204 (2013).