
# SPACES WITH COMPLEXITY ONE

ALYSON BITTNER

ABSTRACT. An A-cellular space is a space built from A and its suspensions, analogously to the way that CW-complexes are built from  $S^0$  and its suspensions. The A-cellular approximation of a space X is an A-cellular space  $CW_A X$  which is closest to X among all A-cellular spaces. The A-complexity of a space X is an ordinal number that quantifies how difficult it is to build an A-cellular approximation of X. In this paper, we study spaces with low complexity. In particular we show that if A is a sphere localized at a set of primes then the A-complexity of each space X is at most 1.

# 1. INTRODUCTION

Let A be a pointed CW-complex. An A-cellular space is a space built out of copies of A via iterated pointed homotopy colimit construction. Given a pointed space X, the A-cellular approximation of X is an A-cellular space  $CW_A X$  equipped with a map  $CW_A X \to X$ , such that the induced map of pointed mapping complexes  $Map_*(A, CW_A X) \to Map_*(A, X)$  is a weak equivalence. For example,  $S^0$ -cellular spaces are CW-complexes and an  $S^0$ -cellular approximation of a space X is the CW-approximation of X. Farjoun [3] showed that the A-cellular approximation of X exists for any A and X and that it is unique up to homotopy equivalence. The space  $CW_A X$  is the best possible approximation of X in the class of A-cellular spaces: for any A-cellular space Y and a map  $Y \to X$  there exists a map  $Y \to CW_A X$ , unique up to homotopy, such that the following diagram is homotopy commutative.



In [2] Chachólski, Dwyer, and Intermont introduced the concept of the Acomplexity of a space X, which is the minimum ordinal number of homotopy colimits necessary to produce  $CW_A(X)$  from copies of A. More precisely, starting with the full subcategory  $\mathbf{C}_0$  of pointed spaces whose objects have the homotopy type of a retract of wedges of A, one can construct an increasing chain of categories indexed by ordinal numbers  $\alpha$ , where  $\mathbf{C}_{\alpha}$  is the category of all spaces of the homotopy type of hocolim<sub>D</sub> F for some small category **D** and a functor  $F : \mathbf{D} \to \bigcup_{\beta < \alpha} \mathbf{C}_{\beta}$  and their retracts. The the A-complexity of X is the minimal ordinal  $\kappa_A(X)$  such that  $X \in \mathbf{C}_{\kappa_A(X)}$ .

In the case of  $A = S^0$  the the category  $\mathbf{C}_1$  consists of all space of a homotopy type of a CW-complex, which implies that  $\kappa_{S^0}(X) = 1$  for any space X. This is however not the case for other choices of A, and in general the complication of a

#### ALYSON BITTNER

space need not even be finite. For example, if  $A = M(\mathbb{Z}/p, n)$  is a Moore space for some  $n \geq 1$  and a prime p and  $X = M(\mathbb{Z}/p^{\infty}, n+1)$  then  $\kappa_A(X) = \omega$  (see [2, Proposition 9.3]). By contrast, Chachólski, Dwyer and Intermont observed that if  $A = S^n$  then  $\kappa_A(X) \leq 2$  for all X and suggested that it should be possible to lower this bound and show that  $\kappa_A(X) \leq 1$  [2, §9.3]. The goal of this note is to show that this is indeed the case, and that this result holds in even greater generality:

**Theorem 1.1.** If A is a sphere, or a sphere localized at a set of primes then  $\kappa_A(X) \leq 1$  for all spaces X.

## Acknowledgements

I want to express my appreciation to my advisor, Bernard Badzioch, for his helpful comments which led to the final version of paper. Additionally, I would like to thank Matthew Sartwell for several useful conversations.

#### 2. Algebras of mapping spaces

Our proof of Theorem 1.1 will be based on the following algebraic description of mapping space obtained by Sartwell [5]. For spaces X, Y let  $\operatorname{Map}^{\Delta}_{*}(X, Y)$  denote the simplicial mapping complex of pointed maps  $X \to Y$ . Given a pointed CW-complex A let  $\mathbf{T}^{A}$  denote a simplicial category on objects  $T_{n}$  for  $n = 0, 1, \ldots$ , and such that

$$\operatorname{Hom}_{\mathbf{T}^{A}}(T_{n},T_{m}) = \operatorname{Map}^{\Delta}_{*}\left(\bigvee^{m} A,\bigvee^{n} A\right)$$

Notice that in this category  $T_n$  is the *n*-fold product of  $T_1$ . A  $\mathbf{T}^A$ -algebra is a product preserving simplicial functor  $\mathbf{T}^A \to \mathbf{sSet}_*$  where  $\mathbf{sSet}_*$  is the category of pointed simplicial sets. Let  $\mathbf{Alg}^{\mathbf{T}^A}$  denote the category of all  $\mathbf{T}^A$ -algebras with natural transformations as morphisms.

Notice that any pointed space X defines a  $\mathbf{T}^A$ -algebra  $\Omega^A(X)$  such that

$$\Omega^{A}(X)(T_{n}) = \operatorname{Map}^{\Delta}_{*}\left(\bigvee^{n} A, X\right)$$

The resulting functor  $\Omega^A \colon \mathbf{Top}_* \to \mathbf{Alg}^{\mathbf{T}^A}$  has a left adjoint  $B^A$ .

The category  $\operatorname{Alg}^{\mathbf{T}^{A}}$  can be equipped with a model category structure where weak equivalences and fibrations are defined as objectwise weak homotopy equivalence and Serre fibrations respectively. Denote also by  $\mathbf{R}^{A}\mathbf{Top}_{*}$  the category  $\mathbf{Top}_{*}$  taken with the model category structure where fibrations are Serre fibrations and a weak equivalences are maps  $f: X \to Y$  that induce a weak equivalences  $f_{*}: \operatorname{Map}^{\Delta}_{*}(A, X) \to \operatorname{Map}^{\Delta}_{*}(A, Y)$ . In other words,  $\mathbf{R}^{A}\mathbf{Top}_{*}$  is obtained by taking the right Bousfield localization of the usual model category structure on  $\mathbf{Top}_{*}$  with respect to the space A as in [4, 5.1.1]. Sartwell showed that the following holds:

**Theorem 2.1.** [5] Let A be a sphere or a sphere localized at a set of primes. The adjoint pair

$$B^A \colon \mathbf{Alg}^{\mathbf{T}^A} \rightleftharpoons \mathbf{R}^A \mathbf{Top}_* \colon \Omega^A$$

is a Quillen equivalence.

### 3. Proof of Theorem 1.1

Directly from the definition of an A-cellular approximation it follows that if spaces X and Y are weakly equivalent in  $R^A$ **Top** then  $CW_A X$  and  $CW_A Y$  are weakly homotopy equivalent, and so  $\kappa_A(X) = \kappa_A(Y)$ . Let A be a space as in Theorem 1.1. By Theorem 2.1 for any space X we get a weak equivalence  $X \simeq B^A Q \Omega^A X$ in  $R^A$ **Top**, where  $Q\Omega^A X$  denoted a cofibrant replacement in the category  $\mathbf{Alg}^{\mathbf{T}^A}$ . Therefore it is only necessary to show that for any space X,  $\kappa_A(B^A Q \Omega^A X) \leq 1$ .

The algebra  $Q\Omega^A X$  can be described more explicitly as follows. Let  $U : \mathbf{Alg}^{\mathbf{T}^A} \to \mathbf{sSet}_*$  denote the forgetful functor,  $U(\Phi) = \Phi(T_1)$ . This functor has a left adjoint  $F : \mathbf{sSet}_* \to \mathbf{Alg}^{\mathbf{T}^A}$ . For  $\Phi \in \mathbf{T}^A$ -algebra, let  $FU_{\bullet}\Phi$  be the simplicial  $\mathbf{T}^A$ -algebra defined by adjoint pair (F, U). Explicitly:

$$FU_n\Phi := (FU)^{n+1}\Phi$$

By [5, 2.3.3] the natural map  $|FU_{\bullet}\Phi| \to \Phi$  is a cofibrant replacement of  $\Phi$  in the category  $\mathbf{Alg}^{\mathbf{T}^{A}}$ .

In view of this fact we need to show that for any space X the A-complexity of  $B^A|FU_{\bullet}\Omega^A X|$  is at most one. The left adjoint  $B^A$  can be described explicitly as the coend  $B^A(\Phi) = \int^{T_n \in T^A} \bigvee_n A \wedge |\Phi(T_n)|$ , which is in a similar manner as done in [1, 3.4] but then composing with objectwise singularization and realization to obtain the desired adjunction [4, 18.3.10]. The functor  $B^A F: \mathbf{sSet}_* \to \mathbf{Top}_*$  is left adjoint to  $U\Omega^A$ , and since  $U\Omega^A(X) = \mathrm{Map}^{\Delta}_*(A, X)$ , thus  $B^A F(Y) = A \wedge |Y|$ . This gives:

$$B^{A}|FU_{\bullet}\Omega^{A}X| \simeq |B^{A}FU_{\bullet}\Omega^{A}X| \simeq \operatorname{hocolim}_{n\in\Delta^{op}}A \wedge |(FU)^{n}\Omega^{A}X(T_{1})|$$
$$\simeq \operatorname{hocolim}_{(n,m)\in\Delta^{op}\times\Delta^{op}}A \wedge ((FU)^{n}\Omega^{A}X(T_{1}))_{m}$$
$$\overset{\sigma\in((FU)^{n}\Omega^{A}X(T_{1}))_{m}}{\simeq \operatorname{hocolim}_{(n,m)\in\Delta^{op}\times\Delta^{op}}}\bigvee A$$

### References

- Bernard Badzioch, Kuerak Chung, and Alexander A Voronov. The canonical delooping machine. Journal of Pure and Applied Algebra, 208(2):531–540, 2007.
- Wojciech Chachólski, WG Dwyer, and M Intermont. The a-complexity of a space. Journal of the London Mathematical Society, 65(1):204-222, 2002.
- [3] Emmanuel D. Farjoun. Cellular Spaces, Null Spaces and Homotopy Localization. Springer, 1995.
- [4] Philip S Hirschhorn. Model Categories and Their Localizations. Number 99. American Mathematical Soc., 2009.
- [5] Matthew Sartwell. Detecting Mapping Spaces and Derived Equivalences of Algebraic Theories. PhD thesis, University at Buffalo, 2016.