

# On the geodesics of homogeneous Finsler spaces with some special $(\alpha, \beta)$ -metrics

Gauree Shanker and Kirandeep Kaur

Department of Mathematics and Statistics

School of Basic and Applied Sciences

Central University of Punjab, Bathinda, Punjab-151001, India

Email: gshankar@cup.ac.in, kirandeep.kaur@cup.ac.in

## Abstract

In this paper, we study geodesics and geodesic vectors for homogeneous exponential Finsler space and homogeneous infinite series Finsler space. Further, we find necessary and sufficient condition for a non-zero vector in these homogeneous spaces to be a geodesic vector.

**2010 Mathematics Subject Classification:** 22E60, 53C30, 53C60.

**Keywords and Phrases:** Homogeneous Finsler space, homogeneous geodesic, infinite series  $(\alpha, \beta)$ -metric, exponential metric, g.o. space.

## 1 Introduction

According to S. S. Chern ([6]), Finsler geometry is just the Riemannian geometry without the quadratic restriction. Finsler generalized Riemann's theory in his doctoral thesis ([11]), but his name was established in differential geometry by Cartan ([5]). In 1972, M. Matsumoto ([25]) has introduced the concept of  $(\alpha, \beta)$ -metric in Finsler geometry. A Finsler metric of the form  $F = \alpha\phi(s)$ ,  $s = \frac{\beta}{\alpha}$ , where  $\alpha = \sqrt{a_{ij}(x)y^i y^j}$  is induced by a Riemannian metric  $\tilde{a} = a_{ij}dx^i \otimes dx^j$  on a connected smooth  $n$ -manifold  $M$  and  $\beta = b_i(x)y^i$  is a 1-form on  $M$  is called an  $(\alpha, \beta)$ -metric. There are various applications of  $(\alpha, \beta)$ -metrics in information geometry ([1]), physics and biology ([2]). Some notable contributions on Finsler spaces with exponential and infinite series  $(\alpha, \beta)$ -metrics can be seen in ([26], [27], [28], [29], [30]).

Geodesic in a Finsler manifold is the generalization of notion of a straight line in an Euclidean space. Geodesic can be viewed as a curve that minimizes the distance between two points on the manifold. A geodesic in a homogeneous Finsler space  $(G/H, F)$  is called homogeneous geodesic if it is an orbit of a one-parameter subgroup of  $G$ . Homogeneous geodesics on homogeneous Riemannian manifolds have been studied by many authors ([12], [18], [19], [34]). The existence of homogeneous geodesics in homogeneous Riemannian manifolds is quite an interesting problem. In ([15]), Kajzer prove that there exist atleast one homogeneous geodesic in a Riemannian manifold. Further, in ([31], [32]), Szenthe prove that there exist infinitely many homogeneous geodesics through identity in a Riemannian manifold  $(G, \alpha)$ , where  $G$  is compact semi-simple Lie group of rank  $\geq 2$ . Later, in ([20]), Kowalski and Szenthe prove the existence of atleast one homogeneous geodesic through each point for any homogeneous Riemannian manifold. Also, Kowalski and Vlasek ([21]) have shown that the result proved in ([20]) is not true in general. They have given some examples of homogeneous Riemannian manifolds of dimension  $\geq 4$  admitting only one homogeneous geodesic. There are many applications of Homogeneous geodesics in mechanics. Arnold ([3]) study the geodesics of left invariant Riemannian metrics on Lie groups, extending Euler's theory of rigid body motion and called homogeneous geodesic as "relative equilibria". The equation of motion of many systems in classical mechanics reduces to the geodesic equation in an appropriate Riemannian manifold. For example, Tóth ([33]) study trajectories which are orbits of a one-parameter symmetry group in case of Lagrangian and Hamiltonian systems. Also, Lacomba ([22]) use homogeneous geodesics in the work of Smale's mechanical systems.

Latifi ([23]) extend the concept of homogeneous geodesics in homogeneous Finsler spaces. In ([23]), he has given a criterion for charaterization of geodesic vectors. Latifi and Razavi ([24]) study homogeneous geodesics in a 3-dimensional connected Lie group with a left invariant Randers metric and show that all the geodesics on spaces equipped with such metrics are homogeneous. Habibi, Latifi and Toomanian ([13]) have extended Szenthe's result of homogeneous geodesics for invariant Finsler metrics. Yan and Deng ([36]) generalize this result and prove that there exists atleast one homogeneous geodesic through each point for any compact homogeneous Finsler space and also extend Kowalski and Szenthe's result to the Randers space. Yan ([37]) prove the existence of atleast one homogeneous geodesic through each point for a homogeneous Finsler space of odd dimension. Hosseini and Moghaddam ([14]) study the existence of homogeneous geodesic in homoge-

neous  $(\alpha, \beta)$ -spaces. Dušek ([10]) indicates a gap in the proof of main result of ([37]) and reproves this result and also studies homogeneous geodesics on a homogeneous Berwald space or homogeneous reversible Finsler space. Recently in 2018, Yan and Huang ([38]) prove that any homogeneous Finsler space admits atleast one homogeneous geodesic through each point.

## 2 Preliminaries

**DEFINITION 2.1.** Let  $V$  be an  $n$ -dimensional real vector space. It is called a Minkowski space if there exists a real valued function  $F : V \rightarrow \mathbb{R}$  satisfying the following conditions:

- (a)  $F$  is smooth on  $V \setminus \{0\}$ ,
- (b)  $F(v) \geq 0 \quad \forall v \in V$ ,
- (c)  $F$  is positively homogeneous, i.e.,  $F(\lambda v) = \lambda F(v), \quad \forall \lambda > 0$ ,
- (d) For a basis  $\{v_1, v_2, \dots, v_n\}$  of  $V$  and  $y = y^i v_i \in V$ , the Hessian matrix  $(g_{ij}) = \left( \frac{1}{2} F_{y^i y^j}^2 \right)$  is positive-definite at every point of  $V \setminus \{0\}$ .

In this case,  $F$  is called a Minkowski norm.

**DEFINITION 2.2.** Let  $M$  be a connected smooth manifold. If there exists a function  $F : TM \rightarrow [0, \infty)$  such that  $F$  is smooth on the slit tangent bundle  $TM \setminus \{0\}$  and the restriction of  $F$  to any  $T_p(M)$ ,  $p \in M$ , is a Minkowski norm, then  $(M, F)$  is called a **Finsler space** and  $F$  is called a Finsler metric.

Let  $(M, F)$  be a Finsler space and let  $(x^i, y^i)$  be a standard coordinate system in  $T_x(M)$ . The induced inner product  $g_y$  on  $T_x(M)$  is given by  $g_y(u, v) = g_{ij}(x, y) u^i v^j$ , where  $u = u^i \frac{\partial}{\partial x^i}$ ,  $v = v^i \frac{\partial}{\partial x^i} \in T_x(M)$ . Also note that  $F(x, y) = \sqrt{g_y(y, y)}$ .

Shen ([7]) has given the condition for an  $(\alpha, \beta)$ -metric to be a Finsler metric in following lemma:

**LEMMA 2.1.** Let  $F = \alpha\phi(s)$ ,  $s = \beta/\alpha$ , where  $\alpha$  is a Riemannian metric and  $\beta$  is a 1-form whose length with respect to  $\alpha$  is bounded above, i.e.,  $b := \|\beta\|_\alpha < b_0$ , where  $b_0$  is a positive real number. Then  $F$  is a Finsler

metric if and only if the function  $\phi = \phi(s)$  is a smooth positive function on  $(-b_0, b_0)$  and satisfies the following condition:

$$\phi(s) - s\phi'(s) + (b^2 - s^2)\phi''(s) > 0, \quad |s| \leq b < b_0.$$

**DEFINITION 2.3.** A diffeomorphism  $\phi: M \rightarrow M$  on a Finsler space  $(M, F)$  is called an **isometry** if  $F(p, X) = F(\phi(p), d\phi_p(X))$ , for any  $p \in M$  and  $X \in T_p(M)$ .

**DEFINITION 2.4.** A Lie group  $G$  is called a Lie transformation group of a smooth manifold  $M$  if it has a smooth action on  $M$ .

**DEFINITION 2.5.** Let  $(M, F)$  be a connected Finsler space. If the action of the group of isometries of  $(M, F)$ , denoted by  $I(M, F)$ , is transitive on  $M$ , then it is called a homogeneous Finsler space.

Let  $(M, Q)$  be a Riemannian manifold. A geodesic  $\sigma: \mathbb{R} \rightarrow M$  is called homogeneous geodesic if there exists a one-parameter group of isometries  $\phi: \mathbb{R} \times M \rightarrow M$  such that  $\sigma(t) = \phi(t, \sigma(0))$ ,  $t \in \mathbb{R}$ . If all the geodesics of a Riemannian manifold are homogeneous, then it is called a g.o. space (geodesic orbit space). Every naturally reductive Riemannian manifold is a g.o. space. The first example of a g.o. space which is not naturally reductive was given by Kaplan ([16]).

In 2014, Yan and Deng ([35]) have studied Finsler g.o. spaces defined as follows:

A Finsler space  $(M, F)$  is called a Finsler g.o. space if every geodesic of  $(M, F)$  is the orbit of a one-parameter subgroup of  $G = I(M, F)$ , i.e., if  $\sigma: \mathbb{R} \rightarrow M$  is a geodesic, then  $\exists$  a non-zero vector  $Z \in \mathfrak{g} = \text{Lie}(G)$  and  $p \in M$  such that  $\sigma(t) = \exp(tZ).p$ . A Finsler g.o. space has vanishing S-curvature. Also, note that every Finsler g.o. space is homogeneous.

A homogeneous Finsler space is a g.o. space if and only if the projections of all the geodesic vectors cover the set  $T_{eH}(G/H) - \{0\}$ .

**DEFINITION 2.6.** Let  $(G/H, F)$  be a homogeneous Finsler space and  $e$  be the identity of  $G$ . A geodesic  $\sigma(t)$  through the origin  $eH$  of  $G/H$  is called homogeneous if it is an orbit of a one-parameter subgroup of  $G$ , i.e., there exists a non-zero vector  $X \in \mathfrak{g} = \text{Lie}(G)$  such that  $\sigma(t) = \exp(tX).eH$ ,  $t \in \mathbb{R}$ .

### 3 Geodesic vector

The problem of studying homogeneous geodesics of a homogeneous space is basically the study of its geodesic vectors.

Let  $(M, F)$  be a homogeneous Finsler space. Then,  $M$  can be written as a coset space  $G/H$ , where  $G = I(M, F)$  is a Lie transformation group of  $M$  and  $H$ , the compact isotropy subgroup of  $I(M, F)$  at some point  $x \in M$  ([9]). Let  $\mathfrak{g}$  and  $\mathfrak{h}$  be the Lie algebras of the Lie groups  $G$  and  $H$  respectively. Also, let  $\mathfrak{m}$  be a subspace of  $\mathfrak{g}$  such that  $\text{Ad}(h)\mathfrak{m} \subset \mathfrak{m} \quad \forall h \in H$ , and  $\mathfrak{g} = \mathfrak{h} + \mathfrak{m}$  be a reductive decomposition of  $\mathfrak{g}$ .

Observe that for any  $Y \in \mathfrak{g}$ , the vector field  $Y^* = \frac{d}{dt} (\exp(tY)H) \Big|_{t=0}$  is called the fundamental Killing vector field on  $G/H$  generated by  $Y$  ([17]). The canonical projection  $\pi: G \longrightarrow G/H$  induces an isomorphism between the subspace  $\mathfrak{m}$  and the tangent space  $T_{eH}(G/H)$  through the following map:

$$\begin{aligned} \mathfrak{m} &\longrightarrow T_{eH}(G/H) \\ v &\longrightarrow \frac{d}{dt} (\exp(tv)H)|_{t=0}. \end{aligned}$$

We have  $d\pi(Y_{\mathfrak{m}}) = Y_{eH}^*$ . Using the natural identification and scalar product  $g_{Y_{eH}^*}$  on  $T_{eH}(G/H)$ , we get a scalar product  $g_{Y_{\mathfrak{m}}}$  on  $\mathfrak{m}$ .

**DEFINITION 3.1.** *Let  $(G/H, F)$  be a homogeneous Finsler space and  $e$  be the identity of  $G$ . A non-zero vector  $X \in \mathfrak{g}$  is called a geodesic vector if the curve  $\exp(tX).eH$  is a geodesic of  $(G/H, F)$ .*

The following result proved in ([23]) gives a criterion for a non-zero vector to be a geodesic vector in a homogeneous Finsler space.

**LEMMA 3.1.** *A non-zero vector  $Y \in \mathfrak{g}$  is a geodesic vector if and only if*

$$g_{Y_{\mathfrak{m}}}(Y_{\mathfrak{m}}, [Y, Z]_{\mathfrak{m}}) = 0 \quad \forall Z \in \mathfrak{g}.$$

Next, we deduce necessary and sufficient condition for a nonzero vector in a homogeneous Finsler space with infinite series  $(\alpha, \beta)$ -metric to be a geodesic vector.

**Theorem 3.1.** *Let  $G/H$  be a homogeneous Finsler space with infinite series metric  $F = \frac{\beta^2}{\beta - \alpha}$  given by an invariant Riemannian metric  $\tilde{a}$  and an invariant vector field  $\tilde{X}$  such that  $\tilde{X}(H) = X$ . Then, a non-zero vector  $Y \in \mathfrak{g}$*

is a geodesic vector if and only if

$$\begin{aligned}
& \frac{\langle X, Y_m \rangle^3}{(\langle X, Y_m \rangle - \sqrt{\langle Y_m, Y_m \rangle})^4} \times \\
& \left[ \langle X, [Y, Z]_m \rangle \left\{ \langle X, Y_m \rangle^2 - 4 \langle Y_m, Y_m \rangle^{3/2} + \langle X, Y_m \rangle \sqrt{\langle Y_m, Y_m \rangle} + 2 \langle Y_m, Y_m \rangle \right\} \right. \\
& \left. + \langle Y_m, [Y, Z]_m \rangle \left\{ \frac{\langle X, Y_m \rangle^2}{\sqrt{\langle Y_m, Y_m \rangle}} - \langle X, Y_m \rangle \right\} \right] = 0. \tag{1}
\end{aligned}$$

*Proof.* Using lemma 3.1 of ([28]), we can write

$$F(Y) = \frac{\langle X, Y \rangle^2}{\langle X, Y \rangle - \sqrt{\langle Y, Y \rangle}}.$$

Also, we know that

$$g_Y(U, V) = \frac{1}{2} \frac{\partial^2}{\partial s \partial t} F^2(Y + sU + tV) \Big|_{s=t=0}.$$

After some calculations, we get

$$\begin{aligned}
g_Y(U, V) &= \frac{\langle X, Y \rangle^2}{(\langle X, Y \rangle - \sqrt{\langle Y, Y \rangle})^4} \left[ \langle X, Y \rangle^2 \langle X, V \rangle \langle X, U \rangle - 4 \langle Y, Y \rangle^{3/2} \langle X, V \rangle \langle X, U \rangle \right. \\
&\quad + 6 \langle Y, Y \rangle \langle X, V \rangle \langle X, U \rangle + \frac{\langle X, Y \rangle^2 \langle X, V \rangle \langle U, Y \rangle}{\sqrt{\langle Y, Y \rangle}} - 4 \langle X, Y \rangle \langle X, V \rangle \langle U, Y \rangle \\
&\quad - \frac{\langle X, Y \rangle^3 \langle U, Y \rangle \langle V, Y \rangle}{\langle Y, Y \rangle^{3/2}} + \frac{\langle X, Y \rangle^3 \langle U, V \rangle}{\sqrt{\langle Y, Y \rangle}} + \frac{4 \langle X, Y \rangle^2 \langle U, Y \rangle \langle V, Y \rangle}{\langle Y, Y \rangle} \\
&\quad \left. - \langle X, Y \rangle^2 \langle U, V \rangle + \frac{\langle X, Y \rangle^2 \langle X, U \rangle \langle V, Y \rangle}{\sqrt{\langle Y, Y \rangle}} - 4 \langle X, Y \rangle \langle X, U \rangle \langle V, Y \rangle \right]. \tag{2}
\end{aligned}$$

From above equation, we can write

$$\begin{aligned}
g_{Y_m}(Y_m, [Y, Z]_m) &= \frac{\langle X, Y_m \rangle^2}{(\langle X, Y_m \rangle - \sqrt{\langle Y_m, Y_m \rangle})^4} \left[ \langle X, Y_m \rangle^2 \langle X, [Y, Z]_m \rangle \langle X, Y_m \rangle \right. \\
&\quad - 4 \langle Y_m, Y_m \rangle^{3/2} \langle X, [Y, Z]_m \rangle \langle X, Y_m \rangle + 6 \langle Y_m, Y_m \rangle \langle X, [Y, Z]_m \rangle \langle X, Y_m \rangle \\
&\quad + \frac{\langle X, Y_m \rangle^2 \langle X, [Y, Z]_m \rangle \langle Y_m, Y_m \rangle}{\sqrt{\langle Y_m, Y_m \rangle}} - 4 \langle X, Y_m \rangle \langle X, [Y, Z]_m \rangle \langle Y_m, Y_m \rangle \\
&\quad - \frac{\langle X, Y_m \rangle^3 \langle Y_m, Y_m \rangle \langle [Y, Z]_m, Y_m \rangle}{\langle Y_m, Y_m \rangle^{3/2}} + \frac{\langle X, Y_m \rangle^3 \langle Y_m, [Y, Z]_m \rangle}{\sqrt{\langle Y_m, Y_m \rangle}} \\
&\quad + \frac{4 \langle X, Y_m \rangle^2 \langle Y_m, Y_m \rangle \langle [Y, Z]_m, Y_m \rangle}{\langle Y_m, Y_m \rangle} - \langle X, Y_m \rangle^2 \langle Y_m, [Y, Z]_m \rangle \\
&\quad \left. + \frac{\langle X, Y_m \rangle^2 \langle X, Y_m \rangle \langle [Y, Z]_m, Y_m \rangle}{\sqrt{\langle Y_m, Y_m \rangle}} - 4 \langle X, Y_m \rangle \langle X, Y_m \rangle \langle [Y, Z]_m, Y_m \rangle \right] \\
&= \frac{\langle X, Y_m \rangle^3}{(\langle X, Y_m \rangle - \sqrt{\langle Y_m, Y_m \rangle})^4} \times \\
&\quad \left[ \langle X, [Y, Z]_m \rangle \left\{ \langle X, Y_m \rangle^2 - 4 \langle Y_m, Y_m \rangle^{3/2} + \langle X, Y_m \rangle \sqrt{\langle Y_m, Y_m \rangle} + 2 \langle Y_m, Y_m \rangle \right\} \right. \\
&\quad \left. + \langle Y_m, [Y, Z]_m \rangle \left\{ \frac{\langle X, Y_m \rangle^2}{\sqrt{\langle Y_m, Y_m \rangle}} - \langle X, Y_m \rangle \right\} \right]. \tag{3}
\end{aligned}$$

Now, from lemma (3.1),  $Y \in \mathfrak{g}$  is a geodesic vector if and only if

$$g_{Y_m}(Y_m, [Y, Z]_m) = 0, \forall Z \in \mathfrak{m}.$$

Therefore  $g_{Y_m}(Y_m, [Y, Z]_m) = 0$  if and only if equation (1) holds.  $\square$

**COROLLARY 3.1.** *Let  $(G/H, F)$  be a homogeneous Finsler space with infinite series metric  $F = \frac{\beta^2}{\beta - \alpha}$  defined by an invariant Riemannian metric  $\langle \cdot, \cdot \rangle$  and an invariant vector field  $\tilde{X}$  such that  $\tilde{X}(H) = X$ . Let  $Y \in \mathfrak{g}$  be a vector such that  $\langle X, [Y, Z]_m \rangle = 0 \forall Z \in \mathfrak{m}$ . Then  $Y$  is a geodesic vector of  $(G/H, \langle \cdot, \cdot \rangle)$  if and only if  $Y$  is a geodesic vector of  $(G/H, F)$ .*

*Proof.* From equation (3), we can write

$$\begin{aligned}
g_{Y_m}(Y_m, [Y, Z]_m) &= \frac{\langle X, Y_m \rangle^3}{\left( \langle X, Y_m \rangle - \sqrt{\langle Y_m, Y_m \rangle} \right)^4} \times \\
&\quad \left[ \langle X, [Y, Z]_m \rangle \left\{ \langle X, Y_m \rangle^2 - 4 \langle Y_m, Y_m \rangle^{3/2} + \langle X, Y_m \rangle \sqrt{\langle Y_m, Y_m \rangle} + 2 \langle Y_m, Y_m \rangle \right\} \right. \\
&\quad \left. + \langle Y_m, [Y, Z]_m \rangle \left\{ \frac{\langle X, Y_m \rangle^2}{\sqrt{\langle Y_m, Y_m \rangle}} - \langle X, Y_m \rangle \right\} \right] \\
&= \frac{\langle X, Y_m \rangle^4 \langle Y_m, [Y, Z]_m \rangle}{\left( \langle X, Y_m \rangle - \sqrt{\langle Y_m, Y_m \rangle} \right)^3 \sqrt{\langle Y_m, Y_m \rangle}}, \text{ because } \langle X, [Y, Z]_m \rangle = 0.
\end{aligned}$$

Therefore  $g_{Y_m}(Y_m, [Y, Z]_m) = 0$  if and only if  $\langle Y_m, [Y, Z]_m \rangle = 0$ .  $\square$

**Theorem 3.2.** Let  $(G/H, F)$  be a homogeneous Finsler space with infinite series metric  $F = \frac{\beta^2}{\beta - \alpha}$  defined by an invariant Riemannian metric  $\langle \cdot, \cdot \rangle$  and an invariant vector field  $\tilde{X}$  such that  $\tilde{X}(H) = X$ . Then  $X$  is a geodesic vector of  $(G/H, \langle \cdot, \cdot \rangle)$  if and only if  $X$  is a geodesic vector of  $(G/H, F)$ .

*Proof.* From equation (2), we can write

$$\begin{aligned}
g_X(X, [X, Z]_{\mathfrak{m}}) &= \frac{\langle X, X \rangle^2}{\left(\langle X, X \rangle - \sqrt{\langle X, X \rangle}\right)^4} \left[ \langle X, X \rangle^2 \langle X, [X, Z]_{\mathfrak{m}} \rangle \langle X, X \rangle \right. \\
&\quad - 4 \langle X, X \rangle^{3/2} \langle X, [X, Z]_{\mathfrak{m}} \rangle \langle X, X \rangle + 6 \langle X, X \rangle \langle X, [X, Z]_{\mathfrak{m}} \rangle \langle X, X \rangle \\
&\quad + \frac{\langle X, X \rangle^2 \langle X, [X, Z]_{\mathfrak{m}} \rangle \langle X, X \rangle}{\sqrt{\langle X, X \rangle}} - 4 \langle X, X \rangle \langle X, [X, Z]_{\mathfrak{m}} \rangle \langle X, X \rangle \\
&\quad - \frac{\langle X, X \rangle^3 \langle X, X \rangle \langle [X, Z]_{\mathfrak{m}}, X \rangle}{\langle X, X \rangle^{3/2}} + \frac{\langle X, X \rangle^3 \langle X, [X, Z]_{\mathfrak{m}} \rangle}{\sqrt{\langle X, X \rangle}} \\
&\quad + \frac{4 \langle X, X \rangle^2 \langle X, X \rangle \langle [X, Z]_{\mathfrak{m}}, X \rangle}{\langle X, X \rangle} - \langle X, X \rangle^2 \langle X, [X, Z]_{\mathfrak{m}} \rangle \\
&\quad \left. + \frac{\langle X, X \rangle^2 \langle X, X \rangle \langle [X, Z]_{\mathfrak{m}}, X \rangle}{\sqrt{\langle X, X \rangle}} - 4 \langle X, X \rangle \langle X, X \rangle \langle [X, Z]_{\mathfrak{m}}, X \rangle \right] \\
&= \frac{\langle X, X \rangle^3 \left( \langle X, X \rangle^2 + \langle X, X \rangle - 2 \langle X, X \rangle \sqrt{\langle X, X \rangle} \right) \langle X, [X, Z]_{\mathfrak{m}} \rangle}{\left( \langle X, X \rangle - \sqrt{\langle X, X \rangle} \right)^4} \\
&= \frac{\langle X, X \rangle^3 \langle X, [X, Z]_{\mathfrak{m}} \rangle}{\left( \langle X, X \rangle - \sqrt{\langle X, X \rangle} \right)^2}.
\end{aligned}$$

Therefore  $g_X(X, [X, Z]_{\mathfrak{m}}) = 0$  if and only if  $\langle X, [X, Z]_{\mathfrak{m}} \rangle = 0$ .  $\square$

Now, we give necessary and sufficient condition for a nonzero vector in a homogeneous Finsler space with exponential metric to be a geodesic vector.

**Theorem 3.3.** *Let  $G/H$  be a homogeneous Finsler space with exponential metric  $F = \alpha e^{\beta/\alpha}$  given by an invariant Riemannian metric  $\tilde{a}$  and an invariant vector field  $\tilde{X}$  such that  $\tilde{X}(H) = X$ . Then, a non-zero vector  $Y \in \mathfrak{g}$  is a geodesic vector if and only if*

$$\left\langle X + \left( \frac{\sqrt{\langle Y_{\mathfrak{m}}, Y_{\mathfrak{m}} \rangle} - \langle X, Y_{\mathfrak{m}} \rangle}{\langle Y_{\mathfrak{m}}, Y_{\mathfrak{m}} \rangle} \right) Y_{\mathfrak{m}}, [Y, Z]_{\mathfrak{m}} \right\rangle = 0. \quad (4)$$

*Proof.* Using lemma 3.4 of ([28]), we can write

$$F(Y) = \langle X, Y \rangle e^{\langle X, Y \rangle / \sqrt{\langle Y, Y \rangle}}.$$

Also, we know that

$$g_Y(U, V) = \frac{1}{2} \frac{\partial^2}{\partial s \partial t} F^2(Y + sU + tV) \Big|_{s=t=0}.$$

After some calculations, we get

$$\begin{aligned} g_Y(U, V) &= e^{2\langle X, Y \rangle / \sqrt{\langle Y, Y \rangle}} \left[ \langle U, V \rangle + 2 \langle X, U \rangle \langle X, V \rangle - \frac{\langle X, Y \rangle \langle Y, U \rangle \langle Y, V \rangle}{\langle Y, Y \rangle^{3/2}} \right. \\ &\quad + \frac{1}{\sqrt{\langle Y, Y \rangle}} \left\{ \langle X, U \rangle \langle Y, V \rangle + \langle X, V \rangle \langle Y, U \rangle - \langle X, Y \rangle \langle U, V \rangle \right\} \\ &\quad \left. + \frac{2 \langle X, Y \rangle}{\langle Y, Y \rangle} \left\{ \frac{\langle X, Y \rangle \langle Y, U \rangle \langle Y, V \rangle}{\langle Y, Y \rangle} - \langle Y, U \rangle \langle X, V \rangle - \langle X, U \rangle \langle Y, V \rangle \right\} \right]. \end{aligned} \quad (5)$$

From above equation, we can write

$$\begin{aligned} g_{Y_m}(Y_m, [Y, Z]_m) &= e^{2\langle X, Y_m \rangle / \sqrt{\langle Y_m, Y_m \rangle}} \times \\ &\left[ \langle Y_m, [Y, Z]_m \rangle + 2 \langle X, Y_m \rangle \langle X, [Y, Z]_m \rangle - \frac{\langle X, Y_m \rangle \langle Y_m, Y_m \rangle \langle Y_m, [Y, Z]_m \rangle}{\langle Y_m, Y_m \rangle^{3/2}} \right. \\ &\quad + \frac{1}{\sqrt{\langle Y_m, Y_m \rangle}} \left\{ \langle X, Y_m \rangle \langle Y_m, [Y, Z]_m \rangle + \langle X, [Y, Z]_m \rangle \langle Y_m, Y_m \rangle - \langle X, Y_m \rangle \langle Y_m, [Y, Z]_m \rangle \right\} \\ &\quad \left. + \frac{2 \langle X, Y_m \rangle}{\langle Y_m, Y_m \rangle} \left\{ \frac{\langle X, Y_m \rangle \langle Y_m, Y_m \rangle \langle Y_m, [Y, Z]_m \rangle}{\langle Y_m, Y_m \rangle} - \langle Y_m, Y_m \rangle \langle X, [Y, Z]_m \rangle - \langle X, Y_m \rangle \langle Y_m, [Y, Z]_m \rangle \right\} \right] \\ &= e^{2\langle X, Y_m \rangle / \sqrt{\langle Y_m, Y_m \rangle}} \left[ \langle Y_m, [Y, Z]_m \rangle \left\{ 1 - \frac{\langle X, Y_m \rangle}{\sqrt{\langle Y_m, Y_m \rangle}} \right\} + \langle X, [Y, Z]_m \rangle \sqrt{\langle Y_m, Y_m \rangle} \right] \\ &= \sqrt{\langle Y_m, Y_m \rangle} e^{2\langle X, Y_m \rangle / \sqrt{\langle Y_m, Y_m \rangle}} \left\langle X + \left( \frac{\sqrt{\langle Y_m, Y_m \rangle} - \langle X, Y_m \rangle}{\langle Y_m, Y_m \rangle} \right) Y_m, [Y, Z]_m \right\rangle. \end{aligned} \quad (6)$$

Now, from lemma (3.1),  $Y \in \mathfrak{g}$  is a geodesic vector if and only if

$$g_{Y_m}(Y_m, [Y, Z]_m) = 0, \quad \forall Z \in \mathfrak{m}.$$

Therefore  $g_{Y_m}(Y_m, [Y, Z]_m) = 0$  if and only if equation (4) holds.  $\square$

**COROLLARY 3.2.** *Let  $(G/H, F)$  be a homogeneous Finsler space with exponential metric  $F = \alpha e^{\beta/\alpha}$  defined by an invariant Riemannian metric  $\langle \cdot, \cdot \rangle$*

and an invariant vector field  $\tilde{X}$  such that  $\tilde{X}(H) = X$ . Let  $Y \in \mathfrak{g}$  be a vector such that  $\langle X, [Y, Z]_{\mathfrak{m}} \rangle = 0 \forall Z \in \mathfrak{m}$ . Then  $Y$  is a geodesic vector of  $(G/H, \langle \cdot, \cdot \rangle)$  if and only if  $Y$  is a geodesic vector of  $(G/H, F)$ .

*Proof.* From equation (6), we can write

$$\begin{aligned} g_{Y_{\mathfrak{m}}}(Y_{\mathfrak{m}}, [Y, Z]_{\mathfrak{m}}) &= e^{2\langle X, Y_{\mathfrak{m}} \rangle / \sqrt{\langle Y_{\mathfrak{m}}, Y_{\mathfrak{m}} \rangle}} \left[ \langle Y_{\mathfrak{m}}, [Y, Z]_{\mathfrak{m}} \rangle \left\{ 1 - \frac{\langle X, Y_{\mathfrak{m}} \rangle}{\sqrt{\langle Y_{\mathfrak{m}}, Y_{\mathfrak{m}} \rangle}} \right\} + \langle X, [Y, Z]_{\mathfrak{m}} \rangle \sqrt{\langle Y_{\mathfrak{m}}, Y_{\mathfrak{m}} \rangle} \right] \\ &= e^{2\langle X, Y_{\mathfrak{m}} \rangle / \sqrt{\langle Y_{\mathfrak{m}}, Y_{\mathfrak{m}} \rangle}} \left\{ 1 - \frac{\langle X, Y_{\mathfrak{m}} \rangle}{\sqrt{\langle Y_{\mathfrak{m}}, Y_{\mathfrak{m}} \rangle}} \right\} \langle Y_{\mathfrak{m}}, [Y, Z]_{\mathfrak{m}} \rangle, \quad \text{because } \langle X, [Y, Z]_{\mathfrak{m}} \rangle = 0. \end{aligned}$$

Therefore  $g_{Y_{\mathfrak{m}}}(Y_{\mathfrak{m}}, [Y, Z]_{\mathfrak{m}}) = 0$  if and only if  $\langle Y_{\mathfrak{m}}, [Y, Z]_{\mathfrak{m}} \rangle = 0$ .  $\square$

**Theorem 3.4.** Let  $(G/H, F)$  be a homogeneous Finsler space with exponential metric  $F = \alpha e^{\beta/\alpha}$  defined by an invariant Riemannian metric  $\langle \cdot, \cdot \rangle$  and an invariant vector field  $\tilde{X}$  such that  $\tilde{X}(H) = X$ . Then  $X$  is a geodesic vector of  $(G/H, \langle \cdot, \cdot \rangle)$  if and only if  $X$  is a geodesic vector of  $(G/H, F)$ .

*Proof.* From equation (5), we can write

$$\begin{aligned} g_X(X, [X, Z]_{\mathfrak{m}}) &= e^{2\langle X, X \rangle / \sqrt{\langle X, X \rangle}} \times \\ &\quad \left[ \langle X, [X, Z]_{\mathfrak{m}} \rangle + 2 \langle X, X \rangle \langle X, [X, Z]_{\mathfrak{m}} \rangle - \frac{\langle X, X \rangle \langle X, X \rangle \langle X, [X, Z]_{\mathfrak{m}} \rangle}{\langle X, X \rangle^{3/2}} \right. \\ &\quad \left. + \frac{1}{\sqrt{\langle X, X \rangle}} \left\{ \langle X, X \rangle \langle X, [X, Z]_{\mathfrak{m}} \rangle + \langle X, [X, Z]_{\mathfrak{m}} \rangle \langle X, X \rangle - \langle X, X \rangle \langle X, [X, Z]_{\mathfrak{m}} \rangle \right\} \right. \\ &\quad \left. + \frac{2 \langle X, X \rangle}{\langle X, X \rangle} \left\{ \frac{\langle X, X \rangle \langle X, X \rangle \langle X, [X, Z]_{\mathfrak{m}} \rangle}{\langle X, X \rangle} - \langle X, X \rangle \langle X, [X, Z]_{\mathfrak{m}} \rangle - \langle X, X \rangle \langle X, [X, Z]_{\mathfrak{m}} \rangle \right\} \right] \\ &= e^{2\langle X, X \rangle / \sqrt{\langle X, X \rangle}} \langle X, [X, Z]_{\mathfrak{m}} \rangle. \end{aligned}$$

Therefore  $g_X(X, [X, Z]_{\mathfrak{m}}) = 0$  if and only if  $\langle X, [X, Z]_{\mathfrak{m}} \rangle = 0$ .  $\square$

## References

- [1] S. I. Amari and H. Nagaoka, *Methods of information geometry*, Translations of Mathematical Monographs, AMS, 191, Oxford Univ. Press, 2000.
- [2] P. L. Antonelli, R. S. Ingarden, and M. Matsumoto, *The Theory of Sprays and Finsler spaces with Applications in Physics and Biology*, Kluwer Academic Publishers, Netherlands, 58, 1993.

- [3] V. I. Arnold, Sur la géométrie différentielle des groupes de Lie de dimension infinie et ses applications à l'hydrodynamique des fluides parfaits, *Ann. Inst. Fourier (Grenoble)*, **16** (1960), 319-361.
- [4] D. Bao, S. S. Chern and Z. Shen, *An Introduction to Riemann-Finsler Geometry*, GTM-200, Springer-Verlag 2000.
- [5] E. Cartan, *Les espaces de Finsler*, Actualites Scientifiques et Industrielles no. 79, Paris, Hermann, 1934.
- [6] S. S. Chern, Finsler geometry is just Riemannian geometry without the quadratic restriction, *Notices Amer. Math. Soc.*, **43** (9) (1996), 959-963.
- [7] S. S. Chern and Z. Shen, *Riemann-Finsler Geometry*, Nankai Tracts in Mathematics, Vol. 6, World Scientific Publishers, 2005.
- [8] S. Deng, *Homogeneous Finsler Spaces*, Springer Monographs in Mathematics, New York, 2012.
- [9] S. Deng and Z. Hou, The group of isometries of a Finsler space, *Pacific J. Math.*, **207** (2002), 149-155.
- [10] Z. Dušek, The affine approach to homogeneous geodesics in homogeneous Finsler spaces, *preprint*, arXiv:1703.01199.
- [11] P. Finsler, Über Kurven und Flächen in allgemeinen Räumen, (Dissertation, Göttingen, 1918), *Birkhäuser Verlag*, Basel, 1951.
- [12] C. Gordon, Homogeneous Riemannian manifolds whose geodesics are Orbits, *Prog. Nonlinear Differ. Eq. App.*, **20** (1996), 155-174.
- [13] P. Habibi, D. Latifi and M. Toomanian, Homogeneous geodesics and the critical points of the restricted Finsler function, *J. Cont. Math.*, **4** (6) (2011), 12–16.
- [14] M. Hosseini and H. R. Salimi Moghaddam, On the existence of homogeneous geodesics in homogeneous  $(\alpha, \beta)$ -spaces, ?
- [15] V.V. Kajzer, Conjugate points of left invariant metrics on Lie group, *Sov. Math.*, **34** (1990), 32–44 [translation from. Izv. Vyssh. Uchebn. Zaved. Mat. **342** (1990), 27–37].
- [16] A. Kaplan, On the geometry of groups of Heisenberg type, *Bull. Lond. Math. Soc.*, **15** (1983), 35–42.

- [17] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry*, Interscience Publishers, Vol. 1, 1963; Vol. 2, 1969.
- [18] B. Kostant, Holonomy and Lie algebra of motions in Riemannian manifolds, *Trans. Am. Math. Soc.*, **80** (1995), 520–542.
- [19] O. Kowalski and L. Vanhecke, Riemannian manifolds with homogeneous geodesics, *Boll. Un. Math. Ital.* **5** (1991), 189-246.
- [20] O. Kowalski and J. Szenthe, On the existence of homogeneous geodesics in homogeneous Riemannian manifolds, *Geom. Dedicata*, **81** (2000), 209–214 [Erratum: *Geom. Dedicata* **84** (2001), 331–332].
- [21] O. Kowalski and Z. Vlášek, Homogeneous Riemannian manifolds with only one homogeneous geodesic, *Publ. Math. Debr.*, **62** (3–4) (2003), 437–446.
- [22] E. A. Lacomba, Mechanical Systems with Symmetry on Homogeneous Spaces, *Trans. Amer. Math. Soc.*, **185** (1973), 477-491.
- [23] D. Latifi, Homogeneous geodesics in homogeneous Finsler spaces, *J. Geom. Phys.*, **57** (2007), 1421-1433.
- [24] D. Latifi and A. Razavi, Homogeneous geodesics of left invariant Randers metrics on a three-dimensional Lie group, *Int. J. Cont. Math. Sci.*, **4** (2009), 873–881.
- [25] M. Matsumoto, On a C-reducible Finsler space, *Tensor N. S.*, **24** (1972), 29-37.
- [26] G. Shanker and S. A. Baby, The  $L$ -dual of a Finsler space with infinite series  $(\alpha, \beta)$ -metric, *Bull. Cal. Math. Soc.*, **107** (4) (2015), 337-356.
- [27] G. Shanker and S. A. Baby, On the projective flatness of a Finsler space with infinite series  $(\alpha, \beta)$ -metric, *South East Asian J. of Math & Math. Sci.*, **11** (1) (2015), 17-24.
- [28] G. Shanker and K. Kaur, Homogeneous Finsler spaces with some special  $(\alpha, \beta)$ -metrics, *Preprint*, arXiv:1712.06328v2[math.DG].
- [29] G. Shanker and Kirandeep Kaur, On the flag curvature of homogeneous Finsler space with some special  $(\alpha, \beta)$ -metrics, *Preprint*, arXiv:1801.05324 [math.DG].

- [30] G. Shanker and Ravindra, On Randers change of exponential metric, *Applied Sciences*, **15** (2013), 94–103.
- [31] J. Szenthe, Homogeneous geodesics of left invariant metrics, *Univ. Iagel. Acta Math.*, **38** (2000), 99–103.
- [32] J. Szenthe, Stationary geodesics of left invariant Lagrangians, *J. Phys. A Math. Gen.*, **34** (2001), 165–175.
- [33] G. Z. Tóth, On Lagrangian and Hamiltonian systems with homogeneous trajectories, *J. Phys. A: Math. Theor.*, **43** (2010), 385206 (19pp).
- [34] E.B. Vinberg, Invariant linear connections in a homogeneous manifold, *Trudy MMO*, **9** (1960), 191–210.
- [35] Z. Yan and S. Deng, Finsler spaces whose geodesics are orbits, *Differ. Geom. Appl.*, **36** (2014), 1–23.
- [36] Z. Yan and S. Deng, Existence of homogeneous geodesics on homogeneous Randers spaces, *Houston J. Math.*, (2016) (in preparation) ?
- [37] Z. Yan, Existence of homogeneous geodesics on homogeneous Finsler spaces of odd dimension, *Monatsh Math.*, **182** (2017), 165–171.
- [38] Z. Yan and Libing Huang, On the existence of homogeneous geodesic in homogeneous Finsler spaces, *Journal of Geometry and Physics*, **124** (2018), 264–267.