Path integral approach to one-dimensional discrete-time quantum walk

Karthik S. Joshi,¹ S. K. Srivatsa,^{2,1} and R. Srikanth^{1,*}

¹Poornaprajna Institute of Scientific Research, Bengaluru, India.

²Dept. of Electrical and Electronics Engineering, PES University, Bengaluru, India.

Discrete-time quantum walk in one-dimension is studied from a path-integral perspective. This enables derivation of a closed-form expression for amplitudes corresponding to any coin-position basis of the state vector of the quantum walker at an arbitrary step of the walk. This provides a new approach to the foundations and applications of quantum walks.

I. INTRODUCTION

An approach to quantum mechanics via the generalization of the action principle in classical mechanics was established by Feynman [1], known as the *path integral* formulation. It assumes that a particle can take all possible paths to travel between two fixed points, say O, the origin, and T, the terminal point respectively. A probability amplitude is assigned to each path the particle can take and the propagator is obtained by taking the sum of all amplitudes corresponding to all paths for the particle to propagate from O to T.

Let C be a path from O to T, with an assigned probability amplitude P(C), which in general is a complex number. The probability that the particle will take the path C is $|P(C)|^2$. The joint probability for the particle to take paths C_1 and C_2 is then $|P(C_1) + P(C_2)|^2$. Notice that the rule for obtaining joint probability deviates from its classical counterpart by an interference term of the form $P(C_1)P^*(C_2) + P^*(C_1)P(C_2)$. Further, in a path C made of two sub - paths say C_1 and C_2 meeting at an intermediate point M, the probability amplitude $P(C) = P(C_1)P(C_2)$.

The path integral approach has been applied to study various physical processes, including the quantum dynamics involving scalar potentials with singularities which were not amenable to the standard operator approach of Schrdinger [2]. It has also been extensively applied in the study of quantum field theory particularly for quantization, gauge fixing and phenomenon concerning elementary particles [3]. The problem of decoherence, central for realization of a quantum computer has been discussed via the path integral approach in [4].

A candidate system to study via Feynman approach is the quantum walk (QW). The QW is a process wherein the evolution of the walker is driven by a fixed unitary operator. QW process begins with the walker at the origin, say, moving backward or forward conditioned on the outcome of a (quantum) coin toss. This process is repeated n - times for a n - step QW. Broad classification of QW include discrete time QW (DTQW) and continuous time QW. Many variants of QW have been studied; alternating [5], coinless [6] and staggered model [7]. QW has also been studied in the presence of noise, for instance [8]. A detailed review of quantum walk is presented in [9].

Realization of QW have been carried out in a variety of physical systems and have been discussed in detail in [10]. Schemes to implement QW have been proposed; ion [11] and optical traps [12], quantum circuits [13] and cavity QED [14] to name a few.

It has been shown that QW serves as an effective tool to design quantum algorithms. Algorithmic applications of quantum walk include *element search* [15] and *triangle finding* (finding a triangle on an undirected graph and also indicate if the graph is triangle free) [16] among others. Also, it has been shown that any quantum computation process can be implemented by performing quantum walk in both continuous and discrete cases [17] [18]. This powerful result only adds more thrust to the study of quantum walk.

Further, close ties of the QW framework and quantum field theory has recently come to fore. Among others, QW as a model to perform quantum simulation of relativistic particles has been studied [19]. With the demonstration of implementation of QW using a Dirac like Hamiltonian [20], modelling the phenomena of neutrino oscillations [21], etc the study of QW has only gained further impetus.

While the rich and varied applications of QW indicates the success of the framework, analytical results are but few. This can be attributed to the many interferences occurring simultaneously, which seemingly makes it analytically intractable. However, the maxima of the probability distribution arising from the QW process was obtained analytically in [22]. In this work, we study QW in one dimension via the Feynman path integral approach, obtaining in the process a closed form expression for the probability amplitude for an arbitrary time step in the case of a one dimensional DTQW.

II. FEYNMAN PATH APPROACH TO QW

We briefly present the Discrete Quantum walk scheme for completeness. Consider the walker to be initially in the state

$$|\psi(t=0)\rangle = (\alpha |0\rangle + e^{i\phi}\beta |1\rangle) \otimes |k\rangle.$$
(1)

We assume α and β to be real numbers without loss of generality. The Hilbert space of the walker is $\mathcal{H} = \mathcal{H}_{coin} \otimes \mathcal{H}_{position}$. One-step evolution of the walker is

^{*}Electronic address: srik@poornnaprajna.org

achieved by applying a unitary operator $U \equiv C \otimes S$ on $|\psi(0)\rangle$, with coin operator C, given by:

$$C = \begin{pmatrix} \cos\theta & \sin\theta\\ \sin\theta & -\cos\theta \end{pmatrix}, \tag{2}$$

and the shift operator S given by:

$$S = \Sigma_k \left| 0 \right\rangle \left\langle 0 \right| \otimes \left| k - 1 \right\rangle \left\langle k \right| + \Sigma_k \left| 1 \right\rangle \left\langle 1 \right| \otimes \left| k + 1 \right\rangle \left\langle k \right|.$$
(3)

Application of U^n carries out a n - step QW. Evolving the initial state Eq.(1) by one step we obtain

$$\begin{aligned} |\psi(1)\rangle &= \alpha \cos\theta \,|0,k-1\rangle + \alpha \sin\theta \,|1,k+1\rangle \\ &+ e^{i\phi}\beta \sin\theta \,|0,k-1\rangle - e^{i\phi}\beta \cos\theta \,|1,k+1\rangle \,. \end{aligned}$$
(4)

The basis vectors of $|\psi(1)\rangle$ represent the final state of the walker. Let the transitions $|k\rangle \rightarrow |k+1\rangle$ and $|k\rangle \rightarrow |k-1\rangle$ be labelled as *forward* (F) and *backward* (B) transition respectively. All possible one-step transitions are deduced from Eq.(4) and are illustrated in Fig. (1).



FIG. 1: Figure shows all possible one step transitions and the corresponding probability amplitudes of the walker starting with initial states $\alpha |0, k\rangle$ and $\beta e^{i\phi} |1, k\rangle$

Note that the probability amplitudes are independent of the instantaneous position of the walker, $|k\rangle$ and the step number, t.

Two steps of the QW can be obtained by combining one-step transitions. For instance, a possible two step evolution can be obtained by combining B and F onestep transitions- a BF transition, which indicates that the walker underwent a $|k\rangle \rightarrow |k-1\rangle$ transition at time t = 1 followed by a $|k-1\rangle \rightarrow |k\rangle$ transition at time t = 2. Similarly, an n - step walk can be represented by a string of Fs and Bs of length n, which prompts the following definition:

Definition 1 (Feynman string). Any n-bit string $S \in \{B, F\}^{\times n}$ represents the n-1 (one step) transitions in a Feynman path of an n - step QW.

Strings starting with a B are called B - strings, likewise, strings starting with F are called F - strings. Given

a string S, its dual \overline{S} , is obtained by interchanging Fs and Bs. Evidently, the dual of a B-string is an F-string and vice-versa.

For any path starting at the origin and terminating at position $|x\rangle$ (say), the corresponding Feynman string S has $\frac{n+|x|}{2}$ Fs and $\frac{n-|x|}{2}$ Bs. Hence, the number of strings N representing paths which start at the origin and terminate at $|x\rangle$ is given by the number of distinct permutations of S. Therefore:

$$N = \frac{n!}{\left(\frac{n\pm|x|}{2}\right)! \left(\frac{n\mp|x|}{2}\right)!} = \binom{n}{\frac{n\pm|x|}{2}}.$$
 (5)

The walker has a certain initial coin state as shown in Eq.(1). This obviously affects the evolution of the walker, as coin states $|0\rangle$ and $|1\rangle$ effect B and F translations respectively. This gives rise to the mapping $B \rightarrow |0\rangle$ and $F \rightarrow |1\rangle$. Each of the above N strings are thus concatenated with a B or an F, effectively giving rise to strings B * S and F * S for all S, leading to the following definition:

Definition 2 (Feynman extended string). An (n + 1)bit string, denoted S^+ , consisting of a Feynman string preceded by the initial coin state of the walker $B(|0\rangle)$ or $F(|1\rangle)$.

Note that each of the extended strings S^+ of an n-step walk has n one step transitions. Furthermore, B * S and F * S for all N number of strings S, will result in 2N strings.

In the Feynman framework, each of the 2N paths are associated with a probability amplitude. Let the m^{th} path string have a probability amplitude $P(S_m)$. The net probability amplitude, P(x), for the walker to be found at position $|x\rangle$ is the sum of all these amplitudes:

$$P(x) = \sum_{m=1}^{2N} P(S_m).$$
 (6)

On the other hand, for a given path represented by an n + 1 - bit string S, the corresponding probability amplitude P(S) is given by the product of all the n one step probability amplitudes P(k):

$$P(S) = \prod_{k=1}^{n} P(k).$$
 (7)

The discussion presented in the remainder of the article pertains to both Feynman string and its extended counterpart. They will be collectively referred to as Feynman strings unless otherwise specified.

Note that each one step transition introduces a $\sin \theta$ or a $\pm \cos \theta$ factor to the probability amplitude, therefore it is proportional to $\cos^i \theta \sin^j \theta$, with i + j = n.

Definition 3 (Switch). In any n-bit Feynman string, $S \in \{B, F\}^{\times n}$, switch is defined as a $F \leftrightarrow B$ transition in the string. It is easily seen from Fig. (1) that number of switches fixes the degree of $\sin \theta$ in the probability amplitude. Evidently, there are multiple strings, say $\eta(n, x, j)$, with same number of switches, which needs to be accounted.

The full path string of length n and terminating at position |x| has $N_F(n,x) = \frac{n+|x|}{2}$ dominant (say, forward) shifts and $N_B(n,x) = \frac{n-|x|}{2}$ minority (say, backward) shifts. Note that for a fixed time step t, n and |x| are both even or both odd.

The number of strings, for a fixed j switches, is equivalent to the problem of filling (i) $N_F(n, x)$ balls into $\mu \equiv \lfloor (j+1)/2 \rfloor$ urns and of $N_B(n, x)$ balls into $\nu \equiv \lceil (j+1)/2 \rceil$ urns; plus (ii) $N_F(n, x)$ balls into ν urns and of $N_B(n, x)$ balls into μ urns;

At the value of j where $N_B(n, x) < \nu$ (i.e., when j = n - |x|), contribution (i) drops out. At precisely the next value of j, $N_B(n, x) < \mu$ (i.e., j = n - |x| + 1), both (i) and (ii) have no more contributions. Thus, both (i) and (ii) contribute up to j = n - |x| - 1, and then at j = n - |x| there is a contribution from (ii) alone. For larger values of j, there aren't any more contributions.

Recall that the problem of filling n balls into k urns, such that no urn is empty, is the combinatoric problem of strong composition, and the number of ways are given by $\binom{n-1}{k-1}$.

The total number of n-Feynman paths with j switches is given by:

$$\eta(n, x, j) \equiv {\binom{n+|x|}{2} - 1} {\binom{j+1}{2} - 1} {\binom{\frac{j+1}{2}}{2} - 1} {\binom{\frac{j+1}{2}}{2} - 1} + {\binom{n-|x|}{2} - 1} {\binom{\frac{j+1}{2}}{2} - 1} {\binom{\frac{j+1}{2}}{2} - 1} {\binom{\frac{j+1}{2}}{2} - 1}, \quad (8)$$

assuming $j \le n - |x| - 1$. If j = n - |x|, then only the first summand in Eq. (8) contributes. The following identity can easily be verified:

$$\binom{n}{\frac{n\pm|x|}{2}} = \sum_{j=1}^{n-|x|-1} \eta(n,x,j) + \binom{\frac{n+|x|}{2}-1}{\lceil\frac{n-|x|+1}{2}\rceil-1}.$$
 (9)

Recall that the LHS of Eq. (9) is same as the total number of Feynman paths obtained in Eq. (5).

Note that in the case of equal forward and backward paths, where we set x = 0, the second summand in the RHS of Eq. (9) vanishes.

Further, when a B - string is concatenated with a F (or vice - versa), an additional switch is introduced. For example, a string with two switches BBFFB, when concatenated with a F to obtain the extended string F * BBFFB, has three switches. The additional switch so introduced in the extended needs to be accounted for. Let $\eta^*(n+1, x, j)$ denote the number of extended strings with j switches, then:

$$\eta^*(n+1, x, j) = \eta(n, x, j) + \eta(n, x, j-1),$$
(10)

for 1 < j < n - |x|. For j = 1 and j = n - |x| + 1, we have:

$$\eta^*(n+1, x, j) = \eta(n, x, j).$$
(11)

Definition 4 (Parity). Given an n bit Feynman (extended) string, the parity ϵ takes the value +1 (resp., -1) when the number of $F \rightarrow F$ transitions in the string are even (resp. odd).

The parity of a Feynman extended string is obtained in the following lemma.

Lemma 1. The parity $\epsilon(n, x, j)$ of Feynman B or Fstrings with j switches is given by

$$\epsilon(n,x,j) = (-1)^{\frac{n+x}{2} - \lfloor \frac{j+1}{2} \rfloor}$$
(12)

and

$$\epsilon(n, x, j) = (-1)^{\frac{n+x}{2} - \lceil \frac{j+1}{2} \rceil + 1}$$
(13)

respectively.

Proof. Let S be a B - string with j switches. Cut the string at points where switches occur. This process creates j + 1 sub-strings with $\lceil \frac{j+1}{2} \rceil$ B-type sub-strings and $\lfloor \frac{j+1}{2} \rfloor$, F-type sub-strings. Here, the B-type sub-strings do not alter the parity of the string S and hence are ignored. Distribute an F bit to each of the F-type sub-strings as all sub-strings must be at least of size one. This leaves us with $\frac{n+x}{2} - \lfloor \frac{j+1}{2} \rfloor$, Fs which can then be redistributed among the sub-strings arbitrarily. Independent of the distribution of Fs among the F-type sub-strings, the number of $F \to F$ transitions are unaltered when the sub-strings are conjoined. Hence, if $\frac{n+x}{2} - \lfloor \frac{j+1}{2} \rfloor$ is even, $\epsilon = -1$, as there are odd number of $F \to F$ transitions, similarly, $\epsilon = 1$ when $\frac{n+x}{2} - \lfloor \frac{j+1}{2} \rfloor$ is odd.

The result can be proved analogously for F- strings.

The above lemma also implies that the parity of a Feynman extended string is invariant under permutations which preserve the number of switches.

Lemma 2. A Feynman B - string (resp. F - string) with j switches determines the coin state in an eigen basis $|b,x\rangle$: $b = j \pmod{2}$ (resp. $b = j + 1 \pmod{2}$).

Proof. Let S be a B - string with j switches, where j is even. Cut the string at points where switches occur as described in lemma (1). As S is a B - string, the first of the j + 1 sub-strings contains only Bs. This sub - string is followed by F sub - string which in turn is followed by a B sub-string and so on. It follows from this arrangement that the last sub-string contains Bs. Every transition terminating with a B results in the coin state $|0\rangle$ as evident from Fig. (1). For an odd number of switches, j, in a B - string, the above process of cutting and arranging the j+1 sub - strings, where j+1 is even, results in the $(j + 1)^{th}$ sub-string to contain only Fs.

Every transition terminating with a F results in the coin state $|1\rangle$ as evident from table Fig. (1)).

The result can be similarly shown for F - strings.

Hence, using the above results, the state vector component corresponding to *a given* Feynman extended string can expressly be written as:

$$|\xi(n, x, j, c)\rangle = \epsilon(n+1, x, j) \cos^{i} \theta \sin^{j} \theta \times |(j+c \pmod{2}), x\rangle, \qquad (14)$$

where, c = 0 for *B*- strings and c = 1 for *F*- strings.

We note that a B- string with j switches and its dual

have *like* parity when j is even and *unlike* parity when j is odd according to Eqs. (12) and (13).

Defining the summing operations:

$$\begin{split} \sum_{1} &\equiv \sum_{x=-n}^{n} \frac{1+(-1)^{n+x}}{2} \sum_{m=1}^{\frac{n+|x|}{2}} \frac{\epsilon(n,x,2m)}{\eta^{*}(n+1,x,2m)} \\ \sum_{2} &\equiv \sum_{x=-n}^{n} \frac{1+(-1)^{n+x}}{2} \sum_{m=1}^{\frac{n+|x|+1}{2}} \frac{\epsilon(n,x,2m-1)}{\eta^{*}(n+1,x,2m-1)}, \end{split}$$
(15)

we obtain the final state vector to be:

$$|\psi(n)\rangle = \left(\sum_{1} \alpha N_B(n, x)(\cos\theta)^{n-2m} (\sin\theta)^{2m} + \sum_{2} \beta e^{i\phi} N_F(n, x)(\cos\theta)^{n-2m+1} (\sin\theta)^{2m-1}\right) |0, x\rangle + \left(\sum_{1} \beta e^{i\phi} N_F(n, x)(\cos\theta)^{n-2m} (\sin\theta)^{2m} - \sum_{2} \alpha N_B(n, x)(\cos\theta)^{n-2m+1} (\sin\theta)^{2m-1}\right) |1, x\rangle,$$
 (16)

which may be considered as an explicit representation of a QW parametrized by position. This allows us to circumvent a direct calculation of amplitudes at each position. Its application will be explored elsewhere.

III. CONCLUSION

In this work, a closed form expression was obtained for the state vector corresponding to arbitrary time step for an n - step QW in one dimension. The Feynman path integral approach was used to derive all possible (backward and forward) transition rules of the walker. The forward and backward transitions were mapped to a string of Fs and Bs respectively. A recipe for calculating the probability amplitude for a given string was provided using the properties of these strings.

The formalism presented in this work can be extended to QW in higher dimensions including walks by augmenting the QW "Feynman rules" presented in Fig. (1), which would allow us to accommodate QW in higher dimensions as well QW in arbitrary graphs. One can in principle extend this formalism to any Markovian process evolving under a fixed unitary operator.

- R. P. Feynman. Space-time approach to non-relativistic quantum mechanics. *Rev. Mod. Phys.*, 20:367–387, Apr 1948.
- [2] E. Nelson. Feynman Integrals and the Schrödinger Equation. Journal of Mathematical Physics, 5:332–343, March 1964.
- [3] Das Ashok. Field Theory: A Path Integral Approach, volume 75. World Scientific, 2006.
- [4] Sergio Albeverio, Laura Cattaneo, Sonia Mazzucchi, and Luca Di Persio. A rigorous approach to the feynmanvernon influence functional and its applications. i. Journal of Mathematical Physics, 48(10):102109, 2007.
- [5] C. Di Franco, M. Mc Gettrick, and Th. Busch. Mimicking the probability distribution of a two-dimensional grover walk with a single-qubit coin. *Phys. Rev. Lett.*, 106:080502, Feb 2011.
- [6] Apoorva Patel, KS Raghunathan, and Pranaw Rungta.

Quantum random walks do not need a coin toss. *Physical Review A*, 71(3):032347, 2005.

- [7] Renato Portugal, Raqueline AM Santos, Tharso D Fernandes, and Demerson N Gonçalves. The staggered quantum walk model. *Quantum Information Processing*, 15(1):85–101, 2016.
- [8] R Srikanth, Subhashish Banerjee, and CM Chandrashekar. Quantumness in a decoherent quantum walk using measurement-induced disturbance. *Physical Re*view A, 81(6):062123, 2010.
- [9] SalvadorElas Venegas-Andraca. Quantum walks: a comprehensive review. *Quantum Information Processing*, 11(5):1015–1106, 2012.
- [10] Kia Manouchehri and Jingbo Wang. Physical Implementation of Quantum Walks. Springer-Verlag, Berlin Heidelberg, 2014.
- [11] B. C. Travaglione and G. J. Milburn. Implementing the

quantum random walk. Phys. Rev. A, 65:032310, Feb 2002.

- [12] C. M. Chandrashekar. Implementing the one-dimensional quantum (hadamard) walk using a bose-einstein condensate. *Phys. Rev. A*, 74:032307, Sep 2006.
- [13] Jozef Košík and Vladimír Bužek. Scattering model for quantum random walks on a hypercube. *Phys. Rev. A*, 71:012306, Jan 2005.
- [14] G. S. Agarwal and P. K. Pathak. Quantum random walk of the field in an externally driven cavity. *Phys. Rev. A*, 72:033815, Sep 2005.
- [15] Neil Shenvi, Julia Kempe, and K. Birgitta Whaley. Quantum random-walk search algorithm. *Phys. Rev. A*, 67:052307.
- [16] Frdric Magniez, Miklos Santha, and Mario Szegedy. Quantum algorithms for the triangle problem. SIAM Journal on Computing, 37(2):413–424, 2007.
- [17] Andrew M Childs. Universal computation by quantum walk. *Physical review letters*, 102(18):180501, 2009.

- [18] Neil B Lovett, Sally Cooper, Matthew Everitt, Matthew Trevers, and Viv Kendon. Universal quantum computation using the discrete-time quantum walk. *Physical Review A*, 81(4):042330, 2010.
- [19] Pablo Arrighi, Vincent Nesme, and Marcelo Forets. The dirac equation as a quantum walk: higher dimensions, observational convergence. *Journal of Physics A: Mathematical and Theoretical*, 47(46):465302, 2014.
- [20] C. M. Chandrashekar. Two-component dirac-like hamiltonian for generating quantum walk on one-, two- and three-dimensional lattices. *Scientific Reports*, 3:2829, Oct 2013.
- [21] Arindam Mallick, Sanjoy Mandal, and C. M. Chandrashekar. Neutrino oscillations in discrete-time quantum walk framework. *The European Physical Journal C*, 77(2):85, Feb 2017.
- [22] Ashwin Nayak and Ashvin Vishwanath. Quantum walk on the line. arXiv preprint quant-ph/0010117, 2000.