

Nonperturbative spectrum of nonlocal gravity

Gianluca Calcagni,¹ Leonardo Modesto,² and Giuseppe Nardelli^{3,4}

¹*Instituto de Estructura de la Materia, CSIC, Serrano 121, 28006 Madrid, Spain*

²*Department of Physics, Southern University of Science and Technology, Shenzhen 518055, China*

³*Dipartimento di Matematica e Fisica, Università Cattolica del Sacro Cuore, via Musei 41, 25121 Brescia, Italy*

⁴*TIFPA – INFN c/o Dipartimento di Fisica, Università di Trento, 38123 Povo (Trento), Italy*

(Dated: March 21, 2018)

We investigate the nonperturbative degrees of freedom of a class of weakly nonlocal gravitational theories that have been proposed as an ultraviolet completion of general relativity. At the perturbative level, it is known that the degrees of freedom of nonlocal gravity are the same of the Einstein–Hilbert theory around any maximally symmetric spacetime. We prove that, at the nonperturbative level, the degrees of freedom are actually eight in four dimensions, contrary to what one might guess on the basis of the “infinite number of derivatives” present in the action. Six of these degrees of freedom may play a role at large scales and on curved backgrounds. We also propose a criterion to select the form factor almost uniquely.

Introduction. A quantum theory of gravity [1–3] should be able to solve, or say something constructive about, some problems left open in general relativity, such as the singularity problem (there exist spacetime points where the laws of physics break down, as in the big bang at the beginning of the Universe or inside black holes), the cosmological constant problem (two thirds of the content of our patch of the cosmos is made of a “dark energy” component not adequately described by general relativity or particle physics), and the mystery surrounding the birth and first stage of development of the Universe (the actual origin of the inflaton is unknown). In recent years, a new perturbative quantum field theory of gravity has rapidly emerged as a promising and accessible framework where the gravitational force consistently obeys the laws of quantum mechanics and all infinities seem to be tamed [4–14]. This proposal adapts ordinary techniques of perturbative field theory to an action with nonlocal operators. Fulfilling initial expectations based on naïve power-counting arguments, the theory turns out to be unitary and super-renormalizable or finite [10] at the quantum level thanks to the nonlocal nature of its dynamics. It may resolve the big bang [15–23] and black-hole singularities [24–29], and its cosmological solutions may unravel interesting bottom-up scenarios in the early universe (inflation) and at late times (dark energy).

Surprisingly, these encouraging features are accompanied by a number of appalling gaps of knowledge on basic questions on the classical theory, such as how to find solutions of the dynamics and whether they match the singularity-free geometries found when linearizing the equations of motion. The dynamics is usually solved with approximations or assumptions which do not give access to all admissible solutions [30]. Also, when considering nonlinear interactions (gravity is as nonlinear as it can be!) it becomes unclear how many initial conditions one must specify and how many degrees of freedom (d.o.f.) populate the spectrum of physical particles of the theory. The purpose of this Letter is to pave the way to fill these gaps and infer the total number of nonperturbative d.o.f. in minimal nonlocal gravity [7, 9] (super-renormalizable

in $D = 4$ dimensions; finite in $D = 5$).

Main results and consequences. We can summarize our findings in three statements. (A) For the types of nonlocality giving rise to a renormalizable theory, the number of field d.o.f. is finite and equal to $D(D - 2) = 8$ in four dimensions. Two of these d.o.f. correspond to the graviton, they propagate on flat spacetime at short distances, and are a familiar acquaintance in the nonlocal linearized dynamics. The other d.o.f. are a novelty because they emerge only when the fully nonlinear nonlocal dynamics is considered. Since they are not visible in a perturbative treatment on Minkowski spacetime, these d.o.f. are *nonperturbative*, typical of curved backgrounds, and, hence, may be important in the description of *large-scale, long-range physics*, such as at astrophysical or cosmological scales. (B) Also the number of initial conditions to specify in order to find classical solutions is finite. This contributes to, or even settles, a seventy-year-old debate about whether nonlocal theories are predictive at all, due to the peculiarities of their problem of initial conditions. The answer is Yes, for the specific nonlocalities considered here. Knowing how to construct dynamical solutions makes a crucial step in the understanding of the capabilities of the specific theory under examination, both at the classical and the quantum level. (C) The system can be recast as a set of finite-order differential equations, and hence the number of initial conditions is finite, only for two specific nonlocal form factors among those circulating in the literature. This constrains the ambiguity on the choice of form factors, i.e. of nonlocal theory.

The most important consequence of identifying new d.o.f. and of knowing how to formulate the initial-condition problem is that now one can, on one hand, construct nontrivial cosmological and black-hole classical solutions previously inaccessible with current methods and, on the other hand, build a robust, rigorous, and systematic set of phenomenological models that can be tested against experiments and observations, a vital task for any candidate theory of natural phenomena. Therefore, the results of the present work may be of interest for the applied mathematician, the quantum-gravity the-

TABLE I. Form factors in nonlocal gravity.

$H(\square)$	$P(\square)$	$f(\omega)$	Form factor name
$H^{\text{pol}}(\square) := \alpha\{\ln P(\square) + \Gamma[0, P(\square)] + \gamma_E\}$	$-l^2\square$	$e^{-\omega}$	Kuz'min [5]
	$O(\square^n)$		Tomboulis [6, 7]
$H^{\text{exp}}(\square) := \alpha P(\square)$	$-l^2\square$	$1 - \omega$	string-related [9, 15]
	$l^4\square^2$		Krasnikov [4]

oretician and phenomenologist, the particle-physics theoretician, the cosmologist, and the astrophysicist.

After a brief overview of the most prominent nonlocal gravities, we show that the number of initial conditions and d.o.f. is finite and we count them explicitly. The logic to follow is simple: (i) we write the nonlocal dynamics with infinitely many derivatives as a system of “master equations” which are second order in spacetime derivatives, hence the Cauchy problem is well defined; (ii) from this reformulation, we extract the number of initial conditions and the d.o.f.. The proof is self-contained and may be skipped by the reader interested only in the physical consequences of the theory, which are discussed above and in the final section.

Master equations. Consider the classical action

$$S = \frac{1}{2\kappa^2} \int d^D x \sqrt{-g} [R + G_{\mu\nu} \gamma(\square) R^{\mu\nu}], \quad (1)$$

where $G_{\mu\nu}$ is the Einstein tensor and $\gamma(\square)$ is a nonlocal form factor, an entire function of \square having special asymptotic properties [5–7, 10, 12, 13, 31]. It can be parametrized as

$$\gamma(\square) = \frac{e^{H(\square)} - 1}{\square}, \quad (2)$$

where $H(\square)$ depends on the dimensionless combination $l^2\square$ and l is a fixed length scale. The four principal theories are shown in Tab. I. In the first case (“pol”: asymptotically polynomial), $P(z)$ is a real positive polynomial of degree n ($2n$ derivatives) with $P(0) = 0$, Γ is Euler function, and γ_E is the Euler–Mascheroni constant. In the second case (“exp”) the form factor is asymptotically exponential. Quantum gravity with Kuz'min or Tomboulis form factor is renormalizable [5–7]; with the stringy profile $H^{\text{exp}}(\square) = -l^2\square$, it is renormalizable if perturbative expansions with the resummed propagator are allowed [11]; with Krasnikov profile $H^{\text{exp}}(\square) = l^4\square^2$, it is believed to be renormalizable, but the proof is not complete [4].

The physics stemming from the action (1) is a hard nut to crack. Even before quantizing the theory, two fundamental questions arise. How many degrees of freedom are there? Can one solve the dynamics once a finite number of initial conditions are given? While it is easy to see that linear systems have a finite number of d.o.f. and of initial conditions, the case with nonlinear interactions is highly nontrivial. Directly manipulating the infinitely many derivatives of Eq. (2) makes very difficult to answer the above questions.

To put the main result of this Letter in the most direct terms, we trade Eq. (2) for kernel functions that obey finite-order differential equations (*master equations*). Equation (1) becomes

$$S = \frac{1}{2\kappa^2} \int d^D x \sqrt{-g} R + \frac{1}{2\kappa^2} \int d^D x \sqrt{-g(x)} \times \int d^D y \sqrt{-g(y)} \int d^D z \sqrt{-g(z)} G_{\mu\nu}(x) \times [\mathcal{G}(x, y; \sigma) - 1] \tilde{G}(y, z) R^{\mu\nu}(z), \quad (3)$$

where \tilde{G} is the Green function solving [36]

$$\square \tilde{G}(y, z) = [-g(z)]^{-1/2} \delta^D(y - z). \quad (4)$$

Intuitively, \square^{-1} is expressed in terms of \tilde{G} , while $\exp H$ is replaced by \mathcal{G} , everything being integrated three times instead of one. To complete the system, we need to write down the master equations for \mathcal{G} , but before doing so we need to specify H . The profile $H(\square)$ can be defined through the integral

$$H_\sigma(\square) := \alpha \int_0^{\sigma P(\square)} d\omega \frac{1 - f(\omega)}{\omega} \xrightarrow{\sigma \rightarrow 1} H(\square), \quad (5)$$

where $\alpha > 0$ is real, $P(\square)$ is a generic function of $l^2\square$, and $f(\omega)$ is arbitrary. The parameter σ is fictitious and will be commented upon shortly. The kernel \mathcal{G} is formally given by $\mathcal{G}(x, y; \sigma) = e^{H_\sigma(\square_x)} \mathcal{G}(x, y; 0)$, but this expression is especially difficult to deal with because in nonlocal quantum gravities the propagator is suppressed in the ultraviolet, so that its inverse (the form factor $\exp H$) explodes in Euclidean momentum space in all realistic cases.¹ To avoid this, we consider the inverse $\mathcal{F} = \mathcal{G}^{-1}$, for which the nonlocal operator in

$$\mathcal{F}(x, y; \sigma) = e^{-H_\sigma(\square_x)} \mathcal{F}(x, y; 0) \quad (6)$$

is damped at high energies. Once \mathcal{F} is found, one can determine $\mathcal{G} = e^{2H_\sigma} \mathcal{F}$ with deconvolution methods [33]. However, Eq. (6) still carries all the above-mentioned unknowns related to infinitely many derivatives.

To solve this issue, we make a crucial observation: any form factor can be written in terms of a kernel \mathcal{F} governed by a simple system of renormalization-group-like equations, which determine how \mathcal{F} varies in the *space of all possible functionals* $P(\square)$. This space is parametrized by σ , where $\sigma = 1$ corresponds to the end of the flow. The exact form of these master equations depends on the choice of $H(\square)$, which is limited by renormalizability and unitarity.

For the general class $f(\omega) = \exp(-\omega)$ (asymptotically polynomial H^{pol} ; includes Kuz'min and Tomboulis form

¹ For instance, for Kuz'min Euclidean form factor (see below) $H_\sigma(-k^2) = \alpha[\ln(l^2 k^2) + \Gamma(0, l^2 k^2) + \gamma_E] \rightarrow +\infty$ as $|k| \rightarrow \infty$; for the string form factor, $H_\sigma(-k^2) = \alpha \sigma l^2 k^2$; and so on.

factors), we can finally write the finite-order differential equations:

$$\sigma \partial_\sigma \mathcal{F}(x, y; \sigma) = \alpha [(\mathcal{K} \star \mathcal{F})(x, y; \sigma) - \mathcal{F}(x, y; \sigma)], \quad (7a)$$

$$(\mathcal{K} \star \mathcal{F})(x, y; \sigma) := \int d^D x' \sqrt{-g} \mathcal{K}(x, x'; \sigma) \mathcal{F}(x', y; \sigma), \quad (7b)$$

$$[\partial_\sigma + P(\square_x)] \mathcal{K}(x, y; \sigma) = 0, \quad (7c)$$

$$\mathcal{F}(x, y; 0) = \mathcal{K}(x, y; 0) = [-g(y)]^{-1/2} \delta^D(x - y), \quad (7d)$$

where \mathcal{K} is the kernel associated with the operator $f[\sigma P(\square)]$. The solution of Eq. (7a) is formally given by Eq. (6). Note that Eq. (7c) corresponds, in quantum gravities, to diffusion in the correct direction and its solution $\mathcal{K}(x, y; \sigma) = e^{-\sigma P(\square_x)} \mathcal{K}(x, y; 0)$ is well defined.

In the much simpler case of the general class $f(\omega) = 1 - \omega$ (exactly monomial H_σ^{exp} ; includes the string-related and Krasnikov exponential form factors), Eqs. (7) are replaced by just one master equation:

$$[\partial_\sigma + \alpha P(\square_x)] \mathcal{F}(x, y; \sigma) = 0, \quad (8)$$

whose solution $\mathcal{F}(x, y; \sigma) = e^{-\alpha \sigma P(\square_x)} \mathcal{F}(x, y; 0)$ is well defined and coincides with Eq. (6). The integro-differential equations (7) or (8) are the generalizations of the diffusion-equation method [32, 34] and of the case with an exactly polynomial $H(\square)$ treated in Ref. [35]. Thus, the original, formidable nonlocal problem is reduced to one with a finite number of derivatives where we can count the initial conditions and the field degrees of freedom. Both turn out to be finite.

Initial conditions for special form factors. Let us consider the system (7) with $P(\square) = -l^2 \square$. The system given by Eqs. (7d), (7a), and $(\partial_\sigma - l^2 \square_x) \mathcal{K}(x, x'; \sigma) = 0$ is second order in spacetime coordinates and first order in the diffusion parameter σ . In synchronous gauge, the metric in D dimensions simplifies to $ds^2 = -dt^2 + h_{ij}(t, \mathbf{x}) dx^i dx^j$, where $i, j = 1, \dots, D-1$, h_{ij} is the metric of the spatial section, and the covariant d'Alembertian operator on a scalar takes the form

$$\square = -\partial_t^2 - \frac{1}{2} h^{ij} \dot{h}_{ij} \partial_t + \frac{1}{\sqrt{h}} \partial_i (\sqrt{h} h^{ij} \partial_j). \quad (9)$$

Therefore, to solve the second-order master equations we only need to specify the spatial metric and its first time derivative. If we insist in having only second-order (in spacetime coordinates) differential equations with a diffusion-like structure, then *there are only two choices for the form factor (2)*: Kuz'min's profile $H^{\text{pol}}(\square)$ with $P(\square) = -l^2 \square$ or the string-related $H^{\text{exp}}(\square) = -l^2 \square$. By removing one ambiguity of the problem [choice of $P(\square)$] once the other [choice of $f(\omega)$] is fixed by requiring a diffusion equation, the long-standing question about the uniqueness of nonlocal gravity is solved. Since the generalized diffusion-equation method (7) can be applied to any $f(\omega)$, it permits to classify all allowed form factors.

Therefore, when $P(\square) = -l^2 \square$ the whole nonlocality in the action and in the equations of motion (EOM) is completely specified by second-order differential equations,

together with the metric and the first time derivative of the spatial metric. If we impose the retarded boundary condition $\tilde{G}(x, y) = 0$ for $y^0 > x^0$, Eq. (4) defines the retarded Green function \tilde{G} at time x^0 only from the value of h_{ij} and its first time derivative, for times $\leq x^0$.

Equations of motion. Let us comment on the variation of the action with respect to the metric for an asymptotically polynomial profile $H_\sigma = \alpha \int_0^{\sigma P} d\omega \omega^{-1} [1 - \exp(-\omega)] = \alpha \int_0^P d\omega \int_0^\sigma d\sigma' \exp(-\sigma' \omega)$ and a monomial $P(\square) = (l^2 \square)^n$. The most delicate term is the variation of the form factor. Iterating multiple times Duhamel's formula $\delta e^A = \int_0^1 ds e^{sA} \delta A e^{(1-s)A}$ for an operator A and using $\delta \square^{-1} = -\square^{-1} (\delta \square) \square^{-1}$ and $\delta H_\sigma = -\alpha \int_0^\sigma d\sigma' \sigma'^{-1} \delta \exp(-\sigma' P)$, we get

$$\delta \gamma = -\frac{1}{\square} (\delta \square) \gamma + \frac{\alpha l^{2n}}{\square} \sum_{k=1}^n \int_0^1 ds e^{s H_\sigma} \int_0^\sigma d\sigma' \int_0^1 ds' \times e^{-s \sigma' P} \square^{k-1} (\delta \square) \square^{n-k} e^{(s-1) \sigma' P} e^{(1-s) H_\sigma}. \quad (10)$$

The full, explicit EOM are reported in [35]. Note that all nonlocal form factors involved in the variation of $\gamma(\square)$ are of the type \square^{-1} , $\exp \square^n$, and $\exp H$. Therefore, the form factors satisfy Eq. (7d) and (8) for the case of $\exp \square$ or the master equations (7d) and (7) for the case of $\exp H$.

The conclusion is that the EOM can be expressed in terms of the kernels \mathcal{K} and \mathcal{G} living in the space of form factors and the Green function $\tilde{G}(x, y)$. These kernels are nonlocal because they depend on two spacetime points, but they are determined by a set of equations independent of the actual EOM. This set may be more or less difficult to solve, but it is well defined. Therefore, the EOM coming from Eq. (3) are nonlocal integral equations but *finite* differential equations: for $P(\square) = O(\square^n)$, they are of derivative order $2n + 2$ and we must specify $2n + 2$ initial conditions. This result agrees with the counting done in Deser–Woodard nonlocal gravity [36], with that in a nonlocal scalar field theory [34, 37], and with the diffusion method in Lagrangian formalism [35].

Up to this point, we concentrated on a class of super-renormalizable theories, but the main result is insensitive to the introduction of other special operators that make the theory finite in even dimension. Here by special we mean nonlocal “killer operators” [10] at least cubic in the curvature, namely, $\mathcal{R} \gamma_{k,1}(\square) \mathcal{R}^2$, $\mathcal{R}^2 \gamma_{k,2}(\square) \mathcal{R}^2$, \dots , $\mathcal{R}^{D/2} \gamma_{k,\dots}(\square) \mathcal{R}^2$, $\mathcal{R} \square \mathcal{R}^{\frac{D}{2}-2} \gamma_{k,\dots}(\square) \mathcal{R}^2$, where $\gamma_{k,\dots}$ may all differ from one another and, again, \mathcal{R} is the generalized curvature. These operators increase the order of the EOM. In $D = 4$, killer operators are cubic or quartic in the Riemann tensor and the order of the EOM is the same, namely, $2n + 2$. In particular, for $n = 1$ four-dimensional finite nonlocal gravity only needs four initial condition at the nonperturbative level.

Degrees of freedom. In the minimal case $n = 1$ in $D = 4$, the EOM are fourth order (just like in local quadratic gravity) and the degrees of freedom are eight. To show this, we introduce two auxiliary fields, a rank-2 symmetric tensor $\phi^{\mu\nu}$ and a scalar field ψ , to infer the

exact number of d.o.f.. Let us consider the Lagrangian

$$2\kappa^2 \mathcal{L} = R + 2G_{\mu\nu} \gamma(\square) \phi^{\mu\nu} - \phi_{\mu\nu} \gamma(\square) \phi^{\mu\nu} + R \gamma(\square) \psi + \psi \gamma(\square) \psi / (D-2). \quad (11)$$

The EOM for the tensor $\phi^{\mu\nu}$ and the scalar ψ are

$$\phi_{\mu\nu} = G_{\mu\nu}, \quad \psi = \phi = G. \quad (12)$$

ψ is just the trace of $\phi_{\mu\nu}$ and is not an independent degree of freedom [35]. Eliminating the auxiliary fields from the action, we end up with the original action (1). Notice that Eq. (12) implies the transverse condition $\nabla^\mu \phi_{\mu\nu} = 0$ on-shell, as a consequence of Bianchi identity. The EOM for the metric are more involved, but not overly so, and agree with the case without auxiliary fields [35]. It turns out that we only deal with second-order differential equations, up to γ factors that, as we have shown, can be dealt with the diffusion-equation method without increasing the derivative order. Therefore, the d.o.f. are: (I) Graviton $g_{\mu\nu}$: symmetric $D \times D$ matrix with $D(D+1)/2$ independent entries, to which one subtracts D Bianchi identities $\nabla^\mu G_{\mu\nu} = 0$ and D diffeomorphisms (the theory is fully diffeomorphism invariant). Total: $D(D-3)/2$. In $D=4$, there are 2 degrees of freedom, the usual polarization modes. (II) Tensor $\phi_{\mu\nu}$: symmetric $D \times D$ matrix to which one subtracts D transverse conditions $\nabla^\mu \phi_{\mu\nu} = 0$. Total: $D(D-1)/2$. In $D=4$, there are 6 degrees of freedom. The grand total is $D(D-2)$. In $D=4$ dimensions, there are eight degrees of freedom.

Applications and conclusions. Some years ago, the diffusion-equation method revealed that a string-motivated nonlocal scalar field theory with exponential operators has a finite number of initial conditions and nonperturbative degrees of freedom [34]. In this Letter, we converted the infinite number of derivatives of renormalizable nonlocal gravities with highly nontrivial nonlocal operators into integral kernels living in the space of all possible form factors and that do not carry pathological nonlocality. The main action (3) must be placed side by side with the finite-order differential equations given by

the system (7) or (8). The system of EOM for the metric and the kernels is of finite order $2n+2$ for the metric, for a form factor (2) with H given by H^{Pol} with $P(\square) \propto \square^n$. For the special form factor with $n=1$, where kernels satisfy second-order differential equation, the EOM for the metric are fourth order and the number of nonperturbative degrees of freedom is eight in four dimensions. Two of these d.o.f. are perturbative and well known in the literature, and describe completely the theory at small scales on a Minkowski background, thanks to asymptotic freedom [9]. Therefore, the other degrees of freedom are nonperturbative and do not propagate on Minkowski spacetime at very small scales, but they should play a capital role *at large scales and on curved backgrounds*. This is the most direct and important consequence of our findings for the physics of nonlocal gravity.

These results repair the old and bad reputation of nonlocal theories for having an ill-defined Cauchy problem. Armed with the generalized diffusion method, one can construct unambiguous analytic solutions of cosmological and astrophysical backgrounds, previously inaccessible via the typical *Ansatz* $\square R = \lambda R$ in the literature. From these new, fully nonperturbative solutions, one will be able to extract conclusions on the stability or absence of big-bang and black-hole singularities, find predictions on the cosmic evolution induced by nonlocal gravity, and check the theory at the nonperturbative level against present and near-future observations, such as those of PLANCK [38, 39] and LIGO [40, 41], according to a varied battery of phenomenological tests that has already been effective for string cosmology and other quantum gravities [3, 42]. Supported by these, other recent, and upcoming data, the phenomenology and testing of nonlocal quantum gravity is about to bloom.

ACKNOWLEDGMENTS

G.C. is under a Ramón y Cajal contract. G.C. and L.M. are supported by the MINECO I+D grants FIS2014-54800-C2-2-P and FIS2017-86497-C2-2-P.

-
- [1] D. Oriti (Ed.), *Approaches to Quantum Gravity* (Cambridge University Press, Cambridge, U.K., 2009).
 - [2] G.F.R. Ellis, J. Murugan, and A. Weltman (Eds.), *Foundations of Space and Time* (Cambridge University Press, Cambridge, U.K., 2012).
 - [3] G. Calcagni, *Classical and Quantum Cosmology* (Springer, Switzerland, 2017).
 - [4] N.V. Krasnikov, Nonlocal gauge theories, *Theor. Math. Phys.* **73**, 1184 (1987) [*Teor. Mat. Fiz.* **73**, 235 (1987)].
 - [5] Y.V. Kuz'min, The convergent nonlocal gravitation, *Yad. Fiz.* 50 (1989) 1630 [*Sov. J. Nucl. Phys.* 50 (1989) 1011].
 - [6] E.T. Tomboulis, Super-renormalizable gauge and gravitational theories, [arXiv:hep-th/9702146](https://arxiv.org/abs/hep-th/9702146).
 - [7] L. Modesto, Super-renormalizable quantum gravity, *Phys. Rev. D* **86**, 044005 (2012) [[arXiv:1107.2403](https://arxiv.org/abs/1107.2403)].
 - [8] T. Biswas, E. Gerwick, T. Koivisto, and A. Mazumdar, Towards singularity and ghost free theories of gravity, *Phys. Rev. Lett.* **108**, 031101 (2012) [[arXiv:1110.5249](https://arxiv.org/abs/1110.5249)].
 - [9] G. Calcagni and L. Modesto, Nonlocal quantum gravity and M-theory, *Phys. Rev. D* **91**, 124059 (2015) [[arXiv:1404.2137](https://arxiv.org/abs/1404.2137)].
 - [10] L. Modesto and L. Rachwał, Super-renormalizable and finite gravitational theories, *Nucl. Phys. B* **889**, 228 (2014) [[arXiv:1407.8036](https://arxiv.org/abs/1407.8036)].
 - [11] S. Talaganis, T. Biswas, and A. Mazumdar, Towards understanding the ultraviolet behavior of quantum loops in infinite-derivative theories of gravity, *Classical Quantum Gravity* **32**, 215017 (2015) [[arXiv:1412.3467](https://arxiv.org/abs/1412.3467)].
 - [12] L. Modesto and L. Rachwał, Universally finite gravitational and gauge theories, *Nucl. Phys. B* **900**, 147 (2015) [[arXiv:1503.00261](https://arxiv.org/abs/1503.00261)].

- [13] E.T. Tomboulis, Renormalization and unitarity in higher derivative and nonlocal gravity theories, *Mod. Phys. Lett. A* **30**, 1540005 (2015).
- [14] A.S. Koshelev, L. Modesto, L. Rachwał, and A.A. Starobinsky, Occurrence of exact R^2 inflation in non-local UV-complete gravity, *J. High Energy Phys.* 1611 (2016) 067 [[arXiv:1604.03127](#)].
- [15] T. Biswas, A. Mazumdar, and W. Siegel, Bouncing universes in string-inspired gravity, *J. Cosmol. Astropart. Phys.* 0603 (2006) 009 [[arXiv:hep-th/0508194](#)].
- [16] G. Calcagni, G. Nardelli, Nonlocal gravity and the diffusion equation, *Phys. Rev. D* **82**, 123518 (2010) [[arXiv:1004.5144](#)].
- [17] T. Biswas, T. Koivisto, and A. Mazumdar, Towards a resolution of the cosmological singularity in non-local higher derivative theories of gravity, *J. Cosmol. Astropart. Phys.* 1011 (2010) 008 [[arXiv:1005.0590](#)].
- [18] A.S. Koshelev and S.Yu. Vernov, On bouncing solutions in non-local gravity, *Phys. Part. Nucl.* **43**, 666 (2012) [[arXiv:1202.1289](#)].
- [19] T. Biswas, A.S. Koshelev, A. Mazumdar, and S.Yu. Vernov, Stable bounce and inflation in non-local higher derivative cosmology, *J. Cosmol. Astropart. Phys.* 1208 (2012) 024 [[arXiv:1206.6374](#)].
- [20] A.S. Koshelev, Stable analytic bounce in non-local Einstein–Gauss–Bonnet cosmology, *Classical Quantum Gravity* **30**, 155001 (2013) [[arXiv:1302.2140](#)].
- [21] G. Calcagni, L. Modesto, and P. Nicolini, Super-accelerating bouncing cosmology in asymptotically-free non-local gravity, *Eur. Phys. J. C* **74**, 2999 (2014) [[arXiv:1306.5332](#)].
- [22] J. Edholm, A.S. Koshelev, and A. Mazumdar, Behavior of the Newtonian potential for ghost-free gravity and singularity-free gravity, *Phys. Rev. D* **94**, 104033 (2016) [[arXiv:1604.01989](#)].
- [23] L. Modesto and L. Rachwał, Finite conformal quantum gravity and nonsingular spacetimes, [arXiv:1605.04173](#).
- [24] V.P. Frolov, Notes on nonsingular models of black holes, *Phys. Rev. D* **94**, 104056 (2016) [[arXiv:1609.01758](#)].
- [25] C. Bambi, L. Modesto, and L. Rachwał, Spacetime completeness of non-singular black holes in conformal gravity, *J. Cosmol. Astropart. Phys.* 1705 (2017) 003 [[arXiv:1611.00865](#)].
- [26] A.S. Koshelev and A. Mazumdar, Absence of event horizon in massive compact objects in infinite derivative gravity, *Phys. Rev. D* **96**, 084069 (2017) [[arXiv:1707.00273](#)].
- [27] G. Calcagni and L. Modesto, Stability of Schwarzschild singularity in non-local gravity, *Phys. Lett. B* **773**, 596 (2017) [[arXiv:1707.01119](#)].
- [28] Y.S. Myung and Y.J. Park, Stability issues of black hole in non-local gravity, *Phys. Lett. B* **779**, 342 (2018) [[arXiv:1711.06411](#)].
- [29] J. Edholm, Conditions for defocusing around more general metrics in infinite derivative gravity, [arXiv:1802.09063](#).
- [30] G. Calcagni, M. Montobbio, and G. Nardelli, Route to nonlocal cosmology, *Phys. Rev. D* **76**, 126001 (2007) [[arXiv:0705.3043](#)].
- [31] M. Asorey, J.L. López, and I.L. Shapiro, Some remarks on high derivative quantum gravity, *Int. J. Mod. Phys. A* **12**, 5711 (1997) [[arXiv:hep-th/9610006](#)].
- [32] G. Calcagni and G. Nardelli, String theory as a diffusing system, *J. High Energy Phys.* 1002 (2010) 093 [[arXiv:0910.2160](#)].
- [33] W. Ulmer, Inverse problem of linear combinations of Gaussian convolution kernels (deconvolution) and some applications to proton/photon dosimetry and image processing, *Inverse Problems* **26**, 085002 (2010).
- [34] G. Calcagni, M. Montobbio, and G. Nardelli, Localization of nonlocal theories, *Phys. Lett. B* **662**, 285 (2008) [[arXiv:0712.2237](#)].
- [35] G. Calcagni, L. Modesto, and G. Nardelli, Initial conditions and degrees of freedom of non-local gravity, *J. High Energy Phys.* 1805 (2018) 087 [[arXiv:1803.00561](#)].
- [36] S. Deser and R.P. Woodard, Observational viability and stability of nonlocal cosmology, *J. Cosmol. Astropart. Phys.* 1311 (2013) 036 [[arXiv:1307.6639](#)].
- [37] E.T. Tomboulis, Nonlocal and quasilocal field theories, *Phys. Rev. D* **92**, 125037 (2015) [[arXiv:1507.00981](#)].
- [38] P.A.R. Ade et al. [Planck Collaboration], Planck 2015 results. XIII. Cosmological parameters, *Astron. Astrophys.* **594**, A13 (2016) [[arXiv:1502.01589](#)].
- [39] P.A.R. Ade et al. [Planck Collaboration], Planck 2015 results. XX. Constraints on inflation, *Astron. Astrophys.* **594**, A20 (2016) [[arXiv:1502.02114](#)].
- [40] B.P. Abbott et al. [LIGO Scientific and Virgo Collaborations], Observation of gravitational waves from a binary black hole merger, *Phys. Rev. Lett.* **116**, 061102 (2016) [[arXiv:1602.03837](#)].
- [41] B.P. Abbott et al. [LIGO Scientific and Virgo Collaborations], GW151226: observation of gravitational waves from a 22-solar-mass binary black hole coalescence, *Phys. Rev. Lett.* **116**, 241103 (2016) [[arXiv:1606.04855](#)].
- [42] G. Calcagni, Multifractional theories: an unconventional review, *J. High Energy Phys.* 1703 (2017) 138 [[arXiv:1612.05632](#)].