

GIANT DIPOLE RESONANCE PARAMETERS OF GROUND-STATE PHOTOABSORPTION: EXPERIMENTAL VALUES WITH UNCERTAINTIES

V. A. Plujko^a, O. M. Gorbachenko^a, R. Capote^{b,*}, P. Dimitriou^b

^a*Nuclear Physics Department, Taras Shevchenko National University, Kyiv, Ukraine*

^b*NAPC-Nuclear Data Section, International Atomic Energy Agency, Vienna, Austria*

Abstract

Updated values and corresponding uncertainties of Isovector Giant Dipole Resonance (GDR) parameters which are obtained by the least-squares fitting of theoretical photoabsorption cross sections to experimental data are presented. The theoretical photoabsorption cross sections are taken as a sum of the components corresponding to the excitation of the GDR and quasideuteron photodisintegration. The current compilation is an extension and improvement of the earlier compilations of Lorentzian parameters for ground-state photoneutron and photoabsorption cross sections and covers experimental data made available up to June 2017.

arXiv:1804.04445v1 [nucl-th] 12 Apr 2018

*Corresponding author.

Email address: R.CapoteNoy@iaea.org (R. Capote)

Contents

1. Introduction and general description of the updates	2
2. Theoretical basis and assumptions	4
3. Discussion and Conclusion	7
References	9
Explanation of Tables	12
Explanation of Graphs.....	15
Tables	
1. Experimental values and uncertainties of GDR parameters within Standard Lorentzian (SLO) approach.	21
2. Experimental values and uncertainties of GDR parameters within Modified Lorentzian (SMLO) approach.....	29
3. References to experimental and evaluated cross section data taken from EXFOR.	37
Graphs 1 - 162	
.....	43

1. Introduction and general description of the updates

Isovector Giant Dipole Resonances (GDR) are fundamental modes of nuclear collective excitations beyond the binding energy per nucleon. Nuclear collective states correspond to vibrations of the nucleons and are strongly manifested in electric dipole (E1) gamma-transitions which dominate over transitions of other multipole order when they occur simultaneously (see, for example, [1–3] for references). This allows us to obtain information on the GDR characteristics (i.e., resonance energy, width and contribution of the GDR to the energy-weighted sum rule (EWSR)) from investigations of the electromagnetic processes of photoabsorption and gamma-decay. However, E1 transitions are dominant near the maximum of the photoabsorption cross sections, but not at lower photon energies closer to the neutron binding energies. There, other physical phenomena like pygmy resonances [4] and scissors mode [5, 6] need to be considered. Those low-energy collective phenomena have little impact on the nuclear GDR characteristics and are neglected in this work. It can be also noted that collective states like GDR have been extensively studied, both experimentally and theoretically, in atomic physics [7, 8], metallic clusters [7, 9, 10], and quantum-dots [4, 11].

A comprehensive experimental database with reliable data for the GDR parameters and their uncertainties is very important for the reliable modeling of E1 gamma-ray cascades in highly excited nuclei, for the study of nuclear reaction mechanisms as well as for verifying different theoretical approaches used to describe the GDR and other nuclear structure properties (deformations, contribution of velocity-dependent forces, shape-transitions, etc), and forms an integral part of modern nuclear reaction computer codes, such as [12, 13]. The experimental values of the GDR parameters in cold atomic nuclei are most reliably deduced from experimental photoabsorption cross sections. Compilations of the parameters of Lorentz curves fitted to the total photoneutron cross section data were presented in Refs. [14–16]. The data from Ref.[16] including GDR parameters for light nuclei ^{12}C , ^{14}N , ^{16}O , ^{27}Al and ^{28}Si were listed in the RIPL-1 database [1] as well as in the RIPL-3 [3] /gamma/gdr-parameters-exp.dat file. Assuming that the contribution of photoproton cross sections to the total photoabsorption cross section is small, then the Lorentzian parameters of the total photoneutron cross sections

in spherical and axially deformed nuclei can be identified with the GDR parameters. However, note that this assumption is poor for light nuclei where photo-charged particle reactions are important.

Updated tables of GDR parameters with estimates of their uncertainties were subsequently given in Ref. [17]. In this compilation, the GDR parameters were treated as variables in the fitting of calculated total photoabsorption cross sections to the experimental data using the least-squares method.

Comprehensive databases of photonuclear reaction parameters are also published in Refs. [18, 19]. The photoproton contribution was included in the database, but the parameters were obtained without performing any specific fit of an analytical function but were extracted directly by digitizing the characteristics of the experimental data peaks. Microscopic predictions of the GDR energies and widths for about 6000 nuclei between the proton and the neutron drip lines are given in the RIPL-3 database [3]. These GDR parameters resulted from a fit of microscopic calculations of the nuclear gamma-strength function to the existing experimental data [20, 21]. The calculations were performed on the basis of the Hartree-Fock-BCS plus quasi-particle random-phase-approximation as well as the microscopic Hartree-Fock-Bogoliubov plus quasi-particle random phase approximation model with a realistic Skyrme interaction.

For heated atomic nuclei, the GDR parameters are determined by gamma-decay data. Compilation and parameterization of the GDR built on excited states are given in Refs. [22, 23].

Tables of updated values of the GDR parameters with estimates of their uncertainties (one-sigma standard deviation) are given in this contribution. This work was performed within an IAEA Coordinated Research Project on Updating the Photonuclear Data Library and generating a Reference Database for Photon Strength Functions [24]. The presented database updates and extends the above-mentioned compilations for the GDR built on cold nuclear states. The parameters are calculated following the prescription of Ref. [17] by fitting Lorentz-like curves to photonuclear cross section data using the least-squares method. For experimental data, the total photoabsorption cross sections retrieved from the EXFOR database [25] or a combination of experimental partial cross sections best suited for approximating the total photoabsorption cross section are used. The evaluated data of [26, 27] are also considered as experimental values. Parameters are given for 144 isotopes from ${}^6\text{Li}$ to ${}^{239}\text{Pu}$ atomic nuclei and for 19 elements of natural isotopic composition (in total 475 entries).

The updated values of the GDR parameters with their associated uncertainties (one-sigma standard deviation) are given in the Tables 1 and 2. They were obtained by fitting the data with two different models of the GDR excitation: a Standard Lorentzian (SLO) model and a Simplified version of the Modified Lorentzian (SMLO) approach [2, 3, 17, 35] (see below Eqs. (2)-(5)). The updated tables include the following improvements with respect to the previous results reported in Ref.[17]:

- GDR parameters were determined for 164 datasets, including the GDR parameters for 14 new isotopes and 13 natural elements, i.e. for ${}^6\text{Li}$, ${}^7\text{Li}$, ${}^9\text{Be}$, ${}^{108}\text{Pd}$, ${}^{112}\text{Sn}$, ${}^{114}\text{Sn}$, ${}^{122}\text{Sn}$, ${}^{151}\text{Eu}$, ${}^{152}\text{Gd}$, ${}^{154}\text{Gd}$, ${}^{158}\text{Gd}$, ${}^{185}\text{Re}$, ${}^{203}\text{Tl}$, ${}^{205}\text{Tl}$ and for ${}^{\text{nat}}\text{C}$, ${}^{\text{nat}}\text{Mg}$, ${}^{\text{nat}}\text{Cu}$, ${}^{\text{nat}}\text{Rb}$, ${}^{\text{nat}}\text{Sr}$, ${}^{\text{nat}}\text{Pd}$, ${}^{\text{nat}}\text{Ag}$, ${}^{\text{nat}}\text{Cd}$, ${}^{\text{nat}}\text{Sb}$, ${}^{\text{nat}}\text{Ba}$, ${}^{\text{nat}}\text{Re}$, ${}^{\text{nat}}\text{Ir}$, ${}^{\text{nat}}\text{U}$.
- GDR parameters were corrected for 23 datasets by revising the energy intervals of the fitting procedure.
- GDR parameters for 7 datasets, namely for isotopes ${}^{12}\text{C}$, ${}^{15}\text{N}$, ${}^{\text{nat}}\text{Ar}$, ${}^{\text{nat}}\text{Fe}$, ${}^{59}\text{Co}$, ${}^{\text{nat}}\text{Zr}$ were excluded due to the poor quality of the fit of the experimental data.

Compared to the RIPL-3 Tables [3], the new tables include the following modifications:

- GDR parameters were added for 271 datasets, including 24 new isotopes that were absent previously, i.e. ${}^6\text{Li}$, ${}^7\text{Li}$, ${}^9\text{Be}$, ${}^{10}\text{B}$, ${}^{65}\text{Zn}$, ${}^{108}\text{Pd}$, ${}^{109}\text{Ag}$, ${}^{112}\text{Sn}$, ${}^{114}\text{Sn}$, ${}^{122}\text{Sn}$, ${}^{151}\text{Eu}$, ${}^{152}\text{Gd}$, ${}^{154}\text{Gd}$, ${}^{158}\text{Gd}$, ${}^{176}\text{Hf}$, ${}^{185}\text{Re}$, ${}^{191}\text{Ir}$, ${}^{193}\text{Ir}$, ${}^{194}\text{Pt}$, ${}^{195}\text{Pt}$, ${}^{196}\text{Pt}$, ${}^{198}\text{Pt}$, ${}^{203}\text{Tl}$, ${}^{205}\text{Tl}$ and 19 elements of natural composition: ${}^{\text{nat}}\text{C}$, ${}^{\text{nat}}\text{O}$, ${}^{\text{nat}}\text{Mg}$, ${}^{\text{nat}}\text{Si}$, ${}^{\text{nat}}\text{S}$, ${}^{\text{nat}}\text{K}$, ${}^{\text{nat}}\text{Ca}$, ${}^{\text{nat}}\text{Cu}$, ${}^{\text{nat}}\text{Rb}$, ${}^{\text{nat}}\text{Sr}$, ${}^{\text{nat}}\text{Ag}$, ${}^{\text{nat}}\text{Pd}$, ${}^{\text{nat}}\text{Cd}$, ${}^{\text{nat}}\text{Sb}$, ${}^{\text{nat}}\text{Ba}$, ${}^{\text{nat}}\text{Re}$, ${}^{\text{nat}}\text{Ir}$, ${}^{\text{nat}}\text{Pb}$, ${}^{\text{nat}}\text{U}$.
- The GDR values for 40 datasets were corrected by revising the energy interval of fitting and modifying the uncertainty estimates.
- The GDR parameters from 8 datasets, namely ${}^{15}\text{N}$, ${}^{19}\text{F}$, ${}^{26}\text{Mg}$, ${}^{51}\text{V}$, ${}^{59}\text{Co}$, ${}^{127}\text{I}$, ${}^{141}\text{Pr}$, ${}^{197}\text{Au}$ were excluded due to the poor quality of the fit to experimental data.

Figures 1–162 show the comparison between the experimental photonuclear cross sections and theoretical calculations for a more extended range of gamma-ray energies ($\lesssim 30$ MeV) compared to those used in the fitting procedure. The theoretical cross sections were calculated using the SLO and SMLO models for the GDR excitation, with parameters taken from Tables 1 and 2, respectively, and including the quasi-deuteron photodisintegration. An estimate of the quality of the description of the experimental data by the two fitted theoretical models is provided by the ratios of the least-square deviations (Eq.(11)) of model SMLO to SLO which are calculated for the different gamma-ray energy intervals and are also indicated in the Table A and in the graphs.

2. Theoretical basis and assumptions

A number of different simplified resonance shapes (consisting of one or a sum of the resonance components) is used for the theoretical description of photoabsorption cross sections. The photonuclear data around the peaks of the GDR can be fitted by Lorentz, Breit-Wigner and Gaussian functions equally well, but their low- and high- energy tails are a different matter [28, 29]. Lorentz (G_L) and Breit-Wigner (G_{BW}) shapes of the photoabsorption cross sections can be derived from different theoretical approaches [30–35]. The Lorentzian shape can be transformed to Breit-Wigner,

$$G_L(\varepsilon; E, \Gamma) \equiv \frac{\varepsilon^2 \Gamma^2}{(\varepsilon^2 - E^2)^2 + (\varepsilon \Gamma)^2} = \frac{\gamma^2(\varepsilon)/4}{(\varepsilon - E)^2 + \gamma^2(\varepsilon)/4} \equiv G_{BW}(\varepsilon; E, \gamma(\varepsilon)), \quad (1)$$

but with shape width $\gamma(\varepsilon) = (2\varepsilon/(\varepsilon + E))\Gamma$ which depends on photon energy ε , resonance energy E and width Γ of the Lorentzian. In the neighborhood of the resonance energy, the width $\gamma(\varepsilon)$ coincides with Γ , i.e. $\gamma(\varepsilon \simeq E) \simeq \Gamma$, however on the wings of the resonance it differs. The Breit-Wigner shape G_{BW} with arbitrary energy dependent width is sometimes referred to as the Energy Dependent Breit-Wigner [28].

A Lorentz shape is preferable for fitting the photonuclear data because the standard Breit-Wigner expression is obtained without account for time reversal invariance and is adequate for describing a strong resonance state when the width is small with respect to the resonance energy [31, 32, 34]. These line shapes correspond to a nuclear response function to an electromagnetic field which proceeds through the excitation of one strong collective state that exhausts the EWSR. However, the photoabsorption and gamma-decay processes on the wings of the GDR are governed by the excitation of states of a different nature and therefore, the fitting of experimental data by Lorentzians should be limited to small energy ranges around the GDR peak [14, 16, 17] in order to obtain reliable values for the GDR parameters.

In this contribution, the theoretical photoabsorption cross section $\sigma_{abs}(\varepsilon_\gamma)$ for a photon with energy ε_γ is taken as a sum of the terms corresponding to the GDR excitation given by $\sigma_{GDR}(\varepsilon_\gamma)$ and the quasi-deuteron photodisintegration:

$$\sigma_{abs}(\varepsilon_\gamma) = \sigma_{GDR}(\varepsilon_\gamma) + \sigma_{QD}(\varepsilon_\gamma), \quad (2)$$

where the component $\sigma_{QD}(\varepsilon_\gamma)$ is a quasi-deuteron contribution, i.e., the photoabsorption cross section on a neutron-proton pair [18, 36]. For a nucleus with N neutrons, Z protons and mass number $A = N + Z$, the latter term is equal to

$$\sigma_{QD}(\varepsilon_\gamma) = 397.8 \frac{NZ}{A} \frac{(\varepsilon_\gamma - 2.224)^{3/2}}{\varepsilon_\gamma^3} \phi(\varepsilon_\gamma) \quad (3)$$

with ε_γ in MeV and σ_{QD} in units of mb. The function $\phi(\varepsilon_\gamma)$ accounts for the Pauli-blocking of the excited neutron-proton pair in the nuclear medium

$$\begin{aligned} \phi(\varepsilon_\gamma < 20 \text{ MeV}) &= \exp(-73.3/\varepsilon_\gamma), \\ \phi(20 < \varepsilon_\gamma < 140 \text{ MeV}) &= 8.3714 \times 10^{-2} - 9.8343 \times 10^{-3} \varepsilon_\gamma + 4.1222 \times 10^{-4} \varepsilon_\gamma^2 - \\ &\quad - 3.4762 \times 10^{-6} \varepsilon_\gamma^3 + 9.3537 \times 10^{-9} \varepsilon_\gamma^4, \\ \phi(\varepsilon_\gamma > 140 \text{ MeV}) &= \exp(-24.2/\varepsilon_\gamma). \end{aligned} \quad (4)$$

The GDR component of the photoabsorption cross section $\sigma_{abs}(\varepsilon_\gamma)$ is considered to be equal to the photoabsorption cross section of electric dipole (E1) gamma-rays and is calculated within framework of two different Lorentzian models: a Standard Lorentzian (SLO) model and a Simplified version of the Modified Lorentzian (SMLO) approach [2, 3, 17, 35]. For deformed nuclei, an approximation applied to axially deformed nuclei is adopted [2, 3, 17]. However, following Ref. [16], some deformed nuclei are treated as spherical ones, in cases where the fit of the experimental data by the one-component Lorentz curve has a lower χ^2 per degree of freedom than the fit by a two-component Lorentzian. Assuming axially symmetric shapes, the expression for σ_{GDR} is a sum of two Lorentz-like curves corresponding to collective vibrations along and perpendicular to the axis of symmetry ($j_m = 2$), respectively (for spherical nuclei $j_m = 1$ or is omitted)

$$\begin{aligned} \sigma_{GDR}(\varepsilon_\gamma) &= \sigma_{GDR}^\alpha(\varepsilon_\gamma) = \sum_{j=1}^{j_m} \sigma_{GDR,j}^\alpha(\varepsilon_\gamma) = \sigma_{TRK} s_j^\alpha \cdot F_j^\alpha(\varepsilon_\gamma), \\ F_j^\alpha(\varepsilon_\gamma) &\equiv \frac{2}{\pi} \frac{G_L(\varepsilon_\gamma; E_{r,j}^\alpha, \Gamma_j^\alpha)}{\Gamma_j^\alpha} = \frac{2}{\pi} \frac{\varepsilon_\gamma^2 \Gamma_j^\alpha}{[\varepsilon_\gamma^2 - (E_{r,j}^\alpha)^2]^2 + [\varepsilon_\gamma \Gamma_j^\alpha]^2}. \end{aligned} \quad (5)$$

Here, $E_{r,j}^\alpha$, Γ_j^α are the resonance energy and shape width, respectively, of the j - mode of the giant dipole excitation for SLO and SMLO models ($\alpha = SLO$ and $SMLO$). Factor s_j^α is the normalized contribution ("weight") of the Lorentzian component F_j^α of model α to the GDR component of the photoabsorption cross section in terms of the Thomas-Reiche-Kuhn (TRK) sum rule σ_{TRK}

$$\sigma_{TRK} = 60 \frac{NZ}{A} = 15A(1 - I^2) \text{ [mb} \cdot \text{MeV]}, \quad (6)$$

where $I = (N - Z)/A$ is the neutron-proton asymmetry factor.

The width Γ_j^{SLO} does not depend on the gamma-ray energy and coincides with GDR width $\Gamma_{r,j}^{SLO}$, while Γ_j^{SMLO} is given by a linear function of the gamma-ray energy

$$\Gamma_j^{SLO} = \Gamma_{r,j}^{SLO} = const, \quad \Gamma_j^{SMLO}(\varepsilon_\gamma) = a_j \varepsilon_\gamma. \quad (7)$$

In the SLO model, the quantities $E_{r,j}^{SLO}$, $\Gamma_j^{SLO} = \Gamma_{r,j}^{SLO}$ and s_j^{SLO} are used as variables in the fitting procedure. In the SMLO model, the parameters $E_{r,j}^{SMLO}$, a_j , s_j^{SMLO} are determined by fitting, and the widths $\Gamma_{r,j}^{SMLO}$ presented in Table 2 are recalculated from obtained a_j by the relationship $\Gamma_{r,j}^{SMLO} = \Gamma_j^{SMLO}(\varepsilon_\gamma = E_{r,j}^{SMLO}) = a_j E_{r,j}^{SMLO}$.

A resonance value $\sigma_{r,j}^\alpha = \sigma_{GDR,j}^\alpha(\varepsilon_\gamma = E_{r,j}^\alpha)$ of the cross section component corresponding to the giant dipole vibration along the j -axis at resonance energy $\varepsilon_\gamma = E_{r,j}^\alpha$ is equal to:

$$\sigma_{r,j}^\alpha = \frac{2}{\pi} \sigma_{TRK} s_j^\alpha / \Gamma_{r,j}^\alpha, \quad (8)$$

where $\Gamma_{r,j}^\alpha = \Gamma_j^\alpha(\varepsilon_\gamma = E_{r,j}^\alpha)$ is width of j -component at corresponding resonance energy.

If integral of function F_j^α over energy is equal to unity, $F_j^{\alpha,int} = \int_0^\infty F_j^\alpha(\varepsilon_\gamma) d\varepsilon_\gamma = 1$, the sum $s^\alpha = s_1^\alpha + s_2^\alpha$ of the weights in Eq.(5) is, by definition, the energy weighted sum-rule in units of the TRK sum rule σ_{TRK} , and s_j^α is the strength of the corresponding mode of the giant dipole excitation. The s^α should be not exceed a value $1 + \Delta$,

$$s^\alpha = \sum_j s_j^\alpha \leq 1 + \Delta, \quad \Delta \approx 0.2 \div 0.3, \quad (9)$$

where Δ is the contribution of the velocity-dependent and exchange forces [37]. Note that the GDR exhausts the EWSR by 84% only within the framework of the hydrodynamic approach to SLO model [33].

In the approximation of equally probable excitation of normal modes of the giant collective vibration, the weights $s_j = s_j^\alpha$ are equal to

$$\begin{cases} s_1 = s/3; & s_2 = 2s/3, \quad \beta > 0, \\ s_1 = 2s/3; & s_2 = s/3, \quad \beta < 0 \end{cases} \quad (10)$$

because of twofold degeneration of the giant collective vibration, which is perpendicular to the axis of symmetry; $s = s^\alpha$.

The β is a parameter of quadrupole deformation and the subindex $j = 1(2)$ in s_j corresponds to the low (high) value component of the GDR energy, respectively, i.e. $E_{r,j=1}^\alpha < E_{r,j=2}^\alpha$.

It should be mentioned that in SLO model $F_j^{SLO,int} = 1$ and the weights s_j are the strengths of the GDR excitation modes in units of the TRK sum rule. For the SMLO and other models with energy-increasing width, $F_j^{\alpha,int} \neq 1$ and the factors s_j can be interpreted to be the GDR strengths only approximately.

In this work a least-squares fitting procedure was employed, in which the data points were weighted according to the inverse square of their uncertainties and a minimum value was sought for a least-square deviation χ_{model}^2 :

$$\chi_{\text{model}}^2 = \frac{1}{N_f} \sum_{i=1}^N \frac{(\sigma_{\text{exp}}(\varepsilon_i) - \sigma_{\text{abs}}^{\text{model}}(\varepsilon_i))^2}{(\Delta\sigma(\varepsilon_i))^2}, \quad (11)$$

where $\sigma_{\text{abs}}^{\text{model}}(\varepsilon_i)$ is the value of the theoretical cross section (2)-(5) with $\sigma_{GDR}(\varepsilon_i)$ for model SLO or SMLO at gamma-ray energy ε_i , $\sigma_{\text{exp}}(\varepsilon_i)$ is the measured value for the total photoabsorption cross section with uncertainty $\Delta\sigma(\varepsilon_i)$ at that energy, and $N_f = N - N_{\text{par}}$ is the number of degrees of freedom for the data set fitted which is equal to the number N of data points within the fitting interval minus the number N_{par} of fitted parameters (3 parameters for each Lorentz-like curve).

If the EXFOR datafiles did not include experimental uncertainties, then they were taken into account according to Ref. [17]. Specifically, two different estimations of errors are adopted in absence of experimental uncertainties to check their impact on the determination of the GDR parameters: a 10% constant error and an energy-dependent error. The former type of error is very often encountered in photoabsorption data files in the EXFOR database, while the latter energy dependence was chosen to simulate the statistical error that is inversely proportional to the counting rate which

is maximum near the GDR. Hence, the energy-dependent relative uncertainties were assumed to take minimum values (10%) near the GDR peaks and maximum values (50%) on the GDR tails. For spherical nuclei, a triangular dependence on gamma-energy was assumed, while for deformed nuclei a trapezoidal dependence with the GDR peaks as the top corners of the trapezium.

For a given nuclide the data were fitted in an energy interval that was determined following the prescription of Refs. [14–19]. The mid-position was chosen near the peak of the photoabsorption cross section and the interval was placed within an energy range defined by the one-neutron and three-neutron separation energies, where the GDR strength is expected to be dominant. Adjustments to this interval were made in cases where there were too few experimental data points, in order to allow for a good fit to be obtained using a Lorentzian curve.

The minimization of the least-square deviation functional was performed by the CERN MINUIT package [38]. The standard deviation of the parameters was estimated using the MINOS procedure of this code. The calculated mode was defined by the following sequence of commands: SEEK 1000, MIGRAD 10000 0.000001, IMPROVE 100, HESSE 1, MINOS 1.

3. Discussion and Conclusion

It should be noted that the energies $E_{r,j}$, widths $\Gamma_{r,j}$ and factors s_j presented in Tables 1 and 2 are, in fact, the parameters of Lorentz-like curves (Lorentz parameters) representing the best fit of the experimental photoabsorption cross sections in the intervals that hold peak energies for the SLO and SMLO models, respectively.

Different studies mentioned in the previous section demonstrate that the Lorentzian parameters can be considered as the GDR parameters or mean values of these parameters if the total photoabsorption cross-section is used in fitting and GDR excitation is dominant photoabsorption mechanism.

The accuracy of the description of GDR parameters by Lorentz parameters of the photoabsorption cross section data depends on many factors. Besides the well-known problems of the selection and verification of the experimental data and estimating the contributions of emission cross sections of different particles to the total photoabsorption cross section, there are also ambiguities in the theoretical description of the GDR component of the E1 photoabsorption cross section. Namely, an unsolved problem with description of nuclear dissipation as well as the approximation of σ_{GDR} by one- or two-component Lorentz-like curves (5) near the beta-stability valley is, as a rule, appropriate for rather heavy ($A \gtrsim 40$) spherical and axially deformed nuclei ($155 \lesssim A \lesssim 190$ or $225 \lesssim A \lesssim 250$) for gamma-ray energies close to the GDR energy.

Furthermore, additional physical effects should also be taken into account for the correct description of the photoabsorption in nuclei. The nuclear responses of the low-lying states (two-phonon states, pygmy dipole resonance) to the E1 field have an impact on the photoabsorption at the low-energy wing of the GDR. The isospin splitting, the neutron excess and the triaxial deformation can affect the shape of the photoabsorption cross section near the GDR and lead to the splitting of the GDR. For medium-mass to heavy nuclei, these effects can be described effectively by an additional broadening of the shape-line. But in some situations, isospin splitting and triaxiality of the nuclear shape have to be considered explicitly in the formulas [28, 39–43].

From the tables 1,2 it can be seen that these uncertainties for GDR energy E_r do not exceed $\sim 10\%$. The values of width Γ_r and the factors s_j are much more sensitive to the shape of the theoretical curve and their systematic

uncertainties can be rather significant. The difference in the magnitude of the parameters obtained from fitting within SLO and SMLO models can be considered as the systematic uncertainty due to ambiguity of the theoretical description of complex mechanism of the GDR relaxation which as yet studied. However, the overall consistency between different sets of parameters for the same nucleus in each table are, as a rule, rather good, while the variation of these parameters can be attributed to systematic uncertainties affecting both measurements and the theoretical approach.

The set of the Lorentzian parameters in the first line (first entry) for each nuclide was determined using, as a rule, the closest experimental data to the total photoabsorption cross sections near GDR and corresponds to minimal least-square deviation in comparison with other entries. Therefore, the quantities in the first entries are the recommended GDR parameters and are used in all our calculations presented below.

The quality of the description of the experimental photonuclear cross sections by the different models is considered by means of the ratio of the least-square deviations (Eq.(11)) for models SMLO and SLO, $Rc_i = \chi_{smlo}^2 / \chi_{slo}^2$, in different intervals ($\Delta\varepsilon_i$). The lower index i indicates the comparison interval: $i = f$ stands for fitting intervals $\Delta\varepsilon_f = \varepsilon_{min} \leq \varepsilon_\gamma \leq \varepsilon_{max}$ from the Tables 1,2 ; $i = l$ for intervals of the GDR left tail $\Delta\varepsilon_l = \varepsilon_0 \leq \varepsilon_\gamma \leq E_{r,1}$ and $i = m$ for whole intervals with the experimental data $\Delta\varepsilon_m = \varepsilon_0 \leq \varepsilon_\gamma \leq 30$ MeV with the energy ε_0 of the first experimental data-point.

The factors Rc_i are indicated in the Graphs and are collected in the Table A. This table gives the ratios of the arithmetic means of least-square values of models SMLO to SLO $\langle \chi_{smlo}^2 \rangle / \langle \chi_{slo}^2 \rangle$ in different gamma-ray intervals and for different sets of nuclei. Here, the isotopes from the mass-ranges $150 \leq A \leq 190$ and $220 \leq A \leq 253$ are named as axially deformed nuclei. The parameters β of effective quadrupole deformation of nuclear surface were taken from file "deflib.dat" packed in "gamma-strength-analytic.tgz", section GAMMA on <http://www-nds.iaea.org/RIPL-2/>. Nuclei with $\beta < 0.1$ were assumed to be spherical.

Table A

The ratio of arithmetical means of least-square deviations for SMLO and SLO models $\langle \chi^2(SMLO) \rangle / \langle \chi^2(SLO) \rangle$

Intervals	Nuclei		
	Spherical	Spherical+Axially deformed	All
$\Delta\varepsilon_f$	0.96	1.11	1.09
$\Delta\varepsilon_l$	0.71	0.73	0.76
$\Delta\varepsilon_m$	0.70	0.69	0.74

The curves in the graphs demonstrate that the low-energy tails of the photoabsorption cross sections within SLO model are, as a rule, higher than the SMLO model and the experimental data. This means that excitations of low-energy states (like pygmy dipole resonance) can not be described as additional peaks to SLO model without re-adjusting the standard values of its GDR parameters that provided description of the GDR range of the photoabsorption cross sections.

As mentioned above, SMLO model is based on nuclear response with the GDR excitation and energy-dependent width $\Gamma(\varepsilon_\gamma)$ that is linearly increasing with gamma-ray energy. The expressions for energy-dependent widths whose energy dependence increases with energy are based on a low-energy approximation of nucleon-nucleon collision cross section in the nuclear medium [44, 45] and are not valid at high energies [46, 47]. Therefore SMLO model can not

be used without modifications of the width above high gamma-ray energies (25-30 MeV) where the QD component is dominant.

Acknowledgments

The authors are very thankful to V.Yu. Denisov, S. Goriely, P. Oblozinský, M.W. Herman , A.V. Ignatyuk, J. Kopecky, and V.V. Varlamov for valuable discussions and comments. We very much appreciate help provided by Katerina Solodovnik in preparing the datafiles. This work is partially supported by the International Atomic Energy Agency through a Coordinated Research Project on Updating the Photonuclear Data Library and generating a Reference Database for Photon Strength Functions (F41032).

References

- [1] P. Oblozinský, M. B. Chadwick, T. Fukahori, A.V. Ignatyuk, S. Kailas, J. Kopecky, G. Molnar, G. Reffo, Z. Su, M. Uhl, P. G. Young, O. Bersillon, E. Běták, R. Capote and V. M. Maslov, Handbook for Calculations of Nuclear Reaction Data: Reference Input Parameter Library), Tech. Rep. IAEA-TECDOC-1034, International Atomic Energy Agency, Vienna, Austria, 1998 (see <http://www-nds.iaea.org/ripl/>).
- [2] T. Belgia, O. Bersillon, R. Capote, T. Fukahori, Zhigang Ge, S. Goriely, M. Herman, A. V. Ignatyuk, S. Kailas, A. Koning, V. Plujko and P. Young, Handbook for calculations of nuclear reaction data, Reference Input Parameter Library- 2, Tech. Rep. IAEA-TECDOC-1506, International Atomic Energy Agency, Vienna, Austria, 2006(see <http://www-nds.iaea.org/RIPL-2/>).
- [3] R. Capote, M. Herman, P. Oblozinský, P. G. Young, S. Goriely, T. Belgia, A. V. Ignatyuk, A. J. Koning, S. Hilaire, V. A. Plujko, M. Avrigeanu, O. Bersillon, M. B. Chadwick, T. Fukahori, Zhigang Ge, Yinlu Han, S. Kailas, J. Kopecky, V. M. Maslov, G. Reffo, M. Sin, E. Sh. Soukhovitskii and P. Talou, Nucl. Data Sheets 110 (2009) 3107; (see <http://www-nds.iaea.org/RIPL-3/>).
- [4] R. Capote, A. Delgado, and A. Gonzalez, Mod. Phys. Lett. B15 (2001) 81.
- [5] J. L. Ullmann, T. Kawano, T. A. Bredeweg, A. Couture, R. C. Haight, M. Jandel, J. M. O'Donnell, R. S. Rundberg, D. J. Vieira, J. B. Wilhelmy, J. A. Becker, A. Chyzh, C. Y. Wu, B. Baramsai, G. E. Mitchell, and M. Krticka, Phys. Rev. C89 (2014) 034603.
- [6] M. R. Mumpower, T. Kawano, J. L. Ullmann, M. Krticka, and T. M. Sprouse, Phys. Rev. C96 (2017) 024612.
- [7] Review articles can be found in Correlations in Clusters and Related Systems, ed. J. P. Connerade (World Scientific, Singapore, 1996).
- [8] E. Lipparini, in Many Body Theory of Correlated Electron Systems, eds. M. I. Gallardo and M. Lozano (World Scientific, Singapore, 1998).
- [9] S. Pollach, C. R. C. Wang and M. M. Kappes, J. Chem. Phys. 94 (1991) 2496.
- [10] G. F. Bertsch and R. A. Broglia, Oscillations in Finite Quantum Systems (Cambridge University Press, Cambridge, 1994).

- [11] A. Delgado, L. Lavin, R. Capote and A. Gonzalez, *Physica E8* (2000) 345.
- [12] M. Herman, R. Capote, B. V. Carlson, P. Obložinský, M. Sin, A. Trkov, H. Wienke and V. Zerkin, *Nucl. Data Sheets* 108 (2007) 2655.
- [13] A. J. Koning, S. Hilaire and M. C. Duijvestijn, TALYS-1.0, Proceedings of the International Conference on Nuclear Data for Science and Technology, April 22-27, 2007, Nice, France, editors O.Bersillon, F.Gunsing, E.Bauge, R.Jacqmin, and S.Leray, EDP Sciences, 2008, p. 211-214 (see www.talys.eu).
- [14] B. L. Berman and S. C. Fultz, *Rev. Mod. Phys.* 47 (1975) 713.
- [15] B. L. Berman, R. G. Pywell, S. S. Dietrich, M. N. Thompson, K. O. McNeill and J. W. Jury, *Phys. Rev. C* 36 (1987) 1286.
- [16] S. S. Dietrich and B. L. Berman, *At. Data Nucl. Data Tables* 38 (1988) 199.
- [17] V. A. Plujko, R. Capote, O. M. Gorbachenko. *At. Data Nucl. Data Tables*, 97 (2011) 567.
- [18] M. B. Chadwick, P. Obložinský, A. I. Blokhin, T. Fukahori, Y. Han, Y. -O. Lee, M. N. Martins, S. F. Mughabghab, V. V. Varlamov, B. Yu and J. Zhang, Handbook on photonuclear data for applications: Cross sections and spectra , Tech. Rep. IAEA-TECDOC-1178, (International Atomic Energy Agency, Vienna, Austria, 2000) (see <http://www-nds.iaea.org/reports-new/tecdocs/iaea-tecdoc-1178.pdf>).
- [19] A. V. Varlamov, V. V. Varlamov, D. S. Rudenko and M. E. Stepanov, Atlas of Giant Dipole Resonances. Parameters and graphs of photonuclear reaction cross sections, Tech. Rep. INDC(NDS)-394, (International Atomic Energy Agency, Vienna, Austria, 1999) (see <http://www-nds.iaea.org/reports-new/indc-reports/indc-nds/indc-nds-0394.pdf>).
- [20] S. Goriely and E. Khan, *Nucl. Phys. A* 706 (2002) 217.
- [21] S. Goriely, E. Khan and M. Samyn, *Nucl. Phys. A* 739 (2004) 331.
- [22] A. Schiller and M. Thoennessen, *At. Data Nucl. Data Tables* 93 (2007) 549.
- [23] D. Kusnezov, Y. Alhassid, K. A. Snover and W. E. Ormand, *Nucl. Phys. A* 687 (2001) 12.
- [24] S.Goriely and P. Dimitriou (Prep.). Summary Report 1st RCM on CRP "Updating Photonuclear Data Library and Generating a Reference Database for Photon Strength Functions", INDC(NDS)-0712, Vienna, 2016.
- [25] International Network of Nuclear Reaction Data Centres: EXFOR/CSISRS database (see <http://www-nds.iaea.org/exfor/>).
- [26] V. V. Varlamov, B. S. Ishkhanov, V. N. Orlin, , K. A. Stopani, *European Phys. Jour. A*, 51 (2015) 67.
- [27] V. V. Varlamov, B. S. Ishkhanov, V. N. Orlin, N. N. Peskov, *Yad. Fiz.* 79 (4) (2016) 315.
- [28] D. G. Gardner, In: Neutron Radiative Capture. (Neutrons physics and nuclear data in science and technology; Vol.3)), Gen. Eds.: A. Michaudon, S. Cierjacks, R. E. Chrien, Pergamon Press, Oxford, New York, 1984; pp.62-118.
- [29] G. A. Bartholomew, E. D. Earle, A. J. Ferguson, J. W. Knowles, and M. A. Lone, *Advances Nucl. Phys.*, 7 (1973) 229.
- [30] D. M. Brink, Ph. D. Thesis, Oxford University, 1955.

- [31] M. Danos, W. Greiner, Phys. Lett., 8 (1964) 113.
- [32] M. Danos, W. Greiner, Phys. Rev., 138B (1965) 876.
- [33] J. M. Eisenberg, W. M. Greiner, Nuclear Theory. Volume I: Nuclear Models, 3rd Ed. (North Holland, Amsterdam 1987).
- [34] C. B. Dover, R. H. Lemmer, F. J. W. Hahne, Ann. Phys. 70 (1972) 458.
- [35] V. A. Plujko, I. M. Kadenko, O. M. Gorbachenko and E. V. Kulich, Int. J. Mod. Phys. E17 (2008) 240.
- [36] M. B. Chadwick, P. Obložinský, P. E. Hodgson and G. Reffo, Phys. Rev. C44 (1991) 814.
- [37] E. Lipparini and S. Stringari, Phys. Rep. 175 (1989) 103.
- [38] CERN Program Library, MINUIT (D506), Function Minimization and Error Analysis; code available online at <http://wwwasdoc.web.cern.ch/wwwasdoc/cernlib.html>; user manual available online at <http://wwwasdoc.web.cern.ch/wwwasdoc/minuit/minmain.html>.
- [39] T. J. Bowles, R. J. Holt, H. E. Jackson, R. M. Laszewski, R. D. McKeon, A. M. Nathan, J. R. Specht, Phys. Rev. C24 (1981) 1940.
- [40] Y. Alhassid, B. Busch, S. Levit, Phys. Rev. Lett., 61 (1988) 1926.
- [41] Y. Alhassid, B. Busch, Nucl. Phys. A509 (1990) 461.
- [42] A. R. Junghans, G. Rusev, R. Schwengner, A. Wagner and E. Grosse, Phys. Lett. B670 (2008) 200.
- [43] E. Grosse, A. R. Junghans, R. Massarczyk, Physics Letters B 739 (2014) 425.
- [44] V. M. Kolomietz, V. A. Plujko and S. Shlomo, Phys. Rev. C54 (1996) 3014.
- [45] V. A. Plujko, O. M. Gorbachenko and M. O. Kavatsyuk, Acta Phys. Slov. 51 (2001) 231.
- [46] C. Kalbach, Phys. Rev., C33 (1986) 818.
- [47] A. J. Koning, M. C. Duijvestijn; Nucl. Phys. A 744 (2004) 15.

Explanation of Tables

Table 1. Experimental values and uncertainties of GDR parameters within the standard Lorentzian (SLO) approach

For experimental data in the fits are used the following cross sections: the total photoabsorption reaction cross section $\sigma(\gamma, abs)$, total photoneutron cross section $\sigma(\gamma, sn)$ and inclusive photoneutron yield cross section $\sigma(\gamma, xn)$ which includes the multiplicity of neutrons emitted in each reaction event:

$$\begin{aligned}\sigma(\gamma, abs) &= \sigma(\gamma, sn) + \sigma(\gamma, cp), \\ \sigma(\gamma, sn) &= \sigma(\gamma, n) + \sigma(\gamma, np) + \sigma(\gamma, 2n) + \sigma(\gamma, 2np) + \sigma(\gamma, 3n) + \dots + \sigma(\gamma, F), \\ \sigma(\gamma, cp) &= \sigma(\gamma, p) + \sigma(\gamma, 2p) + \dots + \sigma(\gamma, d) + \sigma(\gamma, dp) + \dots + \sigma(\gamma, \alpha) + \dots, \\ \sigma(\gamma, xn) &= \sigma(\gamma, 1nx) + 2\sigma(\gamma, 2nx) + 3\sigma(\gamma, 3nx) + \dots + \bar{\nu}\sigma(\gamma, F),\end{aligned}$$

where $\sigma(\gamma, cp)$ is the photo-charged-particle reaction cross section, $\bar{\nu}$ is the average multiplicity of photofission neutrons.

When measured and evaluated data on the total photoabsorption cross section for a given nuclide were absent in the database, the total photoneutron cross section $\sigma(\gamma, sn)$ was taken instead of the photoabsorption cross section $\sigma(\gamma, abs)$. Such an approximation is valid if the contribution of the photo-charged-particle reaction cross sections is small. In the absence of experimental EXFOR data for the $\sigma(\gamma, sn)$, the total photoneutron cross section was evaluated as a combination of the available experimental cross sections for the inclusive photoneutron yield cross section $\sigma(\gamma, xn)$ and photoneutron cross sections with ejection of more than one neutron. It is denoted as $\sigma_{Id}(\gamma, sn)$ with "Id" for type of experimental data used in fitting (see below). For experimental data without uncertainties in the EXFOR base, the relative uncertainties were used either with constant value of 10% or as the energy-dependent quantities with minimal values (10%) near the GDR peaks and maximal values (50%) on the GDR tails as described in Ref.[17] and Sect.2. Throughout the table the italicized numbers refer to the uncertainties in the last digits of the quoted values.

The following designations are used in the table.

Nucl	The target studied (symbol); <i>nat</i> supra-index indicates a natural isotopic composition
Id	Type of experimental data used in fitting:
	0 - experimental $\sigma(\gamma, abs)$ with experimental uncertainties;
	1a - experimental $\sigma(\gamma, abs)$ with constant uncertainties (10 %);
	1b - experimental $\sigma(\gamma, abs)$ with energy-dependent uncertainties;
	2 - evaluated $\sigma(\gamma, abs)$ with experimental uncertainties;
	3a - evaluated $\sigma(\gamma, abs)$ with constant uncertainties (10 %);
	3b - evaluated $\sigma(\gamma, abs)$ with energy-dependent uncertainties;
	4 - experimental $\sigma(\gamma, sn)$ with experimental uncertainties;
	4a - experimental $\sigma(\gamma, sn)$ with constant uncertainties (10 %);
	4b - experimental $\sigma(\gamma, sn)$ with energy-dependent uncertainties;
	5 - evaluated $\sigma(\gamma, sn)$ using experimental data with experimental uncertainties;
	6 - composed $\sigma(\gamma, sn)$ as a combination of selected experimental cross sections: $\sigma_6(\gamma, sn) = (\sigma(\gamma, xn) + \sigma(\gamma, 1n))/2$ with absolute uncertainties: $\Delta\sigma_6(\gamma, sn) = \sqrt{\Delta\sigma^2(\gamma, xn) + \Delta\sigma^2(\gamma, 1n)}/2$;
	7 - composed $\sigma(\gamma, sn)$ as a combination of the experimental cross sections: $\sigma_7(\gamma, sn) = \sigma(\gamma, xn) - \sigma(\gamma, 2n)$ with absolute uncertainties: $\Delta\sigma_7(\gamma, sn) = \sqrt{\Delta\sigma^2(\gamma, xn) + \Delta\sigma^2(\gamma, 2n)}$;
	8 - composed $\sigma(\gamma, sn)$ as a combination of selected experimental cross sections: $\sigma_8(\gamma, sn) = \sigma(\gamma, 1n) + \sigma(\gamma, 2n)$ with absolute uncertainties: $\Delta\sigma_8(\gamma, sn) = \sqrt{\Delta\sigma^2(\gamma, 1n) + \Delta\sigma^2(\gamma, 2n)}$;
	8a - experimental $\sigma_8(\gamma, sn)$ with constant uncertainties (10 %);
	8b - experimental $\sigma_8(\gamma, sn)$ with energy-dependent uncertainties;
	9 - composed $\sigma(\gamma, sn)$ as a combination of selected experimental cross sections: $\sigma_9(\gamma, sn) = \sigma(\gamma, 1n) + \sigma(\gamma, 2n) + \sigma(\gamma, 3n)$ with absolute uncertainties : $\Delta\sigma_9(\gamma, sn) = \sqrt{\Delta\sigma^2(\gamma, 1n) + \Delta\sigma^2(\gamma, 2n) + \Delta\sigma^2(\gamma, 3n)}$;
	10 - composed $\sigma(\gamma, sn)$ as a combination of selected experimental cross sections: $\sigma_{10}(\gamma, sn) = \sigma(\gamma, 1n) + \sigma(\gamma, 2n) + \sigma(\gamma, f)$ with absolute uncertainties: $\Delta\sigma_{10}(\gamma, sn) = \sqrt{\Delta\sigma^2(\gamma, 1n) + \Delta\sigma^2(\gamma, 2n) + \Delta\sigma^2(\gamma, f)}$;
	11 - composed $\sigma(\gamma, sn)$ as a combination of selected experimental cross sections: $\sigma_{11}(\gamma, sn) = (\sigma(\gamma, xn) + \sigma(\gamma, 1n) + \sigma(\gamma, f))/2$ with absolute uncertainties: $\Delta\sigma_{11}(\gamma, sn) = \sqrt{\Delta\sigma^2(\gamma, xn) + \Delta\sigma^2(\gamma, 1n) + \Delta\sigma^2(\gamma, f)}/2$;
	12 - experimental $\sigma(\gamma, 1n)$ with experimental uncertainties.
	13 - experimental $\sigma(\gamma, xn)$ with experimental uncertainties;

Table 1 (continued)

$E_{r,j}, \Gamma_{r,j}, s_j$	parameters of energy, width and weight of Lorentz curves fitted to the corresponding photoabsorption cross sections within the indicated fitting interval. Notation 'spherical' ($j=1$) implies that a one component Lorentz curve gives a better fit to the data than a two-component one, and 'axially deformed' ($j=1,2$) implies the opposite; the values of TRK sum rule for ^{208}Pb are used for $^{\text{nat}}\text{Pb}$.
$E_{r,1}$	energy of the first component of the GDR with uncertainty (one-sigma standard deviation), MeV.
$\Gamma_{r,1}$	width of the first component of the GDR with uncertainty (one-sigma standard deviation), MeV.
s_1	weight of the first component of the GDR with uncertainty (one-sigma standard deviation).
$E_{r,2}$	energy of the second component of the GDR with uncertainty (one-sigma standard deviation), MeV.
$\Gamma_{r,2}$	width of the second component of the GDR with uncertainty (one-sigma standard deviation), MeV.
s_2	weight of the second component of the GDR with uncertainty (one-sigma standard deviation).
s	sum of weights of the first and second component of the GDR ($s = s_1 + s_2$) with uncertainty (one-sigma standard deviation).
$\varepsilon_{\min} (\varepsilon_{\max})$	lower (upper) energy limit of fitting interval, MeV.
Ref	Short references on the experimental data used in the fit.

Table 2. Experimental values and uncertainties of GDR parameters within the modified Lorentzian (SMLO) approach

Same as for Table 1.

Table 3. References to experimental and evaluated cross section data taken from EXFOR

Nucl	The target studied (symbol); <i>nat</i> supra-index indicates a natural isotopic composition.
Id	Type of experimental data used in fitting (without a letter indicating the method of uncertainty estimation employed; with the letters a, b : Ida - calculations with constant uncertainties (10 %) and Idb - calculations with energy-dependent uncertainties). See the explanation for Table 1.
Reaction	Type of reaction.
Ref	Short references on the experimental data.
EXFOR	EXFOR 8-digit entry & subentry number.

Explanation of Graphs

In the graphs presented comparison of experimental photonuclear cross sections near the GDR region from EXFOR database and theoretical calculations for the nuclei from ${}^6\text{Li}$ to ${}^{239}\text{Pu}$.

For experimental data, the photoabsorption cross section $\sigma(\gamma, abs)$ and total photoneutron cross section $\sigma(\gamma, sn)$ were used as it is described in the section “Explanation of Tables”. Short reference for experimental data is indicated in the graphs in accordance with the section “References used in the Tables”. The type of experimental data (ID index) for given nuclide corresponds to that in the first line of the Tables 1,2. These markers allow to locate entry and subentry numbers of EXFOR datafile by the use of information in the Table 3.

The theoretical calculations were performed using Eqs. (2)-(5) with allowance for components of the GDR excitation within SLO and SMLO models and contribution of quasi-deuteron photodisintegration. The theoretical curves are shown for more extended range of gamma-ray energies from 5 to 30 MeV when compared to fitting intervals.

The following designations are used in the graphs.

Title descriptor above the figure.

Notations of the photonuclear reaction type and irradiated target nucleus.

Coordinate axes.

Incident gamma-ray energy in units of MeV is indicated as an abscissa; the cross section in units of mb is indicated as an ordinate.

Vertical short lines.

Two lines normally to the x - coordinate show the energy interval ε_γ from ε_{\min} to ε_{\max} where fitting procedure was performed.

Rc_f, Rc_l, Rc_m .

The $Rc_i = \chi_{\text{smlo}}^2 / \chi_{\text{slo}}^2$ is ratio of the least-square deviations (Eq.(11)) in interval $(\Delta\varepsilon_i)$ indexed by i : $i = f$ stands for fitting intervals $\Delta\varepsilon_f = \varepsilon_{\min} \leq \varepsilon_\gamma \leq \varepsilon_{\max}$ from the Tables 1,2 ; $i = l$ for intervals of the GDR left tail $\Delta\varepsilon_l = \varepsilon_0 \leq \varepsilon_\gamma \leq E_{r,1}$ and $i = m$ for whole intervals with the experimental data $\Delta\varepsilon_m = \varepsilon_0 \leq \varepsilon_\gamma \leq 30$ MeV with the energy ε_0 of the first experimental data-point.

References used in the Tables

- [1958Ful] E. G. Fuller, M. S. Weiss, Phys. Rev. 112 (1958) 560.
- [1962Bog] O. V. Bogdankevich, B. I. Goryachev, V. A. Zapevalov, Zhur. Eksper. Teoret. Fiz. 42 (1962) 150 (Sov. Phys. JETP 15 (1962) 1044).
- [1962Fu1] S. C. Fultz, R. L. Bramblett, J. T. Caldwell, N. E. Hansen, C. P. Jupiter, Phys. Rev. 128 (1962) 2345.
- [1962Fu2] S. C. Fultz, R. L. Bramblett, J. T. Caldwell, N. A. Kerr, Phys. Rev. 127 (1962) 1273.
- [1963Br1] R. L. Bramblett, J. T. Caldwell, G. F. Auchampaugh, S. C. Fultz, Phys. Rev. 129 (1963) 2723.
- [1963Bur] N. A. Burgov, G. V. Danilyan, B. S. Dolbilkin, L. E. Lazareva, F. A. Nikolaev, Zhur. Eksper. Teoret. Fiz. 45,(6) (1963) 1694 (Sov. Phys. JETP 43(1) (1963) 50).
- [1964Baz] E. B. Bazhanov, A. P. Komar, A. V. Kulikov, Zhur. Eksper. Teoret. Fiz. 46 (1964) 1497 (Sov. Phys. JETP 19 (1964) 1014).
- [1964Br1] R. L. Bramblett, J. T. Caldwell, R. R. Harvey, S. C. Fultz, Phys. Rev. 133 (1964) B869.
- [1964Fu1] S. C. Fultz, R. L. Bramblett, J. T. Caldwell, R. R. Harvey, Phys. Rev. 133 (1964) B1149.
- [1964Ha2] R. R. Harvey, J. T. Caldwell, R. L. Bramblett, S. C. Fultz, Phys. Rev. 136 (1964) B126.
- [1964Ric] L. B. Rice, L. N. Bolen, W. D. Whitehead, Phys. Rev. 134 (1964) B557.
- [1965Bel] B. L. Berman, R. L. Bramblett, J. T. Caldwell, R. R. Harvey, S. C. Fultz, Phys. Rev. Lett. 15 (1965) 727.
- [1965Wyc] J. M. Wyckoff, B. Ziegler, H. W. Koch, R. Uhlig, Phys. Rev. 137 (1965) B576.
- [1966Axe] P. Axel, J. Miller, C. Schuhl, G. Tamas, C. Tzara, J. Phys. (Paris) 27 (1966) 262.
- [1966Br1] R. L. Bramblett, J. T. Caldwell, B. L. Berman, R. R. Harvey, S. C. Fultz, Phys. Rev. 148 (1966) 1198.
- [1966Dol] B. S. Dolbilkin, V. A. Zapevalov, V. I. Korin, L. E. Lazareva, F. A. Nikolaev, Izv. Akad. Nauk SSSR, Ser. Fiz 30(2) (1966) 349 (Bull. Acad. Sci. USSR, Phys. Ser. 30(2) (1966) 354).
- [1966Fu1] S. C. Fultz, J. T. Caldwell, B. L. Berman, R. L. Bramblett, R. R. Harvey, Phys. Rev. 143 (1966) 790.
- [1967Ant] G. P. Antropov, I. E. Mitrofanov, B. S. Russkikh Izv. Akad. Nauk SSSR, Ser. Fiz 31 (1967) 336 (Bull. Acad. Sci. USSR, Phys. Ser. 31 (1968) 320).
- [1967Be2] B. L. Berman, J. T. Caldwell, R. R. Harvey, M. A. Kelly, R. L. Bramblett, S. C. Fultz, Phys. Rev. 162 (1967) 1098.
- [1968Be5] R. Bergere, R. Beil, A. Veyssiere, Nucl. Phys. A121 (1968) 463.
- [1968Bez] N. Bezic, D. Jamnik, G. Kernel, J. Krajnik, J. Snajde, Nucl. Phys. A117 (1968) 124.
- [1968Dol] B. S. Dolbilkin, A. I. Isakov, V. I. Korin, L. E. Lazareva, F. A. Nikolaev, Yad. Fiz. 8 (1968) 1080. (Soviet J. Nucl. Phys. 8 (1969) 626.)
- [1968Su1] R. E. Sund, M. P. Baker, L. A. Kull, R. B. Walton, Phys. Rev. 176 (1968) 1366.
- [1968Tom] T. Tomimasu, J. Phys. Soc. Japan 25 (1968) 655.

- [1969Be1] B. L. Berman, R. L. Bramblett, J. T. Caldwell, H. S. Davis, M. A. Kelly, S. C. Fultz, Phys. Rev. 177 (1969) 1745.
- [1969Be6] R. Bergere, H. Beil, P. Carlos, A. Veyssiere, Nucl. Phys. A133 (1969) 417.
- [1969Be8] B. L. Berman, M. A. Kelly, R. L. Bramblett, J. T. Caldwell, H. S. Davis, S. C. Fultz, Phys. Rev. 185 (1969) 1576.
- [1969Bez] N. Bezic, D. Brajnik, D. Jamnik, G. Kernel, Nucl. Phys. A128 (1969) 426.
- [1969Dea] T. K. Deague, E. G. Muirhead, B. M. Spicer, Nucl. Phys. A139 (1969) 501.
- [1969Fu1] S. C. Fultz, B. L. Berman, J. T. Caldwell, R. L. Bramblett, M. A. Kelly, Phys. Rev. 186 (1969) 1255.
- [1969Gor] B. I. Goryachev, B. S. Ishkhanov, I. M. Kapitonov, I. M. Piskarev, V. G. Shevchenko, O. P. Shevchenko, Izv. Akad. Nauk SSSR, Ser. Fiz. 33 (1969) 1736 (Bull. Acad. Sci. USSR, Phys. Ser. 33 (1970) 1588).
- [1969Ish] B. S. Ishkhanov, I. M. Kapitonov, E. V. Lazutin, I. M. Piskarev, O. P. Shevchenko, ZhETF Pisma v Redaktsiyu 10 (1969) 30 (JETP Letters (USSR) 10 (1969) 51).
- [1969Vas] O. V. Vasilev, G. N. Zalesnyi, S. F. Semenko, V. A. Semenov, Yad. Fiz. 10 (1969) 460 (Soviet J. Nucl. Phys. 10 (1969) 263).
- [1970Ant] G. P. Antropov, I. E. Mitrofanov, A. I. Prokofev, V. S. Russkikh Izv. Akad. Nauk SSSR, Ser. Fiz 34 (1970) 116 (Bull. Acad. Sci. USSR, Phys. Ser. 34 (1970) 108).
- [1970Be8] B. L. Berman, S. C. Fultz, J. T. Caldwell, M. A. Kelly, S. S. Dietrich, Phys. Rev. C2 (1970) 2318.
- [1970Gor] B. I. Goryachev, B. S. Ishkhanov, I. M. Kapitonov, I. M. Piskarev, V. G. Shevchenko, O. P. Shevchenko; Yad. Fiz. 11 (1970) 252 (Sov. J. Nucl. Phys. 11 (1970) 141).
- [1970Is1] B. S. Ishkhanov, I. M. Kapitonov, E. V. Lazutin, I. M. Piskarev, O. P. Shevchenko, Izv. Akad. Nauk SSSR, Ser. Fiz. 34 (1970) 2228 (Bull. Acad. Sci. USSR, Phys. Ser. 34 (1971) 1988).
- [1970Is2] B. S. Ishkhanov, I. M. Kapitonov, E. V. Lazutin, I. M. Piskarev, O. P. Shevchenko, Vestnik Moskovskogo Univ., Ser. Fiz. Astron. 6 (1970) 606.
- [1970Is3] B. S. Ishkhanov, I. M. Kapitonov, E. V. Lazutin, I. M. Piskarev, O. P. Shevchenko, Yad. Fiz. 11 (1970) 606 (Sov. J. Nucl. Phys. 11 (1970) 394).
- [1970Su1] R. E. Sund, V. V. Verbinski, H. Weber, L. A. Kull, Phys. Rev. C2 (1970) 1129.
- [1970Ve1] A. Veyssiere, H. Beil, R. Bergere, P. Carlos, A. Lepretre, Nucl. Phys. A159 (1970) 561.
- [1971Alv] R. A. Alvarez, B. L. Berman, D. R. Lasher, T. W. Phillips, S. C. Fultz, Phys. Rev. C4 (1971) 1673.
- [1971Be4] H. Beil, R. Bergere, P. Carlos, A. Lepretre, A. Veyssiere, A. Parlag, Nucl. Phys. A172 (1971) 426.
- [1971Ca1] P. Carlos, H. Beil, R. Bergere, A. Lepretre, A. Veyssiere, Nucl. Phys. A172 (1971) 437.
- [1971Fu2] S. C. Fultz, R. A. Alvarez, B. L. Berman, M. A. Kelly, D. R. Lasher, T. W. Phillips, J. McElhinney, Phys. Rev. C4 (1971) 149.
- [1971Le1] A. Lepretre, H. Beil, R. Bergere, P. Carlos, A. Veyssiere, M. Sugawara, Nucl. Phys. A175 (1971) 609.

- [1971Vas] O. V. Vasilev, V. A. Semenov, S. F. Semenko, *Yad. Fiz.* 13 (1971) 463 (*Sov. J. Nucl. Phys.* 13 (1971) 259).
- [1972Ahr] J. Ahrens, H. Borchert, H. B. Eppler, H. Gimm, H. Gundrum, *Conf. Nucl. Structure Studies*, Sendai, Japan, (1972) 213.
- [1972Ask] H. J. Askin, R. S. Hicks, K. J. F. Allen, R. J. Petty, M. N. Thompson *Nucl. Phys.* A204 (1973) 209.
- [1972De1] T. K. Deague, R. J. Stewart, *Nucl. Phys.* A91 (1972) 305.
- [1972You] L. M. Young, Ph.D. Thesis, University of Illinois(1972), unpublished.
- [1973Bra] R. L. Bramblett, B. L. Berman, M. A. Kelly, J. T. Caldwell, S. C. Fultz, *Int. Conf. on Photonucle. Reactions*, Pacific Grove 1 (1973) 175.
- [1973Gor] A. M. Goryachev, G. N. Zalesnyi, S. F. Semenko, B. A. Tulupov, *Yad. Fiz.* 17 (1973) 463 (*Soviet J. Nucl. Phys.* 17 (1974) 236.).
- [1973Sor] Y. I. Sorokin, V. A. Khrushchev, B. A. Yurev *Izv. Akad. Nauk SSSR, Ser. Fiz.* 37(9) (1973) 1890 (*Bull. Acad. Sci. USSR, Phys. Ser.* 37(9) (1974) 80).
- [1973Ve1] A. Veyssiere, H. Reil, R. Bergere, P. Carlos, A. Lepretre, K. Kernbach, *Nucl. Phys.* A199 (1973) 45.
- [1974Be3] H. Beil, R. Bergere, P. Carlos, A. Lepretre, A. De Miniac, A. Veyssiere, *Nucl. Phys.* A227 (1974) 427.
- [1974Ca5] P. Carlos, H. Beil, R. Bergere, A. Lepretre, A. De Miniac, A. Veyssiere, *Nucl. Phys.* A225 (1974) 171.
- [1974Car] R. Carchon, J. Devos, R. Van de Vyver, C. Van Deynse, H. Ferdinande, *Nucl. Phys.* A223 (1974) 416.
- [1974Fu3] S. C. Fultz, R. A. Alvarez, B. L. Berman, P. Meyer, *Phys. Rev.* C10 (1974) 608.
- [1974Le1] A. Lepretre, H. Beil, R. Bergere, P. Carlos, A. De Miniac, A. Veyssiere, K. Kernbach, *Nucl. Phys.* A219 (1974) 39.
- [1974Sor] Y. I. Sorokin, B. A. Yurev, *Yad. Fiz.* 20 (1974) 233 (*Soviet J. Nucl. Phys.* 20 (1975) 123).
- [1974Ve1] A. Veyssiere, H. Beil, R. Bergere, P. Carlos, A. Lepretre, A. De Miniac, *Nucl. Phys.* A227 (1974) 513.
- [1975Ahr] J. Ahrens, H. Borchert, K. H. Czock, H. B. Eppler, H. Gimm, H. Gundrum, M. Kroning, P. Riehn, G. Sita Ram, A. Zieger, B. Ziegler, *Nucl. Phys.* A251 (1975) 479.
- [1975Mcc] J. J. McCarthy, R. C. Morrison, H. J. Vander-Molen, *Phys. Rev.* C11 (1975) 772.
- [1975Sor] Y. I. Sorokin, B. A. Yurev *Izv. Akad. Nauk SSSR, Ser. Fiz.* 39 (1975) 114 (*Bull. Acad. Sci. USSR, Phys. Ser.* 39(1) (1975) 98).
- [1975Vey] A. Veyssiere, H. Beil, R. Bergere, P. Carlos, A. Lepretre, A. De Miniac, *Jour. de Phys. Lett.* 36 (1975) 267.
- [1976Ca1] P. Carlos, H. Beil, R. Bergere, J. Fagot, A. Lepretre, A. Veyssiere, G. V. Solodukhov, *Nucl. Phys.* A258 (1976) 365.
- [1976Gor] B. I. Goryachev, Y. V. Kuznetsov, V. N. Orlin, N. A. Pozhidaeva, V. G. Shevchenko, *Yad. Fiz.* 23 (1976) 1145.
- [1976Gu1] G. M. Gurevich, L. E. Lazareva, V. M. Mazur, G. V. Solodukhov, B. A. Tulupov, *Nucl. Phys.* A273 (1976) 326.

- [1976Gu2] G. M. Gurevich, L. E. Lazareva, V. M. Mazur, G. V. Solodukhov, *JETP Lett.*(USSR) 23 (1976) 370.
- [1976Le2] A. Lepretre, H. Beil, R. Bergere, P. Carlos, J. Fagot, A. De Miniac, A. Veysiere, H. Miyase, *Nucl. Phys.* A258 (1976) 350.
- [1977Gor] A. M. Goryachev, G. N. Zalesnyi, *Yad. Fiz.* 26 (1977) 465 (*Sov. J. Nucl. Phys.* 26 (1977) 246).
- [1978Go1] A. M. Goryachev, G. N. Zalesnyi, *JETP Letters.* 26 (1978) 99 (*Zhur. Eksp. Teor. Fiz., Pisma v Redakt.* 26 (1978) 107).
- [1978Go2] A. M. Goryachev, G. N. Zalesnyi, *Yad. Fiz.* 27 (1978) 1479 (*Sov. J. Nucl. Phys.* 27 (1978) 779).
- [1978Go3] A. M. Goryachev, G. N. Zalesnyi, *Izvestiya Akademii Nauk KazSSSR, Ser. Fiz. -Mat.* 6 (1978) 8.
- [1978Gur] G. M. Gurevich, L. E. Lazareva, V. M. Mazur, G. V. Solodukhov, *Prob. Yad. Fiz. Kosm. Luch.* 8 (1978) 106.
- [1978Nor] J. W. Norbury, M. N. Thompson, K. Shoda, H. Tsubota, *Australian Jour. of Phys.* 31 (1978) 471.
- [1979Al2] R. A. Alvarez, B. L. Berman, D. D. Faul, F. H. Lewis, Jr., P. Meyer, *Phys. Rev.* C20 (1979) 128.
- [1979Be4] B. L. Berman, D. D. Faul, R. A. Alvarez, P. Meyer, D. L. Olson, *Phys. Rev.* C19 (1979) 1205.
- [1980Ca1] J. T. Caldwell, E. J. Dowdy, B. L. Berman, R. A. Alvarez, P. Meyer, *Phys. Rev.* C21 (1980) 1215.
- [1981Gur] G. M. Gurevich, L. E. Lazareva, V. M. Mazur, S. Yu. Merkulov, G. V. Solodukhov, V. A. Tyutin, *Nucl. Phys.* A351 (1981) 257.
- [1981Ish] B. S. Ishkhanov, I. M. Kapitonov, V. I. Shvedunov, A. I. Gutii, A. M. Parlag, *Yad. Fiz.* 33 (1981) 581 (*Sov. J. Nucl. Phys.* 33 (1981) 303).
- [1985Ahr] J. Ahrens, *Nucl. Phys.* A446 (1985) 229.
- [1986Ass] Y. I. Assafiri, M. N. Thompson, *Nucl. Phys.* A460 (1986) 455.
- [1986Be2] B. L. Berman, J. T. Caldwell, E. J. Dowdy, S. S. Dietrich, P. Meyer, R. A. Alvarez, *Phys. Rev.* C34 (1986) 2201.
- [1986Var] V. V. Varlamov, V. V. Surgutanov, A. P. Chernyaev, N. G. Efimkin, *Book: Fotojad. Dannye - Photo-disint. of Li, Suppl., Moscow* (1986).
- [1987Ahs] M. H. Ahsan, S. A. Siddiqui, H. H. Thies, *Nucl. Phys.* A469 (1987) 381.
- [1987Ber] B. L. Berman, R. G. Pywell, S. S. Dietrich, M. N. Thompson, K. O. McNeill, J. W. Jury, *Phys. Rev.* C36 (1987) 1286.
- [1987OKe] G. J. O'Keefe, M. N. Thompson, Y. I. Assafiri, R. E. Pywel, *Nucl. Phys.* A469 (1987) 239.
- [1989Ras] R. P. Rassool, M. N. Thompson, *Phys. Rev.* C39 (1989) 1631; *Phys. Rev.* C40 (1989) 506.
- [1995Bel] S. N. Belyaev, O. V. Vasiliev, V. V. Voronov, A. A. Nechkin, V. Yu. Ponomarev, V. A. Semenov, *Yad. Fiz.* 58 (1995) 1940 (*Phys. Atomic Nuclei* 58 (1995) 1883).
- [1999Bel] S. N. Belyaev, V. A. Semenov, V. P. Sinichkin, *Workshop on Beam Dynamics and Optimiz, Saratov* (1997) 97.
- [2001Bel] S. N. Belyaev, V. P. Sinichkin, *Conf.: Workshop on Beam Dynamics and Optimiz, Saratov* (2001) 81.

- [2002Ish] B. S. Ishkhanov, I. M. Kapitonov, E. I. Lileeva, E. V. Shirokov, V. A. Erokhova, M. A. Elkin, A. V. Izotova, Moscow State Univ. Inst. of Nucl. Phys. Report No.2002 27/711 (2002).
- [2003Ero] V. A. Erokhova, M. A. Elkin, A. V. Izotova, B. S. Ishkhanov, I. M. Kapitonov, E. I. Lileeva, E. V. Shirokov, *Izv. Ros. Akad. Nauk, Ser. Fiz.* 67 (2003) 1479.
- [2003Rod] T. E. Rodrigues, J. D. T. Arruda-Neto, Z. Carvaheiro, J. Mesa, *Phys. Rev. C* 68 (2003) 68.
- [2003Var] V. V. Varlamov, M. E. Stepanov, V. V. Chesnokov, *Izv. Ros. Akad. Nauk, Ser. Fiz.* 67 (2003) 656 (*Bull. Rus. Acad. Sci. Phys.* 67 (2003) 724).
- [2006Var] V. V. Varlamov, B. S. Ishkhanov, I. V. Makarenko, V. N. Orlin, N. N. Peskov, Moscow State Univ. Inst. of Nucl. Phys. Reports, 9 (2006) 808.
- [2007Var] V. V. Varlamov, N. N. Peskov; Moscow State Univ. Inst. of Nucl. Phys. Reports, 8 (2007) 829.
- [2009Var] V. V. Varlamov, B. S. Ishkhanov, V. N. Orlin, V. A. Tchetvertkova, Moscow State Univ. Inst. of Nucl. Phys. Reports, 3 (2009) 847 In Russian., *Izv. Rossiiskoi Akademii Nauk, Ser. Fiz* 74 (2010) 875 (*Bull. Rus. Acad. Sci. Phys.* 74 (2010) 833).
- [2010Var] V. V. Varlamov, B. S. Ishkhanov, V. N. Orlin, V. A. Tchetvertkova, *Izv. Ros. Akad. Nauk, Ser. Fiz.* 74 (2010) 884 (*Bull. Rus. Acad. Sci. Phys.* 74 (2010) 842).
- [2011Var] V. V. Varlamov, B. S. Ishkhanov, V. N. Orlin, T. S. Polevich, M. E. Stepanov, Moscow State Univ. Inst. of Nucl. Phys. Reports, 5 (2011) 869 In Russian. ; *Yad. Fiz.* 75(11) (2012) 1414 (*Phys. At. Nucl.* 75(11) (2012) 1339), *Eur. Phys. J. A* 50 (2014) 114.
- [2012Var] V. V. Varlamov, V. N. Orlin, N. N. Peskov, T. S. Polevich; Moscow State Univ. Inst. of Nucl. Phys. Reports, 1 (2012) 879 In Russian., *Yad. Fiz.* 76(11) (2013) 1484 (*Phys. At. Nucl.* 76(11) (2013) 1403), *Yad. Fiz.* 77(12) (2014) 1563 (*Phys. At. Nucl.* 77(12) (2014) 1369).
- [2013Ish] B. S. Ishkhanov, V. N. Orlin, N. N. Peskov, M. E. Stepanov, V. V. Varlamov, Moscow State Univ. Inst. of Nucl. Phys. Reports, 1 (2013) 884 In Russian., *Izv. Akad. Nauk RAS, Ser. Fiz* 77 (2013) 433 (*Bull. Rus. Acad. Sci. Phys.* 77 (2013) 388).
- [2013Var] V. V. Varlamov, V. N. Orlin, N. N. Peskov, M. E. Stepanov; *Izv. Ros. Akad. Nauk, Ser. Fiz.* 77 (2013) 433 (*Bull. Rus. Acad. Sci. Phys.* 77 (2013) 388).
- [2014Var] V. V. Varlamov, M. A. Makarov, N. N. Peskov, M. E. Stepanov; *Izv. Ros. Akad. Nauk, Ser. Fiz.* 78 (2014) 599 (*Bull. Russ. Acad. Sci. Phys.* 78(5) (2014) 412).
- [2015Var] V. V. Varlamov, M. A. Makarov, N. N. Peskov, M. E. Stepanov, *Yad. Fiz.* 78 (2015) 678.
- [2016Va1] V. V. Varlamov, A. I. Davydov, M. A. Makarov, V. N. Orlin, N. N. Peskov, *Izv. Ros. Akad. Nauk, Ser. Fiz.* 80 (2016) 351 (*Bull. Rus. Acad. Sci. Phys.* 80 (2016) 317).
- [2016Va2] V. V. Varlamov, B. S. Ishkhanov, V. N. Orlin, N. N. Peskov, *Yad. Fiz.* 79 (2016) 315 (*Journ. Phys. Atom. Nuclei* 79(4) (2016) 501).

Table 1

Experimental values and uncertainties of GDR parameters within Standard Lorentzian (SLO) approach.

Nucl	Id	$E_{r,1}$ (MeV)	$\Gamma_{r,1}$ (MeV)	s_1	$E_{r,2}$ (MeV)	$\Gamma_{r,2}$ (MeV)	s_2	s	$\varepsilon_{min} - \varepsilon_{max}$ (MeV)	Ref							
⁶ Li	2	23.69	16	5.26	79	0.341	31	0.341	31	21.5 - 27.0	1986Var						
	4	12.18	5	11.58	1	0.334	150	0.334	4	5.6 - 24.7	1965Be1						
⁷ Li	0	18.59	21	16.28	116	0.907	51	0.907	51	13.2 - 25.6	1985Ahr						
	2a	16.39	11	16.03	42	0.858	22	0.858	22	13.5 - 25.0	1986Var						
	2b	16.40	11	16.03	42	0.858	22	0.858	22	13.5 - 25.0	1986Var						
	4a	16.34	7	10.42	24	0.236	4	0.236	4	10.1 - 24.7	1973Bra						
	4b	16.26	10	11.17	41	0.248	7	0.248	7	10.1 - 24.7	1973Bra						
⁹ Be	0	23.75	10	9.47	33	0.577	17	0.577	17	17.5 - 26.0	1975Ahr						
¹⁰ B	4a	21.72	10	9.08	16	0.441	8	0.441	8	8.5 - 24.9	1987Ahs						
	4b	22.54	32	10.81	63	0.518	26	0.518	26	8.5 - 24.9	1987Ahs						
¹² C	0	22.86	2	3.61	7	0.671	10	0.671	10	20.1 - 25.0	1969Bez						
	1	22.88	2	3.06	1	0.573	70	0.573	10	13.2 - 24.0	1975Ahr						
	3a	23.14	5	2.89	11	0.546	13	0.546	13	20.2 - 25.0	2002Ish						
	3b	23.09	7	2.95	15	0.549	21	0.549	21	20.2 - 25.0	2002Ish						
	4	22.79	7	3.62	28	0.494	27	0.494	27	14.0 - 24.9	1963Bur						
¹³ C	3a	24.60	13	8.43	41	0.868	22	0.868	22	14.5 - 29.0	2002Ish						
	3b	24.40	17	7.70	63	0.828	36	0.828	36	14.5 - 29.0	2002Ish						
¹⁴ C	3a	15.41	26	5.82	90	0.333	47	26.13	15	7.78	92	0.483	46	0.816	66	14.5 - 30.0	2002Ish
	3b	16.68	57	3.10	67	0.177	23	25.87	17	6.84	84	0.490	46	0.667	51	9.0 - 30.0	2002Ish
natC	0	23.12	2	4.19	6	0.705	7	0.705	7	19.5 - 25.6	1985Ahr						
	0	23.38	1	4.17	4	0.728	5	0.728	5	19.0 - 26.0	1975Ahr						
	0	23.14	2	4.03	7	0.678	8	0.678	8	19.1 - 25.8	1972Ahr						
	1a	23.61	11	2.56	17	0.442	13	0.442	13	19.1 - 26.0	1965Wyc						
	1b	24.04	20	3.40	36	0.480	27	0.480	27	19.1 - 26.0	1965Wyc						
¹⁴ N	0	23.05	3	6.95	13	1.193	16	1.193	16	18.2 - 28.0	1969Bez						
	3a	23.39	8	4.83	17	0.729	15	0.729	15	15.0 - 28.0	2002Ish						
	3b	23.13	18	4.83	35	0.725	47	0.725	47	15.0 - 28.0	2002Ish						
¹⁵ N	3a	24.78	26	12.82	63	1.242	56	1.242	56	14.5 - 28.0	2002Ish						
	3b	24.96	39	12.47	134	1.259	111	1.259	111	14.5 - 28.0	2002Ish						
¹⁶ O	0	23.70	4	5.36	12	0.981	16	0.981	16	18.1 - 26.0	1975Ahr						
	0	23.66	4	5.44	12	0.985	16	0.985	16	18.2 - 25.9	1972Ahr						
	0	23.37	4	5.54	14	0.957	19	0.957	19	18.5 - 26.0	1969Bez						
	0	22.92	5	4.73	17	1.021	28	1.021	28	18.9 - 25.9	1963Bur						
	1a	23.92	10	2.58	11	0.444	15	0.444	15	18.2 - 25.6	1965Wyc						
	1b	24.04	13	2.90	18	0.469	22	0.469	22	18.2 - 25.6	1965Wyc						
	3a	23.71	7	3.98	12	0.762	17	0.762	17	18.2 - 26.0	2002Ish						
	3b	23.89	11	4.59	23	0.820	29	0.820	29	18.2 - 26.0	2002Ish						
¹⁷ O	3a	23.40	10	5.48	31	0.739	24	0.739	24	18.5 - 26.5	2002Ish						
	3b	23.38	13	5.57	49	0.740	36	0.740	36	18.5 - 26.5	2002Ish						
¹⁸ O	3a	19.08	14	2.12	57	0.064	18	24.10	16	5.25	83	0.417	49	0.481	52	18.5 - 26.0	2002Ish
natO	0	23.60	4	5.82	12	1.031	15	1.031	15	18.9 - 27.9	1985Ahr						
	0	23.84	3	6.08	9	1.101	12	1.101	12	18.6 - 27.8	1975Ahr						
¹⁹ F	3a	21.61	36	12.57	7	1.152	580	1.152	67	10.0 - 24.0	2002Ish						
²³ Na	0	17.43	13	3.10	42	0.175	35	21.13	10	4.51	52	0.555	63	0.730	72	14.2 - 23.0	1981Ish
	1a	19.56	27	6.99	65	0.516	50	24.77	14	2.81	85	0.135	41	0.650	65	14.5 - 25.9	1965Wyc
	1b	19.76	30	7.55	90	0.549	63	24.80	13	2.45	77	0.112	37	0.661	73	14.5 - 25.9	1965Wyc
	3a	17.45	15	3.08	33	0.178	34	20.98	12	4.34	49	0.492	55	0.670	65	14.3 - 23.0	2002Ish
	3b	17.39	22	2.90	69	0.161	48	20.94	14	4.55	59	0.519	71	0.680	86	14.3 - 23.0	2002Ish
	4	17.88	9	4.59	21	0.151	11	24.34	20	9.22	55	0.368	25	0.518	27	12.6 - 25.7	1971Alv
²⁴ Mg	0	19.51	12	2.71	40	0.253	58	23.88	36	8.86	144	0.974	163	1.227	173	16.5 - 27.0	1966Dol
	3a	19.81	5	2.72	12	0.359	17	24.42	9	4.76	34	0.416	27	0.775	32	16.3 - 27.0	2003Var
	3b	19.79	6	2.98	18	0.383	24	24.54	9	4.85	44	0.415	36	0.798	43	16.3 - 27.0	2003Var
	3a	19.88	7	2.81	18	0.371	26	24.69	11	4.74	58	0.451	46	0.822	53	15.2 - 27.0	2002Ish
	3b	19.85	7	2.79	19	0.373	25	24.71	10	4.69	53	0.451	44	0.824	50	15.2 - 27.0	2002Ish
²⁵ Mg	3a	22.06	10	6.09	15	0.894	20	0.894	20	9.0 - 24.2	2002Ish						
	3b	22.09	16	6.30	30	0.902	34	0.902	34	9.0 - 24.2	2002Ish						
	4	23.26	22	9.56	44	0.947	52	0.947	52	14.1 - 23.9	1971Alv						
²⁶ Mg	3a	17.38	5	2.15	17	0.144	11	23.64	8	7.25	31	1.149	38	1.294	39	16.1 - 26.5	2003Var
	3b	17.37	12	1.68	29	0.113	17	23.58	8	7.75	37	1.222	45	1.335	48	16.1 - 26.5	2003Var
	3a	17.48	11	1.89	33	0.117	19	23.65	18	6.93	64	1.023	74	1.140	77	16.4 - 26.0	2002Ish
	3b	17.50	24	1.41	50	0.088	27	23.61	19	7.47	76	1.095	89	1.183	93	16.4 - 26.0	2002Ish
	4	17.42	3	2.53	11	0.164	7	22.86	12	5.05	29	0.428	27	0.592	27	16.0 - 25.7	1971Fu2
natMg	1a	22.55	21	7.97	37	0.726	26	0.726	26	15.1 - 26.6	1965Wyc						
	1b	20.08	16	4.04	36	0.499	31	0.499	31	15.1 - 26.6	1965Wyc						
²⁷ Al	0	20.73	8	7.45	26	1.128	30	1.128	30	14.0 - 24.1	1985Ahr						

Table 1 (continued)

Nucl	Id	$E_{r,1}$ (MeV)	$\Gamma_{r,1}$ (MeV)	s_1	$E_{r,2}$ (MeV)	$\Gamma_{r,2}$ (MeV)	s_2	s	$\varepsilon_{min} - \varepsilon_{max}$ (MeV)	Ref
	3b	17.10	4.34	1.227				1.227	14.4 - 19.0	2003Var
	4	16.82	4.11	0.756				0.756	14.0 - 18.9	1974Be3
	4a	16.55	5.03	0.944				0.944	14.5 - 19.0	1970Is3
	4b	16.51	4.99	0.950				0.950	14.5 - 19.0	1970Is3
^{94}Mo	4	16.53	5.12	1.113				1.113	9.6 - 18.9	1974Be3
^{96}Mo	4	16.11	5.64	1.155				1.155	13.2 - 17.0	1974Be3
^{98}Mo	4	15.79	5.90	1.211				1.211	13.2 - 18.9	1974Be3
	4a	15.54	5.32	1.300				1.300	13.1 - 19.0	1970Is3
	4b	15.53	5.44	1.346				1.346	13.1 - 19.0	1970Is3
^{100}Mo	9	15.72	7.68	1.404				1.404	12.1 - 20.0	1974Be3
^{103}Rh	3a	16.24	7.49	1.486				1.486	13.1 - 19.0	2003Var
	3b	16.23	7.60	1.500				1.500	13.1 - 19.0	2003Var
	4	16.14	7.22	1.414				1.414	13.2 - 18.9	1974Le1
	4	16.55	5.92	1.576				1.576	10.0 - 19.0	1962Bog
^{108}Pd	0	14.97	5.42	0.771	18.13	4.03	0.413	1.184	10.0 - 19.0	1969Dea
^{nat}Pd	4	15.89	6.30	1.279				1.279	10.2 - 17.8	1971Lep
^{107}Ag	4	15.83	6.49	1.193				1.193	9.5 - 19.0	1969Ish
	4	15.89	6.65	0.986				0.986	13.1 - 18.7	1969Be1
	4a	15.68	5.56	1.384				1.384	10.0 - 19.0	1962Bog
	4b	15.60	5.43	1.343				1.343	10.0 - 19.0	1962Bog
^{109}Ag	4	13.54	3.49	0.275	16.62	4.41	0.490	0.765	13.1 - 19.0	1969Ish
^{nat}Ag	4	16.06	7.33	1.441				1.441	13.2 - 18.9	1971Lep
^{nat}Cd	4	15.77	5.70	1.227				1.227	10.2 - 17.8	1971Lep
^{115}In	4	15.72	5.57	1.273				1.273	13.2 - 17.8	1974Le1
	4	15.63	5.22	1.285				1.285	13.1 - 17.8	1969Fu1
	4	15.39	4.67	1.374				1.374	10.0 - 18.0	1962Bog
	5	15.53	4.79	1.180				1.180	13.1 - 17.8	2013Var
^{112}Sn	4	15.62	5.01	1.246				1.246	10.9 - 18.0	1974Sor
	5	15.63	5.30	1.300				1.300	13.1 - 18.0	2009Var
^{114}Sn	4	15.82	6.08	1.378				1.378	13.1 - 18.0	1975Sor
	5	15.77	5.91	1.337				1.337	13.1 - 18.0	2009Var
^{116}Sn	4	15.55	5.06	1.254				1.254	13.1 - 17.9	1974Le1
	4	15.67	4.17	1.017				1.017	13.0 - 18.0	1969Fu1
	4a	15.73	5.59	1.299				1.299	13.1 - 18.0	1975Sor
	4b	15.73	5.61	1.304				1.304	13.1 - 18.0	1975Sor
	5	15.58	4.73	1.162				1.162	13.1 - 18.0	2009Var
^{117}Sn	4	15.64	5.02	1.181				1.181	13.2 - 17.8	1974Le1
	4	15.65	5.00	1.153				1.153	13.1 - 17.9	1969Fu1
	4a	15.60	5.16	1.216				1.216	13.1 - 18.0	1975Sor
	4b	15.63	5.27	1.242				1.242	13.1 - 18.0	1975Sor
	5	15.64	4.98	1.166				1.166	13.1 - 18.0	2009Var
^{118}Sn	4	15.43	4.84	1.221				1.221	13.1 - 17.9	1974Le1
	4	15.59	4.75	1.097				1.097	13.1 - 17.9	1969Fu1
	4a	15.37	6.02	1.478				1.478	13.1 - 18.0	1974Sor
	4b	15.38	6.08	1.491				1.491	13.1 - 18.0	1974Sor
	5	15.48	4.56	1.086				1.086	13.1 - 18.0	2009Var
^{119}Sn	4	15.53	4.78	1.085				1.085	13.0 - 17.9	1969Fu1
	4a	14.74	4.75	0.971				0.971	13.1 - 17.9	1975Sor
	4b	14.72	4.60	0.945				0.945	13.1 - 17.9	1975Sor
	5	15.58	5.23	1.190				1.190	13.1 - 18.0	2009Var
^{120}Sn	4	15.37	5.08	1.295				1.295	13.1 - 17.9	1974Le1
	4	15.40	4.86	1.219				1.219	13.1 - 17.9	1969Fu1
	4a	15.30	5.84	1.465				1.465	13.1 - 18.0	1974Sor
	4b	15.27	5.72	1.435				1.435	13.1 - 18.0	1974Sor
	5	15.34	4.79	1.206				1.206	13.1 - 18.0	2009Var
^{122}Sn	4a	15.34	4.73	1.120				1.120	13.1 - 18.0	1975Sor
	4b	15.35	4.74	1.125				1.125	13.1 - 18.0	1975Sor
	5	15.25	4.72	1.082				1.082	13.1 - 18.0	2009Var
^{124}Sn	3a	15.31	4.94	1.173				1.173	13.1 - 18.0	2003Var
	3b	15.29	4.86	1.162				1.162	13.1 - 18.0	2003Var
	4	15.27	4.77	1.150				1.150	13.2 - 17.8	1974Le1
	4	15.18	4.79	1.183				1.183	13.1 - 17.8	1969Fu1
	4a	15.19	5.45	1.254				1.254	13.1 - 18.0	1975Sor
	4b	15.19	5.43	1.253				1.253	13.1 - 18.0	1975Sor
	5	15.17	4.87	1.163				1.163	13.1 - 18.0	2009Var
^{nat}Sb	4	15.48	5.01	1.242				1.242	10.2 - 17.8	1971Lep
^{124}Te	6	15.23	5.50	1.337				1.337	12.0 - 18.9	1976Le2

Table 1 (continued)

Nucl	Id	$E_{r,1}$ (MeV)	$\Gamma_{r,1}$ (MeV)	s_1	$E_{r,2}$ (MeV)	$\Gamma_{r,2}$ (MeV)	s_2	s	$\varepsilon_{min} - \varepsilon_{max}$ (MeV)	Ref
^{126}Te	6	15.15 2	5.36 7	1.358 13				1.358 13	12.0 - 18.9	1976Le2
^{128}Te	6	15.12 2	5.30 8	1.367 14				1.367 14	12.0 - 18.9	1976Le2
^{130}Te	6	15.11 2	4.98 7	1.334 13				1.334 13	12.0 - 18.9	1976Le2
^{127}I	4	14.77 3	4.09 5	0.795 23	17.30 10	3.69 30	0.173 24	0.968 33	12.0 - 20.0	1999Bel
	4	13.91 9	1.10 63	0.052 38	15.20 10	4.73 10	1.088 49	1.140 62	12.2 - 20.0	1989Ras
	4	14.45 20	3.72 37	0.713 215	16.58 46	4.72 61	0.565 234	1.278 318	8.8 - 19.8	1969Be6
	4	14.25 15	3.28 39	0.340 122	16.30 30	5.15 34	0.632 142	0.972 187	12.1 - 19.9	1966Br1
	5	14.20 28	3.17 79	0.367 259	16.21 53	5.22 48	0.810 298	1.177 395	12.1 - 19.9	2006Var
	8	14.61 20	2.62 104	0.163 196	15.72 50	6.19 115	1.012 142	1.175 243	12.1 - 16.9	1987Ber
^{133}Cs	4	15.33 1	5.28 2	1.351 4				1.351 4	12.0 - 19.0	1974Le1
	4	15.24 2	4.97 8	1.155 12				1.155 12	12.1 - 18.7	1969Be1
	5	15.16 13	4.83 49	1.241 74				1.241 74	12.1 - 19.0	2016Va2
^{138}Ba	5	15.13 9	4.51 32	1.126 46				1.126 46	12.1 - 18.7	2016Va2
	4	15.25 1	4.58 5	1.176 9				1.176 9	12.1 - 18.7	1970Be8
$^{\text{nat}}\text{Ba}$	4	15.29 1	4.93 3	1.370 6				1.370 6	10.1 - 17.8	1971Be4
^{139}La	4	15.24 2	4.82 9	1.379 15				1.379 15	12.0 - 18.9	1972De1
	4	15.23 1	4.45 4	1.159 8				1.159 8	12.0 - 18.9	1971Be4
	4	15.13 2	4.45 8	1.253 17				1.253 17	12.1 - 19.0	1968Be5
	4a	14.43 7	3.60 20	0.783 23				0.783 23	12.5 - 19.0	1964Ric
	4b	14.45 12	3.38 35	0.767 52				0.767 52	12.5 - 19.0	1964Ric
^{140}Ce	6	15.03 1	4.39 4	1.292 9				1.292 9	12.0 - 18.9	1976Le2
^{142}Ce	6	14.85 2	5.08 8	1.284 15				1.284 15	12.0 - 18.9	1976Le2
^{141}Pr	4	15.19 1	4.23 8	1.106 18				1.106 18	12.1 - 16.9	1987Ber
	4	15.14 2	4.40 6	1.083 10				1.083 10	12.1 - 18.7	1966Br1
	4a	14.66 7	3.82 21	0.844 25				0.844 25	12.5 - 19.0	1964Ric
	4b	14.70 14	3.75 40	0.840 62				0.840 62	12.5 - 19.0	1964Ric
	12	15.35 2	4.05 4	1.021 8				1.021 8	12.0 - 18.1	1972You
	12	15.04 1	4.47 3	1.179 5				1.179 5	12.0 - 16.9	1971Be4
	12	15.23 1	3.98 4	1.031 7				1.031 7	12.1 - 19.0	1970Su1
	13	15.39 2	3.80 6	1.039 12				1.039 12	12.0 - 18.9	1972De1
^{142}Nd	3a	14.95 5	4.46 17	1.215 30				1.215 30	13.1 - 18.0	2003Var
	3b	14.94 12	4.48 47	1.218 106				1.218 106	13.1 - 18.0	2003Var
	4	14.94 1	4.41 3	1.195 7				1.195 7	12.0 - 18.9	1971Ca1
^{143}Nd	4	15.00 2	4.73 8	1.236 14				1.236 14	12.0 - 19.0	1971Ca1
^{144}Nd	4	15.04 2	5.25 7	1.239 12				1.239 12	12.0 - 18.9	1971Ca1
^{145}Nd	4	14.94 4	6.27 18	1.378 28				1.378 28	12.0 - 18.9	1971Ca1
^{146}Nd	4	14.73 2	5.74 10	1.314 17				1.314 17	12.0 - 18.9	1971Ca1
	4a	13.94 3	3.89 9	1.002 14				1.002 14	12.2 - 19.0	1971Vas
	4b	13.92 4	3.86 13	0.994 28				0.994 28	12.2 - 19.0	1971Vas
^{148}Nd	4	12.78 26	4.03 69	0.326 126	15.49 19	5.22 39	0.827 143	1.153 191	10.8 - 18.6	1971Ca1
	4	13.67 3	2.68 19	0.855 53	15.96 4	1.22 18	0.137 26	0.992 59	13.0 - 17.0	1969Vas
^{150}Nd	4	12.30 6	3.38 26	0.432 46	16.03 9	5.12 31	0.821 62	1.253 77	10.8 - 18.6	1971Ca1
^{144}Sm	4	15.31 2	4.42 7	1.251 13				1.251 13	12.1 - 18.9	1974Ca5
^{148}Sm	4	14.82 1	5.06 5	1.242 9				1.242 9	12.1 - 18.9	1974Ca5
	4a	14.01 2	3.80 1	0.919 50				0.919 11	7.7 - 19.0	1971Vas
	4b	13.99 3	3.76 1	0.907 60				0.907 15	7.7 - 19.0	1971Vas
^{150}Sm	4	14.59 3	5.92 11	1.324 18				1.324 18	12.1 - 18.9	1974Ca5
	4	13.92 3	4.49 10	1.160 19				1.160 19	12.2 - 19.0	1969Vas
^{152}Sm	4	12.39 3	2.99 12	0.377 21	15.73 4	5.15 13	0.853 28	1.230 35	10.9 - 18.8	1974Ca5
	4	11.69 2	2.15 7	0.703 22	14.99 3	2.50 10	0.747 25	1.449 33	11.0 - 18.8	1969Vas
^{154}Sm	0	12.17 20	2.80 95	0.360 187	15.63 49	5.89 151	0.873 286	1.233 342	10.9 - 18.6	1981Gur
	4	12.27 3	2.97 15	0.381 26	15.94 6	5.62 20	0.845 37	1.226 45	11.0 - 18.6	1974Ca5
	4	11.28 4	1.92 14	0.270 20	14.90 3	4.06 9	0.995 21	1.265 29	11.0 - 18.8	1969Vas
^{151}Eu	4a	13.88 5	4.69 14	0.850 23	14.45 7	0.75 34	0.035 19	0.885 30	10.2 - 18.0	1971Vas
	4b	13.93 6	4.71 13	0.873 28	14.44 6	0.62 32	0.026 15	0.899 32	10.2 - 18.0	1971Vas
^{153}Eu	4	12.33 4	2.77 17	0.305 27	15.78 7	5.76 19	0.892 40	1.197 48	10.9 - 18.7	1969Be8
	4a	11.94 2	2.02 10	0.246 12	15.23 4	3.31 13	0.588 17	0.834 21	11.0 - 18.8	1971Vas
	4b	11.96 3	2.09 10	0.262 14	15.16 6	2.96 17	0.531 29	0.792 32	11.0 - 18.8	1971Vas
^{152}Gd	4a	11.79 4	3.03 15	0.312 21	14.72 4	3.16 13	0.566 24	0.878 32	10.2 - 18.0	1971Vas
	4b	11.80 4	3.05 16	0.319 22	14.72 5	3.08 16	0.552 31	0.870 38	10.2 - 18.0	1971Vas
^{154}Gd	4a	11.97 4	2.65 15	0.304 19	15.05 4	3.35 14	0.575 25	0.879 32	10.2 - 18.0	1971Vas
	4b	11.99 4	2.71 15	0.318 21	15.02 5	3.14 18	0.538 33	0.856 39	10.2 - 18.0	1971Vas
^{156}Gd	0	12.46 19	3.14 74	0.501 162	15.79 32	4.56 100	0.683 201	1.184 258	10.9 - 18.7	1981Gur
	4a	11.83 4	3.57 23	0.485 31	15.06 4	2.31 12	0.402 22	0.887 38	11.0 - 18.8	1971Vas
	4b	11.81 5	3.90 35	0.526 50	15.04 6	2.26 21	0.379 42	0.904 65	11.0 - 18.8	1971Vas
^{158}Gd	4a	11.86 4	2.94 15	0.368 21	15.16 5	3.31 20	0.559 26	0.927 34	10.2 - 18.0	1971Vas
	4b	11.92 4	3.09 16	0.415 25	15.03 8	2.44 27	0.438 47	0.853 53	10.2 - 18.0	1971Vas

Table 1 (continued)

Nucl	Id	$E_{r,1}$ (MeV)	$\Gamma_{r,1}$ (MeV)	s_1	$E_{r,2}$ (MeV)	$\Gamma_{r,2}$ (MeV)	s_2	s	$\varepsilon_{min} - \varepsilon_{max}$ (MeV)	Ref
^{237}Np	4	10.99 <i>4</i>	2.20 <i>12</i>	0.316 <i>27</i>	14.08 <i>6</i>	4.66 <i>18</i>	1.157 <i>43</i>	1.473 <i>50</i>	9.4 - 17.8	1986Be2
	4	11.01 <i>15</i>	2.91 <i>63</i>	0.347 <i>108</i>	14.05 <i>18</i>	4.52 <i>58</i>	0.820 <i>137</i>	1.167 <i>174</i>	9.5 - 16.5	1973Ve1
	5	10.89 <i>10</i>	2.28 <i>31</i>	0.250 <i>55</i>	13.98 <i>16</i>	4.71 <i>43</i>	0.905 <i>86</i>	1.156 <i>102</i>	9.3 - 17.7	2007Var
	10	10.98 <i>5</i>	2.17 <i>16</i>	0.312 <i>38</i>	14.06 <i>9</i>	4.64 <i>27</i>	1.160 <i>65</i>	1.472 <i>75</i>	9.4 - 17.8	1986Be2
^{239}Pu	10	11.01 <i>25</i>	2.92 <i>89</i>	0.343 <i>170</i>	14.10 <i>32</i>	4.76 <i>112</i>	0.863 <i>252</i>	1.206 <i>304</i>	9.2 - 16.6	1973Ve1
	0	11.07 <i>28</i>	3.29 <i>100</i>	0.344 <i>185</i>	14.00 <i>38</i>	5.51 <i>74</i>	0.906 <i>227</i>	1.250 <i>293</i>	9.3 - 18.7	1976Gu1
	5	11.27 <i>10</i>	2.75 <i>27</i>	0.380 <i>80</i>	13.68 <i>19</i>	4.24 <i>46</i>	0.658 <i>107</i>	1.038 <i>134</i>	9.2 - 17.7	2007Var
	10	11.31 <i>10</i>	2.48 <i>21</i>	0.385 <i>76</i>	13.90 <i>22</i>	4.36 <i>43</i>	0.766 <i>105</i>	1.151 <i>130</i>	9.1 - 17.8	1986Be2

Table 2 (continued)

Nucl	Id	$E_{r,1}$ (MeV)	$\Gamma_{r,1}$ (MeV)	s_1	$E_{r,2}$ (MeV)	$\Gamma_{r,2}$ (MeV)	s_2	s	$\epsilon_{min} - \epsilon_{max}$ (MeV)	Ref					
	0	20.98	8	8.77	27			1.272	33	14.2 - 24.4	1975Ahr				
	0	21.08	8	8.19	29			1.189	34	14.2 - 24.4	1972Ahr				
	0	21.11	7	8.60	26			1.247	32	14.2 - 24.4	1975Ahr				
	3a	20.62	8	5.32	14			0.876	21	14.2 - 23.0	2002Ish				
	4	21.01	3	7.15	5			0.377	3	14.0 - 24.4	1966Fu1				
²⁸ Si	3a	19.88	16	2.91	22	21.84	20	3.39	54	0.464	128	0.882	165	16.7 - 23.0	2003Var
	3b	19.78	36	2.68	97	21.68	45	4.12	113	0.630	358	0.935	454	16.7 - 23.0	2003Var
	3a	18.13	3	0.55	9	20.61	6	3.38	13	0.655	20	0.714	22	16.8 - 23.0	2002Ish
	3b	18.85	30	1.73	49	20.85	20	3.65	33	0.595	91	0.729	117	16.8 - 23.0	2002Ish
²⁹ Si	3a	20.97	11	6.88	25					0.941	29	0.941	29	14.2 - 23.0	2002Ish
	3b	20.91	16	6.69	46					0.923	47	0.923	47	14.2 - 23.0	2002Ish
³⁰ Si	3a	21.32	21	9.16	48					0.916	48	0.916	48	14.2 - 23.0	2002Ish
	3b	21.25	29	8.88	82					0.899	75	0.899	75	14.2 - 23.0	2002Ish
natSi	0	20.45	3	4.85	10	25.24	19	2.10	59	0.063	17	0.992	24	16.4 - 25.8	1975Ahr
³² S	0	19.57	9	5.10	35					0.583	28	0.583	28	14.7 - 23.0	1968Dol
	3a	21.21	12	6.08	16					1.098	36	1.098	36	14.4 - 23.0	2002Ish
	3b	21.77	20	7.04	41					1.288	78	1.288	78	14.4 - 23.0	2002Ish
³⁴ S	0	21.66	76	13.43	148					1.963	299	1.963	299	12.0 - 25.0	1986Ass
	3a	22.32	22	17.94	75					2.333	98	2.333	98	14.1 - 25.1	2003Var
	3b	22.78	41	16.62	136					2.318	198	2.318	198	12.3 - 25.1	2003Var
	3a	21.23	16	8.94	29					1.337	43	1.337	43	11.0 - 25.0	2002Ish
	3b	21.95	36	10.44	107					1.551	144	1.551	144	11.0 - 25.0	2002Ish
natS	0	20.42	11	5.74	40					0.887	50	0.887	50	17.2 - 23.6	1965Wyc
⁴⁰ Ar	3a	19.91	21	11.50	51					1.648	62	1.648	62	10.5 - 25.0	2002Ish
	3b	20.54	59	13.98	160					1.857	204	1.857	204	10.5 - 25.0	2002Ish
natK	13	21.28	3	7.40	10					0.443	5	0.443	5	16.0 - 25.9	1974Ve1
⁴⁰ Ca	0	20.72	11	6.44	51					1.755	102	1.755	102	17.2 - 23.7	1966Dol
	3a	20.14	5	5.05	20					1.201	28	1.201	28	17.2 - 24.0	2003Ero
	3b	20.11	10	5.09	34					1.202	56	1.202	56	17.2 - 24.0	2003Ero
⁴² Ca	3a	20.53	16	9.75	58					1.678	83	1.678	83	15.2 - 23.0	2003Ero
	3b	20.59	24	8.95	95					1.622	151	1.622	151	15.2 - 23.0	2003Ero
⁴⁴ Ca	3a	20.08	19	12.45	102					1.858	110	1.858	110	15.5 - 26.0	2003Ero
	3b	20.38	47	12.76	154					1.892	207	1.892	207	12.5 - 26.0	2003Ero
⁴⁸ Ca	0	19.90	18	6.42	96					1.512	199	1.512	199	17.9 - 21.6	1987OKe
	3a	19.95	18	7.82	67					1.461	90	1.461	90	15.5 - 23.0	2003Ero
	3b	19.80	28	7.11	100					1.375	160	1.375	160	15.5 - 23.0	2003Ero
natCa	0	20.09	2	5.07	8					1.241	13	1.241	13	15.1 - 24.0	1975Ahr
	0	19.98	3	5.07	9					1.198	15	1.198	15	15.2 - 24.0	1972Ahr
	0	19.62	3	4.65	9					1.045	14	1.045	14	15.0 - 24.0	1968Bez
	1a	20.42	7	3.44	16					0.713	18	0.713	18	15.4 - 24.0	1965Wyc
	1b	20.25	12	3.36	24					0.701	34	0.701	34	15.4 - 24.0	1965Wyc
	2	20.02	4	5.04	13					1.171	22	1.171	22	9.4 - 24.0	1985Ahr
⁴⁶ Ti	3a	19.95	9	7.99	24					1.393	30	1.393	30	13.2 - 25.0	2002Ish
	3b	19.98	25	8.77	70					1.451	103	1.451	103	13.2 - 25.0	2002Ish
⁴⁸ Ti	3a	20.13	25	9.98	75					1.352	88	1.352	88	14.5 - 23.0	2002Ish
	3b	20.43	59	11.23	187					1.472	246	1.472	246	14.5 - 23.0	2002Ish
⁵¹ V	4	18.18	6	5.29	14	21.37	15	3.36	63	0.127	30	0.828	41	14.1 - 22.9	1962Fu1
	3a	17.67	16	3.08	135	22.71	57	16.27	236	2.443	452	2.634	476	15.2 - 25.0	2003Var
	3b	17.96	36	4.91	123	22.87	65	8.20	396	1.053	638	1.699	700	15.2 - 25.0	2003Var
	4a	17.29	4	2.59	6	19.90	2	1.04	4	0.138	5	0.442	8	11.0 - 24.5	1969Gor
	4b	15.58	122	4.69	963	18.40	23	3.67	43	0.549	44	0.597	48	11.0 - 24.5	1969Gor
⁵² Cr	3a	19.16	8	6.70	24					1.075	27	1.075	27	14.3 - 23.0	2002Ish
	3b	19.36	22	7.52	63					1.146	84	1.146	84	14.3 - 23.0	2002Ish
	4a	18.21	4	5.09	9					0.782	12	0.782	12	14.1 - 23.0	1969Gor
	4b	18.65	6	6.36	16					0.977	23	0.977	23	14.1 - 23.0	1969Gor
⁵⁵ Mn	4	16.43	8	2.91	47	20.13	23	11.28	60	1.107	79	1.229	87	14.0 - 23.0	1979Al2
	4	17.02	25	3.88	63	20.05	63	7.60	140	0.693	245	1.061	299	14.1 - 22.8	1974Car
	4a	16.47	10	5.16	18	19.26	8	3.75	31	0.223	33	0.778	49	10.1 - 23.0	1970Is1
	4b	16.45	11	5.14	18	19.26	9	3.90	40	0.235	39	0.782	56	10.1 - 23.0	1970Is1
⁵⁴ Fe	0	19.39	8	5.69	30					1.606	54	1.606	54	16.0 - 23.0	1978Nor
⁵⁹ Co	4	16.44	9	2.42	37	18.68	22	8.63	46	0.922	35	1.009	45	14.0 - 20.9	1979Al2
	4	15.50	16	3.53	49	18.56	21	5.32	80	0.525	120	0.901	151	14.1 - 22.8	1964Baz
	4a	16.64	9	4.68	14	19.43	10	5.96	29	0.466	45	0.882	59	12.1 - 24.0	1969Gor
	4b	16.63	9	4.67	14	19.43	10	5.99	29	0.469	44	0.883	58	12.1 - 24.0	1969Gor
⁵⁸ Ni	3a	18.87	6	6.16	17					0.958	19	0.958	19	14.1 - 22.0	2003Var
	3b	18.78	12	6.03	36					0.942	47	0.942	47	14.1 - 22.0	2003Var
	3a	19.14	13	7.22	38					1.078	44	1.078	44	14.4 - 22.0	2002Ish

Table 2 (continued)

Nucl	Id	$E_{r,1}$ (MeV)	$\Gamma_{r,1}$ (MeV)	s_1	$E_{r,2}$ (MeV)	$\Gamma_{r,2}$ (MeV)	s_2	s	$\varepsilon_{min} - \varepsilon_{max}$ (MeV)	Ref	
	3b	17.21	17	4.62	62	1.293	160	1.293	160	14.4 - 19.0	2003Var
	4	16.89	1	4.28	4	0.781	6	0.781	6	14.0 - 18.9	1974Be3
	4a	16.67	4	5.13	22	0.961	30	0.961	30	14.5 - 19.0	1970Is3
	4b	16.64	5	5.06	28	0.964	39	0.964	39	14.5 - 19.0	1970Is3
⁹⁴ Mo	4	16.73	2	6.04	6	1.268	12	1.268	12	9.6 - 18.9	1974Be3
⁹⁶ Mo	4	16.42	7	6.52	20	1.315	43	1.315	43	13.2 - 17.0	1974Be3
⁹⁸ Mo	4	15.96	4	6.17	18	1.257	28	1.257	28	13.2 - 18.9	1974Be3
	4a	15.65	5	5.45	23	1.317	37	1.317	37	13.1 - 19.0	1970Is3
	4b	15.67	9	5.65	42	1.385	82	1.385	82	13.1 - 19.0	1970Is3
¹⁰⁰ Mo	9	16.02	4	8.44	18	1.515	26	1.515	26	12.1 - 20.0	1974Be3
¹⁰³ Rh	3a	16.59	11	8.44	51	1.638	82	1.638	82	13.1 - 19.0	2003Var
	3b	16.62	25	8.56	107	1.658	193	1.658	193	13.1 - 19.0	2003Var
	4	16.47	5	8.02	21	1.543	35	1.543	35	13.2 - 18.9	1974Le1
	4	16.83	27	7.60	56	1.913	148	1.913	148	10.0 - 19.0	1962Bog
¹⁰⁸ Pd	0	16.15	35	8.26	48	1.390	160	1.465	176	10.0 - 19.0	1969Dea
^{nat} Pd	4	16.24	5	7.63	14	1.502	26	1.502	26	10.2 - 17.8	1971Lep
¹⁰⁷ Ag	4	16.05	5	7.51	17	1.338	26	1.338	26	9.5 - 19.0	1969Ish
	4	16.14	5	7.18	22	1.053	29	1.053	29	13.1 - 18.7	1969Be1
	4a	15.78	8	6.40	24	1.528	44	1.528	44	10.0 - 19.0	1962Bog
	4b	15.87	13	6.64	31	1.581	74	1.581	74	10.0 - 19.0	1962Bog
¹⁰⁹ Ag	4	13.74	22	3.81	154	0.337	167	0.760	209	13.1 - 19.0	1969Ish
^{nat} Ag	4	16.39	2	8.03	10	1.555	16	1.555	16	13.2 - 18.9	1971Lep
^{nat} Cd	4	16.03	2	6.69	6	1.400	10	1.400	10	10.2 - 17.8	1971Lep
¹¹⁵ In	4	15.91	2	6.00	7	1.353	13	1.353	13	13.2 - 17.8	1974Le1
	4	15.79	1	5.58	6	1.357	12	1.357	12	13.1 - 17.8	1969Fu1
	4	15.54	17	5.66	38	1.584	100	1.584	100	10.0 - 18.0	1962Bog
	5	15.63	7	5.00	31	1.221	53	1.221	53	13.1 - 17.8	2013Var
¹¹² Sn	4	15.82	6	5.87	18	1.411	37	1.411	37	10.9 - 18.0	1974Sor
	5	15.81	7	5.74	27	1.388	54	1.388	54	13.1 - 18.0	2009Var
¹¹⁴ Sn	4	16.08	6	6.74	24	1.500	48	1.500	48	13.1 - 18.0	1975Sor
	5	16.01	6	6.50	22	1.448	44	1.448	44	13.1 - 18.0	2009Var
¹¹⁶ Sn	4	15.69	1	5.29	6	1.302	12	1.302	12	13.1 - 17.9	1974Le1
	4	15.74	2	4.34	8	1.049	13	1.049	13	13.0 - 18.0	1969Fu1
	4a	15.94	4	6.14	16	1.403	32	1.403	32	13.1 - 18.0	1975Sor
	4b	15.96	4	6.20	17	1.417	35	1.417	35	13.1 - 18.0	1975Sor
	5	15.69	1	4.94	4	1.203	7	1.203	7	13.1 - 18.0	2009Var
¹¹⁷ Sn	4	15.77	2	5.29	11	1.232	20	1.232	20	13.2 - 17.8	1974Le1
	4	15.77	2	5.31	7	1.208	11	1.208	11	13.1 - 17.9	1969Fu1
	4a	15.76	4	5.61	15	1.298	31	1.298	31	13.1 - 18.0	1975Sor
	4b	15.85	6	5.88	20	1.360	44	1.360	44	13.1 - 18.0	1975Sor
	5	15.75	1	5.22	3	1.209	5	1.209	5	13.1 - 18.0	2009Var
¹¹⁸ Sn	4	15.55	1	5.02	6	1.259	12	1.259	12	13.1 - 17.9	1974Le1
	4	15.70	1	5.02	5	1.148	8	1.148	8	13.1 - 17.9	1969Fu1
	4a	15.60	5	6.42	22	1.560	48	1.560	48	13.1 - 18.0	1974Sor
	4b	15.65	6	6.62	28	1.604	63	1.604	63	13.1 - 18.0	1974Sor
	5	15.56	1	4.69	2	1.109	4	1.109	4	13.1 - 18.0	2009Var
¹¹⁹ Sn	4	15.65	2	5.09	7	1.144	14	1.144	14	13.0 - 17.9	1969Fu1
	4a	14.85	3	4.76	14	0.968	22	0.968	22	13.1 - 17.9	1975Sor
	4b	14.84	4	4.70	17	0.956	32	0.956	32	13.1 - 17.9	1975Sor
	5	15.72	1	5.48	6	1.236	9	1.236	9	13.1 - 18.0	2009Var
¹²⁰ Sn	4	15.50	1	5.26	7	1.334	13	1.334	13	13.1 - 17.9	1974Le1
	4	15.53	1	5.03	4	1.256	9	1.256	9	13.1 - 17.9	1969Fu1
	4a	15.49	5	6.12	24	1.522	51	1.522	51	13.1 - 18.0	1974Sor
	4b	15.50	9	6.15	34	1.528	82	1.528	82	13.1 - 18.0	1974Sor
	5	15.45	1	4.85	2	1.221	4	1.221	4	13.1 - 18.0	2009Var
¹²² Sn	4a	15.45	4	4.98	13	1.158	26	1.158	26	13.1 - 18.0	1975Sor
	4b	15.50	6	5.12	18	1.194	44	1.194	44	13.1 - 18.0	1975Sor
	5	15.35	2	4.72	7	1.087	11	1.087	11	13.1 - 18.0	2009Var
¹²⁴ Sn	3a	15.41	5	5.06	24	1.198	37	1.198	37	13.1 - 18.0	2003Var
	3b	15.40	9	5.00	39	1.189	73	1.189	73	13.1 - 18.0	2003Var
	4	15.39	2	4.90	9	1.174	16	1.174	16	13.2 - 17.8	1974Le1
	4	15.30	2	4.98	8	1.222	16	1.222	16	13.1 - 17.8	1969Fu1
	4a	15.35	4	5.81	20	1.308	36	1.308	36	13.1 - 18.0	1975Sor
	4b	15.39	8	5.95	28	1.342	62	1.342	62	13.1 - 18.0	1975Sor
	5	15.28	1	4.90	3	1.173	5	1.173	5	13.1 - 18.0	2009Var
^{nat} Sb	4	15.62	1	5.59	4	1.356	8	1.356	8	10.2 - 17.8	1971Lep
¹²⁴ Te	6	15.36	2	5.81	9	1.394	18	1.394	18	12.0 - 18.9	1976Le2

Table 2 (continued)

Nucl	Id	$E_{r,1}$ (MeV)	$\Gamma_{r,1}$ (MeV)	s_1	$E_{r,2}$ (MeV)	$\Gamma_{r,2}$ (MeV)	s_2	s	$\varepsilon_{min} - \varepsilon_{max}$ (MeV)	Ref
^{126}Te	6	15.27 2	5.62 8	1.407 15				1.407 15	12.0 - 18.9	1976Le2
^{128}Te	6	15.23 2	5.55 8	1.414 16				1.414 16	12.0 - 18.9	1976Le2
^{130}Te	6	15.21 2	5.19 8	1.377 15				1.377 15	12.0 - 18.9	1976Le2
^{127}I	4	15.05 2	4.76 4	0.979 12	17.33 8	1.83 29	0.035 8	1.014 14	12.0 - 20.0	1999Bel
	4	14.64 22	4.00 27	0.860 179	16.51 39	3.21 91	0.215 166	1.075 244	12.2 - 20.0	1989Ras
	4	14.98 20	4.93 27	1.228 158	17.03 35	2.65 169	0.106 125	1.334 201	8.8 - 19.8	1969Be6
	4	14.65 18	4.34 26	0.650 117	16.80 22	4.24 64	0.311 119	0.961 167	12.1 - 19.9	1966Br1
	5	14.62 37	4.31 53	0.760 279	16.76 43	4.30 107	0.397 279	1.157 395	12.1 - 19.9	2006Var
	8	14.72 26	2.33 157	0.098 205	15.81 65	6.91 195	1.206 80	1.304 220	12.1 - 16.9	1987Ber
^{133}Cs	4	15.44 1	5.50 2	1.397 5				1.397 5	12.0 - 19.0	1974Le1
	4	15.32 2	5.11 8	1.186 14				1.186 14	12.1 - 18.7	1969Be1
	5	15.19 14	4.95 53	1.267 83				1.267 83	12.1 - 19.0	2016Va2
^{138}Ba	5	15.16 9	4.68 35	1.150 52				1.150 52	12.1 - 18.7	2016Va2
	4	15.31 2	4.76 6	1.208 10				1.208 10	12.1 - 18.7	1970Be8
$^{\text{nat}}\text{Ba}$	4	15.46 1	5.57 4	1.511 8				1.511 8	10.1 - 17.8	1971Be4
^{139}La	4	15.30 3	5.11 10	1.436 18				1.436 18	12.0 - 18.9	1972De1
	4	15.15 1	4.09 4	1.068 8				1.068 8	12.0 - 18.9	1971Be4
	4	15.19 3	4.60 9	1.283 19				1.283 19	12.1 - 19.0	1968Be5
	4a	14.44 6	3.34 19	0.751 21				0.751 21	12.5 - 19.0	1964Ric
	4b	14.48 12	3.18 33	0.741 48				0.741 48	12.5 - 19.0	1964Ric
^{140}Ce	6	15.09 1	4.51 5	1.317 10				1.317 10	12.0 - 18.9	1976Le2
^{142}Ce	6	14.95 2	5.24 9	1.313 16				1.313 16	12.0 - 18.9	1976Le2
^{141}Pr	4	15.33 2	4.49 9	1.165 21				1.165 21	12.1 - 16.9	1987Ber
	4	15.20 2	4.63 6	1.127 11				1.127 11	12.1 - 18.7	1966Br1
	4a	14.65 7	3.62 20	0.812 24				0.812 24	12.5 - 19.0	1964Ric
	4b	14.74 13	3.59 39	0.816 60				0.816 60	12.5 - 19.0	1964Ric
	12	15.42 2	4.39 5	1.081 10				1.081 10	12.0 - 18.1	1972You
	12	15.17 1	4.88 3	1.262 6				1.262 6	12.0 - 16.9	1971Be4
	12	15.26 1	4.09 4	1.049 8				1.049 8	12.1 - 19.0	1970Su1
	13	15.43 2	4.02 7	1.081 13				1.081 13	12.0 - 18.9	1972De1
^{142}Nd	3a	15.02 4	4.41 18	1.205 30				1.205 30	13.1 - 18.0	2003Var
	3b	15.04 12	4.50 49	1.221 111				1.221 111	13.1 - 18.0	2003Var
	4	15.01 1	4.56 4	1.226 7				1.226 7	12.0 - 18.9	1971Ca1
^{143}Nd	4	15.08 2	4.99 9	1.281 16				1.281 16	12.0 - 19.0	1971Ca1
^{144}Nd	4	15.17 2	5.56 7	1.295 14				1.295 14	12.0 - 18.9	1971Ca1
^{145}Nd	4	15.14 5	6.75 22	1.456 36				1.456 36	12.0 - 18.9	1971Ca1
^{146}Nd	4	14.88 3	6.04 11	1.361 20				1.361 20	12.0 - 18.9	1971Ca1
	4a	13.89 3	3.78 9	0.956 15				0.956 15	12.2 - 19.0	1971Vas
	4b	13.94 5	3.98 13	0.996 30				0.996 30	12.2 - 19.0	1971Vas
^{148}Nd	4	13.34 38	5.41 70	0.623 184	15.79 11	4.57 56	0.552 165	1.175 247	10.8 - 18.6	1971Ca1
	4	13.73 3	2.71 18	0.870 50	15.96 4	1.09 18	0.113 23	0.982 55	13.0 - 17.0	1969Vas
^{150}Nd	4	12.49 9	3.93 29	0.556 55	16.23 7	4.85 34	0.696 65	1.252 85	10.8 - 18.6	1971Ca1
^{144}Sm	4	15.37 2	4.53 7	1.274 15				1.274 15	12.1 - 18.9	1974Ca5
^{148}Sm	4	14.91 1	5.15 6	1.260 10				1.260 10	12.1 - 18.9	1974Ca5
	4a	14.02 2	4.04 5	0.947 11				0.947 11	7.7 - 19.0	1971Vas
	4b	14.17 4	4.38 8	1.029 19				1.029 19	7.7 - 19.0	1971Vas
^{150}Sm	4	14.76 3	6.01 12	1.340 20				1.340 20	12.1 - 18.9	1974Ca5
	4	13.99 3	4.40 11	1.137 19				1.137 19	12.2 - 19.0	1969Vas
^{152}Sm	4	12.56 4	3.53 13	0.511 25	15.97 3	4.77 14	0.704 30	1.215 39	10.9 - 18.8	1974Ca5
	4	11.74 2	2.24 8	0.751 23	15.00 3	2.31 10	0.672 24	1.423 33	11.0 - 18.8	1969Vas
^{154}Sm	0	12.31 29	3.27 1.04	0.483 217	15.95 36	5.53 176	0.729 305	1.212 374	10.9 - 18.6	1981Gur
	4	12.42 4	3.43 16	0.495 30	16.20 4	5.29 22	0.716 39	1.211 49	11.0 - 18.6	1974Ca5
	4	11.34 4	2.09 15	0.322 23	14.98 3	3.86 9	0.926 23	1.248 32	11.0 - 18.8	1969Vas
^{151}Eu	4a	13.43 34	5.18 31	0.623 163	14.52 7	2.28 60	0.258 133	0.880 210	10.2 - 18.0	1971Vas
	4b	13.27 68	5.04 67	0.545 285	14.54 7	2.59 80	0.327 226	0.872 364	10.2 - 18.0	1971Vas
^{153}Eu	4	12.47 5	3.26 18	0.416 31	16.07 5	5.48 21	0.769 42	1.185 52	10.9 - 18.7	1969Be8
	4a	11.99 3	2.19 11	0.282 13	15.26 4	3.12 13	0.536 18	0.819 22	11.0 - 18.8	1971Vas
	4b	12.00 3	2.23 11	0.292 14	15.19 6	2.88 17	0.496 29	0.788 33	11.0 - 18.8	1971Vas
^{152}Gd	4a	11.96 6	3.49 18	0.397 26	14.75 4	2.87 14	0.475 26	0.872 37	10.2 - 18.0	1971Vas
	4b	11.96 6	3.49 18	0.396 27	14.75 5	2.88 17	0.477 32	0.873 42	10.2 - 18.0	1971Vas
^{154}Gd	4a	12.09 5	3.08 17	0.380 23	15.10 4	3.09 15	0.493 27	0.873 35	10.2 - 18.0	1971Vas
	4b	12.11 5	3.11 17	0.389 24	15.07 5	2.93 18	0.467 33	0.856 41	10.2 - 18.0	1971Vas
^{156}Gd	0	12.63 24	3.61 80	0.632 177	15.97 23	4.10 105	0.537 191	1.169 260	10.9 - 18.7	1981Gur
	4a	12.00 4	3.50 17	0.501 25	15.00 4	2.06 11	0.343 19	0.844 31	11.0 - 18.8	1971Vas
	4b	12.01 6	4.19 37	0.583 52	15.02 6	2.09 20	0.322 39	0.905 65	11.0 - 18.8	1971Vas
^{158}Gd	4a	12.00 5	3.37 18	0.449 26	15.19 5	2.97 19	0.473 27	0.921 38	10.2 - 18.0	1971Vas
	4b	12.06 6	3.50 18	0.487 29	15.02 9	2.21 26	0.368 45	0.855 54	10.2 - 18.0	1971Vas

Table 2 (continued)

Nucl	Id	$E_{r,1}$ (MeV)	$\Gamma_{r,1}$ (MeV)	s_1	$E_{r,2}$ (MeV)	$\Gamma_{r,2}$ (MeV)	s_2	s	$\varepsilon_{min} - \varepsilon_{max}$ (MeV)	Ref							
^{237}Np	4	11.10	5	2.72	12	0.462	32	14.27	5	4.27	19	0.986	46	1.448	56	9.4 - 17.8	1986Be2
	4	11.20	22	3.50	71	0.483	136	14.24	14	4.22	64	0.682	149	1.165	202	9.5 - 16.5	1973Ve1
	5	11.04	13	2.87	32	0.376	64	14.20	13	4.30	47	0.758	90	1.133	110	9.3 - 17.7	2007Var
	10	11.09	7	2.68	16	0.452	45	14.26	8	4.32	30	1.000	70	1.452	83	9.4 - 17.8	1986Be2
^{239}Pu	10	11.21	37	3.59	98	0.494	209	14.32	26	4.46	125	0.710	265	1.204	337	9.2 - 16.6	1973Ve1
	0	11.47	46	4.50	108	0.626	260	14.37	26	4.76	98	0.609	253	1.234	362	9.3 - 18.7	1976Gu1
	5	11.76	47	4.03	108	0.759	319	14.08	17	2.41	164	0.252	258	1.012	410	9.2 - 17.7	2007Var
	10	11.41	16	3.05	24	0.515	111	14.05	24	4.23	72	0.641	150	1.156	187	9.1 - 17.8	1986Be2

Table 3

References to experimental and evaluated cross section data taken from EXFOR.

Nucl	Id	Reaction	Ref	EXFOR
⁶ Li	2	γ , abs	1986Var	M0140016
	4	γ , sn	1965Be1	L0008005
⁷ Li	0	γ , abs	1985Ahr	M0188012
	2	γ , abs	1986Var	M0140024
	4	γ , sn	1973Bra	L0030005
⁹ Be	0	γ , abs	1975Ahr	M0825004
¹⁰ B	4	γ , sn	1987Ahs	M0207002
¹² C	0	γ , abs	1969Bez	L0064002
	1	γ , abs	1975Ahr	M0372004
	3	γ , abs	2002Ish	M0648002
	4	γ , sn	1963Bur	M0160002
¹³ C	3	γ , abs	2002Ish	M0648003
¹⁴ C	3	γ , abs	2002Ish	M0648004
^{nat} C	0	γ , abs	1985Ahr	M0188014
	0	γ , abs	1975Ahr	M0825006
	0	γ , abs	1972Ahr	L0083005
	1	γ , abs	1965Wyc	L0122003
¹⁴ N	0	γ , abs	1969Bez	L0064003
	3	γ , abs	2002Ish	M0648005
¹⁵ N	3	γ , abs	2002Ish	M0648006
¹⁶ O	0	γ , abs	1975Ahr	M0372005
	0	γ , abs	1972Ahr	L0083004
	0	γ , abs	1969Bez	L0064004
	0	γ , abs	1963Bur	L0127002
	1	γ , abs	1965Wyc	L0122004
	3	γ , abs	2002Ish	M0648007
¹⁷ O	3	γ , abs	2002Ish	M0648008
¹⁸ O	3	γ , abs	2002Ish	M0648009
^{nat} O	0	γ , abs	1985Ahr	M0188015
	0	γ , abs	1975Ahr	M0825008
¹⁹ F	3	γ , abs	2002Ish	M0648010
²³ Na	0	γ , abs	1981Ish	M0043025
	1	γ , abs	1965Wyc	L0122005
	3	γ , abs	2002Ish	M0648011
	4	γ , sn	1971Alv	L0022008
²⁴ Mg	0	γ , abs	1966Dol	M0727003
	3	γ , abs	2003Var	M0656003
	3	γ , abs	2002Ish	M0648012
²⁵ Mg	3	γ , abs	2002Ish	M0648013
	4	γ , sn	1971Alv	L0022009
²⁶ Mg	3	γ , abs	2003Var	M0656004
	3	γ , abs	2002Ish	M0648014
	4	γ , sn	1971Fu2	L0026006
^{nat} Mg	1	γ , abs	1965Wyc	L0122006
²⁷ Al	0	γ , abs	1985Ahr	M0188016
	0	γ , abs	1975Ahr	M0825010
	0	γ , abs	1972Ahr	L0083006
	0	γ , abs	1975Ahr	M0372006
	3	γ , abs	2002Ish	M0648015
	4	γ , sn	1966Fu1	L0010007
²⁸ Si	3	γ , abs	2003Var	M0656005
	3	γ , abs	2002Ish	M0648016
²⁹ Si	3	γ , abs	2002Ish	M0648017
³⁰ Si	3	γ , abs	2002Ish	M0648018
^{nat} Si	0	γ , abs	1975Ahr	M0372007
³² S	0	γ , abs	1968Dol	M0824002
	3	γ , abs	2002Ish	M0648019
³⁴ S	0	γ , abs	1986Ass	M0510006
	3	γ , abs	2003Var	M0656006
	3	γ , abs	2002Ish	M0648020
^{nat} S	0	γ , abs	1965Wyc	L0122009
⁴⁰ Ar	3	γ , abs	2002Ish	M0648021
^{nat} K	13	γ , xn	1974Ve1	L0039036
⁴⁰ Ca	0	γ , abs	1966Dol	M0727004
	3	γ , abs	2003Ero	M0653002
⁴² Ca	3	γ , abs	2003Ero	M0653003
⁴⁴ Ca	3	γ , abs	2003Ero	M0653004

Table 3 (continued)

Nucl	Id	Reaction	Ref	EXFOR
⁴⁸ Ca	0	γ , abs	1987OKe	M0636010
	3	γ , abs	2003Ero	M0653005
natCa	0	γ , abs	1975Ahr	M0372008
	0	γ , abs	1972Ahr	L0083003
	0	γ , abs	1968Bez	M0840003
	1	γ , abs	1965Wyc	L0122010
	2	γ , abs	1985Ahr	M0188017
⁴⁶ Ti	3	γ , abs	2002Ish	M0648026
⁴⁸ Ti	3	γ , abs	2002Ish	M0648027
⁵¹ V	4	γ , sn	1962Fu1	L0001008
	3	γ , abs	2003Var	M0656007
	4	γ , sn	1969Gor	M0093005
⁵² Cr	3	γ , abs	2002Ish	M0648028
	4	γ , sn	1969Gor	M0093006
⁵⁵ Mn	4	γ , sn	1979Al2	L0028011
	4	γ , sn	1974Car	M0835002
	4	γ , sn	1970Is1	M0726003
⁵⁴ Fe	0	γ , abs	1978Nor	M0507004
⁵⁹ Co	4	γ , sn	1979Al2	L0028012
	4	γ , sn	1964Baz	M0106003
	4	γ , sn	1969Gor	M0093007
⁵⁸ Ni	3	γ , abs	2003Var	M0656008
	3	γ , abs	2002Ish	M0648029
	4	γ , sn	1974Fu3	L0034002
	4	γ , sn	1970Gor	M0597002
⁶⁰ Ni	3	γ , abs	2003Var	M0656002
	4	γ , sn	1974Fu3	L0034008
	4	γ , sn	1970Gor	M0597003
⁶³ Cu	3	γ , abs	2003Var	M0656010
	3	γ , abs	2002Ish	M0648031
	4	γ , sn	1964Fu1	L0006012
	5	γ , sn	2016Va1	M0920002
	6	γ , sn	1968Su1	L0013002,L0013003
	3	γ , abs	2003Var	M0656011
⁶⁵ Cu	3	γ , abs	2002Ish	M0648032
	4	γ , sn	1964Fu1	L0006013
	5	γ , sn	2016Va1	M0920005
	1	γ , abs	1965Wyc	L0122013
natCu	4	γ , sn	1964Ful	L0006011
	4	γ , sn	1970Is2	M0537005
	4	γ , sn	1968Tom	L0108004
	8	γ , sn	1976Ca1	L0043002,L0043003
	1	γ , abs	2003Rod	M0652007
⁶⁴ Zn	8	γ , sn	1975Mcc	M0496004
⁷⁰ Ge	0	γ , abs	1975Mcc	M0496010
	8	γ , sn	1976Ca1	L0043008,L0043009
⁷² Ge	0	γ , abs	1975Mcc	M0496010
	8	γ , sn	1976Ca1	L0043011,L0043012
	8	γ , sn	1976Ca1	L0043014,L0043015
⁷⁴ Ge	0	γ , abs	1975Mcc	M0496013
	8	γ , sn	1976Ca1	L0043017,L0043018
⁷⁶ Ge	0	γ , abs	1975Mcc	M0496007
	4	γ , sn	1969Be1	L0014012
⁷⁵ As	8	γ , sn	1976Ca1	L0043020,L0043021
⁷⁶ Se	0	γ , abs	1978Gur	M0023002
	8	γ , sn	1976Ca1	L0043023,L0043024
⁷⁸ Se	8	γ , sn	1976Ca1	L0043026,L0043027
⁸⁰ Se	5	γ , sn	2016Va1	M0920008
	8	γ , sn	1976Ca1	L0043029,L0043030
⁸² Se	0	γ , abs	1978Gur	M0023003
	8	γ , sn	1976Ca1	L0043032,L0043033
	4	γ , sn	1971Lep	L0027017
natRb	4	γ , sn	1971Lep	L0027018
natSr	4	γ , sn	1971Le1	L0027019
⁸⁹ Y	4	γ , sn	2003Var	M0656012
	3	γ , abs	1999Bel	M0706002
	4	γ , sn	1967Be2	L0011018
	4	γ , sn	1972You	L0059002
	12	γ , ln	2003Var	M0656013
⁹⁰ Zr	3	γ , abs	2003Var	M0656013

Table 3 (continued)

Nucl	Id	Reaction	Ref	EXFOR
	4	γ , sn	1972Ask	M0360003
	4	γ , sn	1967Be2	L0011019
	8	γ , sn	1971Le1	L0027012,L0027013
⁹¹ Zr	4	γ , sn	1967Be2	L0011020
	5	γ , sn	2015Var	M0914002
⁹² Zr	4	γ , sn	1967Be2	L0011021
⁹⁴ Zr	4	γ , sn	1967Be2	L0011022
	5	γ , sn	2015Var	M0914005
⁹³ Nb	8	γ , sn	1971Le1	L0027015,L0027016
⁹² Mo	3	γ , abs	2003Var	M0656014
	4	γ , sn	1974Be3	L0032020
	4	γ , sn	1970Is3	M0719003
⁹⁴ Mo	4	γ , sn	1974Be3	L0032021
⁹⁶ Mo	4	γ , sn	1974Be3	L0032022
⁹⁸ Mo	4	γ , sn	1974Be3	L0032023
	4	γ , sn	1970Is3	M0719006
¹⁰⁰ Mo	9	γ , sn	1974Be3	L0032017,L0032018,L0032019
¹⁰³ Rh	3	γ , abs	2003Var	M0656015
	4	γ , sn	1974Le1	L0035041
	4	γ , sn	1962Bog	M0769006
¹⁰⁸ Pd	0	γ , abs	1969Dea	M0839004
^{nat} Pd	4	γ , sn	1971Lep	L0035042
¹⁰⁷ Ag	4	γ , sn	1969Ish	M0524002
	4	γ , sn	1969Be1	L0014013
	4	γ , sn	1962Bog	M0769011
¹⁰⁹ Ag	4	γ , sn	1969Ish	M0524003
^{nat} Ag	4	γ , sn	1971Lep	L0035043
^{nat} Cd	4	γ , sn	1971Lep	L0035044
¹¹⁵ In	4	γ , sn	1974Le1	L0035045
	4	γ , sn	1969Fu1	L0017029
	4	γ , sn	1962Bog	M0769003
	5	γ , sn	2013Var	M0863005
¹¹² Sn	4	γ , sn	1974Sor	M0745005
	5	γ , sn	2009Var	M0768038
¹¹⁴ Sn	4	γ , sn	1975Sor	M0746008
	5	γ , sn	2009Var	M0768039
¹¹⁶ Sn	4	γ , sn	1974Le1	L0035046
	4	γ , sn	1969Fu1	L0017030
	4	γ , sn	1975Sor	M0746009
	5	γ , sn	2009Var	M0768040
¹¹⁷ Sn	4	γ , sn	1974Le1	L0035047
	4	γ , sn	1969Fu1	L0017031
	4	γ , sn	1975Sor	M0746010
	5	γ , sn	2009Var	M0768041
¹¹⁸ Sn	4	γ , sn	1974Le1	L0035048
	4	γ , sn	1969Fu1	L0017032
	4	γ , sn	1974Sor	M0745006
	5	γ , sn	2009Var	M0768042
¹¹⁹ Sn	4	γ , sn	1969Fu1	L0017033
	4	γ , sn	1975Sor	M0746011
	5	γ , sn	2009Var	M0768043
¹²⁰ Sn	4	γ , sn	1974Le1	L0035049
	4	γ , sn	1969Fu1	L0017034
	4	γ , sn	1974Sor	M0745007
	5	γ , sn	2009Var	M0768044
¹²² Sn	4	γ , sn	1975Sor	M0746012
	5	γ , sn	2009Var	M0768045
¹²⁴ Sn	3	γ , abs	2003Var	M0656016
	4	γ , sn	1974Le1	L0035050
	4	γ , sn	1969Fu1	L0017035
	4	γ , sn	1975Sor	M0746013
	5	γ , sn	2009Var	M0768046
^{nat} Sb	4	γ , sn	1971Lep	L0035051
¹²⁴ Te	6	γ , sn	1976Le2	L0042004,L0042002
¹²⁶ Te	6	γ , sn	1976Le2	L0042007,L0042005
¹²⁸ Te	6	γ , sn	1976Le2	L0042010,L0042008
¹³⁰ Te	6	γ , sn	1976Le2	L0042013,L0042011
¹²⁷ I	4	γ , sn	1999Bel	M0706004

Table 3 (continued)

Nucl	Id	Reaction	Ref	EXFOR
	4	γ , sn	1989Ras	M0511002
	4	γ , sn	1969Be6	L0015022
	4	γ , sn	1966Br1	L0009009
	5	γ , sn	2006Var	M0693003
	8	γ , sn	1987Ber	L0057005,L0057006
¹³³ Cs	4	γ , sn	1974Le1	L0035053
	4	γ , sn	1969Be1	L0014014
	5	γ , sn	2016Va2	M0922002
¹³⁸ Ba	5	γ , sn	2016Va2	M0922006
	4	γ , sn	1970Be8	L0019008
^{nat} Ba	4	γ , sn	1971Be4	L0024015
¹³⁹ La	4	γ , sn	1972De1	M0398004
	4	γ , sn	1971Be4	L0024016
	4	γ , sn	1968Be5	L0012018
	4	γ , sn	1964Ric	M0928003
¹⁴⁰ Ce	6	γ , sn	1976Le2	L0042016,L0042014
¹⁴² Ce	6	γ , sn	1976Le2	L0042019,L0042017
¹⁴¹ Pr	4	γ , xn	1987Ber	L0057015
	4	γ , sn	1966Br1	L0009010
	4	γ , sn	1964Ric	M0928005
	12	γ , ln	1972You	L0059003
	12	γ , ln	1971Be4	L0024011
	12	γ , ln	1970Su1	L0020002
	13	γ , xn	1972De1	M0398002
¹⁴² Nd	3	γ , abs	2003Var	M0656017
	4	γ , sn	1971Ca1	L0025023
¹⁴³ Nd	4	γ , sn	1971Ca1	L0025024
¹⁴⁴ Nd	4	γ , sn	1971Ca1	L0025025
¹⁴⁵ Nd	4	γ , sn	1971Ca1	L0025026
¹⁴⁶ Nd	4	γ , sn	1971Ca1	L0025027
	4	γ , sn	1971Vas	M0817003
¹⁴⁸ Nd	4	γ , sn	1971Ca1	L0025028
	4	γ , sn	1969Vas	M0816003
¹⁵⁰ Nd	4	γ , sn	1971Ca1	L0025029
¹⁴⁴ Sm	4	γ , sn	1974Ca5	L0033017
¹⁴⁸ Sm	4	γ , sn	1974Ca5	L0033018
	4	γ , sn	1971Vas	M0817005
¹⁵⁰ Sm	4	γ , sn	1974Ca5	L0033019
	4	γ , sn	1969Vas	M0816007
¹⁵² Sm	4	γ , sn	1974Ca5	L0033020
	4	γ , sn	1969Vas	M0816009
¹⁵⁴ Sm	0	γ , abs	1981Gur	M0073002
	4	γ , sn	1974Ca5	L0033021
	4	γ , sn	1969Vas	M0816011
¹⁵¹ Eu	4	γ , sn	1971Vas	M0817015
¹⁵³ Eu	4	γ , sn	1969Be8	L0016018
	4	γ , sn	1971Vas	M0817017
¹⁵² Gd	4	γ , sn	1971Vas	M0817007
¹⁵⁴ Gd	4	γ , sn	1971Vas	M0817009
¹⁵⁶ Gd	0	γ , abs	1981Gur	M0073003
	4	γ , sn	1971Vas	M0817011
¹⁵⁸ Gd	4	γ , sn	1971Vas	M0817013
¹⁶⁰ Gd	3	γ , abs	2003Var	M0656018
	4	γ , sn	1969Be8	L0016019
¹⁵⁹ Tb	4	γ , sn	1976Gor	M0057002
	4	γ , sn	1968Be5	L0012019
	4	γ , sn	1964Br1	L0005006
	4	γ , sn	1962Bog	M0769014
	4	γ , sn	1958Ful	L0111002
	5	γ , sn	2011Var	M0831005
¹⁶⁵ Ho	0	γ , abs	1981Gur	M0073004
	0	γ , abs	1976Gu2	M0056002
	4	γ , sn	1969Be8	L0016020
	4	γ , sn	1968Be5	L0012020
	4	γ , sn	1966Axe	M0844003
	4	γ , sn	1976Gor	M0057006
¹⁶⁸ Er	0	γ , abs	1981Gur	M0073005
¹⁷⁴ Yb	0	γ , abs	1981Gur	M0073006

Table 3 (continued)

Nucl	Id	Reaction	Ref	EXFOR
¹⁷⁵ Lu	4	γ , sn	1969Be6	L0015026
¹⁷⁶ Hf	4	γ , sn	1977Gor	M0007002
¹⁷⁸ Hf	0	γ , abs	1981Gur	M0073007
	0	γ , abs	1976Gu2	M0056003
	4	γ , sn	1977Gor	M0007003
	4	γ , sn	1976Gor	M0057012
¹⁸⁰ Hf	0	γ , abs	1981Gur	M0073008
	0	γ , abs	1976Gu2	M0056004
	4	γ , sn	1977Gor	M0007004
¹⁸¹ Ta	0	γ , abs	1981Gur	M0073009
	0	γ , abs	1976Gu2	M0056005
	3	γ , abs	2003Var	M0656019
	4	γ , sn	2001Bel	M0707003
	4	γ , sn	1968Be5	L0012021
	4	γ , sn	1963Br1	L0003005
	4	γ , sn	1962Bog	M0769009
	4	γ , sn	1969Ish	M0737003
	4	γ , sn	1967Ant	M0539005
	4	γ , sn	1958Ful	L0111003
	5	γ , sn	2012Var	M0850006
¹⁸² W	0	γ , abs	1981Gur	M0073010
	0	γ , abs	1976Gu2	M0056006
	4	γ , sn	1978Go3	M0025002
¹⁸⁴ W	0	γ , abs	1981Gur	M0073011
	4	γ , sn	1978Go3	M0025003
	4	γ , sn	1973Gor	M0645007
¹⁸⁶ W	0	γ , abs	1981Gur	M0073012
	4	γ , sn	1978Go3	M0025004
	4	γ , sn	1969Be8	L0016021
	4	γ , sn	1973Gor	M0645008
¹⁸⁵ Re	4	γ , sn	1973Gor	M0645009
^{nat} Re	4	γ , sn	1975Vey	M0845013
¹⁸⁶ Os	5	γ , sn	2015Var	M0915002
	7	γ , sn	1979Be4	L0046004,L0046002
¹⁸⁸ Os	5	γ , sn	2014Var	M0916005
	9	γ , sn	1979Be4	L0046005,L0046006,L0046007
¹⁸⁹ Os	5	γ , sn	2014Var	M0916009
	9	γ , sn	1979Be4	L0046009,L0046010,L0046011
¹⁹⁰ Os	5	γ , sn	2015Var	M0915007
	9	γ , sn	1979Be4	L0046013,L0046014,L0046015
¹⁹² Os	5	γ , sn	2015Var	M0915011
	9	γ , sn	1979Be4	L0046017,L0046018,L0046019
¹⁹¹ Ir	4	γ , sn	1978Go1	M0008002
	4	γ , sn	1978Go2	M0049002
¹⁹³ Ir	4	γ , sn	1978Go1	M0008003
^{nat} Ir	4	γ , sn	1975Vey	M0845014
¹⁹⁴ Pt	4	γ , sn	1978Go1	M0008004
	13	γ , xn	1978Go2	M0049006
¹⁹⁵ Pt	4	γ , sn	1978Go1	M0008005
	13	γ , xn	1978Go2	M0049008
¹⁹⁶ Pt	4	γ , sn	1978Go1	M0008006
¹⁹⁸ Pt	4	γ , sn	1978Go1	M0008007
¹⁹⁷ Au	0	γ , abs	1981Gur	M0073013
	0	γ , abs	1976Gu2	M0056007
	4	γ , sn	1970Ve1	L0021010
	4	γ , sn	1962Fu2	L0002005
	4	γ , sn	1973Sor	M0650003
	4	γ , sn	1958Ful	L0111004
	8	γ , sn	1987Ber	L0057009,L0057010
	8	γ , sn	2010Var	M0798003
²⁰³ Tl	4	γ , sn	1970Ant	L0109002
²⁰⁵ Tl	4	γ , sn	1970Ant	L0109003
²⁰⁶ Pb	4	γ , sn	1964Ha2	L0007014
	4	γ , sn	1973Sor	L0107002
²⁰⁷ Pb	4	γ , sn	1964Ha2	L0007015
²⁰⁸ Pb	3	γ , abs	2003Var	M0656020
	4	γ , sn	1995Bel	M0702003
	4	γ , sn	1970Ve1	L0021011

Table 3 (continued)

Nucl	Id	Reaction	Ref	EXFOR
natPb	4	γ , sn	1964Ha2	L0007016
	4	γ , sn	1973Sor	L0107004
	5	γ , sn	2013Ish	M0864028
	12	γ , ln	1972You	L0059004
	2	γ , abs	1985Ahr	M0188018
²⁰⁹ Bi	8	γ , sn	1987Ber	L0057012,L0057013
	0	γ , abs	1976Gu2	M0056008
	4	γ , sn	1995Bel	M0702004
	4	γ , sn	1964Ha2	L0007017
	4	γ , sn	1973Sor	M0650005
²³² Th	4	γ , sn	1967Ant	M0539008
	5	γ , sn	2016Va2	M0922010
	12	γ , sn	1972You	L0059005
	0	γ , abs	1976Gu1	M0090002
	4	γ , sn	1980Ca1	L0050030
²³³ U	4	γ , sn	1973Ve1	L0031014
	5	γ , sn	2007Var	M0722003
	10	γ , sn	1980Ca1	L0050002,L0050003,L0050004
²³⁴ U	4	γ , sn	1986Be2	L0058026
	11	γ , sn	1986Be2	L0058004,L0058003,L0058002
²³⁵ U	4	γ , sn	1986Be2	L0058027
	11	γ , sn	1986Be2	L0058007,L0058006,L0058005
²³⁶ U	0	γ , abs	1976Gu1	M0090003
	4	γ , sn	1980Ca1	L0050032
²³⁸ U	4	γ , sn	1980Ca1	L0050033
	10	γ , sn	1980Ca1	L0050010,L0050011,L0050012
	0	γ , abs	1976Gu1	M0090004
natU	4	γ , sn	1980Ca1	L0050031
	4	γ , sn	1973Ve1	L0031015
	5	γ , sn	2007Var	M0722008
	2	γ , abs	1985Ahr	M0188019
	²³⁷ Np	4	γ , sn	1986Be2
4		γ , sn	1973Ve1	L0031017
5		γ , sn	2007Var	M0722013
10		γ , sn	1986Be2	L0058008,L0058009,L0058010
10		γ , sn	1973Ve1	L0031007,L0031008,L0031009g
²³⁹ Pu	0	γ , abs	1976Gu1	M0090005
	5	γ , sn	2007Var	M0722018
	10	γ , sn	1986Be2	L0058012,L0058013,L0058014

Graphs 1 – 162





















































