NEW INTEGRAL REPRESENTATIONS FOR RANKIN–SELBERG L-FUNCTIONS

ANDREW R. BOOKER, M. KRISHNAMURTHY, AND MIN LEE

ABSTRACT. We derive integral representations for the Rankin–Selberg L-functions on $\mathrm{GL}(3) \times \mathrm{GL}(1)$ and $\mathrm{GL}(3) \times \mathrm{GL}(2)$ by a process of unipotent averaging at archimedean places. A key feature of our result is that it allows one to fix the choice of test vector at finite places, irrespective of ramification. This enables a new proof of the functional equation for $\mathrm{GL}(3) \times \mathrm{GL}(2)$ Rankin–Selberg L-functions in many cases.

1. Introduction

Let $f(z) = \sum_{n=1}^{\infty} a_n e^{2\pi i n z}$ be a holomorphic newform of weight k and level $\Gamma_0(N)$. Let χ be a primitive Dirichlet character of conductor q and set $f_{\chi}(z) = \sum_{n=1}^{\infty} a_n \chi(n) e^{2\pi i n z}$. Then, by [20, Lemma 4.3.10], we have

$$f_{\chi} = \frac{\tau(\chi)}{q} \sum_{q=1}^{q} \overline{\chi(-a)} f \Big|_{k} \left(\begin{smallmatrix} 1 & a/q \\ & 1 \end{smallmatrix} \right),$$

where $\tau(\chi)$ is the associated Gauss sum. Taking the Mellin transform of both sides, we obtain

(1.1)
$$\Lambda(s, f \times \chi) = \frac{\tau(\chi)}{q} \sum_{a=1}^{q} \overline{\chi(-a)} \Lambda(s, f, a/q),$$

where $\Gamma_{\mathbb{C}}(s) = 2(2\pi)^{-s}\Gamma(s)$,

$$\Lambda(s,f,a/q) = \Gamma_{\mathbb{C}}(s) \sum_{n=1}^{\infty} a_n e^{2\pi i a n/q} n^{-s} \quad \text{and} \quad \Lambda(s,f \times \chi) = \Gamma_{\mathbb{C}}(s) \sum_{n=1}^{\infty} a_n \chi(n) n^{-s}.$$

The above notion of unipotent averaging is a key ingredient in Weil's proof of the converse theorem [20, Theorem 4.3.15] as well as in our prior work [1, 3] on GL(2) converse theorems. The goal of the present paper is to prove an analogue of (1.1) for higher-rank Rankin-Selberg L-functions using a similar process of unipotent averaging; see Theorem 3.4. Our result can be interpreted as an integral representation for the Rankin-Selberg L-function in terms of the associated (Whittaker) essential functions, and we speculate that this may have applications to the study of algebraicity properties and nonvanishing of special values. We outline some necessary notation and explain our result in greater detail below.

Let F be a number field with adèle ring \mathbb{A}_F , and let $\pi \cong \bigotimes \pi_v$ (resp. $\tau \cong \bigotimes \tau_v$) be an irreducible generic (with respect to some fixed additive character $\psi = \bigotimes \psi_v$ of $F \setminus \mathbb{A}_F$) automorphic representation of $\mathrm{GL}_n(\mathbb{A}_F)$ (resp. $\mathrm{GL}_m(\mathbb{A}_F)$), with m < n. For each place v of F, let $L(s, \pi_v \boxtimes \tau_v)$ be the local Rankin–Selberg factor [11, 12] and set

$$\Lambda(s, \pi \boxtimes \tau) = \prod_{v} L(s, \pi_v \boxtimes \tau_v).$$

A. R. B. was partially supported by EPSRC Grant EP/K034383/1. M. L. was supported by a Royal Society University Research Fellowship. No data were created in the course of this study.

The completed L-function $\Lambda(s, \pi \boxtimes \tau)$, definied initially for $\Re(s) \gg 1$, continues to a meromorphic function of $s \in \mathbb{C}$ and satisfies a functional equation relating $\Lambda(s, \pi \boxtimes \tau)$ to $\Lambda(1 - s, \widetilde{\pi} \boxtimes \widetilde{\tau})$ [11]. The collection of such functions for a fixed π and suitably varying τ plays a central role in converse theorems, whose purpose is to characterize automorphic representations π in terms of the analytic behavior of $\Lambda(s, \pi \boxtimes \tau)$.

For each non-archimedean place v of F, let ξ_v^0 (resp. φ_v^0) denote the "essential vector" in the space of π_v (resp. τ_v) [10, 13, 19], and let $W_{\xi_v^0} \in \mathcal{W}(\pi_v, \psi_v)$ (resp. $W_{\varphi_v^0} \in \mathcal{W}(\tau_v, \psi_v^{-1})$) be the associated essential Whittaker functions (as in loc. cit.) formed relative to a ψ_v whose conductor is \mathfrak{o}_v . When τ_v is unramified, it follows from [19, Corollary 3.3] that

$$(1.2) L(s, \pi_v \boxtimes \tau_v) = \int_{U_m(F_v) \backslash \operatorname{GL}_m(F_v)} W_{\xi_v^0} \begin{pmatrix} h_v \\ I_{n-m} \end{pmatrix} W_{\varphi_v^0}(h_v) \| \det h_v \|_v^{s - \frac{n-m}{2}} dh_v$$

for a suitable normalization of the measure on $U_m(F_v)\backslash GL_m(F_v)$. For m=n-1, the above equality is a part of the characterization of the essential vector in [10, 13], which we review in the next section; the fact that it is true for any $m \leq n-1$ is a result of a concrete realization of essential functions in [19]. It follows from this that, if τ is such that τ_v is unramified at every finite v, then the Rankin–Selberg L-function $\Lambda(s, \pi \boxtimes \tau)$ can be represented by a global integral of the corresponding automorphic forms, which unfolds to a product of local integrals of the above shape.

However, when τ is ramified at a place v, the local integral can vanish unless ξ_v^0 is adjusted in a suitable way. In fact in general one has that there is an element $t_v^0 \in \mathcal{W}(\pi_v, \psi_v) \otimes \mathcal{W}(\tau_v, \psi_v^{-1})$ whose local zeta integral is precisely the L-function $L(s, \pi_v \boxtimes \tau_v)$. Now, let S be the finite set of finite places where τ_v is ramified. For $v \in S$, let t_v^0 be as described above, and for a finite place v not in S set $t_v^0 = W_{\xi_v^0} \otimes W_{\varphi_v^0}$. For any choice of a pair of archimedean Whittaker functions (W_∞, W_∞') , writing each t_v^0 for $v \in S$ as a sum of pure tensors, we get a finite sum of pure (global) tensors such that

$$\sum_{j} W_{j} \otimes W'_{j} = (W_{\infty} \otimes W'_{\infty}) \prod_{v < \infty} t_{v}^{0}.$$

Fixing such a decomposition, let (ϕ_j, φ_j) be the automorphic forms associated with (W_j, W_j) . Then

$$(1.3) P_{\infty}(s)\Lambda(s,\pi\boxtimes\tau) = \sum_{j} \int_{\mathrm{GL}_{m}(F)\backslash\mathrm{GL}_{m}(\mathbb{A}_{F})} \mathbb{P}_{m}^{n}(\phi_{j}) \left(\begin{pmatrix} h \\ I_{n-m} \end{pmatrix} \right) \varphi_{j}(h) \|\det h\|^{s-\frac{n-m}{2}} dh,$$

where $P_{\infty}(s)$ is an entire function depending only on the pair $(W_{\infty}, W'_{\infty})$, and \mathbb{P}_m^n is the projection operator defined in (2.4) below.

In this paper, we study integral representations for $L(s,\pi\boxtimes\tau)$ in the case $n=3,\,m\in\{1,2\}$. We show that (in the above notation) one can take $t_v^0=W_{\xi_v^0}\otimes W_{\varphi_v^0}$ at all finite places v, regardless of the ramification of τ , after going through a process of unipotent averaging on the right-hand side of (1.3). To illustrate our main theorem (Theorem 3.4), let us consider the classical case, $F=\mathbb{Q}$. Let $N,q\geq 1$ be the conductors of π and τ , respectively, and assume for simplicity that (N,q)=1 and τ_v is a twist-minimal principal series representation for every finite place v. We fix the essential vector $\xi_f^0=\bigotimes_{v<\infty}\xi_v^0$ and consider pure tensors ξ in the space of π of the form $\xi=\xi_\infty\otimes\xi_f^0$. Let U_ξ be the function on $\mathrm{GL}_3(\mathbb{A}_\mathbb{Q})$ obtained by summing the associated Whittaker function W_ξ as in (2.3). Then U_ξ may be viewed as an automorphic function on $\mathrm{GL}_2(\mathbb{A}_\mathbb{Q})$ by restriction via $h\mapsto \binom{h}{1}$. It in fact corresponds to a classical cusp form of level 1. If we write Φ to denote the restriction of U_ξ to $\mathrm{GL}_3(\mathbb{R})$, then the corresponding classical automorphic form on $\mathrm{GL}_2(\mathbb{R})$ is given by $h\mapsto \Phi\binom{h}{1}$ for $h\in \mathrm{GL}_2(\mathbb{R})$. Our key observation is that by shifting this function by a rational unipotent element, one can produce automorphic forms (on $\mathrm{GL}_2(\mathbb{R})$) of arbitrary level. To

be precise, let $(\beta_1, \beta_2) \in \mathbb{Q}^2$ with common denominator q, and consider the function

$$h \mapsto \Phi \begin{pmatrix} h & {}^t \beta \\ & 1 \end{pmatrix}, \text{ for } h \in \mathrm{GL}_2(\mathbb{R}).$$

As we show in Section 3.1, this is an automorphic form with respect to the principal congruence subgroup $\Gamma(q)$ of level q. If φ is an automorphic function in the space of τ that corresponds to a pure tensor whose component at every finite place v is φ_n^0 , put

$$\Lambda(s,\pi,\tau,\xi_{\infty},\varphi_{\infty},\beta) = \int_{\Gamma(q)\backslash\operatorname{GL}_2(\mathbb{R})} \Phi\begin{pmatrix} h & t\beta \\ & 1 \end{pmatrix} \varphi(h) \|\det\,h\|_{\mathbb{R}}^{s-\frac{1}{2}} \, dh.$$

We can paraphrase Theorem 3.4 in this context as saying that

$$P_{\infty}(s)\Lambda(s,\pi\boxtimes\tau)=\Lambda(s,\pi,\tau,\xi_{\infty},\varphi_{\infty},(0,1/q)),$$

for a certain entire function $P_{\infty}(s)$, depending on the pair $(\xi_{\infty}, \varphi_{\infty})$. (Notice the similarity with (1.3), albeit here the integral on the right-hand side involves classical forms.) Further, applying Theorem 3.4 we deduce the analytic continuation and functional equation of $\Lambda(s, \pi \boxtimes \tau)$ (cf. Theorem 3.9). Our proof is different from the proof of Jacquet, Piatetski-Shapiro and Shalika [11], in that it does not use the local functional equations at ramified places. The argument is analogous to the proof of the functional equation for additive twists on GL(2) (see [18, §A.3–A.4]), and it is based on a higher-rank matrix identity appearing in [16].

Theorem 3.4 is similar in spirit to the global Birch lemma of [15, 17], which plays a central role in the study of \mathfrak{p} -adic interpolation of special values of $L(s, (\pi \otimes \chi) \boxtimes \tau)$ for a cuspidal pair (π, τ) and a Hecke character χ of finite order and non-trivial \mathfrak{p} -power conductor. While in loc. cit. the unipotent averaging is done at the finite place corresponding to \mathfrak{p} , here we do it over the archimedean places as in [1], which allows us to retain the essential vector at all finite places. In addition, it should be emphasized that χ can be taken to be trivial in our setting.

Acknowledgements. The second author (M. K.) would like to thank A. Raghuram for pointing him to [15].

2. Preliminaries

Let \mathfrak{o}_F denote the ring of integers of F. For each place v of F, let F_v denote the completion of F at v. For $v < \infty$, let \mathfrak{o}_v denote the ring of integers in F_v , \mathfrak{p}_v the unique maximal ideal in \mathfrak{o}_v and q_v the cardinality of $\mathfrak{o}_v/\mathfrak{p}_v$. Also, for every finite v, fix a generator ϖ_v of \mathfrak{p}_v with absolute value $\|\varpi_v\|_v = q_v^{-1}$. Let $F_\infty = \prod_{v \mid \infty} F_v$ and let $\mathbb{A}_F = F_\infty \times \mathbb{A}_{F,f}$ denote the ring of adèles of F. Throughout the paper, we fix an unramified additive character $\psi = \bigotimes \psi_v$ of $F \setminus \mathbb{A}_F$ whose local conductor at every finite place v is \mathfrak{o}_v , i.e., ψ_v is trivial on \mathfrak{o}_v but non-trivial on \mathfrak{p}_v^{-1} . We write ψ_∞ to denote the character $\prod_{v \mid \infty} \psi_v$ of F_∞ .

For any n > 1, let $B_n = T_n U_n$ be the Borel subgroup of GL_n consisting of upper triangular matrices; let $P'_n \supset B_n$ be the standard parabolic subgroup of type (n-1,1) with Levi decomposition $P'_n = M_n N_n$. Then $M_n \cong GL_{n-1} \times GL_1$ and

$$N_n = \left\{ \begin{pmatrix} I_{n-1} & * \\ & 1 \end{pmatrix} \right\}.$$

If R is any F-algebra and H is any algebraic F-group, we write H(R) to denote the corresponding group of R-points. Let $P_n(R) \subset P'_n(R)$ denote the mirabolic subgroup consisting of matrices whose last row is of the form $(0, \ldots, 0, 1)$, i.e.,

$$P_n(R) = \left\{ \begin{pmatrix} h & y \\ & 1 \end{pmatrix} : h \in GL_{n-1}(R), y \in R^{n-1} \right\} \cong GL_{n-1}(R) \ltimes N_n(R).$$

Let w_n denote the long Weyl element in $GL_n(R)$. For $1 \le m \le n-1$, let α_m denote the permutation matrix

$$\alpha_m = \begin{pmatrix} 1 & & \\ I_m & & \\ & & I_{n-m-1} \end{pmatrix}.$$

Note that $\alpha_{n-1} = w_n({w_{n-1} \choose 1})$. The character

$$u \mapsto \psi\left(\sum u_{i,i+1}\right) \quad \text{for } u \in U_n(\mathbb{A})$$

defines a generic character of $U_n(F) \setminus U_n(\mathbb{A}_F)$, and by abuse of notation we continue to denote this character by ψ . Similarly, for each v, we also obtain a generic character ψ_v of $U_n(F_v)$. Further, for any algebraic subgroup $V \subseteq U_n$, ψ and ψ_v define characters of $V(\mathbb{A}_F)$ and $V(F_v)$, respectively, via restriction. In particular, we may consider the character $\psi|_{N_n(\mathbb{A}_F)}$; its stabilizer in $M_n(\mathbb{A}_F)$ is then $P_{n-1}(\mathbb{A}_F)$, where we regard P_{n-1} as a subgroup of M_n via $h \mapsto \binom{h}{1}$.

For each $v < \infty$, we will consider certain compact open subgroups of $GL_n(F_v)$; namely, let $K_v = GL_n(\mathfrak{o}_v)$, and for any integer $m \geq 0$, set

$$K_{1,v}(\mathfrak{p}_v^m) = \left\{ g \in \operatorname{GL}_n(\mathfrak{o}_v) : g \equiv \begin{pmatrix} & * \\ * & \vdots \\ 0 \cdots 0 & 1 \end{pmatrix} (\operatorname{mod} \mathfrak{p}_v^m) \right\},$$

$$K_{0,v}(\mathfrak{p}_v^m) = \left\{ g \in \operatorname{GL}_n(\mathfrak{o}_v) : g \equiv \begin{pmatrix} & * \\ * & \vdots \\ 0 \cdots 0 & * \end{pmatrix} (\operatorname{mod} \mathfrak{p}_v^m) \right\},$$

so that $K_{1,v}(\mathfrak{p}_v^m)$ is a normal subgroup of $K_{0,v}(\mathfrak{p}_v^m)$, with quotient $K_{0,v}(\mathfrak{p}_v^m)/K_{1,v}(\mathfrak{p}_v^m) \cong (\mathfrak{o}_v/\mathfrak{p}_v^m)^{\times}$. Let $K_f = \prod_{v < \infty} K_v$, and for an integral ideal \mathfrak{a} of F, set

$$K_i(\mathfrak{a}) = \prod_{v < \infty} K_{i,v}(\mathfrak{p}_v^{m_v}) \quad \text{for } i = 0, 1,$$

where m_v are the unique non-negative integers such that $\mathfrak{a} = \prod_v (\mathfrak{p}_v \cap \mathfrak{o}_F)^{m_v}$. Then $K_1(\mathfrak{a}) \subseteq K_0(\mathfrak{a}) \subseteq K_f$ are compact open subgroups of $GL_n(\mathbb{A}_{F,f})$. We may also form the corresponding principal congruence subgroups of $GL_n(F_{\infty})$, embedded diagonally, namely,

$$\Gamma_i(\mathfrak{a}) = \{ \gamma \in \mathrm{GL}_n(F) : \gamma_f \in K_i(\mathfrak{a}) \} \subset \mathrm{GL}_n(F_\infty) \quad \text{for } i = 0, 1,$$

where γ_f denotes the image of γ in $GL_n(\mathbb{A}_{F,f})$.

From strong approximation for GL_n , it follows that the set $GL_n(F) \setminus GL_n(\mathbb{A}_F) / GL_n(F_{\infty}) K_1(\mathfrak{a})$ of double cosets is finite of cardinality h, where h is the class number of F. Let us write

(2.1)
$$\operatorname{GL}_n(\mathbb{A}_F) = \coprod_{j=1}^h \operatorname{GL}_n(F)g_j \operatorname{GL}_n(F_\infty)K_1(\mathfrak{a}),$$

where each $g_i \in GL_n(\mathbb{A}_{F,f})$. Then

(2.2)
$$\operatorname{GL}_n(F) \backslash \operatorname{GL}_n(\mathbb{A}_F) / K_1(\mathfrak{a}) \cong \coprod_{j=1}^h \Gamma_{1,j}(\mathfrak{a}) \backslash \operatorname{GL}_n(F_{\infty}),$$

where $\Gamma_{1,j}(\mathfrak{a}) = \{ \gamma \in \mathrm{GL}_n(F) : \gamma_f \in g_j K_1(\mathfrak{a}) g_j^{-1} \} \subset \mathrm{GL}_n(F_{\infty}),$ embedded diagonally. Replacing $K_1(\mathfrak{a})$ by $K_0(\mathfrak{a})$ in this definition, we get the corresponding groups $\Gamma_{0,j}(\mathfrak{a})$.

Before proceeding further, let us introduce our choice of measures. For each place v of F and $n \geq 1$, we normalize the Haar measure on $\operatorname{GL}_n(F_v)$ and K_v so that $\operatorname{vol}(K_v) = 1$. We fix the Haar measure on $\operatorname{U}_n(F_v)$ for which $\operatorname{vol}(\operatorname{U}_n(F_v) \cap K_v) = 1$. From these measures, we obtain a right-invariant measure on $\operatorname{U}_n(F_v) \setminus \operatorname{GL}_n(F_v)$. Globally, on $\operatorname{GL}_n(\mathbb{A}_F)$ and $\operatorname{U}_n(\mathbb{A}_F)$, respectively, we take the corresponding product measure. On the compact quotient $\operatorname{U}(F) \setminus \operatorname{U}_n(\mathbb{A}_F)$, we choose the

right-invariant measure of unit volume. Since $U_2(\mathbb{A}_F) \cong \mathbb{A}_F$ and $\mathbb{A}_F = F + F_{\infty} + \prod_{v < \infty} \mathfrak{o}_v$, it follows from our normalization that $\operatorname{vol}(\mathfrak{o}_F \backslash F_{\infty}) = 1$.

Suppose (π, V_{π}) is an irreducible admissible representation of $\operatorname{GL}_n(\mathbb{A}_F)$ (not necessarily automorphic) whose central character ω_{π} is an idèle class character. We write $\pi \cong \bigotimes \pi_v$ as a restricted tensor product with respect to a distinguished set of spherical vectors, where for each finite v, π_v is an irreducible admissible representation of $\operatorname{GL}_n(F_v)$ on a complex vector space V_{π_v} , and for each $v \mid \infty, \pi_v$ is an irreducible admissible Harish-Chandra module. However, for $v \mid \infty$, we can pass to the canonical completion of π_v in the sense of Casselman and Wallach (see [12]) to obtain a smooth representation of moderate growth of $\operatorname{GL}_n(F_v)$. By abuse of notation we continue to write (π_v, V_{π_v}) to denote the canonical completion at an archimedean place v and let $(\pi_\infty, V_{\pi_\infty})$ denote the topological tensor product of these representations. Thus we may (and we will) take V_{π} to be the restricted tensor product $V_{\pi_\infty} \otimes \bigotimes_{v < \infty} V_{\pi_v}$ and π the corresponding representation of $\operatorname{GL}_n(\mathbb{A}_F)$.

Now with $\pi\cong \bigotimes \pi_v$ as in the previous paragraph, let us further assume that each π_v is generic, meaning there is a non-zero linear form (continuous when $v\mid \infty$) $\lambda_v:V_{\pi_v}\to \mathbb{C}$ satisfying $\lambda_v(\pi_v(n)w)=\psi_v(n)\lambda_v(w)$ for $w\in V_{\pi_v}, n\in \mathrm{N}_n(F_v)$. Let $\mathcal{W}(\pi_v,\psi_v)$ denote the Whittaker model of π_v , viz. the space of functions on $\mathrm{GL}_n(F)$ defined by $g\mapsto \lambda_v(\pi_v(g)w)$ for fixed $w\in V_{\pi_v}$. By taking the tensor product of λ_{π_v} for $v\mid \infty$, we can also form the space $\mathcal{W}(\pi_\infty,\psi_\infty)$. We note that for $v<\infty$ where π_v is unramified, there is a unique vector $W^0_{\pi_v}\in \mathcal{W}(\pi_v,\psi_v)$ that is fixed under K_v and takes the value 1 at the identity. The global Whittaker model $\mathcal{W}(\pi,\psi)$ of π is the space spanned by the functions $W_\infty\prod_{v<\infty}W_v$ with $W_\infty\in\mathcal{W}(\pi_\infty,\psi_\infty)$, $W_v\in\mathcal{W}(\pi_v,\psi_v)$ and $W_v=W^0_{\pi_v}$ for almost all v. For every $\eta\in V_\pi$ we denote by W_η the corresponding element of $\mathcal{W}(\pi,\psi)$.

Following [6, 14], for each $\eta \in V_{\pi}$ we set

(2.3)
$$U_{\eta}(g) = \sum_{\gamma \in U_n(F) \backslash P_n(F)} W_{\eta}(\gamma g),$$

which can also be written as

$$U_{\eta}(g) = \sum_{\gamma \in U_{n-1}(F) \backslash \operatorname{GL}_{n-1}(F)} W_{\eta} \left(\begin{pmatrix} \gamma & \\ & 1 \end{pmatrix} g \right).$$

It is shown in [14, Section 12] that this sum converges absolutely and uniformly on compact subsets to a continuous function on $GL_n(\mathbb{A}_F)$, and that it is cuspidal along the unipotent radical of any standard maximal parabolic subgroup of GL_n . Further, U_η is left invariant under both $P_n(F)$ and the center $Z_n(F)$. For $W \in \mathcal{W}(\pi, \psi)$, let $\widetilde{W}(g) = W(w_n^t g^{-1}) \in \mathcal{W}(\tilde{\pi}, \psi^{-1})$. For each place v, if $W_v \in \mathcal{W}(\pi_v, \psi_v)$, we similarly define $\widetilde{W}_v \in \mathcal{W}(\pi_v, \psi_v^{-1})$. Let V_η (cf. [6]) denote the function

$$V_{\eta}(g) = \sum_{\gamma \in \mathrm{U}'_n(F) \backslash \mathrm{Q}_n(F)} W_{\eta}(\alpha_{n-1}\gamma g),$$

where $Q_n = \overline{P}_n$ is the opposite mirabolic subgroup and $U'_n = \alpha_{n-1}^{-1} U_n \alpha_{n-1}$. If we put

$$\widetilde{U}_{\eta}(g) = \sum_{\gamma \in U_{n-1}(F) \backslash \operatorname{GL}_{n-1}(F)} \widetilde{W}_{\eta} \begin{pmatrix} \gamma & \\ & 1 \end{pmatrix} g$$
.

Then as explained in [4], one has the equality $\widetilde{U}_{\eta}(^tg^{-1}) = V_{\eta}(g)$. The following is a basic result of Cogdell and Piatetski-Shapiro [6, 7]:

Proposition 2.1. Suppose π is as above with automorphic central character ω_{π} . Then π is cuspidal automorphic if and only if $U_{\eta} = V_{\eta}$ for all $\eta \in V_{\pi}$.

For m < n, let $Y = Y_{n,m}$ be the unipotent radical of the standard parabolic subgroup of GL_n of type $(m+1,1,\ldots,1)$. (In this notation, $N_n = Y_{n,n-2}$.) For a function f on $GL_n(\mathbb{A}_F)$ that is left

invariant under Y(F), we set

(2.4)
$$\mathbb{P}_{m}^{n}(f)(g) = \int_{Y(F)\backslash Y(\mathbb{A}_{F})} f(yg)\overline{\psi(y)} \, dy,$$

where dy is the Haar measure on $Y(\mathbb{A}_F)$ for which the quotient $Y(F)\backslash Y(\mathbb{A}_F)$ has unit volume. In particular, we can consider $\mathbb{P}_m^n(U_\eta)(g)$, with the following Fourier series expansion:

(2.5)
$$\mathbb{P}_{m}^{n}(U_{\eta})(g) = \sum_{\gamma \in \mathcal{U}_{m+1}(F) \backslash \mathcal{P}_{m+1}(F)} W_{\eta} \begin{pmatrix} \gamma & \\ & I_{n-m-1} \end{pmatrix} g$$

$$= \sum_{\gamma \in \mathcal{U}_{m}(F) \backslash \operatorname{GL}_{m}(F)} W_{\eta} \begin{pmatrix} \gamma & \\ & I_{n-m} \end{pmatrix} g .$$

Viewed as a function on $P_{m+1}(\mathbb{A}_F)$, $\mathbb{P}_m^n(U_\eta)(g)$ is left invariant under $P_{m+1}(F)$ and is cuspidal in the sense that all the relevant unipotent integrals are zero. Observe that \mathbb{P}_{n-1}^n is the identity.

Suppose τ is an automorphic subrepresentation of $GL_m(\mathbb{A}_F)$, meaning V_{τ} is a subspace of the space of automorphic forms on $GL_m(\mathbb{A}_F)$. Let ϕ be an automorphic form in V_{τ} and set

$$(2.6) I(s; U_{\eta}, \phi) = \int_{\mathrm{GL}_m(F)\backslash \mathrm{GL}_m(\mathbb{A}_F)} \mathbb{P}_m^n(U_{\eta}) \begin{pmatrix} h \\ I_{n-m} \end{pmatrix} \phi(h) \|\det h\|^{s-\frac{n-m}{2}} dh.$$

This integral is absolutely convergent for all s, as we are integrating a cusp form against an automorphic form. For $\Re(s) \gg 1$ we can replace $\mathbb{P}^n_m(U_\eta)$ by its Fourier expansion and unfold, to get

$$I(s; U_{\eta}, \phi) = \int_{U_m(\mathbb{A}_E) \backslash \operatorname{GL}_m(\mathbb{A}_E)} W_{\eta} \begin{pmatrix} h \\ I_{n-m} \end{pmatrix} W_{\phi}(h) \| \det h \|^{s - \frac{n-m}{2}} dh,$$

where W_{ϕ} is the Whittaker on $GL_m(\mathbb{A}_F)$ associated to ϕ with respect to ψ^{-1} , i.e.

$$W_{\phi}(h) = \int_{\mathrm{U}_m(F)\backslash \mathrm{U}_m(\mathbb{A}_F)} \phi(uh)\psi(u) \, du.$$

Now, if η (resp. ϕ) corresponds to a pure tensor under $\pi \cong \bigotimes \pi_v$ (resp. $\tau \cong \bigotimes \tau_v$), then by the uniqueness of the Whittaker model we have

$$W_{\eta}(g) = \prod_{v} W_{\eta_v}(g), \quad W_{\phi}(g) = \prod_{v} W_{\phi_v}(g_v),$$

and the integral now factors as

$$I(s; U_{\eta}, \phi) = \prod_{v} \Psi_{v}(s; W_{\eta_{v}}, W_{\phi_{v}}),$$

where the local integrals are given by

$$\Psi_v(s; W_{\eta_v}, W_{\phi_v}) = \int_{\mathrm{U}_m(F_v) \backslash \operatorname{GL}_m(F_v)} W_{\eta_v} \begin{pmatrix} h_v & \\ & I_{n-m} \end{pmatrix} W_{\phi_v}(h_v) \, \| \det h_v \|_v^{s - \frac{n-m}{2}} \, dh_v.$$

It follows from the Rankin-Selberg theory of local factors that

$$E_v(s) := \frac{\Psi_v(s; W_{\eta_v}, W_{\phi_v})}{L(s, \pi_v \boxtimes \tau_v)}$$

is entire for all v [11, 12]. If v is non-archimedean then $E_v(s) \in \mathbb{C}[q_v^s, q_v^{-s}]$, and $E_v(s) = 1$ for almost all finite v. Setting $E(s) = \prod_v E_v(s)$, for pure tensors η and ϕ as above one has

$$I(s; U_{\eta}, \phi) = E(s) \prod_{v} L(s, \pi_{v} \boxtimes \tau_{v}).$$

One also has the analogous integrals involving the function V_{η} . Namely, put $\alpha'_{m} = \alpha_{n-1}^{-1}\alpha_{m}$, $1 \leq m \leq n-1$, and consider the function (cf. [8])

$$V_n^m(g) = V_\eta(\alpha_m'g)$$
 for $g \in GL_n(\mathbb{A}_F)$.

This function is clearly left invariant under $\alpha'_m^{-1}Q_n(F)\alpha'_m$ which contains the group $Y_{n,m}(F)$. Hence $\mathbb{P}^n_m(V^m_n)$ is well defined, and we put

$$(2.7) I(s; V_{\eta}, \phi) = \int_{\mathrm{GL}_m(F)\backslash \mathrm{GL}_m(\mathbb{A}_F)} \mathbb{P}_m^n(V_{\eta}^m) \begin{pmatrix} h \\ I_{n-m} \end{pmatrix} \phi(h) \|\det h\|^{s-\frac{n-m}{2}} dh.$$

As explained in loc. cit., this integral converges for $-\Re(s) \gg 1$ and unfolds to give

$$I(s; V_{\eta}, \phi) = \prod_{v} \widetilde{\Psi}_{v}(1 - s; \rho(w_{n,m})\widetilde{W}_{\eta_{v}}, \widetilde{W}_{\phi_{v}}),$$

where ρ denotes right translation, $w_{n,m} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, and

$$\widetilde{\Psi}_{v}(s; W, W') = \int_{\mathrm{U}_{m}(F_{v})\backslash \mathrm{GL}_{m}(F_{v})} \int_{M_{n-m-1,m}(F_{v})} W \begin{pmatrix} h \\ x & I_{n-m-1} \\ & 1 \end{pmatrix} W'(h) \|\det h\|_{v}^{s-\frac{n-m}{2}} dx dh.$$

Further, the local ϵ -factor $\epsilon(s, \pi_v \boxtimes \tau_v, \psi_v)$ is defined so that the following local functional equation [8, Theorem 3.2] holds:

(2.8)
$$\frac{\widetilde{\Psi}(1-s;\rho(w_{n,m})\widetilde{W}_{\eta_v},\widetilde{W}_{\phi_v})}{L(1-s,\widetilde{\pi}_v\boxtimes\widetilde{\tau}_v)} = \omega_{\tau_v}(-1)^{n-1}\epsilon(s,\pi_v\boxtimes\tau_v,\psi_v)\frac{\Psi_v(s;W_{\eta_v},W_{\phi_v})}{L(s,\pi_v\boxtimes\tau_v)}.$$

If $\tau = \mathbf{1}$ is the trivial representation of $\mathrm{GL}_1(\mathbb{A}_F)$, then for each v, we write $L(s, \pi_v)$ and $\epsilon(s, \pi_v, \psi_v)$ to denote $L(s, \pi_v \times \mathbf{1}_v)$ and $\epsilon(s, \pi_v \times \mathbf{1}_v, \psi_v)$, respectively. These are known to be the same as the standard local factors of Godement–Jacquet attached to the representation π_v and the character ψ_v .

With $\pi \cong \bigotimes \pi_v$ as above, let us now recall the notion of the essential vector of π_v . For each finite v, according to [10] (see also [13]), there is a unique positive integer $m(\pi_v)$ such that the space of $K_1(\mathfrak{p}_v^{m(\pi_v)})$ -fixed vectors is of dimension 1. Further, as mentioned in the introduction, by loc. cit., there is a unique vector ξ_v^0 in this space, called the essential vector, with the associated essential function $W_{\xi_v^0} = W_{\pi_v}$ in $\mathcal{W}(\pi_v, \psi_v)$ satisfying the condition $W_{\xi_v^0}(g^h_1) = W_{\xi_v^0}(g^h_1)$ for all $h \in \mathrm{GL}_{n-1}(\mathfrak{o}_v)$ and $g \in \mathrm{GL}_{n-1}(F_v)$. For $v < \infty$ and any unramified representation τ_v of $\mathrm{GL}_m(F_v)$, let $W_{\tau_v}^{0,\psi_v} \in W(\tau_v,\psi_v)$ denote the normalized spherical function (cf. [13, p. 2]). If $m(\pi_v) = 0$ then by uniqueness of essential functions one has the equality $W_{\xi_v^0} = W_{\pi_v}^{0,\psi_v}$ in $W(\pi_v,\psi_v)$. The integral ideal $\mathfrak{n} = \prod_{v < \infty} (\mathfrak{p}_v \cap \mathfrak{o}_F)^{m(\pi_v)} \subseteq \mathfrak{o}_F$ is called the conductor of π . We write \mathfrak{n} to denote a finite idèle corresponding to \mathfrak{n} . In fact, we may fix \mathfrak{n} so that $\mathfrak{n}_v = \varpi_v^{m(\pi_v)}$ for all finite v. The integer $m(\pi_v)$ can also be characterized as the degree of the monomial in the local ϵ -factor $\epsilon(s,\pi_v,\psi_v)$ [10]; we write

(2.9)
$$\epsilon(s, \pi_v, \psi_v) = \epsilon(\pi_v, \psi_v) q_v^{m(\pi_v)(\frac{1}{2} - s)}.$$

A crucial property of the conductor is that $K_0(\mathfrak{n})$ acts on the space of $K_1(\mathfrak{n})$ -fixed vectors via the central character $\omega = \omega_{\pi}$ (cf. [6, Section 8]). Precisely, for $g_v = (g_{i,j}) \in K_{0,v}(\mathfrak{p}_v^{m(\pi_v)})$, define

$$\chi_{\pi_v}(g_v) = \begin{cases} 1 & \text{if } m(\pi_v) = 0, \\ \omega_v(g_{n,n}) & \text{if } m(\pi_v) > 0, \end{cases}$$

and put $\chi_{\pi} = \bigotimes_{v < \infty} \chi_{\pi_v}$. Let $\xi_f^0 = \bigotimes_{v < \infty} \xi_v^0$, which forms a basis for the space of $K_1(\mathfrak{n})$ -fixed vectors in V_{π} . Then it is shown in loc. cit. that χ_{π} is a character of $K_0(\mathfrak{n})$ trivial on $K_1(\mathfrak{n})$, and that for any finite v,

$$\pi_v(g)\xi_v^0 = \chi_{\pi_v}(g)\xi_v^0 \quad \text{for all } g \in K_{0,v}(\mathfrak{p}_v^{m(\pi_v)}).$$

For each j, χ_{π} also determines a character of $\Gamma_{0,j}(\mathfrak{n})$ that is trivial on $\Gamma_{1,j}(\mathfrak{n})$, which we continue to denote by χ_{π} .

We will require the next two lemmas later in §3.4. To that end, for any v and any ξ_v in the space of π_v , put

(2.10)
$$\tilde{\xi}_v = \epsilon(\pi_v, \psi_v)^{1-n} \pi_v \binom{\mathsf{n}_v I_{n-1}}{\mathsf{1}} \xi_v.$$

(Note that for $v \mid \infty$, $\mathsf{n}_v = 1$.) First, we identify the essential function $W_{\widetilde{\pi}_v} \in \mathcal{W}(\widetilde{\pi}_v, \psi_v^{-1})$ associated to the contragredient representation $\widetilde{\pi}_v$ in the following lemma.

Lemma 2.2. For $v < \infty$,

$$\widetilde{W}_{\tilde{\xi}_{v}^{0}}(g) = W_{\tilde{\pi}_{v}}(g).$$

Proof. It is a straightforward calculation to check that the function $g \mapsto \widetilde{W}_{\tilde{\xi}_{v}^{0}}(g)$ is right $K_{1,v}(\mathfrak{p}_{v}^{m(\pi_{v})})$ -invariant. Indeed one checks that

$$\widetilde{W}_{\tilde{\xi}_{v}^{0}}(gk) = \chi_{\pi_{v}}(k)^{-1}\widetilde{W}_{\tilde{\xi}_{v}^{0}}(g), \quad \text{for } k \in K_{0,v}(\mathfrak{p}_{v}^{m(\pi_{v})}).$$

Since the space of such functions in $W(\widetilde{\pi}_v, \psi_v^{-1})$ is 1-dimensional, there exists a constant c so that $\widetilde{W}_{\widetilde{\epsilon}\underline{0}}(g) = cW_{\widetilde{\pi}_v}(g)$.

It remains to show that c=1. For that, we apply the local functional equation (2.8) with $\eta_v = \tilde{\xi}_v^0$ and $W_{\phi_v} = W_{\tau_v}^{0,\overline{\psi}_v}$ the normalized spherical function (with respect to ψ_v^{-1}) corresponding to the unramified (unitary) principal series representation $\tau_v = \operatorname{Ind}(\chi_1, \ldots, \chi_{n-1})$ of $\operatorname{GL}_{n-1}(F_v)$. Here the χ_j are unramified characters of F_v^{\times} of the form $\chi_j(x) = \|x\|_v^{it_j}$ for $t_j \in \mathbb{R}$. With these choices, since $W_{\tilde{\xi}_v^0} = cW_{\tilde{\pi}_v}$ and $W_{\tilde{\pi}_v}$ is the essential function for the representation $\tilde{\pi}_v$, the left-hand side of (2.8) reduces to the following equality:

$$c = \epsilon(s, \pi_v \boxtimes \tau_v, \psi_v) \frac{\Psi_v(s; W_{\tilde{\xi}_v^0}, W_{\tau_v}^{0, \overline{\psi}_v})}{L(s, \pi_v \boxtimes \tau_v)}.$$

Through the change of variable $g \mapsto g(\mathsf{n}_v^{-1}I_{n-1})$ in the integral defining $\Psi_v(s; W_{\tilde{\xi}_v^0}, W_{\tau_v}^{0,\overline{\psi}_v})$, one checks that

$$\Psi_v(s; W_{\tilde{\epsilon}^0}, W_{\tau_v}^{0, \overline{\psi}_v}) = \omega_{\tau_v}^{-1}(\mathbf{n}_v) \|\mathbf{n}_v^{-1}\|_v^{s - \frac{1}{2}} \epsilon(\pi_v, \psi_v)^{1 - n} \Psi_v(s; W_{\xi_v^0}, W_{\tau_v}^{0, \overline{\psi}_v}).$$

On the other hand, according to (1.2), $\Psi_v(s; W_{\xi_v^0}, W_{\tau_v}^{0, \overline{\psi}_v}) = L(s, \pi_v \boxtimes \tau_v)$. We also have

$$\epsilon(s, \pi_v \boxtimes \tau_v, \psi_v) = \prod_{j=1}^{n-1} \epsilon(s + it_j, \pi_v, \psi_v)$$

and $\omega_{\tau_v}(\mathsf{n}_v) = q_v^{m(\pi_v)\sum_j it_j}$. Putting these together we obtain c = 1.

Let ξ be a pure tensor in the space of π such that $\xi_v = \xi_v^0$ for all $v < \infty$. We also define its dual counterpart; namely, for $W' = \widetilde{W}_{\tilde{\xi}_{\infty}} \times \prod_{v < \infty} W_{\tilde{\pi}_v} \in \mathcal{W}(\tilde{\pi}, \psi^{-1})$, let

$$U'_{\xi}(g) = \sum_{\gamma \in \mathcal{U}_{n-1}(F) \backslash \operatorname{GL}_{n-1}(F)} W'\left(\begin{pmatrix} \gamma & \\ & 1 \end{pmatrix} g\right).$$

Lemma 2.3. Suppose π is a cuspidal automorphic representation of $GL_n(\mathbb{A}_F)$ and ξ is a pure tensor in the space of π such that $\xi_v = \xi_v^0$ for all finite v. Then

$$U_{\xi}\begin{pmatrix} tg^{-1}\begin{pmatrix} \mathsf{n}I_{n-1} & \\ & 1 \end{pmatrix} \end{pmatrix} = \epsilon_{\pi}^{n-1}U'_{\xi}(g),$$

where $\epsilon_{\pi} = \prod_{v} \epsilon(\pi_{v}, \psi_{v})$.

Proof. Keeping the above notation, let $\xi' = \xi_{\infty} \otimes \bigotimes_{v < \infty} \tilde{\xi}_{v}^{0}$. Then

$$U_{\xi}\begin{pmatrix} tg^{-1}\begin{pmatrix} \mathsf{n}I_{n-1} & \\ & 1 \end{pmatrix} \end{pmatrix} = \prod_{v < \infty} \epsilon(\pi_v, \psi_v)^{n-1} \cdot U_{\xi'}(^tg^{-1})$$
$$= \prod_{v < \infty} \epsilon(\pi_v, \psi_v)^{n-1} \cdot \widetilde{U}_{\xi'}(g)$$
$$= \epsilon_{\pi}^{n-1} U_{\xi}'(g).$$

Here, the first equality is a consequence of the definition of $U_{\mathcal{E}'}$, the second equality follows from Proposition 2.1, and the last equality follows from Lemma 2.2.

We will also need a certain auxiliary function related to the local Rankin–Selberg L-factor $L(s, \pi_v \boxtimes \tau_v)$. In order to introduce this we pass to a more general setting and take π_v (resp. τ_v) to be an irreducible admissible representation of $\mathrm{GL}_n(F_v)$ (resp. $\mathrm{GL}_m(F_v)$), not necessarily the local component of a global representation. It is known that the local L-function $L(s, \pi_v)$ is of the form $P_{\pi_v}(q_v^{-s})^{-1}$, where $P_{\pi_v} \in \mathbb{C}[X]$ has degree at most n and satisfies $P_{\pi_v}(0) = 1$. We may then find n complex numbers $\{\alpha_{v,i}\}$ (allowing some of them to be zero) such that

$$L(s, \pi_v) = \prod_{i=1}^{n} (1 - \alpha_{v,i} q_v^{-s})^{-1}.$$

We call the set $\{\alpha_{v,i}\}$ the Langlands parameters of π_v ; if π_v is spherical, they are the usual Satake parameters. Let $\{\beta_{v,j}\}$ be the Langlands parameters of τ_v , and set

$$L(s, \pi_v \times \tau_v) = \prod_{i,j} (1 - \alpha_{v,i} \beta_{v,j} q_v^{-s})^{-1}.$$

Of course, $L(s, \pi_v \times \tau_v) = L(s, \pi_v \boxtimes \tau_v)$ if both π_v and τ_v are spherical.

In the following lemma, we explore the connection between $L(s, \pi_v \times \tau_v)$ and $L(s, \pi_v \boxtimes \tau_v)$ in more detail, but let us first recall the full classification of irreducible admissible representations of $\mathrm{GL}_n(F_v)$. Let \mathcal{A}_n denote the set of equivalence classes of such representations and put $\mathcal{A} = \bigcup \mathcal{A}_n$. The essentially square-integrable representations of $GL_n(F_v)$ have been classified by Bernstein and Zelevinsky; they are given as follows: If σ_v is an essentially square-integrable representation of $\mathrm{GL}_n(F_v)$, then there is divisor $a \mid n$ and a supercuspidal representation η_v of $\mathrm{GL}_a(F_v)$ such that if $b = \frac{n}{a}$ and Q is the standard (upper) parabolic subgroup of $GL_n(F_v)$ of type (a, \ldots, a) , then σ_v can be realized as the unique quotient of the (normalized) induced representation

$$\operatorname{Ind}_{Q}^{\operatorname{GL}_{n}(F_{v})}(\eta_{v},\eta_{v}\|\cdot\|_{v},\ldots,\eta_{v}\|\cdot\|_{v}^{b-1});$$

the integer a and the class of η_v are uniquely determined by σ_v . In short, σ_v is parametrized by b and η_v , and we denote this by $\sigma_v = \sigma_b(\eta_v)$; further, σ_v is square-integrable (also called "discrete series") if and only if the representation $\eta_v \| \cdot \|_v^{\frac{b-1}{2}}$ of $\mathrm{GL}_a(F_v)$ is unitary. Now, let P be an upper parabolic subgroup of $\mathrm{GL}_n(F_v)$ of type (n_1,\ldots,n_r) . For each $i=1,\ldots,r$,

let $\tau_{i,v}^0$ be a discrete series representation of $\mathrm{GL}_{n_i}(F_v)$. Let (s_1,\ldots,s_r) be a sequence of real numbers

satisfying $s_1 \ge \cdots \ge s_r$, and put $\tau_{i,v} = \tau_{i,v}^0 \otimes \|\cdot\|_v^{s_i}$ (an essentially square-integrable representation), for $i = 1, \ldots, r$. Then the induced representation

$$\xi_v = \operatorname{Ind}_P^{\operatorname{GL}_n(F_v)}(\tau_{1,v} \otimes \cdots \otimes \tau_{r,v})$$

is said to be a representation of $GL_n(F_v)$ of Langlands type. If $\tau_v \in \mathcal{A}_n$, then it is well known that it is uniquely representable as the quotient of an induced representation of Langlands type. We write $\tau_v = \tau_{1,v} \boxplus \cdots \boxplus \tau_{r,v}$ to denote this realization of τ_v . Thus one obtains a sum operation on the set \mathcal{A} [11, (9.5)]. It follows easily from the definition that $L(s, \pi_v \times \tau_v)$ is bi-additive:

$$L(s, \pi_v \times (\tau_v \boxplus \tau'_v)) = L(s, \pi_v \times \tau_v) L(s, \pi_v \times \tau'_v)$$

$$L(s, (\pi_v \boxplus \pi'_v) \times \tau_v) = L(s, \pi_v \times \tau_v) L(s, \pi'_v \times \tau_v)$$

for all $\pi_v, \pi'_v, \tau_v, \tau'_v \in \mathcal{A}$. The local factor $L(s, \pi_v \boxtimes \tau_v)$ is also bi-additive in the above sense, according to [11, (9.5) Theorem].

Lemma 2.4. Given $v < \infty$, suppose $\pi_v \in \mathcal{A}_n$ and $\tau_v \in \mathcal{A}_m$. Then

$$L(s, \pi_v \times \tau_v) = P(q_v^{-s})L(s, \pi_v \boxtimes \tau_v)$$

for some polynomial $P \in \mathbb{C}[X]$.

Proof. Since π_v is a sum of essentially square-integrable representations and both $L(s, \pi_v \times \tau_v)$ and $L(s, \pi_v \boxtimes \tau_v)$ are additive with respect to this sum, it suffices to prove the lemma for essentially square-integrable representations π_v . So for the remainder of the proof we assume π_v is an essentially square-integrable representation of the form $\pi_v = \sigma_b(\eta_v)$. We proceed by induction on m. If m = 1, then $\tau_v = \chi_v$ is a quasi-character of F_v^{\times} and $L(s, \pi_v \boxtimes \chi_v) = L(s, \pi_v \boxtimes \chi_v)$, where $\pi_v \boxtimes \chi_v$ is the representation of $GL_n(F_v)$ defined by $g \mapsto \pi_v(g)\chi_v(\det g)$. If χ_v is unramified, then

$$L(s, \pi_v \otimes \chi_v) = L(s, \pi_v \times \chi_v)$$

and consequently P=1; on the other hand if χ_v is ramified then $L(s, \pi_v \times \chi_v)=1$, and the assertion follows since $L(s, \pi_v \otimes \chi_v)^{-1}$ is a polynomial in q_v^{-s} .

We now assume m > 1. Suppose τ_v is an essentially square-integrable representation of $\mathrm{GL}_m(F_v)$, say $\tau_v = \sigma_{b'}(\eta_v')$, $\eta_v' \in \mathcal{A}_{a'}$ is supercuspidal, and a'b' = m. Then the standard L-factor $L(s, \tau_v)$ is given by $L(s, \tau_v) = L(s + b' - 1, \eta_v')$ (see [11]). Consequently, $L(s, \tau_v) = 1$ unless a' = 1 and $\eta_v' = \chi_v$ is an unramified quasi-character of F_v^{\times} . However, if $L(s, \tau_v) = 1$ then $L(s, \pi_v \times \tau_v) = 1$, and the assertion of the lemma follows. Hence we may assume $\tau_v = \sigma_m(\chi_v)$ for an unramified quasi-character χ_v of F_v^{\times} , in which case

(2.11)
$$L(s, \pi_v \times \tau_v) = L(s, \pi_v \otimes \chi_v \| \cdot \|_v^{m-1})$$
$$= L(s, \sigma_b(\eta_v) \otimes \chi_v \| \cdot \|_v^{m-1})$$
$$= L(s + m - 1 + b - 1, \eta_v \otimes \chi_v).$$

On the other hand, it also follows from [11, (8.2) Theorem] that

(2.12)
$$L(s, \pi_v \boxtimes \tau_v) = \begin{cases} \prod_{j=0}^{m-1} L(s+j+b-1, \eta_v \otimes \chi_v) & \text{if } m \le n, \\ \prod_{i=0}^{b-1} L(s+m-1+i, \eta_v \otimes \chi_v) & \text{if } m > n. \end{cases}$$

Now, from (2.11) and (2.12), one sees that the ratio $\frac{L(s,\pi_v \times \tau_v)}{L(s,\pi_v \boxtimes \tau_v)}$ is a polynomial in q_v^{-s} , thus proving the lemma when τ_v is an essentially square-integrable representation.

If τ_v is not essentially square-integrable, then as discussed above there is a partition $\sum_{i=1}^k m_i = m$, with each $m_i < m$, and essentially square-integrable representations $\tau_{i,v}$ of $\mathrm{GL}_{m_i}(F_v)$, $i = 1, \ldots, k$, so that $\tau_v = \coprod_i \tau_{i,v}$. We may then use additivity and apply the induction hypothesis to get the desired conclusion.

Corollary 2.5. If $L(s, \pi_v \boxtimes \tau_v) = 1$, then either $L(s, \pi_v) = 1$ or $L(s, \tau_v) = 1$.

Proof. If $L(s, \pi_v \boxtimes \tau_v) = 1$, then Lemma 2.4 implies that $L(s, \pi_v \times \tau_v)$ is a polynomial in q_v^{-s} and hence must be 1. This in turn implies the conclusion.

3.
$$GL_3 \times GL_2$$

We keep the notation of the previous section and take n=3, m=2. Thus π is an irreducible admissible generic representation of $\mathrm{GL}_3(\mathbb{A}_F)$ with conductor \mathfrak{n} and central character ω_{π} , and τ is an automorphic subrepresentation of $GL_2(\mathbb{A}_F)$ with central character ω_{τ} . We fix nonzero integral ideals $\mathfrak{q}' \subseteq \mathfrak{q}$ and consider only those τ of conductor \mathfrak{q} .

Choose ideals $\mathfrak{a}_i \subseteq \mathfrak{o}_F$ representing the ideal classes of F so that $\prod_i \mathfrak{a}_i$ is coprime to \mathfrak{q}' . Let a_j be a finite idèle such that $a_j = (a_j)$, and take $a_1 = 1$. For each j, put $h_j = \begin{pmatrix} a_j \\ 1 \end{pmatrix}$ and $g_j = \binom{h_j}{1} \in GL_3(\mathbb{A}_{F,f})$. Then $\{g_j\}$ is a valid set of representatives in (2.1) (with n=3). In what follows, we will also consider strong approximation for $GL_2(\mathbb{A}_F)$, i.e.

$$\operatorname{GL}_2(\mathbb{A}_F) = \coprod_{j=1}^h \operatorname{GL}_2(F)h_j \operatorname{GL}_2(F_\infty)K,$$

where the compact open subgroup $K \subset GL_2(\mathbb{A}_{F,f})$ is either $K_f = GL_2(\mathfrak{o}_F)$, $K_1(\mathfrak{q})$ or $K_0(\mathfrak{q})$. We then have the group $G_j = \Gamma_{1,j}(\mathfrak{o}_F) = \Gamma_{0,j}(\mathfrak{o}_F)$ along with the congruence subgroups $\Gamma_{0,j}(\mathfrak{q})$, $\Gamma_{1,j}(\mathfrak{q})$ and $\Gamma_i(q)$. Explicitly,

$$G_j = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, d \in \mathfrak{o}_F, c \in \mathfrak{a}_j^{-1}, b \in \mathfrak{a}_j, ad - bc \in \mathfrak{o}_F^{\times} \right\},\,$$

which is precisely the stabilizer (acting on the right) of the lattice $\mathfrak{o}_F \oplus \mathfrak{a}_i \subset F \oplus F$, and

$$\Gamma_{0,j}(\mathfrak{q}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G_j : c \in \mathfrak{q}\mathfrak{a}_j^{-1} \right\},
\Gamma_{1,j}(\mathfrak{q}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_{0,j}(\mathfrak{q}) : d - 1 \in \mathfrak{q} \right\},
\Gamma_j(\mathfrak{q}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_{1,j}(\mathfrak{q}) : a - 1 \in \mathfrak{q}, b \in \mathfrak{q}\mathfrak{a}_j \right\}.$$

Note that $\Gamma_j(\mathfrak{q}) \subseteq \Gamma_{1,j}(\mathfrak{q}) \subseteq \Gamma_{0,j}(\mathfrak{q}) \subseteq G_j$, and that $\Gamma_j(\mathfrak{q})$ is normal in G_j . For any $\alpha \in F \hookrightarrow F_{\infty}$ embedded diagonally, we form the adèle $(\alpha,0) \in F_{\infty} \times \mathbb{A}_{F,f}$ which we continue to denote by α . In this way, we view $\beta = (\beta_1, \beta_2) \in F^2$ as an element of \mathbb{A}^2_F . For $\xi_\infty \in V_{\pi_\infty}$, note that

$$h \mapsto U_{\xi_{\infty} \otimes \xi_f^0} \begin{pmatrix} h \\ 1 \end{pmatrix}, \text{ for } h \in GL_2(\mathbb{A}_F),$$

is a rapidly decreasing automorphic function on $GL_2(\mathbb{A}_F)$ (cf. [8, Theorem 1.1]) which allowed us to consider integrals of the type (2.6) while defining $\Lambda(s, \pi \boxtimes \tau)$. We now follow [2] in order to define additive twists (see loc. cit.) in the current context. To that end, we modify the above function by inserting a unipotent element. To be precise, with $\beta = (\beta_1, \beta_2)$ as above, consider the function

$$h \mapsto U_{\xi_{\infty} \otimes \xi_f^0} \begin{pmatrix} h & {}^t \beta \\ & 1 \end{pmatrix}, \text{ for } h \in GL_2(\mathbb{A}_F);$$

it is rapidly decreasing (by the gauge estimates in loc. cit.), but as one can easily check, it is not in general $GL_2(F)$ -invariant, i.e., not an automorphic function on $GL_2(\mathbb{A}_F)$.

3.1. The function $\Phi_{\xi_{\infty},j}$ and its Fourier expansion. For each $j=1,\ldots,h$, put

$$\Phi_{\xi_{\infty},j}(g) = U_{\xi_{\infty} \otimes \xi_f^0}((g,g_j)), \text{ for } g \in \mathrm{GL}_n(F_{\infty}),$$

and for $\beta_1 \in \mathfrak{aq}^{-1}$ and $\beta_2 \in \mathfrak{q}^{-1}$, consider the function

(3.1)
$$h \mapsto \Phi_{\xi_{\infty},j} \begin{pmatrix} h & {}^{t}\beta \\ 1 \end{pmatrix}, \quad \text{for } h \in \mathrm{GL}_{2}(F_{\infty}).$$

Our goal in this section is to derive its Fourier series expansion (see (3.5)), which will also reveal it to be (classical) automorphic form on $GL_2(F_{\infty})$. For $k \in \{1, \ldots, h\}$, choose distinct prime ideals \mathfrak{p}_k and \mathfrak{p}'_k such that $\mathfrak{p}_k \sim \mathfrak{q}'\mathfrak{p}'_k \sim \mathfrak{a}_k$ and \mathfrak{p}_k is coprime to $\mathfrak{q}'\mathfrak{a}_1 \cdots \mathfrak{a}_h$, and let α_k be a generator of the principal fractional ideal $\mathfrak{p}_k^{-1}\mathfrak{q}'\mathfrak{p}'_k$. Then for each fixed j, $\alpha_k\mathfrak{a}_j + \mathfrak{o}_F = \mathfrak{p}_k^{-1}$ runs through a set of representatives of the class group of F. Let p_k be a finite idèle such that $(p_k) = p_k$.

Lemma 3.1. Let R_1 , R_2 be sets of representatives for $F^{\times}/\mathfrak{o}_F^{\times}$. Then for any fixed j,

$$\left\{ \begin{pmatrix} \gamma_1 \gamma_2 & \\ & \gamma_1 \end{pmatrix} \begin{pmatrix} 1 & \\ \alpha_k & 1 \end{pmatrix} : \gamma_1 \in R_1, \gamma_2 \in R_2, 1 \le k \le h \right\}$$

is a set of representatives for $U_2(F) \setminus GL_2(F)/G_i$.

Proof. For $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2(F)$, let $I_j(M)$ denote the fractional ideal $c\mathfrak{a}_j + d\mathfrak{o}_F$. It is easy to see that $I_j(M)$ depends only on the coset of M in $U_2(F)\backslash \operatorname{GL}_2(F)/G_j$, i.e. $I_j(uMg)=I_j(M)$ for all $u\in U_2(F), g\in G_j$. For a given M there is a unique choice of $k\in\{1,\ldots,h\}$ and $\gamma_1\in R_1$ such that $I_j(M)=\gamma_1\mathfrak{p}_k^{-1}$. Thus, $I_j(\gamma_1^{-1}M)=\mathfrak{p}_k^{-1}=\alpha_k\mathfrak{a}_j+\mathfrak{o}_F$. Suppose that $\gamma_1^{-1}M$ has bottom row (c-d). Then $c\mathfrak{a}_j+d\mathfrak{o}_F=\alpha_k\mathfrak{a}_j+\mathfrak{o}_F$, which in turn implies

 $c\mathfrak{o}_F + d\mathfrak{a}_i^{-1} = \alpha_k\mathfrak{o}_F + \mathfrak{a}_i^{-1}$. Thus, we have

$$(3.2) \qquad (\alpha_k \quad 1) = \begin{pmatrix} c & d \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

for some $A \in \mathfrak{o}_F$, $B \in \mathfrak{a}_j$, $C \in \mathfrak{a}_j^{-1}$, $D \in \mathfrak{o}_F$. It follows that the determinant AD - BC is an element of \mathfrak{o}_F , but it need not be a unit. However, the choice of $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ in (3.2) is not unique. For any particular solution $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$, it is straightforward to see that $\begin{pmatrix} A' & B' \\ C' & D' \end{pmatrix}$ is another solution if and only if $\binom{A'}{C'} = \binom{A}{C} + s\binom{a}{-c}$ and $\binom{B'}{D'} = \binom{B}{D} + t\binom{-d}{c}$ for some $s \in \mathfrak{p}_k$ and $t \in \mathfrak{a}_j \mathfrak{p}_k$. Thus,

$$\det\begin{pmatrix} A' & B' \\ C' & D' \end{pmatrix} - \det\begin{pmatrix} A & B \\ C & D \end{pmatrix} = s \det\begin{pmatrix} d & B \\ -c & D \end{pmatrix} + t \det\begin{pmatrix} A & -d \\ C & c \end{pmatrix} = s + t\alpha_k,$$

so by appropriate choice of s and t we may adjust the determinant by any element of $\mathfrak{p}_k + \alpha_k \mathfrak{a}_i \mathfrak{p}_k =$

Let $\epsilon \in \mathfrak{o}_F^{\times}$ be the unique unit such that $\epsilon \gamma_1^{-2} \det M \in R_2$. We choose $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ in the above with $\det g = \epsilon$. Then $g \in G_j$ and $\gamma_1^{-1}Mg$ has bottom row $(\alpha_k - 1)$. Multiplying on the left by a suitable $u \in U_2(F)$, we can make it lower-triangular, i.e. $u\gamma_1^{-1}Mg = \begin{pmatrix} \gamma_2 \\ \alpha_k \end{pmatrix}$ for some $\gamma_2 \in F^{\times}$. In fact, evaluating the determinant, we have $\gamma_2 = \epsilon \gamma_1^{-2} \det M \in R_2$.

Thus, $uMg = \binom{\gamma_1 \gamma_2}{\gamma_1} \binom{1}{\alpha_k 1}$ is of the required form. Moreover, although the pair (u, g) is not unique in the above construction, it is clear that k, γ_1 and γ_2 are. This completes the proof. \square

Lemma 3.2. We have

$$W_{\xi_f^0}\left(\begin{pmatrix} \gamma_1\gamma_2\mathsf{a}_j & \\ & \gamma_1 \\ & & 1 \end{pmatrix}\begin{pmatrix} 1 \\ \mathsf{a}_j\alpha_k & 1 \\ & & 1 \end{pmatrix}\right) = \psi_{v_k}\big(\alpha_k^{-1}\gamma_2\big)W_{\xi_f^0}\begin{pmatrix} \gamma_1\gamma_2\mathsf{a}_j\mathsf{p}_k & \\ & \gamma_1\mathsf{p}_k^{-1} \\ & & 1 \end{pmatrix},$$

and both sides vanish unless $\gamma_1 \in \mathfrak{p}_k$, $\gamma_2 \in \mathfrak{a}_j^{-1} \mathfrak{p}_k^{-2}$.

Proof. Suppose $v \neq v_k$ is a finite place of F. Then the corresponding local factors agree on both sides since ξ_v^0 is fixed by matrices of the form $\begin{pmatrix} g \\ 1 \end{pmatrix}$ for $g \in \mathrm{GL}_2(\mathfrak{o}_v)$. Hence, it is enough to prove the local equality for $v = v_k$. To that end, let us write the v-component of $\mathsf{a}_j \alpha_k$ as u/ϖ_v for $u \in \mathfrak{o}_v^\times$. We have

(3.3)
$$\begin{pmatrix} 1 \\ u/\varpi_v & 1 \end{pmatrix} = \begin{pmatrix} 1 & u^{-1}\varpi_v \\ & 1 \end{pmatrix} \begin{pmatrix} \varpi_v \\ & \varpi_v^{-1} \end{pmatrix} \begin{pmatrix} -u^{-1} \\ u & \varpi_v \end{pmatrix},$$

and consequently

$$W_{\xi_v^0}\left(\begin{pmatrix} \gamma_1\gamma_2\mathsf{a}_j & & \\ & \gamma_1 & \\ & & 1 \end{pmatrix}\begin{pmatrix} 1 & & \\ u/\varpi_v & 1 & \\ & & 1 \end{pmatrix}\right) = \psi_v\big(\gamma_2\mathsf{a}_ju^{-1}\varpi_v\big)W_{\xi_v^0}\begin{pmatrix} \gamma_1\gamma_2\mathsf{a}_j\varpi_v & & \\ & & \gamma_1\varpi_v^{-1} & \\ & & & 1 \end{pmatrix}.$$

This concludes the proof of the first assertion. Next, for any finite place v of F and $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2(F_v)$, one has

$$W_{\xi_v^0}\begin{pmatrix} g \\ 1 \end{pmatrix} \neq 0 \Longrightarrow c, d \in \mathfrak{o}_v.$$

It follows from this that $\gamma_1 \in \mathfrak{p}_k$. On the other hand, for $x \in \mathfrak{o}_v$, note that

$$\begin{split} W_{\xi_v^0} \begin{pmatrix} \gamma_1 \gamma_2 \mathsf{a}_j \varpi_v \\ & \gamma_1 \varpi_v^{-1} \\ & 1 \end{pmatrix} &= W_{\xi_v^0} \begin{pmatrix} \begin{pmatrix} \gamma_1 \gamma_2 \mathsf{a}_j \varpi_v \\ & \gamma_1 \varpi_v^{-1} \\ & 1 \end{pmatrix} \begin{pmatrix} 1 & x \\ & 1 \\ & & 1 \end{pmatrix} \end{pmatrix} \\ &= \psi_v \big(\gamma_2 \mathsf{a}_j \varpi_v^2 x \big) W_{\xi_v^0} \begin{pmatrix} \gamma_1 \gamma_2 \mathsf{a}_j \varpi_v \\ & \gamma_1 \varpi_v^{-1} \\ & & 1 \end{pmatrix}. \end{split}$$

Thus, if $W_{\xi_v^0}\begin{pmatrix} \gamma_1 \gamma_2 \mathsf{a}_j \varpi_v \\ \gamma_1 \varpi_v^{-1} \\ 1 \end{pmatrix} \neq 0$ then $\gamma_2 \mathsf{a}_j \varpi_v^2 \in \mathfrak{o}_v$, or in other words (globally) $\gamma_2 \in \mathfrak{a}_j^{-1} \mathfrak{p}_k^{-2}$. \square

In what follows, we use the notation G^x to denote $x^{-1}Gx$. Now

$$\begin{split} \Phi_{\xi_{\infty},j} \begin{pmatrix} h & \beta \\ & 1 \end{pmatrix} &= \sum_{\gamma \in \mathrm{U}_2(F) \backslash \mathrm{GL}_2(F)} W_{\xi_{\infty}} \begin{pmatrix} \gamma_{\infty} \\ & 1 \end{pmatrix} \begin{pmatrix} h & \beta \\ & 1 \end{pmatrix} \end{pmatrix} W_{\xi_f^0} \begin{pmatrix} \gamma_f \\ & 1 \end{pmatrix} g_j \\ &= \sum_{\gamma \in \mathrm{U}_2(F) \backslash \mathrm{GL}_2(F)} \psi_{\infty} ((\gamma_{\infty}\beta)_2) W_{\xi_{\infty}} \begin{pmatrix} \gamma_{\infty}h \\ & 1 \end{pmatrix} W_{\xi_f^0} \begin{pmatrix} \gamma_f h_j \\ & 1 \end{pmatrix} \\ &= \sum_{\gamma \in \mathrm{U}_2(F) \backslash \mathrm{GL}_2(F) / G_j} W_{\xi_f^0} \begin{pmatrix} \gamma_f h_j \\ & 1 \end{pmatrix} \sum_{\eta \in \mathrm{U}_2(F) \backslash \mathrm{GG}_j \backslash G_j} \psi_{\infty} ((\gamma\eta\beta)_2) W_{\xi_{\infty}} \begin{pmatrix} \gamma\eta h \\ & 1 \end{pmatrix}, \end{split}$$

which by Lemmas 3.1 and 3.2 can be written as

$$(3.4) \sum_{k} \frac{1}{n_{jk}(\mathfrak{q}')} \sum_{\gamma_{1} \in R_{1}, \gamma_{2} \in R_{2}} \psi_{v_{k}} (\alpha_{k}^{-1} \gamma_{2}) W_{\xi_{f}^{0}} \begin{pmatrix} \gamma_{1} \gamma_{2} \mathsf{a}_{j} \mathsf{p}_{k} \\ \gamma_{1} \mathsf{p}_{k}^{-1} \\ 1 \end{pmatrix} \cdot \sum_{\eta \in \mathcal{U}_{2}(F)^{\ell_{k}} \cap \Gamma_{j}(\mathfrak{q}') \backslash G_{j}} \psi_{\infty} (\gamma_{1} (\ell_{k} \eta \beta)_{2}) W_{\xi_{\infty}} \begin{pmatrix} \gamma_{1} \gamma_{2} \\ \gamma_{1} \\ 1 \end{pmatrix} \begin{pmatrix} \ell_{k} \eta h \\ 1 \end{pmatrix} ,$$

where $\ell_k = \begin{pmatrix} 1 \\ \alpha_k \end{pmatrix}$ and $n_{jk}(\mathfrak{q}') = [U_2(F)^{\ell_k} \cap \Gamma_j(\mathfrak{q}') : U_2(F)^{\ell_k} \cap G_j]$. One can check that

$$U_2(F)^{\ell_k} \cap G_j = \{\ell_k^{-1}(1, x)\ell_k : x \in \mathfrak{a}_j\mathfrak{p}_k^2\}.$$

From this and Lemma 3.2, it follows that the final summand in (3.4), viewed as a function of η , is constant on left cosets of $U_2(F)^{\ell_k} \cap G_j$ in G_j , so (3.4) is well defined. Similarly, we have

$$U_2(F)^{\ell_k} \cap \Gamma_j(\mathfrak{q}') = \left\{ \ell_k^{-1} \begin{pmatrix} 1 & x \\ 1 & 1 \end{pmatrix} \ell_k : x \in \mathfrak{a}_j \mathfrak{q}' \mathfrak{p}_k^2 \right\}.$$

In particular, $n_{jk}(\mathfrak{q}') = N(\mathfrak{q}')$.

Since $\beta_1 \in \mathfrak{a}_j \mathfrak{q}^{-1}$ and $\beta_2 \in \mathfrak{q}^{-1}$, the function $\eta \mapsto \psi_{\infty}(\gamma_1(\ell_k \eta \beta)_2)$ is constant on left cosets of $\Gamma_j(\mathfrak{q}')$ in G_j . Thus (3.4) may be rewritten as

$$(3.5) \quad \sum_{k} \frac{1}{N(\mathfrak{q}')} \sum_{\gamma_{1} \in R_{1}, \gamma_{2} \in R_{2}} \psi_{v_{k}} (\alpha_{k}^{-1} \gamma_{2}) W_{\xi_{f}^{0}} \begin{pmatrix} \gamma_{1} \gamma_{2} \mathsf{a}_{j} \mathsf{p}_{k} & \gamma_{1} \mathsf{p}_{k}^{-1} \\ & \gamma_{1} \mathsf{p}_{k}^{-1} \end{pmatrix} \\ \cdot \sum_{\eta \in \Gamma_{j}(\mathfrak{q}') \backslash G_{j}} \psi_{\infty} (\gamma_{1} (\ell_{k} \eta \beta)_{2}) \sum_{\epsilon \in \mathrm{U}_{2}(F)^{\ell_{k}} \cap \Gamma_{j}(\mathfrak{q}') \backslash \Gamma_{j}(\mathfrak{q}')} W_{\xi_{\infty}} \begin{pmatrix} \gamma_{1} \gamma_{2} & \gamma_{1} \\ & \gamma_{1} \end{pmatrix} \begin{pmatrix} \ell_{k} \epsilon \eta h & 1 \end{pmatrix} .$$

In particular, since we are free to take $\mathfrak{q}' = \mathfrak{q}$, it follows that (3.1) is left invariant under $\Gamma_i(\mathfrak{q})$.

3.2. Additive twists. We continue with the above setup. Recall that at every finite place v we have the essential vector ξ_v^0 in the space of π_v with the associated essential function $W_{\xi_v^0}$ which satisfies

$$W_{\xi_v^0}egin{pmatrix}1&&&\&1&\&&1\end{pmatrix}=1.$$

Let $\varphi \in V_{\tau}$ be any decomposable vector which is locally the new vector φ_v^0 at any finite place v of F and put $\varphi_j(h) = \varphi(h, h_j), h \in \mathrm{GL}_2(F_{\infty})$, so that $\varphi_j \in \mathcal{A}(\Gamma_{0,j}(\mathfrak{q}) \setminus \mathrm{GL}_2(F_{\infty}), \omega_{\tau_{\infty}}, \chi_{\tau}^{-1})$. Here again the essential function $W_{\varphi_v^0}$ satisfies

$$W_{\varphi_v^0} \begin{pmatrix} 1 & \\ & 1 \end{pmatrix} = 1.$$

If τ_v is unramified then (1.2) with n=3 reads as

(3.6)
$$\int_{\mathrm{U}_2(F_v)\backslash \mathrm{GL}_2(F_v)} W_{\xi_v^0} \begin{pmatrix} g_v \\ 1 \end{pmatrix} W_{\varphi_v^0}(g_v) \| \det g_v \|_v^{s-\frac{1}{2}} dg_v = L(s, \pi_v \boxtimes \tau_v).$$

For each $j = 1, \ldots, h$, put

$$(3.7) \quad \Lambda_{j}(s,\pi,\tau,\xi_{\infty},\varphi_{\infty},\beta) = \frac{N(\mathfrak{a}_{j})^{\frac{1}{2}-s}}{[\Gamma_{1,j}(\mathfrak{q}):\Gamma_{j}(\mathfrak{q})]} \int_{\Gamma_{j}(\mathfrak{q})\backslash \operatorname{GL}_{2}(F_{\infty})} \Phi_{\xi_{\infty},j} \begin{pmatrix} h & t\beta \\ & 1 \end{pmatrix} \varphi_{j}(h) \|\det h\|_{\infty}^{s-\frac{1}{2}} dh.$$

In view of (3.5), this integral is well defined; we call it a (generalized) additive twist.

Lemma 3.3. For $x \in \mathfrak{a}_i$, $j = 1, \ldots, h$, we have

$$\Lambda_j(s, \pi, \tau, \xi_{\infty}, \varphi_{\infty}, (\beta_1, \beta_2)) = \Lambda_j(s, \pi, \tau, \xi_{\infty}, \varphi_{\infty}, (\beta_1 + x, \beta_2)).$$

Proof. Since $U_{\xi_{\infty} \otimes \xi_f^0}$ is invariant under $Z_3(F) P_3(F)$ and ξ_f^0 is $K_1(\mathfrak{n})$ -invariant, it follows that each $\Phi_{\xi_{\infty},j}$ is left invariant by

$$Z_3(F) \operatorname{P}_3(F) \cap \operatorname{GL}_3(F_\infty) g_j K_1(\mathfrak{n}) g_j^{-1}.$$

From this we see that

$$\Lambda_j(s, \pi, \tau, \xi_{\infty}, \varphi_{\infty}, (\beta_1 + x, \beta_2)) = \Lambda_j(s, \pi, \tau, \xi_{\infty}, \varphi_{\infty}, (\beta_1, \beta_2))$$

for all $x \in \mathfrak{a}_i$.

If $(\beta_1, \beta_2) = (0, 0)$ then strong approximation implies that

(3.8)
$$\sum_{j} \Lambda_{j}(s, \pi, \tau, \xi_{\infty}, \varphi_{\infty}, \beta) = \Lambda(s, \pi, \tau, \xi_{\infty}, \varphi_{\infty}),$$

where

$$\Lambda(s,\pi,\tau,\xi_{\infty},\varphi_{\infty}):=\frac{1}{\operatorname{vol}(K_{1}(\mathfrak{q}))}\int_{\operatorname{GL}_{2}(F)\backslash\operatorname{GL}_{2}(\mathbb{A}_{F})}U_{\xi_{\infty}\otimes\xi_{f}^{0}}\begin{pmatrix}h&\\&1\end{pmatrix}\varphi(h)\|\det h\|^{s-\frac{1}{2}}\,dh.$$

If, in addition, τ is unramified (so that $\mathfrak{q} = \mathfrak{o}_F$), then it follows from (3.6) that

$$\Lambda(s, \pi, \tau, \xi_{\infty}, \varphi_{\infty}) = \Psi_{\infty}(s, \xi_{\infty}, \varphi_{\infty}) L(s, \pi \boxtimes \tau) = \Psi_{\infty}(s, \xi_{\infty}, \varphi_{\infty}) L(s, \pi \times \tau),$$

where

$$L(s, \pi \boxtimes \tau) = \prod_{v < \infty} L(s, \pi_v \boxtimes \tau_v)$$
 and $L(s, \pi \times \tau) = \prod_{v < \infty} L(s, \pi_v \times \tau_v).$

We now state our main result, which is convenient to formulate in terms of $L(s, \pi \times \tau)$ rather than $L(s, \pi \boxtimes \tau)$.

Theorem 3.4. Let notation be as above, and set $\mathfrak{a} = \bigcap_{j=1}^h \mathfrak{a}_j$. Let \mathfrak{c} be the conductor of χ_{τ} , and choose integral ideals \mathfrak{q}_1 and \mathfrak{q}_2 satisfying $\mathfrak{c} \cap \prod_{\mathfrak{p} \mid \mathfrak{q}} \mathfrak{p} \subseteq \mathfrak{q}_2 \subseteq \mathfrak{c}$ and $\mathfrak{q} \subseteq \mathfrak{q}_1 \subseteq \prod_{\mathfrak{p} \mid \mathfrak{q}\mathfrak{q}_2^{-1}} \mathfrak{p}^{\operatorname{ord}_{\mathfrak{p}}(\mathfrak{q})}$. Then for any $\beta_2 \in \mathfrak{q}_2^{-1} \setminus \bigcup_{\mathfrak{p} \mid \mathfrak{q}_2} \mathfrak{p}\mathfrak{q}_2^{-1}$, we have

$$\tau_{\mathfrak{q}}(\chi_{\tau},\beta_{2})\Psi_{\infty}(s,\xi_{\infty},\varphi_{\infty})L(s,\pi\times\tau) = \frac{1}{N(\mathfrak{q}_{1})}\sum_{\beta_{1}\in\mathfrak{a}\mathfrak{q}_{1}^{-1}/\mathfrak{a}}\sum_{j=1}^{h}\Lambda_{j}(s,\pi,\tau,\xi_{\infty},\varphi_{\infty},(\beta_{1},\beta_{2})),$$

where

$$\Psi_{\infty}(s,\xi_{\infty},\varphi_{\infty}) = \int_{\mathrm{U}_{2}(F_{\infty})\backslash \mathrm{GL}_{2}(F_{\infty})} W_{\xi_{\infty}} \begin{pmatrix} h & \\ & 1 \end{pmatrix} W_{\varphi_{\infty}}(h) \|\det h\|_{\infty}^{s-\frac{1}{2}} dh$$

and

$$\tau_{\mathfrak{q}}(\chi_{\tau},\beta_2) = \sum_{d \in (\mathfrak{o}_F/\mathfrak{q})^{\times}} \chi_{\tau}(d) \psi_{\infty}(d\beta_2).$$

In particular, if $\mathfrak{q} = \mathfrak{c} \cap \prod_{\mathfrak{p} \mid \mathfrak{q}} \mathfrak{p}$, then with $\mathfrak{q}_2 = \mathfrak{q}$ we have

$$\Psi_{\infty}(s,\xi_{\infty},\varphi_{\infty})L(s,\pi\times\tau) = \tau_{\mathfrak{q}}(\chi_{\tau},\beta_{2})^{-1}\sum_{j=1}^{h}\Lambda_{j}(s,\pi,\tau,\xi_{\infty},\varphi_{\infty},(0,\beta_{2})).$$

Remark. The condition $\mathfrak{c} \cap \prod_{\mathfrak{p}|\mathfrak{q}} \mathfrak{p} \subseteq \mathfrak{q}_2 \subseteq \mathfrak{c}$ ensures that $\tau_{\mathfrak{q}}(\chi_{\tau}, \beta_2) \neq 0$, by [22, Corollary 1].

3.3. **Proof of Theorem 3.4.** Let ζ be a finite idèle with $\zeta_v \in \mathfrak{o}_v$ for all $v < \infty$, $\zeta_v \in \mathfrak{o}_v^{\times}$ whenever $\operatorname{ord}_v(\mathfrak{q}) = 0$, and $\zeta = 1$ if $\mathfrak{c} \neq \mathfrak{q}$. Set $\mathfrak{q}' = \mathfrak{q} \cap (\zeta)$ and $h'_j = \begin{pmatrix} \mathsf{a}_j \zeta \\ 1 \end{pmatrix}$. We will consider the more general integral (3.10)

$$\Lambda_{j}(s,\pi,\tau,\xi_{\infty},\varphi_{\infty},\beta,\zeta) := \frac{N(\mathfrak{a}_{j})^{\frac{1}{2}-s}}{\left[\Gamma_{1,j}(\mathfrak{q}):\Gamma_{j}(\mathfrak{q}')\right]} \int_{\Gamma_{j}(\mathfrak{q}')\backslash\operatorname{GL}_{2}(F_{\infty})} \Phi_{\xi_{\infty},j} \begin{pmatrix} h & t\beta \\ & 1 \end{pmatrix} \varphi(h,h'_{j}) \|\det h\|_{\infty}^{s-\frac{1}{2}} dh,$$

where $\beta_1 \in \mathfrak{a}_j \mathfrak{q}_1^{-1}$, $\beta_2 \in \mathfrak{q}_2^{-1} \setminus \bigcup_{\mathfrak{p} \mid \mathfrak{q}_2} \mathfrak{p} \mathfrak{q}_2^{-1}$.

First, substituting (3.5) into (3.10) and changing $h \mapsto \eta^{-1}h$, we get

$$(3.11) N(\mathfrak{a}_{j})^{\frac{1}{2}-s} \frac{N(\mathfrak{q}')^{-1}}{[\Gamma_{1,j}(\mathfrak{q}):\Gamma_{j}(\mathfrak{q}')]} \sum_{k} \sum_{\gamma_{1} \in R_{1}, \gamma_{2} \in R_{2}} \psi_{v_{k}} (\alpha_{k}^{-1}\gamma_{2}) W_{\xi_{f}^{0}} \begin{pmatrix} \gamma_{1}\gamma_{2} \mathsf{a}_{j} \mathsf{p}_{k} \\ \gamma_{1} \mathsf{p}_{k}^{-1} \end{pmatrix}$$

$$\cdot \sum_{\eta \in \Gamma_{j}(\mathfrak{q}') \backslash G_{j}} \psi_{\infty} (\gamma_{1}(\ell_{k}\eta\beta)_{2}) \int_{\Gamma_{j}(\mathfrak{q}') \backslash \operatorname{GL}_{2}(F_{\infty})} \sum_{\epsilon \in \operatorname{U}_{2}(F)^{\ell_{k}} \cap \Gamma_{j}(\mathfrak{q}') \backslash \Gamma_{j}(\mathfrak{q}')}$$

$$W_{\xi_{\infty}} \begin{pmatrix} \gamma_{1}\gamma_{2} & \\ \gamma_{1} & \\ 1 \end{pmatrix} \begin{pmatrix} \ell_{k}\epsilon h & \\ 1 \end{pmatrix} \varphi(\eta^{-1}h, h'_{j}) \| \det h \|_{\infty}^{s-\frac{1}{2}} dh.$$

Since $h \mapsto \varphi(\eta^{-1}h, h'_j)$ is left-invariant under $\Gamma_j(\mathfrak{q}')$, we may collapse the integral and the sum over ϵ , and make the change of variable $h \mapsto \ell_k^{-1}h$ to get

$$(3.12) N(\mathfrak{a}_{j})^{\frac{1}{2}-s} \frac{N(\mathfrak{q}')^{-1}}{[\Gamma_{1,j}(\mathfrak{q}):\Gamma_{j}(\mathfrak{q}')]} \sum_{k} \sum_{\gamma_{1} \in R_{1}, \gamma_{2} \in R_{2}} \psi_{v_{k}} (\alpha_{k}^{-1} \gamma_{2}) W_{\xi_{j}^{0}} \begin{pmatrix} \gamma_{1} \gamma_{2} \mathsf{a}_{j} \mathsf{p}_{k} \\ \gamma_{1} \mathsf{p}_{k}^{-1} \end{pmatrix}$$

$$\sum_{\eta \in \Gamma_{j}(\mathfrak{q}') \setminus G_{j}} \psi_{\infty} (\gamma_{1}(\ell_{k} \eta \beta)_{2}) \int_{U_{2}(F) \cap \Gamma_{j}(\mathfrak{q}')^{\ell_{k}^{-1}} \setminus GL_{2}(F_{\infty})} W_{\xi_{\infty}} \begin{pmatrix} \gamma_{1} \gamma_{2} \\ \gamma_{1} \\ 1 \end{pmatrix} \begin{pmatrix} h \\ 1 \end{pmatrix}$$

$$\cdot \varphi (\eta^{-1} \ell_{k}^{-1} h, h'_{j}) \| \det h \|_{\infty}^{s-\frac{1}{2}} dh.$$

Let us now consider the archimedean integral in (3.12). We change $h \mapsto \gamma_1^{-1}h$ to get

$$\omega_{\tau_{\infty}}^{-1}(\gamma_1)\|\gamma_1\|_{\infty}^{1-2s} \int_{\mathrm{U}_2(F)\cap\Gamma_j(\mathfrak{q}')^{\ell_k^{-1}}\backslash \mathrm{GL}_2(F_{\infty})} W_{\xi_{\infty}}\left(\begin{pmatrix} \gamma_2 & & \\ & 1 & \\ & & 1 \end{pmatrix} \begin{pmatrix} h & \\ & & 1 \end{pmatrix}\right) \varphi\left(\eta^{-1}\ell_k^{-1}h, h_j'\right)\|\det h\|_{\infty}^{s-\frac{1}{2}} dh.$$

Since $\varphi(\eta^{-1}\ell_k^{-1}h,h_j')=\varphi(h,\ell_k\eta h_j'),$ this becomes

(3.13)
$$\omega_{\tau_{\infty}}^{-1}(\gamma_{1}) \|\gamma_{1}^{2}\|_{\infty}^{\frac{1}{2}-s} \int_{\mathrm{U}_{2}(F_{\infty})\backslash \mathrm{GL}_{2}(F_{\infty})} W_{\xi_{\infty}} \begin{pmatrix} \gamma_{2} \\ 1 \end{pmatrix} h \\ 1 \end{pmatrix} W_{\varphi}'(h) \|\det h\|_{\infty}^{s-\frac{1}{2}} dh,$$

where

$$(3.14) W_{\varphi}'(h) = \int_{\mathrm{U}_2(F)\cap\Gamma_j(\mathfrak{q}')^{\ell_k^{-1}}\setminus\mathrm{U}_2(F_\infty)} \varphi(u(x)h, \ell_k \eta h_j') \psi_\infty(\gamma_2 x) \, du(x),$$

and u(x) denotes the upper unipotent matrix $\begin{pmatrix} 1 & x \\ 1 \end{pmatrix}$. Then it is clear that the function $h \mapsto W'_{\varphi}\left(\begin{pmatrix} \gamma_2^{-1} \\ 1 \end{pmatrix}h\right)$ belongs to the Whittaker model of τ_{∞} with respect to the character ψ_{∞}^{-1} . Hence, by the local multiplicity one theorem, there exists a number $a_{jk}^{\varphi}(\gamma_2, \eta)$ so that

(3.15)
$$W_{\varphi}'(h) = a_{jk}^{\varphi}(\gamma_2, \eta) W_{\varphi_{\infty}} \left(\begin{pmatrix} \gamma_2 \\ 1 \end{pmatrix} h \right), \text{ for all } h \in GL_2(F_{\infty}),$$

where $W_{\varphi_{\infty}}$ is the Whittaker function (with respect to ψ_{∞}^{-1}) associated to φ_{∞} . Consequently, (3.13) can now be written as

$$\omega_{\tau_{\infty}}^{-1}(\gamma_{1})a_{jk}^{\varphi}(\gamma_{2},\eta)\|\gamma_{1}^{2}\|_{\infty}^{\frac{1}{2}-s}\int_{\mathrm{U}_{2}(F_{\infty})\backslash\mathrm{GL}_{2}(F_{\infty})}W_{\xi_{\infty}}\begin{pmatrix} \gamma_{2} \\ 1 \end{pmatrix}h \\ 1 \end{pmatrix}W_{\varphi_{\infty}}\begin{pmatrix} \gamma_{2} \\ 1 \end{pmatrix}h \|\det h\|_{\infty}^{s-\frac{1}{2}}dh,$$

which after the change $h \mapsto \begin{pmatrix} \gamma_2^{-1} \\ 1 \end{pmatrix} h$ becomes

$$(3.16) \quad \omega_{\tau_{\infty}}^{-1}(\gamma_{1})a_{jk}^{\varphi}(\gamma_{2},\eta)\|\gamma_{1}^{2}\gamma_{2}\|_{\infty}^{\frac{1}{2}-s}\|\gamma_{2}\|_{\infty}\int_{\mathrm{U}_{2}(F_{\infty})\backslash\mathrm{GL}_{2}(F_{\infty})}W_{\xi_{\infty}}\begin{pmatrix}h\\&1\end{pmatrix}W_{\varphi_{\infty}}(h)\|\det h\|_{\infty}^{s-\frac{1}{2}}dh.$$

We note that this is precisely the local Rankin–Selberg integral for $\pi \times \tau$ at ∞ , viz. $\prod_{v|\infty} \Psi_v(s; W_{\xi_v}, W_{\varphi_v})$, in the notation of Section 2.

Lemma 3.5. We have

$$a_{jk}^{\varphi}(\gamma_2, \eta) = N(\mathfrak{a}_j \mathfrak{p}_k^2 \mathfrak{q}') W_{\varphi_f} \left(\begin{pmatrix} \gamma_2 & \\ & 1 \end{pmatrix} \ell_k \eta h_j' \right).$$

Proof. We have $\varphi(g) = \sum_{\gamma \in F^{\times}} W_{\varphi}(({}^{\gamma}_{1})g)$ for all $g \in GL_{2}(\mathbb{A}_{F})$, and thus

$$\varphi(h, \ell_k \eta h_j') = \sum_{\gamma \in F^{\times}} W_{\varphi_f} \begin{pmatrix} \gamma & \\ & 1 \end{pmatrix} \ell_k \eta h_j' W_{\varphi_{\infty}} \begin{pmatrix} \gamma & \\ & 1 \end{pmatrix} h.$$

Further, a calculation similar to that in the proof of Lemma 3.2 shows that if $W_{\varphi_f}({\gamma_1})\ell_k\eta h_j'$ is non-zero then $\gamma \in \mathfrak{p}_k^{-2}\mathfrak{q}_1^{-1}\mathfrak{a}_j^{-1}(\zeta^{-1})$. Plugging this into (3.14), we get

$$\begin{split} W_{\varphi}'(h) &= \int_{\mathrm{U}_2(F)\cap\Gamma_j(\mathfrak{q}')^{\ell_k^{-1}}\backslash\,\mathrm{U}_2(F_\infty)} \sum_{\gamma\in F^\times} W_{\varphi_f}\bigg(\binom{\gamma}{1}\,\ell_k\eta h_j'\bigg)\,W_{\varphi_\infty}\bigg(\binom{\gamma}{1}\,u(x)h\bigg)\,\psi_\infty(\gamma_2x)\,du(x) \\ &= \sum_{\gamma\in F^\times} W_{\varphi_f}\bigg(\binom{\gamma}{1}\,\ell_k\eta h_j'\bigg)\,W_{\varphi_\infty}\bigg(\binom{\gamma}{1}\,h\bigg)\int_{\mathrm{U}_2(F)\cap\Gamma_j(\mathfrak{q}')^{\ell_k^{-1}}\backslash\,\mathrm{U}_2(F_\infty)} \psi_\infty\big((\gamma_2-\gamma)x\big)\,du(x) \\ &= N(\mathfrak{a}_j\mathfrak{p}_k^2\mathfrak{q}')W_{\varphi_f}\bigg(\binom{\gamma_2}{1}\,\ell_k\eta h_j'\bigg)\,W_{\varphi_\infty}\bigg(\binom{\gamma_2}{1}\,h\bigg)\,. \end{split}$$

Here we have used the fact that the integral vanishes unless $\gamma = \gamma_2$, and that $\operatorname{vol}(U_2(F) \cap \Gamma_j(\mathfrak{q}')^{\ell_k^{-1}} \setminus U_2(F_\infty)) = N(\mathfrak{a}_j \mathfrak{p}_k^2 \mathfrak{q}')$, since the additive measure is normalized so that $\operatorname{vol}(F_\infty/\mathfrak{o}_F) = 1$. Our assertion now follows from (3.15).

By (3.16) and Lemma 3.5, we now have (3.17)

$$\begin{split} \Lambda_{j}(s,\pi,\tau,\xi_{\infty},\varphi_{\infty},\beta,\zeta) &= N(\mathfrak{a}_{j}\mathfrak{p}_{k}^{2})N(\mathfrak{a}_{j})^{\frac{1}{2}-s}\prod_{v\mid\infty}\Psi_{v}(s;W_{\xi_{v}},W_{\varphi_{v}}) \\ & \cdot \sum_{k}\sum_{\gamma_{1},\gamma_{2}\in F^{\times}/\mathfrak{o}_{F}^{\times}} \|\gamma_{1}^{2}\gamma_{2}\|_{\infty}^{\frac{1}{2}-s} \|\gamma_{2}\|_{\infty}\psi_{v_{k}}\big(\alpha_{k}^{-1}\gamma_{2}\big)W_{\xi_{f}^{0}}\begin{pmatrix} \gamma_{1}\gamma_{2}\mathsf{a}_{j}\mathsf{p}_{k} & & & \\ & \gamma_{1}\mathsf{p}_{k}^{-1} & & \\ & & 1 \end{pmatrix} \\ & \cdot \frac{1}{[\Gamma_{1,j}(\mathfrak{q}):\Gamma_{j}(\mathfrak{q}')]}\sum_{\eta\in\Gamma_{j}(\mathfrak{q}')\backslash G_{j}}\psi_{\infty}\big(\gamma_{1}(\ell_{k}\eta\beta)_{2}\big)W_{\varphi_{f}}\bigg(\begin{pmatrix} \gamma_{1}\gamma_{2} & & & \\ & \gamma_{1}\end{pmatrix}\ell_{k}\eta h_{j}'\bigg) \,. \end{split}$$

Next we study the average of the inner sum over $\beta_1 \in \mathfrak{a}_j \mathfrak{q}_1^{-1}/\mathfrak{a}_j$:

$$(3.18) \qquad \frac{1}{N(\mathfrak{q}_1)} \sum_{\beta_1 \in \mathfrak{a}_j \mathfrak{q}_1^{-1}/\mathfrak{a}_j} \frac{1}{[\Gamma_{1,j}(\mathfrak{q}) : \Gamma_j(\mathfrak{q}')]} \sum_{\eta \in \Gamma_j(\mathfrak{q}') \setminus G_j} \psi_{\infty} (\gamma_1(\ell_k \eta \beta)_2) W_{\varphi_f} \begin{pmatrix} \gamma_1 \gamma_2 & \\ & \gamma_1 \end{pmatrix} \ell_k \eta h_j' \end{pmatrix}.$$

If $\eta = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ then $(\ell_k \eta \beta)_2 = (A\alpha_k + C)\beta_1 + (B\alpha_k + D)\beta_2$. Since $\gamma_1 \alpha_k \in \mathfrak{q}$, it follows that $\psi_{\infty}(\gamma_1(\ell_k \eta \beta)_2) = \psi_{\infty}(\gamma_1(C\beta_1 + D\beta_2))$.

Next note that $W_{\varphi_f}\left(\begin{pmatrix} \gamma_1 \gamma_2 & \gamma_1 \end{pmatrix} \ell_k \eta h_j' \right)$ is invariant under replacing η by $\eta(\begin{smallmatrix} 1 & x \\ 1 \end{pmatrix}$ for $x \in (\zeta)\mathfrak{a}_j$, while $\psi_{\infty}\left(\gamma_1(\ell_k \eta \beta)_2\right)$ changes by a factor of $\psi_{\infty}(C\gamma_1\beta_2 x)$. Hence, averaging over $x \in (\zeta)\mathfrak{a}_j/\mathfrak{q}'\mathfrak{a}_j$, we see that (3.18) vanishes unless $C\gamma_1 \in \mathfrak{q}_2(\zeta^{-1})\mathfrak{a}_j^{-1}$. Further, computing the average over β_1 , we get 0 unless $C\gamma_1 \in \mathfrak{q}_1\mathfrak{a}_j^{-1}$. Therefore, we can restrict η to $\Gamma_{0,j}(\mathfrak{m})$, where $\mathfrak{m} = \mathfrak{q}_1(\gamma_1^{-1}) \cap \mathfrak{q}_2(\zeta^{-1})(\gamma_1^{-1}) \cap \mathfrak{o}_F$. Since $\Gamma_j(\mathfrak{q}')$ is normal in $\Gamma_{0,j}(\mathfrak{m})$, we may swap left and right cosets, so (3.18) becomes

$$(3.19) \qquad \frac{1}{\left[\Gamma_{1,j}(\mathfrak{q}):\Gamma_{j}(\mathfrak{q}')\right]} \sum_{\eta \in \Gamma_{0,j}(\mathfrak{m})/\Gamma_{j}(\mathfrak{q}')} \psi_{\infty}(\gamma_{1}\beta_{2}\eta_{22}) W_{\varphi_{f}}\left(\begin{pmatrix} \gamma_{1}\gamma_{2} & \\ & \gamma_{1} \end{pmatrix} \ell_{k}\eta h_{j}'\right),$$

where η_{22} denotes the lower-right entry of η .

Now set $\mathfrak{m}' = \mathfrak{q}_1 \cap \mathfrak{q}_2(\zeta^{-1}),$

$$X_0 = \left\{ \left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix} \right) \in \Gamma_{0,j}(\mathfrak{m}') : b \in (\zeta)\mathfrak{a}_j \right\}, \quad X_1 = \left\{ \left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix} \right) \in X_0 : d-1 \in \mathfrak{q} \right\}.$$

Then as a function of η , (3.19) is right invariant under X_1 and acts via the nebentypus character χ_{τ} under X_0 . Hence, (3.19) equals

$$(3.20) \qquad \frac{[X_1:\Gamma_j(\mathfrak{q}')]}{[\Gamma_{1,j}(\mathfrak{q}):\Gamma_j(\mathfrak{q}')]} \sum_{\mu \in \Gamma_{0,j}(\mathfrak{m})/X_0} W_{\varphi_f} \left(\begin{pmatrix} \gamma_1 \gamma_2 & \\ & \gamma_1 \end{pmatrix} \ell_k \mu h_j' \right) \sum_{\nu \in X_0/X_1} \chi_\tau(\nu) \psi_\infty(\gamma_1 \beta_2 \mu_{22} \nu_{22}).$$

Next, note that for a fixed γ_1 , the ideal $\gamma_1\mu_{22}\mathfrak{o}_F + \mathfrak{q}$ is independent of the choice of representative for μ . We will show that the sum over those μ for which $\gamma_1\mu_{22}\mathfrak{o}_F + \mathfrak{q} \neq \mathfrak{o}_F$ vanishes. First, if $\gamma_1\mu_{22}\mathfrak{o}_F + \mathfrak{c} \neq \mathfrak{o}_F$ then the Gauss sum

$$\sum_{\nu \in X_0/X_1} \chi_{\tau}(\nu) \psi_{\infty}(\gamma_1 \beta_2 \mu_{22} \nu_{22}) = \sum_{d \in (\mathfrak{o}_F/\mathfrak{q})^{\times}} \chi_{\tau}(d) \psi_{\infty}(\gamma_1 \beta_2 \mu_{22} d)$$

vanishes. Hence, we may assume that $(\zeta) = \mathfrak{o}_F$, so $\mathfrak{m}' = \mathfrak{q}$ and $X_0 = \Gamma_{0,j}(\mathfrak{q})$.

If $\gamma_1 \mathfrak{o}_F + \mathfrak{q} = \mathfrak{o}_F$ then $\mathfrak{m} = \mathfrak{q}$, whence $D\mathfrak{o}_F + \mathfrak{q} = \mathfrak{o}_F$, so we may assume that $\gamma_1 \mathfrak{o}_F \subseteq \mathfrak{p}$ for some prime ideal $\mathfrak{p} \supseteq \mathfrak{q}$. We split the coset representatives for μ as $g_1 g_2$, where $g_1 = \begin{pmatrix} A_1 & B_1 \\ C_1 & D_1 \end{pmatrix}$ runs through representatives for $\Gamma_{0,j}(\mathfrak{m})/\Gamma_{0,j}(\mathfrak{p}^{-1}\mathfrak{q})$ and $g_2 = \begin{pmatrix} A_2 & B_2 \\ C_2 & D_2 \end{pmatrix}$ runs through representatives for $\Gamma_{0,j}(\mathfrak{p}^{-1}\mathfrak{q})/\Gamma_{0,j}(\mathfrak{q})$. Then $\psi_{\infty}(\gamma_1 \mu_{22}\beta_2) = \psi_{\infty}(\gamma_1 D_1 D_2\beta_2)$. Defining

$$f_{\gamma_1,g_1}(g_2) = \sum_{d \in (\mathfrak{o}_F/\mathfrak{q}_2)^{\times}} \chi_{\tau}(d) \psi_{\infty}(\gamma_1 D_1 D_2 d\beta_2),$$

it is plain that $f_{\gamma_1,g_1}(kg_2k') = f_{\gamma_1,g_1}(g_2)\overline{\chi_{\tau}(k')}$ for any $k \in \Gamma_{1,j}(\mathfrak{p}^{-1}\mathfrak{q}), k' \in \Gamma_{0,j}(\mathfrak{q})$. As the following lemma shows, these terms therefore contribute nothing to (3.20).

Lemma 3.6. Suppose \mathfrak{p} is prime ideal dividing \mathfrak{q} . Let $f:\Gamma_{0,j}(\mathfrak{p}^{-1}\mathfrak{q})\to\mathbb{C}$ be left invariant under $\Gamma_{1,j}(\mathfrak{p}^{-1}\mathfrak{q})$ and transform by $\overline{\chi}_{\tau}$ on the right under $\Gamma_{0,j}(\mathfrak{q})$. Then for any $x\in \mathrm{GL}_2(\mathbb{A}_{F,f})$,

$$\sum_{g\in \Gamma_{0,j}(\mathfrak{p}^{-1}\mathfrak{q})/\Gamma_{0,j}(\mathfrak{q})}W_{\varphi_f}(xgh_j)f(g)=0.$$

Proof. Let v be the place corresponding to \mathfrak{p} , and put $m = \operatorname{ord}_{\mathfrak{p}}(\mathfrak{q})$. For any subgroup $G \subseteq \operatorname{GL}_2$, let G^1 denote $G \cap \operatorname{SL}_2$. The natural map

$$\Gamma^1_{1,j}(\mathfrak{p}^{-1}\mathfrak{q}) \to K^1_{1,v}(\mathfrak{p}^{m-1})/K^1_v(\mathfrak{p}^m)$$

is surjective by strong approximation for $\mathrm{SL}_2(\mathbb{A}_F)$. Fix $k_1 \in K^1_{1,v}(\mathfrak{p}^{m-1})$ and choose $k \in \Gamma^1_{1,j}(\mathfrak{p}^{-1}\mathfrak{q})$ that maps to the coset $k_1K^1_v(\mathfrak{p}^m)$ above. We have

$$\sum_{g \in \Gamma_{0,j}(\mathfrak{p}^{-1}\mathfrak{q})/\Gamma_{0,j}(\mathfrak{q})} W_{\varphi_f}(xgh_j)f(g) = \sum_{g \in \Gamma_{0,j}(\mathfrak{p}^{-1}\mathfrak{q})/\Gamma_{0,j}(\mathfrak{q})} W_{\varphi_f}(xkgh_j)f(g).$$

For any finite place $w \neq v$, we have

$$W_{\varphi_w}(xkgh_j) = W_{\varphi_w}(xgh_j),$$

since $(gh_j)^{-1}k_w(gh_j) \in K_{1,w}(\mathfrak{p}_w^{\operatorname{ord}_{\mathfrak{p}_w}(\mathfrak{q})})$. At v we have $k_v = k_1k'$ for some $k' \in K_v^1(\mathfrak{p}^m)$, so that

$$W_{\varphi_v}(xkgh_j) = W_{\varphi_v}(xk_1k'g) = W_{\varphi_v}(xk_1gg^{-1}k'g) = W_{\varphi_v}(xk_1g) = W_{\varphi_v}(xgg^{-1}k_1g),$$

since $g^{-1}k'g \in g^{-1}K_v^1(\mathfrak{p}^m)g = K_v^1(\mathfrak{p}^m)$.

Now, if k_1 runs through a set of representatives for $K_{1,v}^1(\mathfrak{p}_v^{m-1})/K_v^1(\mathfrak{p}_v^m)$, so does $g^{-1}k_1g$, since g normalizes both $K_{1,v}^1(\mathfrak{p}_v^{m-1})$ and $K_v^1(\mathfrak{p}_v^m)$. Thus, averaging over k_1 , our sum becomes

$$\sum_{g \in \Gamma_{0,j}(\mathfrak{p}^{-1}\mathfrak{q})/\Gamma_{0,j}(\mathfrak{q})} f(g) \prod_{\substack{w < \infty \\ w \neq v}} W_{\varphi_w}(xgh_j) \cdot \frac{1}{[K^1_{1,v}(\mathfrak{p}^{m-1}_v) : K^1_{1,v}(\mathfrak{p}^m_v)]} \sum_{k_1} W_{\varphi_v}(xgk_1).$$

Consider the inner sum $\sum_{k_1} W_{\varphi_v}(tk_1)$ as a function of t. Fixing $k \in K_{1,v}(\mathfrak{p}_v^{m-1})$, we have $k \in \binom{y}{1} K_{1,v}^1(\mathfrak{p}_v^{m-1})$ for some $y \in \mathfrak{o}_v^{\times}$. Hence,

$$\sum_{k_1} W_{\varphi_v}(tkk_1) = \sum_{k_1} W_{\varphi_v}(t({}^{y}_{1})k_1) = \sum_{k_1} W_{\varphi_v}(tk_1({}^{y}_{1})) = \sum_{k_1} W_{\varphi_v}(tkk_1),$$

since $\begin{pmatrix} y \\ 1 \end{pmatrix} \in K_{1,v}(\mathfrak{p}_v^m)$. Therefore $\sum_{k_1} W_{\varphi_v}(tk_1) = 0$.

Hence, we may assume that $\gamma_1 \mu_{22} \mathfrak{o}_F + \mathfrak{q} = \mathfrak{o}_F$. In this case, we can choose representatives of the form $\mu = \begin{pmatrix} 1 & x \\ 1 & 1 \end{pmatrix}$, where $x \in \mathfrak{a}_j \mathfrak{p}_k^2 / (\zeta) \mathfrak{a}_j \mathfrak{p}_k^2$, and we have

$$\sum_{\nu \in X_0/X_1} \chi_{\tau}(\nu) \psi_{\infty}(\gamma_1 \beta_2 \mu_{22} \nu_{22}) = \overline{\chi_{\tau}(\gamma_1)} \tau_{\mathfrak{q}}(\chi_{\tau}, \beta_2).$$

Further, we compute that

$$W_{\varphi_f}\left(\begin{pmatrix} \gamma_1 \gamma_2 & \\ & \gamma_1 \end{pmatrix} \ell_k \eta h_j' \right) = \omega_{\tau_\infty}^{-1}(\gamma_1) W_{\varphi_f}\left(\begin{pmatrix} \gamma_2 & \\ & 1 \end{pmatrix} \ell_k h_j' \right).$$

Consequently the finite part of $\frac{1}{N(\mathfrak{q}_1)} \sum_{\beta_1 \in \mathfrak{a}\mathfrak{q}_1^{-1}/\mathfrak{a}} \Lambda_j(s,\pi,\tau,\xi_\infty,\varphi_\infty,\beta,\zeta)$ takes the form

$$\begin{split} &\frac{N(\zeta)[X_1:\Gamma_j(\mathfrak{q}')]}{[\Gamma_{1,j}(\mathfrak{q}):\Gamma_j(\mathfrak{q}')]}N(\mathfrak{a}_j)^{\frac{1}{2}-s}\sum_kN(\mathfrak{a}_j\mathfrak{p}_k^2)\|\gamma_2\|_{\infty} \\ &\cdot \sum_{\gamma_1\in\mathfrak{o}_F^{\times}\backslash\mathfrak{p}_k\cap F^{\times}}\sum_{\gamma_2\in\mathfrak{o}_F^{\times}\backslash\mathfrak{a}_j^{-1}\mathfrak{p}_k^{-2}\cap F^{\times}}\|\gamma_1^2\gamma_2\|_{\infty}^{\frac{1}{2}-s}W_{\xi_f^0}\begin{pmatrix}\gamma_1\gamma_2\mathsf{a}_j\mathsf{p}_k&\\&\gamma_1\mathsf{p}_k^{-1}&\\&1\end{pmatrix} \\ &\cdot \tau_{\mathfrak{q}}(\chi_{\tau},\beta_2)\psi_{v_k}\big(\alpha_k^{-1}\gamma_2\big)W_{\varphi_f}\bigg(\begin{pmatrix}\gamma_2&\\&1\end{pmatrix}\ell_kh_j'\bigg)\,\omega_{\tau_{\infty}}^{-1}(\gamma_1)\overline{\chi_{\tau}(\gamma_1)}. \end{split}$$

For γ_1, γ_2 as above, let $\mathfrak{m}_1 = (\gamma_1)\mathfrak{p}_k^{-1}$ and $\mathfrak{m}_2 = (\gamma_2)\mathfrak{a}_j\mathfrak{p}_k^2$. Then $\|\gamma_1^2\gamma_2\|_{\infty} = N(\mathfrak{m}_1^2\mathfrak{m}_2\mathfrak{a}_j^{-1})$ and $(\gamma_1) = \mathfrak{p}_k\mathfrak{m}_1$.

Lemma 3.7. We have

$$\psi_{v_k} (\alpha_k^{-1} \gamma_2) W_{\varphi_f} \begin{pmatrix} \begin{pmatrix} \gamma_2 \mathsf{p}_k & \\ & \mathsf{p}_k \end{pmatrix} \ell_k h_j' \end{pmatrix} = W_{\varphi_f} \begin{pmatrix} \zeta & \\ & 1 \end{pmatrix} W_{\varphi_f} \begin{pmatrix} \begin{pmatrix} \gamma_2 \mathsf{p}_k^2 & \\ & 1 \end{pmatrix} h_j \end{pmatrix}.$$

Proof. We check this locally at every place v by the same computation as in the proof of Lemma 3.2. The proof here splits naturally into two cases. First, suppose $\mathfrak{c} \neq \mathfrak{q}$. Then $\zeta = 1$ and the lemma is trivial to check at any finite place $v \neq v_k$. For $v = v_k$, we write the v-component of $\alpha_k \mathsf{a}_j = u/\varpi_v$ as in Lemma 3.2 and use the decomposition (3.3) to re-write the left hand side (locally at v) as

$$\begin{split} \psi_{v_k} \left(\alpha_k^{-1} \gamma_2 \right) W_{\varphi_v^0} \left(\begin{pmatrix} \gamma_2 \varpi_v & \\ & \varpi_v \end{pmatrix} \ell_k h_j' \right) &= \psi_{v_k} \left(\alpha_k^{-1} \gamma_2 \right) W_{\varphi_v^0} \left(\begin{pmatrix} \gamma_2 \mathbf{a}_j \varpi_v & \\ & \varpi_v \end{pmatrix} \begin{pmatrix} 1 & \\ u/\varpi_v & 1 \end{pmatrix} \right) \\ &= W_{\varphi_v^0} \left(\begin{pmatrix} \gamma_2 \mathbf{a}_j \varpi_v^2 & \\ & 1 \end{pmatrix} \begin{pmatrix} 0 & -u^{-1} \\ u & \varpi_v \end{pmatrix} \right). \end{split}$$

Since the prime ideal \mathfrak{p}_k corresponding to the place $v = v_k$ is co-prime to \mathfrak{q} , we see that $W_{\varphi_v^0}$ is right $\mathrm{GL}_2(\mathfrak{o}_v)$ -invariant and thus the conclusion in the case at hand.

Next, suppose $\mathfrak{c} = \mathfrak{q}$. For $v < \infty$ such that $\operatorname{ord}_v(\mathfrak{q}) = 0$, since by choice $\zeta_v \in \mathfrak{o}_v^{\times}$, it follows that $W_{\varphi_v^0}$ is right invariant under translation by $\binom{\zeta_v}{1}$. Consequently the proof for such v is similar to the above argument. Henceforth we assume v is such that $\operatorname{ord}_v(\mathfrak{q}) > 0$. Then $v \neq v_k$, and we need to show

$$W_{\varphi_v^0}\left(\begin{pmatrix} \gamma_2 & \\ & 1 \end{pmatrix} \ell_k h'_{j,v} \right) = W_{\varphi_v^0}\begin{pmatrix} \zeta_v & \\ & 1 \end{pmatrix} W_{\varphi_v^0}\left(\begin{pmatrix} \gamma_2 & \\ & 1 \end{pmatrix} h_{j,v} \right).$$

Moving $h'_{i,v}$ past ℓ_k to the left, we re-write the left hand side of the above equation as

$$W_{\varphi_v^0}\left(\begin{pmatrix} \gamma_2\zeta_v\mathsf{a}_{j,v} & \\ & 1 \end{pmatrix}\begin{pmatrix} 1 & \\ \alpha_k\mathsf{a}_{j,v}\zeta_v & 1 \end{pmatrix}\right),$$

which in turn equals

$$W_{\varphi_v^0} \begin{pmatrix} \gamma_2 \zeta_v \mathsf{a}_{j,v} & \\ & 1 \end{pmatrix},$$

since $\operatorname{ord}_v(\alpha_k \mathbf{a}_{j,v} \zeta_v) \geq \operatorname{ord}_v(\mathfrak{q})$ by choice of α_k . It therefore remains to prove

$$(3.21) \qquad \qquad W_{\varphi_v^0} \begin{pmatrix} \gamma_2 \zeta_v \mathsf{a}_{j,v} & \\ & 1 \end{pmatrix} = W_{\varphi_v^0} \begin{pmatrix} \zeta_v & \\ & 1 \end{pmatrix} W_{\varphi_v^0} \begin{pmatrix} \gamma_2 \mathsf{a}_{j,v} & \\ & 1 \end{pmatrix}.$$

Let St denote the Steinberg representation of $GL_2(F_v)$, i.e., the unique Langlands quotient of the induced module $Ind(\|\cdot\|_v^{-1/2}, \|\cdot\|_v^{1/2})$. It is a well-known fact that τ_v is either supercuspidal, or a twist of the Steinberg representation $St \otimes \chi$, or an irreducible principal series representation $Ind(\chi_1, \chi_2)$, where χ, χ_1 and χ_2 are characters of F_v^{\times} . Since $ord_v(\mathfrak{c}) = ord_v(\mathfrak{q}) > 0$, it follows from [23, Proposition 3.4] that τ_v cannot be supercuspidal. Further, using the formula for the conductor of τ_v in the remaining cases (see [9, Remark 4.25]), we conclude that $\tau_v \cong Ind(\chi_1, \chi_2)$ with either χ_1 or χ_2 (but not both) ramified. Thus one of the Langlands parameters of τ_v is zero; say $(\alpha, \beta) = (\alpha, 0)$ are the Langlands parameters of τ_v . Now, we may appeal to the explicit description of the essential function $W = W_{\varphi_v^0}$ (cf. [21, Theorem 4.1]) which in our case (n = 2) reads as

$$W\begin{pmatrix} \varpi_v^f \\ 1 \end{pmatrix} = \begin{cases} q_v^{-f/2} s_f(\alpha, \beta) & \text{if } f \ge 0, \\ 0 & \text{otherwise,} \end{cases}$$

where $s_f(\alpha, \beta)$ is the Schur polynomial $\frac{\alpha^{f+1}-\beta^{f+1}}{\alpha-\beta}$. Since $\beta=0$ and $\operatorname{ord}_v(\zeta_v), \operatorname{ord}_v(\gamma_2 \mathsf{a}_{j,v}) \geq 0$, our assertion (3.21) follows.

We define

(3.22)
$$\lambda_{\pi}(\mathfrak{m}_{1},\mathfrak{m}_{2}) = N(\mathfrak{m}_{1}\mathfrak{m}_{2})W_{\xi_{f}^{0}}\begin{pmatrix} \mathsf{m}_{1}\mathsf{m}_{2} & \\ & \mathsf{m}_{1} \\ & & 1 \end{pmatrix}$$

and

$$\lambda_{\tau}(\mathfrak{m}_2) = \sqrt{N(\mathfrak{m}_2)} W_{\varphi_f} \begin{pmatrix} \mathsf{m}_2 \\ & 1 \end{pmatrix},$$

where m_1 and m_2 are finite idèles such that $\mathfrak{m}_1=(m_1)$ and $\mathfrak{m}_2=(m_2)$. Then

$$\begin{split} \frac{1}{N(\mathfrak{q}_1)} \sum_{\beta_1 \in \mathfrak{a}\mathfrak{q}_1^{-1}/\mathfrak{a}} \Lambda_j(s,\pi,\tau,\xi_\infty,\varphi_\infty,\beta,\zeta) &= \frac{\sqrt{N(\zeta)}[X_1:\Gamma_j(\mathfrak{q}')]}{[\Gamma_{1,j}(\mathfrak{q}):\Gamma_j(\mathfrak{q}')]} \tau_{\mathfrak{q}}(\chi_\tau,\beta_2) \lambda_\tau((\zeta)) \prod_{v|\infty} \Psi_v(s;W_{\xi_v},W_{\varphi_v}) \\ &\quad \cdot \sum_k \sum_{\mathfrak{m}_1 \sim \mathfrak{p}_k^{-1}} \sum_{\mathfrak{m}_2 \sim \mathfrak{a}_j \mathfrak{p}_k^2} \lambda_\pi(\mathfrak{m}_1,\mathfrak{m}_2) \lambda_\tau(\mathfrak{m}_2) \chi_{\omega_\tau}(\mathfrak{m}_1) N(\mathfrak{m}_1^2\mathfrak{m}_2)^{-s}, \end{split}$$

where $\chi_{\omega_{\tau}}$ is the Größencharakter of modulus \mathfrak{q} associated to ω_{τ} (which need not be primitive). Noting that $\frac{[X_1:\Gamma_j(\mathfrak{q}')]}{[\Gamma_{1,j}(\mathfrak{q}):\Gamma_j(\mathfrak{q}')]} = \frac{N(\mathfrak{q})}{N(\mathfrak{q}_1(\zeta)\cap\mathfrak{q}_2)}$, we can now sum over j to get

$$\frac{1}{N(\mathfrak{q}_1)} \sum_{\beta_1} \sum_j \Lambda_j(s,\pi,\tau,\xi_\infty,\varphi_\infty,\beta,\zeta) = \frac{\sqrt{N(\mathfrak{q}^2(\zeta))}}{N(\mathfrak{q}_1(\zeta)\cap\mathfrak{q}_2)} \tau_{\mathfrak{q}}(\chi_\tau,\beta_2) \lambda_\tau((\zeta)) \prod_{v\mid\infty} \Psi_v(s;W_{\xi_v},W_{\varphi_v}) \\ \cdot \sum_{\mathfrak{m}_1,\mathfrak{m}_2} \lambda_\pi(\mathfrak{m}_1,\mathfrak{m}_2) \lambda_\tau(\mathfrak{m}_2) \chi_{\omega_\tau}(\mathfrak{m}_1) N(\mathfrak{m}_1^2\mathfrak{m}_2)^{-s}.$$

It remains to identify the Dirichlet series in the above expression.

Lemma 3.8. We have

(3.23)
$$\sum_{\mathfrak{m}_1,\mathfrak{m}_2} \lambda_{\pi}(\mathfrak{m}_1,\mathfrak{m}_2) \lambda_{\tau}(\mathfrak{m}_2) \chi_{\omega_{\tau}}(\mathfrak{m}_1) N(\mathfrak{m}_1^2 \mathfrak{m}_2)^{-s} = L(s, \pi \times \tau).$$

Proof. For a fixed choice of τ and a non-zero integral ideal \mathfrak{a} , define

$$c_{\pi,\tau}(\mathfrak{a}) = \sum_{\mathfrak{m}_1^2\mathfrak{m}_2 = \mathfrak{a}} \lambda_{\pi}(\mathfrak{m}_1,\mathfrak{m}_2) \lambda_{\tau}(\mathfrak{m}_2) \chi_{\omega_{\tau}}(\mathfrak{m}_1).$$

Then, for any unramified idèle class character ω , we have

$$\lambda_{\tau \otimes \omega}(\mathfrak{m}_2) = \lambda_{\tau}(\mathfrak{m}_2)\chi_{\omega}(\mathfrak{m}_2)$$
 and $\chi_{\omega_{\tau \otimes \omega}}(\mathfrak{m}_1) = \chi_{\omega_{\tau}}(\mathfrak{m}_1)\chi_{\omega}(\mathfrak{m}_1)^2$,

so that

$$(3.24) c_{\pi,\tau\otimes\omega}(\mathfrak{a}) = c_{\pi,\tau}(\mathfrak{a})\chi_{\omega}(\mathfrak{a}).$$

Next define $\lambda_{\pi \times \tau}$ to be the Dirichlet coefficients of $L(s, \pi \times \tau)$, so that, for any unramified ω , we have

(3.25)
$$L(s, \pi \times (\tau \otimes \omega)) = \sum_{\mathfrak{a}} \lambda_{\pi \times \tau}(\mathfrak{a}) \chi_{\omega}(\mathfrak{a}) N(\mathfrak{a})^{-s}.$$

Note that both $c_{\pi,\tau}$ and $\lambda_{\pi\times\tau}$ are multiplicative, so it suffices to show that they agree at prime powers.

Given integers

$$(3.26) \lambda_1 \ge \lambda_2 \ge \lambda_3 \ge 0,$$

the Schur polymial $s_{\lambda_1,\lambda_2,\lambda_3}(x_1,x_2,x_3)$ is the ratio

(3.27)
$$\frac{\det \begin{pmatrix} x_1^{\lambda_1+2} & x_2^{\lambda_1+2} & x_3^{\lambda_1+2} \\ x_1^{\lambda_2+1} & x_2^{\lambda_2+1} & x_3^{\lambda_2+1} \\ x_1^{\lambda_3} & x_2^{\lambda_3} & x_3^{\lambda_3} \end{pmatrix}}{\det \begin{pmatrix} x_1^2 & x_2^2 & x_3^2 \\ x_1 & x_2 & x_3 \\ 1 & 1 & 1 \end{pmatrix}}.$$

By the Cauchy identity [5, Theorem 43.3], for sufficiently small $\alpha_1, \alpha_2, \alpha_3, \gamma_1, \gamma_2, \gamma_3 \in \mathbb{C}$, we have

$$\prod_{i=1}^{3} \prod_{j=1}^{3} \frac{1}{1 - \alpha_i \gamma_j} = \sum_{\lambda} s_{\lambda}(\alpha_1, \alpha_2, \alpha_3) s_{\lambda}(\gamma_1, \gamma_2, \gamma_3),$$

where the sum runs over all $\lambda = (\lambda_1, \lambda_2, \lambda_3)$ satisfying (3.26). Note that if $\gamma_3 = 0$ then only terms with $\lambda_3 = 0$ contribute to the above sum. Writing $\lambda_2 = k_1$, $\lambda_1 = k_1 + k_2$ and replacing $(\gamma_1, \gamma_2, \gamma_3)$ by $(x\gamma_1, x\gamma_2, 0)$ for a small $x \in \mathbb{C}$, we obtain

$$\prod_{i=1}^{3} \prod_{j=1}^{2} \frac{1}{1 - \alpha_{i} \gamma_{j} x} = \sum_{k_{1}=0}^{\infty} \sum_{k_{2}=0}^{\infty} s_{k_{1} + k_{2}, k_{1}, 0}(\alpha_{1}, \alpha_{2}, \alpha_{3}) s_{k_{1} + k_{2}, k_{1}, 0}(\gamma_{1}, \gamma_{2}, 0) x^{2k_{1} + k_{2}}.$$

Now fix a prime ideal \mathfrak{p} , and let v be the corresponding place of F; then $q_v = N(\mathfrak{p})$. Let $\{\alpha_1, \alpha_3, \alpha_3\}$ (resp. $\{\gamma_1, \gamma_2\}$) denote the Langlands parameters of π_v (resp. τ_v), so that

(3.28)
$$L(s, \pi_v) = \prod_{i=1}^{3} \frac{1}{1 - \alpha_i N(\mathfrak{p})^{-s}} \quad \text{and} \quad L(s, \tau_v) = \prod_{i=1}^{2} \frac{1}{1 - \gamma_i N(\mathfrak{p})^{-s}}.$$

Then by the above we have

$$L(s, \pi_v \times \tau_v) = \prod_{i=1}^3 \prod_{j=1}^2 \frac{1}{1 - \alpha_i \gamma_j N(\mathfrak{p})^{-s}}$$

$$= \sum_{k_1=0}^\infty \sum_{k_2=0}^\infty s_{k_1+k_2, k_1, 0}(\alpha_1, \alpha_2, \alpha_3) s_{k_1+k_2, k_1, 0}(\gamma_1, \gamma_2, 0) N(\mathfrak{p})^{-(2k_1+k_2)s}.$$

On the other hand, we have

$$L(s, \pi_v \times \tau_v) = \sum_{k=0}^{\infty} \lambda_{\pi \times \tau}(\mathfrak{p}^k) N(\mathfrak{p})^{-ks},$$

whence

(3.29)
$$\lambda_{\pi \times \tau}(\mathfrak{p}^k) = \sum_{2k_1 + k_2 = k} s_{k_1 + k_2, k_1, 0}(\alpha_1, \alpha_2, \alpha_3) s_{k_1 + k_2, k_1, 0}(\gamma_1, \gamma_2, 0).$$

Further, by (3.28) and the identity $L(s, \tau_v) = \sum_{k=0}^{\infty} \lambda_{\tau}(\mathfrak{p}^k) N(\mathfrak{p})^{-ks}$, we have

$$\lambda_{\tau}(\mathfrak{p}^{k_2}) = \sum_{j=0}^{k_2} \gamma_1^j \gamma_2^{k_2 - j} = \frac{\gamma_1^{k_2 + 1} - \gamma_2^{k_2 + 1}}{\gamma_1 - \gamma_2}.$$

Moreover, $\chi_{\omega_{\tau}}(\mathfrak{p}^{k_1}) = (\gamma_1 \gamma_2)^{k_1}$, so that

(3.30)
$$\lambda_{\tau}(\mathfrak{p}^{k_2})\chi_{\omega_{\tau}}(\mathfrak{p}^{k_1}) = (\gamma_1\gamma_2)^{k_1} \frac{\gamma_1^{k_2+1} - \gamma_2^{k_2+1}}{\gamma_1 - \gamma_2} = s_{k_1+k_2,k_1,0}(\gamma_1, \gamma_2, 0),$$

by (3.27). Thus,

(3.31)
$$c_{\pi,\tau}(\mathfrak{p}^k) = \sum_{2k_1 + k_2 = k} \lambda_{\pi}(\mathfrak{p}^{k_1}, \mathfrak{p}^{k_2}) s_{k_1 + k_2, k_1, 0}(\gamma_1, \gamma_2, 0).$$

Suppose now that τ is unramified. Then we have $\mathfrak{q}_1 = \mathfrak{o}_F$, so we may take $\beta_1 = \beta_2 = 0$. In that case, we have (cf. (3.9))

$$\sum_{j} \Lambda_{j}(s, \pi, \tau, \xi_{\infty}, \varphi_{\infty}, \beta) = \Psi_{\infty}(s, \xi_{\infty}, \varphi_{\infty}) L(s, \pi \times \tau).$$

Choosing ξ_{∞} and φ_{∞} such that $\Psi_{\infty}(s, \xi_{\infty}, \varphi_{\infty}) \neq 0$, we conclude that (3.23) holds. Replacing τ by an unramified twist $\tau \otimes \omega$, by (3.24) and (3.25), we have

$$\sum_{\mathfrak{a}} c_{\pi,\tau}(\mathfrak{a}) \chi_{\omega}(\mathfrak{a}) N(\mathfrak{a})^{-s} = \sum_{\mathfrak{a}} \lambda_{\pi \times \tau}(\mathfrak{a}) \chi_{\omega}(\mathfrak{a}) N(\mathfrak{a})^{-s}$$

for any unramified ω . By [4, Lemma 4.2] it follows that $c_{\pi,\tau}(\mathfrak{a}) = \lambda_{\pi\times\tau}(\mathfrak{a})$ for all \mathfrak{a} . In particular, taking $\mathfrak{a} = \mathfrak{p}^k$, by (3.29) and (3.31) we find that (3.32)

$$\sum_{2k_1+k_2=k} s_{k_1+k_2,k_1,0}(\alpha_1,\alpha_2,\alpha_3)s_{k_1+k_2,k_1,0}(\gamma_1,\gamma_2,0) = \sum_{2k_1+k_2=k} \lambda_{\pi}(\mathfrak{p}^{k_1},\mathfrak{p}^{k_2})s_{k_1+k_2,k_1,0}(\gamma_1,\gamma_2,0),$$

whenever τ is unramified. Applying this with $\tau = \|\cdot\|^{it_1} \boxplus \|\cdot\|^{it_2}$ for arbitrary $t_1, t_2 \in \mathbb{R}$, the Satake parameters $(\gamma_1, \gamma_2) = (N(\mathfrak{p})^{-it_1}, N(\mathfrak{p})^{-it_2})$ are Zariski-dense in \mathbb{C}^2 , so (3.32) holds for arbitrary $\gamma_1, \gamma_2 \in \mathbb{C}$. Further, from (3.30) it is easy to see that the polynomials $s_{k_1+k_2,k_1,0}(x_1, x_2, 0)$, for k_1, k_2 ranging over all non-negative integers, are linearly independent. Therefore, from (3.32) we conclude that $\lambda_{\pi}(\mathfrak{p}^{k_1}, \mathfrak{p}^{k_2}) = s_{k_1+k_2,k_1,0}(\alpha_1, \alpha_2, \alpha_3)$.

Finally, applying this together with (3.29) and (3.31) for an arbitrary τ (not necessarily unramified), we conclude that $c_{\pi,\tau}(\mathfrak{p}^k) = \lambda_{\pi \times \tau}(\mathfrak{p}^k)$, as desired.

Hence, we obtain

$$\frac{1}{N(\mathfrak{q}_1)} \sum_{\beta_1} \sum_{i} \Lambda_j(s, \pi, \tau, \xi_\infty, \varphi_\infty, \beta, \zeta) = \frac{\sqrt{N(\mathfrak{q}^2(\zeta))}}{N(\mathfrak{q}_1(\zeta) \cap \mathfrak{q}_2)} \tau_{\mathfrak{q}}(\chi_\tau, \beta_2) \lambda_\tau((\zeta)) \Psi_\infty(s, \xi_\infty, \varphi_\infty) L(s, \pi \times \tau).$$

Taking $\zeta = 1$, we obtain the conclusion of Theorem 3.4.

3.4. Functional equation. In this section we assume both π and τ are cuspidal automorphic representations. We then show how to use Theorem 3.4 to give a new proof of the analytic properties of $L(s, \pi \boxtimes \tau)$ when π and τ have coprime conductors and τ_v is a twist-minimal principal series representation for all finite v.

Let $\xi = \bigotimes_v \xi_v$ be a pure tensor in the space of π such that $\xi_v = \xi_v^0$ for every finite v. Let (U_ξ, U'_ξ) be the *automorphic pair* attached to such a ξ as in Lemma 2.3. Likewise, let (U_φ, U'_φ) be the automorphic pair associated with a pure tensor $\bigotimes_v \varphi_v$ in the space of τ satisfying $\varphi_v = \varphi_v^0$ for all $v < \infty$.

Theorem 3.9. Keep the above notation as well as that of Theorem 3.4. Assume that π and τ are cuspidal automorphic, with $\mathfrak{n} + \mathfrak{q} = \mathfrak{o}_F$ and $\mathfrak{c} = \mathfrak{q}$. Then $L(s, \pi \boxtimes \tau)$ continues to an entire function and satisfies the functional equation

$$\Psi_{\infty}(s, \xi_{\infty}, \varphi_{\infty}) L(s, \pi \boxtimes \tau) = \epsilon N(\mathfrak{n}^2 \mathfrak{q}^3)^{\frac{1}{2} - s} \widetilde{\Psi}_{\infty}(1 - s, \xi_{\infty}, \varphi_{\infty}) L(1 - s, \widetilde{\pi} \boxtimes \widetilde{\tau}),$$

where

$$\widetilde{\Psi}_{\infty}(s, \xi_{\infty}, \varphi_{\infty}) = \prod_{v \mid \infty} \Psi_{v}(s; \widetilde{W}_{\widetilde{\xi}_{v}}, \widetilde{W}_{\widetilde{\varphi}_{v}}) \quad (\text{cf. } (2.10))$$

and ϵ is as defined in (3.35).

Proof. For any finite-index subgroup $H \leq \Gamma_i(\mathfrak{q})$, we have

 $\Lambda_j(s,\pi,\tau,\xi_\infty,\varphi_\infty,(0,\beta_2))$

$$=\frac{N(\mathfrak{a}_{j})^{\frac{1}{2}-s}}{\left[\Gamma_{1,j}(\mathfrak{q}):H\right]}\int_{H\backslash\operatorname{GL}_{2}(F_{\infty})}U_{\xi}\left(\begin{pmatrix}1&&&\\&1&\beta_{2}\\&&1\end{pmatrix}\begin{pmatrix}h&&\\&1\end{pmatrix},\begin{pmatrix}\mathsf{a}_{j}&&\\&&1\end{pmatrix}\right)U_{\varphi}(h,h_{j})\|\det h\|^{s-\frac{1}{2}}dh.$$

Let n and q be idèles such that n = (n) and q = (q). We make the change of variables $h \mapsto {}^t h^{-1}$ and use the identities

$$U_{\xi}(g) = \epsilon_{\pi}^2 U_{\xi}' \begin{pmatrix} {}^t g^{-1} \begin{pmatrix} \mathsf{n} & & \\ & \mathsf{n} & \\ & & 1 \end{pmatrix} \end{pmatrix}, \quad U_{\varphi}(g) = \epsilon_{\tau} U_{\varphi}' \begin{pmatrix} {}^t g^{-1} \begin{pmatrix} \mathsf{q} & \\ & 1 \end{pmatrix} \end{pmatrix},$$

from Lemma 2.3 to get

$$\frac{\epsilon_{\pi}^{2} \epsilon_{\tau} N(\mathfrak{a}_{j})^{\frac{1}{2}-s}}{[\Gamma_{1,j}(\mathfrak{q}):H]} \int_{{}^{t}H \backslash \operatorname{GL}_{2}(F_{\infty})} U_{\xi}' \begin{pmatrix} 1 & & \\ & 1 & \\ & -\beta_{2} & 1 \end{pmatrix} \begin{pmatrix} h & \\ & 1 \end{pmatrix}, \begin{pmatrix} \mathsf{a}_{j}^{-1} \mathsf{n} & \\ & & \mathsf{n} \\ & & 1 \end{pmatrix} \end{pmatrix} \\ \cdot U_{\varphi}' \begin{pmatrix} h, \begin{pmatrix} \mathsf{a}_{j}^{-1} \mathsf{q} & \\ & & 1 \end{pmatrix} \end{pmatrix} \| \det h \|^{\frac{1}{2}-s} dh.$$

Next we make the change of variables $h \mapsto \gamma h$ for some $\gamma \in \mathrm{GL}_2(F)$ and use automorphy of U'_{ξ} and U'_{φ} to get

$$\frac{\epsilon_{\pi}^{2}\epsilon_{\tau}(N(\mathfrak{a}_{j})\|\det\gamma\|_{\infty})^{\frac{1}{2}-s}}{[\Gamma_{1,j}(\mathfrak{q}):H]} \int_{\gamma^{-1}(^{t}H)\gamma\backslash\operatorname{GL}_{2}(F_{\infty})} U'_{\xi}\begin{pmatrix}\begin{pmatrix}h\\&1\end{pmatrix},\begin{pmatrix}\gamma^{-1}\\&1\end{pmatrix}\begin{pmatrix}1\\&\beta_{2}&1\end{pmatrix}\begin{pmatrix}\mathsf{a}_{j}^{-1}\mathsf{n}\\&&1\end{pmatrix}\end{pmatrix} \\ \cdot U'_{\varphi}\begin{pmatrix}h,\gamma^{-1}\begin{pmatrix}\mathsf{a}_{j}^{-1}\mathsf{q}\\&&1\end{pmatrix}\end{pmatrix} \|\det h\|^{\frac{1}{2}-s}\,dh.$$

Put $r = q\beta_2$ and choose $\gamma \in GL_2(F)$, a_k and $\kappa \in K(\mathfrak{q}^2)$ such that $\binom{\mathsf{a}_j^{-1}\mathsf{nq}}{\mathsf{nq}^2} = \gamma \binom{\mathsf{a}_k}{1} \kappa$. Then we have

$$\begin{pmatrix} \gamma^{-1} & \\ & 1 \end{pmatrix} \begin{pmatrix} 1 & \\ & 1 \\ & \beta_2 & 1 \end{pmatrix} \begin{pmatrix} \mathsf{a}_j^{-1} \mathsf{n} & \\ & & \mathsf{n} \\ & & & 1 \end{pmatrix} = \mathsf{q}^{-1} \begin{pmatrix} 1 & & -\beta_1' \\ & 1 & -\beta_2' \\ & & & 1 \end{pmatrix} \begin{pmatrix} \mathsf{a}_k & \\ & 1 \\ & & 1 \end{pmatrix} \begin{pmatrix} \kappa & \\ & 1 \end{pmatrix} \begin{pmatrix} 1 & \mathsf{nru} & \mathsf{qu} \\ & \frac{\mathsf{nrv}+1}{\mathsf{q}} & \mathsf{v} \\ & \mathsf{nr} & \mathsf{q} \end{pmatrix},$$

where $\binom{\beta_1'}{\beta_2'} = \mathsf{nq} \gamma^{-1} \binom{\mathsf{a}_j^{-1} \mathsf{u}}{\mathsf{v}} = \binom{\mathsf{a}_k}{1} \kappa \binom{\mathsf{u}}{\mathsf{q}^{-1} \mathsf{v}}$. We may choose $\mathsf{u}, \mathsf{v} \in \prod_{\mathfrak{p}} \mathfrak{o}_{\mathfrak{p}}$ so that $\frac{\mathsf{nrv} + 1}{\mathsf{q}} \in \prod_{\mathfrak{p}} \mathfrak{o}_{\mathfrak{p}}$, (nqv) and $(\mathsf{nqa}_j^{-1} \mathsf{u})$ are principal, and $\beta_1' = 0$. Taking determinants, we find that $(\det \gamma) = \mathfrak{n}^2 \mathfrak{q}^3 \mathfrak{a}_j^{-1} \mathfrak{a}_k^{-1}$. Also,

$$\gamma^{-1} \begin{pmatrix} \mathsf{a}_j^{-1} \mathsf{q} & \\ & 1 \end{pmatrix} = (\mathsf{n} \mathsf{q}^2)^{-1} \begin{pmatrix} \mathsf{a}_k \mathsf{q}^2 & \\ & 1 \end{pmatrix} \cdot \begin{pmatrix} \mathsf{q}^{-2} & \\ & 1 \end{pmatrix} \kappa \begin{pmatrix} \mathsf{q}^2 & \\ & 1 \end{pmatrix} \in (\mathsf{n} \mathsf{q}^2)^{-1} \begin{pmatrix} \mathsf{a}_k \mathsf{q}^2 & \\ & 1 \end{pmatrix} K_1(\mathfrak{q}^4).$$

Hence, we obtain

$$\frac{cN(\mathfrak{n}^2\mathfrak{q}^3)^{\frac{1}{2}-s}N(\mathfrak{a}_k)^{s-\frac{1}{2}}}{[\Gamma_{1,j}(\mathfrak{q}):H]} \int_{\gamma^{-1}({}^tH)\gamma\backslash\operatorname{GL}_2(F_\infty)} U'_{\xi} \begin{pmatrix} 1 & & \\ & 1 & \beta'_2 \\ & & 1 \end{pmatrix} \begin{pmatrix} h & \\ & 1 \end{pmatrix}, \begin{pmatrix} \mathsf{a}_k & \\ & 1 \\ & & 1 \end{pmatrix} \end{pmatrix} \\ \cdot U'_{\varphi} \begin{pmatrix} h, \begin{pmatrix} \mathsf{a}_k \mathsf{q}^2 & \\ & 1 \end{pmatrix} \end{pmatrix} \|\det h\|^{\frac{1}{2}-s} \, dh,$$

where $c = \epsilon_{\pi}^2 \epsilon_{\tau} \omega_{\widetilde{\pi}_f}(\mathsf{q}^{-1}) \chi_{\widetilde{\pi}}(\mathsf{q}) \omega_{\widetilde{\tau}_f}((\mathsf{n}\mathsf{q}^2)^{-1}) = \epsilon_{\pi}^2 \epsilon_{\tau} \chi_{\omega_{\pi}}(\mathsf{q}) \omega_{\tau_f}(\mathsf{n}\mathsf{q}^2).$

We choose H so that $\gamma^{-1}({}^tH)\gamma \subseteq \Gamma_k(\mathfrak{q}^2)$. Then $[\Gamma_{1,j}(\mathfrak{q}):H]=[\Gamma_{1,k}(\mathfrak{q}):\gamma^{-1}({}^tH)\gamma]$, and by (3.10), the above is

$$(3.34) cN(\mathfrak{n}^2\mathfrak{q}^3)^{\frac{1}{2}-s}\Lambda_k(1-s,\widetilde{\pi},\widetilde{\tau},\xi_{\infty}',\varphi_{\infty}',(0,\beta_2'),\mathfrak{q}^2),$$

where ξ_{∞}' and φ_{∞}' are the vectors in $V_{\widetilde{\eta}_{\infty}}$ and $V_{\widetilde{\tau}_{\infty}}$ such that $W_{\xi_{\infty}'} = \widetilde{W}_{\widetilde{\xi}_{\infty}}$ and $W_{\varphi_{\infty}'} = \widetilde{W}_{\widetilde{\varphi}_{\infty}}$. Applying Theorem 3.4 with $\mathfrak{q}_1 = \mathfrak{o}_F$ and $\mathfrak{q}_2 = \mathfrak{q}$, we have

$$\sum_{j=1}^{h} \Lambda_{j}(s, \pi, \tau, \xi_{\infty}, \varphi_{\infty}, (0, \beta_{2})) = \tau_{\mathfrak{q}}(\chi_{\tau}, \beta_{2}) \Psi_{\infty}(s, \xi_{\infty}, \varphi_{\infty}) L(s, \pi \times \tau).$$

On the other hand, summing (3.34) over k and applying (3.33) with $\zeta = q^2$ and π, τ, ψ_{∞} replaced by their duals, we get

$$cN(\mathfrak{n}^2\mathfrak{q}^3)^{\frac{1}{2}-s}\overline{\tau_{\mathfrak{q}}(\chi_{\tau},\beta_2')}\lambda_{\widetilde{\tau}}(\mathfrak{q}^2)\Psi_{\infty}(1-s,\xi_{\infty}',\varphi_{\infty}')L(1-s,\widetilde{\pi}\times\widetilde{\tau})$$

$$=\epsilon N(\mathfrak{n}^2\mathfrak{q}^3)^{\frac{1}{2}-s}\widetilde{\Psi}_{\infty}(1-s,\xi_{\infty},\varphi_{\infty})L(1-s,\widetilde{\pi}\times\widetilde{\tau}),$$

where

(3.35)
$$\epsilon = \frac{\epsilon_{\pi}^2 \epsilon_{\tau} \chi_{\omega_{\pi}}(\mathfrak{q}) \omega_{\tau_f}(\mathsf{nq}^2) \lambda_{\widetilde{\tau}}(\mathfrak{q}^2) \overline{\tau_{\mathfrak{q}}(\chi_{\tau}, \beta_2')}}{\tau_{\mathfrak{q}}(\chi_{\tau}, \beta_2)}.$$

Finally, since $\mathfrak{n} + \mathfrak{q} = \mathfrak{o}_F$, we have $L(s, \pi \boxtimes \tau) = L(s, \pi \times \tau)$ and $L(s, \widetilde{\pi} \boxtimes \widetilde{\tau}) = L(s, \widetilde{\pi} \times \widetilde{\tau})$. This concludes the proof.

4.
$$GL_3 \times GL_1$$

In this section, $\tau = \omega$ is an idèle class character of F of conductor \mathfrak{q} , and π is an irreducible admissible generic representation of $\mathrm{GL}_3(\mathbb{A}_F)$. For each finite place v of F, the essential vector ξ_v^0 and the corresponding $W_{\xi_v^0}$ are as described in the previous section. For $\beta \in F^\times$, we embed β in \mathbb{A}_F via $\beta \mapsto (\beta, 0) \in F_\infty \times \mathbb{A}_{F,f}$ as before, and for $\xi_\infty \in V_{\pi_\infty}$, $j = 1, \ldots, h$, we define the function $\Phi_{\xi_\infty,j}$ on \mathbb{A}_F^\times via

$$\Phi_{\xi_{\infty},j}(y) = \mathbb{P}_1^3(U_{\xi}) \begin{pmatrix} (\beta y, \mathsf{a}_j) & (\beta, 0) \\ & 1 \\ & & 1 \end{pmatrix}, \quad \text{for } \xi = \xi_{\infty} \otimes \xi_f^0, y \in \mathbb{A}_F^{\times}.$$

This function is not invariant under F^{\times} due to the presence of $(\beta, 0)$, but as before it will follow from its Fourier expansion (see below) that it is invariant under a suitable congruence subgroup when viewed as a function on F_{∞}^{\times} . We will from here onwards take $\beta \in \mathfrak{aq}^{-1}$, where \mathfrak{a} as before is the product $\prod_{j} \mathfrak{a}_{j}$.

4.1. The Fourier expansion of $\Phi_{\xi_{\infty},j}$. By definition, for $y \in F_{\infty}^{\times}$, we have

$$\Phi_{\xi_{\infty},j}(y) = \sum_{\gamma \in F^{\times}} W_{\xi_{\infty} \otimes \xi_f^0} \begin{pmatrix} \gamma(\beta y, \mathsf{a}_j) & \gamma(\beta, 0) \\ & 1 \\ & & 1 \end{pmatrix}.$$

One checks that

$$W_{\xi_f^0}\begin{pmatrix} \gamma \mathsf{a}_j & \\ & 1 \\ & & 1 \end{pmatrix} \neq 0 \implies \gamma \in \mathfrak{a}_j^{-1} \cap F^\times,$$

so that

$$\Phi_{\xi_{\infty},j}(y) = \sum_{\gamma \in \mathfrak{a}_{j}^{-1} \cap F^{\times}} W_{\xi_{j}^{0}} \begin{pmatrix} \gamma \mathsf{a}_{j} & & \\ & 1 & \\ & & 1 \end{pmatrix} W_{\xi_{\infty}} \begin{pmatrix} \gamma \beta y & \gamma \beta & \\ & 1 & \\ & & 1 \end{pmatrix}.$$

Let us put $a_{\xi_f^0}(\mathsf{a}_j,\gamma) = W_{\xi_f^0}(\gamma^{\mathsf{a}_j})_1$, and note that it is invariant under $\gamma \mapsto \eta \gamma$ for $\eta \in \mathfrak{o}_F^{\times}$. Thus the above becomes

$$\Phi_{\xi_{\infty},j}(y) = \sum_{\gamma \in \mathfrak{o}_F^{\times} \backslash \mathfrak{a}_i^{-1} \cap F^{\times}} a_{\xi_f^0}(\mathsf{a}_j,\gamma) \sum_{\eta \in \mathfrak{o}_F^{\times}} \psi_{\infty}(\eta \gamma \beta) W_{\xi_{\infty}} \begin{pmatrix} \eta \gamma \beta y & \\ & 1 \\ & & 1 \end{pmatrix}.$$

Since $\gamma\beta \in \mathfrak{q}^{-1}$, the map $\eta \mapsto \psi_{\infty}(\eta\gamma\beta)$ factors through $\Gamma_q = \{\epsilon \in \mathfrak{o}_F^{\times} : \epsilon \equiv 1 \pmod{\mathfrak{q}}\}$, and consequently

$$(4.1) \qquad \Phi_{\xi_{\infty},j}(y) = \sum_{\gamma \in \mathfrak{o}_{F}^{\times} \setminus \mathfrak{a}_{j}^{-1} \cap F^{\times}} a_{\xi_{j}^{0}}(\mathsf{a}_{j},\gamma) \sum_{\eta \in \Gamma_{\mathfrak{q}} \setminus \mathfrak{o}_{F}^{\times}} \psi_{\infty}(\eta \gamma \beta) \sum_{\epsilon \in \Gamma_{\mathfrak{q}}} W_{\xi_{\infty}} \begin{pmatrix} \epsilon \eta \gamma \beta y & \\ & 1 & \\ & & 1 \end{pmatrix}.$$

In particular, it follows that $\Phi_{\xi_{\infty},j}(y)$ is $\Gamma_{j}(\mathfrak{q})$ invariant.

4.2. Additive twists. For $\beta \in \mathfrak{a}_j\mathfrak{q}^{-1}$, $\xi_{\infty} \in V_{\pi_{\infty}}$, and with the rest of the notation as above, we define additive twist (the analogue of (3.7)) in this situation to be

$$\Lambda_{j}(s,\pi,\omega,\xi_{\infty},\beta) = \frac{N(\mathfrak{a}_{j})^{1-s}\|\beta\|_{\infty}^{s-1}}{[\mathfrak{o}_{F}^{\times}:\Gamma_{\mathfrak{q}}]} \int_{\Gamma_{\mathfrak{q}}\backslash F_{\infty}^{\times}} \Phi_{\xi_{\infty},j}(y)\omega_{\infty}(y)\|y\|_{\infty}^{s-1}d^{\times}y,$$

where ω_{∞} is the archimedean component of ω . We insert (4.1) into the above expression and collapse the integral and the sum over ϵ to get

$$\begin{split} \Lambda_j(s,\pi,\omega,\xi_\infty,\beta) &= \frac{N(\mathfrak{a}_j)^{1-s}\|\beta\|_\infty^{s-1}}{[\mathfrak{o}_F^\times : \Gamma_{\mathfrak{q}}]} \sum_{\gamma \in \mathfrak{o}_F^\times \backslash \mathfrak{a}_j^{-1} \cap F^\times} a_{\xi_f^0}(\mathsf{a}_j,\gamma) \sum_{\eta \in \Gamma_{\mathfrak{q}} \backslash \mathfrak{o}_F^\times} \psi_\infty(\eta\gamma\beta) \\ & \cdot \int_{F_\infty^\times} W_{\xi_\infty} \begin{pmatrix} \eta\gamma\beta y & 1 \\ & 1 \end{pmatrix} \omega_\infty(y) \|y\|_\infty^{s-1} d^\times y. \end{split}$$

Changing $y \mapsto (\gamma \eta \beta)^{-1} y$, we obtain

$$\begin{split} \Lambda_j(s,\pi,\omega,\xi_\infty,\beta) &= N(\mathfrak{a}_j)^{1-s} \sum_{\gamma \in \mathfrak{o}_F^\times \backslash \mathfrak{a}_j^{-1} \cap F^\times} \|\gamma\|_\infty^{1-s} a_{\xi_f^0}(\mathsf{a}_j,\gamma) \\ & \cdot \frac{1}{[\mathfrak{o}_F^\times \colon \Gamma_{\mathfrak{q}}]} \sum_{\eta \in \Gamma_{\mathfrak{q}} \backslash \mathfrak{o}_F^\times} \psi_\infty(\eta\gamma\beta) \omega_\infty(\eta\gamma\beta)^{-1} \int_{F_\infty^\times} W_{\xi_\infty} \begin{pmatrix} y & & \\ & 1 & \\ & & 1 \end{pmatrix} \omega_\infty(y) \|y\|_\infty^{s-1} d^\times y. \end{split}$$

In the notation of [2], the average over η is $e_{\mathfrak{q}}((\gamma\beta),\omega_{\infty}^{-1})$. Putting all of this together we obtain

$$\Lambda_j(s,\pi,\omega,\xi_\infty,\beta) = \sum_{\gamma \in \mathfrak{o}_F^\times \backslash \mathfrak{a}_i^{-1} \cap F^\times} N((\gamma)\mathfrak{a}_j)^{1-s} a_{\xi_j^0}(\mathsf{a}_j,\gamma) e_{\mathfrak{q}}((\gamma\beta),\omega_\infty^{-1}) \prod_{v \mid \infty} \Psi_v(s;W_{\xi_v},\omega_v).$$

For $\gamma \in \mathfrak{o}_F^{\times} \setminus \mathfrak{a}_i^{-1} \cap F^{\times}$, let $\mathfrak{a} = (\gamma)\mathfrak{a}_j$, then note that

$$\lambda_{\pi}(\mathfrak{o}_F, \mathfrak{a}) = \underset{26}{a_{\xi_f^0}}(\mathsf{a}_j, \gamma) N(\mathfrak{a}).$$

Consequently

$$\Lambda_j(s,\pi,\omega,\xi_\infty,\beta) = \sum_{\mathfrak{a}\sim\mathfrak{a}_j} \frac{\lambda_\pi(\mathfrak{o}_F,\mathfrak{a})e_{\mathfrak{q}}(\mathfrak{a}\mathfrak{a}_j^{-1}(\beta),\omega_\infty^{-1})}{N(\mathfrak{a})^s} \cdot \prod_{v\mid\infty} \Psi_v(s;W_{\xi_v},\omega_v).$$

Lemma 4.1. Let $\lambda_{\pi}(\mathfrak{a})$ denote the Dirichlet coefficients of $L(s,\pi)$, so that

$$L(s, \pi \otimes \omega) = \sum_{\mathfrak{a} \subset \mathfrak{o}_E} \frac{\lambda_{\pi}(\mathfrak{a}) \chi_{\omega}(\mathfrak{a})}{N(\mathfrak{a})^s}$$

for every unramified idèle class character ω . Then $\lambda_{\pi}(\mathfrak{a}) = \lambda_{\pi}(\mathfrak{o}_F, \mathfrak{a})$.

Proof. This essentially follows from Lemma 3.8, where we have identified the double Dirichlet coefficient $\lambda_{\pi}(\mathfrak{m}_1,\mathfrak{m}_2)$ of π . To be precise, by multiplicativity it suffices to verify the desired identity at prime powers. To that end, fix a prime \mathfrak{p} and let v be the corresponding place of F. Let $\alpha_1,\alpha_2,\alpha_3$ be the Langlands parameters of π_v and let $k \geq 0$ be any integer. From the proof of Lemma 3.8 we have $\lambda_{\pi}(\mathfrak{o}_F,\mathfrak{p}^k) = s_{k,0,0}(\alpha_1,\alpha_2,\alpha_3)$. On the other hand

$$s_{k,0,0}(\alpha_1,\alpha_2,\alpha_3) = \sum_{m_1+m_2+m_3=k} \alpha_1^{m_1} \alpha_2^{m_2} \alpha_3^{m_3},$$

where m_1, m_2 and m_3 are non-negative integers. The right-hand side of this expression is precisely $\lambda_{\pi}(\mathfrak{p}^k)$ thus proving our claim at prime powers.

Let $L_j(s, \pi, \omega, \xi_{\infty}, \beta)$ denote the finite part of $\Lambda_j(s, \pi, \omega, \xi_{\infty}, \beta)$. The following analogue of Theorem 3.4 for $GL(3) \times GL(1)$ is a consequence of [1, Proposition 3.1].

Corollary 4.2. For any idèle class character character ω , there are numbers $\beta_i \in F$ and $c_{ij} \in \mathbb{C}$ (depending on ω) such that

$$L(s, \pi \times \omega) = \sum_{i,j} c_{ij} L_j(s, \pi, \omega, \xi_{\infty}, \beta_i).$$

To end, we note the following corollary, which is another consequence of identifying the Dirichlet coefficient of $L(s,\pi)$ in terms of the associated essential function. Although this directly follows from (1.2) by a local calculation (see [19, Corollary 3.3]), the argument here is global in nature. Moreover, it can be extended to $GL(n) \times GL(m)$ for arbitrary n > m, i.e. we can identify the Dirichlet coefficients of $L(s, \pi \times \tau)$ for any pair (π, τ) in terms of the associated essential functions. We will investigate this for n > 3 in a forthcoming paper.

Corollary 4.3. Let $\xi \in V_{\pi}$ be a decomposable vector with $\xi_v = \xi_v^0$ for all finite v. Then

$$\int_{F^{\times}\backslash\mathbb{A}_{F}^{\times}} \mathbb{P}_{1}^{3}(U_{\xi}) \begin{pmatrix} h & & \\ & 1 & \\ & & 1 \end{pmatrix} \|h\|^{s-1} d^{\times}h = L(s,\pi) \prod_{v \mid \infty} \Psi_{v}(s; W_{\xi_{v}}),$$

where $\Psi_v(s; W_{\xi_v}) = \int_{F_v^{\times}} W_{\xi_v} \binom{a}{1} \|a\|_v^{s-1} d^{\times} a$.

Proof. We have the Fourier expansion $\mathbb{P}_1^3(U_\xi)(g) = \sum_{\gamma \in F^{\times}} W_{\xi}(\binom{\gamma}{1}_1)g$. Inserting this in the left-hand side above, we get

$$\int_{F^{\times}\backslash\mathbb{A}_{F}^{\times}} \sum_{\gamma} W_{\xi} \begin{pmatrix} \gamma h & \\ & 1 \\ & & 1 \end{pmatrix} \|h\|^{s-1} d^{\times} h.$$

Using $\mathbb{A}_F^{\times} = \coprod_i F^{\times} F_{\infty}^{\times} \mathsf{a}_i (\prod_{v < \infty} \mathfrak{o}_v^{\times})$, we may write this as

$$\sum_{j} \int_{\mathfrak{o}^{\times} \backslash F_{\infty}^{\times}} \sum_{\gamma \in \mathfrak{a}_{j}^{-1} \cap F^{\times}} W_{\xi_{f}^{0}} \begin{pmatrix} \gamma \mathsf{a}_{j} & & \\ & 1 & \\ & & 1 \end{pmatrix} W_{\xi_{\infty}} \begin{pmatrix} \gamma h_{\infty} & & \\ & 1 & \\ & & 1 \end{pmatrix} \| \mathsf{a}_{j} \|^{s-1} \| h \|_{\infty}^{s-1} \, d^{\times} h_{\infty},$$

which in turn is the same as

$$\sum_{j} \int_{\mathfrak{o}^{\times} \backslash F_{\infty}^{\times}} \sum_{\gamma \in \mathfrak{o}_{F}^{\times} \backslash \mathfrak{a}_{j}^{-1} \cap F^{\times}} a_{\xi_{f}^{0}}(\mathsf{a}_{j}, \gamma) \sum_{\eta \in \mathfrak{o}_{F}^{\times}} W_{\xi_{\infty}} \begin{pmatrix} \gamma \eta h_{\infty} & \\ & 1 \\ & & 1 \end{pmatrix} \|\mathsf{a}_{j}\|^{s-1} \|h\|_{\infty}^{s-1} \, d^{\times} h_{\infty}.$$

Thus the left-hand side becomes (by a similar calculation to the above)

$$\sum_{j} \sum_{\gamma \in \mathfrak{o}_F^\times \backslash \mathfrak{a}_j^{-1} \cap F^\times} a_{\xi_f^0}(\mathsf{a}_j,\gamma) \|\mathsf{a}_j\|^{s-1} \|\gamma\|_\infty^{1-s} \prod_{v \mid \infty} \Psi_v(s; W_{\xi_v}).$$

Since $a_{\xi_f^0}(a_j, \gamma) = N(\mathfrak{a})^{-1} \lambda_{\pi}(\mathfrak{o}_F, \mathfrak{a})$ for $(\gamma)\mathfrak{a}_j = \mathfrak{a}$, by Lemma 4.1, this is

$$\sum_{j} \sum_{\mathfrak{a} \sim \mathfrak{a}_{j}} \frac{\lambda_{\pi}(\mathfrak{a})}{N(\mathfrak{a})^{s}} \cdot \prod_{v \mid \infty} \Psi_{v}(s; W_{\xi_{v}}),$$

proving our assertion.

REFERENCES

- [1] Andrew R. Booker and M. Krishnamurthy, A strengthening of the GL(2) converse theorem, Compos. Math. 147 (2011), no. 3, 669–715.
- [2] ______, Further refinements of the GL(2) converse theorem, Bull. Lond. Math. Soc. 45 (2013), no. 5, 987–1003.
- [3] ______, Weil's converse theorem with poles, Int. Math. Res. Not. IMRN (2014), no. 19, 5328–5339.
- [4] ______, A converse theorem for GL(n), Adv. Math. **296** (2016), 154–180.
- [5] Daniel Bump, Lie groups, second ed., Graduate Texts in Mathematics, vol. 225, Springer, New York, 2013.
- [6] J. W. Cogdell and I. I. Piatetski-Shapiro, Converse theorems for GL_n, Inst. Hautes Études Sci. Publ. Math. (1994), no. 79, 157–214.
- [7] Cogdell, J. W. and Piatetski-Shapiro, I. I., Converse theorems for GL_n. II, J. Reine Angew. Math., 507 (1999), 165–188.
- [8] James W. Cogdell, L-functions and converse theorems for GL_n, Automorphic forms and applications, IAS/Park City Math. Ser., vol. 12, Amer. Math. Soc., Providence, RI, 2007, pp. 97–177.
- [9] Gelbart, Stephen S., Automorphic forms on adèle groups, Annals of Mathematics Studies, No. 83, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1975, pp. x+267.
- [10] H. Jacquet, I. I. Piatetski-Shapiro, and J. Shalika, Conducteur des représentations du groupe linéaire, Math. Ann. 256 (1981), no. 2, 199–214.
- [11] H. Jacquet, I. I. Piatetskii-Shapiro, and J. A. Shalika, Rankin-Selberg convolutions, Amer. J. Math. 105 (1983), no. 2, 367–464.
- [12] Hervé Jacquet, Archimedean Rankin-Selberg integrals, Automorphic forms and L-functions II. Local aspects, Contemp. Math., vol. 489, Amer. Math. Soc., Providence, RI, 2009, pp. 57–172.
- [13] _____, A correction to conducteur des représentations du groupe linéaire, Pacific J. Math. **260** (2012), no. 2, 515–525.
- [14] Hervé Jacquet, Ilja Iosifovitch Piatetski-Shapiro, and Joseph Shalika, Automorphic forms on GL(3). II, Ann. of Math. (2) 109 (1979), no. 2, 213–258.
- [15] Fabian Januszewski, Modular symbols for reductive groups and p-adic Rankin-Selberg convolutions over number fields, J. Reine Angew. Math. 653 (2011), 1–45.
- [16] Min Lee and Jens Marklof, Effective equidistribution of rational points on expanding horospheres, IMRN, Vol. 2017, No. 00, pp. 1–30
- [17] D. Kazhdan, B. Mazur, and C.-G. Schmidt, Relative modular symbols and Rankin-Selberg convolutions, J. Reine Angew. Math. 519 (2000), 97–141.
- [18] E. Kowalski, P. Michel and J. VanderKam, "Rankin-Selberg L-functions in the level aspect," Duke Math. J. 114 (2002), no. 1, 123–191.

- [19] Nadir Matringe, Essential Whittaker functions for GL(n), Doc. Math. 18 (2013), 1191–1214.
- [20] Toshitsune Miyake, *Modular forms*, English ed., Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2006, Translated from the 1976 Japanese original by Yoshitaka Maeda.
- [21] Miyauchi, Michitaka, Whittaker functions associated to newforms for GL(n) over p-adic fields, J. Math. Soc. Japan, **66** (2014), no. 1, 17–24.
- [22] Jacob Nemchenok, *Imprimitive Gussian sums and theta functions over a number field*, Trans. Amer. Math. Soc. **338** (1993), 465–478.
- [23] Tunnell, Jerrold B., On the local Langlands conjecture for GL(2), Invent. Math., 46, (1978), 2, 179–200.

E-mail address: andrew.booker@bristol.ac.uk

 $E ext{-}mail\ address: muthu-krishnamurthy@uiowa.edu}$

 $E ext{-}mail\ address: min.lee@bristol.ac.uk}$

School of Mathematics, University of Bristol, University Walk, Bristol, BS8 1TW, United Kingdom

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOWA, 14 MACLEAN HALL, IOWA CITY, IA 52242-1419, USA