Bounded Generalized Mixture Functions

Antonio Diego S. Farias^{a,b,**}, Valdigleis S. Costa^{a,*}, Luiz Ranyer de Araújo Lopes^a, Benjamín Bedregal^a, Regivan H. N. Santiago^a

^aGroup for Logic, Language, Information, Theory and Applications
Federal University of Rio Grande do Norte - UFRN, Department of Informatics and Applied Mathematics,
Avenue Senador Salgado Filho, 3000, Universitary Campus of Lagoa Nova, 59.078-970, Natal, RN, Brazil

^bGroup of Theory of Computation, Logic and Fuzzy Mathematics
Federal Rural University of Semi-Arid - UFERSA, Department of Exacts and Naturals Sciences
Highway 226, KM 405, São Geraldo, 59.900-000, Pau dos Ferros, RN, Brazil.

Abstract

In literature, it is common to find problems which require a way to encode a finite set of information into a single data; usually means are used for that. An important generalization of means are the so called Aggregation Functions, with a noteworthy subclass called OWA functions. There are, however, further functions which are able to provide such codification which do not satisfy the definition of aggregation functions; this is the case of pre-aggregation and mixture functions.

In this paper we investigate two special types of functions: *Generalized Mixture* and *Bounded Generalized Mixture* functions. They generalize both: OWA and Mixture functions. Both Generalized and Bounded Generalized Mixture functions are developed in such way that the weight vectors are variables depending on the input vector. A special generalized mixture operator, **H**, is provided and applied in a simple toy example.

Keywords: Aggregation functions, Pre-aggregation functions, OWA functions, Mixture functions, Generalized Mixture functions, Bounded Generalized Mixture functions

1. Introduction

Several applications require the conversion of a finite collection of data (of same type) into a single data [1, 2, 3, 4]. Some abstract tools which deal with that are the

^{*}Corresponding author

^{**}Principal corresponding author

Email addresses: antonio.diego@ufersa.edu.br(Antonio Diego S. Farias), valdigleis@gmail.com(Valdigleis S. Costa), ranyer.lopes@gmail.com(Luiz Ranyer de Araújo Lopes), bedregal@dimap.ufrn.br(Benjamín Bedregal), regivan@dimap.ufrn.br(Regivan H. N. Santiago)

Preprint submited to Information Fusion

Universidade Federal Rural do Semi-Árido - UFERSA, Campus Pau dos Ferros, BR 226, KM 405, São Geraldo, 59.900-000, Pau dos Ferros, RN, Brasil.

so called *Aggregation Functions* and *Mixture Functions* [5]. Yager [6] introduced a special class of aggregation functions, called *Orde Weighted Averaging* - OWA, and ever since several authors have proposed generalizations for them. Two generalizations are: (1) *Mixture functions* [5] and (2) *Generalized Mixture functions* [7, 8, 9]. They are not always aggregation functions, since some of them do not satisfy the monotonicity, however they are also efficient in order to "codify" a set of information into a sigle one. Some other extensions were also proposed and can be found in: [10, 11, 12, 13, 14].

In this paper we investigate *Generalized Mixture functions* (GM), which are weighted averaging means whose weights are dynamic; namely the weights are not fixed beforehand but depend on the input variables. This provides a more flexible usage of weights which is not possible for functions like OWAs. The resulting functions do not require the property of monotonicity, essencial in aggregations, instead they require diretional monotonicity[15]. Further we investigate the weakening of condition $\sum_{i=1}^{n} w_i = 1 \text{ to } \sum_{i=1}^{n} w_i \leq 1 \text{, thereby obtaining another generalization of OWAs, called } Bounded Generalized Mixture - BGM functions. This paper ends with the proposal of a special GM function, H, which have a wide range of properties like: idempotency, symmetry, homogeneity and diretional monotonicity. It is applyied on a simple toy example.$

This work is structured in the following way: The next section provides the basic concepts of Aggregation functions; section 3 introduces the concepts of *Generalized Mixture* (GM) and *Bounded Generalized Mixture* (BGM) operators, it shows properties, constructions, examples and propose a particular GM function: H; section 4 presents the final remarks and section 5 an illustrative application for GM.

2. Aggregation Functions

Aggregation functions are important mathematical tools for applications in various fields, such as: Information fuzzy [2, 16, 17]; Decision making [3, 18, 19]; Image processing [4, 20, 21] and Engineering [22].

Definition 1. An n-ary aggregation function is a mapping $A : [0,1]^n \to [0,1]$, which associates each n-dimensional vector $\mathbf{x} = (x_1, \dots, x_n)$ to a single value $A(\mathbf{x})$ and satisfies conditions of: (A1) mononicity (A2) Boundary:

(A1) If
$$\mathbf{x} \leq \mathbf{y}$$
, i.e., $x_i \leq y_i$, for all $i = 1, 2, ..., n$, then $A(\mathbf{x}) \leq A(\mathbf{y})$;

(A2)
$$A(0,...,0) = 0$$
 and $A(1,...,1) = 1$.

Example 1. Given $\mathbf{x} = (x_1, ..., x_n)$,

(a) Arithmetic Mean:
$$Arith(\mathbf{x}) = \frac{1}{n}(x_1 + x_2... + x_n)$$

(b) Minimum:
$$Min(\mathbf{x}) = min\{x_1, x_2, ..., x_n\};$$

(c) Maximum:
$$Max(\mathbf{x}) = max\{x_1, x_2, ..., x_n\};$$

(d) Product:
$$Prod(\mathbf{x}) = \prod_{i=1}^{n} x_i$$
;

(e) Weighted Average: For $\mathbf{w} = (w_1, \dots, w_n) \in [0, 1]^n$, with $\sum_{i=1}^n w_i = 1$, $WAvg_{\mathbf{w}}(\mathbf{x}) = \sum_{i=1}^n w_i \cdot x_i$.

Remark 1. From now on we will use the short term "aggregation" instead of "n-ary aggregation function".

Aggregations can be divided into four distinct classes: *Averaging, Conjunctive, Disjunctive* and *Mixed*. A wider description about them can be found in [23, 24, 25]. In this work, we only study averaging functions which satisfy the following property:

Definition 2. A function $f:[0,1]^n \longrightarrow [0,1]$ satisfies the **averaging property**, if for all $\mathbf{x} \in [0,1]^n$:

$$Min(\mathbf{x}) \le f(\mathbf{x}) \le Max(\mathbf{x}).$$

When aggregations satisfy the averaging property we say that they are **averaging aggregations**. The functions Min, Max, Arith and WAvg are examples of averaging aggregations. Besides that, an aggregation $A: [0,1]^n \to [0,1]$ can satisfy:

- (1) **Idempotency**, i.e., A(x,...,x) = x for all $x \in [0,1]$;
- (2) **Homogeneity** of order k, i.e., for all $\lambda \in [0,1]$ and $\mathbf{x} \in [0,1]^n$, $f(\lambda x_1, \lambda x_2, ..., \lambda x_n) = \lambda^k f(x_1, x_2, ..., x_n)$. When A is homogeneous of order 1 we simply say that f is homogeneous;
- (3) Shift-invariance, i.e., $f(x_1 + r, x_2 + r, ..., x_n + r) = f(x_1, x_2, ..., x_n) + r$, for all $r \in [-1, 1]$, $\mathbf{x} \in [0, 1]^n$, $(x_1 + r, x_2 + r, ..., x_n + r) \in [0, 1]^n$ and $f(x_1, x_2, ..., x_n) + r \in [0, 1]$;
- (4) **Monotonicity**, i.e., $f(x_1, \dots, x_n) \leq f(y_1, \dots, y_n)$ whenever $x_i \leq y_i$, for all $i \in \{1, \dots, n\}$;
- (5) **Strict monotonicity**, i.e., $f(\mathbf{x}) < f(\mathbf{y})$ whenever $\mathbf{x} < \mathbf{y}$, i.e., $\mathbf{x} \le \mathbf{y}$ and $\mathbf{x} \ne \mathbf{y}$, where $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_n)$;
- (6) **Symmetry**, i.e., its value is not changed under the permutations of the coordinates of x, i.e.,

$$f(x_1, ..., x_n) = f(x_{\sigma_{(1)}}, \cdots, x_{\sigma_{(n)}})$$

for all x and any permutation $\sigma: \{1, ..., n\} \rightarrow \{1, ..., n\}$;

(7) The existence of **neutral element**, i.e., there is $e \in [0, 1]$, such that for $t \in [0, 1]$ at any coordinate of the input vector \mathbf{x} , it has to be:

$$f(e, ..., e, t, e, ..., e) = t;$$

(8) The existence of **absorbing element** or (*annihilator*), i.e., there is $a \in [0,1]$, such that

$$f(x_1,...,x_{i-1},a,x_{i+1},...,x_n)=a;$$

- (9) The existence of **zero divisor**, i.e., there is $a \in]0,1[$, such that for any $\mathbf{x} \in]0,1]^n$, with a in one of its coordinates it is verified that $f(\mathbf{x}) = 0$;
- (10) The existence of **one divisor**, i.e., there is $a \in [0, 1[$, such that for any $\mathbf{x} \in]0, 1]^n$, with a in one of its coordinates it is verified that $f(\mathbf{x}) = 1$.

Example 2.

- (i) The functions: Arith, Min and Max are examples of idempotent, homogeneous, shift-invariant and symmetric aggregations.
- (ii) Min and Max have the elements 0 and 1 as its respective annihilators, whereas Arith does not have annihiladors.
- (iii) Min, Max and Arith does not have zero divisors and one divisors.
- 2.1. Ordered Weighted Averaging OWA Functions

In the field of aggregations there is a very important kind of function in which the aggregation behavior is provided parametrically; they are called: *Ordered Weighted Averaging* or simply OWA [6].

Definition 3. Let be an input vector $\mathbf{x} = (x_1, x_2, \dots, x_n) \in [0, 1]^n$ and a vector of weights $\mathbf{w} = (w_1, \dots, w_n) \in [0, 1]^n$, such that $\sum_{i=1}^n w_i = 1$. Assuming a permutation of \mathbf{x} :

$$sort(\mathbf{x}) = (x_{(1)}, x_{(2)}, \dots, x_{(n)})$$

such that $x_{(1)} \ge x_{(2)} \ge ... \ge x_{(n)}$. The Ordered Weighted Averaging (OWA) function with respect to \mathbf{w} , is the function OWA_w: $[0,1]^n \to [0,1]$ defined by:

$$OWA_{\mathbf{w}}(\mathbf{x}) = \sum_{i=1}^{n} w_i \cdot x_{(i)}$$

Remark 2. In what follows we remove w from $OWA_{\mathbf{w}}(\mathbf{x})$ and write only OWA.

Examples and properties of OWA functions can be found in [5]. Here is important to note that Min, Max, Arith and Median (described bellow) are examples of OWA.

Example 3. Given a vector \mathbf{x} and its ordered permutation $sort(\mathbf{x}) = (x_{(1)}, \dots, x_{(n)})$, the Median function

$$Med(\mathbf{x}) = \begin{cases} \frac{1}{2}(x_{(k)} + x_{(k+1)}), & \text{if } n = 2k\\ x_{(k+1)}, & \text{if } n = 2k+1 \end{cases}$$

is an OWA function in which the vector of weights is defined by:

- If n is odd, then $w_i = 0$ for all $i \neq \frac{n+1}{2}$ and $w_{\frac{n+1}{2}} = 1$.
- If n is even, then $w_i = 0$ for all $i \neq \frac{n}{2}$ and $i \neq \frac{n+2}{2}$, and $w_{\frac{n}{2}} = w_{\frac{n+2}{2}} = \frac{1}{2}$.

In addition there are some special types of OWAs: centered OWA or cOWA[10].

The OWA functions are defined in terms of a predetermined vector of weights; namely this vector of weights is fixed previously by the user. In the next section we present a generalized form of OWA in order to relax this situation. The vector of weights will be in function of the vector of inputs $\mathbf{x}=(x_1,\dots x_n)$. To achieve that we replace, in the OWA expression, the vector of weights by a family of functions, called **weighted functions**.

3. Weighted functions

As mentioned, the OWA functions are means with previously fixed weights. In the literature we can find some kind of functions which overcome this situation, by providing variable weights. This functions are called here *weighted functions*. An important example of that is the Mean of Bajraktarevic, presented in [23], and a particular case called of **Mixture function**, which are functions that have the following form:

$$M(\mathbf{x}) = \frac{\sum_{i=1}^{n} w_i(x_i) \cdot x_i}{\sum_{i=1}^{n} w_i(x_i)}$$
(1)

Generally, the mixture functions are not aggregation functions, since they do not always satisfy monotonicity, however the references [9, 26, 27] provide sufficient conditions to overcome this situation.

Remark 3. Note at equation 1 that each weight $w_i(x_i)$ is the value of single variable function; namely the weight is the value of a function w_i applied to the i-th position of the input vector $\mathbf{x} = (x_1, \dots, x_n)$. However, this restriction can be relaxed in order to obtain a weight $w_i(\mathbf{x})$, i.e., the weight is in function of the whole input. This generalization of mixture operators was done by Pereira [7, 8] and the resulting functions were called of **Generalized Mixture Functions (GMF)**.

Although Pereira has introduced GMFs he did not provided a deep investigation about them. In references [7, 8, 9] studied only GMF's depending on a single variable, that is, functions of the form $W(\mathbf{x}) = \sum_{i=1}^{n} w_i(x_i) \cdot x_i$. In what follows we provide some results about such GMFs; its relation to OWA's, Mixture Functions and Preaggregations. We finally generalize GMF's to the notion of Bounded Generalized Mixture Functions (BGMF) and provide some relations to the notions of monotonicity, directional monotonicity, Weak-dual and Weak-conjugate functions.

3.1. Weighted Averaging Functions

Definition 4. A finite family of functions $\Gamma = \{f_i : [0,1]^n \to [0,1] \mid 1 \le i \le n\}$ such that $\sum_{i=1}^n f_i(\mathbf{x}) = 1$ is called **family of weight-functions (FWF)**. A **Generalized**

Mixture Function or simply GM associated to a FWF Γ is a function $GM_{\Gamma}:[0,1]^n \to [0,1]$ given by:

$$GM_{\Gamma}(\mathbf{x}) = \sum_{i=1}^{n} f_i(\mathbf{x}) \cdot x_i$$

Example 4.

- 1. The GM operator associated to $\Gamma = \{f_i(\mathbf{x}) = \frac{1}{n} | 1 \le i \le n\}$, $GM_{\Gamma}(\mathbf{x})$, is $Arith(\mathbf{x})$;
- 2. The function Minimum can be obtained from $\Gamma = \{f_i \mid 1 \leq i \leq n\}$, where for all $\mathbf{x} \in [0,1]^n$, $f_{(n)}(\mathbf{x}) = 1$ and $f_i(\mathbf{x}) = 0$, if $i \neq (n)$, where $(\cdot) : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ is a permutation such that $sort(\mathbf{x}) = (x_{(1)}, \dots, x_{(n)})$;
- *3. Similarly, the function Maximum is also of type GM with* Γ *dually defined.*

Theorem 1. For any vector of weights $\mathbf{w} = (w_1, w_2, ..., w_n)$, the function OWA_w is a GM function.

Proof. Define $f_i(\mathbf{x}) = w_{p(i)}$, where $p: \{1, 2, \dots, n\} \longrightarrow \{1, 2, \dots, n\}$ is the inverse permutation of q(i) = (i). Then,

$$\mathsf{GM}_{\Gamma}(x_1, \dots, x_n) = \sum_{i=1}^n f_i(x_1, \dots, x_n) \cdot x_i$$
$$= \sum_{i=1}^n w_{p(i)} x_i$$

but as p is the inverse of q, it follows that q(p(i)) = i, that is, (p(i)) = i. Thus,

$$\mathsf{GM}_{\Gamma}(x_1,\cdots,x_n) = \sum_{i=1}^n w_{p(i)} x_{(p(i))}$$

Making the necessary changes in this sum, we have:

$$\begin{aligned}
\mathsf{GM}_{\Gamma}(x_1, \cdots, x_n) &= \sum_{i=1}^n w_i x_{(i)} \\
&= \mathsf{OWA}_{\mathbf{w}}(x_1, \cdots, x_n)
\end{aligned}$$

Example 5. If $\mathbf{w} = (0.3, 0.4, 0.3)$, then for $\mathbf{x} = (0.1, 1.0, 0.9)$ we have $x_1 = x_{(3)}$, $x_2 = x_{(1)}$ and $x_3 = x_{(2)}$. Thus, $f_1(\mathbf{x}) = 0.3$, $f_2(\mathbf{x}) = 0.3$, $f_3(\mathbf{x}) = 0.4$, and $GM(\mathbf{x}) = 0.3 \cdot 0.1 + 0.3 \cdot 1.0 + 0.4 \cdot 0.9 = 0.69$.

Note that, by Theorem 1, any OWA function is GM. However, there are GM functions which are not OWA:

Example 6. Let $\Gamma = \{\sin(x) \cdot y, 1 - \sin(x) \cdot y\}$. The respective GM function is $GM(x,y) = (\sin(x) \cdot y) \cdot x + (1 - \sin(x) \cdot y) \cdot y$, which is not an OWA function.

Proposition 1. Mixture functions are a particular case of GM.

Proof. A mixture operator $M(\mathbf{x})$ can be see as a GM function, with weight-functions given by $f_i(\mathbf{x}) = \frac{w_i(x_i)}{\sum\limits_{i=1}^n w_i(x_i)}$.

Remark 4. The GM function at Example 6 cannot be characterized as a mixture function, since w_1 is not a function which depends only on variable x and w_2 is not a function which depends only on variable y.

Below, we propose a new generalized form of GM, which is obtained by relaxing the condition $\sum\limits_{i=1}^n f_i(\mathbf{x}) = 1$ to $\sum\limits_{i=1}^n f_i(\mathbf{x}) \leq 1$. This new family is called Bounded Generalized Mixture functions.

Definition 5. A family of weak weight-functions (wFWF) is a finite family of functions

$$\Gamma = \{f_i : [0,1]^n \to [0,1] | 1 \le i \le n \}$$
 such that:

(I)
$$\sum_{i=1}^{n} f_i(\mathbf{x}) \leq 1$$
, and

(II)
$$\sum_{i=1}^{n} f_i(1,\dots,1) = 1$$
, for all $i \in \{1,2,\dots,n\}$.

A Bounded Ganeralized Mixture (BGM) operator associated to a wFWF Γ is a function $BGM_{\Gamma}: [0,1]^n \to [0,1]$ given by:

$$BGM_{\Gamma}(\mathbf{x}) = \sum_{i=1}^{n} f_i(\mathbf{x}) \cdot x_i$$

Remark 5.

1. Since BGM is a generalized form of GM, then both OWAs and mixture functions are instances of BGM. Moreover, GM functions are BGM operators subject to the condition:

(III)
$$\sum_{i=1}^{n} f_i(\mathbf{x}) = 1$$
, for any $\mathbf{x} \in [0, 1]^n$,

2. Let $\Gamma = \{f_i(x,y) = \frac{x}{n} | 1 \le i \le n\}$. Then, $BGM_{\Gamma}(\mathbf{x}) = \sum_{i=1}^n \frac{x_i^2}{n}$ is not a GM operator.

3.2. Properties of GM and BGM functions

Although GM and BGM are generalized forms of OWA, we cannot always guarantee that a BGM is an averaging function, although GM functions are averaging function. The next proposition gives us a sufficient condition to achieve that.

Proposition 2. If Γ is a FWF, then GM_{Γ} is averaging, i.e.:

$$Min(\mathbf{x}) \leq GM_{\Gamma}(\mathbf{x}) \leq Max(\mathbf{x})$$

Proof. For all $\mathbf{x} = (x_1, ..., x_n)$,

$$Min(\mathbf{x}) \le x_i \le Max(\mathbf{x}), \ \forall i = 1, 2, ..., n.$$

So,

$$\sum_{i=1}^{n} f_i(\mathbf{x}) \cdot Min(\mathbf{x}) \le \sum_{i=1}^{n} f_i(\mathbf{x}) \cdot x_i \le \sum_{i=1}^{n} f_i(\mathbf{x}) \cdot Max(\mathbf{x}),$$

but as $\sum_{i=1}^{n} f_i(\mathbf{x}) = 1$, it follows that

$$Min(\mathbf{x}) \le \sum_{i=1}^{n} f_i(\mathbf{x}) \cdot x_i \le Max(\mathbf{x})$$

Remark 6. Note that FWF cannot simply be replaced by wFWF, since for $f_1(x,y) = \frac{x}{2}$ and $f_2(x,y) = \frac{y}{2}$, we have BGM(0.5,0.5) = 0.25 < Min(0.5,0.5), although we can guarantee that:

$$BGM_{\Gamma}(\mathbf{x}) < Max(\mathbf{x})$$

Proposition 3. Let Γ be a wFWF. Then, the BGM $_{\Gamma}$ is idempotent if, and only, if $\sum_{i=1}^{n} f_i(x, \dots, x) = 1$ for any $x \in [0, 1]$.

Proof. If $\sum_{i=1}^{n} f_i(\mathbf{x}) = 1$ and $\mathbf{x} = (x, ..., x)$, then:

$$\mathsf{BGM}_{\Gamma}(\mathbf{x}) = \sum_{i=1}^{n} f_i(\mathbf{x}) \cdot x = x \cdot \sum_{i=1}^{n} f_i(\mathbf{x}) = x$$

Reciprocally, if BGM $_{\Gamma}$ is a idempotent function and $\sum_{i=1}^{n} f_i(x, \dots, x) < 1$ for some $x \in [0, 1]$ we have to

$$\mathsf{BGM}_{\Gamma}(\mathbf{x}) = \sum_{i=1}^{n} f_i(\mathbf{x}) \cdot x < x \cdot 1 = x.$$

Thus, the condition $\sum_{i=1}^{n} f_i(x, \dots, x) = 1$ cannot be removed.

Corollary 1. Any GM function is idempotent.

Remark 7. The BGM function described in Remark 6, is not idempotent, since BGM(0.5, 0.0) = 0.25. Thus, we cannot always guarantee that a BGM is idempotent

Proposition 4. If Γ is a FWF invariant under translations³, then GM_{Γ} is shift-invariant.

Proof. Let $\mathbf{x} = (x_1, ..., x_n) \in [0, 1]^n$ and $\lambda \in [-1, 1]$ such that $\mathbf{y} = (x_1 + \lambda, x_2 + \lambda, ..., x_n + \lambda) \in [0, 1]^n$. then,

$$\begin{aligned}
\mathsf{GM}_{\Gamma}(\mathbf{y}) &= \sum_{i=1}^{n} f_i(\mathbf{y}) \cdot (x_i + \lambda) \\
&= \sum_{i=1}^{n} f_i(\mathbf{y}) \cdot x_i + \sum_{i=1}^{n} f_i(\mathbf{y}) \cdot \lambda \\
&= \sum_{i=1}^{n} f_i(\mathbf{x}) \cdot x_i + \lambda \\
&= \mathsf{GM}_{\Gamma}(\mathbf{x}) + \lambda
\end{aligned}$$

Remark 8. The condition FWF is also important to preserve shift-invariance, since if we define $f_1(x,y) = f_2(x,y) = \frac{|x-y|}{2}$, for $(x,y) \neq (1,1)$, and $f_1(1,1) = f_2(1,1) = \frac{1}{2}$, then f_1 and f_2 are invariant under translations, but BGM(0,0.1) = 0.005 and $BGM(0+0.1,0.1+0.1) = 0.015 \neq 0.005+0.1$.

Proposition 5. If Γ is a wFWF and each $f_i \in \Gamma$ is homogeneous of order k, then $BGM_{\Gamma}(\mathbf{x})$ is homogeneous of order k+1.

Proof. The case $\lambda = 0$ is trivial. Now, to $\lambda \neq 0$ we have:

$$\begin{aligned} \mathsf{BGM}_{\Gamma}(\lambda x_1,...,\lambda x_n) &= \sum_{i=1}^n f_i(\lambda x_1,...,\lambda x_n) \cdot \lambda x_i \\ &= \lambda \cdot \sum_{i=1}^n \lambda^k f_i(x_1,...,x_n) x_i \\ &= \lambda^{k+1} \cdot \mathsf{BGM}_{\Gamma}(x_1,...,x_n) \end{aligned}$$

Remark 9. Note that if Γ is a FWF, i.e, $\sum_{i=1}^{n} f_i(\mathbf{x}) = 1$, then f_i cannot be homogeneous of order k > 0, since

$$1 = \sum_{i=1}^{n} f_i(\lambda x_1, \dots, \lambda x_n) = \lambda^k \sum_{i=1}^{n} f_i(\mathbf{x}) = \lambda^k,$$

This means that $f_i(x_1+\lambda,x_2+\lambda,...,x_n+\lambda)=f_i(x_1,x_2,...,x_n)$ for any $\mathbf{x}\in[0,1]^n$, for $i\in\{1,2,\cdots,n\}$ and $\lambda\in[-1,1]$ such that $(x_1+\lambda,x_2+\lambda,...,x_n+\lambda)\in[0,1]^n$.

i.e., there are no GM's homogeneous of order k > 1. However, if we remove this restriction, then we can have Γ with homogeneous f_i s of order k > 0. For example, $f_i(\mathbf{x}) = \frac{x_i}{n}$ is homogeneous of order 1, and so, according to Proposition 5, BGM $_{\Gamma}$ is homogeneous of order 2.

The next example shows a GM function which is not a mixture operator, since f_i does not depend exclusively from x_i . This GM_Γ is idempotent, homogeneous and shift-invariant, but is not monotonic, since $\mathsf{GM}_\Gamma(0.5, 0.2, 0.1) = 0.375$ and $\mathsf{GM}_\Gamma(0.5, 0.22, 0.2) = 0.368$.

Example 7. The family Γ defined by

$$f_i(x_1, ..., x_n) = \begin{cases} \frac{1}{n}, & \text{if } x_1 = \dots = x_n = 0\\ \frac{x_i}{\sum\limits_{j=1}^n x_j}, & \text{otherwise} \end{cases}$$

if a FWF and,

$$\mathit{GM}_{\Gamma}(\mathbf{x}) = \left\{ \begin{array}{ll} 0, & \textit{if } x_1, ..., x_n = 0 \\ \frac{\sum\limits_{i=1}^n x_i^2}{\sum\limits_{i=1}^n x_i}, & \textit{otherwise} \end{array} \right.$$

Proposition 6. The N-dual⁴, with respect to stantard fuzzy negation⁵, of a GM function is also a GM function.

Proof. If Γ is a wFWF, then

$$\begin{aligned}
\mathsf{GM}_{\Gamma}^{N}(x_{1}, \dots, x_{n}) &= 1 - \sum_{i=1}^{n} f_{i}(1 - x_{1}, \dots, 1 - x_{n}) \cdot (1 - x_{i}) \\
&= 1 - \sum_{i=1}^{n} f_{i}(1 - x_{1}, \dots, 1 - x_{n}) + \sum_{i=1}^{n} f_{i}(1 - x_{1}, \dots, 1 - x_{n}) \cdot x_{i} \\
&= \sum_{i=1}^{n} f_{i}(1 - x_{1}, \dots, 1 - x_{n}) \cdot x_{i} \\
&= \sum_{i=1}^{n} g_{i}(x_{1}, \dots, x_{n}) \cdot x_{i},
\end{aligned}$$

where $g_i(x_1, \dots, x_n) = f_i(1 - x_1, \dots, 1 - x_n)$.

Remark 10.

The N-dual of a function $F:[0,1]^n \longrightarrow [0,1]$ is $F^N(x_1,\cdots,x_n)=N(F(N(x_1),\cdots,N(x_n)),$ where N is a fuzzy negation, i.e., a decreasing function $N:[0,1] \longrightarrow [0,1]$ with N(0)=1 and N(1)=0.

The standard fuzzy negation if N(x)=1-x

1. The N-dual of a BGM with respect to stantard fuzzy negation it will not be a BGM function, but it will be of the form

$$BGM_{\Gamma}^{N}(\mathbf{x}) = BGM_{\Gamma'}(\mathbf{x}) + h(\mathbf{x}),$$

where
$$h:[0,1]^n \to [0,1]$$
 is given by $h(\mathbf{x}) = 1 - \sum_{i=1}^n f_i(\mathbf{x})$.

2. The dual of a GM function will not always be a GM function, for example: If $N(x) = 1 - x^n$, then

$$GM_{\Gamma}^{N}(x_{1}, \dots, x_{n}) = \sum_{i=1}^{n} f_{i}(1 - x_{1}^{n}, \dots, 1 - x_{n}^{n}) \cdot x_{i}^{n}$$

$$= \sum_{i=1}^{n} \left(f_{i}(1 - x_{1}^{n}, \dots, 1 - x_{n}^{n}) \cdot x_{i}^{n-1} \right) \cdot x_{i}$$

and $\Gamma' = \{g_i(\mathbf{x}) = f_i(1 - x_1^n, \dots, 1 - x_n^n) \cdot x_i^{n-1} | 1 \le i \le n \}$ not is a FWF, but is a wFWF. So, this N-dual is a BGM function.

This provides a motivation to define the **weak dual** of a GM function, as follow:

Definition 6. If GM_{Γ} is a generalized mixture function, then the **weak dual** of a GM_{Γ} with respect to fuzzy negations N is the function:

$$GM^{wN}_{\Gamma}(x_1,\cdots,x_n)=\sum_{i=1}^n f_i(N(x_1),\cdots,N(x_n))\cdot x_i$$

It is obvious that the weak dual of a GM functions is a GM function. Futhermore, we can define the weak dual of a BGM function, which is also a BGM function.

Example 8. The weak dual of GM defined in example 6 with respect to fuzzy negation $N(x) = 1 - x^{\alpha}$ is

$$GM_{\Gamma}^{wN}(x_1, \cdots, x_n) = \begin{cases} 0, & \text{if } x_1, \cdots, x_n = 0\\ \frac{n - \sum\limits_{i=1}^{n} (1 - x_i)^{\alpha} \cdot x_i}{n - \sum\limits_{i=1}^{n} (1 - x_i)^{\alpha}}, & \text{otherwise} \end{cases}$$

The construction of weak duals can be generalized, as follow:

Proposition 7. Let $\gamma_1, \dots, \gamma_n : [0,1] \longrightarrow [0,1]$ be functions. If GM_{Γ} is a GM (or BGM) function, then

$$GM_{\Gamma}^{\gamma_1,\dots,\gamma_n}(x_1,\dots,x_n) = \sum_{i=1}^n f_i(\gamma_1(x_1),\dots,\gamma_n(x_n)) \cdot x_i$$

is a GM (or BGM) function.

Proof. Straightforward.

When $\gamma_1 = \cdots = \gamma_n = \gamma$ are automorphism we say that $\mathsf{GM}_{\Gamma}^{\gamma_1, \dots, \gamma_n}$ is a **weak conjugate** of GM_{Γ} and we denote by $\mathsf{GM}_{\Gamma}^{\gamma}$.

Proposition 8. If $\Gamma=\{f_1,\cdots,f_n\}$ is a wFWF, then $\Gamma^R=\{g_1,\cdots,f_n\}$, whete $g_i=f_{n-i+1}$, also is a wFWF. Besides that, $\mathsf{BGM}_{\Gamma}^R=\mathsf{BGM}_{\Gamma^R}$

Proof. Sraightforward.

In Pereira [7, 8, 9] some criteria were considered to generate monotone GM. However, in this work we will not give a deep exposition of monotonicity, instead we present a brief discussion on a more weakened form, called **weak monotonicity** or **directional monotonicity**.

3.3. Directional Monotonicity

There are many *n*-ary functions that do not satisfy the monotonicity although, they are monotone with respect to certain directions. In this sence, Wilkin and Beliakov (in [28, 29]) introduceed the concept of **weakly monotonicity**, that was generalized by Bustince *et al.* in [15], which define the notion of **diretional monotonicity**.

Definition 7. Let $\mathbf{r} = (r_1, \dots, r_n)$ be a nnt null n-dimentional vector. A function $F: [0,1]^n \longrightarrow [0,1]$ is **r-increasing** if for all $\mathbf{x} = (x_1, \dots, x_n)$ and t > 0 such that $(x_1 + tr_1, \dots, x_n + tr_n) \in [0,1]^n$, we have

$$F(x_1, \dots, x_n) \le F(x_1 + tr_1, \dots, x_n + tr_n),$$

that is, F is increasing in the direction of vector \mathbf{r} .

Definition 8. A function $F: [0,1]^n \longrightarrow [0,1]$ is an n-ary **pre-aggregation** function (or simply pre-aggregation) if satisfies the boundary condition, $F(0,\dots,0)=0$ and $F(1,\dots,1)=1$, and is \mathbf{r} -increasing for some direction $\mathbf{r}\in [0,1]^n$.

Lucca *et al.* [30] presented properties, constructions and application for pre-aggregations. They show that the following functions are pre-aggregations:

Example 9.

- 1. $Mode(x_1, \dots, x_n)$ is (1, 1)-increasing;
- 2. $F(x,y) = x (max\{0, x y\})^2$ is (0,1)-increasing;
- *3. The weighted Lehmer mean (with convention* 0/0 = 0)

$$L_{\lambda}(x,y) = \frac{\lambda x^2 + (1-\lambda)y^2}{\lambda x + (1-\lambda)y}, \text{ where } 0 < \lambda < 1$$

is $(1 - \lambda, \lambda)$ -increasing;

4.

$$A(x,y) = \begin{cases} x(1-x), & if \ y \le 3/4\\ 1, & otherwise \end{cases}$$

is (0, a)-increasing for any a > 0, but for no other direction;

5.

$$B(x,y) = \begin{cases} y(1-y), & \text{if } x \le 3/4\\ 1, & \text{otherwise} \end{cases}$$

is (b,0)-increasing for any b>0, but for no other direction.

Remark 11. Any aggregation functions is also a pre-aggregation function.

Proposition 9. *If* BGM_{Γ} *is shift-invariant, is a pre-aggregation* (k, \dots, k) *-increasing.*

Proof. Note that for all $\mathbf{x} = (x_1, x_2, \dots, x_n) \in [0, 1]^n$ and any t > 0 such that $(x_1 + tk, x_2 + tk, \dots, x_n + tk) \in [0, 1]$ we have

$$\mathsf{BGM}_{\Gamma}(x_1+tk,\cdots,x_n+tk) = \mathsf{BGM}_{\Gamma}(x_1,\cdots,x_n)+tk,$$

and so

$$\mathsf{BGM}_{\Gamma}(x_1,\cdots,x_n) \leq \mathsf{BGM}(x_1+tk,\cdots,x_n+tk)$$

П

Corollary 2. If Γ is a FWF invariant under translations, i.e, $f_i(x_1 + \lambda, x_2 + \lambda, ..., x_n + \lambda) = f_i(x_1, x_2, ..., x_n)$, for $i \in \{1, 2, \cdots, n\}$, for any $\mathbf{x} = (x_1, \cdots, x_n) \in [0, 1]^n$ and $\lambda \in [0, 1]$ such that $(x_1 + \lambda, x_2 + \lambda, ..., x_n + \lambda) \in [0, 1]^n$, then BGM_{Γ} is a preaggregation (k, \cdots, k) -increasing.

Proof. By Proposition 5 BGM $_{\Gamma}$ is shift-invariant, and so, by Proposition 9, BGM $_{\Gamma}$ is a pre-aggregation function (k, k, \dots, k) -increasing.

In fact, the conditions required by Corollary 2 are very strong. In the following proposition, we relax these conditions:

Proposition 10. If Γ is a rFWF with $f_i(x_1, \dots, x_n) \leq f_i(x_1 + \lambda, \dots, x_i + \lambda)$, for $i \in \{1, 2, \dots, n\}$, for any $\mathbf{x} = (x_1, \dots, x_n) \in [0, 1]^n$ and $\lambda \in [0, 1]$ such that $(x_1 + \lambda, x_2 + \lambda, \dots, x_n + \lambda) \in [0, 1]^n$, then BGM_{Γ} is a pre-aggregation function (k, k, \dots, k) -increasing.

Proof. For any $\mathbf{x} = (x_1, \dots, x_n) \in [0, 1]^n$ and $\lambda \in [0, 1]$ such that $\mathbf{y} = (x_1 + \lambda, x_2 + \lambda, \dots, x_n + \lambda) \in [0, 1]^n$ we observe that

$$\begin{split} \mathsf{BGM}_{\Gamma}(\mathbf{y}) &= \sum_{i=1}^n f_i(\mathbf{y}) \cdot (x_i + \lambda) \\ &= \sum_{i=1}^n f_i(\mathbf{y}) \cdot x_i + \sum_{i=1}^n f_i(\mathbf{y}) \cdot \lambda \\ &\geq \sum_{i=1}^n f_i(\mathbf{x}) \cdot x_i + \lambda \\ &\geq \mathsf{BGM}_{\Gamma}(\mathbf{x}) \end{split}$$

Example 10. Let Γ be the family of functions:

$$f_i(x_1, \dots, x_n) = \begin{cases} \frac{1}{n}, & \text{if } x_1 = \dots = x_n \\ \frac{x_{(1)} - x_i}{\sum_{j=1}^n (x_{(1)} - x_j)}, & \text{otherwise} \end{cases}.$$

We can easily prove that all those functions satisfy:

$$f_i(x_1 + \lambda, x_2 + \lambda, \dots, x_n + \lambda) = f_i(x_1, x_2, \dots, x_n).$$

More generally, for any $\alpha \geq 1$

$$f_i(x_1, \dots, x_n) = \begin{cases} \frac{1}{n}, & \text{if } x_1 = \dots = x_n \\ \frac{x_{(1)} - x_i}{\sum_{j=1}^n (x_{(1)} - x_j)^{\alpha}}, & \text{otherwise} \end{cases}$$

is such that

$$f_i(x_1, x_2, \cdots, x_n) \le f_i(x_1 + \lambda, x_2 + \lambda, \cdots, x_n + \lambda).$$

Thus, the corresponding BGM is (k, \dots, k) -increasing. In additon, note that, for $\alpha > 1$, $\Gamma = \{f_i\}$ does not satisfies $\sum_{i=1}^n f_i(\mathbf{x}) = \mathbf{1}$.

We can also establish a criterion analogous to Proposition 10, by replacing the vector (k, \dots, k) by direction \mathbf{r} , as follow:

Proposition 11. If Γ is a FWF such that there is a directional vector $\mathbf{r} = (r_1, r_2, \cdots, r_n) \in [0, 1]^n$ with $f_i(x_1, \cdots, x_n) \leq f_i(x_1 + \lambda \cdot r_1, \cdots, x_i + \lambda \cdot r_n)$, for $i \in \{1, 2, \cdots, n\}$, for any $\mathbf{x} = (x_1, \cdots, x_n) \in [0, 1]^n$ and $\lambda \in [0, 1]$ such that $(x_1 + \lambda \cdot r_1, x_2 + \lambda \cdot r_2, ..., x_n + \lambda \cdot r_n) \in [0, 1]^n$, then BGM_{Γ} is a pre-aggregation function \mathbf{r} -increasing.

Proof. Similar to Proposition 10.

Corollary 3. If Γ is a wFWF such that there is a directional vector \mathbf{r} with $\frac{\partial f_i}{\partial \mathbf{r}}(\mathbf{x}) \geq 0$ for any $f_i \in \Gamma$ and $\mathbf{x} \in [0,1]^n$, then BGM_{Γ} is a pre-aggregation function \mathbf{r} -increasing.

Example 11. If $f_i = w_i$ is constant, then BGM_{Γ} is **r**-increasing for any direction **r**. Now, given a direction $\mathbf{r} = (r_1, \dots, r_n) \in [0, 1]^n$ we can build a **r**-increasing BGM function defining:

$$f_i(x_1,\dots,x_n) = \begin{cases} 0, & \text{if } x_{(n)} = 0\\ \frac{\min\left\{\frac{x_i}{r_i},1\right\}}{n}, & \text{otherwise} \end{cases}$$

In what follows we propose a new GM function which will be investigated and applied in this paper.

Example 12. *Definition: Let* Γ *be the following family of functions:*

$$f_i(\mathbf{x}) = \begin{cases} \frac{1}{n}, & \text{if } \mathbf{x} = (x, ..., x) \\ \frac{1}{n-1} \left(1 - \frac{|x_i - Med(\mathbf{x})|}{\sum\limits_{j=1}^{n} |x_j - Med(\mathbf{x})|} \right), & \text{otherwise} \end{cases}$$

 Γ is a FWF, whose GM function, denoted by H. The computation of H can be performed using the following expressions:

$$\mathbf{H}(\mathbf{x}) = \begin{cases} x, & \text{if } \mathbf{x} = (x, ..., x) \\ \frac{1}{n-1} \sum_{i=1}^{n} \left(x_i - \frac{x_i | x_i - Med(\mathbf{x})|}{\sum\limits_{j=1}^{n} | x_j - Med(\mathbf{x})|} \right), & \text{otherwise} \end{cases}$$

An interesting property that function is that:

$$\mathbf{H}(\mathbf{x}) = \mathsf{OWA}_{(f_1(\mathbf{x}), \cdots, f_n(\mathbf{x}))}(\mathbf{x})$$

In the next subsection we discuss others properties of the function H.

3.4. Properties of H

In this part of paper we will discuss about the properties of operator **H**. It is easy to check that $\sum_{i=1}^n f_i(\mathbf{x}) = 1$ for any $\mathbf{x} \in [0,1]^n$ and therefore, **H** is an idempotent averaging function. Furthermore, its weight-functions are invariant under translations and is also homogeneous of order 0, because:

1. For $\mathbf{y} = (x_1 + \lambda, ..., x_n + \lambda)$ we have $Med(\mathbf{x}') = Med(\mathbf{x}) + \lambda$ and for $\mathbf{x} \neq (x, ..., x)$. So,

$$f_{i}(\mathbf{y}) = \frac{1}{n-1} \left(1 - \frac{|x_{i} + \lambda - Med(\mathbf{y})|}{\sum_{j=1}^{n} |x_{j} + \lambda - Med(\mathbf{x}')|} \right)$$

$$= \frac{1}{n-1} \left(1 - \frac{|x_{i} + \lambda - (Med(\mathbf{x}) + \lambda)|}{\sum_{j=1}^{n} |x_{j} + \lambda - (Med(\mathbf{x}) + \lambda)|} \right)$$

$$= \frac{1}{n-1} \left(1 - \frac{|x_{i} - Med(\mathbf{x})|}{\sum_{j=1}^{n} |x_{j} - Med(\mathbf{x})|} \right)$$

$$= f_{i}(\mathbf{x}).$$

Therefore, $(f_1(y), ..., f_n(y)) = (f_1(x), ..., f_n(x)).$

- 2. The case in which $\mathbf{x} = (x, ..., x)$ is immediate.
- 3. To check the second property, make $\mathbf{y}=(\lambda x_1,...,\lambda x_n)$, note that $Med(\mathbf{y})=\lambda Med(\mathbf{x})$ and for $\mathbf{x}\neq(x,...,x)$

$$f_{i}(\mathbf{y}) = \frac{1}{n-1} \left(1 - \frac{|\lambda x_{i} - Med(\lambda \mathbf{x})|}{\sum\limits_{j=1}^{n} |\lambda x_{j} - Med(\lambda \mathbf{x})|} \right)$$

$$= \frac{1}{n-1} \left(1 - \frac{|\lambda x_{i} - \lambda Med(\mathbf{x})|}{\sum\limits_{j=1}^{n} |\lambda x_{j} - \lambda Med(\mathbf{x})|} \right)$$

$$= \frac{1}{n-1} \left(1 - \frac{|\lambda| \cdot |x_{i} - Med(\mathbf{x})|}{|\lambda| \cdot \sum\limits_{j=1}^{n} |x_{j} - Med(\mathbf{x})|} \right)$$

$$= f_{i}(\mathbf{x})$$

Therefore, $(f_1(\mathbf{x''}), ..., f_n(\mathbf{x''})) = (f_1(\mathbf{x}), ..., f_n(\mathbf{x})) = f(\mathbf{x}).$

4. The case in which $\mathbf{x}=(x,...,x)$ is also immediately. Note that the case in which $\lambda=0$ is obvious.

Corollary 4. H is shift-invariant and homogeneous.

In addition to idempotency, homogeneity and shift-invariance ${\bf H}$ has the following properties.

Proposition 12. H has no neutral element.

Proof. Suppose **H** has a neutral element e, find the vector of weight for $\mathbf{x} = (e, ..., e, x, e, ..., e)$. Note that if $n \geq 3$, then $Med(\mathbf{x}) = e$ and therefore,

$$f_i(\mathbf{x}) = \frac{1}{n-1} \left(1 - \frac{|x_i - Med(\mathbf{x})|}{\sum\limits_{j=1}^n |x_j - Med(\mathbf{x})|} \right)$$
$$= \frac{1}{n-1} \left(1 - \frac{|x_i - e|}{\sum\limits_{j=1}^n |x_j - e|} \right)$$
$$= \frac{1}{n-1} \left(1 - \frac{|x_i - e|}{|x - e|} \right)$$

therefore,

$$f_i(\mathbf{x}) = \left\{ \begin{array}{l} \frac{1}{n-1}, \text{ if } x_i = e \\ 0, \text{ if } x_i = x \end{array} \right., \text{ to } n \geq 3$$

i.e.,

$$f(\mathbf{x}) = \left(\frac{1}{n-1}, ..., \frac{1}{n-1}, 0, \frac{1}{n-1}, ..., \frac{1}{n-1}\right)$$

and

$$\mathbf{H}(\mathbf{x}) = (n-1) \cdot \frac{e}{n-1} = e$$

But since e is a neutral element of \mathbf{H} , $\mathbf{H}(\mathbf{x}) = x$. Absurd, since we can always take $x \neq e$.

For n=2, we have $Med(\mathbf{x})=\frac{x+e}{2}$, where $\mathbf{x}=(x,e)$ or $\mathbf{x}=(e,x)$. In both cases it is not difficult to show that $f(\mathbf{x})=(0.5,0.5)$ and $\mathbf{H}(\mathbf{x})=\frac{x+e}{2}$. Thus, taking $x\neq e$, again we have $\mathbf{H}(x,e)\neq x$.

Proposition 13. H has no absorbing elements.

Proof. To n=2, we have $\mathbf{H}(\mathbf{x})=\frac{x_1+x_2}{2}$, which has no absorbing elements. Now for $n\geq 3$ we have to $\mathbf{x}=(a,0,...,0)$ with $Med(\mathbf{x})=0$ therefore,

$$f_1(\mathbf{x}) = \frac{1}{n-1} \left(1 - \frac{a}{a} \right) = 0$$
 and $f_i = \frac{1}{n-1}, \forall i = 2, ..., n$.

therefore,

$$\mathbf{H}(a,0,...,0) = 0 \cdot a + \frac{1}{n-1} \cdot 0 + ... + \frac{1}{n-1} \cdot 0 = a \Rightarrow a = 0,$$

but to $\mathbf{x} = (a, 1, ..., 1)$ we have to $Med(\mathbf{x}) = 1$. Furthermore,

$$f_1(\mathbf{x}) = \frac{1}{n-1} \left(1 - \frac{1-a}{1} - a \right) = 0$$

and

$$f_i = \frac{1}{n-1}$$
 para $i = 2, 3, ..., n$.

therefore,

$$\mathbf{H}(a, 1, ..., 1) = 0 \cdot a + \frac{1}{n-1} \cdot 1 + ... + \frac{1}{n-1} \cdot 1 = a \Rightarrow a = 1.$$

With this we prove that **H** does not have absorbing elements.

Proposition 14. H has no zero divisors.

Proof. Let $a \in]0,1[$ and consider $\mathbf{x}=(a,x_2,...,x_n)\in]0,1]^n$. In order to have $\mathbf{H}(\mathbf{x})=\sum_{i=1}^n f_i(\mathbf{x})\cdot x_i=0$ we have $f_i(\mathbf{x})\cdot x_i=0$ for all i=1,2,...,n. But as $a\neq 0$ and we can always take $x_2,x_3,...,x_n$ also different from zero, then for each i=1,2,...,n there remains only the possibility of terms:

$$f_i(\mathbf{x}) = 0$$
 para $i = 1, 2, ..., n$.

This is an absurd, for $f_i(\mathbf{x}) \in [0,1]$ e $\sum_{i=1}^n f_i(\mathbf{x}) = 1$. like this, \mathbf{H} has no zero divisors.

Proposition 15. H does not have one divisors

Proof. Just to see that $a \in [0, 1[$, we have to $\mathbf{H}(a, 0, ..., 0) = f_1(\mathbf{x}).a \leq a < 1.$

Proposition 16. H is symmetric.

Proof. Let $P:\{1,2,...,n\} \rightarrow \{1,2,...,n\}$ be a permutation. So we can easily see that

$$Med(x_{P(1)}, x_{P(2)}, ..., x_{P(n)}) = Med(x_1, x_2, ..., x_n)$$

for all $\mathbf{x} = (x_1, x_2, ..., x_n) \in [0, 1]^n$. We also have to $\sum_{i=1}^n |x_{P(i)} - Med(x_{P(1)}, x_{P(2)}, ..., x_{P(n)})| = \sum_{i=1}^n |x_i - Med(\mathbf{x})|$. Thus, it suffices to consider the case where $\mathbf{y} = (x_{P(1)}, x_{P(2)}, ..., x_{P(n)}) \neq (x, x, ..., x)$. But for $\mathbf{y} \neq (x, x, ..., x)$ we have to:

$$\begin{aligned} \mathbf{H}(\mathbf{y}) &= \frac{1}{n-1} \sum_{i=1}^{n} \left(x_{P(i)} - \frac{x_{P(i)}|x_{P(i)} - Med(\mathbf{y})|}{\sum\limits_{j=1}^{n} |x_{P(i)} - Med(\mathbf{y})|} \right) \\ &= \frac{\sum\limits_{i=1}^{n} x_{P(i)}}{n-1} - \frac{1}{n-1} \cdot \sum\limits_{i=1}^{n} \frac{x_{P(i)}|x_{P(i)} - Med(\mathbf{x})|}{\sum\limits_{j=1}^{n} |x_{P(i)} - Med(\mathbf{x})|} \\ &= \frac{\sum\limits_{i=1}^{n} x_{i}}{n-1} - \frac{1}{n-1} \cdot \sum\limits_{i=1}^{n} \frac{x_{P(i)}|x_{P(i)} - Med(\mathbf{x})|}{\sum\limits_{j=1}^{n} |x_{i} - Med(\mathbf{x})|} \\ &= \frac{\sum\limits_{i=1}^{n} x_{i}}{n-1} - \frac{1}{n-1} \cdot \sum\limits_{i=1}^{n} \frac{x_{i}|x_{i} - Med(\mathbf{x})|}{\sum\limits_{j=1}^{n} |x_{i} - Med(\mathbf{x})|} \\ &= \mathbf{H}(\mathbf{x}). \end{aligned}$$

Proposition 17. If $N:[0,1] \longrightarrow [0,1]$ is the standard fuzzy negation, then $\mathbf{H}^N = \mathbf{H}$. Proof. If $\mathbf{x} = (x, \dots, x)$, then

$$\mathbf{H}^{N}(\mathbf{x}) = 1 - \mathbf{H}(1 - x, 1 - x, \dots, 1 - x)$$

= $1 - (1 - x) = x = \mathbf{H}(\mathbf{x})$

For $\mathbf{x} \neq (x, \dots, x)$ and $\mathbf{y} = (1 - x_1, \dots, 1 - x_n)$, we have:

$$\mathbf{H}^{N}(\mathbf{x}) = 1 - \frac{1}{n-1} \sum_{i=1}^{n} \left(1 - x_{i} - \frac{(1-x_{i})|1-x_{i}-Med(\mathbf{y})|}{\sum_{j=1}^{n} |1-x_{i}-Med(\mathbf{y})|} \right)$$

$$= 1 - \frac{1}{n-1} \sum_{i=1}^{n} \left(1 - x_{i} - \frac{(1-x_{i})|1-x_{i}-1+Med(\mathbf{x})|}{\sum_{j=1}^{n} |1-x_{i}-1+Med(\mathbf{x})|} \right)$$

$$= 1 - \frac{1}{n-1} \sum_{i=1}^{n} \left(1 - x_{i} - \frac{(1-x_{i})|-x_{i}+Med(\mathbf{x})|}{\sum_{j=1}^{n} |-x_{i}+Med(\mathbf{x})|} \right)$$

$$= 1 - \frac{1}{n-1} \sum_{i=1}^{n} \left(1 - x_i - \frac{(1-x_i)|x_i - Med(\mathbf{x})|}{\sum_{j=1}^{n} |x_i - Med(\mathbf{x})|} \right)$$

$$= 1 - \frac{1}{n-1} \left[n - \sum_{i=1}^{n} \left(x_i - \frac{x_i|x_i - Med(\mathbf{x})|}{\sum_{j=1}^{n} |x_i - Med(\mathbf{x})|} \right) - \sum_{i=1}^{n} \frac{|x_i - Med(\mathbf{x})|}{\sum_{j=1}^{n} |x_i - Med(\mathbf{x})|} \right]$$

$$= 1 - \frac{1}{n-1} \left[n - 1 - \sum_{i=1}^{n} \left(x_i - \frac{x_i|x_i - Med(\mathbf{x})|}{\sum_{j=1}^{n} |x_i - Med(\mathbf{x})|} \right) \right]$$

$$= \frac{1}{n-1} \sum_{i=1}^{n} \left(x_i - \frac{x_i|x_i - Med(\mathbf{x})|}{\sum_{j=1}^{n} |x_i - Med(\mathbf{x})|} \right)$$

$$= \mathbf{H}(\mathbf{x})$$

Proposition 18. If k > 0, then **H** is (k, \dots, k) -increasing.

Proof. As **H** is shift-invariant, its follow that **H** is (k, \dots, k) -increasing.

Corollary 5. H is a pre-aggregation function.

Therefore, **H** satisfies the following properties:

- Idempotency;
- Homogeneity;
- Shift-invariance;
- Symmetry;
- has no neutral element;
- has no absorbing elements;
- has no zero divisors;
- does not have one divisors;
- is self dual;
- is a preagregation (k, \dots, k) -increasing.

Aggregation functions are very important for Computer Science, since in many applications the expected result is a single data, and therefore they usually use aggregation functions to convert this set of data into a unique output. In fact, pre-aggregation can also be applied. In this sense, the Appendix contains a simple application which apply GM functions on the problem of image reduction.

4. Final remarks

In this paper we study two generalized forms of Ordered Weighted Averaging function and Mixture function, called **Generalized Mixture** and **Bounded Generalized Mixture** functions. This functions are defined by weights, which are obtained dynamically from of each input vector $\mathbf{x} \in [0,1]^n$. We demonstrated, among other results, that OWA and mixture functions are particular cases of GM and BGM functions, and thus functions likes *Arithmetic Mean*, *Median*, *Maximum*, *Minimum* and cOWA are also instances of GM function.

In the second part of this work, we present some properties as well as constructs and examples of GM functions. In particular we define a special GM function, called \mathbf{H} . We show that \mathbf{H} satisfies important properties like: Idempotency, symmetry, homogeneity, shift-invariance; it does not have neither zero and one divisors nor neutral elements. We further prove that \mathbf{H} is a pre-aggregation (k, \dots, k) -increasing.

A illustrative application is presented in the Appendix. A further insight into the applications of these functions will be addressed in future works.

References

References

- [1] D. Dubois, H. Prade, On the use of aggregation operations in information fusion processes, Fuzzy Sets and Systems 142 (2004) 143 161. Aggregation Techniques.
- [2] A. D. S. Farias, L. R. A. Lopes, B. C. Bedregal, R. H. N. Santiago, Closure properties for fuzzy recursively enumerable languages and fuzzy recursive languages,, Journal of Intelligent & Fuzzy Systems 31 (2016) 1795–1806.
- [3] D. Paternain, A. Jurio, E. Barrenechea, H. Bustince, B. Bedregal, E. Szmidt, An alternative to fuzzy methods in decision-making problems, Expert Systems with Applications 39 (2012) 7729 7735.
- [4] D. Paternain, J. Fernandez, H. Bustince, R. Mesiar, G. Beliakov, Construction of image reduction operators using averaging aggregation functions, Fuzzy Sets and Systems 261 (2015) 87 111. Theme: Aggregation operators.
- [5] G. Beliakov, H. Bustince, T. Calvo, A Practical Guide to Averaging Functions, volume 329 of *Studies in Fuzziness and Soft Computing*, Springer, 2016.
- [6] R. R. Yager, Ordered weighted averaging aggregation operators in multicriteria decision making, IEEE Transactions on Systems, Man, and Cybernetics 18 (1988) 183 – 190.
- [7] R. A. M. Pereira, G. Pasi, On non-monotonic aggregation: mixture operators, in: Proc. 4th Meeting of the EURO Working Group on Fuzzy Sets (EUROFUSE'99) and 2nd Internat. Conf. on Soft and Intelingent Computing (SIC'99), Budapest, Hungary, 1999.

- [8] R. A. M. Pereira, The orness of mixture operators: the exponential case, in: Proc. 8th Internat. Conf. on Information Processing and Management of Uncertainty in Knowledge-Based Systems (IPMU'200), Madrid, Spain, 2000.
- [9] R. A. M. Pereira, R. A. Ribeiro, Aggregation with generalized mixture operators using weighting functions, Fuzzy Sets and Systems 137 (2003) 43 58. Preference Modelling and Applications.
- [10] R. R. Yager, Centered OWA operators, Soft Computing 11 (2006) 631–639.
- [11] I. Lizasoain, C. Moreno, Owa operators defined on complete lattices, Fuzzy Sets and Systems 224 (2013) 36 52. Theme: Aggregation functions and implications.
- [12] B. Llamazares, Constructing choquet integral-based operators that generalize weighted means and owa operators, Information Fusion 23 (2015) 131 138.
- [13] Z. Xu, Induced uncertain linguistic owa operators applied to group decision making, Information Fusion 7 (2006) 231 238.
- [14] R. R. Yager, Lexicographic ordinal owa aggregation of multiple criteria, Information Fusion 11 (2010) 374 380.
- [15] H. Bustince, J. Fernandez, A. Kolesárová, R. Mesiar, Directional monotonicity of fusion functions, European Journal of Operational Research 244 (2015) 300 308.
- [16] E. Hancer, B. Xue, M. Zhang, D. Karaboga, B. Akay, A multi-objective artificial bee colony approach to feature selection using fuzzy mutual information, in: Evolutionary Computation (CEC), 2015 IEEE Congress on, 2015, pp. 2420–2427.
- [17] L. Lingling, Z. Xian, H. Pengju, L. Zhigang, The research on the method of fuzzy information processing, in: System Science, Engineering Design and Manufacturing Informatization (ICSEM), 2012 3rd International Conference on, volume 2, 2012, pp. 47–50.
- [18] H. Bustince, M. Galar, B. Bedregal, A. Kolesárová, R. Mesiar, A new approach to interval-valued choquet integrals and the problem of ordering in interval-valued fuzzy set applications, IEEE Transactions on Fuzzy Systems 21 (2013) 1150– 1162.
- [19] R. R. Yager, G. Gumrah, M. Z. Reformat, Using a web personal evaluation tool pet for lexicographic multi-criteria service selection, Knowledge-Based Systems 24 (2011) 929 942.
- [20] G. Beliakov, H. Bustince, D. Paternain, Image reduction using means on discrete product lattices, IEEE Transactions on Image Processing 21 (2012).
- [21] R. P. Joseph, C. S. Singh, M. Manikandan, Brain tumor MRI image segmentation and detection in image processing, International Journal of Research and Tecnology 3 (2014).

- [22] X. Liang, W. Xu, Aggregation method for motor drive systems, Electric Power Systems Research 117 (2014) 27 35.
- [23] G. Beliakov, H. Bunstince, T. Calvo, A Practical Guide to Averaging Functions, volume 329 of *Studies in Fuzziness and Soft Computing*, 1 ed., Springer, Switzeland, 2016.
- [24] D. Dubois, H. Prade, Fundamentals of Fuzzy Sets, volume 7 of *The Handbooks of Fuzzy Sets*, 1 ed., Springer, New York, 2000.
- [25] M. Grabisch, J. L. Marichal, R. Mesiar, E. Pap, Aggregation Functions, volume 127 of *Encyclopedia of Mathematics and its Applications*, University Press Cambridge, 2009.
- [26] R. Mesiar, J. Špirková, Weighted means and weighting functions, Kybernetika 42 (2006) 151-160.
- [27] R. Mesiar, J. Špirková, L. Vavríková, Weighted aggregation operators based on minimization, Information Sciences 178 (2008) 1133 1140.
- [28] T. Wilkin, G. Beliakov, Weakly monotone averaging functions, in: IPMU'2014, volume 444, 2014, pp. 364–373.
- [29] T. Wilkin, G. Beliakov, Weakly monotone aggregation functions, International Journal of Intelligent. Systems 30 (2015) 144–169.
- [30] G. Lucca, J. A. Sanz, G. P. Dimuro, B. Bedregal, R. Mesiar, A. Kolesárová, H. Bustince, Preaggregation functions: Construction and an application, IEEE Transactions on Fuzzy Systems 24 (2016) 260–272.
- [31] R. C. Gonzales, R. E. Woods, Digital Image Processing, 3rd ed., Pearson, New Jersey, 2008.
- [32] R. Keys, Cubic convolution interpolation for digital image processing, IEEE Transactions on Acoustics, Speech, and Signal Processing 29 (1981) 1153–1160.
- [33] T. M. Lehmann, C. Gonner, K. Spitzer, Survey: interpolation methods in medical image processing, IEEE Transactions on Medical Imaging 18 (1999) 1049–1075.
- [34] P. Thevenaz, T. Blu, M. Unser, Interpolation revisited, IEEE Transactions on Medical Imaging 19 (2000) 739–758.

5. Appendix: Illustrative example of application (Image reduction)

In this part of our work we use the GM functions Min, Max, Arith, Med, cOWA and H to build image reduction operators. A broad discussion on this field of application can be found in [31]. Here, we are just interested to shows a possibility of application for our functions.

5.1. Methodological process

The methodological process is the same as that used in [4], it consists of:

- 1. Reduce the input images using the Min, Max, Arith, Med, cOWA and H;
- 2. Magnify the reduced images to the original size using three different method: (1) The nearest neighbor interpolation, (2) The bilinear interpolation and (2) the bicubic interpolation (see [31, 32, 33, 34]);
- 3. Compare the output images with the original one using the measure PSNR.

We use ten original images, in grayscale, of size 512×512 . The obtained results are shown below (The bold value represents the high quality image, and the italic value represents the second high quality image):

5.2. Results

USING 2 \times 2 BLOCKS USING 4 \times 4 BLOCKS

	Min	Max	Med	Arith	cOWA	H	Min	Max	Med	Arith	cOWA	H
Img 0	1 26,68848	26,60371	30,66996	30,89667	30,73823	30,75448	21,37117	20,83960	26,73708	27,07854	27,01270	27,07067
Img 0	2 33,50403	33,46846	37,51525	37,64240	37,57713	37,58138	19,70858	19,54290	23,92198	24,07786	24,05762	24,07478
Img 0	3 26,80034	26,74460	30,47904	30,55504	30,52128	30,51564	20,46198	20,82576	25,64113	26,16092	26,08186	26,14607
Img 0	4 28,90415	28,83284	32,88120	33,01225	32,94828	32,94146	22,59335	22,24354	27,94347	28,26449	28,19574	28,25700
Img 0:	5 25,04896	25,04438	28,75582	28,85475	28,81506	28,79901	18,86628	19,55278	24,12507	24,68962	24,58713	24,67322
Img 0	6 38,10156	38,07248	42,08612	42,13003	42,12316	42,11653	29,48308	29,26559	34,89670	35,11481	35,09436	35,11023
Img 0	7 24,48520	24,38872	28,31229	28,45667	28,35114	28,37668	18,95771	18,72670	24,18918	24,55073	24,48373	24,54269
Img 0	8 23,69576	23,73464	27,41557	27,51579	27,46383	27,45864	17,71071	18,59348	23,11305	23,54332	23,43522	23,53119
Img 0	9 26,19262	26,09448	30,06427	30,22940	30,11893	30,13332	20,97846	20,44416	26,23824	26,53197	26,42064	26,52562
Img 1	0 21,48459	21,41350	25,37475	25,58054	25,43016	25,45073	16,47636	16,22205	21,89755	22,22614	22,10356	22,21825
Avg	27.49057	27.43978	31.35543	31.48735	31.40872	31.41279	20.66077	20.62565	25.87034	26,22384	26.14726	26.21497

Table 1: PSNR values of reconstruction of imagens by nn interpolation.

 ${\rm USING}~2~\times~2~{\rm BLOCKS}~~{\rm USING}~4~\times~4~{\rm BLOCKS}$

	Min	Max	Med	Arith	cOWA	н	Min	Max	Med	Arith	cOWA	н
Img 01	27,25658	27,41249	31,70137	31,66148	31,64818	31,70944	21,84394	21,46624	28,12885	28,03911	28,13262	28,08806
Img 02	29,07393	29,09065	29,98667	30,00618	29,99790	29,99295	20,22210	19,99324	24,09349	24,09114	24,09696	24,10058
Img 03	28,07377	27,53953	31,96271	31,87901	31,87085	31,94673	21,36383	21,65788	27,34577	27,53279	27,57114	27,56163
Img 04	29,70934	29,78913	34,39128	34,28215	34,31414	34,37504	23,23057	22,96007	29,81717	29,65596	29,77096	29,71475
Img 05	26,30684	25,74955	30,17965	30,08193	30,05530	30,16533	19,54307	20,06159	25,32192	25,47922	25,51400	25,51442
Img 06	40,09734	39,94107	48,99047	48,55730	48,52986	48,86710	30,92215	30,60188	42,72668	41,77064	41,99358	41,97442
Img 07	25,10689	25,04408	28,93328	28,92340	28,89276	28,94254	19,43662	19,19604	24,96897	25,00413	25,05911	25,02899
Img 08	24,63619	24,10410	28,19100	28,17758	28,16818	28,19312	18,28578	18,86696	23,87169	24,09781	24,07356	24,10310
Img 09	26,60297	26,71398	30,54028	30,56126	30,52693	30,55733	21,32747	20,91360	27,09762	27,10526	27,16280	27,13073
Img 10	21,93973	21,90280	25,71329	25,74295	25,69402	25,73353	16,77848	16,57833	22,58040	22,61488	22,63949	22,63987
Avg	27,88036	27,72874	32,05900	31.98732	31.96981	32.04831	21,29540	21,22958	27.59525	27,53909	27,60142	27,58566

Table 2: PSNR values of reconstruction of imagens by bilinear interpolation.

USING 2 \times 2 BLOCKS USING 4 \times 4 BLOCKS

	Min	Max	Med	Arith	cOWA	н	Min	Max	Med	Arith	cOWA	Н
Img 01	27,39667	27,45993	32,53367	32,62657	32,52946	32,58602	21,83423	21,39364	28,64265	28,74908	28,80893	28,78768
Img 02	30,06149	30,00816	31,28820	31,31873	31,30611	31,29877	20,20038	19,88701	24,49596	24,56989	24,56761	24,57359
Img 03	28,09952	27,62931	32,92967	32,90897	32,87767	32,93859	21,25132	21,55589	27,82091	28,31402	28,28961	28,32229
Img 04	29,92114	29,94430	35,70586	35,70361	35,68906	35,73313	23,22310	22,89860	30,47704	30,54773	30,60332	30,59348
Img 05	26,38597	25,93655	31,32017	31,30790	31,25508	31,33640	19,45423	20,06391	25,74518	26,18606	26,15139	26,20092
Img 06	40,05229	40,02173	51,35284	51,07478	51,01447	51,31081	30,81953	30,48357	44,31891	43,83439	44,03526	44,05492
Img 07	25,23188	25,16984	29,85564	29,93609	29,85733	29,89915	19,36949	19,11221	25,29211	25,49221	25,49999	25,50641
Img 08	24,72669	24,32047	29,10402	29,15066	29,11737	29,12822	18,21007	18,91559	24,17857	24,57330	24,49174	24,56575
Img 09	26,73252	26,79140	31,27454	31,38274	31,29368	31,32452	21,32252	20,85345	27,41366	27,56839	27,55860	27,58354
Img 10	22,04218	21,98136	26,39147	26,52171	26,41585	26,44659	16,76501	16,53815	22,82004	23,00025	22,96201	23,01459
Avg	28,06504	27,92630	33,17561	33,19318	33,13561	33,20022	21,24499	21,17020	28,12050	28,28353	28,29685	28,32032

Table 3: PSNR values of reconstruction of imagens by bicubic interpolation.