

The Probability Distribution Function of Gas Surface Density in M33

Edvige Corbelli¹, Bruce G. Elmegreen², Jonathan Braine³, and David Thilker⁴

¹ INAF-Osservatorio Astrofisico di Arcetri, Largo E. Fermi, 5, 50125 Firenze, Italy

e-mail: edvige@arcetri.astro.it

² IBM Research Division, T.J. Watson Research Center, 1101 Kitchawan Road, Yorktown Hts., NY 10598

e-mail: bge@us.ibm.com

³ Laboratoire d'Astrophysique de Bordeaux, Univ. Bordeaux, CNRS, B18N, allée Geoffroy Saint-Hilaire, 33615 Pessac, France

e-mail: jonathan.braine@u-bordeaux.fr

⁴ Center for Astrophysical Sciences, The Johns Hopkins University, 3400 N. Charles Street, Baltimore, MD 21218, USA

e-mail: dthilker@pha.jhu.edu

Received; accepted

ABSTRACT

Aims. We examine the interstellar medium (ISM) of M33 to unveil fingerprints of self-gravitating gas clouds across the star-forming disk.

Methods. The probability distribution functions (PDFs) for atomic, molecular, and total gas surface densities are determined at a resolution of about 50 pc from available HI and CO emission line data over regions that share coherent morphological properties and considering cloud samples at different evolutionary stages along the star formation cycle.

Results. Most of the total gas PDFs from the central region to the edge of the star-forming disk are well-fitted by log-normal functions whose width decreases radially outwards. Because the HI velocity dispersion is approximately constant across the disk, the decrease of the PDF width is consistent with a lower Mach number for the turbulent ISM at large galactocentric radii where a higher fraction of HI is in the warm phase. The atomic gas is found mostly at face-on column densities below $N_H^{lim} = 2.5 \cdot 10^{21} \text{ cm}^{-2}$, with small radial variations of N_H^{lim} . The molecular gas PDFs do not show strong deviations from log-normal functions in the central region where molecular fractions are high. Here the high pressure and rate of star formation shapes the PDF as a log-normal function dispersing self-gravitating complexes with intense feedback at all column densities that are spatially resolved. Power law PDFs for the molecules are found near and above N_H^{lim} , in the well defined southern spiral arm and in a continuous dense filament extending at larger galactocentric radii. In the filament nearly half of the molecular gas departs from a log-normal PDF and power laws are also observed in pre-star forming molecular complexes. The slope of the power law is between -1 and -2 . This slope, combined with maps showing where the different parts of the power law PDFs come from, suggest a power-law stratification of density within molecular cloud complexes, which is consistent with the dominance of self-gravity.

Key words. ISM: clouds – Galaxies: individual (M33) – Galaxies: ISM –

1. Introduction

The formation and evolution of self-gravitating structures in the interstellar medium (hereafter ISM) is a key ingredient of star formation. Stars form in the dense parts of interstellar clouds where self-gravity overwhelms local gas pressure, causing collapse (McKee & Ostriker 2007). The connections between these dense parts and their surrounding gas can reveal the processes involved. If the cloud is part of a shell (Palmeirim, et al. 2017), converging flow (Whitworth et al. 2018) or spiral arm shock front (Elmegreen et al. 2014), then the clouds are likely to have been formed or influenced by the associated pressures, after which gravity could have made the central parts dense enough for star formation. If disk gravitational instabilities generate spiral arms and filaments that fragment into molecular clouds, then even the birth of these complexes might be driven by gravity out to the edges of their HI envelopes.

The column density and density where environmental processes lose their influence and self-gravity takes over are indicators of the driving pressures and timescales for cloud evolution. The column density at the threshold for strong gravity in a medium of pressure P is $\Sigma_{\text{thres}} \sim (2P/\pi G)^{0.5}$, and the dynam-

ical time for evolution is proportional to the idealized free fall time at density ρ , which is $t_{\text{ff}} = (32G\rho/3\pi)^{-0.5}$. In interacting galaxies where pressure is high, the column density and density at the transition should both be high, making star-forming clouds opaque, fully molecular and quickly evolving. Conversely, in the far-outer parts of spiral galaxies and in dwarf irregulars where the pressure is low, the transition should be at a low density, making the timescales long and possibly placing the first stages of collapse in the atomic medium (e.g., Michalowski et al. 2015; Meidt 2016; Elmegreen 2018).

The best way to determine the transition point from environmental to internal cloud control is by mapping the dynamical state of the cloud and its surroundings (i.e. which region is collapsing, accreting, or in virial equilibrium, and the type of support against gravity). Our galaxy offers the possibility to study individual star forming regions, identify dense clumps, filaments, shocks, and map their column densities and velocity fields. This has been done in several local regions (e.g., Deharveng et al. 2015; Figueira et al. 2017) but it is difficult to do in external galaxies (Hirota et al. 2011; Donovan Meyer et al. 2013; Colombo et al. 2014; Tosaki et al. 2017; Baba et al.

2017; Faesi et al. 2018) because of a lack of spatial resolution needed to identify high column density clumps.

An alternative to cloud-scale mapping is the use of column density probability distribution functions (PDFs) from maps of gas emission or absorption. The PDF for gas in a large-scale map gives the probability distribution of column density averaged over all of the clouds and intercloud regions. This does not allow the history of individual star-forming clouds to be determined, but it can give the average column density where cloud self-gravity begins to dominate over environmental pressure as a driving force. This is possible because self-gravity gives a cloud an internal density structure that is a power law with radius, such as $\rho(r) \propto r^{-1.5}$ or r^{-2} (e.g. Shu 1977), as observed, e.g., by Mueller et al. (2002). Then the PDF has a matching power law at high density even if the cloud has locally a log-normal PDF from compressible turbulence (Elmegreen 2011). Small scale collapse inside a turbulent cloud also produces a power-law tail which extends to lower densities as time proceeds, while turbulent motions broaden the PDF (Girichidis et al. 2014). Computer simulations showing this power law extension are in Klessen (2000), Vázquez-Semadeni et al (2008), Kritsuk et al. (2011), Federrath & Klessen (2013), Pan et al. (2016) and elsewhere. Log-normal distributions without self-gravity were first discussed and simulated by Vázquez-Semadeni (1994), Passot & Vázquez-Semadeni (1998), and Padoan, Nordlund & Jones (1997). The lower limit to the power law part of the PDF corresponds approximately to the column density at the edge of the self-gravitating part of the cloud, and therefore gives the pressure at that place (e.g., Kainulainen et al. 2011). Sensitivity limits, however, are important and should be taken into account since, for example, the total PDF may not curve over at low column density if the detection limits for gas are close to the apparent PDF peak (Alves et al. 2017).

PDFs have been extensively studied and observed in the local interstellar medium using also extinction or dust emission. In the Milky Way the resolution is sufficiently high to trace the molecular clumps that form individual stars (Kainulainen et al. 2009, 2011; Froebrich & Rowles 2010; Lombardi, Lada & Alves 2010; Schneider et al. 2012, 2015b). Lombardi, Lada & Alves (2010) found a log-normal PDF using extinction maps in a large region around Perseus and Taurus in the solar neighborhood, with a slight excess over the log-normal above $A_V \sim 2$ mag. Power law PDF extensions were also found in dust emission (Schneider et al. 2013, 2015a,c; Lombardi et al. 2015) and molecular line emission (Schneider et al. 2016) up to $A_V \sim 100$ mag (Schneider et al. 2015b). Schneider et al. (2012) measured the PDFs of dust emission in the Rosette molecular cloud and found it became a power law above $A_V = 9$ mag in the central regions and above 4 mag in other regions; it was a log-normal far from the Rosette nebula. It is still not clear, however, whether non star-forming clouds have power-law extensions. Although in most external galaxies we cannot yet resolve the cores of individual star-forming clumps, we may still trace where gravity takes over ISM pressure on a larger scale, and it can be useful to compare the PDFs in various environments, such as galaxy centers, spiral arms and interarms, and as a function of galactocentric distance and cloud type.

M33 is a good candidate to examine for these questions because it is the nearest isolated spiral galaxy. Recent surveys of the ISM throughout the whole star-forming disk of M33 have mapped the atomic and molecular gas phases with high resolution (Gratier et al. 2012; Druard et al. 2014; Corbelli et al. 2014). The H α and GALEX ultraviolet data complement the Spitzer mid-infrared survey to trace star formation from the em-

bedded to the exposed Young Stellar Clusters (Verley et al. 2009; Sharma et al. 2011). The possible association between 566 Giant Molecular Clouds (hereafter GMCs) and 600 Young Stellar Cluster Candidates (hereafter YSCCs) has helped in defining the duration of the lifecycle of GMCs in M33 (Corbelli et al. 2017). Still, it is unclear what drives the transition from inactive gas to a star-forming cloud. Also, if stellar feedback disrupts these clouds, it is of interest to see if this disruption affects only the small scale structure of the ISM or if it also contributes to the global disk morphology (Wada et al. 2000). The presence of turbulence in the ISM of M33 has been investigated via power spectra using optical or far-infrared images (Elmegreen et al. 2003; Combes et al. 2012). A recent numerical simulation of the large scale structure of the M33 disk (Dobbs et al. 2018) points out the role of both stars and gas in triggering the formation of high-density filaments by gravitational instabilities. At the same time, the presence of very low density cavities in the interarm regions underlines the need for a high level of feedback (e.g., $\sim 10\%$ of the total energy of massive stars) to deposit turbulent energy. The purpose of this paper is to understand the role of turbulence and gravity in triggering the formation of GMCs and star formation throughout the disk of M33. Due to the strong evidence of non-negligible turbulent motion in the M33 disk we shall interpret the log-normal PDFs that we detect in the ISM of this galaxy as due to turbulence although there might be other features associated to clouds that can produce log-normal column density distributions (Tassis et al. 2010).

The analysis of the PDF for the molecular gas in M33 by Druard et al. (2014) has shown deviations from a log-normal shape for gas column density above $\sim 1.7 \times 10^{21} \text{ cm}^{-2}$. The authors attributed the deviations of a galaxy-wide PDF to self-gravity, but do not draw any definitive conclusion on the cloud scale affected by gravitational contraction. The coarse spatial resolution of CO observations in external galaxies in fact does not allow analysis of the details of individual clouds and hence to distinguish a free-fall contraction of the whole GMC complex from self-gravity of clumps and cores inside them. The publication of a complete GMC catalogue in the star forming disk of M33 and of further analysis of the GMC properties, that followed the complete census of molecular gas presented by Druard et al. (2014), open the possibility to investigate the role of gravity in different types of GMCs which are likely GMCs at different evolutionary stages. Furthermore, making use of high angular resolution HI data products presented by Corbelli et al. (2014) we can examine the PDFs of atomic and total gas. For the molecular component we extend the analysis of Druard et al. (2014) to specific disk areas or cloud types where self-gravity or turbulence plays a major role, and determine if the PDFs depend on star formation rate or galactocentric distance and what this implies on the resolved scale of ~ 50 pc. This is particularly relevant for M33 given that both disk gravitational instabilities and high levels of stellar feedbacks play a role in shaping the atomic gas distribution in the disk (Dobbs et al. 2018).

In what follows, the data and cloud types are summarized in Section 2, the PDFs and power law slopes are presented in Section 3. Results are discussed in Section 4 and summarized in Section 5.

2. The data

In analyzing the data of M33, we assume a distance of 840 kpc (Freedman et al. 1991; Gieren et al. 2013) which implies a linear scale of 4.1 pc per arcsec and 0.24 kpc per arcmin. As position angle and inclination of the gaseous disk we use PA=20° and

$i=54^\circ$ based on the averages for a tilted ring model fitted to the 21-cm velocity field at $R < 7$ kpc (Corbelli et al. 2014). The PDF is analyzed for face-on values of H-atoms column densities.

2.1. The neutral atomic gas traced by HI 21-cm line emission

To analyze the distribution of the neutral atomic gas, we use the high resolution Very Large Array (VLA) and the Green Bank Telescope (GBT) observations of the 21-cm HI line emission described by Corbelli et al. (2014). We generally choose an angular resolution that is less than the limit in the VLA+GBT survey in order to trace more sensitively the faint diffuse emission, and a spectral resolution of 1.29 km s^{-1} . The moment task in MIRIAD has been used to generate the moment maps. The moment-0 is the integrated intensity map of the 21-cm line, the moment-1, or mean velocity, is the map of the weighted mean velocity along the line of sight, and the moment-2, or velocity dispersion, is the map of the intensity-weighted mean deviation of the velocity along the line of sight. For each pixel i the values of integrated intensity, I_i , mean velocity, V_i , and velocity dispersion, $\sigma_{21,i}$, are computed using $S_{i,j}$, the 21-cm line intensity in the velocity channel centered on v_j , as follows:

$$I_i = \sum_j S_{i,j} \quad V_i = \frac{\sum_j S_{i,j} \times v_j}{\sum_j S_{i,j}} \quad (1)$$

$$\sigma_{21,i} = \sqrt{\frac{\sum_j S_{i,j} \times (v_j - V_i)^2}{\sum_j S_{i,j}}} \quad (2)$$

where the sum extends over all spectral channels with detectable emission. The images we use in this paper mostly have a 10 arcsec (41 pc) restored beam size and are masked to mitigate contributions from foreground Milky Way emission. The pixel size is 4 arcsec, smaller than the beam size. To check the high column density PDF we also use images at 5 arcsec spatial resolution with pixel size of 2 arcsec. The radial trend of the HI line width is inferred from the moment-2 map at a spatial resolution of 10 arcsec.

Analysis of the moment-0 map at 10 arcsec resolution indicates a noise standard deviation $\sigma_{HI} = 0.0049 \text{ Jy/beam km s}^{-1} = 5.4 \times 10^{19} \text{ H cm}^{-2}$ on a pixel by pixel basis. We consider only pixels with a flux above $3\sigma_{HI}$, and trace the statistical analysis down to this limiting flux value, which corresponds to a face-on column density of about $10^{20} \text{ H cm}^{-2}$.

The line of sight velocity dispersion σ_{21} , defined above, is assumed to be isotropic. The map of the velocity dispersion of the M33 disk at 21-cm, can be an indicator of the turbulence and kinetic energy of the HI gas, especially for nearby galaxies which can be imaged at high resolution to avoid line broadening due to disk dynamics. We show the moment-2 map in the left panel of Figure 1 using the black contours to outline the high density HI filaments or arms, as shown by the 21-cm moment-0 map in the right panel of the same figure. In the dispersion map we masked out all pixels corresponding to the two regions with large infall/outflow of gas: the stream towards the central region just south of the nucleus, and a smaller region above the northern arm. In these areas the signal is a superposition of emission lines

at two velocities, one corresponding to gas in the disk and the other being due to extraplanar gas.

2.2. The molecular gas and the GMCs population

We use molecular data of the deep $^{12}\text{CO}(J=2-1)$ whole-disk survey carried out with the IRAM-30m telescope (Druard et al. 2014) at $10.7''$ resolution (44 pc). From the spectral cube of this database at a spatial resolution of $12''$ and at a spectral resolution of 2.6 km s^{-1} , a catalogue of 566 giant molecular clouds (GMCs) has been presented by Corbelli et al. (2017), following a procedure described and used for M33 by Gratier et al. (2012). The main beam temperature noise for the dataset at $12''$ resolution is 0.033 K in each channel. The moment-0, or brightness map that we use has 3 arcsec pixels and on average $\sigma_{CO} = 0.27 \text{ K km s}^{-1}$, considering a typical spectral window of 25 km s^{-1} per beam. Using a CO-to- H_2 conversion factor $X = N(\text{H}_2)/I_{CO(1-0)} = 4 \times 10^{20} \text{ cm}^{-2}/(\text{K km s}^{-1})$ (Gratier et al. 2017), twice the galactic value, and an intrinsic line ratio $R_{1-0}^{2-1} = I_{2-1}/I_{1-0} = 0.8$ (Druard et al. 2014), we estimate $\sigma_{\text{H}_2} = 1.3 \times 10^{20} \text{ H}_2 \text{ cm}^{-2}$. This is an average estimate of the map noise because there are variations of the main beam temperature noise and of the signal width across the disk. To trace the PDF we consider only pixels with flux above $3\sigma_{CO}$ and trace the statistical analysis down to the corresponding limiting column density, which in term of total H atoms reads $3\sigma_H \approx 7.8 \times 10^{20} \text{ H cm}^{-2}$ (i.e. $4.6 \times 10^{20} \text{ H cm}^{-2}$ face-on value).

Molecular clouds are classified in three broad categories: clouds without obvious star formation (A), clouds with embedded star formation (B) and clouds with exposed star formation (C) by Corbelli et al. (2017). An on-line catalogue lists several cloud properties such as cloud radius and luminous mass computed by converting the total CO line luminosity of a cloud into mass as given by equation (1) of Corbelli et al. (2017). Type A,B,C clouds have average luminous masses of 1.3, 2.6 and $3.1 \times 10^5 M_\odot$. As noticed by Corbelli et al. (2017) and by Braine et al. (2018), going from non-star-forming to exposed star-forming clouds, the average cloud mass grows. The total H_2 mass in A,B,C-type clouds is, respectively, 0.16 , 0.14 and $0.79 \times 10^8 M_\odot$. Adding the unclassified D-type clouds, we have a total molecular hydrogen mass of $1.12 \times 10^8 M_\odot$ in GMCs or a total GMC gas mass (considering He and heavy elements, i.e., taking a hydrogen fraction of 0.73 %) of $1.53 \times 10^8 M_\odot$. That is about half of the total molecular ISM mass. The rest is in lower-mass molecular clouds (below the survey resolution limit) which we shall call ‘‘diffuse molecular gas’’, even though some of this gas may be self-gravitating. By selecting 630 mid-infrared (hereafter MIR) sources from the list of Sharma et al. (2011) that are YSCCs in the early formation and evolutionary phases, Corbelli et al. (2017) determined that they are strongly correlated with the GMCs (the correlation length is of order 17 pc, smaller than the typical GMC radius). Since age estimates of the YSCCs are between 3.5 and 8 Myrs, given the fractions of GMCs in each class (i.e., presumably in each evolutionary phase) the GMC lifetime was estimated to be 14.2 Myrs, with the longest phase being the C-type phase, when the YSCCs break out of their clouds before total gas dispersal.

To sample the surface density distribution for each cloud, we can either use the corresponding pixels identified by the cloud extraction algorithm, or use the cloud radius r_c to find the associated pixels. Although most of GMCs are rather spherical, there a few of them which have an elongated shape and look filamentary. We have carried out the analysis using both selec-




Fig. 1. Velocity dispersion (moment-2) map of the 21-cm line emission in M33 at 10 arcsec resolution (*left panel*). The black contours underline the location of the bright HI filaments or arms, they are drawn at $0.2 \text{ Jy beam}^{-1} \text{ km s}^{-1}$ and overlaid also on the moment-0 map in the *right panel*. The two areas that have been masked out for the likely presence of gas infall/outflow (the stream towards the central region in the southern half and a smaller region above the northern arm) have anomalously large velocity dispersions and have yellow-red colours in the velocity dispersion map. The colour legend for the velocity dispersion map (*left panel*) is in units of km s^{-1} , for the intensity map (*right panel*) is in $\text{Jy beam}^{-1} \text{ km s}^{-1}$. Coordinates of the map centers are written in the bottom part of the Figure.

tion criteria at first but found no sensible differences in the resulting PDFs. We therefore decided to use the cloud radii listed by Corbelli et al. (2017) to select pixels corresponding to each GMC. This simplifies the procedure to find the atomic gas at and around cloud positions. The average GMC radius decreases from 26 to 22 to 18 pc from A- to B- to C-type clouds, even though the mass increases in this sequence. This seem to suggest that GMCs are more compact as they evolve, likely collapsing and shrinking as they collect more material and begin to form stars. It is not clear however if GMCs of type-C are more bound and dominated by gravity since the presence of young stars in these molecular complexes might increase the local turbulence. It is therefore of interest to investigate the PDFs of GMC at each evolutionary stage.

2.3. Selected regions and total gas maps

Using the cold gas images we have selected four regions which might be of interest for cloud formation and destruction and have enough pixels with CO emission above the thresholds to ensure a statistically significant analysis. These are: the central region (CR) and three other regions that have prominent HI overdensities and non-negligible H_2 fractions, namely the northern arm

(NA), the southern arm (SA) and the eastern filaments (EF). The latter feature is the HI overdensity that starts at NGC604 (the brightest HII region in M33 located at 9.1 arcmin to the north and 7.5 arcmin to the east in Figure 1) and extend radially outwards (as an extension of the northern arm towards the south). Figure 2 shows these regions on the 21-cm emission map of M33. We mark all pixels with CO J=2-1 integrated emission above $3\sigma_{CO}$ using one colour for each selected region and using the green colour for the interarm regions (IA). We shall refer to interarm region when we analyze everything that is not in one of the selected areas despite there are filament segments which are included in it. We do not have the sensitivity to select only the truly interarm ISM.

By analyzing individual regions or radial bins separately, we minimize mixing data relative to gas with different scale heights and disk gravity. The southern arm looks less disturbed than the northern arm and it is the only arm that fits well with density wave models, as underlined by Humphreys & Sandage (1980). This justifies a separate analysis of the PDFs relative to each arm. Results for the eastern filament have been checked by considering also the same region without the area in and around NGC604. This has been done in order to make sure that it is not




Fig. 2. Map of regions selected for this study. The grey background image is the 21-cm map of M33 and the 6 yellow ellipses indicate the boundary of the radial bins: 1,2,3,4,5,7 kpc.. The colored dots indicate all pixels where CO J=2-1 integrated emission is above $3\sigma_{CO}$ and are color coded according to the different selected regions described in the text: cyan for central region (CR), blue for northern arm (NA), magenta for southern arm (SA), red for eastern filament (EF), green for interarm (IA).

only the presence of the large star forming region driving the detected properties.

When we analyze GMCs in selected regions we require that the center of the GMCs to be within the boundaries of the region (not the whole GMC or its HI envelope). To collect pixels with CO or HI emission above the threshold associated with the n -cloud, we require that the distance $d_{i,n}^{CO}$ and $d_{i,n}^{HI}$ between the i -pixel center and the n -cloud center satisfies

$$d_{i,n}^{CO} \leq 1.7 \times r_c^n \quad d_{i,n}^{HI} \leq 2.5 \times r_c^n. \quad (3)$$

For gaussian shaped radial distribution of cloud brightness we recover 90% of the total cloud molecular mass with this selection. For the HI envelope the factor 2.5 encompasses pixels next to the GMC without appreciable CO emission.

The total hydrogen gas map is the sum of the molecular and atomic gas surface densities pixel by pixel. The CO J=2-1 map was aligned with the HI 21-cm map and the HI and CO line brightnesses converted to H atom column densities. Before adding the molecular gas surface density to the atomic one, we masked out all 4 arcsec wide pixels with H_2 column density below $2\sigma_{H_2}$ (We mask pixels below 2 and not $3\sigma_{H_2}$ because the

aligned CO map with 4 arcsec pixels has a lower noise than the original σ_{H_2} value. Given the subsolar metallicities of M33 it is likely that there is not much molecular gas at lower surface densities because of radiative dissociation. Because the HI image has a better sensitivity than the CO image in tracing the hydrogen column density, the lower limit for sampling the total gas column density will be the same as that of the HI gas.

3. The global and local gas PDFs across the M33 disk

PDFs of inclination-corrected gas surface densities were made by summing pixels in logarithmic bins for HI, H_2 , or total gas using units of H atoms cm^{-2} . One set of PDFs was made for different radial intervals and another set of PDFs was made by summing pixels in four selected regions. In this last case, the PDFs corresponding to three types of GMCs and the diffuse CO are also shown (diffuse CO is where CO emission is associated with unresolved low mass clouds or is truly diffuse). The three GMC types are: GMC-A (containing no trace of star formation), GMC-B (with only embedded star formation), and GMC-C (with exposed HII regions).

In order to localize deviations of the PDF from a log-normal distribution we investigate the local slopes defined as the derivative of the PDF in every bin of column density (logarithmically spaced). Local slopes are computed fitting a line segment through two adjacent bins on each side. A log-normal PDF will have a linearly decreasing slope while a power law PDF will have a constant slope. Fitted log-normal PDFs to the data are also shown to complement the analysis for the most relevant cases. As discussed in the next Section and mentioned in Section 1, compressible turbulence shapes the PDF as a log-normal while self-gravity gives a cloud an internal density structure and a power law PDF for the gas surface density.

3.1. The PDF across arms, filaments, clouds, and the galaxy center

Figure 3 shows the molecular gas PDFs: in the bottom panels for gas in the different GMCs types and diffuse (color coded using different colours), and with thick lines for the total molecular gas (regardless of GMC type or diffuse). The columns correspond to six different regions in the galaxy, as indicated at the top. The full disk is on the left and includes all pixels with CO brightness above $3\sigma_{CO}$, followed by the central region, the northern and southern spiral arms, and the eastern filament. The combination of molecular gas which is not in CR, NA, SA, EF is labeled “interarm” (IA) and it is shown on the right. In the top 5 panels, Figure 3 shows the local slopes of these PDFs: S_{CO} (black dots) are the PDF slopes for total molecular emission, S_{GMC-A} , S_{GMC-B} , S_{GMC-C} , S_{diff} , are the PDF slopes for molecular gas in GMC-type A,B,C and diffuse. The log-normal parts of the PDFs in the bottom panels have linearly decreasing slopes while the power law parts of the PDFs have constant slopes and in this case the point distributions in the slope panels flatten. The panels show linearly decreasing trends in the center and in the northern arm while for the eastern filament constant slopes are present. For the total CO the PDFs have power law parts in the southern arm, eastern filament and interarm regions. The southern arm and filament show also power law parts for individual GMC-types. Elsewhere, the PDFs are approximately log-normal, or too irregular to detect a deviation from log-normal.




Fig. 3. (*Bottom panels*) The PDFs of molecular gas column density in units H-atoms cm^{-2} for the total molecular gas in the disk (black heavy curves), and for the gas in three GMC types: A (non-star forming), B (embedded star formation), and C (exposed star formation) in blue, red, green, respectively in the on-line version. The PDF for diffuse molecular gas (meaning low mass clouds or truly diffuse) is marked in magenta in the on-line version. All PDFs have been normalized to the number of pixels at the peak of the total molecular gas distribution ($n_{\text{max}}^{\text{CO}}$) and refer to face-on values. (*Top panels*) The local slopes S in the corresponding PDFs for the total molecular gas, S_{CO} , for the surface density of type A-,B-,C-GMCs, and for the diffuse molecular gas. The columns correspond to different regions in the galaxy: the full disk is on the left, followed by the central region, the northern and southern spiral arms, and the eastern filament. The combination of molecular gas from the rest of the galaxy is labeled “interarm” and it is shown on the right.

For the total molecular gas in the whole disk (black colors, left panel), the power law part of the PDF starts at a column density of $\sim 10^{21.1} \text{ cm}^{-2}$, which corresponds to about 0.7 magnitudes of visual extinction; $A_V = 0.7 \text{ mag}$ is obtained using the conversion factor in Bohlin et al. (1978) but it can have slightly lower values given the metallicities of M33 (Magrini et al. 2010). We would like to remark that the whole disk PDF shown in Figure 3 is the same as that presented by Druard et al. (2014),

being drawn from the same data. However, we use the slope to infer the column density interval in which the PDF deviates from a log-normal distribution, instead of fitting a log-normal function, given the limited range of column density sampled. The ascending side of the PDF, at low column densities, is not traced because of the sensitivity limits of the survey and of molecular gas dissociation thresholds. This leaves large uncertainties in the parameters of the best fitting log-normal functions, which

in the next subsection are only shown for a qualitative comparison with the data. The analysis of the slopes shows a power law behaviour which starts at slightly lower column densities than those found by Druard et al. (2014) (1.3×10^{21} H atoms cm^{-2} versus 3.4×10^{21} H atoms cm^{-2}). The spatial resolution of the survey limits the analysis to $N_H < 10^{22}$ H atoms cm^{-2} since these high column density regions tend to be small in physical size and cover only a small portion of the beam. This explains the sharp drop in the molecular gas PDF at high column density.

In Figure 4 we show the atomic gas PDF obtained from HI in the proximity of GMCs i.e. 21-cm column densities at a distance less than 2.5 cloud radii with respect to the cloud centers. It is evident that the PDF drops at high column density, above $N_H^{\text{lim}} = 2.5 \times 10^{21} \text{ cm}^{-2}$. In the central region the drop of the HI PDF in cloud envelopes happens at slightly lower column densities. To interpret N_H^{lim} we recall that the molecular fraction as a function of the total gas column density has been investigated in several galaxies and also in M33 (Figure 13 of Gardan et al. (2007)). The gas mass fraction in molecular form increases as the gas surface density increases, with lower fractions found for gas at large galactocentric distances. However, the trend of molecular gas mass fractions with atomic gas surface densities has not been explicitly investigated in M33. With the sensitivity and resolution of the current surveys we find average molecular mass fractions of order 0.5 as the HI column density approaches N_H^{lim} at galactocentric distances less than 4 kpc; lower average fractions, of order 0.3, are found in the outermost regions. Hence $N_{\text{HI}} \approx 2-4 N_{\text{H}_2}$ when the HI column density is of order N_H^{lim} , and we interpret N_H^{lim} as the maximum sustainable atomic column density in the M33 disk. Molecules can form where the atomic gas column density is lower than N_H^{lim} , especially in the inner disk where gas volume densities and dust abundances are high. However, molecules always form where N_{HI} is close to N_H^{lim} or the total gas column density is higher than $2 N_H^{\text{lim}}$ throughout the whole star forming disk of M33 (or $A_v > 2.4$ if we consider metallicities lower than about 0.4 dex than solar in M33).

The HI PDF is approximately a power law at low column density. This is particularly evident in the interarm PDF and in the total HI PDF (labelled All in Figure 4) which both extend towards lower column densities. Figure 4 clearly shows deviations from a log-normal distribution in the HI PDF of All, EF and IA PDFs. Slopes are constant for $N_H < 5 \times 10^{20} \text{ cm}^{-2}$. What causes the deviation from a log-normal distribution at low column density? Numerical simulations of density fluctuations in compressible polytropic turbulence by Passot & Vázquez-Semadeni (1998) suggest that at low density the PDF can approach a power law if the polytropic index $\gamma > 1$. Values of $\gamma > 1$ imply an equation of state where the Mach number increases with decreasing density. Then the PDF gets broader than the isothermal case at low density, producing the power law instead of a log-normal. This is a picture that may apply to the interarm and other low-density regions if the voids produced by feedback or divergent flows in a spiral arm have higher Mach numbers than the clouds. We can exclude that the power law PDF observed for the HI at low column density is due to noise or incomplete sampling in interferometric observations. Ossenkopf-Okada et al. (2016) have in fact shown that the impact of the observational noise needs to be at least 10% of the PDF peak column density to affect the shape of the PDF at column densities 5-10 times lower than the peak. But the noise level in the HI map is about 5% of the PDF peak column density and deviations from a log-normal are detected starting from column densities only 2-3 times lower than the peak. Moreover, our use of sin-




Fig. 4. The PDF (left panels) and the local PDF slope (right panels) of the HI column density in and around the population of GMCs in the whole disk of M33 (labelled All, bottom panel), in the four selected regions (intermediate panels) and in the interarm (top panel), labelled as in Section 2.3 and Figure 2 caption.

gle disk data to complement VLA interferometric observations and achieve a complete uv -plane coverage should avoid possible PDF distortions. However, before drawing any definite conclusion on what drives the low column density PDF, one should consider possible resolution limits which can be relevant if, for example, the low density gas were made of unresolved low mass clouds.

Figure 5 plots the PDFs for HI (blue), H_2 (red) and total H (black) in the left two columns, and the slopes with the corresponding colors in the right two columns. The six rows are different bins of galactocentric distance, which are, from bottom to top, 0-1, 1-2, 2-3, 3-4, 4-5, and 5-7 kpc. The 4th bin from the bottom shows the clearest power law structure for the molecules, with index -2.2, starting at a column density of $\sim 10^{21} \text{ cm}^{-2}$. In this radial bin the power law is even present for the total gas, starting at $\sim 10^{21.4} \text{ cm}^{-2}$, and the PDF is highly symmetric between the high and low density side. The large stellar cluster NGC604 and the gas in its proximity lie in this radial interval. Approximate power laws are also present in the two outermost radial bins in the molecular PDFs. Here some caution should be used in interpreting the sharper PDF fall-off of the molecular gas which can be caused by dark CO i.e. by a population of compact molecular clouds, undetected by the CO survey, or by radial variations of the CO-to- H_2 conversion factor. A higher CO-to- H_2 conversion factor at large galactocentric radii would increase the PDF at high column densities and give a more symmetric total gas PDF as well.

The atomic gas PDFs in separate radial bins, as displayed in Figure 5, show similar drops at high column densities as the PDFs of HI cloud envelopes in Figure 4. On a linear scale the HI column density is distributed smoothly around a peak value

at about $0.5 N_H^{lim}$ with only a few pixels extending above N_H^{lim} . The galaxy has no HI gas with face-on column densities above $4 \times 10^{21} \text{ cm}^{-2}$. We have used the 21-cm map at a higher spatial resolution (half of what has been normally used in this paper) to check this finding, but no significant variations have been detected. We cannot exclude that there are localized opacity corrections to the HI column density above N_H^{lim} (Braun et al. 2009; Braun 2012). As in Figure 4, the HI PDFs drop more slowly than a log-normal at low column densities and they are close to a power law. The power law slopes are between ~ 2 and ~ 4 outward of 2 kpc, increasing towards larger galactocentric radii.

3.2. PDF widths and the 21-cm line velocity dispersions

The widths of total gas PDFs get narrower going from 1 kpc to large galactocentric radii. If the possible presence of dark CO keeps uncertain the widths of the molecular PDFs at large radii, this radial trend can be inferred also from the low column density side of the PDF, dominated by the atomic gas. However, there is no radial trend for the HI velocity dispersion, as shown by the 21-cm moment-2 map in Figure 1. In Figure 6 the filled squares indicate radial averages of the 21-cm velocity dispersion, σ_{21} , with their standard deviations. The flatness of the distribution is remarkable and the resolution of the 21-cm survey for this galaxy is excellent to avoid line broadening due to large scale motion in the disk, such as a rising rotation curve in the center or spiral arm dynamical perturbations. For several galaxies the observed 21-cm line velocity dispersion decreases towards large galactocentric radii (Tamburro et al. 2009). We would like to point out that if we lower the spatial resolution of the HI survey by a factor 10 we also detect a decrease of the velocity dispersion from a central value of 21 km s^{-1} to about 14 km s^{-1} at the edge of the star forming disk. The resolution and sensitivity of the 21-cm dataset used in this paper, together with the regular morphology and low inclination of this nearby spiral galaxy, seems appropriate for using the 21-cm line broadening as tracer of turbulent motion. There are some variations in the HI dispersion across the disk, and this can be seen in Figure 1.

In Figure 6 we use the open symbols to plot radial averages of the kinetic energy per unit area estimated as $E_k = 1.5\sigma_{21}^2 \Sigma_{HI}$ in units of $10^{45} \text{ ergs/pc}^2$. Radial averages of the velocity dispersion and of the HI surface densities are about constant across the star forming disk (Corbelli et al. 2014), but since their small variations are correlated in the inner disk, variations in the kinetic energy per unit area are more evident, even though they are less than a factor 2. The energy density E_k peaks around 2 kpc due to the presence of the southern and northern arms where the kinetic energy density in the atomic gas is about $5 \times 10^{46} \text{ ergs pc}^{-2}$. The energy density decreases beyond 3 kpc and drops at the edge of the star forming disk at about 7 kpc because of the low gas surface density there. Between the four selected regions, the interarm and the eastern filament regions have the lowest kinetic energy densities for the atomic phase. The eastern filament is an HI overdensity but the HI velocity dispersion is lower by about 2 km s^{-1} than in the arms.

Values of the velocity dispersion are well above $6\text{--}8 \text{ km s}^{-1}$, which is the expected thermal width for the warm phase of the ISM, hence turbulent velocities dominate across the whole disk. The decrease of PDF width for total and atomic hydrogen at large galactocentric radii could then be due to a decrease in the Mach number (which is of order unity). Because the velocity dispersion is constant, this would imply that the average thermal temperature of the gas increases with radius, possibly as a result

of a transition to a higher fraction of HI in the warm phase as the stellar surface density, and hence the disk pressure, decreases.

In Figure 7 we show the best fitting log-normal functions to the molecular gas PDFs in the selected regions. It is clear that log-normal functions fit better the data in the center and northern arm region than in the southern arm and eastern filament region. The peak and the width of log-normal functions have been considered free parameters in the intervals $[-0.4, 0.4]$ and $[0, 1]$ respectively (uncertainties have been estimated using Poisson statistics). Even though the significance of the deviations are larger for the all-disk and interarm distribution, we show the four regions here to underline some differences between their PDFs. For example in the regions where the PDFs are truly log-normal, namely the center and northern arm, the GMCs of type C show clearly a wider log-normal distribution than GMCs of types A and B, likely because the newly born massive stars in type C GMCs increase the turbulent motion.

3.3. Interarm and gravity dominated filaments

In this Section we investigate some details of the power law parts of the molecular gas PDFs across the disk that has been shown in Figure 3. The all-disk, the southern arm, eastern filament and interarm all show a molecular gas PDF which deviates from a log normal at $N_H > 10^{21} \text{ cm}^{-2}$ and it closely resembles a power law. To prove that the power law behavior of the molecular gas PDF along the eastern filaments is a property of the gas along the filament and not a result of the gas around the bright stellar cluster NGC 604 located at one extreme, we have also carried out the analysis excluding the pixels around NGC 604 from the analysis but the same conclusions apply.

The power law PDF for molecules in the southern arm or in the eastern filament could be the result of self-gravity in GMCs, which gives them a power law internal density profile (Klessen 2000; Vázquez-Semadeni et al 2008; Kritsuk et al. 2011; Girichidis et al. 2014). A similar interpretation has been suggested by Druard et al. (2014) when they detected an excess in the galaxy-wide PDF at high column densities. To better investigate this possibility, in Figure 8 we color the different parts of the molecular power law to see where they arise in the southern arm. We display the HI map of the southern arm in the background and underline in yellow all pixels with molecular densities that have $10^{21} < N_H < 2.5 \times 10^{21} \text{ cm}^{-2}$, and in cyan all pixels that have $2.5 \times 10^{21} < N_H < 4 \times 10^{21} \text{ cm}^{-2}$ and finally in blue where $N_H > 4 \times 10^{21} \text{ cm}^{-2}$. The blue regions tend to be in the cores of the yellow and cyan regions, indicating that the power law PDF results from a power law internal structure inside GMCs. This is consistent with the model where cloud self-gravity makes the power law PDF in these regions. Most of the yellow regions with no cyan or blue pixels in the center are type-A GMCs which are located in the inner part of the arm. These pristine clouds presumably get compressed and accrete mass as they cross the arm, developing a high density core that fragments and form stars changing their status to type-C. We estimate that about 70 % of the molecular gas in the southern arm and eastern filament has column densities above 10^{21} cm^{-2} where it deviates from a log-normal distribution. At the resolution of our survey the slope or power law index can be as low as -1.2 to -1.5 for CO in the eastern filament or southern arm while it is -2 for the overall CO PDF and -2.8 for interarms.

Power law PDFs are also observed for galactic GMCs and they have similar internal density gradients. For example Schneider et al. (2015c) determined power law PDF tails above 4 to 5 magnitudes of visual extinction for four local molecular




Fig. 5. The PDFs (left) and PDF slopes (right) of the atomic (blue), molecular (red) and total (black) hydrogen gas in 6 radial bins increasing from the bottom to top and separated by 1 kpc intervals for all but the top row, where the radial interval is 2 kpc. The label in the upper left corner of the leftmost panels indicate the radial boundaries of each bin in kpc. Each PDF has been normalized to their peak value. The 4th bin from the bottom (radius 3-4 kpc) has a clear power-law structure in the molecules and total gas.

clouds, using dust emission maps, and showed that the power law slopes in the PDF corresponded to the slopes of the density distributions inside the clouds, as expected for self-gravitating clouds. Schneider et al. (2015a) measured PDFs for four infrared dark clouds using dust emission and found them to be power-law throughout, regardless of their internal star formation rates, with a transition to log-normal in the lower column density regions surrounding the dark clouds. Cloud core infall was also observed in CO, demonstrating that the power law structure for these clouds is from self-gravity. Evidently, power-law PDFs are common and galactic studies show that the origin of those power laws can often be traced to power-law internal density structures inside individual clouds. Some of the excess dense gas can be from local compression too, as demonstrated for particular regions near bright nebulae (Schneider et al. 2012).

The power law index increases when we average data from regions where some gas follows a power law and some gas follows a log-normal distribution. However, we would like to point out that a power law PDF for column density might have another

explanation that applies to the interarm region. Here the presence of small unresolved cloudlets implies that the gas that is observed in each resolution element can be described as a random collection of individual clouds which follow a power law mass distribution function, $dP/dM \propto M^{-\kappa}$. For $\kappa \sim 1.5$ (e.g., Solomon et al. 1987), the distribution function of column density is a power law with a slope of about -1.5 on a log-log plot if the average gas density is about constant and cloud surface density and size scale as $M^{1/3}$. Steeper mass function slopes such as $\kappa \sim 1.8$ have been observed for molecular clouds in our Galaxy or in external galaxies (Heithausen et al. 1998; Rosolowsky 2005) and these would be better in agreement with the observed slope of -2.8 for the power law of the interarm PDF if clouds are unresolved. More recently Braine et al. (2018), have shown that in M33 the slope of the cloud mass spectrum steepens from $\kappa = 1.4$ to $\kappa = 1.9$ going radially outward. The mass distribution of star clusters in interarms suggests also a steeper cloud mass spectrum ($\kappa \sim 2$) in these regions (Messa et al. 2018). In the interarm, there appears to be no correspondence between different parts




Fig. 6. The radial average values of the 21-cm velocity dispersion and their standard deviations are shown with filled square symbols with errorbars. The open circles indicate the average energy per square parsecs in the same radial bins; energy values have been normalized to 10^{45} ergs pc^{-2} . The asterisks connected by a dashed line (in red in the on-line version) indicate radial averages of the atomic gas surface density in solar masses per square parsecs.




Fig. 7. The best fitting log-normal functions to the PDF of total CO, of CO in GMCs of type A,B,C and diffuse in the four selected regions of the M33 disk. Each PDF is normalized to its maximum value.




Fig. 8. The image of the southern arm with the HI map in the background and colours underlining the CO core-envelope cloud structure which follow a power law PDF. For the CO emission we plot: in yellow all pixels with molecular surface densities $10^{21} < N_H < 2.5 \times 10^{21} \text{ cm}^{-2}$, in cyan all pixels that have $2.5 \times 10^{21} < N_H < 4 \times 10^{21} \text{ cm}^{-2}$ and finally in blue the core of GMCs with $N_H > 4 \times 10^{21} \text{ cm}^{-2}$.

of the PDF power law and different parts of individual clouds, such as a core-envelope structure. Also, samples of each cloud type in the interarm seem to have log-normal PDF, representing turbulent conditions. Still the PDF for all of the clouds is close to a power-law, and that may be a stochastic effect from cloud sampling when the cloud size is smaller than the beam. A comprehensive analysis of theoretical predictions for the observed PDF of HI and H_2 clouds below the spatial resolution limits of the surveys needs further investigation and will be addressed in forthcoming paper.

4. Discussion

Figures 3 and 5 indicate that the gas with high surface density in M33 has a PDF that is log-normal in the central regions and northern arm, and power-law for the molecular emission in the southern arm and filaments and in the interarm regions. The molecular gas has a power law PDF in general at mid-radii, and even the total gas PDF is somewhat power law at 3 to 4 kpc radius. The total gas has more of a log-normal PDF, presumably because it includes both the atomic and molecular phases with the latter being dominant at high column densities. The question we set out to address in this paper is where and how the transition from weakly self-gravitating gas to strongly self-gravitating clouds occurs. The most direct interpretation of the data presented in the previous Section is that this transition occurs at the same time as the gas converts from atoms to molecules, because the signature of self-gravity, the power law PDF, is primarily in the molecules at column densities close to Σ_{lim} . However, this is not necessarily the case. It could be that much of the HI is also part of the power-law cloud structure, perhaps even an extension of the same power law, in a shielding layer around the molecules, and we cannot see that because the turbulent compressions in the HI are much larger than the radial density range.

The power law PDF is usually interpreted as the result of a turbulent cloud with a radial gradient in the average density (Klessen 2000; Vázquez-Semadeni et al 2008; Kritsuk et al. 2011). The turbulence presumably gives each small region a log-normal PDF in volume density, but the centroid density of that PDF varies with radius in proportion to the average density (for a constant compressive Mach number). The sum of all the local log-normals in a whole cloud becomes a power law, over-

whelming the local log-normals, when the total density range from the radial gradient exceeds the fluctuating density range in each local piece of turbulence (Elmegreen 2011). This situation is fairly easy to achieve for the molecular gas because the density range in the radial gradient can be large, a factor of 100 or more, and that range exceeds the local compressions from turbulence at a modest Mach number. For an HI envelope the radial density range can be much smaller, perhaps only a factor of 10 from the cloud edge where the turbulent pressure equals the ambient pressure to the base of the envelope where molecules appear. If the density variations from turbulent compressions in the HI envelope exceed the average density variation over radius, then the power law component of the HI PDF may not be seen. The HI PDF at higher than average density mostly shows the sudden drop from the conversion to molecules, so the radial structure of the gas cannot be determined this way.

This model where turbulent fluctuations dominate density gradients in the atomic envelopes of clouds explains why the power law parts of the PDFs generally disappear in the total hydrogen gas, which approximately follows a log-normal with smaller widths at larger galactocentric radii. The HI and total gas are close to a log-normal form around the peak, even though the molecular parts of the clouds and perhaps each entire cloud has a power-law radial profile in density. The PDF slope levels off to a constant value at the high density, molecule-dominated end, where the internal power law density structure of the cloud dominates the turbulent density fluctuations in the molecules. For small clouds, the transition from atoms to molecules happens at scales that are not sampled by the 50 pc resolution of the present survey.

Power law PDFs at high column density can also appear even without self-gravity if the polytropic index, γ , is less than 1. According to Passot & Vázquez-Semadeni (1998), low γ leads to an increasing Mach number at higher density because the thermal temperature drops, and this causes higher density compressions than in an isothermal gas. The result is a broadening of the PDF at high density and a power law structure for the case they considered. However, cloud temperature gradients might reverse when star formation is taking place in the cloud core, like for C-type GMCs. While this γ effect may be present to some extent, numerical simulations of cloud assembly and collapse also show power laws just from self-gravity. For example, the PDF of the eastern filament (Fig. 3) resembles the PDF in Figure 2 of Ballesteros-Paredes et al. (2011) (panels relative to 13-14 Myrs) for numerical simulations of cloud formation and evolution by converging flows by Heitsch & Hartmann (2008). In this model GMCs assembly through thermal and gravity-driven instabilities in the atomic gas and they start collapsing around 10 Myrs with self-gravity driving departures of the PDF from a log-normal. Indeed, the eastern filament lies in an atomic-dominated region, similar to that of the numerical simulation, and the resolution and its inflow Mach number are also similar to that of our data (i.e., the Mach number is 1.5 resulting from a velocity dispersion of $\sigma_{21} = 11 \text{ km s}^{-1}$ in warm HI gas).

The shape of the column density PDF on large scales may also depend on the relative times spent in the turbulence-dominated phase and in the gravity-dominated phase. If gravitational contraction to a power-law density structure occurs during cloud formation, and feedback-driven turbulence occurs during cloud disruption, then regions with a gradual formation and a fast disruption should have a higher fraction of the mass in the pre- and early star formation phases when the power-law PDF is still apparent. Whether these phases show up in molecules or atoms depends also on the ease of forming molecules. In the

central regions of galaxies where the pressure is high, a high fraction of the ISM can be molecular, including even some non-gravitating clouds. This distinction may be relevant to our observation that regions close to the galaxy center have a molecular PDF that looks like a log-normal, while filaments further out have a molecular PDF that looks more like a power law. Near the center, the molecular PDF apparently includes more of the turbulent ISM and in that sense is like the atomic-gas PDF further out. The high pressure in the central regions increases the column density threshold for strong gravity, making this regime still undetectable at the resolution of the actual surveys. The filaments further out, on the other hand, tend to form molecules at the same time as the gas becomes self-gravitating, so the molecular PDF shows the power law density structure in that gas.

For the center and northern arm we notice that type-C GMCs show a wider PDF than type-A or B GMCs with a possible sign of gravity-driven power laws at very high column densities, of order 10^{22} cm^{-2} . For the southern arm, departures from a log-normal are present for all 3 types of GMCs, which share similar widths, for $N_H \approx 3 \times 10^{21} \text{ cm}^{-2}$. Clearly in these regions the power law regime driven by gravity is a short lived phase, prior to the birth of stars that stir again the GMCs, and molecular hydrogen forms rapidly without self-gravity (Dobbs et al. 2008).

What needs still to be investigated is the power law detected in the interarm PDF. The slopes are steeper there than in individual over-density regions, and the PDFs of individual GMC classes do not show strong deviations from a log-normal distribution. The number of pixels with high column density is low in these regions, so most for the detected CO does not show the core-halo cloud structure like in the southern arm (Figure 8). It might be that in the interarm regions, heating is more effective in warming up the outer envelopes of clouds, which are also of lower mass, so that we have $\gamma < 1$ without strong gravity. Or, for smaller mass clouds, the resolution of the survey is not enough to measure surface densities and detect the self-gravitating parts of individual clouds.

5. Summary

We examined in detail the probability distribution functions of the atomic and molecular ISM of M33 at a resolution of about 50 pc on a cloud-by-cloud basis and in the overall gas distribution in selected disk areas, to possibly unveil fingerprints of self-gravitating gas at these spatial scales. In particular we focused on spatially resolved GMC complexes and on sensitive high resolution atomic gas imaging to sample different regions across the star forming disk. As shown by Dobbs et al. (2018) gaseous spiral features, i.e. filamentary overdensities in the atomic gas, form as a result of gravitational instabilities in the disk of M33. At the same time, feedback must take place in order to form large cavities between the filaments, which can break as shock fronts propagates in the ISM. As the galactocentric distance increases, the rate of star formation per unit area, as well as the gravity due to the stellar surface density, decreases. It is not clear a priori where and if recycled turbulent gas driven by shocks in the ISM makes a fast transition to a gravity-driven regime, or else if the disk dynamical instabilities trigger H_2 formation and new star formation episodes. A first analysis of the PDF across the M33 disk, shown in this paper, suggest that both regimes can actually take place. We have sampled different regions to possibly trace non-gravitating turbulent complexes as well as gravity bound clouds in a statistical way using high resolution all-disk survey of the ISM in M33. We summarize below the main results:

- The face-on atomic gas column density distribution drops beyond $N_H^{lim} = 2.5 \times 10^{21} \text{ cm}^{-2}$ at all galactocentric radii; no HI is detected at $N_H > 2N_H^{lim}$. The peak of the atomic gas PDF is at about $10^{21} \text{ H cm}^{-2}$ (or about $0.5 N_H^{lim}$), with a slight increase of this value if we sample the envelope of molecular complexes. Around the PDF peak the HI here has a log-normal distribution. The slope of the PDF at low column densities becomes constant with a power-law structure, and this might be connected with the presence of cavities in the interarms where the warm/cold interface takes place, particularly if $\gamma > 1$.
- The width of the HI PDF has its maximum value at a distance of around 2-3 kpc from the center, likely due to the presence of the two main arms. Here the kinetic energy density in the atomic gas has its maximum value. The dispersion of the HI gas varies by about a factor 2 across the disk but radial averages show a constant trend with values of order 13 km s^{-1} and a radial decrease of only 1 km s^{-1} beyond 4 kpc. This constant trend suggest a ubiquitous presence of turbulence. The smaller width of the PDF at large galactocentric radii can be explained as a decrease of the Mach number due to disk flaring and to warm HI filling the ISM.
- The PDF of the molecular gas, as traced by CO, follows a log-normal in the central region. This applies also to the gas associated with GMCs at various evolutionary stages. The molecular PDF of the northern arm looks very similar to that of the central region. In the southern arm instead a power law distribution is detected around and beyond N_H^{lim} likely due to gravity shaping the structure of GMCs at all evolutionary stages, prior to and during star formation.
- The PDF of the molecular hydrogen in the outermost filament that is rich in CO shows a power law with indexes between -1 and -1.5. This is detected also in all the evolutionary phases of the GMCs which presumably assemble from atomic gas through thermal and gravitationally-driven dynamical instabilities and turbulence is less effective in shaping the cloud structure. There is a correspondence between different parts of the PDF power law and different parts of individual clouds, such as a core-envelope structure, for GMCs located in the outer filament and in the southern arm.

The analysis presented in this paper suggests that on intermediate scales self-gravity plays an early role in molecular cloud assembly and dominates the internal cloud structure in regions that are free of excessive feedback. The interpretation of the PDF shape for unresolved clouds in the diffuse molecular gas or interarm, which shows deviations from a log-normal distribution, requires some further investigation which will be presented elsewhere.

References

- Alves, J. Lombardi, M., Lada, C.J. 2017, *A&A*, 606, L2
 Baba, J., Morokuma-Matsui, K., Saitoh, T.R. 2017, *MNRAS*, 464, 246
 Ballesteros-Paredes, J., Vázquez-Semadeni, E., Gazol, A., et al. 2011, *MNRAS*, 416, 1436
 Bohlin, R. C., Savage, B. D., & Drake, J. F. 1978, *ApJ*, 224, 132
 Braine, J., Rosolowsky, E., Gratier, P., Corbelli, E., & Schuster, K.-F. 2018, *A&A*, 612, A51
 Braun, R. 2012, *ApJ*, 749, 87
 Braun, R., Thilker, D. A., Walterbos, R. A. M., & Corbelli, E. 2009, *ApJ*, 695, 937
 Colombo, D., Hughes, A., Schinnerer, E. 2014, *ApJ*, 784, 3
 Combes, F., Boquien, M., Kramer, C., et al. 2012, *A&A*, 539, A67
 Corbelli, E., Thilker, D., Zibetti, S., Giovanardi, C., & Salucci, P. 2014, *A&A*, 572, A23
 Corbelli, E., Braine, J., Bandiera, R., et al. 2017, *A&A*, 601, A146
 Deharveng, L., Zavagno, A., Samal, M. R. et al. 2015, *A&A*, 582, A1
 Dobbs, C. L., Glover, S. C. O., Clark, P. C., & Klessen, R. S. 2008, *MNRAS*, 389, 1097
 Dobbs, C. L., Pettitt, A. R., Corbelli, E., & Pringle, J. E. 2018, *MNRAS*, in press
 Donovan Meyer, J., Koda, J., Momose, R. et al. 2013, *ApJ*, 772, 107
 Druard, C., Braine, J., Schuster, K.F., et al. 2014, *A&A*, 567, A118
 Elmegreen, B. G., Leitner, S. N., Elmegreen, D. M., & Cuillandre, J.-C. 2003, *ApJ*, 593, 333
 Elmegreen, B.G. 2011, *ApJ*, 737, 10
 Elmegreen, D. M., Elmegreen, B.G., Erroz-Ferrer, S., et al. 2014, *ApJ*, 780, 32
 Elmegreen, B.G. 2018, *ApJ*, 854, 16
 Faesi, C.M., Lada, C.J., & Forbrich, J. 2018, arXiv180106238
 Federrath C. & Klessen R. S., 2013, *ApJ*, 763, 51
 Figueira, M., Zavagno, A., Deharveng, L., et al., 2017, *A&A*, 600, A93
 Froebrich, D., & Rowles, J. 2010, *MNRAS*, 406, 1350
 Freedman, W. L., Wilson, C. D., & Madore, B. F. 1991, *ApJ*, 372, 455
 Gardan, E., Braine, J., Schuster, K. F., Brouillet, N., & Sievers, A. 2007, *A&A*, 473, 91
 Gieren, W., Górski, M., Pietrzyński, G., et al. 2013, *ApJ*, 773, 69
 Girichidis, P., Konstandin, L., Whitworth, A. P., & Klessen, R. S. 2014, *ApJ*, 781, 91
 Gratier, P., Braine, J., Rodríguez-Fernandez, N. J., et al. 2012, *A&A*, 542, A108
 Gratier, P., Braine, J., Schuster, K., et al. 2017, *A&A*, 600, A27
 Heithausen, A., Bensch, F., Stutzki, J., Falgarone, E., & Panis, J. F. 1998, *A&A*, 331, L65
 Heitsch, F. & Hartmann, L. 2008, *ApJ*, 689, 290
 Hirota, A., Kuno, N., Sato, N., Nakanishi, H., Tosaki, T., & Sorai, K. 2011, *ApJ*, 737, 40
 Humphreys, R. M. & Sandage, A. 1980, *ApJS*, 44, 319
 Kainulainen, J., Beuther, H., Henning, T., & Plume, R. 2009, *A&A*, 508, L35
 Kainulainen, J., Beuther, H., Banerjee, R., et al. 2011, *A&A*, 530, A64
 Klessen, R. S. 2000, *ApJ*, 535, 869
 Kritsuk, A. G., Norman, M. L., & Wagner, R. 2011, *ApJL*, 727, L20
 Lombardi, M., Lada, C. J., & Alves, J. 2010, *A&A*, 512, A67
 Lombardi, M., Alves, J., Lada, C.J. 2015, *A&A*, 576, L1
 Magrini, L., Stanghellini, L., Corbelli, E., Galli, D., & Villaver, E. 2010, *A&A*, 512, A63
 Messa, M., Adamo, A., Calzetti, D., et al. 2018, *MNRAS*, 477, 1683
 Michalowski, M. J., Gentile, G., Hjorth, J., et al. 2015, *A&A*, 582, A78
 Mueller, K. E., Shirley, Y. L., Evans, N. J., II, & Jacobson, H. R. 2002, *ApJS*, 143, 469
 McKee, C.F., & Ostriker, E.C. 2007, *ARA&A*, 45, 565
 Meidt, S.E. 2016, *ApJ*, 818, 69
 Myers, P.C. 2015, *ApJ*, 806, 226
 Ossenkopf-Okada, V., Csengeri, T., Schneider, N., Federrath, C., & Klessen, R. S. 2016, *A&A*, 590, A104
 Padoan P., Nordlund A., Jones B. J. T., 1997, *MNRAS*, 288, 145
 Palmeirim, P., Zavagno, A., Elia, D. et al. 2017, *A&A*, 605, A35
 Pan, L., Padoan, P., Haugbølle, T., & Nordlund, Å., 2016, *ApJ*, 825, 30
 Passot, T., & Vázquez-Semadeni, E. 1998, *Phys.Rev.E*, 58, 4501
 Rosolowsky, E. 2005, *PASP*, 117, 1403
 Schneider, N. Csengeri, T., Hennemann, M. et al. 2012, *A&A*, 540, L11
 Schneider, N., André, Ph., Könyves, V., et al. 2013, *ApJ*, 766, L17
 Schneider, N. Csengeri, T., Klessen, R.S., Tremblin, P., Ossenkopf, V., Peretto, N., Simon, R., Bontemps, S., & Federrath, C. 2015a, *A&A*, 578, A29
 Schneider, N., Bontemps, S. Girichidis, P. et al. 2015b, *MNRAS*, 453, L41
 Schneider, N., Ossenkopf, V., Csengeri, T., Klessen, R.S., Federrath, C., Tremblin, P. Girichidis, P., Bontemps, S., & André, Ph. 2015, *A&A*, 575, A79
 Schneider, N., Bontemps, S. Motte, F., Ossenkopf, V., Klessen, R.S., Simon, R., Fechtenbaum, S., Herpin, F., Tremblin, P., Csengeri, T., Myers, P.C., Hill, T., Cunningham, M. & Federrath, C. 2016, *A&A*, 587, A74
 Sharma, S., Corbelli, E., Giovanardi, C., Hunt, L. K., & Palla, F. 2011, *A&A*, 534, A96
 Shu, F.H. 1977, *ApJ*, 214, 488
 Solomon, P. M., Rivolo, A. R., Barrett, J., & Yahil, A. 1987, *ApJ*, 319, 730
 Tamburro, D., Rix, H.-W., Leroy, A. K., et al. 2009, *AJ*, 137, 4424
 Tassis, K., Christie, D. A., Urban, A., et al. 2010, *MNRAS*, 408, 1089
 Tosaki, T., Kohno, K., Harada, N. et al. 2017, *PASJ*, 69, 18
 Vázquez-Semadeni, E. 1994, *ApJ*, 423, 681
 Vázquez-Semadeni E., González, R.F., Ballesteros-Paredes, J., Gazol, A., & Kim, J. 2008, *MNRAS*, 390, 769
 Verley, S., Corbelli, E., Giovanardi, C., & Hunt, L. K. 2009, *A&A*, 493, 453
 Wada, K., Spaans, M., & Kim, S. 2000, *ApJ*, 540, 797
 Whitworth, A., Lomax, O., Balfour, S., et al. 2018, *PASJ*, 70, S55