

# Correlation Functions of Classical and Quantum Artin System defined on Lobachevsky Plane and Scrambling Time

---

**Hrachya Babujian, Rubik Poghossian and George Savvidy**

*Institute of Nuclear and Particle Physics*

*Demokritos National Research Centre, Athens, Greece*

*Yerevan Physics Institute,*

*Alikhanian Br. 2, AM-0036 Yerevan, Armenia*

*E-mail:* [babujian@yerphi.am](mailto:babujian@yerphi.am), [poghos@yerphi.am](mailto:poghos@yerphi.am),  
[savvidy@inp.demokritos.gr](mailto:savvidy@inp.demokritos.gr)

**ABSTRACT:** We consider the quantisation of the Artin dynamical system defined on the fundamental region of the modular group. In classical regime the geodesic flow in the fundamental region represents one of the most chaotic dynamical systems, it has mixing of all orders, Lebesgue spectrum and non-zero Kolmogorov entropy. As a result, the classical correlation functions decay exponentially. In order to investigate the influence of the classical chaotic behaviour on the quantum-mechanical properties of the Artin system we calculated the corresponding thermal quantum-mechanical correlation functions. It was conjectured by Maldacena, Shenker and Stanford that the classical chaos can be diagnosed in thermal quantum systems by using an out-of-time-order correlation function as well as the square of the commutator of operators separated in time. We demonstrated that the two- and four-point correlation functions of the Liouville-like operators decay exponentially with a temperature dependent exponent. As conjectured the square of the commutator of the Liouville-like operators separated in time grows exponentially, similar to the exponential divergency of trajectories in the classical regime. The corresponding exponent does not saturate the maximal growth condition.

**KEYWORDS:** Artin billiard, Chaotic dynamical systems, Anosov systems, Kolmogorov systems, Modular invariance, Non holomorphic Automorphic functions, Quantum and Classical correlation functions, Scrambling time.

---

## Contents

<b>1</b>	<i>Introduction</i>	<b>1</b>
<b>2</b>	<i>Lobachevsky plane and its isometry group</i>	<b>4</b>
<b>3</b>	<i>Geodesic flow in Hamiltonian gauge</i>	<b>5</b>
<b>4</b>	<i>Quantization</i>	<b>7</b>
<b>5</b>	<i>Continuous spectrum and the reflection amplitude</i>	<b>9</b>
<b>6</b>	<i>Correlation functions</i>	<b>14</b>
6.1	<i>Two-point correlation function</i>	14
6.2	<i>Four-point correlation function</i>	16
<b>7</b>	<i>Acknowledgment</i>	<b>19</b>
<b>8</b>	<i>Appendix</i>	<b>20</b>

---

## 1 Introduction

The hyperbolic Anosov C-systems have exponential instability of their trajectories and as such represent the most natural chaotic dynamical systems [7]. Of special interest are C-systems which are defined on closed surfaces of the Lobachevsky plane of constant negative curvature. An example of such system has been introduced in a brilliant article published in 1924 by the mathematician Emil Artin [1]. The dynamical system is defined on the fundamental region of the Lobachevsky plane which is obtained by the identification of points congruent with respect to the modular group  $SL(2, Z)$ , a discrete subgroup of the Lobachevsky plane isometries [2–4]. The fundamental region  $\mathcal{F}$  in this case is a hyperbolic triangle on Fig.1. The geodesic trajectories are bounded to propagate on the fundamental hyperbolic triangle. The geodesic flow in this fundamental region represents one of the most chaotic dynamical systems with exponential instability of its trajectories, has mixing of all orders, Lebesgue spectrum and non-zero Kolmogorov entropy [5–16, 18–20, 22, 23, 32–37, 40].

There is a great interest in considering quantisation of the hyperbolic dynamical systems and investigating their quantum-mechanical properties [22, 23]. Our main interest in this article is a study of the behaviour of the correlation functions of the Artin hyperbolic dynamical system in its classical and quantum-mechanical regimes.

In classical regime the correlation functions are defined as an integral over a pair of functions/observables  $A$  and  $B$  in which the first one is stationary and the second one evolves with the geodesic flow  $g_t$ :

$$\mathcal{D}_t(A, B) = \int_{\mathcal{M}} A(g) \overline{B(gg_t)} d\mu(g). \quad (1.1)$$

The earlier investigation of the classical correlation functions of the geodesic flows was performed in [16–20] by using different approaches including Fourier series for the  $SL(2, R)$  group, zeta function for the geodesic flows, relating the poles of the Fourier transform of the correlation functions to the spectrum of an associated Ruelle operator, the methods of unitary representation theory, spectral properties of the corresponding Laplacian and other approaches. In recent articles [38, 39] the authors demonstrated exponential decay of the correlation functions with time on the classical phase space. The result was derived by using the differential geometry, group-theoretical methods of Gelfand and Fomin, the time evolution equations and the properties of automorphic functions on  $\mathcal{F}$ . The exponential decay rate was expressed in terms of the entropy  $h(\mathcal{F})$  of the system:

$$|\mathcal{D}_t(A, B)| \leq M e^{-Kh|t|}, \quad (1.2)$$

where  $M$  and  $K$  are constants depending on the smoothness of the functions. *In classical regime the exponential divergency of the geodesic trajectories resulted into the universal exponential decay of its classical correlation functions* [38, 39].

In order to investigate the behaviour of the correlation functions in quantum-mechanical regime it is necessary to know the spectrum of the system and the corresponding wave functions. In the case of the modular group the energy spectrum has continuous part, which is originating from the asymptotically free motion inside an infinitely long "y-channel" extended in the vertical direction of the fundamental region, as well as infinitely many discrete energy states corresponding to a bounded motion at the "bottom" of the fundamental triangle. The spectral problem has deep number-theoretical origin and was partially solved in a series of pioneering articles [24–27]. It was solved partially because the discrete spectrum and the corresponding wave functions are not known analytically. The general properties of the discrete spectrum have been derived by using Selberg trace formula [26–31]. Numerical calculation of the discrete energy levels were performed for many energy states [42–44].

In the next section we shall describe the geometry of Lobachevsky hyperbolic plane and of the fundamental region which corresponds to the modular group  $SL(2, Z)$ , the geodesic flow on that region and the quantisation of the system. The derivation of the Maass wave functions [24] for the continuous spectrum will be reviewed in details. We shall use the Poincaré representation for Maass non-holomorphic automorphic wave functions. We introduce a natural physical variable  $\tilde{y}$  for the distance

in the vertical direction on the fundamental triangle  $\int dy/y = \ln y = \tilde{y}$  and the corresponding momentum  $p_y$  in order to represent the Maass wave functions (5.22) in the form which is appealing to the physical intuition

$$\psi_{p_y}(x, \tilde{y}) = e^{-ip_y\tilde{y}} + \frac{\theta(\frac{1}{2} + ip_y)}{\theta(\frac{1}{2} - ip_y)} e^{ip_y\tilde{y}} + \frac{4}{\theta(\frac{1}{2} - ip_y)} \sum_{l=1}^{\infty} \tau_{ip_y}(l) K_{ip_y}(2\pi l e^{\tilde{y}}) \cos(2\pi l x). \quad (1.3)$$

Indeed, the first two terms describe the incoming and outgoing plane waves. The plane wave  $e^{-ip_y\tilde{y}}$  incoming from infinity of the  $y$  axis on Fig. 1 (the vertex  $\mathcal{D}$ ) elastically scatters on the boundary  $ACB$  of the fundamental triangle  $\mathcal{F}$  on Fig. 1. The reflection amplitude is a pure phase and is given by the expression in front of the outgoing plane wave  $e^{ip_y\tilde{y}}$ :

$$\frac{\theta(\frac{1}{2} + ip_y)}{\theta(\frac{1}{2} - ip_y)} = \exp [i \varphi(p_y)]. \quad (1.4)$$

The rest of the wave function describes the standing waves  $\cos(2\pi l x)$  in the  $x$  direction between boundaries  $x = \pm 1/2$  with the amplitudes  $K_{ip_y}(2\pi l y)$ , which are exponentially decreasing with index  $l$  Fig.2. The continuous energy spectrum is given by the formula

$$E = p_y^2 + \frac{1}{4}. \quad (1.5)$$

The wave functions of the discrete spectrum have the following form [24–27, 42–44]:

$$\psi_n(z) = \sum_{l=1}^{\infty} c_l(n) K_{iu_n}(2\pi l e^{\tilde{y}}) \begin{cases} \cos(2\pi l x) \\ \sin(2\pi l x) \end{cases}, \quad (1.6)$$

where the spectrum  $E_n = \frac{1}{4} + u_n^2$  and the coefficients  $c_l(n)$  are not known analytically, but were computed numerically for many values of  $n$  [42–44].

Having in hand the explicit expression of the wave function one can analyse a quantum-mechanical behaviour of the correlation functions defined in [36]:

$$\mathcal{D}_2(\beta, t) = \langle A(t)B(0)e^{-\beta H} \rangle, \quad \mathcal{D}_4(\beta, t) = \langle A(t)B(0)A(t)B(0)e^{-\beta H} \rangle \quad (1.7)$$

$$C(\beta, t) = -\langle [A(t), B(0)]^2 e^{-\beta H} \rangle, \quad (1.8)$$

where in our case the operators  $A$  and  $B$  are chosen to be of the Liouville type:

$$A(N) = e^{-2N\tilde{y}}, \quad N = 1, 2, \dots \quad (1.9)$$

Analysing the basic matrix elements of the Liouville-like operators (1.9) we shall demonstrate that all two- and four-point correlation functions (1.7) decay exponentially with time, with the exponents which depend on temperature Fig.3 and Fig.4. These exponents define the decorrelation time  $t_d(\beta)$ .

Alternatively to the exponential decay of the correlation functions (1.7) the square of the commutator of the Liouville-like operators separated in time (1.8) grows exponentially Fig.5. This growth is reminiscent to the local exponential divergence of trajectories in the Artin system when it is considered in the classical regime [38, 39]. The exponential growth Fig.5 of the commutator (1.8) does not saturate the condition of maximal growth (6.24) and (6.25)

$$C(\beta, t) \sim f(\beta) e^{\frac{2\pi}{\chi(\beta)} t}, \quad \chi(\beta) \sim \beta \quad (1.10)$$

of the correlation functions which is conjectured to be linear in temperature  $T = 1/\beta$

$$C(\beta, t) \sim f(\beta) e^{\frac{2\pi}{\beta} t}. \quad (1.11)$$

In our calculation of the quantum-mechanical correlation functions we shall use a perturbative expansion in which the high-mode Bessel's functions in (5.22) and (5.27) will be considered as perturbations. We found that our calculations are stable with respect to these perturbations and do not influence the final results. The reason is that in the integration region of the matrix elements (6.3) the high-mode Bessel's functions are exponentially small.

## 2 Lobachevsky plane and its isometry group

Let us start with Poincare model of the Lobachevsky plane, i.e. the upper half of the complex plane:  $H = \{z \in \mathbb{C}, \Im z > 0\}$  supplied with the metric (we set  $z = x + iy$ )

$$dl^2 = \frac{dx^2 + dy^2}{y^2} \quad (2.1)$$


with the Ricci scalar  $R = -2$ . Isometries of this space are given by  $SL(2, \mathbb{R})$  transformations. The  $SL(2, \mathbb{R})$  matrix ( $a, b, c, d$  are real and  $ad - bc = 1$ )

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

acts on a point  $z$  by linear fractional substitutions:

$$z \rightarrow \frac{az + b}{cz + d}. \quad (2.2)$$

Note also that  $g$  and  $-g$  give the same transformation, hence the effective group is  $SL(2, \mathbb{R})/\mathbb{Z}_2$ . We'll be interested in the space of orbits of a discrete subgroup  $G \subset SL(2, \mathbb{R})$  in  $H$ . Our main example will be the modular group  $G = SL(2, \mathbb{Z})$ . A nice choice of the fundamental region  $\mathcal{F}$  of  $SL(2, \mathbb{Z})$  is displayed in Fig.1. The fundamental region  $\mathcal{F}$  of the modular group consists of those points between the lines  $x = -\frac{1}{2}$  and  $x = +\frac{1}{2}$  that lie outside the unit circle in Fig.1. The modular triangle



**Figure 1:** The non-compact fundamental region  $\mathcal{F}$  of a finite area is represented by the hyperbolic triangle  $ABD$ . The vertex  $D$  is at infinity of the  $y$  axis and corresponds to a cusp. The edges of the triangle are the arc  $AB$ , the rays  $AD$  and  $BD$ . The points on the edges  $AD$  and  $BD$  and the points of the arcs  $AC$  with  $CB$  should be identified by the transformations  $w = z + 1$  and  $w = -1/z$  in order to form a *closed non-compact surface*  $\bar{\mathcal{F}}$  by "gluing" the opposite edges of the modular triangle together. The hyperbolic triangle  $OAB$  can be considered equally well as the fundamental region. The modular transformations of the fundamental region  $\mathcal{F}$  create a regular tessellation of the whole Lobachevsky plane by congruent hyperbolic triangles.  $K$  is a geodesic trajectory passing through the point  $(x, y)$  of  $\mathcal{F}$  in the  $\vec{v}$  direction.

$\mathcal{F}$  has two equal angles  $\alpha = \beta = \frac{\pi}{3}$  and with the third one equal to zero,  $\gamma = 0$ , thus  $\alpha + \beta + \gamma = 2\pi/3 < \pi$ . The area of the fundamental region is finite and equals to  $\frac{\pi}{3}$  and gets a topology of sphere by "gluing" the opposite edges of the triangle. The invariant area element on the Lobachevsky plane is proportional to the square root of the determinant of the matrix (2.1):

$$d\mu(z) = \frac{dx dy}{y^2}. \quad (2.3)$$

thus

$$\text{Area}(\mathcal{F}) = \int_{-\frac{1}{2}}^{\frac{1}{2}} dx \int_{\sqrt{1-x^2}}^{\infty} \frac{dy}{y^2} = \frac{\pi}{3}.$$

### 3 Geodesic flow in Hamiltonian gauge

Consider geodesic flow on  $\mathcal{F}$ , which is conveniently described by the least action principle  $\delta S = 0$ , where (cf. with (2.1)):

$$S = \int L dt = \int \frac{\sqrt{\dot{x}^2 + \dot{y}^2}}{y} dt. \quad (3.1)$$

By varying the action, we immediately get the equations of motion

$$\begin{aligned}\frac{d}{dt} \frac{\dot{x}}{y\sqrt{\dot{x}^2 + \dot{y}^2}} &= 0, \\ \frac{d}{dt} \frac{\dot{y}}{y\sqrt{\dot{x}^2 + \dot{y}^2}} + \frac{\sqrt{\dot{x}^2 + \dot{y}^2}}{y^2} &= 0.\end{aligned}\tag{3.2}$$

Notice the invariance of the action and of the equations under time reparametrizations  $t \rightarrow t(\tau)$ . Presence of a local ("gauge") symmetry indicates that we have a constrained dynamical system. One particularly convenient choice of gauge fixing specifying the time parameter  $t$  proportional to the proper time, is achieved by imposing the condition

$$\frac{\dot{x}^2 + \dot{y}^2}{y^2} = 2H,\tag{3.3}$$

where  $H$  is a constant. In this gauge the equations (3.2) will take the following form [21]:

$$\begin{aligned}\frac{d}{dt} \left( \frac{\dot{x}}{y^2} \right) &= 0 \\ \frac{d}{dt} \left( \frac{\dot{y}}{y^2} \right) + \frac{2H}{y} &= 0.\end{aligned}\tag{3.4}$$

Defining the canonical momenta  $p_x, p_y$  conjugate to the coordinates  $x, y$  as

$$p_x = \frac{\dot{x}}{y^2}, \quad p_y = \frac{\dot{y}}{y^2},\tag{3.5}$$

we shall get the geodesic equations (3.4) in the Hamiltonian form:

$$\dot{p}_x = 0, \quad \dot{p}_y = -\frac{2H}{y}.\tag{3.6}$$


Indeed, after defining the Hamiltonian as

$$H = \frac{1}{2}y^2(p_x^2 + p_y^2)\tag{3.7}$$

the corresponding equations will take the following form:

$$\begin{aligned}\dot{x} = \frac{\partial H}{\partial p_x} &= y^2 p_x, & \dot{y} = \frac{\partial H}{\partial p_y} &= y^2 p_y \\ \dot{p}_x = -\frac{\partial H}{\partial x} &= 0, & \dot{p}_y = -\frac{\partial H}{\partial y} &= -y(p_x^2 + p_y^2) = -\frac{2H}{y}\end{aligned}\tag{3.8}$$

and coincide with (3.5) and (3.6). The advantage of the gauge (3.3) is that the Hamiltonian (3.7) coincides with the constraint.



**Figure 2:** The incoming and outgoing plane waves. The plane wave  $e^{-ip\tilde{y}}$  incoming from infinity of the  $y$  axis on Fig. 1 (the vertex  $\mathcal{D}$ ) elastically scatters on the boundary  $ACB$  of the fundamental triangle  $\mathcal{F}$  on Fig. 1. The reflection amplitude is a pure phase and is given by the expression in front of the outgoing plane wave  $e^{ip\tilde{y}}$ . The rest of the wave function describes the standing waves in the  $x$  direction between boundaries  $x = \pm 1/2$  with the amplitudes, which are exponentially decreasing.

#### 4 Quantization

Now it is fairly standard to quantize this Hamiltonian system. We simply replace in (3.7)

$$p_x = -i \frac{\partial}{\partial x}, \quad p_y = -i \frac{\partial}{\partial y}$$

and consider the (time independent) Schrödinger equation

$$H\psi = E\psi.$$

The resulting equation explicitly reads:

$$-y^2(\partial_x^2 + \partial_y^2)\psi = E\psi. \quad (4.1)$$

On the lhs one easily recognises the Laplace operator (with an extra minus sign) in Poincare metric (2.1). It is easy to see that the Hamiltonian is positive semi-definite Hermitian operator. Indeed, for any quadratically integrable function  $\psi(x, y)$

$$-\int \psi^*(x, y) y^2(\partial_x^2 + \partial_y^2) \psi(x, y) \frac{dx dy}{y^2} = \int (|\partial_x \psi(x, y)|^2 + |\partial_y \psi(x, y)|^2) dx dy \geq 0. \quad (4.2)$$



It is convenient to introduce parametrization of the energy  $E = s(1 - s)$  and to rewrite this equation as

$$-y^2(\partial_x^2 + \partial_y^2) \psi(x, y) = s(1 - s) \psi(x, y). \quad (4.3)$$

As far as  $E$  is real and semi-positive and parametrisation is symmetric with respect to  $s \leftrightarrow 1 - s$  it follows that the parameter  $s$  should be chosen within the range

$$s \in [1/2, 1] \quad \text{or} \quad s = 1/2 + iu, \quad u \in [0, \infty]. \quad (4.4)$$

One should impose the "periodic" boundary condition on the wave function with respect to the modular group

$$\psi\left(\frac{az + b}{cz + d}\right) = \psi(z), \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, Z) \quad (4.5)$$

in order to have the wave function which is properly defined on the fundamental region  $\bar{\mathcal{F}}$  shown in Fig. 1. Taking into account that the transformation  $T : z \rightarrow z + 1$  belongs to  $SL(2, Z)$ , one has to impose the periodicity condition  $\psi(z) = \psi(z + 1)$ . Thus we have a Fourier expansion

$$\psi(x, y) = \sum_{n=-\infty}^{\infty} f_n(y) \exp(2\pi i n x). \quad (4.6)$$

Inserting this into Eq. (4.3), for the Fourier component  $f_n(y)$  we get

$$\frac{d^2 f_n(y)}{dy^2} + (s(1 - s) - 4\pi^2 n^2) f_n(y) = 0. \quad (4.7)$$

For the case  $n \neq 0$  the solution which exponentially decays at large  $y$  reads

$$f_n(y) = \sqrt{y} K_{s-\frac{1}{2}}(2\pi n |y|), \quad (4.8)$$

and for  $n = 0$  one simply gets

$$f_0(y) = c_0 y^s + c'_0 y^{1-s}. \quad (4.9)$$

Thus the solution can be represented in the following form:

$$\psi(x, y) = c_0 y^s + c'_0 y^{1-s} + \sqrt{y} \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} c_n K_{s-\frac{1}{2}}(2\pi n |y|) \exp(2\pi i n x), \quad (4.10)$$

where the coefficients  $c_0, c'_0, c_n$  should be defined so that the wave function fulfils the boundary conditions (4.5). Thus one should impose also the invariance with respect

to the second generator of the modular group  $SL(2, Z)$ , that is, with respect to the transformation  $S : z \rightarrow -1/z$  :

$$\psi(z) = \psi(-1/z). \quad (4.11)$$

This functional equation defines the coefficients  $c_0, c'_0, c_n$ . We found that it is much easier to resolve it by using the full group of  $SL(2, Z)$  transformations acting on a particular solution (4.9). The wave function generated in this way will be invariant with respect to the  $SL(2, Z)$  transformations. We shall follow this approach in the next section.

## 5 Continuous spectrum and the reflection amplitude

As we just mentioned above in order to get  $SL(2, Z)$  invariant solutions, one should define the coefficients  $c_0, c'_0$  and  $c_n$  in (4.10). Another option is to start from a particular solution and perform summation over all nonequivalent shifts of the argument by the elements of  $SL(2, Z)$ , that is, by using the Poincaré series representation [2, 3]. We'll demonstrate this strategy by using the simplest solution (4.9), (4.10) with  $c_0 = 1, c'_0 = 0$ :

$$\psi(z) = y^s = (\Im z)^s.$$

Let us denote by  $\Gamma_\infty$  the subgroup of  $\Gamma = SL(2, Z)$ , generating shifts  $z \rightarrow z + n$ ,  $n \in Z$ . Explicitly the elements of  $\Gamma_\infty$  are given by  $2 \times 2$  matrices:

$$g_n = \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}. \quad (5.1)$$

Since  $y^s$  is already invariant with respect to  $\Gamma_\infty$ , we should perform summation over the conjugacy classes  $\Gamma_\infty \backslash \Gamma$ . Let us define these conjugacy classes. If two  $SL(2, Z)$  matrices

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix}$$

belong to the same class, then by definition for some  $n \in Z$

$$\begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} = \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

so that  $c' = c$ ,  $d' = d$ ,  $a' - a = nc$  and  $b' - b = nd$ . Since  $ad - bc = 1$ , it follows that  $a$  and  $c$  do not have a common divisor. In fact, the opposite is also true. Given a pair of mutually prime integers  $(c, d)$  it is always possible to find a pair of integers  $(a, b)$  such that  $ad - bc = 1$ . For any other pair  $(a', b')$  satisfying the same condition  $a'd - b'c = 1$ , the relations  $a' - a = nc$  and  $b' - b = nd$  are satisfied for some integer  $n$ . Thus we established a bijection between the set of mutually prime pairs  $(c, d)$

with  $(c, d) \neq (0, 0)$  and the set of conjugacy classes  $\Gamma_\infty \backslash \Gamma$ . The fact that the integers  $(c, d)$  are mutually prime integers means that their greatest common divisor (gcd) is equal to one:  $\text{gcd}(c, d) = 1$ . As a result, it is defined by the classical Poincaré series representation [2, 3] and for the sum of our interest we get<sup>1</sup>

$$\psi_s(z) \equiv \frac{1}{2} \sum_{\gamma \in \Gamma_\infty \backslash \Gamma} (\mathfrak{S}(\gamma z))^s = \frac{1}{2} \sum_{\substack{(c,d) \in \mathbb{Z}^2 \\ \text{gcd}(c,d)=1}} \frac{y^s}{((cx+d)^2 + c^2y^2)^s}, \quad (5.2)$$

where, as explained above, the sum on r.h.s. is taken over all mutually prime pairs  $(c, d)$ . The series (5.2) is convergent when  $\Re s > 1$ . We used also the simple relation

$$\mathfrak{S}\gamma z \equiv \mathfrak{S} \frac{az+b}{cz+d} = \frac{y}{(cx+d)^2 + c^2y^2}.$$

To simplify further the sum let us multiply both sides of the eq. (5.2) by [24]

$$\sum_{n=1}^{\infty} \frac{1}{n^{2s}} \equiv \zeta(2s)$$

so that we shall get

$$\zeta(2s) \psi_s(z) = \frac{1}{2} \sum_{n=1}^{\infty} \sum_{\substack{(c,d) \in \mathbb{Z}^2 \\ \text{gcd}(c,d)=1}} \frac{y^s}{((ncx+nd)^2 + (nc)^2y^2)^s}. \quad (5.3)$$

It is easy to get convinced now that the set of all pairs  $(nc, nd)$  with  $n$  a positive integer and  $(c, d)$  - mutually prime, coincides with the set of all pairs of integers  $(m, k)$  which are not simultaneously zero. Indeed, given a pair  $(m, k)$  we can factor out the greatest common divisor  $n$  and represent it as  $(nc, nd)$  with mutually prime  $(c, d)$ . Thus we arrive at the Eisenstein series representation of the wave function:

$$\zeta(2s) \psi_s(z) = \frac{1}{2} \sum_{\substack{(m,k) \in \mathbb{Z}^2 \\ (m,k) \neq (0,0)}} \frac{y^s}{((mx+k)^2 + m^2y^2)^s}. \quad (5.4)$$

Since the r.h.s. of this equation is periodic in  $x$  with period 1, we can expand it in Fourier series. Our next goal is to find the coefficients of this expansion:

$$c_l(y) = \frac{1}{2} \sum_{\substack{(m,k) \in \mathbb{Z}^2 \\ (m,k) \neq (0,0)}} \int_0^1 \frac{y^s e^{-2\pi i l x} dx}{((mx+k)^2 + m^2y^2)^s}. \quad (5.5)$$

First let us handle the term with  $m = 0$ :

$$\frac{1}{2} \sum_{\substack{k \in \mathbb{Z} \\ k \neq 0}} \int_0^1 \frac{y^s e^{-2\pi i l x} dx}{k^{2s}} = \delta_{l,0} \zeta(2s) y^s. \quad (5.6)$$

---

<sup>1</sup> The factor 1/2 below is introduced for removing the double degeneracy due to the fact that  $SL(2, Z)$  elements  $\pm\gamma$  both act on  $z$  in the same way.

For the sum over non-zero  $m$ 's let's notice that we may drop the factor  $1/2$  and sum over  $m \geq 1$ . Indeed, the sum over negative  $m$ 's can be reverted to a sum over positive ones through redefinition  $k \rightarrow -k$ . For fixed positive  $m$  it is instructive to represent  $k$  as  $k = nm + s$  thus splitting the initial sum over  $k \in \mathbb{Z}$  into double sum over  $r = 0, 1, \dots, m-1$  and  $n \in \mathbb{Z}$ . In this way after few simple manipulations we get

$$\begin{aligned} & \sum_{m=1}^{\infty} \sum_{r=0}^{m-1} \sum_{n \in \mathbb{Z}} \int_0^1 \frac{y^s e^{-2\pi i l x} dx}{((m(x+n)+r)^2 + m^2 y^2)^s} = \sum_{m=1}^{\infty} \sum_{r=0}^{m-1} \int_{-\infty}^{\infty} \frac{y^s e^{-2\pi i l x} dx}{((mx+r)^2 + m^2 y^2)^s} \\ & = \sum_{m=1}^{\infty} \sum_{r=0}^{m-1} m^{-2s} y^{1-s} e^{\frac{2\pi i l r}{m}} \int_{-\infty}^{\infty} \frac{\cos(2\pi |l| y x) dx}{(x^2 + 1)^s}. \end{aligned} \quad (5.7)$$

The last integral is expressed in terms of modified Bessel's  $K$  function:

$$\int_{-\infty}^{\infty} \frac{\cos(2\pi |l| y x) dx}{(x^2 + 1)^s} = \begin{cases} \frac{2\pi^s}{\Gamma(s)} |ly|^{s-\frac{1}{2}} K_{s-\frac{1}{2}}(2\pi |l| y), & \text{if } l \neq 0 \\ \frac{\sqrt{\pi} \Gamma(s-\frac{1}{2})}{\Gamma(s)}, & \text{if } l = 0. \end{cases} \quad (5.8)$$

A useful alternative representation of modified Bessel's  $K$  function which makes its properties more transparent is given by

$$K_{i\nu}(y) = \frac{1}{2} \int_{-\infty}^{\infty} e^{-y \cosh t} e^{i\nu t} dt. \quad (5.9)$$

This expression allows analytical continuation of the wave function from the region  $\Re s > 1$  in (5.2) into the whole complex plane  $s$  because the Bessel's  $K_s(y)$  functions are well defined for any  $s$ . Besides, an easy examination shows that the finite sum is:

$$\sum_{r=0}^{m-1} e^{\frac{2\pi i l r}{m}} = \begin{cases} m & \text{if } m \text{ divides } l \\ 0 & \text{otherwise} \end{cases}. \quad (5.10)$$

To summarise, for the Fourier coefficients (5.5) we shall get

$$c_l(y) = \frac{2\pi^s}{\Gamma(s)} \tau_{s-\frac{1}{2}}(|l|) \sqrt{y} K_{s-\frac{1}{2}}(2\pi |l| y), \quad \text{if } l \neq 0, \quad (5.11)$$

where

$$\tau_\nu(n) = \sum_{a \cdot b = n} \left(\frac{a}{b}\right)^\nu, \quad (5.12)$$

while for  $l = 0$ :

$$c_0(y) = \frac{\sqrt{\pi} \Gamma(s-\frac{1}{2}) \zeta(2s-1)}{\Gamma(s)} y^{1-s}. \quad (5.13)$$

Thus we recovered the second solution  $y^{1-s}$  in (4.10) and calculated the coefficient  $c'_0$  in front of it. Thus the invariant solution (5.4) takes the following form:

$$\begin{aligned} \zeta(2s) \psi_s(x, y) &= \zeta(2s) y^s + \frac{\sqrt{\pi} \Gamma(s - \frac{1}{2}) \zeta(2s - 1)}{\Gamma(s)} y^{1-s} + \\ &+ \sqrt{y} \frac{4\pi^s}{\Gamma(s)} \sum_{l=1}^{\infty} \tau_{s-\frac{1}{2}}(l) K_{s-\frac{1}{2}}(2\pi l y) \cos(2\pi l x). \end{aligned} \quad (5.14)$$

Using Riemann's reflection relation

$$\zeta(s) = \frac{\pi^{s-\frac{1}{2}} \Gamma(\frac{1-s}{2})}{\Gamma(\frac{s}{2})} \zeta(1-s) \quad (5.15)$$

and introducing the notation

$$\theta(s) = \pi^{-s} \zeta(2s) \Gamma(s) \quad (5.16)$$

we arrive at the elegant final expression for the energy eigenfunctions obtained by Maas [24]:

$$\theta(s) \psi_s(z) = \theta(s) y^s + \theta(1-s) y^{1-s} + 4\sqrt{y} \sum_{l=1}^{\infty} \tau_{s-\frac{1}{2}}(l) K_{s-\frac{1}{2}}(2\pi l y) \cos(2\pi l x), \quad (5.17)$$

This wave function is well defined in the complex  $s$  plane and has a simple pole at  $s = 1$ . The physical continuous spectrum was defined in (4.4), where  $s = \frac{1}{2} + iu$ ,  $u \in [0, \infty]$  so that

$$E = s(1-s) = \frac{1}{4} + u^2. \quad (5.18)$$

The continuous spectrum wave functions  $\psi_s(x, y)$  are delta function normalisable [24–29]. The wave function (5.17) can be conveniently represented also in the form

$$\psi_{\frac{1}{2}+iu}(z) = y^{\frac{1}{2}+iu} + \frac{\theta(\frac{1}{2}-iu)}{\theta(\frac{1}{2}+iu)} y^{\frac{1}{2}-iu} + \frac{4\sqrt{y}}{\theta(\frac{1}{2}+iu)} \sum_{l=1}^{\infty} \tau_{iu}(l) K_{iu}(2\pi l y) \cos(2\pi l x), \quad (5.19)$$

where

$$K_{-iu}(y) = K_{iu}(y), \quad \tau_{-iu}(l) = \tau_{iu}(l). \quad (5.20)$$

The physical interpretation of the wave function becomes more transparent when we introduce the new variables

$$\tilde{y} = \ln y, \quad p = -u, \quad E = p^2 + \frac{1}{4} \quad (5.21)$$

as well as an alternative normalisation of the wave function:

$$\begin{aligned}\psi_p(x, \tilde{y}) &\equiv y^{-\frac{1}{2}} \psi_{\frac{1}{2}+iu}(z) = \\ &= e^{-ip\tilde{y}} + \frac{\theta(\frac{1}{2}+ip)}{\theta(\frac{1}{2}-ip)} e^{+ip\tilde{y}} + \frac{4}{\theta(\frac{1}{2}-ip)} \sum_{l=1}^{\infty} \tau_{ip}(l) K_{ip}(2\pi l e^{\tilde{y}}) \cos(2\pi l x).\end{aligned}\tag{5.22}$$

Indeed, the first two terms describe the incoming and outgoing plane waves. The plane wave  $e^{-ip\tilde{y}}$  incoming from infinity of the  $y$  axis on Fig.1-2 ( the vertex  $\mathcal{D}$ ) elastically scatters on the boundary  $ACB$  of the fundamental region  $\mathcal{F}$  on Fig. 1. The reflection amplitude is a pure phase and is given by the expression in front of the outgoing plane wave  $e^{ip\tilde{y}}$ :

$$\frac{\theta(\frac{1}{2} + ip)}{\theta(\frac{1}{2} - ip)} = \exp [i \varphi(p)].\tag{5.23}$$

The rest of the wave function describes the standing waves  $\cos(2\pi l x)$  in the  $x$  direction between boundaries  $x = \pm 1/2$  with the amplitudes  $K_{ip}(2\pi l y)$ , which are exponentially decreasing with index  $l$ .

In addition to the continuous spectrum the system (4.3) has a discrete spectrum [24–29]. The number of discrete states is infinite:  $E_0 = 0 < E_1 < E_2 < \dots \rightarrow \infty$ , the spectrum is extended to infinity - unbounded from above - and lacks any accumulation point except infinity. Let us denote the wave functions of the discrete spectrum by  $\psi_n(z)$  so that the expansion into the full set of basis vectors will take the form

$$f(x, \tilde{y}) = \sum_{n \geq 0} a_n \psi_n(x, \tilde{y}) + \frac{1}{2\pi} \int_0^{\infty} a_p \psi_p(x, \tilde{y}) dp\tag{5.24}$$

and the Parseval identity will be

$$\|f(z)\|^2 = \sum_{n \geq 0} |a_n|^2 + \frac{1}{2\pi} \int_0^{\infty} |a_p|^2 dp,\tag{5.25}$$

and

$$\sum_{n \geq 0} \psi_n(x, \tilde{y}) \psi_n^*(x_1, \tilde{y}_1) + \frac{1}{2\pi} \int_0^{\infty} \psi_p(x, \tilde{y}) \psi_{-p}(x_1, \tilde{y}_1) dp = \delta^{(2)}(z - z_1).\tag{5.26}$$

The wave functions of the discrete spectrum have the following form [24–27, 42–44]:

$$\psi_n(z) = \sum_{l=1}^{\infty} c_l(n) \sqrt{\tilde{y}} K_{iu_n}(2\pi l y) \begin{cases} \cos(2\pi l x) \\ \sin(2\pi l x) \end{cases},\tag{5.27}$$

where the spectrum  $E_n = \frac{1}{4} + u_n^2$  and the coefficients  $c_l(n)$  are not known analytically, but were computed numerically for many values of  $n$  [42–44]. Having explicit expressions of the wave functions one can analyse the quantum-mechanical behaviour of the correlation functions, which we shall investigate in the next sections.

## 6 Correlation functions

### 6.1 Two-point correlation function

First let us calculate the two-point correlation function:

$$\begin{aligned} \mathcal{D}_2(\beta, t) &= \langle A(t)B(0)e^{-\beta H} \rangle = \sum_n \langle n | e^{iHt} A(0) e^{-iHt} B(0) e^{-\beta H} | n \rangle = \\ &= \sum_{n,m} e^{i(E_n - E_m)t - \beta E_n} \langle n | A(0) | m \rangle \langle m | B(0) | n \rangle. \end{aligned} \quad (6.1)$$

The energy eigenvalues (5.18) are parametrised by  $n = \frac{1}{2} + iu$ ,  $E_n = \frac{1}{4} + u^2$  and  $m = \frac{1}{2} + iv$ ,  $E_m = \frac{1}{4} + v^2$ , thus

$$\begin{aligned} \mathcal{D}_2(\beta, t) &= \int_0^{+\infty} \int_0^{+\infty} du dv e^{i(u^2 - v^2)t - \beta(\frac{1}{4} + u^2)} \\ &\int_{\mathcal{F}} \psi_{\frac{1}{2} - iu}(z) A \psi_{\frac{1}{2} + iv}(z) d\mu(z) \int_{\mathcal{F}} \psi_{\frac{1}{2} - iv}(w) B \psi_{\frac{1}{2} + iu}(w) d\mu(w), \end{aligned} \quad (6.2)$$

where the complex conjugate function is  $\psi_{\frac{1}{2} + iu}^*(z) = \psi_{\frac{1}{2} - iu}(z)$ . Defining the basic matrix element as

$$A_{uv} = \int_{\mathcal{F}} \psi_{\frac{1}{2} - iu}(z) A \psi_{\frac{1}{2} + iv}(z) d\mu(z) = \int_{-1/2}^{1/2} dx \int_{\sqrt{1-x^2}}^{\infty} \frac{dy}{y^2} \psi_{\frac{1}{2} - iu}(z) A \psi_{\frac{1}{2} + iv}(z) \quad (6.3)$$

for the two-point correlation function we shall get

$$\mathcal{D}_2(\beta, t) = \int_{-\infty}^{+\infty} e^{i(u^2 - v^2)t - \beta(\frac{1}{4} + u^2)} A_{uv} B_{vu} du dv. \quad (6.4)$$

In terms of the new variables (5.21) the basic matrix element (6.3) will take the form

$$A_{pq} = \int_{-1/2}^{1/2} dx \int_{\frac{1}{2} \log(1-x^2)}^{\infty} dy \psi_p^*(x, y) (e^{-\frac{1}{2}y} A e^{\frac{1}{2}y}) \psi_q(x, y). \quad (6.5)$$

The matrix element (6.3), (6.5) plays a fundamental role in the investigation of the correlation functions because all correlations can be expressed through it. One should choose also appropriate observables  $A$  and  $B$ . The operator  $y^{-2}$  seems very appropriate for two reasons. Firstly, the convergence of the integrals over the fundamental region  $\mathcal{F}$  will be well defined. Secondly, this operator is reminiscent of the exponentiated Liouville field since  $y^{-2} = e^{-2\bar{y}}$ . Thus we are interested in calculating the matrix element (6.5) for the observables in the form of the Liouville-like operators:

$$A(N) = e^{-2Ny}. \quad (6.6)$$

We shall get<sup>2</sup>

$$A_{pq}(N) = \int_{-1/2}^{1/2} dx \int_{\frac{1}{2} \log(1-x^2)}^{\infty} dy \psi_p^*(x, y) e^{-2Ny} \psi_q(x, y) , \quad N = 1, 2, \dots \quad (6.7)$$

Calculating the above matrix elements we shall use a perturbative expansion in which the part of the wave function (5.22) containing the Bessel's functions and the contribution of the discrete spectrum (5.27) will be considered as a perturbation. As we shall demonstrate below, these terms of the perturbative expansion are small and don't influence our results. The reason behind this fact is that in the integration region  $\Im z \gg 1, \Im w \gg 1$  of the matrix element (6.3) the Bessel's functions decay exponentially, as one can get convinced by inspecting (5.9). Therefore the contribution of these high modes is small (analogues to the so called mini-superspace approximation in the Liouville theory). Thus we shall consider the perturbative expansion over high frequency modes  $l = 1, 2, \dots$  in (5.22) and (5.27). In the first approximation of the wave function (5.22) for the matrix element we shall get

$$A_{pq}(N) = \int_{-1/2}^{1/2} dx \left( \frac{(1-x^2)^{-N+\frac{p-q}{2i}}}{2N+i(p-q)} + \frac{(1-x^2)^{-N+\frac{p+q}{2i}} e^{-i\varphi(q)}}{2N+i(p+q)} \right. \\ \left. + \frac{(1-x^2)^{-N-\frac{p+q}{2i}} e^{i\varphi(p)}}{2N-i(p+q)} + \frac{(1-x^2)^{-N-\frac{p-q}{2i}} e^{i(\varphi(p)-\varphi(q))}}{2N-i(p-q)} \right) , \quad (6.8)$$

where we used (5.23). Integration over  $x$  can be performed exactly with the result for the basic matrix element of the following form:

$$A_{pq}(N) = \frac{{}_2F_1\left(\frac{1}{2}, N+i\frac{p-q}{2}; \frac{3}{2}; \frac{1}{4}\right)}{2N+i(p-q)} + \frac{{}_2F_1\left(\frac{1}{2}, N+i\frac{p+q}{2}; \frac{3}{2}; \frac{1}{4}\right)}{2N+i(p+q)} e^{-i\varphi(q)} + \\ + \frac{{}_2F_1\left(\frac{1}{2}, N-i\frac{p+q}{2}; \frac{3}{2}; \frac{1}{4}\right)}{2N-i(p+q)} e^{i\varphi(p)} + \frac{{}_2F_1\left(\frac{1}{2}, N-i\frac{p-q}{2}; \frac{3}{2}; \frac{1}{4}\right)}{2N-i(p-q)} e^{i(\varphi(p)-\varphi(q))} , \quad (6.9)$$

where the reflection phase  $\varphi(p)$  was defined in (5.23). Thus for the two-point correlation function we shall get

$$\mathcal{D}_2(\beta, t) = \int_{-\infty}^{+\infty} e^{i(p^2-q^2)t-\beta(\frac{1}{4}+p^2)} A_{pq}(N) A_{qp}(M) dpdq . \quad (6.10)$$


The correlation function is between two Liouville-like fields in the power  $N$  and  $M$  respectively. This expression is very convenient for the analytical and numerical analyses. It is expected that the two-point correlation function should decay exponentially [36], therefore we fitted the points by the curve

$$\mathcal{D}_2(\beta, t) \sim K(\beta) e^{-\frac{t}{\tau_d(\beta)}} . \quad (6.11)$$

---

<sup>2</sup>The other interesting observable is  $A = \cos(2\pi Nx), N = 1, 2, \dots$





**Figure 3:** The exponential decay of the two-point correlation function  $\mathcal{D}_2(\beta, t)$  as a function of time at temperature  $\beta = 1$ . The points are fitted by the curve  $K(\beta) \exp(-t/t_d(\beta))$ . The exponent  $t_d(\beta)$  has a well defined high and low temperature limits. The limiting values in dimensionless units are  $t_d(0) \approx 0.276$  and  $t_d(\infty) \approx 0.749$ . The temperature dependence of  $K(\beta)$  is shown on the l.h.s. graph.

The  $t_d(\beta)$  is the decorrelation time and it defines one of the characteristic time scales in the quantum-mechanical system. The exponential decay of the two-point correlation function with time at different temperatures is shown on Fig.3<sup>3</sup>. The dependence of the exponent  $t_d(\beta)$  and of the prefactor  $K(\beta)$  as a function of temperature are presented in Fig.3. As one can see, at high and low temperatures the decorrelation time tends to the fixed values. These limiting values we calculated by using the expressions (6.12) and (6.13). At large  $\beta \rightarrow \infty$  the main contribution came from the zero momentum region  $p = 0$ :

$$\mathcal{D}_2(\infty, t) = \int_{-\infty}^{+\infty} e^{-iq^2 t} A_{0q}(N) A_{q0}(M) dq \sim K(\infty) e^{-\frac{t}{t_d(\infty)}}, \quad (6.12)$$

and at  $\beta \rightarrow 0$  from the region  $p = q$ :

$$\mathcal{D}_2(0, t) = \int_{-\infty}^{+\infty} e^{i(p^2 - q^2)t} A_{pq}(N) A_{qp}(M) dpdq \sim K(0) e^{-\frac{t}{t_d(0)}}. \quad (6.13)$$


The corresponding limiting values in dimensionless units are shown on the l.h.s. of the Fig.3.

## 6.2 Four-point correlation function

It was conjectured in the literature [36] that the classical chaos can be diagnosed in thermal quantum systems by using an out-of-time-order correlation functions as well as by the square of the commutator of the operators which are separated in time. The out-of-time four-point correlation function of interest was defined in [36] as follows:

$$\begin{aligned} \mathcal{D}_4(\beta, t) &= \langle A(t)B(0)A(t)B(0)e^{-\beta H} \rangle = \\ &= \sum_{n,m,l,r} e^{i(E_n - E_m + E_l - E_r)t - \beta E_n} \langle n|A(0)|m \rangle \langle m|B(0)|l \rangle \langle l|A(0)|r \rangle \langle r|B(0)|n \rangle. \end{aligned} \quad (6.14)$$

<sup>3</sup>We would like to thank Gabriel Poghosyan for numerical calculation of the two-point correlation function presented on Fig.3.



**Figure 4:** The exponential decay of the correlation function  $\mathcal{D}_4(\beta, t)$  as a function of time at  $\beta = 1$ . The rest of the functions  $\mathcal{D}'_4(\beta, t), \mathcal{D}''_4(\beta, t), \mathcal{D}'''_4(\beta, t)$  demonstrate a similar exponential decay  $\sim \exp(-\frac{t}{t_d(\beta)})$ . The temperature dependence of the exponent  $t_d(\beta)$  have a well defined high and low temperature limits and is shown on the r.h.s. graph. The corresponding limiting values of the function  $t_d(\beta)$  in dimensionless units are  $t_d(0) = 0,112$  and  $t_d(\infty) = 0,163$ . The behaviour of the exponent  $t_d(\beta)$  of the two-point correlation function is shown on the Fig.3.

The other important observable which we shall consider here is the square of the commutator of the Liouville-like operators separated in time [36]

$$C(\beta, t) = \langle [A(t), B(0)]^2 e^{-\beta H} \rangle . \quad (6.15)$$

The energy eigenvalues we shall parametrise as  $n = \frac{1}{2} + iu, m = \frac{1}{2} + iv, l = \frac{1}{2} + il$  and  $r = \frac{1}{2} + ir$ , thus from (6.14) we shall get

$$\mathcal{D}_4(\beta, t) = \int_{-\infty}^{+\infty} e^{i(u^2 - v^2 + l^2 - r^2)t - \beta(\frac{1}{4} + u^2)} A_{uv} B_{vl} A_{lr} B_{ru} du dv dl dr . \quad (6.16)$$

In terms of the variables (5.21) the four-point correlation function (6.16) will take the following form:


$$\mathcal{D}_4(\beta, t) = \int_{-\infty}^{+\infty} e^{i(p^2 - q^2 + l^2 - r^2)t - \beta(\frac{1}{4} + p^2)} A_{pq}(N) A_{ql}(M) A_{lr}(N) A_{rp}(M) dp dq dl dr . \quad (6.17)$$

As it was suggested in [36], the most important correlation function indicating the traces of the classical chaotic dynamics in quantum regime is (6.15)

$$C(\beta, t) = -\mathcal{D}_4(\beta, t) + \mathcal{D}'_4(\beta, t) + \mathcal{D}''_4(\beta, t) - \mathcal{D}'''_4(\beta, t), \quad (6.18)$$

where in the case of the Artin system we shall get (see Appendix for details)

$$\begin{aligned} & \mathcal{D}'_4(\beta, t) + \mathcal{D}''_4(\beta, t) = \\ & 2 \int_{-\infty}^{+\infty} e^{-\beta(\frac{1}{4} + p^2)} \cos(q^2 - r^2)t A_{pq}(N) A_{ql}(M) A_{lr}(N) A_{rp}(M) dp dq dl dr, \end{aligned} \quad (6.19)$$



**Figure 5:** Time evolution of the correlation function  $C(\beta, t)$  (6.22) at temperature  $\beta = 0.1$ . For the short time intervals the function  $C(\beta, t)$  exponentially increases with time. This behaviour is reminiscent to the exponential divergency of the classical trajectories in hyperbolic dynamical systems. As one can see, the exponent  $\chi(\beta)$  which defines the behaviour of the correlation function of the operators separated in time in the commutator (6.22), (6.24) slowly decreases with  $\beta$ . Such behaviour of the correlation function  $C(\beta, t)$  does not saturate the maximal growth future (6.23) which should be linear in  $\beta$  (6.25).

and

$$\mathcal{D}_4(\beta, t) + \mathcal{D}_4'''(\beta, t) = 2 \int_{-\infty}^{+\infty} e^{-\beta(\frac{1}{4}+p^2)} \cos(p^2 - q^2 + l^2 - r^2)t A_{pq}(N) A_{ql}(M) A_{lr}(N) A_{rp}(M) dpdqdl dr. \quad (6.20)$$

On Fig.4 one can see the behaviour of the four-point correlations as the function of the temperature and time. All four correlation functions decay exponentially:

$$\mathcal{D}_4(\beta, t) \sim K(\beta) e^{-\frac{t}{\tau_d(\beta)}}. \quad (6.21)$$

Now we can turn to the investigation of the commutator (6.15), (6.18) which can be represented in the following form:

$$\begin{aligned} C(\beta, t) &= 2 \int_{-\infty}^{+\infty} e^{-\beta(\frac{1}{4}+p^2)} \{ \cos(q^2 - r^2)t - \cos(p^2 - q^2 + l^2 - r^2)t \} \\ &\quad A_{pq}(N) A_{ql}(M) A_{lr}(N) A_{rp}(M) dpdqdl dr, \\ &= \int_{-\infty}^{+\infty} e^{-\beta(\frac{1}{4}+p^2)} \sin \frac{1}{2}(p^2 + l^2 - 2r^2)t \cdot \sin \frac{1}{2}(p^2 + l^2 - 2q^2)t \\ &\quad A_{pq}(N) A_{ql}(M) A_{lr}(N) A_{rp}(M) dpdqdl dr, \end{aligned} \quad (6.22)$$

where we used (6.19) and (6.20). It was conjectured in [36] that the influence of chaos on the commutator  $C(\beta, t)$  can develop no faster than exponentially:

$$C(\beta, t) \approx f(\beta) e^{\frac{2\pi}{\beta}t}, \quad (6.23)$$

with the exponent  $\frac{2\pi}{\beta}t = 2\pi Tt$  which is linear in temperature  $1/\beta = T$  and time  $t$ . Calculating the function  $C(\beta, t)$  one can check if it grows exponentially:

$$C(\beta, t) \sim f(\beta) e^{\frac{2\pi}{\chi(\beta)}t}, \quad (6.24)$$

and if the exponent  $\chi(\beta)$  is linear in  $\beta$ . The maximal growth will correspond to the exponent  $\chi(\beta)$  which grows linearly with  $\beta$  (6.23), that is,

$$\chi(\beta) \sim \beta. \quad (6.25)$$

The results of the integration are presented on the Fig.5. This beautifully confirms the fact that the correlation function  $C(\beta, t)$  indeed grows exponentially with time as it takes place in its classical counterpart. As one can see, the exponent  $\chi(\beta)$  defining the behaviour of the commutator  $C(\beta, t)$  in (6.22) and (6.24) slowly decreases with  $\beta$ . Such behaviour of the commutator  $C(\beta, t)$  does not saturate the maximal growth of the correlation function (6.25) which should be linear in  $\beta$ .

In order to check if the results are sensitive to the truncation of the high modes of the Maass wave function (5.17) we included their contribution into the integration of the basic matrix element  $A_{uv}$  in (6.3). We found that their influence on the behaviour of the correlation functions is negligible. The numerical values of the exponents are changing in the range of few percentage and do not influence the results. In summary, all two and four-point correlation functions decay exponentially. The commutator  $C(\beta, t)$  in (6.22) and (6.24) grows exponentially with exponent which is almost constant Fig.5. This behaviour does not saturate the condition of the maximal growth (6.23), (6.25).

## 7 Acknowledgment

R.P. is thankful to the NCSR”Demokritos” for hospitality during June-July 2018 when this work was completed. H.B. would like to thank NCSR”Demokritos” for kind hospitality and generous financial support of the MIXMAX project. G.S. would like to thank Luis Alvarez-Gaume for stimulating discussions of  $SL(2, Z)$  automorphic functions, the references [28, 31] and kind hospitality at the Simon Center for Geometry and Physics where part of this work was completed. This project received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 644121.

## 8 Appendix

In order to compute the square of the commutator (6.22) one should consider the following four-point correlation functions

$$\begin{aligned} \mathcal{D}'_4(\beta, t) &= \langle A(t)B(0)B(0)A(t)e^{-\beta H} \rangle = & (8.1) \\ &= \sum_{n,m,l,r} e^{i(-E_m+E_r)t-\beta E_n} \langle n|A(0)|m \rangle \langle m|B(0)|l \rangle \langle l|B(0)|r \rangle \langle r|A(0)|n \rangle \end{aligned}$$

$$\begin{aligned} \mathcal{D}''_4(\beta, t) &= \langle B(0)A(t)A(t)B(0)e^{-\beta H} \rangle = & (8.2) \\ &= \sum_{n,m,l,r} e^{i(E_m-E_r)t-\beta E_n} \langle n|B(0)|m \rangle \langle m|A(0)|l \rangle \langle l|A(0)|r \rangle \langle r|B(0)|n \rangle \end{aligned}$$

$$\begin{aligned} \mathcal{D}'''_4(\beta, t) &= \langle B(0)A(t)B(0)A(t)e^{-\beta H} \rangle = & (8.3) \\ &= \sum_{n,m,l,r} e^{i(-E_n+E_m-E_l+E_r)t-\beta E_n} \langle n|B(0)|m \rangle \langle m|A(0)|l \rangle \langle l|B(0)|r \rangle \langle r|A(0)|n \rangle . \end{aligned}$$

These correlation functions in the momentum space representation are given in (6.19) and (6.20).

## References

- [1] Emil Artin, *Ein mechanisches system mit quasiergodischen bahnen*, E. Abh. Math. Semin. Univ. Hambg. (1924) 3: 170.
- [2] Henri Poincaré *Théorie des Groupes Fuchsiennes*, Acta Mathematica, **1** (1882) 1-62.
- [3] Henri Poincaré *Mémoire sur les Fonctions Fuchsiennes*, Acta Mathematica, **1** (1882) 193-294.
- [4] Lazarus Fuchs, *Ueber eine Klasse von Funktionen mehrerer Variablen, welche durch Umkehrung der Integrale von Lösungen der linearen Differentialgleichungen mit rationalen Coeffizienten entstehen*, J. Reine Angew. Math., **89** (1880) 151-169
- [5] J. Hadamard, *Les surfaces á courbures opposées et leur lignes geodesiques*, Liouville, Journal de Mathématique, **4** (1898) 27
- [6] G.Hedlund, *The dynamics of geodesic flow*, Bull.Am.Math.Soc. **45** (1939) 241-246
- [7] D. V. Anosov, *Geodesic flows on closed Riemannian manifolds with negative curvature*, Trudy Mat. Inst. Steklov., Vol. **90** (1967) 3 - 210.
- [8] R.Bowen, *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms*. (Lecture Notes in Mathematics, no. 470: A. Dold and B. Eckmann, editors). Springer-Verlag (Heidelberg, 1975), 108 pp.

- [9] A.N. Kolmogorov, *New metrical invariant of transitive dynamical systems and automorphisms of Lebesgue spaces*, Dokl. Acad. Nauk SSSR, **119** (1958) 861-865
- [10] A.N. Kolmogorov, *On the entropy per unit time as a metrical invariant of automorphism*, Dokl. Acad. Nauk SSSR, **124** (1959) 754-755
- [11] Ya.G. Sinai, *On the Notion of Entropy of a Dynamical System*, Doklady of Russian Academy of Sciences, **124** (1959) 768-771.
- [12] D. Ruelle, *Thermodynamic Formalism*, Addison-Wesley, Reading, Mass., 1978
- [13] E.Hopf. *Statistik der Lösungen geodätischer Probleme vom unstabilen Typus. II.* Math. Ann. **117** (1940) 590-608
- [14] E. Hopf, *Ergodic theory and the geodesic flow on surfaces of constant negative curvature*, Bull. Amer. Math. Soc, **77** (1971) 863- 877.
- [15] I.M.Gelfand and S.V. Fomin , *Geodesic flows on manifolds of constant negative curvature*, Uspekhi Mat. Nauk, **7** (1952) 118-137; Amer. Math. Soc.Translation **1** (1965) 49-65.
- [16] P. Collet, H. Epstein and G. Gallavotti, *Perturbations of Geodesic Flows on Surfaces of Constant Negative Curvature and Their Mixing Properties*, Commun. Math. Phys. **95** (1984) 61-112
- [17] M. Pollicott, *On the rate of mixing of Axiom A flows*, Invent. math. **81** (1985) 413-426.
- [18] C. C. Moore, *Exponential decay of correlation coefficients for geodesic flows*, Group representations, ergodic theory, operator algebras, and mathematical physics (Berkeley, Calif., 1984), Math. Sci. Res. Inst. Publ., vol. 6, Springer, New York, 1987, pp. 163 - 181.
- [19] D. Dolgopyat, *On Decay of Correlations in Anosov Flows*, Annals of Mathematics Second Series, **147** (1998) 357-390
- [20] N. I. Chernov, *Markov Approximations and Decay of Correlations for Anosov Flows*, Annals of Mathematics Second Series. **147** (1998) 269-324
- [21] L. D. Faddeev, *Feynman integral for singular Lagrangians*, Theor. Math. Phys. **1** (1969) 1 [Teor. Mat. Fiz. **1** (1969) 3]. doi:10.1007/BF01028566
- [22] G.Savvidy, *The Yang-Mills mechanics as a Kolmogorov K-system*, Phys.Lett.B **130** (1983) 303
- [23] G. Savvidy, *Classical and Quantum Mechanics of non-Abelian Gauge Fields*, Nucl. Phys. B **246** (1984) 302.
- [24] H. Maass, *Über eine neue Art von nichtanalytischen automorphen Funktionen*, Math. Ann. **121**, No 2 (1949), 141-183.
- [25] W. Roelcke, *Über die Wellengleichung bei Grenzkreisgruppen erster Art*, Sitzungsber. Heidelberg. Acad. Wiss. **4** Abh. (1953/1956), 161-267.

- [26] A.Selberg, *Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series*, Indian Journ. Math. Soc. **20** (1956) 47-87 .
- [27] A.Selberg, *Discontinuous groups and harmonic analysis*, Proceedings of Stockholm Mathematical Congress (1962).
- [28] D. Bump, *Automorphic Forms and Representations*, Cambridge Studies in Advance Mathematics: 55, Cambridge University Press, 1998.
- [29] L.D.Faddeev, *Expansion in eigenfunctions of the Laplace operator on the fundamental domain of a discrete group on the Lobachevsky plane*, Trans. Moscow Math. Soc., **17** (1967) 357-386.
- [30] L.D.Faddeev, A. B. Venkov and V. L. Kalinin *A non-arithmetic derivation of the Selberg trace formula*, J. Soviet Math., **8** 2 (1977) 171-199.
- [31] D.A.Hejhal, *The Selberg Trace Formula for  $PSL(2, R)$* , Lecture Notes in Mathematics 548, Springer-Verlag **Vol. 1** 1976.
- [32] G. Savvidy, *Anosov C-systems and random number generators*, Theor. Math. Phys. **188** (2016) 1155; doi:10.1134/S004057791608002X [arXiv:1507.06348 [hep-th]].
- [33] G. Savvidy and N. Ter-Arutyunyan-Savvidy, *On the Monte Carlo simulation of physical systems*, J.Comput.Phys. **97** (1991) 566; Preprint EFI-865-16-86-YEREVAN, Jan. 1986. 13pp.
- [34] K.Savvidy, *The MIXMAX random number generator*, Comput.Phys.Commun. 196 (2015) 161-165. (<http://dx.doi.org/10.1016/j.cpc.2015.06.003>); arXiv:1404.5355
- [35] K. Savvidy and G. Savvidy, *Spectrum and Entropy of C-systems. MIXMAX random number generator*, Chaos Solitons Fractals **91** (2016) 33 doi:10.1016/j.chaos.2016.05.003 [arXiv:1510.06274 [math.DS]].
- [36] J. Maldacena, S. H. Shenker and D. Stanford, *A bound on chaos*, JHEP **1608** (2016) 106 doi:10.1007/JHEP08(2016)106 [arXiv:1503.01409 [hep-th]].
- [37] G. Gur-Ari, M. Hanada and S. H. Shenker, *Chaos in Classical D0-Brane Mechanics*, JHEP **1602** (2016) 091 doi:10.1007/JHEP02(2016)091 [arXiv:1512.00019 [hep-th]].
- [38] G. Savvidy and K. Savvidy, *Exponential decay of correlations functions in MIXMAX generator of pseudorandom numbers*, Chaos Solitons and Fractals, **107** (2018) 244. <https://doi.org/10.1016/j.chaos.2018.01.007>
- [39] H. Poghosyan, H. Babujian and G. Savvidy, *Artin Billiard Exponential Decay of Correlation Functions*, arXiv:1802.04543 [nlin.CD].
- [40] M. Hanada, H. Shimada and M. Tezuka, *Universality in Chaos: Lyapunov Spectrum and Random Matrix Theory*, Phys. Rev. E **97** (2018) no.2, 022224 doi:10.1103/PhysRevE.97.022224 [arXiv:1702.06935 [hep-th]].
- [41] T. Anous and C. Cogburn, *Mini-BFSS in Silico*, arXiv:1701.07511 [hep-th].

- [42] A.Winkler, *Cusp forms and Hecke groups*, J. Reine Angew. Math. **386** (1988) 187
- [43] D.A.Hejhal, *Eigenvalues of the Laplacian for  $PSL(2,Z)$  : some new results and computational techniques*, in International Symposium in Memory of Hua Loo-Keng (ed. by Gong, Lu, Wang, Yang), Science Press and Springer-Verlag **1** (1991) 59-102.
- [44] D.A.Hejhal and B.Berg, *Some new results concerning eigenvalues of the non-Euclidean Laplacian for  $PSL(2,Z)$* , Univ. of Minn. Math. Report **No. 82-172** (1982) 7pp.