

The system of partial differential equations for the C_0 function

Tai-Fu Feng^{*a}, Chao-Hsi Chang^{†b,c,d}, Jian-Bin Chen^{e‡}, Hai-Bin Zhang^{a§}

^a*Department of Physics, Hebei University, Baoding, 071002, China*

^b*Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,
Chinese Academy of Science, Beijing, 100190, China*

^c*CCAST (World Laboratory), P.O.Box 8730, Beijing, 100190, China*

^d*School of Physical Sciences, University of Chinese
Academy of Sciences, Beijing 100049, China and*

^e*Department of Physics, Taiyuan University of Technology, Taiyuan, 030024, China*

Abstract

We present an approach to analyze the scalar integrals of any Feynman diagrams in detail here. This method not only completely recovers some well-known results in the literature, but also produces some new results on the C_0 function. The approach can be employed to evaluate the coefficient of arbitrary power of ε in the expansion of a scalar integral, where $D = 4 - 2\varepsilon$ denotes the time-space dimension.

PACS numbers: 02.30.Jr, 11.10.Gh, 12.38.Bx

Keywords: Scalar integral, Mellin-Barnes transformation, Linear partial differential equation

* email:fengt@hbu.edu.cn

† email:zhangzx@itp.ac.cn

‡ email:chenjianbin@tyut.edu.cn

§ email:hbzhang@hbu.edu.cn

I. INTRODUCTION

The discovery of the Higgs particle in the Large Hadron Collider (LHC) implies the great success of the standard model (SM)[1, 2]. With the increasing of luminosity of the collider, one of the targets for particle physics now is to test the SM precisely and to search for new physics (NP) beyond the SM[3–5].

How to evaluate the scalar integrals exactly is an open problem to predict the electroweak observables precisely in the SM. The author of literature [6] presents several methods to evaluate those scalar integrals. Nevertheless each method mentioned in this literature has its blemishes since it can only be applied to the Feynman diagrams with special topology and kinematic invariants.

In dimensional regularization, any scalar integral can be expanded around $\varepsilon = 0$, here the time-space dimension $D = 4 - 2\varepsilon$. For example, the one-loop three-point function C_0 is expanded as

$$C_0 = C_0^{(0)} + \sum_{i=1}^{\infty} C_0^{(i)} \varepsilon^i, \quad (1)$$

around $D = 4$, and the well-known result of $C_0^{(0)}$ is presented in the literature[7]. However, evaluating $C_0^{(i)}$ precisely is also necessary to obtain high order radiative corrections to the amplitude self-consistently when the virtual interactions originate from the counter terms[8]. In addition, the vector and tensor coefficients C_i, C_{00}, C_{ij} , ($i, j = 1, 2$) adopted in the paper [9] can be expressed by the one-point function A_0 , the two-point function B_0 , and the three-point function C_0 , respectively.

There are many analytical results on the scalar integral C_0 already in the references. In Ref. [10] the massless C_0 function is presented as the linear combination of the fourth kind Appell function F_4 whose arguments are the dimensionless ratios among the external momenta squared, and is simplified further as the linear combination of the Gauss function ${}_2F_1$ through the quadratic transformation [11] in the literature [12]. With some special assumptions on the virtual masses, the analytic expressions of the scalar integral C_0 are given by the multiple hypergeometric functions in Ref. [13] through the corresponding Mellin-Barnes representations. Taking the massless C_0 function as an example, the author of Ref [14] presents an algorithm to evaluate the scalar integrals of one-loop vertex-type Feynman diagrams. Certainly, some analytic results of the C_0 function can also be extracted from the

expressions for the scalar integrals of one-loop massive N -point Feynman diagrams [15, 16]. In addition, the literature [17] also provides a geometrical interpretation of the analytic expressions of the scalar integrals from one-loop N -point Feynman diagrams. Using the recurrence relations respecting the time-space dimension, the papers [18, 19] formulate one-loop two-point function B_0 as the linear combination of the Gauss function ${}_2F_1$, one-loop three-point function C_0 with arbitrary external momenta and virtual masses as the linear combination of the Appell function F_1 with two arguments, and one-loop four-point function D_0 with arbitrary external momenta and virtual masses as the linear combination of the Lauricella-Saran function F_s with three arguments, respectively. The expression for the scalar integral C_0 is convenient for analytic continuation and numerical evaluation because continuation of the Appell functions has been analyzed thoroughly. Nevertheless, how to perform continuation of the Lauricella-Saran function F_s outside its convergent domain is still a challenge. Basing on the hypergeometric system of linear partial differential equations (PDEs), we present an approach to evaluate the scalar integral of arbitrary multiloop Feynman diagram systematically. Actually the system of PDEs satisfied by the corresponding scalar integral is a holonomic integrable system which can be transformed into the Pfaffian system of PDEs [20] by the algebra decomposition. Then one can obtain the singularities of the holonomic hypergeometric system and the number of independent solutions in certain parameter space. Together with the analytic expressions in some convergent regions, the holonomic hypergeometric system provides a new way to understand the corresponding scalar integral. Taking the C_0 function as an example, we elucidate how to evaluate the coefficient of arbitrary power of ε in the expansion of a scalar integral around $D = 4$. In fact, the corresponding analytic results on the C_0 function coincide with those well-known results mentioned above. The evaluations of scalar integrals of any Feynman diagrams such as that presented in Refs. [21–23] are given elsewhere.

A holonomic hypergeometric system of linear PDEs is given through the corresponding Mellin-Barnes representation of the concerned scalar integral [24], where the system of linear PDEs is satisfied by the scalar integral in the whole parameter space of the independent variables. The scalar integral is written as the multiple hypergeometric functions [25] of the independent variables for some isolated singularities of the integrand by residue theorem [26]. In addition, the Horn's study of convergence[27] predicts the absolutely and

uniformly convergent regions of those derived hypergeometric functions exactly. Several convergent regions of the hypergeometric functions compose a set in which each convergent region does not intersect with the others. Additionally each convergent region of the set either does not intersect with, or is a proper subset of other convergent regions which do not belong to the set. Within each convergent region in the set, the scalar integral can be written as the sum of those hypergeometric functions whose convergent regions contain the concerned element entirely.

In view of mathematics, the most important point for a given multiple power series is how to get its absolutely and uniformly convergent region, and how to continue it to the whole parameter space. Fortunately, the continuation can be achieved by the system of linear PDEs mentioned above. As stated in our previous work[28], an effective Hamiltonian can be constructed if those linear PDEs are the stationary conditions of the corresponding system. Then we can numerically continue the scalar integral to the entire parameter space with the finite element method.

Our presentation is organized as follows. We briefly mention some typical results of the massive C_0 function in section II at first. Then we present in detail our analyses on the massless C_0 function in section III, the C_0 function with one nonzero virtual mass in section IV, and the C_0 function with three equally virtual masses in section V, respectively. In section VI, we recognize the system of linear PDEs as the stationary conditions of a functional under some restrictions according to Hamilton's principle, which is convenient to numerically continue the scalar integral to the whole parameter space with the finite element method. The conclusions are summarized in section VII and some tedious formulae are presented in the appendixes.

II. SOME RESULTS OF THE C_0 FUNCTION IN GENERAL CASE

The Mellin-Barnes representation of massive C_0 function is generally written as

$$\begin{aligned}
& C_0(p_1^2, p_2^2, p_3^2, m_1^2, m_2^2, m_3^2) \\
&= \int \frac{d^D q}{(2\pi)^D} \frac{1}{(q^2 - m_3^2)((q + p_1)^2 - m_2^2)((q - p_2)^2 - m_1^2)} \\
&= -\frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}(2\pi i)^5} \int_{-i\infty}^{+i\infty} ds_1 \left(-\frac{m_1^2}{p_2^2}\right)^{s_1} \Gamma(-s_1) \int_{-i\infty}^{+i\infty} ds_2 \left(-\frac{m_2^2}{p_3^2}\right)^{s_2} \Gamma(-s_2)
\end{aligned}$$

$$\begin{aligned}
& \times \int_{-i\infty}^{+i\infty} ds_3 \left(-\frac{m_3^2}{p_3^2} \right)^{s_3} \Gamma(-s_3) \int_{-i\infty}^{+i\infty} dz_1 \int_{-i\infty}^{+i\infty} dz_2 \Gamma(-z_1) \Gamma(-z_2) \\
& \times \Gamma\left(3 - \frac{D}{2} + s_1 + s_2 + s_3 + z_1 + z_2\right) \left(\frac{p_1^2}{p_3^2}\right)^{z_1} \left(\frac{p_2^2}{p_3^2}\right)^{z_2} \\
& \times \frac{\Gamma\left(\frac{D}{2} - 2 - s_1 - s_3 - z_2\right) \Gamma\left(\frac{D}{2} - 2 - s_2 - s_3 - z_1\right) \Gamma(1 + s_3 + z_1 + z_2)}{\Gamma(D - 3 - s_1 - s_2 - s_3)} \\
& = \frac{i(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} F_{p_3}(\xi_{13}, \xi_{23}, \xi_{33}, x_{13}, x_{23}) \\
& = \frac{i(m_3^2)^{D/2-3}}{(4\pi)^{D/2}} F_{m_3}(\eta_{13}, \eta_{23}, \eta_{33}, y_{13}, y_{23}) \tag{2}
\end{aligned}$$

with $p_3^2 = (p_1 + p_2)^2$, $\xi_{ij} = -m_i^2/p_j^2$, $\eta_{ij} = -p_i^2/m_j^2 = 1/\xi_{ji}$, $x_{ij} = p_i^2/p_j^2$, and $y_{ij} = m_i^2/m_j^2$ ($i, j = 1, 2, 3$), respectively. Here the dimensionless functions F_{p_3} , $\xi_{33}^{D/2-3} F_{m_3}$ comply with the holonomic hypergeometric system of linear PDEs

$$\begin{aligned}
& \left\{ \left[3 - \frac{D}{2} + \sum_{i=1}^3 \hat{\vartheta}_{\xi_{i3}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} \right] \left[1 + \hat{\vartheta}_{\xi_{33}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} \right] \left[4 - D + \sum_{i=1}^3 \hat{\vartheta}_{\xi_{i3}} \right] \right. \\
& \left. - \frac{1}{\xi_{33}} \hat{\vartheta}_{\xi_{33}} \prod_{i=1}^3 \left[2 - \frac{D}{2} + \hat{\vartheta}_{\xi_{i3}} + \hat{\vartheta}_{\xi_{33}} + \hat{\vartheta}_{x_{(3-i)3}} \right] \right\} F_{p_3} = 0, \\
& \left\{ \left[3 - \frac{D}{2} + \sum_{i=1}^3 \hat{\vartheta}_{\xi_{i3}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} \right] \left[4 - D + \sum_{i=1}^3 \hat{\vartheta}_{\xi_{i3}} \right] \right. \\
& \left. + \frac{1}{\xi_{j3}} \hat{\vartheta}_{\xi_{j3}} \left[2 - \frac{D}{2} + \hat{\vartheta}_{\xi_{33}} + \hat{\vartheta}_{\xi_{j3}} + \hat{\vartheta}_{x_{(3-j)3}} \right] \right\} F_{p_3} = 0, \\
& \left\{ \left[3 - \frac{D}{2} + \sum_{i=1}^3 \hat{\vartheta}_{\xi_{i3}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} \right] \left[1 + \hat{\vartheta}_{\xi_{33}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} \right] \right. \\
& \left. - \frac{1}{x_{j3}} \hat{\vartheta}_{x_{j3}} \left[2 - \frac{D}{2} + \hat{\vartheta}_{\xi_{33}} + \hat{\vartheta}_{\xi_{(3-j)3}} + \hat{\vartheta}_{x_{j3}} \right] \right\} F_{p_3} = 0, \quad (j = 1, 2), \tag{3}
\end{aligned}$$

with the Euler operator $\hat{\vartheta}_x = x\partial/\partial x$ etc. Or equivalently, the functions F_{m_3} , $\eta_{33}^{D/2-3} F_{p_3}$ satisfy the following holonomic system of linear PDEs

$$\begin{aligned}
& \left\{ \left[3 - \frac{D}{2} + \sum_{i=1}^2 \hat{\vartheta}_{y_{i3}} + \sum_{i=1}^3 \hat{\vartheta}_{\eta_{i3}} \right] \prod_{i=1}^2 \left[1 + \hat{\vartheta}_{\eta_{i3}} + \hat{\vartheta}_{\eta_{33}} + \hat{\vartheta}_{y_{(3-i)3}} \right] \right. \\
& \left. - \frac{1}{\eta_{33}} \hat{\vartheta}_{\eta_{33}} \left[\frac{D}{2} - 1 + \sum_{i=1}^3 \hat{\vartheta}_{\eta_{i3}} \right] \left[2 - \frac{D}{2} + \sum_{i=1}^2 \hat{\vartheta}_{y_{i3}} + \hat{\vartheta}_{\eta_{33}} \right] \right\} F_{m_3} = 0, \\
& \left\{ \left[3 - \frac{D}{2} + \sum_{i=1}^2 \hat{\vartheta}_{y_{i3}} + \sum_{i=1}^3 \hat{\vartheta}_{\eta_{i3}} \right] \left[1 + \hat{\vartheta}_{\eta_{j3}} + \hat{\vartheta}_{\eta_{33}} + \hat{\vartheta}_{y_{(3-j)3}} \right] \right. \\
& \left. + \frac{1}{\eta_{j3}} \hat{\vartheta}_{\eta_{j3}} \left[\frac{D}{2} - 1 + \sum_{i=1}^3 \hat{\vartheta}_{\eta_{i3}} \right] \right\} F_{m_3} = 0,
\end{aligned}$$

$$\begin{aligned} & \left\{ \left[3 - \frac{D}{2} + \sum_{i=1}^2 \hat{\vartheta}_{y_{i3}} + \sum_{i=1}^3 \hat{\vartheta}_{\eta_{i3}} \right] \left[1 + \hat{\vartheta}_{\eta_{(3-j)3}} + \hat{\vartheta}_{\eta_{33}} + \hat{\vartheta}_{y_{j3}} \right] \right. \\ & \left. - \frac{1}{y_{j3}} \hat{\vartheta}_{y_{j3}} \left[2 - \frac{D}{2} + \sum_{i=1}^2 \hat{\vartheta}_{y_{i3}} + \hat{\vartheta}_{\eta_{33}} \right] \right\} F_{m_3} = 0, \quad (j = 1, 2). \end{aligned} \quad (4)$$

Here we emphasize that the two systems in Eq.(3) and Eq.(4) are equivalent. Actually the Mellin-Barnes representation presented here is equivalent to that of Eq. (3.4) in Ref. [15]. In addition, the author of Ref. [15] also presents a single hypergeometric function originating from the Mellin-Barnes representation in Eq. (3.6). With our notation, the corresponding hypergeometric function is rewritten as

$$\begin{aligned} C_{m_3}(p_1^2, p_2^2, p_3^2, m_1^2, m_2^2, m_3^2) &= \frac{i(m_3^2)^{D/2-3}}{(4\pi)^{D/2}} \sum_{j_1=0}^{\infty} \sum_{j_2=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \sum_{n_3=0}^{\infty} \\ &\times \frac{(-)^{n_1+n_2+n_3}}{j_1! j_2! n_1! n_2! n_3!} \left[1 - y_{13} \right]^{j_1} \left[1 - y_{23} \right]^{j_2} \eta_{13}^{n_1} \eta_{23}^{n_2} \eta_{33}^{n_3} \\ &\times \frac{\Gamma(3 - \frac{D}{2} + j_1 + j_2 + \sum_{i=1}^3 n_i) \Gamma(1 + j_1 + n_2 + n_3)}{\Gamma(3 + j_1 + j_2 + 2 \sum_{i=1}^3 n_i)} \\ &\times \Gamma(1 + j_2 + n_1 + n_3) \Gamma(1 + n_1 + n_2), \end{aligned} \quad (5)$$

which obviously satisfies the holonomic hypergeometric systems in Eq.(3) and Eq.(4). Using the adjacent ratios of the coefficients, we define

$$\begin{aligned} \Phi_{j_k}(j_1, j_2, n_1, n_2, n_3) &= \frac{(j_1 + j_2 + n_1 + n_2 + n_3)(j_k + n_{3-k} + n_3)}{j_k(j_1 + j_2 + 2n_1 + 2n_2 + 2n_3)}, \\ \Phi_{n_k}(j_1, j_2, n_1, n_2, n_3) &= \frac{(j_1 + j_2 + n_1 + n_2 + n_3)(j_{3-k} + n_k + n_3)(n_1 + n_2)}{n_k(j_1 + j_2 + 2n_1 + 2n_2 + 2n_3)^2}, \\ \Phi_{n_3}(j_1, j_2, n_1, n_2, n_3) &= \frac{(j_1 + j_2 + n_1 + n_2 + n_3) \prod_{i=1}^2 (j_{3-i} + n_i + n_3)}{n_3(j_1 + j_2 + 2n_1 + 2n_2 + 2n_3)^2}, \quad (k = 1, 2). \end{aligned} \quad (6)$$

The Cartesian hypersurface of the multiple hypergeometric series in Eq.(5) is written as

$$\begin{aligned} & r_1^2 r_3 - r_1 r_2 r_3 + r_3^2 - r_1 r_3^2 - r_1 r_2 r_4 + r_2^2 r_4 - 2r_3 r_4 + r_1 r_3 r_4 + r_2 r_3 r_4 \\ & + r_4^2 - r_2 r_4^2 + r_1 r_2 r_5 - 2r_3 r_5 + r_1 r_3 r_5 - 2r_4 r_5 + r_2 r_4 r_5 + r_3 r_4 r_5 + r_5^2 = 0. \end{aligned} \quad (7)$$

with $r_i = |y_{i3} - 1|$, $r_{2+i} = |\eta_{i3}|$, ($i = 1, 2$), $r_5 = |\eta_{33}|$, respectively. Correspondingly the absolutely and uniformly convergent domain of the series is

$$\Omega_1^{',a} = \left\{ (\eta_{13}, \eta_{23}, \eta_{33}, y_{13}, y_{23}) : \begin{array}{l} |\eta_{13}| < 4, |\eta_{23}| < 4, |\eta_{33}| < 8, \\ |y_{13} - 1| < 1, |y_{23} - 1| < 1 \end{array} \right\}. \quad (8)$$

Through the recurrence relations with respect to the time-space dimension [18, 19], the scalar integral C_0 can also be presented by the linear combination of the first kind Appell hypergeometric function F_1 . Where the independent variables of the hypergeometric functions are dimensionless ratios among the parameters r_{12} , r_{23} , r_{13} , r_{123} , and m_i , ($i = 1, 2, 3$), and the linear combination coefficients depend on the parameters λ_{12} , λ_{23} , λ_{13} , and g_{123} . With our notation above, those parameters are expressed as

$$\begin{aligned}
\lambda_{12} &= \kappa^2(m_1^2, m_2^2, p_3^2), & \lambda_{13} &= \kappa^2(m_1^2, p_2^2, m_3^2), \\
\lambda_{23} &= \kappa^2(p_1^2, m_2^2, m_3^2), & g_{123} &= -2\kappa^2(p_1^2, p_2^2, p_3^2), \\
\lambda_{123} &= 8m_1^2m_2^2m_3^2 - 2m_1^2(m_2^2 + m_3^2 - p_1^2)^2 \\
&\quad - 2m_2^2(m_1^2 + m_3^2 - p_2^2)^2 - 2m_3^2(m_1^2 + m_2^2 - p_3^2)^2 \\
&\quad + 2(m_1^2 + m_2^2 - p_3^2)(m_1^2 + m_3^2 - p_2^2)(m_2^2 + m_3^2 - p_1^2), \\
r_{12} &= -\frac{\lambda_{12}}{4p_3^2}, & r_{13} &= -\frac{\lambda_{13}}{4p_2^2}, & r_{23} &= -\frac{\lambda_{23}}{4p_1^2}, & r_{123} &= \frac{\lambda_{123}}{g_{123}},
\end{aligned} \tag{9}$$

with the Källèn function squared

$$\kappa^2(x, y, z) = 2xy + 2yz + 2zx - x^2 - y^2 - z^2. \tag{10}$$

Using the preparation above, one can check that the analytic expression in Eq. (74) of Ref. [19] complies with the holonomic hypergeometric systems in Eq.(3).

In order to apply the residue theorem, we should choose five Γ functions in the numerator of the integrand to determine the poles because the Mellin-Barnes representation in Eq. (2) is a five-fold contour integral. There are $C_9^5 = 126$ different choices in all. When the poles are determined by some specific choices of Γ functions, for example, $\Gamma(-s_3)$, $\Gamma(-z_1)$, $\Gamma(-z_2)$, $\Gamma(1 + s_3 + z_1 + z_2)$, and $\Gamma(3 - \frac{D}{2} + s_1 + s_2 + s_3 + z_1 + z_2)$, the Jacobian of the transformation

$$\begin{aligned}
-s'_3 &= -s_3, \\
-s'_1 &= 3 - \frac{D}{2} + s_1 + s_2 + s_3 + z_1 + z_2, \\
-s'_2 &= 1 + s_3 + z_1 + z_2, \\
-z'_1 &= -z_1, \\
-z'_2 &= -z_2
\end{aligned} \tag{11}$$

is zero. Correspondingly this special choice of poles in the numerator of the integrand does not induce correction to the Feynman integral at all [26]. For other choices of poles that the

Jacobian of the corresponding transformation is not zero, the Feynman integral is written as a quintuple hypergeometric function.

In the hypergeometric function mentioned above, the coefficient of the power function of independent variables is a fraction, whose numerator and denominator are both products of several Γ functions. Where the independent variable of each Γ function is a linear combination of the time-space dimension and the summation indices, and the coefficients of the summation indices are either positive or negative integers. Because of this reason, the hypergeometric function is decomposed into the sum of several hypergeometric series, so that the independent variables of all Γ functions of each 'new' hypergeometric series have definite signs at $D = 4$. After the transformation of summation indices, we obtain the absolutely and uniformly convergent region of each 'new' hypergeometric series with Horn's study of convergence [27]. The intersection of all convergent regions is the absolutely and uniformly convergent region of the original hypergeometric function.

Some convergent regions of the quintuple hypergeometric functions compose a set, and each convergent region does not intersect with the others in the set. Additionally each convergent region in the set either does not intersect with, or is a proper subset of the other convergent region which does not belong to the set. In each convergent region of the set, the Feynman integral can be written as the sum of those hypergeometric functions whose convergent regions contain the concerned region of the set entirely.

In order to shorten the length of text, we apply the approach to the Feynman integrals of the C_0 function with some special assumptions on the virtual masses.

III. THE SCALAR INTEGRAL OF MASSLESS C_0 FUNCTION

Generally the Mellin-Barnes representation of massless C_0 function is simplified as

$$\begin{aligned}
C_0(p_1^2, p_2^2, p_3^2) &= \int \frac{d^D q}{(2\pi)^D} \frac{1}{q^2(q+p_1)^2(q-p_2)^2} \\
&= \frac{i(-)^{D/2}}{(4\pi)^{D/2}\Gamma(D-3)(2\pi i)^2} \int_{-i\infty}^{+i\infty} dz_1 \int_{-i\infty}^{+i\infty} dz_2 (p_1^2)^{z_1} (p_2^2)^{z_2} \\
&\quad \times (p_3^2)^{D/2-3-z_1-z_2} \Gamma(-z_1)\Gamma(-z_2)\Gamma(3-\frac{D}{2}+z_1+z_2) \\
&\quad \times \Gamma(1+z_1+z_2)\Gamma(\frac{D}{2}-2-z_1)\Gamma(\frac{D}{2}-2-z_2), \tag{12}
\end{aligned}$$

which coincides with Eq. (24) of Ref. [13] (i.e. Eq. (2.5) of Ref. [14]). The analytic expression of the massless C_0 function is obtained by summing over residues of the integrand, and the corresponding results are enumerated as following.

- 1(a): After summing over the residues of $\Gamma(-z_1)$ and $\Gamma(-z_2)$, we obtain the double hypergeometric function as

$$\begin{aligned}
C_{0,1a}(p_1^2, p_2^2, p_3^2) &= \frac{i(-)^{D/2}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}\Gamma(D-3)} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \frac{(-)^{n_1+n_2}}{n_1!n_2!} \left(\frac{p_1^2}{p_3^2}\right)^{n_1} \left(\frac{p_2^2}{p_3^2}\right)^{n_2} \\
&\quad \times \Gamma(1+n_1+n_2)\Gamma(3-\frac{D}{2}+n_1+n_2)\Gamma(\frac{D}{2}-2-n_1) \\
&\quad \times \Gamma(\frac{D}{2}-2-n_2) \\
&= \frac{i(-)^{D/2}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma^2(2-\frac{D}{2})\Gamma^2(\frac{D}{2}-1)}{\Gamma(D-3)} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \\
&\quad \times A_{n_1, n_2}^{(1a)} x_{13}^{n_1} x_{23}^{n_2} .
\end{aligned} \tag{13}$$

The necessary condition $|p_3^2| > \max(|p_1^2|, |p_2^2|)$ (i.e. $|x_{i3}| < 1$) should be satisfied to guarantee the uniform absolute-convergence of the double series. Here the adjacent ratios of the coefficients are

$$\begin{aligned}
\Phi'_1(n_1, n_2) &= \frac{A_{1+n_1, n_2}^{(1a)}}{A_{n_1, n_2}^{(1a)}} = \frac{(1+n_1+n_2)(3-D/2+n_1+n_2)}{(1+n_1)(3-D/2+n_1)} , \\
\Phi'_2(n_1, n_2) &= \frac{A_{n_1, 1+n_2}^{(1a)}}{A_{n_1, n_2}^{(1a)}} = \frac{(1+n_1+n_2)(3-D/2+n_1+n_2)}{(1+n_2)(3-D/2+n_2)} .
\end{aligned} \tag{14}$$

In order to investigate the absolutely and uniformly convergent region of the double series in Eq.(13), we define

$$\begin{aligned}
\Phi_1(n_1, n_2) &= \lim_{\lambda \rightarrow \infty} \Phi'_1(\lambda n_1, \lambda n_2) = \frac{(n_1+n_2)^2}{n_1^2} , \\
\Phi_2(n_1, n_2) &= \lim_{\lambda \rightarrow \infty} \Phi'_2(\lambda n_1, \lambda n_2) = \frac{(n_1+n_2)^2}{n_2^2} .
\end{aligned} \tag{15}$$

The Cartesian curve of the double power series in Eq.(13) is

$$\sqrt{r_1} + \sqrt{r_2} = 1 , \tag{16}$$

with $r_i = |x_{i3}|$. For convenience we denote the region surrounded by the coordinate axes and the Cartesian curve in the positive quadrant of the plane Or_1r_2 by C , and denote the rectangle by D in the positive quadrant of the plane Or_1r_2 bounded by the coordinate axes and the straight lines parallel to the coordinate axes $r_1 = 1/\Phi_1(1, 0) = 1$,

and $r_2 = 1/\Phi_2(0, 1) = 1$. According to Horn's study of convergence of the hypergeometric series [27], one finds the well-known conclusion [25] that the double power series in Eq.(13) absolutely and uniformly converges in the intersection of the regions C and D in the plane Or_1r_2 . In other words, the absolutely and uniformly convergent region of the double power series in Eq.(13) is written as

$$\Omega'_1 = \left\{ (x_{13}, x_{23}) : \left| |x_{13}| < 1, |x_{23}| < 1, \sqrt{|x_{13}|} + \sqrt{|x_{23}|} < 1 \right. \right\}. \quad (17)$$

- 1(b): For the isolated singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(\frac{D}{2} - 2 - z_2)$, the corresponding double hypergeometric function is written as

$$\begin{aligned} C_{0,1b}(p_1^2, p_2^2, p_3^2) &= -\frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D - 3)} x_{23}^{D/2-2} \\ &\times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \frac{\Gamma(1 + n_1 + n_2)\Gamma(\frac{D}{2} - 1 + n_1 + n_2)}{n_1!n_2!\Gamma(3 - \frac{D}{2} + n_1)\Gamma(\frac{D}{2} - 1 + n_2)} x_{13}^{n_1} x_{23}^{n_2}. \end{aligned} \quad (18)$$

Similarly the condition $|p_3^2| > \max(|p_1^2|, |p_2^2|)$ should be satisfied to guarantee the double series converging. Using Horn's theory of convergence, one obtains that the double hypergeometric series above is absolutely and uniformly convergent in the region Eq.(17).

- 1(c): For the isolated singularities defined by the poles of $\Gamma(-z_2)$, $\Gamma(\frac{D}{2} - 2 - z_1)$, the corresponding double hypergeometric function is written as

$$\begin{aligned} C_{0,1c}(p_1^2, p_2^2, p_3^2) &= -\frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D - 3)} x_{13}^{D/2-2} \\ &\times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \frac{\Gamma(1 + n_1 + n_2)\Gamma(\frac{D}{2} - 1 + n_1 + n_2)}{n_1!n_2!\Gamma(\frac{D}{2} - 1 + n_1)\Gamma(3 - \frac{D}{2} + n_2)} x_{13}^{n_1} x_{23}^{n_2}. \end{aligned} \quad (19)$$

Certainly the essential condition $|p_3^2| > \max(|p_1^2|, |p_2^2|)$ should be satisfied to guarantee the convergence of the double series.

- 1(d): For the isolated singularities defined by the poles of $\Gamma(\frac{D}{2} - 2 - z_1)$, $\Gamma(\frac{D}{2} - 2 - z_2)$, the corresponding double hypergeometric function is written as

$$\begin{aligned} C_{0,1d}(p_1^2, p_2^2, p_3^2) &= \frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D - 3)} (x_{13}x_{23})^{D/2-2} \\ &\times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \frac{\Gamma(D - 3 + n_1 + n_2)\Gamma(\frac{D}{2} - 1 + n_1 + n_2)}{n_1!n_2!\Gamma(\frac{D}{2} - 1 + n_1)\Gamma(\frac{D}{2} - 1 + n_2)} x_{13}^{n_1} x_{23}^{n_2}. \end{aligned} \quad (20)$$

Correspondingly the necessary condition $|p_3^2| > \max(|p_1^2|, |p_2^2|)$ should be satisfied to guarantee the convergence of the double series. The absolutely and uniformly convergent region of Eq.(19) and Eq.(20) is also provided in Eq.(17) concretely.

- 2(a): For the isolated singularities defined by the poles of $\Gamma(1 + z_1 + z_2)$, $\Gamma(-z_2)$, the corresponding double hypergeometric function is written as

$$C_{0,2a}(p_1^2, p_2^2, p_3^2) = C_{0,1c}(p_3 \rightarrow p_1, x_{13} \rightarrow x_{31}, x_{23} \rightarrow x_{21}) . \quad (21)$$

The necessary condition $|p_1^2| > \max(|p_2^2|, |p_3^2|)$ should be satisfied to guarantee the convergence of the double series.

- 2(b): For the isolated singularities defined by the poles of $\Gamma(3 - \frac{D}{2} + z_1 + z_2)$, $\Gamma(-z_2)$, the corresponding double hypergeometric function is written as

$$C_{0,2b}(p_1^2, p_2^2, p_3^2) = C_{0,1a}(p_3 \rightarrow p_1, x_{13} \rightarrow x_{31}, x_{23} \rightarrow x_{21}) . \quad (22)$$

- 2(c): For the isolated singularities defined by the poles of $\Gamma(1 + z_1 + z_2)$, $\Gamma(\frac{D}{2} - 2 - z_2)$, the corresponding double hypergeometric function is written as

$$C_{0,2c}(p_1^2, p_2^2, p_3^2) = C_{0,1d}(p_3 \rightarrow p_1, x_{13} \rightarrow x_{31}, x_{23} \rightarrow x_{21}) . \quad (23)$$

- 2(d): For the isolated singularities defined by the poles of $\Gamma(3 - \frac{D}{2} + z_1 + z_2)$, $\Gamma(\frac{D}{2} - 2 - z_2)$, the corresponding double hypergeometric function is written as

$$C_{0,2d}(p_1^2, p_2^2, p_3^2) = C_{0,1b}(p_3 \rightarrow p_1, x_{13} \rightarrow x_{31}, x_{23} \rightarrow x_{21}) . \quad (24)$$

- 3(a): For the isolated singularities defined by the poles of $\Gamma(-z_1)$, and $\Gamma(1 + z_1 + z_2)$, the corresponding double hypergeometric function is written as

$$C_{0,3a}(p_1^2, p_2^2, p_3^2) = C_{0,1b}(p_3 \rightarrow p_2, x_{13} \rightarrow x_{12}, x_{23} \rightarrow x_{32}) . \quad (25)$$

The condition $|p_2^2| > \max(|p_1^2|, |p_3^2|)$ is necessary to guarantee the double series uniformly converging.

- 3(b): For the isolated singularities defined by the poles of $\Gamma(-z_1)$, and $\Gamma(3 - \frac{D}{2} + z_1 + z_2)$, the corresponding double hypergeometric function is written as

$$C_{0,3b}(p_1^2, p_2^2, p_3^2) = C_{0,1a}(p_3 \rightarrow p_2, x_{13} \rightarrow x_{12}, x_{23} \rightarrow x_{32}) . \quad (26)$$

- 3(c): For the isolated singularities defined by the poles of $\Gamma(\frac{D}{2} - 2 - z_1)$, and $\Gamma(1 + z_1 + z_2)$, the corresponding double hypergeometric function is written as

$$C_{0,3c}(p_1^2, p_2^2, p_3^2) = C_{0,1d}(p_3 \rightarrow p_2, x_{13} \rightarrow x_{12}, x_{23} \rightarrow x_{32}). \quad (27)$$

- 3(d): For the isolated singularities defined by the poles of $\Gamma(\frac{D}{2} - 2 - z_1)$, and $\Gamma(3 - \frac{D}{2} + z_1 + z_2)$, the corresponding double hypergeometric function is written as

$$C_{0,3d}(p_1^2, p_2^2, p_3^2) = C_{0,1c}(p_3 \rightarrow p_2, x_{13} \rightarrow x_{12}, x_{23} \rightarrow x_{32}). \quad (28)$$

Applying Horn's study of convergence to Eq.(21), Eq.(22), Eq.(23), and Eq.(24), we obtain the absolutely and uniformly convergent region of the double power series as

$$\Omega'_2 = \Omega'_1(x_{13} \rightarrow x_{31}, x_{23} \rightarrow x_{21}). \quad (29)$$

Similarly the absolutely and uniformly convergent region of the double power series in Eq.(25), Eq.(26), Eq.(27), and Eq.(28) is described by

$$\Omega'_3 = \Omega'_1(x_{13} \rightarrow x_{12}, x_{23} \rightarrow x_{32}). \quad (30)$$

Obviously the intersection $\Omega'_i \cap \Omega'_j$, $i, j = 1, 2, 3$, $i \neq j$ is empty set, i.e.

$$\Omega'_1 \cap \Omega'_2 = \emptyset, \Omega'_1 \cap \Omega'_3 = \emptyset, \Omega'_2 \cap \Omega'_3 = \emptyset. \quad (31)$$

Based on the above-mentioned analyses, the scalar integral C_0 in the convergent region Ω'_1 is presented as

$$\begin{aligned} C_0(p_1^2, p_2^2, p_3^2) &= C_{0,1a}(p_1^2, p_2^2, p_3^2) + C_{0,1b}(p_1^2, p_2^2, p_3^2) \\ &\quad + C_{0,1c}(p_1^2, p_2^2, p_3^2) + C_{0,1d}(p_1^2, p_2^2, p_3^2) \\ &= \frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D - 3)} f(x_{13}, x_{23}), \end{aligned} \quad (32)$$

with

$$\begin{aligned} f(x, y) &= \frac{1}{\Gamma(3 - \frac{D}{2})} F_4 \left(\begin{matrix} 1, & 3 - \frac{D}{2} \\ 3 - \frac{D}{2}, & 3 - \frac{D}{2} \end{matrix} \middle| x, y \right) \\ &\quad - \frac{y^{D/2-2}}{\Gamma(3 - \frac{D}{2})} F_4 \left(\begin{matrix} 1, & \frac{D}{2} - 1 \\ 3 - \frac{D}{2}, & \frac{D}{2} - 1 \end{matrix} \middle| x, y \right) \end{aligned}$$

$$\begin{aligned}
& -\frac{x^{D/2-2}}{\Gamma(3-\frac{D}{2})} F_4 \left(\begin{matrix} 1, \frac{D}{2}-1 \\ \frac{D}{2}-1, 3-\frac{D}{2} \end{matrix} \middle| x, y \right) \\
& + \frac{\Gamma(D-3)}{\Gamma(\frac{D}{2}-1)} (xy)^{D/2-2} F_4 \left(\begin{matrix} D-3, \frac{D}{2}-1 \\ \frac{D}{2}-1, \frac{D}{2}-1 \end{matrix} \middle| x, y \right), \tag{33}
\end{aligned}$$

which is consistent with the Eq. (7) of Ref. [10] exactly, where F_4 denotes the Appell function. Similarly in the convergent region Ω'_2 , the massless C_0 function is given as

$$\begin{aligned}
C_0(p_1^2, p_2^2, p_3^2) &= C_{0,2a}(p_1^2, p_2^2, p_3^2) + C_{0,2b}(p_1^2, p_2^2, p_3^2) \\
&+ C_{0,2c}(p_1^2, p_2^2, p_3^2) + C_{0,2d}(p_1^2, p_2^2, p_3^2) \\
&= \text{Eq. (32)}(p_3 \rightarrow p_1, x_{13} \rightarrow x_{31}, x_{23} \rightarrow x_{21}). \tag{34}
\end{aligned}$$

In the parameter space Ω'_3 , the massless C_0 function is presented as

$$\begin{aligned}
C_0(p_1^2, p_2^2, p_3^2) &= C_{0,3a}(p_1^2, p_2^2, p_3^2) + C_{0,3b}(p_1^2, p_2^2, p_3^2) \\
&+ C_{0,3c}(p_1^2, p_2^2, p_3^2) + C_{0,3d}(p_1^2, p_2^2, p_3^2) \\
&= \text{Eq. (32)}(p_3 \rightarrow p_2, x_{13} \rightarrow x_{12}, x_{23} \rightarrow x_{32}). \tag{35}
\end{aligned}$$

The emphasized point here is that the functions $f(x_{13}, x_{23})$, $x_{13}^{D/2-3} f(x_{31}, x_{21})$, and $x_{23}^{D/2-3} f(x_{12}, x_{32})$ all satisfy the system of PDEs[24, 29–31]

$$\left\{ \left(\sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} + 1 \right) \left(\sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} + 3 - \frac{D}{2} \right) - \frac{1}{x_{j3}} \hat{\vartheta}_{x_{j3}} \left(\hat{\vartheta}_{x_{j3}} + 2 - \frac{D}{2} \right) \right\} f = 0, \quad (j = 1, 2). \tag{36}$$

How to continue the solution to the whole parameter space has already been presented in our previous work [28]. Actually the function $f(x, y)$ can be written in term of the Gauss function ${}_2F_1$ [12] by the well-known reduction of the Appell function of the fourth kind [11], then the massless scalar integral is analytic continued through the transformations of the Gauss function. Continuation through the holonomic hypergeometric system of PDEs here can be applied to evaluate the scalar integral of any multiloop Feynman diagram.

IV. THE C_0 FUNCTION WITH ONE NONZERO VIRTUAL MASS

In this case, the Mellin-Barnes representation of the scalar integral is

$$C_0^a(p_1^2, p_2^2, p_3^2, m^2) = \int \frac{d^D q}{(2\pi)^D} \frac{1}{(q^2 - m^2)(q + p_1)^2(q - p_2)^2}$$

$$\begin{aligned}
&= -\frac{i(-)^{D/2-3}}{(4\pi)^{D/2}(2\pi i)^3} \int_{-i\infty}^{+i\infty} ds (-m^2)^s \Gamma(-s) \\
&\quad \times \int_{-i\infty}^{+i\infty} dz_1 (p_1^2)^{z_1} \Gamma(-z_1) \int_{-i\infty}^{+i\infty} dz_2 (p_2^2)^{z_2} \Gamma(-z_2) \\
&\quad \times (p_3^2)^{D/2-3-s-z_1-z_2} \Gamma\left(3 - \frac{D}{2} + s + z_1 + z_2\right) \\
&\quad \times \Gamma\left(\frac{D}{2} - 2 - s - z_2\right) \Gamma\left(\frac{D}{2} - 2 - s - z_1\right) \\
&\quad \times \frac{\Gamma(1 + s + z_1 + z_2)}{\Gamma(D - 3 - s)}. \tag{37}
\end{aligned}$$

which coincides with Eq. (26) of Ref. [13] (i.e. Eq. (4.3) of Ref. [15]). Applying the residue theorem to the singularities of Γ functions in the numerator of integrand in the Mellin-Barnes representation, we formulate the scalar integral as various hypergeometric functions whose absolutely and uniformly convergent regions are analyzed through the Horn's study of convergence [27]. If the scalar integral is expressed as

$$C_0^a(p_1^2, p_2^2, p_3^2, m^2) = \frac{i(-)^{D/2}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} F_{a,p_3}(\xi_{33}, x_{13}, x_{23}) \tag{38}$$

with $\xi_{33} = -m^2/p_3^2$, then the dimensionless function F_{a,p_3} satisfies the holonomic hypergeometric system of linear PDEs

$$\begin{aligned}
&\left\{ \left(3 - \frac{D}{2} + \hat{\vartheta}_{\xi_{33}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}}\right) \left(1 + \hat{\vartheta}_{\xi_{33}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}}\right) (4 - D + \hat{\vartheta}_{\xi_{33}}) \right. \\
&\quad \left. - \frac{1}{\xi_{33}} \hat{\vartheta}_{\xi_{33}} \prod_{i=1}^2 \left(2 - \frac{D}{2} + \hat{\vartheta}_{\xi_{33}} + \hat{\vartheta}_{x_{i3}}\right) \right\} F_{a,p_3} = 0, \\
&\left\{ \left(3 - \frac{D}{2} + \hat{\vartheta}_{\xi_{33}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}}\right) \left(1 + \hat{\vartheta}_{\xi_{33}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}}\right) \right. \\
&\quad \left. - \frac{1}{x_{j3}} \hat{\vartheta}_{x_{j3}} \left(2 - \frac{D}{2} + \hat{\vartheta}_{\xi_{33}} + \hat{\vartheta}_{x_{j3}}\right) \right\} F_{a,p_3} = 0, \quad (j = 1, 2). \tag{39}
\end{aligned}$$

Setting $m_1 = m_2 = 0$ in the corresponding system of linear PDEs of Eq. (3), we find the first, fourth and fifth PDEs composing the system presented above. Now we present our results in detail below.

A. The hypergeometric functions and their convergent regions

- 1: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(-z_1)$, and $\Gamma(-z_2)$ in the numerator of integrand of the Mellin-Barnes representation, the derived hypergeomet-

ric function is written as

$$\begin{aligned}
C_{0,1}^a(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3)\Gamma(4-D)} \\
&\times \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{33}^j x_{13}^{n_1} x_{23}^{n_2} \frac{\Gamma(4-D+j)}{j!n_1!n_2!} \\
&\times \frac{\Gamma(3 - \frac{D}{2} + j + n_1 + n_2)\Gamma(1+j+n_1+n_2)}{\Gamma(3 - \frac{D}{2} + j + n_1)\Gamma(3 - \frac{D}{2} + j + n_2)} \\
&= \frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3)} \\
&\times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} x_{13}^{n_1} x_{23}^{n_2} \\
&\times \frac{\Gamma(3 - \frac{D}{2} + n_1 + n_2)\Gamma(1+n_1+n_2)}{n_1!n_2!\Gamma(3 - \frac{D}{2} + n_1)\Gamma(3 - \frac{D}{2} + n_2)} \\
&+ \frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3)\Gamma(4-D)} \\
&\times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{33}^j x_{13}^{n_1} x_{23}^{n_2} \frac{\Gamma(4-D+j)}{j!n_1!n_2!} \\
&\times \frac{\Gamma(3 - \frac{D}{2} + j + n_1 + n_2)\Gamma(1+j+n_1+n_2)}{\Gamma(3 - \frac{D}{2} + j + n_1)\Gamma(3 - \frac{D}{2} + j + n_2)}. \quad (40)
\end{aligned}$$

The condition $|p_3^2| > \max(m^2, |p_1^2|, |p_2^2|)$ is necessary to guarantee the convergence of the hypergeometric series. In addition, $D = 4$ is the second order pole of the first double hypergeometric function, and the first order pole of the second triple hypergeometric function, respectively. Certainly the expression of Eq. (40) complies with the system of the PDEs in Eq. (39) explicitly.

The absolutely and uniformly convergent region of the hypergeometric functions is given by

$$\Omega_1^a = \left\{ (\xi_{33}, x_{13}, x_{23}) : \left| \begin{array}{l} \sqrt{|x_{13}|} + \sqrt{|x_{23}|} < 1, \quad |\xi_{33}| + |x_{13}| < 1, \\ |\xi_{33}| + |x_{23}| < 1 \end{array} \right. \right\}. \quad (41)$$

- 2: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(-z_1)$, and $\Gamma(\frac{D}{2}-2-s-z_2)$ in the numerator of integrand of the Mellin-Barnes representation, the corresponding hypergeometric series is given as

$$C_{0,2}^a(p_1^2, p_2^2, p_3^2, m^2) = -\frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3)\Gamma(4-D)}$$

$$\begin{aligned}
& \times \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{33}^j x_{13}^{n_1} x_{23}^{D/2-2-j+n_2} \frac{\Gamma(4-D+j)}{j!n_1!n_2!} \\
& \times \frac{\Gamma(1+n_1+n_2)\Gamma(\frac{D}{2}-1+n_1+n_2)}{\Gamma(3-\frac{D}{2}+j+n_1)\Gamma(\frac{D}{2}-1-j+n_2)} \\
& = -\frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2-\frac{D}{2})\Gamma^2(\frac{D}{2}-1)}{(4\pi)^{D/2} \Gamma(D-3)} \\
& \times \sum_{n_1=0}^{\infty} \sum_{n_2=1}^{\infty} x_{13}^{n_1} x_{23}^{D/2-2+n_2} \\
& \times \frac{\Gamma(1+n_1+n_2)\Gamma(\frac{D}{2}-1+n_1+n_2)}{n_1!n_2!\Gamma(3-\frac{D}{2}+n_1)\Gamma(\frac{D}{2}-1+n_2)} \\
& -\frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2-\frac{D}{2})\Gamma^2(\frac{D}{2}-1)}{(4\pi)^{D/2} \Gamma(D-3)\Gamma(4-D)} \\
& \times \sum_{n_1=0}^{\infty} \sum_{j=1}^{\infty} \sum_{n_2=1}^{\infty} \xi_{33}^j x_{13}^{n_1} x_{23}^{D/2-2+n_2} \frac{\Gamma(4-D+j)}{j!n_1!(j+n_2)!} \\
& \times \frac{\Gamma(1+j+n_1+n_2)\Gamma(\frac{D}{2}-1+j+n_1+n_2)}{\Gamma(3-\frac{D}{2}+j+n_1)\Gamma(\frac{D}{2}-1+n_2)} \\
& -\frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma(2-\frac{D}{2})\Gamma(\frac{D}{2}-1)}{(4\pi)^{D/2} \Gamma(D-3)\Gamma(4-D)} \\
& \times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \sum_{j=1}^{\infty} \xi_{33}^{j+n_2} x_{13}^{n_1} x_{23}^{D/2-2-j} \\
& \times \frac{(-)^j \Gamma(4-D+j+n_2)\Gamma(2-\frac{D}{2}+j)}{(j+n_2)!n_1!n_2!} \\
& \times \frac{\Gamma(1+n_1+n_2)\Gamma(\frac{D}{2}-1+n_1+n_2)}{\Gamma(3-\frac{D}{2}+j+n_1+n_2)} \\
& -\frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2-\frac{D}{2})\Gamma(\frac{D}{2}-1)}{(4\pi)^{D/2} \Gamma(D-3)\Gamma(4-D)} \\
& \times \sum_{n_1=0}^{\infty} \sum_{j=1}^{\infty} \xi_{33}^j x_{13}^{n_1} x_{23}^{D/2-2} \frac{\Gamma(4-D+j)}{(j!)^2 n_1!} \\
& \times \frac{\Gamma(1+j+n_1)\Gamma(\frac{D}{2}-1+j+n_1)}{\Gamma(3-\frac{D}{2}+j+n_1)} \\
& -\frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2-\frac{D}{2})\Gamma(\frac{D}{2}-1)}{(4\pi)^{D/2} \Gamma(D-3)} \\
& \times \sum_{n_1=0}^{\infty} x_{13}^{n_1} x_{23}^{D/2-2} \frac{\Gamma(\frac{D}{2}-1+n_1)}{\Gamma(3-\frac{D}{2}+n_1)}. \tag{42}
\end{aligned}$$

The essential condition $|p_3^2| > \max(m^2, |p_1^2|, |p_2^2|)$, $|p_2^2| > m^2$ should be satisfied to guarantee the hypergeometric series converging. In the above equation, one derives the

first expression after applying the residue theorem to the Mellin-Barnes representation. The second expression decomposes the summation indices of the hypergeometric function in the first expression, so that $D = 4$ is the pole of the same fixed order for each term of every hypergeometric series in the second expression. Furthermore, we can check the second expression satisfying the system of linear PDEs in Eq. (39) explicitly, this fact implies that the decomposition of summation indices above is correct and reasonable.

Horn's theory of convergence predicts the absolutely and uniformly convergent region of the hypergeometric functions as

$$\Omega_2^a = \left\{ (\xi_{33}, x_{13}, x_{23}) : \begin{cases} |\xi_{33}| < |x_{23}|, \sqrt{|x_{13}|} + \sqrt{|x_{23}|} < 1, \\ |\xi_{33}| + |x_{13}| < 1, |\xi_{33}| + |x_{23}| < 1 \end{cases} \right\}. \quad (43)$$

- 3: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(-z_1)$, and $\Gamma(1+s+z_1+z_2)$ in the numerator of integrand of the Mellin-Barnes representation, the corresponding power series is

$$\begin{aligned} C_{0,3}^a(p_1^2, p_2^2, p_3^2, m^2) &= -\frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3)\Gamma(4-D)} \\ &\quad \times \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{33}^j x_{13}^{n_1} x_{23}^{-1-j-n_1-n_2} \frac{(-)^j \Gamma(4-D+j)}{j!n_1!n_2!} \\ &\quad \times \frac{\Gamma(1+j+n_1+n_2)\Gamma(\frac{D}{2}-1+n_1+n_2)}{\Gamma(3-\frac{D}{2}+j+n_1)\Gamma(\frac{D}{2}-1+n_2)} \\ &= -\frac{i(-)^{D/2}(p_2^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3)} \\ &\quad \times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} x_{12}^{n_1} x_{32}^{D/2-2+n_2} \\ &\quad \times \frac{\Gamma(1+n_1+n_2)\Gamma(\frac{D}{2}-1+n_1+n_2)}{n_1!n_2!\Gamma(3-\frac{D}{2}+n_1)\Gamma(\frac{D}{2}-1+n_2)} \\ &\quad - \frac{i(-)^{D/2}(p_2^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3)\Gamma(4-D)} \\ &\quad \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{32}^j x_{12}^{n_1} x_{32}^{D/2-2+n_2} \frac{(-)^j \Gamma(4-D+j)}{j!n_1!n_2!} \\ &\quad \times \frac{\Gamma(1+j+n_1+n_2)\Gamma(\frac{D}{2}-1+n_1+n_2)}{\Gamma(3-\frac{D}{2}+j+n_1)\Gamma(\frac{D}{2}-1+n_2)}. \end{aligned} \quad (44)$$

The necessary condition $|p_2^2| > \max(|p_1^2|, |p_3^2|, m^2)$ should be satisfied to guarantee the

power series converging. In addition, $D = 4$ is the second order pole of the first double hypergeometric function, and the first order pole of the second triple hypergeometric function, respectively. Certainly the expression of Eq. (44) complies with the system of linear PDEs in Eq. (39) explicitly.

Through Horn's study of convergence, the absolutely and uniformly convergent region of the series is written as

$$\Omega_3^a = \left\{ (\xi_{32}, x_{12}, x_{32}) : \left| |\xi_{32}| + |x_{32}| < 1, \sqrt{|x_{12}|} + \sqrt{|x_{32}|} < 1 \right. \right\}. \quad (45)$$

- 4: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(-z_1)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$ in the numerator of integrand of the Mellin-Barnes representation, the corresponding hypergeometric function is written as

$$\begin{aligned} C_{0,4}^a(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(-)^{D/2} (p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} \\ &\quad \times \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{33}^j x_{13}^{n_1} x_{23}^{D/2-3-j-n_1-n_2} \frac{(-)^j \Gamma(4-D+j)}{j! n_1! n_2!} \\ &\quad \times \frac{\Gamma(3 - \frac{D}{2} + j + n_1 + n_2) \Gamma(1 + n_1 + n_2)}{\Gamma(3 - \frac{D}{2} + j + n_1) \Gamma(3 - \frac{D}{2} + n_2)} \\ &= \frac{i(-)^{D/2} (p_2^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3)} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \\ &\quad \times x_{12}^{n_1} x_{32}^{n_2} \frac{\Gamma(3 - \frac{D}{2} + n_1 + n_2) \Gamma(1 + n_1 + n_2)}{n_1! n_2! \Gamma(3 - \frac{D}{2} + n_1) \Gamma(3 - \frac{D}{2} + n_2)} \\ &\quad + \frac{i(-)^{D/2} (p_2^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \\ &\quad \times \xi_{32}^j x_{12}^{n_1} x_{32}^{n_2} \frac{(-)^j \Gamma(4-D+j)}{j! n_1! n_2!} \\ &\quad \times \frac{\Gamma(3 - \frac{D}{2} + j + n_1 + n_2) \Gamma(1 + n_1 + n_2)}{\Gamma(3 - \frac{D}{2} + j + n_1) \Gamma(3 - \frac{D}{2} + n_2)}. \end{aligned} \quad (46)$$

The necessary condition $|p_2^2| > \max(|p_3^2|, |p_1^2|, m^2)$ should be satisfied to guarantee the hypergeometric series converging. The convergent region of the term is consistent with Ω_3^a in Eq.(45) exactly. Furthermore, $D = 4$ is the second order pole of the first double hypergeometric function, and the first order pole of the second triple hypergeometric function, respectively. Certainly the expression of Eq. (46) complies with the system of linear PDEs in Eq. (39) explicitly.

- 6: For the isolated singularities originating from the poles of $\Gamma(-s)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(-z_2)$ in the numerator of integrand of the Mellin-Barnes representation, the corresponding hypergeometric series is

$$C_{0,6}^a(p_1^2, p_2^2, p_3^2, m^2) = C_{0,2}^a(x_{13} \leftrightarrow x_{23}). \quad (47)$$

The condition $|p_3^2| > \max(m^2, |p_1^2|, |p_2^2|)$, $|p_1^2| > m^2$ should be met to guarantee the hypergeometric functions converging. Applying Horn's study of convergence, one gets the absolutely and uniformly convergent region of the hypergeometric series as

$$\Omega_6^a = \left\{ (\xi_{33}, x_{13}, x_{23}) : \begin{cases} |\xi_{33}| < |x_{13}|, \sqrt{|x_{13}|} + \sqrt{|x_{23}|} < 1, \\ |\xi_{33}| + |x_{13}| < 1, |\xi_{33}| + |x_{23}| < 1 \end{cases} \right\}. \quad (48)$$

- 7: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(\frac{D}{2} - 2 - s - z_2)$ in the numerator of integrand of the Mellin-Barnes representation, the corresponding hypergeometric series $C_{0,7}^a$ is presented in Eq. (A1), where the condition $|p_3^2| > \max(|p_1^2|, |p_2^2|)$, $\min(|p_1^2|, |p_2^2|) > m^2$, $|p_1^2 p_2^2| > m^2 |p_3^2|$ should be met to guarantee the power series converging. After implementing the residue theorem, one derives the first expression in Eq. (A1). The second expression decomposes the summation indices of the hypergeometric series in the first expression, so that $D = 4$ is the pole of the same fixed order for each term of every hypergeometric series in the second expression. In addition, the second expression also satisfies the system of linear PDEs in Eq. (39).

The absolutely and uniformly convergent region of the term is written as

$$\Omega_7^a = \left\{ (\xi_{33}, x_{13}, x_{23}) : \begin{cases} \sqrt{|x_{13}|} + \sqrt{|x_{23}|} < 1, |\xi_{33}| + |x_{13}| < 1, \\ |\xi_{33}| + |x_{23}| < 1, |\xi_{33}|(1 + |x_{23}|) < |x_{13} x_{23}|, \\ |\xi_{33}|(1 + |x_{13}|) < |x_{13} x_{23}|, \\ |\xi_{33}| < (|x_{13} - \xi_{33}|)(|x_{23} - \xi_{33}|) \end{cases} \right\}. \quad (49)$$

- 8: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(1 + s + z_1 + z_2)$ in the numerator of integrand of the Mellin-Barnes representation, correspondingly the tedious hypergeometric series $C_{0,8}^a$ is given in Eq. (A2), where the necessary condition $|p_2^2| > \max(|p_1^2|, |p_3^2|)$, $|p_1^2| > m^2$ should be satisfied to guarantee

the triple series converging. Applying the residue theorem, one derives the first expression of Eq. (A2). The second expression decomposes the summation indices of the hypergeometric series in the first expression, so that $D = 4$ is the pole of the same fixed order for each term of every hypergeometric series in the second expression. In addition, the second expression also satisfies the system of linear PDEs in Eq. (39).

The absolutely and uniformly convergent region of the series is given by

$$\Omega_8^a = \left\{ (\xi_{32}, x_{12}, x_{32}) : \begin{cases} \sqrt{|x_{12}|} + \sqrt{|x_{32}|} < 1, & |\xi_{32}| + |x_{32}| < 1, \\ |\xi_{32}| + |x_{12}| < 1, & |\xi_{32}|(1 + |x_{32}|) < |x_{12}|, \\ |\xi_{32}|(|x_{12}| + |x_{32}|) < |x_{12}|, \\ |\xi_{32}x_{32}| < (|x_{12}| - |\xi_{32}|)(1 - |\xi_{32}|) \end{cases} \right\}. \quad (50)$$

- 9: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, the corresponding power series $C_{0,9}^a$ is given by Eq. (A3), where the condition $|p_2^2| > \max(|p_1^2|, |p_3^2|)$, $|p_1^2| > m^2$ should be satisfied to guarantee the triple series converging. Applying the residue theorem, one derives the first expression of Eq. (A3). The second expression decomposes the summation indices of the hypergeometric series in the first expression, so that $D = 4$ is the pole of the same fixed order for each term of every hypergeometric series in the second expression. In addition, the second expression also satisfies the system of linear PDEs in Eq. (39). The convergent region of the series coincides with the region Ω_8^a in Eq.(50) exactly.

- 10: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(1 + s + z_1 + z_2)$, and $\Gamma(-z_2)$, the corresponding hypergeometric series is

$$C_{0,10}^a(p_1^2, p_2^2, p_3^2, m^2) = C_{0,3}^a(p_2 \rightarrow p_1, \xi_{32} \rightarrow \xi_{31}, x_{12} \rightarrow x_{21}, x_{32} \rightarrow x_{31}). \quad (51)$$

The condition $|p_1^2| > \max(|p_2^2|, |p_3^2|, m^2)$ should be satisfied to guarantee the hypergeometric series converging. Horn's theory of convergence gives the absolutely and uniformly convergent region of the series as

$$\Omega_{10}^a = \left\{ (\xi_{31}, x_{21}, x_{31}) : \left| |\xi_{31}| + |x_{31}| < 1, \sqrt{|x_{21}|} + \sqrt{|x_{31}|} < 1 \right. \right\}. \quad (52)$$

Additionally $D = 4$ is the second order pole of the first double hypergeometric function, and the first order pole of the second triple hypergeometric function, respectively.

Certainly the expression of Eq. (46) also complies with the system of linear PDEs in Eq. (39).

- 11: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(1 + s + z_1 + z_2)$, and $\Gamma(\frac{D}{2} - 2 - s - z_2)$, the derived hypergeometric series $C_{0,11}^a$ is

$$C_{0,11}^a(p_1^2, p_2^2, p_3^2, m^2) = C_{0,8}^a(p_2 \rightarrow p_1, \xi_{32} \rightarrow \xi_{31}, x_{12} \rightarrow x_{21}, x_{32} \rightarrow x_{31}), \quad (53)$$

where the condition $|p_1^2| > \max(|p_2^2|, |p_3^2|)$, $|p_2^2| > m^2$ should be met to guarantee the hypergeometric series converging. The absolutely and uniformly convergent region of the hypergeometric series is given by

$$\Omega_{11}^a = \left\{ (\xi_{31}, x_{21}, x_{31}) : \begin{cases} \sqrt{|x_{21}|} + \sqrt{|x_{31}|} < 1, & |\xi_{31}| + |x_{31}| < 1, \\ |\xi_{31}| + |x_{21}| < 1, & |\xi_{31}|(1 + |x_{31}|) < |x_{21}|, \\ |\xi_{31}|(|x_{21}| + |x_{31}|) < |x_{21}|, \\ |\xi_{31}x_{31}| < (|x_{21}| - |\xi_{31}|)(1 - |\xi_{31}|) \end{cases} \right\}. \quad (54)$$

- 13: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, and $\Gamma(-z_2)$, the induced hypergeometric series is

$$C_{0,13}^a(p_1^2, p_2^2, p_3^2, m^2) = C_{0,4}^a(p_2 \rightarrow p_1, \xi_{32} \rightarrow \xi_{31}, x_{12} \rightarrow x_{21}, x_{32} \rightarrow x_{31}). \quad (55)$$

The condition $|p_1^2| > \max(|p_2^2|, |p_3^2|, m^2)$ should be met to guarantee the triple series converging. The convergent region of this series is characterized by Eq.(52) also.

- 14: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, and $\Gamma(\frac{D}{2} - 2 - s - z_2)$, the corresponding hypergeometric series $C_{0,14}^a$ is

$$C_{0,14}^a(p_1^2, p_2^2, p_3^2, m^2) = C_{0,9}^a(p_2 \rightarrow p_1, \xi_{32} \rightarrow \xi_{31}, x_{12} \rightarrow x_{21}, x_{32} \rightarrow x_{31}), \quad (56)$$

where the condition $|p_1^2| > \max(|p_2^2|, |p_3^2|)$, $|p_2^2| > m^2$ should be satisfied to guarantee the hypergeometric series converging. The convergent region of the series is also defined by Ω_{11}^a in Eq.(54).

- 16: For the isolated singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(-z_2)$, and $\Gamma(\frac{D}{2} - 2 - s - z_1)$, the scalar integral contains some unknown linear combining parameters originating from the non-isolating singularities. The essential condition $|p_3^2| > \max(|p_2^2|, m^2)$, $m^2 > |p_1^2|$ should be satisfied to guarantee the corresponding hypergeometric series converging.

- 17: For the isolated singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(-z_2)$, and $\Gamma(\frac{D}{2} - 2 - s - z_2)$, the corresponding hypergeometric function contains some unknown linear combining parameters originating from the non-isolating singularities. The condition $|p_3^2| > \max(|p_1^2|, m^2)$, $m^2 > |p_2^2|$ should be satisfied to guarantee the derived hypergeometric series converging.
- 18: For the isolated singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(-z_2)$, and $\Gamma(1 + s + z_1 + z_2)$, the hypergeometric series is

$$\begin{aligned}
C_{0,18}^a(p_1^2, p_2^2, p_3^2, m^2) &= -\frac{i(m^2)^{D/2-3}\eta_{33}^{D/2-2}}{(4\pi)^{D/2}}\Gamma(2 - \frac{D}{2})\Gamma(\frac{D}{2} - 1) \\
&\times \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \frac{(-)^{n_1+n_2}}{j!n_1!n_2!} \eta_{33}^j \eta_{13}^{n_1} \eta_{23}^{n_2} \Gamma(\frac{D}{2} - 1 + j + n_1) \\
&\times \frac{\Gamma(1 + j + n_1 + n_2)\Gamma(\frac{D}{2} - 1 + j + n_2)}{\Gamma(D - 2 + j + n_1 + n_2)\Gamma(\frac{D}{2} - 1 + j)}, \quad (57)
\end{aligned}$$

the condition $m^2 > \max(|p_1^2|, |p_2^2|, |p_3^2|)$ should be satisfied to guarantee the power series converging. The absolutely and uniformly convergent region of the series of Eq.(57) is written as

$$\Omega_{18}^a = \left\{ (\eta_{13}, \eta_{23}, \eta_{33}) : \begin{cases} |\eta_{13}| + |\eta_{33}| < 1, & |\eta_{23}| + |\eta_{33}| < 1, \\ |\eta_{33}| < (1 - |\eta_{13}|)(1 - |\eta_{23}|) \end{cases} \right\}. \quad (58)$$

Certainly the scalar integral of Eq. (57) also satisfies the system of PDEs in Eq. (39).

- 19: For the isolated singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(-z_2)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, correspondingly the hypergeometric series is

$$\begin{aligned}
C_{0,19}^a(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(m^2)^{D/2-3}}{(4\pi)^{D/2}}\Gamma(2 - \frac{D}{2})\Gamma(\frac{D}{2} - 1) \\
&\times \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \frac{(-)^{n_1+n_2}}{j!n_1!n_2!} \eta_{33}^j \eta_{13}^{n_1} \eta_{23}^{n_2} \Gamma(1 + j + n_1) \\
&\times \frac{\Gamma(3 - \frac{D}{2} + j + n_1 + n_2)\Gamma(1 + j + n_2)}{\Gamma(\frac{D}{2} + j + n_1 + n_2)\Gamma(3 - \frac{D}{2} + j)}. \quad (59)
\end{aligned}$$

The convergent region of Eq.(59) totally coincides with Ω_{18}^a in Eq.(58).

- 20: For the isolated singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(\frac{D}{2} - 2 - s - z_2)$, the derived hypergeometric function contains some unknown linear

combination parameters originating from the non-isolated singularities. The condition $|p_3^2| > \max(|p_1^2|, |p_2^2|, m^2)$, $|p_2^2| > m^2$, $|p_3^2 m^2| > |p_1^2 p_2^2|$ should be satisfied to guarantee the corresponding triple series converging.

- 23: For the isolated singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(1 + s + z_1 + z_2)$, the corresponding hypergeometric series is

$$\begin{aligned}
C_{0,23}^a(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(-)^{D/2}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1) \\
&\times \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{33}^{D/2-2+j-n_1} x_{13}^{n_1} x_{23}^{1-D/2-j-n_2} \\
&\times \frac{(-)^{j+n_1} \Gamma(\frac{D}{2} - 1 + n_1 + n_2) \Gamma(\frac{D}{2} - 1 + j + n_2)}{j! n_1! n_2! \Gamma(\frac{D}{2} - 1 + n_2)} \\
&\times \frac{\Gamma(2 - \frac{D}{2} - j + n_1)}{\Gamma(\frac{D}{2} - 1 - j + n_1)} \\
&= \frac{i(-)^{D/2}(p_2^2)^{D/2-3}}{(4\pi)^{D/2}} \Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1) \\
&\times \sum_{j=0}^{\infty} \sum_{n_1=1}^{\infty} \sum_{n_2=0}^{\infty} \xi_{32}^{D/2-2-n_1} x_{12}^{j+n_1} x_{32}^{D/2-2+n_2} \\
&\times \frac{(-)^{n_1} \Gamma(\frac{D}{2} - 1 + j + n_1 + n_2) \Gamma(\frac{D}{2} - 1 + j + n_2)}{j!(j+n_1)! n_2! \Gamma(\frac{D}{2} - 1 + n_2)} \\
&\times \frac{\Gamma(2 - \frac{D}{2} + n_1)}{\Gamma(\frac{D}{2} - 1 + n_1)} \\
&+ \frac{i(-)^{D/2}(p_2^2)^{D/2-3}}{(4\pi)^{D/2}} \Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1) \\
&\times \sum_{n_1=0}^{\infty} \sum_{j=1}^{\infty} \sum_{n_2=0}^{\infty} \xi_{32}^{D/2-2+j} x_{12}^{n_1} x_{32}^{D/2-2+n_2} \\
&\times \frac{(-)^j \Gamma(\frac{D}{2} - 1 + n_1 + n_2) \Gamma(\frac{D}{2} - 1 + j + n_1 + n_2)}{(j+n_1)! n_1! n_2! \Gamma(\frac{D}{2} - 1 + n_2)} \\
&\times \frac{\Gamma(2 - \frac{D}{2} + j)}{\Gamma(\frac{D}{2} - 1 + j)} \\
&+ \frac{i(-)^{D/2}(p_2^2)^{D/2-3}}{(4\pi)^{D/2}} \Gamma^2(2 - \frac{D}{2}) \\
&\times \sum_{j=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{32}^{D/2-2} x_{12}^j x_{32}^{D/2-2+n_2} \\
&\times \frac{\Gamma(\frac{D}{2} - 1 + j + n_2) \Gamma(\frac{D}{2} - 1 + j + n_2)}{(j!)^2 n_2! \Gamma(\frac{D}{2} - 1 + n_2)}. \tag{60}
\end{aligned}$$

The condition $|p_2^2| > \max(|p_1^2|, |p_3^2|, m^2)$, $m^2 > |p_1^2|$ is essential to guarantee the convergence of the hypergeometric series. By implementing the residue theorem, one derives the first expression above. The second expression decomposes the summation indices of the hypergeometric series in the first expression, so that $D = 4$ is the pole of the same fixed order for each term of every hypergeometric series in the second expression above. In addition, the second expression also satisfies the system of linear PDEs in Eq. (39).

Correspondingly the convergent region of the series is given by

$$\Omega_{23}^a = \left\{ (\xi_{32}, x_{12}, x_{32}) : \begin{cases} |x_{12}| < |\xi_{32}|, \sqrt{|x_{12}|} + \sqrt{|x_{32}|} < 1, \\ |\xi_{32}| + |x_{32}| < 1, |\xi_{32}| + |x_{12}| < 1, \\ |x_{12}| + |\xi_{32}x_{32}| < |\xi_{32}|, |x_{12}|(1 + |\xi_{32}|) < |\xi_{32}| \end{cases} \right\}. \quad (61)$$

- 24: For the isolated singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, the corresponding hypergeometric series is

$$\begin{aligned} C_{0,24}^a(p_1^2, p_2^2, p_3^2, m^2) &= -\frac{i(-)^{D/2}(p_2^2)^{D/2-3}}{(4\pi)^{D/2}}\Gamma(2 - \frac{D}{2})\Gamma(\frac{D}{2} - 1) \\ &\times \sum_{j=0}^{\infty} \sum_{n_1=1}^{\infty} \sum_{n_2=0}^{\infty} \xi_{32}^{D/2-2-n_1} x_{12}^{j+n_1} x_{32}^{n_2} \\ &\times \frac{(-)^{n_1} \Gamma(1 + j + n_1 + n_2) \Gamma(1 + j + n_2)}{j!(j + n_1)!n_2! \Gamma(3 - \frac{D}{2} + n_2)} \\ &\times \frac{\Gamma(2 - \frac{D}{2} + n_1)}{\Gamma(\frac{D}{2} - 1 + n_1)} \\ &- \frac{i(-)^{D/2}(p_2^2)^{D/2-3}}{(4\pi)^{D/2}}\Gamma(2 - \frac{D}{2})\Gamma(\frac{D}{2} - 1) \\ &\times \sum_{n_1=0}^{\infty} \sum_{j=1}^{\infty} \sum_{n_2=0}^{\infty} \xi_{32}^{D/2-2+j} x_{12}^{n_1} x_{32}^{n_2} \\ &\times \frac{(-)^j \Gamma(1 + n_1 + n_2) \Gamma(1 + j + n_1 + n_2)}{(j + n_1)!n_1!n_2! \Gamma(3 - \frac{D}{2} + n_2)} \\ &\times \frac{\Gamma(2 - \frac{D}{2} + j)}{\Gamma(\frac{D}{2} - 1 + j)} \\ &- \frac{i(-)^{D/2}(p_2^2)^{D/2-3}}{(4\pi)^{D/2}}\Gamma^2(2 - \frac{D}{2}) \\ &\times \sum_{j=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{32}^{D/2-2} x_{12}^j x_{32}^{n_2} \end{aligned}$$

$$\times \frac{\Gamma^2(1+j+n_2)}{(j!)^2 n_2! \Gamma(3 - \frac{D}{2} + n_2)}. \quad (62)$$

The condition $|p_2^2| > \max(|k^2|, |p_1^2|, m^2)$, $m^2 > |p_1^2|$ is essential to guarantee the multiple power series converging. In addition, $D = 4$ is the first order pole of the first two triple hypergeometric functions, and the second order pole of the third double hypergeometric function, respectively. Certainly the expression satisfies the system of linear PDEs in Eq. (39).

Applying Horn's study of convergence, one finds that the absolutely and uniformly convergent region of Eq.(62) is given by the domain Ω_{23}^a exactly.

- 26: For the isolated singularities defined by the poles of $\Gamma(-z_2)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(\frac{D}{2} - 2 - s - z_2)$, the derived hypergeometric function contains some unknown linear combining parameters originating from the non-isolating singularities. The condition $|p_3^2| > \max(|p_1^2|, |p_2^2|, m^2)$, $|p_1^2| > m^2$, $|p_3^2 m^2| > |p_1^2 p_2^2|$ is necessary to guarantee the hypergeometric series converging.
- 29: For the isolated singularities defined by the poles of $\Gamma(-z_2)$, $\Gamma(\frac{D}{2} - 2 - s - z_2)$, and $\Gamma(1 + s + z_1 + z_2)$, the corresponding hypergeometric series is written as

$$C_{0,29}^a(p_1^2, p_2^2, p_3^2, m^2) = C_{0,23}^a(p_2 \rightarrow p_1, \xi_{32} \rightarrow \xi_{31}, x_{12} \rightarrow x_{21}, x_{32} \rightarrow x_{31}). \quad (63)$$

The convergent region of the series is similarly given by

$$\Omega_{29}^a = \left\{ (\xi_{31}, x_{21}, x_{31}) : \begin{cases} |x_{21}| < |\xi_{31}|, \sqrt{|x_{21}|} + \sqrt{|x_{31}|} < 1, \\ |\xi_{31}| + |x_{21}| < 1, |\xi_{31}| + |x_{31}| < 1, \\ |\xi_{31} x_{31}| + |x_{21}| < |\xi_{31}|, |x_{21}|(1 + |\xi_{31}|) < |\xi_{31}| \end{cases} \right\}. \quad (64)$$

- 30: When the isolated singularities are defined by the poles of $\Gamma(-z_2)$, $\Gamma(\frac{D}{2} - 2 - s - z_2)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, the corresponding hypergeometric series is

$$C_{0,30}^a(p_1^2, p_2^2, p_3^2, m^2) = C_{0,24}^a(p_2 \rightarrow p_1, \xi_{32} \rightarrow \xi_{31}, x_{12} \rightarrow x_{21}, x_{32} \rightarrow x_{31}). \quad (65)$$

The necessary condition $|p_1^2| > \max(|p_2^2|, |p_3^2|, m^2)$, $m^2 > |p_2^2|$ should be satisfied to guarantee the multiple power series converging. The convergent region of this series is also described by the region Ω_{29}^a in Eq.(64).

- 32: For the isolated singularities defined by the poles of $\Gamma(\frac{D}{2} - 2 - s - z_2)$, $\Gamma(\frac{D}{2} - 2 - s - z_2)$, and $\Gamma(1 + s + z_1 + z_2)$, the corresponding analytic expression is zero,

$$C_{0,32}^a(p_1^2, p_2^2, p_3^2, m^2) \equiv 0. \quad (66)$$

- 33: For the isolated singularities defined by the poles of $\Gamma(\frac{D}{2} - 2 - s - z_2)$, $\Gamma(\frac{D}{2} - 2 - s - z_2)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, correspondingly the derived hypergeometric function is

$$\begin{aligned} C_{0,33}^a(p_1^2, p_2^2, p_3^2, m^2) = & -\frac{i(-)^{D/2}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}}\Gamma(2 - \frac{D}{2})\Gamma(\frac{D}{2} - 1)\sum_{j=0}^{\infty}\sum_{n_1=0}^{\infty}\sum_{n_2=0}^{\infty} \\ & \times \frac{(-)^{n_1+n_2}}{j!n_1!n_2!}\xi_{33}^{D/2-1+j+n_1+n_2}x_{13}^{-1-j-n_1}x_{23}^{-1-j-n_2} \\ & \times \frac{\Gamma(3 - \frac{D}{2} + j + n_1 + n_2)\Gamma(1 + j + n_1)\Gamma(1 + j + n_2)}{\Gamma(\frac{D}{2} + j + n_1 + n_2)\Gamma(3 - \frac{D}{2} + j)} \end{aligned} \quad (67)$$

The necessary condition $m^2 < \min(|p_1^2|, |p_2^2|)$, $|p_3^2|m^2 < |p_1^2p_2^2|$ should be satisfied to guarantee the triple series converging. The convergent region of Eq (67) is

$$\Omega_{33}^a = \left\{ (\xi_{33}, x_{13}, x_{23}) : \begin{cases} |\xi_{33}| < |x_{13}x_{23}|, |\xi_{33}| < |x_{13}|, |\xi_{33}| < |x_{23}|; \\ |\xi_{33}|(1 + |x_{13}|) < |x_{13}x_{23}|, |\xi_{33}|(1 + |x_{23}|) < |x_{13}x_{23}|, \\ |\xi_{33}| < (|x_{13}| - |\xi_{33}|)(|x_{23}| - |\xi_{33}|) \end{cases} \right\} \quad (68)$$

B. The scalar integral in different convergent regions

In order to pursue our analysis, we summarize the results presented above. The absolutely and uniformly convergent domain of the hypergeometric function $C_{0,1}^a$ is the domain Ω_1^a , that of the hypergeometric function $C_{0,2}^a$ is the domain Ω_2^a , that of the hypergeometric functions $C_{0,3}^a$, $C_{0,4}^a$ is the domain Ω_3^a , that of the hypergeometric function $C_{0,6}^a$ is the domain Ω_6^a , that of the hypergeometric function $C_{0,7}^a$ is the domain Ω_7^a , that of the hypergeometric functions $C_{0,8}^a$ and $C_{0,9}^a$ is the domain Ω_8^a , that of the hypergeometric functions $C_{0,10}^a$ and $C_{0,13}^a$ is the domain Ω_{10}^a , that of the hypergeometric functions $C_{0,11}^a$ and $C_{0,14}^a$ is the domain Ω_{11}^a , that of the hypergeometric functions $C_{0,18}^a$ and $C_{0,19}^a$ is the domain Ω_{18}^a , that of the hypergeometric functions $C_{0,23}^a$ and $C_{0,24}^a$ is the domain Ω_{23}^a , that of the hypergeometric functions $C_{0,29}^a$ and $C_{0,30}^a$ is the domain Ω_{29}^a , as well as that of the hypergeometric function $C_{0,33}^a$ is the domain Ω_{33}^a , respectively. The scalar integral of other parameter space contains some unknown

linear combination parameters originating from the non-isolated singularities. Among those convergent regions, the following domains compose a set

$$\{\Omega_7^a, \Omega_8^a, \Omega_{11}^a, \Omega_{18}^a, \Omega_{23}^a, \Omega_{29}^a\}, \quad (69)$$

whose element satisfies the following constraints simultaneously,

$$\Omega_i^a \cap \Omega_j^a = \emptyset, \quad i, j = 7, 8, 11, 18, 23, 29, \quad i \neq j. \quad (70)$$

In addition, the relations between the elements of the set and those convergent regions which do not belong to the set are

$$\begin{aligned} \Omega_7^a &\subset \Omega_{\alpha_1}^a, \quad \alpha_1 = 1, 2, 6, 33; \quad \Omega_7^a \cap \Omega_{\beta_1}^a = \emptyset, \quad \beta_1 = 3, 10; \\ \Omega_{11}^a &\subset \Omega_{\alpha_2}^a, \quad \alpha_2 = 10, 33; \quad \Omega_{11}^a \cap \Omega_{\beta_2}^a = \emptyset, \quad \beta_2 = 1, 2, 3, 6; \\ \Omega_{29}^a &\subset \Omega_{10}^a, \quad \Omega_{29}^a \cap \Omega_{\beta_3}^a = \emptyset, \quad \beta_3 = 1, 2, 3, 6, 33; \\ \Omega_8^a &\subset \Omega_{\alpha_4}^a, \quad \alpha_4 = 3, 33; \quad \Omega_8^a \cap \Omega_{\beta_4}^a = \emptyset, \quad \beta_4 = 1, 2, 6, 10; \\ \Omega_{23}^a &\subset \Omega_3^a, \quad \Omega_{23}^a \cap \Omega_{\beta_5}^a = \emptyset, \quad \beta_5 = 1, 2, 6, 10, 33; \\ \Omega_{18}^a &\cap \Omega_{\beta_6}^a = \emptyset, \quad \beta_6 = 1, 2, 3, 6, 10, 33. \end{aligned} \quad (71)$$

With the preparation above the concrete expression of C_0^a in different parameter space is respectively presented as following.

- 1: In the convergent region Ω_7^a , the scalar integral C_0^a is written as the sum

$$C_0^a(p_1^2, p_2^2, p_3^2, m^2) = \{C_{0,1}^a + C_{0,2}^a + C_{0,6}^a + C_{0,7}^a + C_{0,33}^a\}(p_1^2, p_2^2, p_3^2, m^2). \quad (72)$$

In the limit $m^2 \rightarrow 0$, the expression of Eq.(72) recovers that presented in Eq.(32). Furthermore, we expand the hypergeometric functions $C_{0,i}^a$, ($i = 1, 2, 6, 7, 33$) around $\varepsilon = 0$, and find that $D = 4$ is the pole of the second order for each hypergeometric function individually. Nevertheless the sum presented in Eq.(72) is an analytic function of the dimension in the neighborhood of $D = 4$ because those singularities are canceled clearly, and the final result is given as

$$\begin{aligned} C_0^a(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \{f_\varepsilon(x_{13}, x_{23}) \\ &\quad + 2\varepsilon \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{33}^j x_{13}^{n_1} x_{23}^{n_2} A'_{j,n_1,n_2}{}^{(a)}(x_{13}, x_{23}) \} \end{aligned}$$

$$\begin{aligned}
& \times \frac{\Gamma(j)\Gamma^2(1+j+n_1+n_2)}{j!n_1!n_2!\Gamma(1+j+n_1)\Gamma(1+j+n_2)} \\
& + 2\varepsilon \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{32}^j x_{13}^{n_1} \xi_{33}^{n_2} B_{j,n_1,n_2}^{(a)} \\
& \times \frac{\Gamma(j)\Gamma(j+n_2)\Gamma^2(1+n_1+n_2)}{n_1!n_2!\Gamma(1+j+n_1+n_2)\Gamma(1+j+n_2)} \\
& + 2\varepsilon \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{31}^j \xi_{33}^{n_1} x_{23}^{n_2} C_{j,n_1,n_2}^{(a)} \\
& \times \frac{\Gamma(j)\Gamma(j+n_1)\Gamma^2(1+n_1+n_2)}{n_1!n_2!\Gamma(1+j+n_1+n_2)\Gamma(1+j+n_1)} \\
& + \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \left(\frac{\xi_{33}}{x_{13}x_{23}} \right)^j \xi_{31}^{n_1} \xi_{32}^{n_2} \\
& \times \frac{(-)^{n_1+n_2} \Gamma(j+n_1+n_2) \Gamma(j+n_1) \Gamma(j+n_2)}{n_1!n_2! \Gamma^2(j) \Gamma(1+j+n_1+n_2)} \\
& \times \left[\psi(j+n_1) + \psi(j+n_2) - 2\psi(j) \right. \\
& \left. - \frac{1}{j+n_1+n_2} + \ln \frac{\xi_{33}}{x_{13}x_{23}} + \frac{\varepsilon}{2} D_{j,n_1,n_2}^{(a)} \right] \\
& + 2\varepsilon \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \sum_{n_2=0}^{\infty} \xi_{31}^j \xi_{32}^{n_1} \xi_{33}^{n_2} \\
& \times \frac{(-)^{j+n_1} \Gamma(j) \Gamma(j+n_1+n_2) \Gamma(n_1) \Gamma^2(1+n_2)}{\Gamma(1+j+n_1+n_2) \Gamma(1+n_1+n_2) \Gamma(1+j+n_2)} \\
& + \mathcal{O}(\varepsilon^2) \}. \tag{73}
\end{aligned}$$

with

$$\begin{aligned}
f_\varepsilon(x, y) &= \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} x^{n_1} y^{n_2} \frac{\Gamma^2(1+n_1+n_2)}{n_1!n_2!\Gamma(1+n_1)\Gamma(1+n_2)} \\
& \times \left[\ln x \ln y - 2 \ln x \psi(1+n_2) - 2 \ln y \psi(1+n_1) \right. \\
& + 2 \ln(xy) \psi(1+n_1+n_2) + 4\psi(1+n_1)\psi(1+n_2) \\
& - 4\{\psi(1+n_1) + \psi(1+n_2)\}\psi(1+n_1+n_2) \\
& \left. + 4\psi^2(1+n_1+n_2) + 2\psi^{(1)}(1+n_1+n_2) - \frac{\varepsilon}{2} A_{n_1,n_2}^{(a)}(x, y) \right]. \tag{74}
\end{aligned}$$

Here the coefficients of ε above $A_{n_1,n_2}^{(a)}(x, y)$, $A'_{j,n_1,n_2}(x, y)$, $B_{j,n_1,n_2}^{(a)}$, $C_{j,n_1,n_2}^{(a)}$, and $D_{j,n_1,n_2}^{(a)}$ can be found in Eq.(B1). The absolutely and uniformly converging region Ω_7^a is a proper subset of the cube

$$\Xi_K^a = \left\{ (\xi_{33}, x_{13}, x_{23}) : \left| \xi_{33} \right| < 1, |x_{13}| < 1, |x_{23}| < 1 \right\}. \tag{75}$$

In order to continue the scalar integral C_0^a from the region Ω_7^a to the whole cube Ξ_K^a , one can employ the system of linear PDEs in Eq.(39) with the finite element method. We will consider this point in detail in section VI.

- 2(a): In the convergent region Ω_{11}^a , the scalar integral C_0^a is written as the sum

$$\begin{aligned} C_0^a(p_1^2, p_2^2, p_3^2, m^2) &= \{C_{0,10}^a + C_{0,11}^a + C_{0,13}^a + C_{0,14}^a + C_{0,33}^a\}(p_1^2, p_2^2, p_3^2, m^2) \\ &= \frac{i(-)^{D/2}(p_1^2)^{D/2-3}}{(4\pi)^{D/2}} F_{a,p_1}(\xi_{31}, x_{31}, x_{21}). \end{aligned} \quad (76)$$

In the limit $m^2 \rightarrow 0$, the expression of Eq.(76) recovers that presented in Eq.(34). Additionally we expand the hypergeometric functions $C_{0,i}^a$, ($i = 10, 11, 13, 14, 33$) around $\varepsilon = 0$, and find that $D = 4$ is the pole of the second order for each hypergeometric function individually. Nevertheless the sum presented in Eq.(76) is an analytic function of the dimension in the neighborhood of $D = 4$, and expansion of C_0^a around $\varepsilon = 0$ is

$$\begin{aligned} C_0^a(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(p_1^2)^{D/2-3}}{(4\pi)^{D/2}} \left\{ f_\varepsilon(x_{21}, x_{31}) \right. \\ &\quad + 2\varepsilon \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{31}^j x_{31}^{n_1} x_{21}^{n_2} E_{j,n_1,n_2}^{(a)} \\ &\quad \times \frac{\Gamma(j)\Gamma(1+j+n_1+n_2)\Gamma(1+n_1+n_2)}{\Gamma(1+j)\Gamma(1+j+n_2)\Gamma^2(1+n_1)\Gamma(1+n_2)} \\ &\quad - 2\varepsilon \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \sum_{n_2=0}^{\infty} \xi_{31}^{j+n_2} x_{31}^{j+n_1} x_{21}^{-j} \\ &\quad \times \frac{(-)^{n_2}\Gamma(j)\Gamma(1+j+n_1+n_2)\Gamma(j+n_2)\Gamma(1+n_1)}{\Gamma^2(1+j+n_1)\Gamma(1+j+n_2)\Gamma(1+n_2)} \\ &\quad \times \left(\ln x_{31} + \psi(1+j+n_1+n_2) + \psi(1+n_1) \right) \\ &\quad + \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{31}^{j+n_1+n_2} x_{31}^{n_1} x_{21}^{-j-n_1} \\ &\quad \times \frac{(-)^{j+n_2}\Gamma(j+n_1+n_2)\Gamma(j+n_1)\Gamma(1+n_1+n_2)}{(j+n_1+n_2)!n_1!n_2!\Gamma(j)\Gamma(1+n_1)} \\ &\quad \times \left[\left(\frac{1}{j+n_1+n_2} - \psi(j+n_1) - \psi(1+n_1+n_2) \right) \right. \\ &\quad \left. + 2\psi(1+n_1) + \ln \frac{x_{21}}{x_{31}\xi_{31}} + \frac{\varepsilon}{2} G_{j,n_1,n_2}^{(a)} \right] + \mathcal{O}(\varepsilon^2) \left. \right\}, \end{aligned} \quad (77)$$

where the coefficients of ε above $E_{j,n_1,n_2}^{(a)}$ and $G_{j,n_1,n_2}^{(a)}$ are given in Eq.(B2) also.

- 2(b): In the convergent region Ω_{29}^a , the scalar integral C_0^a is formulated as

$$\begin{aligned} C_0^a(p_1^2, p_2^2, p_3^2, m^2) &= \{C_{0,10}^a + C_{0,13}^a + C_{0,29}^a + C_{0,30}^a\}(p_1^2, p_2^2, p_3^2, m^2) \\ &= \frac{i(-)^{D/2}(p_1^2)^{D/2-3}}{(4\pi)^{D/2}} F'_{a,p_1}(\xi_{31}, x_{21}, x_{31}). \end{aligned} \quad (78)$$

In addition, $D = 4$ is the pole of the second order for each hypergeometric function of $C_{0,i}^a$ ($i = 10, 13, 29, 30$) individually. Nevertheless the sum presented in Eq.(78) is an analytic function of the dimension in the neighborhood of $D = 4$, and expansion of C_0^a around $\varepsilon = 0$ is

$$\begin{aligned} C_0^a(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(p_1^2)^{D/2-3}}{(4\pi)^{D/2}} \left\{ f_\varepsilon(x_{21}, x_{31}) \right. \\ &\quad + \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{31}^j x_{31}^{n_1} x_{21}^{n_2} \\ &\quad \times \frac{(-)^j \Gamma(1+n_1+n_2) \Gamma(1+j+n_1+n_2) \Gamma(j)}{j! n_1! n_2! \Gamma(1+n_1) \Gamma(1+j+n_2)} \\ &\quad \times \left[\psi(1+j+n_1+n_2) + \psi(1+n_1+n_2) \right. \\ &\quad \left. - 2\psi(1+n_1) + \ln y_1 - \frac{\varepsilon}{2} H_{j,n_1,n_2}^{(a)} \right] \\ &\quad + \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=1}^{\infty} \xi_{31}^{-n_2} x_{31}^{j+n_1+n_2} x_{21}^{j+n_2} \\ &\quad \times \frac{(-)^{n_2} \Gamma(1+j+n_1+n_2) \Gamma(1+j+n_1) \Gamma(n_2)}{j! n_1! (j+n_2)! \Gamma(1+n_2) \Gamma(1+n_1)} \\ &\quad \times \left[2\psi(1+n_1) - \psi(1+j+n_1+n_2) - \psi(1+j+n_1) \right. \\ &\quad \left. - \ln y_1 + \frac{\varepsilon}{2} P_{j,n_1,n_2}^{(a)} \right] + \mathcal{O}(\varepsilon^2) \left. \right\}, \end{aligned} \quad (79)$$

where the coefficients of ε above $H_{j,n_1,n_2}^{(a)}$ and $P_{j,n_1,n_2}^{(a)}$ are presented in Eq.(B1). In addition, the functions $x_{13}^{D/2-3} F_{a,p_1}$, $x_{13}^{D/2-3} F'_{a,p_1}$ both comply with the system of linear PDEs in Eq (39). Or equivalently, the functions F_{a,p_1} , F'_{a,p_1} satisfy the following system of linear PDEs

$$\begin{aligned} &\left\{ \left(3 - \frac{D}{2} + \hat{\theta}_{\xi_{31}} + \hat{\theta}_{x_{21}} + \hat{\theta}_{x_{31}} \right) (4 - D + \hat{\theta}_{\xi_{31}}) \right. \\ &\quad \left. + \frac{1}{\xi_{31}} \hat{\theta}_{\xi_{31}} \left(2 - \frac{D}{2} + \hat{\theta}_{\xi_{31}} + \hat{\theta}_{x_{21}} \right) \right\} F_{a,p_1}(\xi_{31}, x_{21}, x_{31}) = 0, \\ &\left\{ \left(3 - \frac{D}{2} + \hat{\theta}_{\xi_{31}} + \hat{\theta}_{x_{21}} + \hat{\theta}_{x_{31}} \right) (1 + \hat{\theta}_{x_{21}} + \hat{\theta}_{x_{31}}) \right. \\ &\quad \left. - \frac{1}{x_{21}} \hat{\theta}_{x_{21}} \left(2 - \frac{D}{2} + \hat{\theta}_{\xi_{31}} + \hat{\theta}_{x_{21}} \right) \right\} F_{a,p_1}(\xi_{31}, x_{21}, x_{31}) = 0, \end{aligned}$$

$$\left\{ \left(3 - \frac{D}{2} + \hat{\theta}_{\xi_{31}} + \hat{\theta}_{x_{21}} + \hat{\theta}_{x_{31}} \right) (1 + \hat{\theta}_{x_{21}} + \hat{\theta}_{x_{31}}) - \frac{1}{x_{31}} \hat{\theta}_{x_{31}} \left(2 - \frac{D}{2} + \hat{\theta}_{x_{31}} \right) \right\} F_{a,p_1}(\xi_{31}, x_{21}, x_{31}) = 0. \quad (80)$$

Obviously the union of the regions Ω_{11}^a and Ω_{29}^a is a proper subset of the cube

$$\Xi_{p_1}^a = \left\{ (\xi_{31}, x_{21}, x_{31}) : \left| \xi_{31} \right| < 1, |x_{21}| < 1, |x_{31}| < 1 \right\}. \quad (81)$$

In order to continue the scalar integral C_0^a from the region $\Omega_{11}^a \cup \Omega_{29}^a$ to the whole cube $\Xi_{p_1}^a$, one employs the system of linear PDEs of Eq. (80).

- 3(a): In the convergent region Ω_8^a , the scalar integral C_0^a is

$$\begin{aligned} C_0^a(p_1^2, p_2^2, p_3^2, m^2) &= \left\{ C_{0,3}^a + C_{0,4}^a + C_{0,8}^a + C_{0,9}^a + C_{0,33}^a \right\} (p_1^2, p_2^2, p_3^2, m^2) \\ &= \frac{i(p_2^2)^{D/2-3}}{(4\pi)^{D/2}} F_{a,p_2}(\xi_{32}, x_{12}, x_{32}). \end{aligned} \quad (82)$$

In the limit $m^2 \rightarrow 0$, the expression of Eq.(82) recovers that presented in Eq.(35), and the expansion of C_0^a around $\varepsilon = 0$ is

$$C_0^a(p_1^2, p_2^2, p_3^2, m^2) = \text{Eq.}(77)(p_1 \rightarrow p_2, \xi_{31} \rightarrow \xi_{32}, x_{21} \rightarrow x_{12}, x_{31} \rightarrow x_{32}). \quad (83)$$

- 3(b): In the convergent region Ω_{23}^a , the scalar integral C_0^a is written as

$$\begin{aligned} C_0^a(p_1^2, p_2^2, p_3^2, m^2) &= \left\{ C_{0,3}^a + C_{0,4}^a + C_{0,23}^a + C_{0,24}^a \right\} (p_1^2, p_2^2, p_3^2, m^2) \\ &= \frac{i(p_2^2)^{D/2-3}}{(4\pi)^{D/2}} F'_{a,p_2}(\xi_{32}, x_{12}, x_{32}). \end{aligned} \quad (84)$$

Correspondingly the expansion of C_0^a around $\varepsilon = 0$ is given by

$$C_0^a(p_1^2, p_2^2, p_3^2, m^2) = \text{Eq.}(79)(p_1 \rightarrow p_2, \xi_{31} \rightarrow \xi_{32}, x_{21} \rightarrow x_{12}, x_{31} \rightarrow x_{32}). \quad (85)$$

Additionally the functions $x_2^{D/2-3} F_{a,p_2}$, $x_2^{D/2-3} F'_{a,p_2}$ both comply with the system of linear PDEs in Eq (39). Or equivalently, the functions F_{a,p_2} , F'_{a,p_2} satisfy the system which is obtained from Eq.(80) through the interchanging $1 \leftrightarrow 2$.

Obviously the union of the regions Ω_8^a and Ω_{23}^a is a proper subset of the cube

$$\Xi_{p_2}^a = \left\{ (\xi_{32}, x_{12}, x_{32}) : \left| \xi_{32} \right| < 1, |x_{12}| < 1, |x_{32}| < 1 \right\}. \quad (86)$$

In order to continue the scalar integral C_0^a from the region $\Omega_8^a \cup \Omega_{23}^a$ to the whole cube $\Xi_{p_1}^a$, one employs the correspondingly holonomic hypergeometric system of linear PDEs.

- 4: In the parameter space Ω_{18}^a , the scalar integral C_0^a is written as

$$\begin{aligned} C_0^a(p_1^2, p_2^2, p_3^2, m^2) &= \{C_{0,18}^a + C_{0,19}^a\}(p_1^2, p_2^2, p_3^2, m^2) \\ &= \frac{i(m^2)^{D/2-3}}{(4\pi)^{D/2}} F_{a,m}(\eta_{13}, \eta_{23}, \eta_{33}). \end{aligned} \quad (87)$$

The function of Eq (87) is absolutely and uniformly convergent in the connect region Ω_{18}^a . Except a convenient factor, the expression coincides with Eq (27) in the literature [13]. We expand the hypergeometric functions $C_{0,i}^a$, ($i = 18, 19$) around $\varepsilon = 0$, and find that $D = 4$ is the pole of the first order for each hypergeometric function individually. Nevertheless the sum presented in Eq.(87) is an analytic function of the dimension in the neighborhood of $D = 4$, and the expansion of C_0 around $\varepsilon = 0$ is

$$\begin{aligned} C_0^a(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(m^2)^{D/2-3}}{(4\pi)^{D/2}} \left\{ \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \eta_{33}^j \eta_{13}^{n_1} \eta_{23}^{n_2} \right. \\ &\quad \times \frac{(-)^{n_1+n_2} \Gamma(1+j+n_1+n_2) \Gamma(1+j+n_1) \Gamma(1+j+n_2)}{j! n_1! n_2! \Gamma(2+j+n_1+n_2) \Gamma(1+j)} \\ &\quad \times [2\psi(1+j) - \psi(1+j+n_1) - \psi(1+j+n_2) \\ &\quad \left. - \ln \eta_{33} - \frac{1}{1+j+n_1+n_2} + \frac{\varepsilon}{2} Q_{j,n_1,n_2}^{(a)} \right] + \mathcal{O}(\varepsilon^2) \Big\}, \end{aligned} \quad (88)$$

where the concrete expression of $Q_{j,n_1,n_2}^{(a)}$ is presented in Eq.(B1). Additionally the function $x_m^{D/2-3} F_{a,m}$ complies with the system of linear PDEs in Eq (39). Or equivalently, the function $F_{a,m}$ satisfies the following system of linear PDEs

$$\begin{aligned} &\left\{ \left(3 - \frac{D}{2} + \sum_{i=1}^3 \hat{\theta}_{\eta_{i3}} \right) (1 + \hat{\theta}_{\eta_{33}} + \hat{\theta}_{\eta_{13}}) (1 + \hat{\theta}_{\eta_{33}} + \hat{\theta}_{\eta_{23}}) \right. \\ &\quad \left. - \frac{1}{\eta_{33}} \hat{\theta}_{\eta_{33}} \left(\frac{D}{2} - 1 + \sum_{i=1}^3 \hat{\theta}_{\eta_{i3}} \right) \left(2 - \frac{D}{2} + \hat{\theta}_{\eta_{33}} \right) \right\} F_{a,m}(\eta_{13}, \eta_{13}, \eta_{23}) = 0, \\ &\left\{ \left(3 - \frac{D}{2} + \sum_{i=1}^3 \hat{\theta}_{\eta_{i3}} \right) (1 + \hat{\theta}_{\eta_{33}} + \hat{\theta}_{\eta_{j3}}) \right. \\ &\quad \left. + \frac{1}{\eta_{j3}} \hat{\theta}_{\eta_{j3}} \left(\frac{D}{2} - 1 + \sum_{i=1}^3 \hat{\theta}_{\eta_{i3}} \right) \right\} F_{a,m}(\eta_{13}, \eta_{13}, \eta_{23}) = 0, \quad (j = 1, 2). \end{aligned} \quad (89)$$

Obviously the absolutely and uniformly convergent region Ω_{18}^a is a proper subset of the cube

$$\Xi_m^a = \left\{ (\eta_{13}, \eta_{13}, \eta_{23}) : \left| \eta_{13} \right| < 1, \left| \eta_{23} \right| < 1, \left| \eta_{33} \right| < 1 \right\}. \quad (90)$$

In order to continue the scalar integral C_0^a from the region Ω_{18}^a to the whole cube Ξ_m^a , one employs the system of PDEs in Eq. (89).

V. THE C_0 FUNCTION WITH THREE EQUAL MASSES

In this case, the Mellin-Barnes representation of the scalar integral is

$$\begin{aligned}
C_0^b(p_1^2, p_2^2, p_3^2, m^2) &= \int \frac{d^D q}{(2\pi)^D} \frac{1}{[q^2 - m^2][(q + p_1)^2 - m^2][(q - p_2)^2 - m^2]} \\
&= -\frac{i(-)^{D/2-3}}{(4\pi)^{D/2}(2\pi i)^3} \int_{-i\infty}^{+i\infty} ds (-m^2)^s \Gamma(-s) \\
&\quad \times \int_{-i\infty}^{+i\infty} dz_1 (p_1^2)^{z_1} \Gamma(-z_1) \int_{-i\infty}^{+i\infty} dz_2 (p_2^2)^{z_2} \Gamma(-z_2) \\
&\quad \times (p_3^2)^{D/2-3-s-z_1-z_2} \Gamma(3 - \frac{D}{2} + s + z_1 + z_2) \\
&\quad \times \frac{\Gamma(\frac{D}{2} - 2 - s - z_2) \Gamma(\frac{D}{2} - 2 - s - z_1) \Gamma(1 + z_1 + z_2)}{\Gamma(D - 3 - 2s)}, \quad (91)
\end{aligned}$$

which is equivalent to the representation of Eq. (36) of Ref. [13], and that of Eq. (4.5) of Ref. [16]. Applying the residue theorem to the singularities of Γ functions in the numerator of integrand, we formulate the Mellin-Barnes representation as various hypergeometric functions whose absolutely and uniformly convergent regions are analyzed through Horn's study of convergence [27]. As the scalar integral is given by

$$C_0^b(p_1^2, p_2^2, p_3^2, m^2) = \frac{i(-)^{D/2}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} F_{b,k}(\xi_{m3}, x_{13}, x_{23}), \quad (92)$$

with $\xi_{mj} = \xi_{ij}$, ($i, j = 1, 2, 3$). In addition, the dimensionless function $F_{b,k}$ satisfies the system of linear PDEs

$$\begin{aligned}
&\left\{ \left(3 - \frac{D}{2} + \hat{\vartheta}_{\xi_{m3}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}}\right) (5 - D + 2\hat{\vartheta}_{\xi_{m3}}) (4 - D + 2\hat{\vartheta}_{\xi_{m3}}) \right. \\
&\quad \left. - \frac{1}{\xi_{m3}} \hat{\vartheta}_{\xi_{m3}} \prod_{i=1}^2 \left(2 - \frac{D}{2} + \hat{\vartheta}_{\xi_{m3}} + \hat{\vartheta}_{x_{i3}}\right) \right\} F_{b,p_3} = 0, \\
&\left\{ \left(3 - \frac{D}{2} + \hat{\vartheta}_{\xi_{m3}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}}\right) \left(1 + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}}\right) \right. \\
&\quad \left. - \frac{1}{x_{j3}} \hat{\vartheta}_{x_{j3}} \left(2 - \frac{D}{2} + \hat{\vartheta}_{\xi_{m3}} + \hat{\vartheta}_{x_{j3}}\right) \right\} F_{b,p_3} = 0, \quad (j = 1, 2). \quad (93)
\end{aligned}$$

Now we present our results below in detail.

A. The hypergeometric functions and their convergent regions

- 1: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(-z_1)$, and $\Gamma(-z_2)$, the corresponding hypergeometric series is written as:

$$\begin{aligned}
C_{0,1}^b(p_1^2, p_2^2, p_3^2, m^2) &= -\frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}\Gamma^2(2-\frac{D}{2})\Gamma^2(\frac{D}{2}-1)}{(4\pi)^{D/2}\Gamma(D-3)} \\
&\times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} x_{13}^{n_1} x_{23}^{n_2} \frac{\Gamma(3-\frac{D}{2}+n_1+n_2)\Gamma(1+n_1+n_2)}{n_1!n_2!\Gamma(3-\frac{D}{2}+n_2)\Gamma(3-\frac{D}{2}+n_1)} \\
&-\frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}\Gamma^2(2-\frac{D}{2})\Gamma^2(\frac{D}{2}-1)}{(4\pi)^{D/2}\Gamma(D-3)\Gamma(4-D)} \\
&\times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m3}^j x_{13}^{n_1} x_{23}^{n_2} \Gamma(4-D+2j) \\
&\times \frac{\Gamma(3-\frac{D}{2}+j+n_1+n_2)\Gamma(1+n_1+n_2)}{j!n_1!n_2!\Gamma(3-\frac{D}{2}+j+n_2)\Gamma(3-\frac{D}{2}+j+n_1)}. \tag{94}
\end{aligned}$$

Here the condition $|p_3^2| > \max(m^2, |p_1^2|, |p_2^2|)$ should be satisfied to guarantee the convergence of hypergeometric function. In addition, $D = 4$ is the second order pole of the first double hypergeometric function, and the first order pole of the second triple hypergeometric function, respectively. Certainly the expression of Eq. (94) complies with the system of linear PDEs in Eq. (93).

The absolutely and uniformly convergent region of the hypergeometric function is

$$\Omega_1^b = \left\{ (\xi_{m3}, x_{13}, x_{23}) : \begin{cases} |\xi_{m3}| < 1/4, \\ \sqrt{|x_{13}|} + \sqrt{|x_{23}|} < 1 \end{cases} \right\}. \tag{95}$$

- 2: When the isolated singularities are defined by the poles of $\Gamma(-s)$, $\Gamma(-z_1)$, and $\Gamma(\frac{D}{2}-2-s-z_2)$, the derived hypergeometric series $C_{0,2}^b$ is given by Eq. (A4), where the necessary condition $|p_3^2| > \max(m^2, |p_1^2|, |p_2^2|)$, $|p_2^2| > m^2$ should be met to guarantee the hypergeometric series converging. By implementing the residue theorem to the Mellin-Barnes representation, one derives the first expression. The second expression decomposes the summation indices of the hypergeometric series in the original expression, so $D = 4$ is the pole of the same fixed order for each term of every hypergeometric series of Eq. (A4). Furthermore, the decomposition of the summation indices is correct and reasonable because the second expression also satisfies the system of linear

PDEs in Eq. (93). Horn's study of convergence predicts the absolutely and uniformly convergent region of the double hypergeometric functions as

$$\Omega_2^b = \left\{ (\xi_{m3}, x_{13}, x_{23}) : \left| \begin{array}{l} |\xi_{m3}x_{13}| < |x_{23}|, |\xi_{m3}| < |x_{23}|/4, \\ \sqrt{|x_{13}|} + \sqrt{|x_{23}|} < 1 \end{array} \right. \right\}. \quad (96)$$

- 3: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(-z_1)$, and $\Gamma(1+z_1+z_2)$, the corresponding hypergeometric series $C_{0,3}^b$ is

$$C_{0,3}^b(p_1^2, p_2^2, p_3^2, m^2) = C_{0,2}^b(p_3 \rightarrow p_2, \xi_{m3} \rightarrow \xi_{m2}, x_{13} \rightarrow x_{12}, x_{23} \rightarrow x_{32}), \quad (97)$$

where the condition $|p_2^2| > \max(m^2, |p_1^2|, |p_3^2|)$, $|p_3^2| > m^2$ is necessary to guarantee the convergence of the hypergeometric functions. The absolutely and uniformly convergent region of Eq.(97) is similarly given as

$$\Omega_3^b = \left\{ (\xi_{m2}, x_{12}, x_{32}) : \left| \begin{array}{l} |\xi_{m2}x_{12}| < |x_{32}|, |\xi_{m2}| < |x_{32}|/4, \\ \sqrt{|x_{12}|} + \sqrt{|x_{32}|} < 1 \end{array} \right. \right\}. \quad (98)$$

- 4: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(-z_1)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, the corresponding hypergeometric series is

$$C_{0,4}^b(p_1^2, p_2^2, p_3^2, m^2) = C_{0,1}^b(p_3 \rightarrow p_2, \xi_{m3} \rightarrow \xi_{m2}, x_{13} \rightarrow x_{12}, x_{23} \rightarrow x_{32}). \quad (99)$$

The condition $|p_2^2| > \max(m^2, |p_1^2|, |p_3^2|)$ is necessary to guarantee the hypergeometric functions converging. In addition, the absolutely and uniformly convergent region of the term is

$$\Omega_4^b = \left\{ (\xi_{m2}, x_{12}, x_{32}) : \left| \begin{array}{l} |\xi_{m2}| < 1/4, \\ \sqrt{|x_{12}|} + \sqrt{|x_{32}|} < 1 \end{array} \right. \right\}. \quad (100)$$

- 6: When the isolated singularities are produced by the poles of $\Gamma(-s)$, $\Gamma(-z_2)$, and $\Gamma(\frac{D}{2} - 2 - s - z_1)$, the corresponding hypergeometric series $C_{0,6}^b$ is presented in Eq. (A5), where the condition $|p_3^2| > \max(m^2, |p_1^2|, |p_2^2|)$, $|p_1^2| > m^2$ should be satisfied to guarantee the convergence of the hypergeometric functions. Applying the residue theorem to the Mellin-Barnes representation, one derives the first expression of Eq. (A5). The second expression decomposes the summation indices of the hypergeometric series in the original expression, so $D = 4$ is the pole of the same fixed order for each term

of every hypergeometric series of the second equation in Eq. (A5). In addition, the decomposition of the summation indices is correct and reasonable because the second expression also complies with the system of linear PDEs in Eq. (93).

The concretely convergent region of the hypergeometric functions is given by

$$\Omega_6^b = \left\{ (\xi_{m_3}, x_{13}, x_{23}) : \left| \begin{array}{l} |\xi_{m_3} x_{23}| < |x_{13}|, |\xi_{m_3}| < |x_{13}|/4, \\ \sqrt{|x_{13}|} + \sqrt{|x_{23}|} < 1 \end{array} \right. \right\}. \quad (101)$$

- 7: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(\frac{D}{2} - 2 - s - z_2)$, the corresponding hypergeometric series $C_{0,7}^b$ is written in Eq. (A6), where the condition $|p_3^2| > \max(|p_1^2|, |p_2^2|)$, $\min(|p_1^2|, |p_2^2|) > m^2$, $|p_1^2 p_2^2| > |p_3^2| m^2$ is essential to guarantee the convergence of the hypergeometric functions. Applying the residue theorem to the Mellin-Barnes representation, one derives the first expression of Eq. (A6). The second expression decomposes the summation indices of the hypergeometric series in the original expression, so $D = 4$ is the pole of the same fixed order for each term of every hypergeometric series of the second equation in Eq. (A6). In addition, the decomposition of the summation indices is correct and reasonable because the subsequent expression also satisfies the system of linear PDEs in Eq. (93). Correspondingly the absolutely and uniformly convergent region of this hypergeometric function is concretely written as

$$\Omega_7^b = \left\{ (\xi_{m_3}, x_{13}, x_{23}) : \left| \begin{array}{l} |\xi_{m_3} x_{13}| < |x_{23}|, |\xi_{m_3} x_{23}| < |x_{13}|, \\ |\xi_{m_3}| < |x_{13} x_{23}|, |\xi_{m_3}| < |x_{13}|/4, \\ |\xi_{m_3}| < |x_{23}|/4, \sqrt{|x_{13}|} + \sqrt{|x_{23}|} < 1 \end{array} \right. \right\}. \quad (102)$$

- 8: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(1 + z_1 + z_2)$, the corresponding hypergeometric series $C_{0,8}^b$ is presented as

$$C_{0,8}^b(p_1^2, p_2^2, p_3^2, m^2) = C_{0,7}^b(p_3 \rightarrow p_2, \xi_{m_3} \rightarrow \xi_{m_2}, x_{13} \rightarrow x_{12}, x_{23} \rightarrow x_{32}). \quad (103)$$

Applying Horn's study of convergence, we write the absolutely and uniformly convergent region of the hypergeometric functions in Eq.(103) as

$$\Omega_8^b = \left\{ (\xi_{m_2}, x_{12}, x_{32}) : \left| \begin{array}{l} |\xi_{m_2}| < |x_{12} x_{32}|, |\xi_{m_2} x_{32}| < |x_{12}|, \\ |\xi_{m_2} x_{12}| < |x_{32}|, |\xi_{m_2}| < |x_{12}|/4, \\ |\xi_{m_2}| < |x_{32}|/4, \sqrt{|x_{12}|} + \sqrt{|x_{32}|} < 1 \end{array} \right. \right\}. \quad (104)$$

- 9: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, the corresponding hypergeometric series $C_{0,9}^b$ is

$$C_{0,9}^b(p_1^2, p_2^2, p_3^2, m^2) = C_{0,6}^b(p_3 \rightarrow p_2, \xi_{m3} \rightarrow \xi_{m2}, x_{13} \rightarrow x_{12}, x_{23} \rightarrow x_{32}). \quad (105)$$

The convergent region of those hypergeometric functions is concretely written as

$$\Omega_9^b = \left\{ (\xi_{m2}, x_{12}, x_{32}) : \begin{array}{l} |\xi_{m2}x_{32}| < |x_{12}|, |\xi_{m2}| < |x_{12}|/4, \\ \sqrt{|x_{12}|} + \sqrt{|x_{32}|} < 1 \end{array} \right\}. \quad (106)$$

- 10: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(1 + z_1 + z_2)$, and $\Gamma(-z_2)$, correspondingly the hypergeometric function $C_{0,10}^b$ is written as

$$C_{0,10}^b(p_1^2, p_2^2, p_3^2, m^2) = C_{0,6}^b(p_3 \rightarrow p_1, \xi_{m3} \rightarrow \xi_{m1}, x_{13} \rightarrow x_{31}, x_{23} \rightarrow x_{21}). \quad (107)$$

where the necessary condition $|p_1^2| > \max(m^2, |p_2^2|, |p_3^2|)$, $|p_3^2| > m^2$ should be satisfied to guarantee the convergence of the hypergeometric functions. Additionally the expression satisfies the system of linear PDEs in Eq. (93) also.

Through Horn's study of convergence, the convergent region of the series is similarly given by

$$\Omega_{10}^b = \left\{ (\xi_{m1}, x_{21}, x_{31}) : \begin{array}{l} |\xi_{m1}x_{21}| < |x_{31}|, |\xi_{m1}| < |x_{31}|/4, \\ \sqrt{|x_{21}|} + \sqrt{|x_{31}|} < 1 \end{array} \right\}. \quad (108)$$

- 11: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(1 + z_1 + z_2)$, and $\Gamma(\frac{D}{2} - 2 - s - z_2)$, the hypergeometric function $C_{0,11}^b$ is

$$C_{0,11}^b(p_1^2, p_2^2, p_3^2, m^2) = C_{0,7}^b(p_3 \rightarrow p_1, \xi_{m3} \rightarrow \xi_{m1}, x_{13} \rightarrow x_{31}, x_{23} \rightarrow x_{21}). \quad (109)$$

Here the necessary condition $|p_1^2| > \max(|p_2^2|, |p_3^2|)$, $|p_2^2p_3^2| > |p_1^2|m^2$ should be met to guarantee the hypergeometric functions converging. Additionally the expression complies with the system of linear PDEs in Eq. (93) also.

Using Horn's study of convergence, one formulates the absolutely and uniformly convergent region of Eq.(109) as

$$\Omega_{11}^b = \left\{ (\xi_{m1}, x_{21}, x_{31}) : \begin{array}{l} |\xi_{m1}| < |x_{21}x_{31}|, |\xi_{m1}x_{31}| < |x_{21}|, \\ |\xi_{m1}x_{21}| < |x_{31}|, |\xi_{m1}| < |x_{21}|/4, \\ |\xi_{m1}| < |x_{31}|/4, \sqrt{|x_{21}|} + \sqrt{|x_{31}|} < 1 \end{array} \right\}. \quad (110)$$

- 13: For the isolated singularities defined by the poles of $\Gamma(-s)$, $\Gamma(-z_2)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, the corresponding hypergeometric series is given as

$$C_{0,13}^b(p_1^2, p_2^2, p_3^2, m^2) = C_{0,1}^b(p_3 \rightarrow p_1, \xi_{m3} \rightarrow \xi_{m1}, x_{13} \rightarrow x_{31}, x_{23} \rightarrow x_{21}) . \quad (111)$$

Similarly the convergent region of the term is

$$\Omega_{13}^b = \left\{ (\xi_{m1}, x_{21}, x_{31}) : \left| |\xi_{m1}| < 1/4, \sqrt{|x_{21}|} + \sqrt{|x_{31}|} < 1 \right. \right\} . \quad (112)$$

- 14: When the isolated singularities are produced by the poles of $\Gamma(-s)$, $\Gamma(\frac{D}{2} - 2 - s - z_2)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, the corresponding hypergeometric series $C_{0,14}^b$ is presented by

$$C_{0,14}^b(p_1^2, p_2^2, p_3^2, m^2) = C_{0,2}^b(p_3 \rightarrow p_1, \xi_{m3} \rightarrow \xi_{m1}, x_{13} \rightarrow x_{31}, x_{23} \rightarrow x_{21}) , \quad (113)$$

where the necessary condition $|p_1^2| > \max(|p_2^2|, |p_3^2|)$, $|p_2^2| > m^2$ should be satisfied to guarantee the convergence of the hypergeometric functions. Similarly the absolutely and uniformly convergent region of the series is concretely written as

$$\Omega_{14}^b = \left\{ (\xi_{m1}, x_{21}, x_{31}) : \left| \begin{array}{l} |\xi_{m1} x_{31}| < |x_{21}|, |\xi_{m1}| < |x_{21}|/4, \\ \sqrt{|x_{21}|} + \sqrt{|x_{31}|} < 1 \end{array} \right. \right\} . \quad (114)$$

- 16: For the singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(-z_2)$, and $\Gamma(\frac{D}{2} - 2 - s - z_1)$, the scalar integral contains some unknown linear combination parameters originating from the non-isolated singularities. The necessary condition $|p_3^2| > \max(|p_1^2|, |p_2^2|, m^2)$, $m^2 > |p_1^2|$ should be satisfied to guarantee the hypergeometric series converging.
- 17: For the singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(-z_2)$, and $\Gamma(\frac{D}{2} - 2 - s - z_2)$, the scalar integral contains some unknown linear combination parameters originating from the non-isolated singularities. The condition $|p_3^2| > \max(|p_1^2|, |p_2^2|, m^2)$, $m^2 > |p_2^2|$ should be satisfied to guarantee the convergence of the hypergeometric series.
- 19: For the isolated singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(-z_2)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, the corresponding hypergeometric series is given as

$$C_{0,19}^b(p_1^2, p_2^2, p_3^2, m^2) = -\frac{i(m^2)^{D/2-3}}{(4\pi)^{D/2}} \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \eta_{3m}^j \eta_{1m}^{n_1} \eta_{2m}^{n_2}$$

$$\begin{aligned} & \times \frac{\Gamma(1+j+n_1)\Gamma(1+j+n_2)\Gamma(1+n_1+n_2)}{j!n_1!n_2!} \\ & \times \frac{\Gamma(3-\frac{D}{2}+j+n_1+n_2)}{\Gamma(3+2j+2n_1+2n_2)}. \end{aligned} \quad (115)$$

The necessary condition $4m^2 > \max(|p_1^2|, |p_2^2|, |p_3^2|)$ should be met to guarantee the convergence of the triple series. In addition, $D = 4$ is the first order pole of the hypergeometric function. Certainly the expression of Eq. (115) complies with the system of linear PDEs in Eq. (93).

Correspondingly the convergent region of the term is

$$\Omega_{19}^b = \left\{ (\eta_{1m}, \eta_{2m}, \eta_{3m}) : \left| \eta_{1m} \right| < 4, \left| \eta_{2m} \right| < 4, \left| \eta_{3m} \right| < 4 \right\}. \quad (116)$$

- 20: For the singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(\frac{D}{2} - 2 - s - z_2)$, and $\Gamma(\frac{D}{2} - 2 - s - z_1)$, the scalar integral contains some unknown linear combination parameters originating from the non-isolated singularities. The condition $|p_3^2| > \max(|p_1^2|, |p_2^2|, m^2)$, $|p_2^2| > m^2$, $|p_3^2|m^2 > |p_1^2p_2^2|$ should be satisfied to guarantee the convergence of the hypergeometric series.
- 21: When the isolated singularities are produced by the poles of $\Gamma(-z_1)$, $\Gamma(\frac{D}{2} - 2 - s - z_2)$, and $\Gamma(1 + z_1 + z_2)$, the corresponding hypergeometric series is given as

$$\begin{aligned} C_{0,21}^b(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \left\{ \Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1) \right\} \frac{\xi_{m3}^{D/2-1}}{x_{23}} \\ & \times \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m3}^{j+n_1+n_2} x_{13}^{n_1} x_{23}^{-n_1-n_2} \Gamma(1+j+n_1) \\ & \times \frac{(-)^{n_1} \Gamma(1+n_1+n_2) \Gamma(2+2j+2n_1+2n_2)}{j!n_1!n_2! \Gamma(\frac{D}{2}+j+n_1+n_2) \Gamma(2+j+2n_1+n_2)} \end{aligned} \quad (117)$$

The condition $\min(|p_2^2|, |p_3^2|) > m^2$, $|p_2^2p_3^2| > |p_1^2|m^2$ is necessary to guarantee the hypergeometric series converging. In addition, $D = 4$ is the first order pole of the hypergeometric function. Certainly the expression of Eq. (117) satisfies the system of linear PDEs in Eq. (93).

Correspondingly the convergent region of the series is

$$\Omega_{21}^b = \left\{ (\xi_{m3}, x_{13}, x_{23}) : \left| \xi_{m3} \right| < 1/4, \left| \xi_{m3}x_{13} \right| < |x_{23}|, \left| \xi_{m3} \right| < |x_{23}|/4 \right\}. \quad (118)$$

- 23: For the singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(1 + z_1 + z_2)$, the scalar integral contains some unknown linear combination parameters originating from the non-isolated singularities. The condition $|p_3^2| > \max(|p_2^2|, m^2)$, $|p_3^2|^2 > |p_1^2|m^2$, $|p_3^2|m^2 > |p_1^2p_2^2|$ should be satisfied to guarantee the convergence of the hypergeometric series.
- 24: For the singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, the scalar integral contains some unknown linear combination parameters originating from the non-isolated singularities. The condition $|p_2^2| > \max(|p_1^2|, |p_3^2|, m^2)$, $m^2 > |p_1^2|$ should be satisfied to guarantee the convergence of the hypergeometric series.
- 25: For the singularities defined by the poles of $\Gamma(-z_1)$, $\Gamma(1 + z_1 + z_2)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, the scalar integral contains some unknown linear combination parameters originating from the non-isolated singularities. The necessary condition $|p_2^2| > \max(|p_1^2|, |p_3^2|, m^2)$, $m^2 > |p_3^2|$ should be satisfied to guarantee the convergence of the hypergeometric series.
- 26: For the singularities defined by the poles of $\Gamma(-z_2)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(\frac{D}{2} - 2 - s - z_2)$, the scalar integral contains some unknown linear combination parameters originating from the non-isolated singularities. The condition $|p_3^2| > \max(|p_1^2|, |p_2^2|, m^2)$, $|p_1^2| > m^2$, $|p_3^2|m^2 > |p_1^2p_2^2|$ should be satisfied to guarantee the convergence of the hypergeometric series.
- 27: For the isolated singularities defined by the pole of $\Gamma(-z_2)$, $\Gamma(\frac{D}{2} - 2 - s - z_1)$, and $\Gamma(1 + z_1 + z_2)$, the corresponding hypergeometric series is

$$\begin{aligned}
C_{0,27}^b(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \left\{ \Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1) \right\} \frac{\xi_{m3}^{D/2-1}}{x_{13}} \\
&\times \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m3}^{j+n_1+n_2} x_{23}^{n_2} x_{13}^{-n_1-n_2} \Gamma(1 + j + n_2) \\
&\times \frac{(-)^{n_2} \Gamma(1 + n_1 + n_2) \Gamma(2 + 2j + 2n_1 + 2n_2)}{j! n_1! n_2! \Gamma(\frac{D}{2} + j + n_1 + n_2) \Gamma(2 + j + n_1 + 2n_2)}. \quad (119)
\end{aligned}$$

The condition $\min(|p_1^2|, |p_3^2|) > m^2$, $|p_1^2p_3^2| > |p_2^2|m^2$ should be satisfied to guarantee the convergence of the hypergeometric series. In addition, $D = 4$ is the first order pole

of the hypergeometric function. Certainly the expression of Eq. (119) complies with the system of PDEs in Eq. (93).

Correspondingly the convergent region of the series is

$$\Omega_{27}^b = \left\{ (\xi_{m3}, x_{13}, x_{23}) : \left| \xi_{m3} \right| < 1/4, \left| \xi_{m3} x_{23} \right| < |x_{13}|, \left| \xi_{m3} \right| < |x_{13}|/4 \right\}. \quad (120)$$

- 29: For the singularities defined by the poles of $\Gamma(-z_2)$, $\Gamma(\frac{D}{2}-2-s-z_2)$, and $\Gamma(1+z_1+z_2)$, the scalar integral contains some unknown linear combination parameters originating from the non-isolated singularities. The essential condition $|p_3^2| > \max(|p_1^2|, m^2)$, $|p_3^2|^2 > |p_2^2|m^2$, $|p_3^2|m^2 > |p_1^2p_2^2|$ should be satisfied to guarantee the convergence of the hypergeometric series.
- 30: For the singularities defined by the poles of $\Gamma(-z_2)$, $\Gamma(\frac{D}{2}-2-s-z_2)$, and $\Gamma(3-\frac{D}{2}+s+z_1+z_2)$, the scalar integral contains some unknown linear combination parameters originating from the non-isolated singularities. The condition $|p_1^2| > \max(|p_2^2|, |p_3^2|, m^2)$, $m^2 > |p_2^2|$ should be satisfied to guarantee the convergence of the hypergeometric series.
- 31: For the singularities defined by the poles of $\Gamma(-z_2)$, $\Gamma(1+z_1+z_2)$, and $\Gamma(3-\frac{D}{2}+s+z_1+z_2)$, the scalar integral contains some unknown linear combination parameters originating from the non-isolated singularities. The condition $|p_1^2| > \max(|p_2^2|, |p_3^2|, m^2)$, $m^2 > |p_3^2|$ should be satisfied to guarantee the convergence of the hypergeometric series.
- 32: For the singularities defined by the poles of $\Gamma(\frac{D}{2}-2-s-z_1)$, $\Gamma(\frac{D}{2}-2-s-z_2)$, and $\Gamma(1+z_1+z_2)$, the scalar integral contains some unknown linear combination parameters originating from the non-isolated singularities. The necessary condition $|p_3^2| > \max(|p_1^2|, |p_2^2|, m^2)$, $|p_1^2p_3^2| > |p_2^2|m^2$, $|p_2^2p_3^2| > |p_1^2|m^2$, $|p_3^2|m^2 > |p_1^2p_2^2|$ should be satisfied to guarantee the convergence of the hypergeometric series.
- 33: When the isolated singularities are produced by the poles of $\Gamma(D/2-2-s-z_1)$, $\Gamma(D/2-2-s-z_2)$, and $\Gamma(3-\frac{D}{2}+s+z_1+z_2)$, the corresponding hypergeometric series is given as

$$C_{0,33}^b(p_1^2, p_2^2, p_3^2, m^2) = \frac{i(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \left\{ \Gamma(2-\frac{D}{2})\Gamma(\frac{D}{2}-1) \right\} \frac{\xi_{m3}^{D/2-1}}{x_{13}x_{23}}$$

$$\begin{aligned} & \times \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m_3}^{j+n_1+n_2} x_{13}^{-j-n_2} x_{23}^{-j-n_1} \Gamma(1+j+n_1) \\ & \times \frac{(-)^j \Gamma(1+j+n_2) \Gamma(2+2j+2n_1+2n_2)}{j! n_1! n_2! \Gamma(\frac{D}{2}+j+n_1+n_2) \Gamma(2+2j+n_1+n_2)}. \end{aligned} \quad (121)$$

The necessary condition $\min(|p_1^2|, |p_2^2|) > m^2$, $|p_1^2 p_2^2| > |p_3^2| m^2$ should be met to guarantee the convergence of the hypergeometric series. In addition, $D = 4$ is the first order pole of the hypergeometric function. Certainly the expression of Eq. (121) complies with the system of linear PDEs in Eq. (93).

The convergent region is apparently written as

$$\Omega_{33}^b = \left\{ (\xi_{m_3}, x_{13}, x_{23}) : \left| \begin{aligned} & |\xi_{m_3}| < |x_{13} x_{23}|, \quad |\xi_{m_3}| < |x_{13}|/4, \quad |\xi_{m_3}| < |x_{23}|/4 \end{aligned} \right. \right\} \quad (122)$$

- 34: For the singularities defined by the poles of $\Gamma(D/2 - 2 - s - z_1)$, $\Gamma(1 + z_1 + z_2)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, the scalar integral contains some undefined linear combination parameters originating from the non-isolated singularities. The necessary condition $|p_2^2| > \max(|p_1^2|, |p_3^2|, m^2)$, $|p_1^2 p_2^2| > |p_3^2| m^2$, $|p_1^2| > m^2$ should be satisfied to guarantee the convergence of the hypergeometric series.
- 35: For the singularities defined by the poles of $\Gamma(D/2 - 2 - s - z_2)$, $\Gamma(1 + z_1 + z_2)$, and $\Gamma(3 - \frac{D}{2} + s + z_1 + z_2)$, the scalar integral contains some undefined linear combination parameters originating from the non-isolated singularities. The necessary condition $|p_1^2| > \max(|p_2^2|, |p_3^2|, m^2)$, $|p_1^2 p_2^2| > |p_3^2| m^2$, $|p_2^2| > m^2$ should be satisfied to guarantee the convergence of the hypergeometric series.

B. The scalar integral in different convergent regions

The absolutely and uniformly convergent region of the hypergeometric function $C_{0,\alpha}$ is the domain Ω_α^b , $\alpha = 1, \dots, 4, 6, \dots, 11, 13, 14, 19, 21, 27, 33$. The scalar integral of other parameter space contains some unknown linear combination parameters originating from the non-isolated singularities. Among those convergent regions, the following domains compose a set

$$\left\{ \Omega_7^b, \Omega_8^b, \Omega_{11}^b, \Omega_{19}^b \right\}, \quad (123)$$

whose element satisfies the following constraints simultaneously,

$$\Omega_i^b \cap \Omega_j^b = \emptyset, \quad i, j = 7, 8, 11, 19, \quad i \neq j. \quad (124)$$

Meanwhile, the relations between the elements of the set and those convergent regions which do not belong to the set are

$$\begin{aligned} \Omega_7^b &\subset \Omega_{\alpha_1}^b, \quad \alpha_1 = 1, 2, 6, 7, 21, 27, 33; \\ \Omega_7^b \cap \Omega_{\beta_1}^b &= \emptyset, \quad \beta_1 = 3, 4, 8, 9, 10, 11, 13, 14, 19; \\ \Omega_8^b &\subset \Omega_{\alpha_2}^b, \quad \alpha_2 = 3, 4, 8, 9, 21, 27, 33; \\ \Omega_8^b \cap \Omega_{\beta_2}^b &= \emptyset, \quad \beta_2 = 1, 2, 6, 7, 10, 11, 13, 14, 19; \\ \Omega_{11}^b &\subset \Omega_{\alpha_3}^b, \quad \alpha_3 = 10, 11, 13, 14, 21, 27, 33; \\ \Omega_{11}^b \cap \Omega_{\beta_3}^b &= \emptyset, \quad \beta_3 = 1, 2, 3, 4, 6, 7, 8, 9, 19; \\ \Omega_{19}^b \cap \Omega_{\beta_4}^b &= \emptyset, \quad \beta_4 = 1, \dots, 4, 6, \dots, 11, 13, 14, 21, 27, 33. \end{aligned} \quad (125)$$

With the preparation above the concrete expression of C_0 in different parameter space is respectively presented as following.

- 1: In the convergent region Ω_7^b , the scalar integral C_0^b is written as

$$\begin{aligned} C_0^b(p_1^2, p_2^2, p_3^2, m^2) &= \{C_{0,1}^b + C_{0,2}^b + C_{0,6}^b + C_{0,7}^b + C_{0,21}^b \\ &\quad + C_{0,27}^b + C_{0,33}^b\}(p_1^2, p_2^2, p_3^2, m^2) \end{aligned} \quad (126)$$

In the limit $m^2 \rightarrow 0$, the expression of Eq.(126) recovers that presented in Eq.(32). Furthermore, $D = 4$ is the pole of the second order for each hypergeometric function individually in the above expression. Nevertheless the scalar integral presented in Eq.(126) is an analytic function of the dimension in the neighborhood of $D = 4$ because those singularities are canceled clearly, and the final result is given as

$$\begin{aligned} C_0^b(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \{f_\varepsilon(x_{13}, x_{23}) \\ &\quad + 2\varepsilon \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m^3}^j x_{13}^{n_1} x_{23}^{n_2} A_{j,n_1,n_2}^{(b)} \\ &\quad \times \frac{\Gamma(2j)\Gamma(1+j+n_1+n_2)\Gamma(1+n_1+n_2)}{j!n_1!n_2!\Gamma(1+j+n_1)\Gamma(1+j+n_2)} \} \end{aligned}$$

$$\begin{aligned}
& + \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m_3}^{j+n_1+n_2} x_{13}^{n_1} x_{23}^{-j-n_1} \\
& \times \frac{(-)^{n_1} \Gamma(1+n_1+n_2) \Gamma(2j+2n_1+2n_2) \Gamma(j+n_1)}{(j+n_1+n_2)! n_1! n_2! \Gamma(1+j+2n_1+n_2) \Gamma(j)} \\
& \times \left[\left(-2\psi(2j+2n_1+2n_2) - \psi(j+n_1) \right. \right. \\
& \left. \left. + \psi(1+j+n_1+n_2) + 2\psi(1+j+2n_1+n_2) + \psi(1+n_1) \right. \right. \\
& \left. \left. - \psi(1+n_1+n_2) - \ln \frac{x_{13} \xi_{m_3}}{x_{23}} \right) + \frac{\varepsilon}{2} B_{j,n_1,n_2}^{(b)} \right] \\
& + \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m_3}^{j+n_1+n_2} x_{13}^{-j-n_2} x_{23}^{n_2} \\
& \times \frac{(-)^{n_2} \Gamma(1+n_1+n_2) \Gamma(2j+2n_1+2n_2) \Gamma(j+n_2)}{(j+n_1+n_2)! n_1! n_2! \Gamma(1+j+n_1+2n_2) \Gamma(j)} \\
& \times \left[\left(-2\psi(2j+2n_1+2n_2) - \psi(j+n_2) \right. \right. \\
& \left. \left. + \psi(1+j+n_1+n_2) + 2\psi(1+j+n_1+2n_2) + \psi(1+n_2) \right. \right. \\
& \left. \left. - \psi(1+n_1+n_2) - \ln \frac{x_{23} \xi_{m_3}}{x_{13}} \right) + \frac{\varepsilon}{2} C_{j,n_1,n_2}^{(b)} \right] \\
& + \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m_3}^{j+n_1+n_2} x_{13}^{-j-n_2} x_{23}^{-j-n_1} \\
& \times \frac{(-)^j \Gamma(j+n_2) \Gamma(j+n_1) \Gamma(2j+2n_1+2n_2)}{\Gamma(j) n_1! n_2! \Gamma(1+j+n_1+n_2) \Gamma(2j+n_1+n_2)} \\
& \times \left[\left(2\psi(2j+2n_1+2n_2) - 2\psi(2j+n_1+n_2) - \psi(j) \right. \right. \\
& \left. \left. + \psi(j+n_1) + \psi(j+n_2) - \psi(1+j+n_1+n_2) \right. \right. \\
& \left. \left. - \ln \frac{x_{13} x_{23}}{\xi_{m_3}} \right) + \frac{\varepsilon}{2} D_{j,n_1,n_2}^{(b)} \right] \\
& + \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \xi_m^{j+n_1+n_2} x_{13}^{-j} x_{23}^{-n_1} \\
& \times \frac{\Gamma(2j+2n_1+2n_2) \Gamma(1+n_2) \Gamma(j) \Gamma(n_1)}{(j+n_1+n_2)! (n_1+n_2)! (j+n_2)! \Gamma(j+n_1)} \\
& \times \left[1 + \varepsilon \left(2\psi(2j+2n_1+2n_2) - 2\psi(j+n_1) \right. \right. \\
& \left. \left. - \psi(1+n_2) + \psi(j) + \psi(n_1) - \ln(x_{13} x_{23}) \right) \right] \\
& - 2\varepsilon \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \sum_{n_2=0}^{\infty} \xi_m^{j+n_2} x_{13}^{j+n_1} x_{23}^{-j} \\
& \times \frac{\Gamma(2j+2n_2) \Gamma(j) \Gamma(1+j+n_1+n_2) \Gamma(1+n_1)}{(2j+n_1+n_2)! (j+n_1)! (j+n_2)! n_2!} \\
& \times \left(\ln x_{13} - \psi(1+2j+n_1+n_2) - \psi(1+j+n_1) \right)
\end{aligned}$$

$$\begin{aligned}
& +\psi(1+j+n_1+n_2) + \psi(1+n_2)) \\
& -2\varepsilon \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=1}^{\infty} \xi_m^{j+n_2} x_{13}^{-j} x_{23}^{j+n_2} \\
& \times \frac{\Gamma(2j+2n_1)\Gamma(j)\Gamma(1+j+n_1+n_2)\Gamma(1+n_2)}{(2j+n_1+n_2)!(j+n_1)!(j+n_2)!n_1!} \\
& \times \left(\ln x_{23} - \psi(1+2j+n_1+n_2) - \psi(1+j+n_2) \right. \\
& \left. +\psi(1+j+n_1+n_2) + \psi(1+n_1) \right) \\
& +2\varepsilon \sum_{j=0}^{\infty} \sum_{n_1=1}^{\infty} \xi_{m3}^{j+n_1} x_{13}^{n_1} x_{23}^{-n_1} \frac{(-)^{n_1} \Gamma(2j+2n_1)\Gamma(n_1)}{n_1!j!\Gamma(1+j+2n_1)} E_{j,n_1}^{(b)}(x_{13}) \\
& +2\varepsilon \sum_{j=0}^{\infty} \sum_{n_2=1}^{\infty} \xi_{m3}^{j+n_2} x_{13}^{-n_2} x_{23}^{n_2} \frac{(-)^{n_2} \Gamma(2j+2n_2)\Gamma(n_2)}{n_2!j!\Gamma(1+j+2n_2)} E_{j,n_2}^{(b)}(x_{23}) \\
& + \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} x_m^{j+n_1} x_1^{-j} x_2^{-n_1} \frac{\Gamma(2j+2n_1)\Gamma(j)\Gamma(n_1)}{j!n_1!(j+n_1)!\Gamma(j+n_1)} \\
& \times \left[1 + G_{j,n_1}^{(b)}(x_{13}x_{23}) \right] \\
& + \mathcal{O}(\varepsilon^2) \}, \tag{127}
\end{aligned}$$

where the coefficients of ε above $A_{j,n_1,n_2}^{(b)}$, $B_{j,n_1,n_2}^{(b)}$, $C_{j,n_1,n_2}^{(b)}$, $D_{j,n_1,n_2}^{(b)}$, $E_{j,n}^{(b)}(x)$, and $G_{j,n}^{(b)}(x)$ can be found in Eq.(B2). The absolutely and uniformly converging region Ω_7^b is a proper subset of the cube

$$\Xi_K^b = \left\{ (\xi_{m3}, x_{13}, x_{23}) : \left| \xi_{m3} \right| < 1, |x_{13}| < 1, |x_{23}| < 1 \right\}. \tag{128}$$

In order to continue the scalar integral C_0^b from the region Ω_7^b to the cube Ξ_K^b , one employs the system of PDEs in Eq.(93) with the finite element method. We will consider this in detail in section VI.

- 2: In the convergent region Ω_8^b , the scalar integral C_0^b is

$$\begin{aligned}
C_0^b(p_1^2, p_2^2, p_3^2, m^2) & = \left\{ C_{0,3}^b + C_{0,4}^b + C_{0,8}^b + C_{0,9}^b + C_{0,21}^b \right. \\
& \left. + C_{0,27}^b + C_{0,33}^b \right\} (p_1^2, p_2^2, p_3^2, m^2) \\
& = \frac{i(p_2^2)^{D/2-3}}{(4\pi)^{D/2}} F_{b,p_2}(\xi_{m2}, x_{12}, x_{32}) \tag{129}
\end{aligned}$$

In the limit $m^2 \rightarrow 0$, the expression of Eq.(129) recovers that presented in Eq.(35) exactly. The expansion of the scalar integral around $\varepsilon = 0$ is

$$C_0^b = \text{Eq.}(127)(p_3 \rightarrow p_2, \xi_{m3} \rightarrow \xi_{m2}, x_{13} \rightarrow x_{12}, x_{23} \rightarrow x_{32}). \tag{130}$$

Furthermore, the function $x_{23}^{D/2-3}F_{b,p_2}$ complies with the system of linear PDEs in Eq (93). Or equivalently, the function F_{b,p_2} satisfies the holonomic hypergeometric system of linear PDEs obtained through the interchanging $2 \leftrightarrow 3$ in Eq.(93).

Obviously the absolutely and uniformly converging region Ω_8^b is a proper subset of the cube

$$\Xi_{p_2}^b = \left\{ (\xi_{m_2}, x_{12}, x_{32}) : \left| \xi_{m_2} \right| < 1, |x_{12}| < 1, |x_{32}| < 1 \right\}. \quad (131)$$

In order to continue the scalar integral C_0^b from the region Ω_8^b to the cube $\Xi_{p_2}^b$, one employs the corresponding system of linear PDEs.

- 3: In the parameter space Ω_{11}^b , the scalar integral C_0^b is

$$\begin{aligned} C_0^b(p_1^2, p_2^2, p_3^2, m^2) &= \left\{ C_{0,10}^b + C_{0,11}^b + C_{0,13}^b + C_{0,14}^b + C_{0,21}^b \right. \\ &\quad \left. + C_{0,27}^b + C_{0,33}^b \right\} (p_1^2, p_2^2, p_3^2, m^2) \\ &= \frac{i(p_1^2)^{D/2-3}}{(4\pi)^{D/2}} F_{b,p_1}(\xi_{m_1}, x_{21}, x_{31}) \end{aligned} \quad (132)$$

In the limit $m^2 \rightarrow 0$, the expression of Eq.(132) recovers that presented in Eq.(34). The expansion of the scalar integral around $\varepsilon = 0$ is

$$C_0^b = \text{Eq.(127)}(p_3 \rightarrow p_1, \xi_{m_3} \rightarrow x_{m_1}, x_{13} \rightarrow x_{31}, x_{23} \rightarrow x_{21}). \quad (133)$$

In addition, the function $x_{13}^{D/2-3}F_{b,p_1}$ complies with the system of linear PDEs in Eq (93). Or equivalently, the function F_{b,p_1} satisfies the holonomic hypergeometric system of linear PDEs obtained through the interchanging $1 \leftrightarrow 3$ in Eq.(93).

Obviously the absolutely and uniformly converging region Ω_{11}^b is a proper subset of the cube

$$\Xi_{p_1}^b = \left\{ (\xi_{m_1}, x_{21}, x_{31}) : \left| \xi_{m_1} \right| < 1, |x_{21}| < 1, |x_{31}| < 1 \right\}. \quad (134)$$

In order to continue the scalar integral C_0^b from the region Ω_{11}^b to the cube $\Xi_{p_1}^b$, one employs the corresponding system of linear PDEs.

- 4: In the parameter space Ω_{19}^b , the scalar integral C_0^b is written as

$$\begin{aligned} C_0^b(p_1^2, p_2^2, p_3^2, m^2) &= C_{0,19}^b(p_1^2, p_2^2, p_3^2, m^2) \\ &= \frac{i(m^2)^{D/2-3}}{(4\pi)^{D/2}} F_{b,m}(\eta_{1m}, \eta_{2m}, \eta_{3m}). \end{aligned} \quad (135)$$

Except a convenient factor, the expression coincides with Eq (37) of the literature [13] exactly. We expand the hypergeometric functions $C_{0,19}^b$ around $\varepsilon = 0$ as

$$\begin{aligned}
C_0^b(p_1^2, p_2^2, p_3^2, m^2) &= -\frac{i(m^2)^{D/2-3}}{(4\pi)^{D/2}} \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \eta_{3m}^j \eta_{1m}^{n_1} \eta_{2m}^{n_2} \\
&\times \frac{\Gamma(1+j+n_1)\Gamma(1+j+n_2)\Gamma(1+n_1+n_2)}{j!n_1!n_2!} \\
&\times \frac{\Gamma(1+j+n_1+n_2)}{\Gamma(3+2j+2n_1+2n_2)} \left[1 + \varepsilon\psi(1+j+n_1+n_2) \right. \\
&\left. + \mathcal{O}(\varepsilon^2) \right]. \tag{136}
\end{aligned}$$

Additionally the function $\xi^{D/2-3}F_{b,m}$ complies with the system of linear PDEs in Eq (93). Or equivalently, the function $F_{b,m}$ satisfies the following system of linear PDEs

$$\begin{aligned}
&\left\{ \left(3 - \frac{D}{2} + \sum_{j=1}^3 \hat{\theta}_{\eta_{jm}} \right) \prod_{j \neq i}^3 (1 + \hat{\theta}_{y_{im}} + \hat{\theta}_{y_{jm}}) \right. \\
&\left. - \frac{1}{y_{im}} \hat{\theta}_{y_{im}} \left(1 + 2 \sum_{j=1}^3 \hat{\theta}_{\eta_{jm}} \right) \left(2 + 2 \sum_{j=1}^3 \hat{\theta}_{\eta_{jm}} \right) \right\} F_{b,m} = 0, \quad (i = 1, 2, 3). \tag{137}
\end{aligned}$$

Obviously the cube

$$\Xi_m^b = \left\{ (\eta_{1m}, \eta_{2m}, \eta_{3m}) : \left| \eta_{1m} \right| < 1, \left| \eta_{2m} \right| < 1, \left| \eta_{3m} \right| < 1 \right\}. \tag{138}$$

is a proper subset of the absolutely and uniformly converging region Ω_{19}^b . In order to continue the scalar integral C_0^b from the region Ω_{19}^b to the relevant parameter space, one employs the system of linear PDEs in Eq. (137).

VI. THE SYSTEM OF LINEAR PDES AS THE STATIONARY CONDITION OF A FUNCTIONAL

As stated above, the C_0 function of one nonzero mass is formulated through the hypergeometric functions in some convergent regions of independent variables. Since there is not the reduction formula for the hypergeometric functions, the scalar integral cannot be analytically continued outside the convergent regions. Nevertheless the continuation of the scalar integrals to the whole parameter space can be done numerically by the systems of

linear PDEs in Eq.(39). In order to proceed with our analysis, we expand F_{a,p_3} around time-space dimensions $D = 4$ as

$$F_{a,p_3}(\xi_{33}, x_{13}, x_{23}) = F_{a,p_3}^{(0)}(\xi_{33}, x_{13}, x_{23}) + \sum_{i=1}^{\infty} \varepsilon^i F_{a,p_3}^{(i)}(\xi_{33}, x_{13}, x_{23}), \quad (139)$$

where the function $F_{a,p_3}^{(n)}(\xi_{33}, x_{13}, x_{23})$ satisfies the system of PDEs

$$\begin{aligned} & \left[\xi_{33} \left(1 + \hat{\vartheta}_{\xi_{33}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} \right)^2 \hat{\vartheta}_{\xi_{33}} - \prod_{i=1}^2 (\hat{\vartheta}_{\xi_{33}} + \hat{\vartheta}_{x_{i3}}) \hat{\vartheta}_{\xi_{33}} \right] F_{a,p_3}^{(n)} + f_{a,0}^{(n)} = 0, \\ & \left[x_{j3} \left(1 + \hat{\vartheta}_{\xi_{33}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} \right)^2 - \hat{\vartheta}_{x_{j3}} (\hat{\vartheta}_{\xi_{33}} + \hat{\vartheta}_{x_{j3}}) \right] F_{a,p_3}^{(n)} + f_{a,j}^{(n)} = 0, \quad (j = 1, 2), \end{aligned} \quad (140)$$

with

$$\begin{aligned} f_{a,0}^{(n)}(\xi_{33}, x_{13}, x_{23}) &= \left[\xi_{33} \left(2 + 5\hat{\vartheta}_{\xi_{33}} + 4 \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} + (3\hat{\vartheta}_{\xi_{33}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}})(\hat{\vartheta}_{\xi_{33}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}}) \right) \right. \\ & \quad \left. - 2\hat{\vartheta}_{\xi_{33}}^2 - \hat{\vartheta}_{\xi_{33}} \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} \right] F_{a,p_3}^{(n-1)}(\xi_{33}, x_{13}, x_{23}) \\ & \quad + \left[2\xi_{33} \left(1 + \hat{\vartheta}_{\xi_{33}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} \right) - \hat{\vartheta}_{\xi_{33}} \right] F_{a,p_3}^{(n-2)}(\xi_{33}, x_{13}, x_{23}), \\ f_{a,j}^{(n)}(\xi_{33}, x_{13}, x_{23}) &= \left[x_{j3} \left(1 + \hat{\vartheta}_{\xi_{33}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} \right) - \hat{\vartheta}_{x_{j2}} \right] F_{a,p_3}^{(n-1)}(\xi_{33}, x_{13}, x_{23}), \quad (j = 1, 2) \end{aligned} \quad (141)$$

Through the transformation of variables

$$\begin{aligned} t_m &= \ln \xi_{33}, \\ t_1 &= \ln x_{13}, \\ t_2 &= \ln x_{23}, \end{aligned} \quad (142)$$

the system of PDEs in Eq.(140) is recognized as stationary conditions of the modified functional

$$\begin{aligned} \Pi^*(F_{a,p_3}^{(n)}) &= \Pi \left(\frac{\partial F_{a,p_3}^{(n)}}{\partial t_m} \right) \\ & \quad + \sum_{i=1}^2 \int_{\Omega} \chi_i \left\{ e^{t_i} \left(1 + \frac{\partial}{\partial t_m} + \frac{\partial}{\partial t_1} + \frac{\partial}{\partial t_2} \right)^2 F_{a,p_3}^{(n)} \right. \\ & \quad \left. - \frac{\partial}{\partial t_i} \left(\frac{\partial}{\partial t_m} + \frac{\partial}{\partial t_i} \right) F_{a,p_3}^{(n)} + f_{a,i}^{(n)} \right\} dt_m dt_1 dt_2. \end{aligned} \quad (143)$$

Here $\chi_1(t_m, t_1, t_2)$, $\chi_2(t_m, t_1, t_2)$ are Lagrange multipliers, Ω represents the parameter space where the continuation of the solution is made numerically, and $\Pi\left(\frac{\partial F_{a,k}^{(n)}}{\partial t_m}\right)$ is the functional of the first PDE in Eq.(141):

$$\begin{aligned} \Pi\left(\frac{\partial F_{a,p_3}^{(n)}}{\partial t_m}\right) &= \int_{\Omega} \left\{ (e^{t_m} - 1) \left(\frac{\partial^2 F_{a,p_3}^{(n)}}{\partial t_m^2}\right)^2 + e^{t_m} \left(\frac{\partial^2 F_{a,p_3}^{(n)}}{\partial t_m \partial t_1}\right)^2 + e^{t_m} \left(\frac{\partial^2 F_{a,p_3}^{(n)}}{\partial t_m \partial t_2}\right)^2 \right. \\ &\quad + \left(e^{t_m} - \frac{1}{2}\right) \frac{\partial^2 F_{a,p_3}^{(n)}}{\partial t_m^2} \frac{\partial^2 F_{a,p_3}^{(n)}}{\partial t_m \partial t_1} + \left(e^{t_m} - \frac{1}{2}\right) \frac{\partial^2 F_{a,p_3}^{(n)}}{\partial t_m^2} \frac{\partial^2 F_{a,p_3}^{(n)}}{\partial t_m \partial t_2} \\ &\quad + \left(e^{t_m} - \frac{1}{2}\right) \frac{\partial^2 F_{a,p_3}^{(n)}}{\partial t_m \partial t_1} \frac{\partial^2 F_{a,p_3}^{(n)}}{\partial t_m \partial t_2} + e^{t_m} \frac{\partial F_{a,p_3}^{(n)}}{\partial t_m} \frac{\partial^2 F_{a,p_3}^{(n)}}{\partial t_m^2} \\ &\quad \left. - f_{a,0}^{(n)} \frac{\partial F_{a,p_3}^{(n)}}{\partial t_m} \right\} dt_m dt_1 dt_2 . \end{aligned} \quad (144)$$

Furthermore, the stationary condition of the second term of Eq.(143) is the second PDEs in Eq.(141), that of the third term of Eq.(143) is the third PDEs in Eq.(141), which are recognized as two restrictions of the system. Once the solutions $F_{a,k}^{(i)}$, ($i \leq n-1, n \geq 1$) are obtained numerically in the parameter space Ω , the solution $F_{a,k}^{(n)}$ can be numerically continued from the convergent region Ω_7^a to the concerned region Ω with the finite element method [32].

The systems of linear PDEs of Eq. (39), Eq. (80), and Eq. (89) are compatible with each other since they all originate from the Mellin-Barnes representation of Eq. (37). Correspondingly a similar functional is constructed with the system of PDEs in Eq. (80). Expansion of F_{a,p_1} around space-time dimensions $D = 4$ is

$$F_{a,p_1}(\xi_{31}, x_{21}, x_{31}) = F_{a,p_1}^{(0)}(\xi_{31}, x_{21}, x_{31}) + \sum_{i=1}^{\infty} \varepsilon^i F_{a,p_1}^{(i)}(\xi_{31}, x_{21}, x_{31}), \quad (145)$$

the solution $F_{a,p_1}^{(n)}$ can be numerically continued from the convergent region $\Omega_{11}^a \cup \Omega_{29}^a$ to the concerned region Ω with the finite element method. The compatibility between the systems of linear PDEs of Eq. (39) and Eq. (80) implies

$$F_{a,p_1}^{(n)}(\xi_{31}, x_{21}, x_{31}) \equiv \sum_{n'=0}^n \frac{(\ln x_{13})^{n'}}{n'!} F_{a,p_3}^{(n-n')}(\xi_{33}, x_{13}, x_{23}), \quad (146)$$

which can be used to check whether the numerical program runs correctly.

Similarly the C_0 function of three equal masses is formulated through the hypergeometric functions in some convergent regions, and the continuation of the scalar integrals to the whole

parameter space can be made numerically with the systems of linear PDEs in Eq.(93). In order to proceed with our analysis, we expand F_{b,p_3} around space-time dimensions $D = 4$ as

$$F_{b,p_3}(\xi_{m_3}, x_{13}, x_{23}) = F_{b,p_3}^{(0)}(\xi_{m_3}, x_{13}, x_{23}) + \sum_{i=1}^{\infty} \varepsilon^i F_{b,p_3}^{(i)}(\xi_{m_3}, x_{13}, x_{23}), \quad (147)$$

where the function $F_{b,p_3}^{(n)}(\xi_{m_3}, x_{13}, x_{23})$ satisfies the system of PDEs

$$\begin{aligned} & \left[2\xi_{m_3} (1 + \hat{\vartheta}_{\xi_{m_3}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}}) (1 + 2\hat{\vartheta}_{\xi_{m_3}}) \hat{\vartheta}_{\xi_{m_3}} \right. \\ & \left. - \prod_{i=1}^2 (\hat{\vartheta}_{\xi_{m_3}} + \hat{\vartheta}_{x_{i3}}) \hat{\vartheta}_{\xi_{m_3}} \right] F_{b,p_3}^{(n)} + f_{b,0}^{(n)} = 0, \\ & \left[x_{j3} (1 + \hat{\vartheta}_{\xi_{m_3}} + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}}) (1 + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}}) \right. \\ & \left. - \hat{\vartheta}_{x_{j3}} (\hat{\vartheta}_{\xi_{m_3}} + \hat{\vartheta}_{x_{j3}}) \right] F_{b,p_3}^{(n)} + f_{b,j}^{(n)} = 0, \quad (j = 1, 2), \end{aligned} \quad (148)$$

with

$$\begin{aligned} f_{b,0}^{(n)}(\xi_{m_3}, x_{13}, x_{23}) &= \left[2\xi_{m_3} (1 + 6\hat{\vartheta}_{\xi_{m_3}} + 4\hat{\vartheta}_{\xi_{m_3}} \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} + 6\hat{\vartheta}_{\xi_{m_3}}^2) \right. \\ & \quad \left. - 2\hat{\vartheta}_{\xi_{m_3}}^2 - \hat{\vartheta}_{\xi_{m_3}} \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}} \right] F_{b,p_3}^{(n-1)}(\xi_{m_3}, x_{13}, x_{23}) \\ & \quad + \left[2\xi_{m_3} (3 + 6\hat{\vartheta}_{\xi_{m_3}} + 2 \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}}) - \hat{\vartheta}_{\xi_{m_3}} \right] F_{b,p_3}^{(n-2)}(\xi_{m_3}, x_{13}, x_{23}) \\ & \quad + \xi_{m_3} F_{b,p_3}^{(n-3)}(\xi_{m_3}, x_{13}, x_{23}), \\ f_{b,j}^{(n)}(\xi_{m_3}, x_{13}, x_{23}) &= \left[x_{j3} (1 + \sum_{i=1}^2 \hat{\vartheta}_{x_{i3}}) - \hat{\vartheta}_{x_{j3}} \right] F_{b,p_3}^{(n-1)}(\xi_{m_3}, x_{13}, x_{23}), \quad (j = 1, 2) \end{aligned} \quad (149)$$

Through the transformation of variables in Eq. (142), the system of PDEs in Eq.(148) is recognized as stationary conditions of the modified functional

$$\begin{aligned} \Pi^*(F_{b,p_3}^{(n)}) &= \Pi\left(\frac{\partial F_{b,p_3}^{(n)}}{\partial t_m}\right) \\ & \quad + \sum_{i=1}^2 \int_{\Omega} \chi_i \left\{ e^{t_i} \left(1 + \frac{\partial}{\partial t_m} + \frac{\partial}{\partial t_1} + \frac{\partial}{\partial t_2} \right) \left(1 + \frac{\partial}{\partial t_1} + \frac{\partial}{\partial t_2} \right) F_{b,p_3}^{(n)} \right. \\ & \quad \left. - \frac{\partial}{\partial t_i} \left(\frac{\partial}{\partial t_m} + \frac{\partial}{\partial t_i} \right) F_{b,p_3}^{(n)} + f_{b,i}^{(n)} \right\} dt_m dt_1 dt_2. \end{aligned} \quad (150)$$

Here $\chi_1(t_m, t_1, t_2)$, $\chi_2(t_m, t_1, t_2)$ are Lagrange multipliers, Ω represents the parameter space where the continuation of the solution is made numerically, and $\Pi\left(\frac{\partial F_{b,p_3}^{(n)}}{\partial t_m}\right)$ is the functional

of the first PDE in Eq.(149):

$$\begin{aligned}
\Pi\left(\frac{\partial F_{b,p_3}^{(n)}}{\partial t_m}\right) &= \int_{\Omega} \left\{ (4e^{t_m} - 1) \left(\frac{\partial^2 F_{b,p_3}^{(n)}}{\partial t_m^2} \right)^2 + \left(2e^{t_m} - \frac{1}{2} \right) \frac{\partial^2 F_{b,p_3}^{(n)}}{\partial t_m^2} \frac{\partial^2 F_{b,p_3}^{(n)}}{\partial t_m \partial t_1} \right. \\
&\quad + \left(2e^{t_m} - \frac{1}{2} \right) \frac{\partial^2 F_{b,p_3}^{(n)}}{\partial t_m^2} \frac{\partial^2 F_{b,p_3}^{(n)}}{\partial t_m \partial t_2} - \frac{1}{2} \frac{\partial^2 F_{b,p_3}^{(n)}}{\partial t_m \partial t_1} \frac{\partial^2 F_{b,p_3}^{(n)}}{\partial t_m \partial t_2} \\
&\quad \left. - e^{t_m} \left(\frac{\partial F_{b,p_3}^{(n)}}{\partial t_m} \right)^2 - f_0^{(n)} \frac{\partial F_{b,p_3}^{(n)}}{\partial t_m} \right\} dt_m dt_1 dt_2 . \tag{151}
\end{aligned}$$

Furthermore, the stationary condition of the second term of Eq.(150) is the second PDEs in Eq.(149), that of the third term of Eq.(143) is the third PDEs in Eq.(149), which are recognised as two restrictions of the system. Once the solutions $F_k^{(i)}$ ($i \leq n-1, n \geq 1$) are obtained numerically in the parameter region Ω , the solution $F_k^{(n)}$ can be numerically continued from the convergent region Ω_7^b to the concerned region Ω with the finite element method [32].

The systems of linear PDEs of Eq. (93) and Eq. (137) are compatible with each other since every one of them originates from the Mellin-Barnes representation of Eq. (91). Correspondingly a similar functional is constructed with the holonomic hypergeometric system of PDEs obtained through $1 \leftrightarrow 3$ in Eq. (93). Expansion of F_{b,p_1} around space-time dimensions $D = 4$ is

$$F_{b,p_1}(\xi_{m1}, x_{21}, x_{31}) = F_{b,p_1}^{(0)}(\xi_{m1}, x_{21}, x_{31}) + \sum_{i=1}^{\infty} \varepsilon^i F_{b,p_1}^{(i)}(\xi_{m1}, x_{21}, x_{31}), \tag{152}$$

the solution $F_{b,p_1}^{(n)}$ can be numerically continued from the convergent region Ω_{11}^b to the concerned region Ω with the finite element method. The compatibility between the holonomic hypergeometric systems of linear PDEs implies

$$F_{b,p_1}^{(n)}(\xi_{m1}, x_{21}, x_{31}) \equiv \sum_{n'=0}^n \frac{(\ln x_{13})^{n'}}{n'!} F_{b,p_3}^{(n-n')}(\xi_{m3}, x_{13}, x_{23}), \tag{153}$$

which can be used to check whether the numerical program runs correctly.

VII. SUMMARY

Using the α - or Feynman-parameterization, we express any scalar integral by its Mellin-Barnes representation, and present the system of linear PDEs satisfied by the scalar

integral. Through the residue theorem and homology, those scalar integrals are written as the multiple hypergeometric functions of the independent variables. Horn's theory of convergence predicts the absolutely and uniformly convergent region of each hypergeometric functions. Several convergent regions of the hypergeometric functions compose a set, where each definition domain does not intersect with the others in the set. Additionally each convergent region of the set either does not intersect with, or is a proper subset of other convergent region which does not belong to the set. In each definition domain of the set, the scalar integral can be written as the sum of those hypergeometric functions whose convergent regions contain the concerned element entirely. With the idea above, some well-known results of Ref. [10, 13] are recovered. Taking the system of PDEs as stationary conditions of a functional under some given restrictions, one continues the scalar integrals to the whole parameter space numerically with the finite element methods.

The approach proposed here can be applied to evaluate any Feynman integral of one-loop multiple point diagrams directly. In some connected regions of absolute and uniform convergence, for example, the D_0 function is formulated as the linear combination of the nonuple hypergeometric functions. The corresponding system of PDEs is composed by nine linear independent PDEs.

In order to apply the method to the scalar integrals of multi-loop diagrams, one introduces some auxiliary parameters [21–23]. For example, the analysis on the scalar integral of two-loop planar vertex introduces two auxiliary kinematic invariants which are linear combinations of p_1^2 , p_2^2 , and p_3^2 , the analysis on the scalar integral of two-loop non-planar vertex introduces five auxiliary kinematic invariants which are linear combinations of p_1^2 , p_2^2 , and p_3^2 , respectively. We will present our analyses elsewhere.

Acknowledgments

The work has been supported partly by the National Natural Science Foundation of China (NNSFC) with Grant No. 11821505, No. 11447601, No. 11675239, No. 11535002, No. 11705045, and No. 11805140. Furthermore, the author (C.-H. Chang) is also supported by Key Research program of Frontier Sciences, CAS, Grant No. QYZDY-SSW-SYS006.

Appendix A: Some derived hypergeometric functions

$$\begin{aligned}
C_{0,7}^a(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \\
&\quad \times \xi_{33}^j x_{13}^{D/2-2-j+n_1} x_{23}^{D/2-2-j+n_2} \frac{\Gamma(4-D+j)}{j!n_1!n_2!} \\
&\quad \times \frac{\Gamma(\frac{D}{2} - 1 - j + n_1 + n_2) \Gamma(D-3-j+n_1+n_2)}{\Gamma(\frac{D}{2} - 1 - j + n_1) \Gamma(\frac{D}{2} - 1 - j + n_2)} \\
&= \frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3)} \\
&\quad \times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} x_{13}^{D/2-2+n_1} x_{23}^{D/2-2+n_2} \\
&\quad \times \frac{\Gamma(\frac{D}{2} - 1 + n_1 + n_2) \Gamma(D-3+n_1+n_2)}{n_1!n_2! \Gamma(\frac{D}{2} - 1 + n_1) \Gamma(\frac{D}{2} - 1 + n_2)} \\
&\quad + \frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} \\
&\quad \times \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \sum_{n_2=0}^{\infty} \xi_{33}^{j+n_2} x_{13}^{D/2-2+n_1} x_{23}^{D/2-2-j} \\
&\quad \times \frac{(-)^j \Gamma(4-D+j+n_2) \Gamma(2 - \frac{D}{2} + j)}{(j+n_2)! (j+n_1+n_2)! n_2!} \\
&\quad \times \frac{\Gamma(\frac{D}{2} - 1 + n_1 + n_2) \Gamma(D-3+n_1+n_2)}{\Gamma(\frac{D}{2} - 1 + n_1)} \\
&\quad + \frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} \\
&\quad \times \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \xi_{33}^j x_{13}^{D/2-2+n_1} x_{23}^{D/2-2+n_2} \Gamma(4-D+j) \\
&\quad \times \frac{\Gamma(\frac{D}{2} - 1 + j + n_1 + n_2) \Gamma(D-3+j+n_1+n_2)}{j!(j+n_1)!(j+n_2)! \Gamma(\frac{D}{2} - 1 + n_1) \Gamma(\frac{D}{2} - 1 + n_2)} \\
&\quad + \frac{i(-)^{D/2}(p_3^2)^{D/2-3} \Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} \\
&\quad \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=1}^{\infty} \xi_{33}^{j+n_1} x_{13}^{D/2-2-j} x_{23}^{D/2-2+n_2} \\
&\quad \times \frac{(-)^j \Gamma(4-D+j+n_1) \Gamma(2 - \frac{D}{2} + j)}{(j+n_1)! n_1! (j+n_1+n_2)!} \\
&\quad \times \frac{\Gamma(\frac{D}{2} - 1 + n_1 + n_2) \Gamma(D-3+n_1+n_2)}{\Gamma(\frac{D}{2} - 1 + n_2)}
\end{aligned}$$

$$\begin{aligned}
& + \frac{i(-)^{D/2}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1) \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{33}^{j+n_1+n_2} x_{13}^{D/2-2-j-n_2} x_{23}^{D/2-2-j-n_1} \\
& \times \frac{(-)^{n_1+n_2} \Gamma(4 - D + j + n_1 + n_2) \Gamma(2 - \frac{D}{2} + j + n_1)}{(j + n_1 + n_2)! n_1! n_2! \Gamma(2 - \frac{D}{2} + j)} \\
& \times \frac{\Gamma(2 - \frac{D}{2} + j + n_2)}{\Gamma(4 - D + j)} \\
& + \frac{i(-)^{D/2}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{1}{\Gamma(D-3)\Gamma(4-D)} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \xi_{33}^{j+n_1+n_2} x_{13}^{D/2-2-j} x_{23}^{D/2-2-n_1} \\
& \times \frac{(-)^{j+n_1} \Gamma(\frac{D}{2} - 1 + n_2) \Gamma(D-3+n_2)}{(j + n_1 + n_2)! (n_1 + n_2)! (j + n_2)!} \\
& \times \Gamma(4 - D + j + n_1 + n_2) \Gamma(2 - \frac{D}{2} + j) \Gamma(2 - \frac{D}{2} + n_1) \\
& + \frac{i(-)^{D/2}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma^2(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1)}{\Gamma(D-3)\Gamma(4-D)} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \xi_{33}^j x_{13}^{D/2-2+n_1} x_{23}^{D/2-2} \frac{\Gamma(4-D+j)}{(j!)^2 (j+n_1)!} \\
& \times \frac{\Gamma(\frac{D}{2} - 1 + j + n_1) \Gamma(D-3+j+n_1)}{\Gamma(\frac{D}{2} - 1 + n_1)} \\
& + \frac{i(-)^{D/2}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma(2 - \frac{D}{2})}{\Gamma(D-3)\Gamma(4-D)} \\
& \times \sum_{n_1=1}^{\infty} \sum_{j=1}^{\infty} \xi_{33}^{j+n_1} x_{13}^{D/2-2-j} x_{23}^{D/2-2} \frac{(-)^j \Gamma(4-D+j+n_1)}{((j+n_1)!)^2 n_1!} \\
& \times \Gamma(\frac{D}{2} - 1 + n_1) \Gamma(D-3+n_1) \Gamma(2 - \frac{D}{2} + j) \\
& + \frac{i(-)^{D/2}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma^2(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1)}{\Gamma(D-3)\Gamma(4-D)} \\
& \times \sum_{j=1}^{\infty} \sum_{n_2=1}^{\infty} \xi_{33}^j x_{13}^{D/2-2} x_{23}^{D/2-2+n_2} \frac{\Gamma(4-D+j)}{(j!)^2 (j+n_2)!} \\
& \times \frac{\Gamma(\frac{D}{2} - 1 + j + n_2) \Gamma(D-3+j+n_2)}{\Gamma(\frac{D}{2} - 1 + n_2)} \\
& + \frac{i(-)^{D/2}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma(2 - \frac{D}{2})}{\Gamma(D-3)\Gamma(4-D)} \sum_{n_2=1}^{\infty} \sum_{j=1}^{\infty} \\
& \times \xi_{33}^{j+n_2} x_{13}^{D/2-2} x_{23}^{D/2-2-j} \frac{(-)^j \Gamma(4-D+j+n_2)}{((j+n_2)!)^2 n_2!}
\end{aligned}$$

$$\begin{aligned}
& \times \Gamma\left(\frac{D}{2} - 1 + n_2\right) \Gamma(D - 3 + n_2) \Gamma\left(2 - \frac{D}{2} + j\right) \\
& + \frac{i(-)^{D/2} (p_3^2)^{D/2-3} \Gamma\left(\frac{D}{2} - 1\right)}{(4\pi)^{D/2} \Gamma(4 - D)} \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \\
& \times \xi_{33}^{j+n_1} x_{13}^{D/2-2-j} x_{23}^{D/2-2-n_1} \frac{(-)^{j+n_1} \Gamma(4 - D + j + n_1)}{(j + n_1)! n_1! j!} \\
& \times \Gamma\left(2 - \frac{D}{2} + j\right) \Gamma\left(2 - \frac{D}{2} + n_1\right) \\
& + \frac{i(-)^{D/2} (p_3^2)^{D/2-3} \Gamma\left(2 - \frac{D}{2}\right) \Gamma\left(\frac{D}{2} - 1\right)}{(4\pi)^{D/2} \Gamma(4 - D)} \sum_{j=1}^{\infty} \xi_{33}^j x_{13}^{D/2-2} \\
& \times x_{23}^{D/2-2-j} \frac{(-)^j \Gamma(4 - D + j) \Gamma\left(2 - \frac{D}{2} + j\right)}{(j!)^2}. \tag{A1}
\end{aligned}$$

$$\begin{aligned}
C_{0,s}^a(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(-)^{D/2} (p_3^2)^{D/2-3} \Gamma^2\left(2 - \frac{D}{2}\right) \Gamma^2\left(\frac{D}{2} - 1\right)}{(4\pi)^{D/2} \Gamma(D - 3) \Gamma(4 - D)} \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \\
& \times \frac{(-)^j}{j! n_1! n_2!} \xi_{33}^j x_{13}^{D/2-2-j+n_1} x_{23}^{1-D/2-n_1-n_2} \\
& \times \frac{\Gamma(D - 3 - j + n_1 + n_2) \Gamma\left(\frac{D}{2} - 1 + n_1 + n_2\right) \Gamma(4 - D + j)}{\Gamma\left(\frac{D}{2} - 1 - j + n_1\right) \Gamma\left(\frac{D}{2} - 1 + n_2\right)} \\
& = \frac{i(-)^{D/2} (p_2^2)^{D/2-3} \Gamma^2\left(2 - \frac{D}{2}\right) \Gamma^2\left(\frac{D}{2} - 1\right)}{(4\pi)^{D/2} \Gamma(D - 3)} \\
& \times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} x_{12}^{D/2-2+n_1} x_{32}^{D/2-2+n_2} \\
& \times \frac{\Gamma(D - 3 + n_1 + n_2) \Gamma\left(\frac{D}{2} - 1 + n_1 + n_2\right)}{n_1! n_2! \Gamma\left(\frac{D}{2} - 1 + n_1\right) \Gamma\left(\frac{D}{2} - 1 + n_2\right)} \\
& + \frac{i(-)^{D/2} (p_2^2)^{D/2-3} \Gamma^2\left(2 - \frac{D}{2}\right) \Gamma^2\left(\frac{D}{2} - 1\right)}{(4\pi)^{D/2} \Gamma(D - 3) \Gamma(4 - D)} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \sum_{n_2=0}^{\infty} \xi_{32}^j x_{12}^{D/2-2+n_1} x_{32}^{D/2-2+n_2} \\
& \times \frac{(-)^j \Gamma\left(\frac{D}{2} - 1 + j + n_1 + n_2\right) \Gamma(4 - D + j)}{j! (j + n_1)! n_2! \Gamma\left(\frac{D}{2} - 1 + n_2\right)} \\
& \times \frac{\Gamma(D - 3 + n_1 + n_2)}{\Gamma\left(\frac{D}{2} - 1 + n_1\right)} \\
& + \frac{i(-)^{D/2} (p_2^2)^{D/2-3} \Gamma\left(2 - \frac{D}{2}\right) \Gamma\left(\frac{D}{2} - 1\right)}{(4\pi)^{D/2} \Gamma(D - 3) \Gamma(4 - D)} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=1}^{\infty} \xi_{32}^{j+n_1} x_{12}^{D/2-2-j} x_{32}^{D/2-1+j+n_2}
\end{aligned}$$

$$\begin{aligned}
& \times \frac{(-)^{n_1} \Gamma(\frac{D}{2} - 1 + j + n_1 + n_2) \Gamma(4 - D + j + n_1)}{(j + n_1)! n_1! (j + n_2)! \Gamma(\frac{D}{2} - 1 + j + n_2)} \\
& \times \Gamma(D - 3 + n_2) \Gamma(2 - \frac{D}{2} + j) \\
& + \frac{i(-)^{D/2} (p_2^2)^{D/2-3}}{(4\pi)^{D/2}} \Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1) \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{32}^{j+n_1+n_2} x_{12}^{D/2-2-j-n_2} x_{32}^{D/2-2+n_2} \\
& \times \frac{(-)^{j+n_1} \Gamma(\frac{D}{2} - 1 + n_1 + n_2) \Gamma(4 - D + j + n_1 + n_2)}{(j + n_1 + n_2)! n_1! n_2! \Gamma(\frac{D}{2} - 1 + n_2)} \\
& \times \frac{\Gamma(2 - \frac{D}{2} + j + n_2)}{\Gamma(4 - D + j)} \\
& + \frac{i(-)^{D/2} (p_2^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1)}{\Gamma(4 - D)} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \xi_{32}^{j+n_1} x_{12}^{D/2-2-j} x_{32}^{D/2-2+j} \\
& \times \frac{(-)^{n_1} \Gamma(\frac{D}{2} - 1 + j + n_1) \Gamma(4 - D + j + n_1)}{(j + n_1)! n_1! j! \Gamma(\frac{D}{2} - 1 + j)} \\
& \times \Gamma(2 - \frac{D}{2} + j) \\
& + \frac{i(-)^{D/2} (p_2^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma^2(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1)}{\Gamma(D - 3) \Gamma(4 - D)} \\
& \times \sum_{j=1}^{\infty} \sum_{n_2=0}^{\infty} \xi_{32}^j x_{12}^{D/2-2} x_{32}^{D/2-2+n_2} \\
& \times \frac{(-)^j \Gamma(\frac{D}{2} - 1 + j + n_2) \Gamma(4 - D + j) \Gamma(D - 3 + n_2)}{(j!)^2 n_2! \Gamma(\frac{D}{2} - 1 + n_2)}. \quad (A2)
\end{aligned}$$

$$\begin{aligned}
C_{0,9}^a(p_1^2, p_2^2, p_3^2, m^2) &= - \frac{i(-)^{D/2} (p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{\Gamma(D - 3) \Gamma(4 - D)} \\
& \times \sum_{n_2=0}^{\infty} \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \frac{(-)^j}{j! n_1! n_2!} \xi_{33}^j x_{13}^{D/2-2-j+n_1} x_{23}^{-1-n_1-n_2} \\
& \times \frac{\Gamma(1 + n_1 + n_2) \Gamma(4 - D + j) \Gamma(\frac{D}{2} - 1 - j + n_1 + n_2)}{\Gamma(3 - \frac{D}{2} + n_2) \Gamma(\frac{D}{2} - 1 - j + n_1)} \\
& = - \frac{i(-)^{D/2} (p_2^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{\Gamma(D - 3) \Gamma(3 - \frac{D}{2})} \sum_{n_1=0}^{\infty} x_{12}^{D/2-2+n_1} \\
& - \frac{i(-)^{D/2} (p_2^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{\Gamma(D - 3) \Gamma(4 - D) \Gamma(3 - \frac{D}{2})}
\end{aligned}$$

$$\begin{aligned}
& \times \frac{\sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \xi_{32}^j x_{12}^{D/2-2-j+n_1} \frac{(-)^j \Gamma(4-D+j)}{j!}}{i(-)^{D/2} (p_2^2)^{D/2-3} \Gamma^2(2-\frac{D}{2}) \Gamma^2(\frac{D}{2}-1)} \\
& \frac{(4\pi)^{D/2}}{\Gamma(D-3) \Gamma(4-D)} \\
& \times \frac{\sum_{n_2=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{j=1}^{\infty} \xi_{32}^{j+n_1+n_2} x_{12}^{D/2-2-j-n_2} x_{32}^{n_2}}{(-)^{j+n_1} \Gamma(1+n_1+n_2) \Gamma(4-D+j+n_1+n_2)} \\
& \frac{(j+n_1+n_2)! n_1! n_2! \Gamma(3-\frac{D}{2}+n_2)}{\Gamma(2-\frac{D}{2}+j+n_2)} \\
& \frac{\Gamma(2-\frac{D}{2}+j)}{\Gamma(2-\frac{D}{2}+j)} \\
& \frac{i(-)^{D/2} (p_2^2)^{D/2-3} \Gamma(2-\frac{D}{2}) \Gamma(\frac{D}{2}-1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} \\
& \times \frac{\sum_{n_2=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{j=1}^{\infty} \xi_{32}^{j+n_1} x_{12}^{D/2-2-j} x_{32}^{j+n_2}}{(-)^{n_1} \Gamma(1+j+n_1+n_2) \Gamma(4-D+j+n_1)} \\
& \frac{(j+n_1)! n_1! (j+n_2)! \Gamma(3-\frac{D}{2}+j+n_2)}{\Gamma(\frac{D}{2}-1+n_2) \Gamma(2-\frac{D}{2}+j)} \\
& \frac{i(-)^{D/2} (p_2^2)^{D/2-3} \Gamma^2(2-\frac{D}{2}) \Gamma^2(\frac{D}{2}-1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} \\
& \times \frac{\sum_{n_2=1}^{\infty} \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \xi_{32}^j x_{12}^{D/2-2+n_1} x_{32}^{n_2}}{(-)^j \Gamma(1+j+n_1+n_2) \Gamma(4-D+j) \Gamma(\frac{D}{2}-1+n_1+n_2)} \\
& \frac{j! (j+n_1)! n_2! \Gamma(3-\frac{D}{2}+n_2) \Gamma(\frac{D}{2}-1+n_1)}{i(-)^{D/2} (p_2^2)^{D/2-3} \Gamma^2(2-\frac{D}{2}) \Gamma^2(\frac{D}{2}-1)} \\
& \frac{(4\pi)^{D/2}}{\Gamma(D-3)} \\
& \times \frac{\sum_{n_2=1}^{\infty} \sum_{n_1=1}^{\infty} x_{12}^{D/2-2+n_1} x_{32}^{n_2}}{\Gamma(1+n_1+n_2) \Gamma(\frac{D}{2}-1+n_1+n_2)} \\
& \frac{n_1! n_2! \Gamma(3-\frac{D}{2}+n_2) \Gamma(\frac{D}{2}-1+n_1)}{i(-)^{D/2} (p_2^2)^{D/2-3} \Gamma^2(2-\frac{D}{2}) \Gamma^2(\frac{D}{2}-1)} \\
& \frac{(4\pi)^{D/2}}{\Gamma(D-3) \Gamma(4-D)} \\
& \times \frac{\sum_{n_2=1}^{\infty} \sum_{n_1=1}^{\infty} \xi_{32}^{n_1} x_{12}^{D/2-2} x_{32}^{n_2}}{(-)^{n_1} \Gamma(1+n_1+n_2) \Gamma(4-D+n_1) \Gamma(\frac{D}{2}-1+n_2)} \\
& \frac{(n_1!)^2 n_2! \Gamma(3-\frac{D}{2}+n_2)}{
\end{aligned}$$

$$\begin{aligned}
& \frac{i(-)^{D/2}(p_2^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D - 3)} \\
& \times \sum_{n_2=1}^{\infty} x_{12}^{D/2-2} x_{32}^{n_2} \frac{\Gamma(\frac{D}{2} - 1 + n_2)}{\Gamma(3 - \frac{D}{2} + n_2)} \\
& \frac{i(-)^{D/2}(p_2^2)^{D/2-3} \Gamma(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D - 3)\Gamma(4 - D)} \\
& \times \sum_{n_2=1}^{\infty} \sum_{n_1=0}^{\infty} \xi_{32}^{n_1+n_2} x_{12}^{D/2-2-n_2} x_{32}^{n_2} \\
& \times \frac{(-)^{n_1} \Gamma(4 - D + n_1 + n_2) \Gamma(2 - \frac{D}{2} + n_2)}{n_1! n_2! \Gamma(3 - \frac{D}{2} + n_2)}. \tag{A3}
\end{aligned}$$

$$\begin{aligned}
C_{0,2}^b(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D - 3)\Gamma(4 - D)} x_{23}^{D/2-2} \\
& \times \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m3}^j x_{13}^{n_1} x_{23}^{-j+n_2} \Gamma(4 - D + 2j) \\
& \times \frac{\Gamma(1 + n_1 + n_2) \Gamma(\frac{D}{2} - 1 - j + n_1 + n_2)}{j! n_1! n_2! \Gamma(3 - \frac{D}{2} + j + n_1) \Gamma(\frac{D}{2} - 1 - j + n_2)} \\
& = \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D - 3)} x_{23}^{D/2-2} \\
& \times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} x_{13}^{n_1} x_{23}^{n_2} \frac{\Gamma(1 + n_1 + n_2) \Gamma(\frac{D}{2} - 1 + n_1 + n_2)}{n_1! n_2! \Gamma(3 - \frac{D}{2} + n_1) \Gamma(\frac{D}{2} - 1 + n_2)} \\
& + \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D - 3)\Gamma(4 - D)} x_{23}^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=1}^{\infty} \xi_{m3}^j x_{13}^{n_1} x_{23}^{n_2} \\
& \times \frac{\Gamma(1 + j + n_1 + n_2) \Gamma(4 - D + 2j) \Gamma(\frac{D}{2} - 1 + n_1 + n_2)}{j! n_1! (j + n_2)! \Gamma(3 - \frac{D}{2} + j + n_1) \Gamma(\frac{D}{2} - 1 + n_2)} \\
& + \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3} \Gamma(2 - \frac{D}{2})\Gamma(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D - 3)\Gamma(4 - D)} x_{23}^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_2=0}^{\infty} \sum_{n_1=1}^{\infty} \xi_{m3}^{j+n_2} x_{13}^{j+n_1} x_{23}^{-j} \Gamma(2 - \frac{D}{2} + j) \Gamma(\frac{D}{2} - 1 + n_1) \\
& \times \frac{(-)^j \Gamma(1 + j + n_1 + n_2) \Gamma(4 - D + 2j + 2n_2)}{(j + n_2)! (j + n_1)! n_2! \Gamma(3 - \frac{D}{2} + 2j + n_1 + n_2)} \\
& + \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D - 3)\Gamma(4 - D)} x_{23}^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m3}^{j+n_1+n_2} x_{13}^{n_1} x_{23}^{-j-n_1} \frac{\Gamma(2 - \frac{D}{2} + j + n_1)}{\Gamma(2 - \frac{D}{2} + j)}
\end{aligned}$$

$$\begin{aligned}
& \times \frac{(-)^{n_1} \Gamma(1 + n_1 + n_2) \Gamma(4 - D + 2j + 2n_1 + 2n_2)}{(j + n_1 + n_2)! n_1! n_2! \Gamma(3 - \frac{D}{2} + j + 2n_1 + n_2)} \\
& + \frac{i(-)^{D/2-3} (p_3^2)^{D/2-3} \Gamma(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} x_{23}^{D/2-2} \\
& \times \sum_{j=0}^{\infty} \sum_{n_1=1}^{\infty} \xi_{m_3}^{j+n_1} x_{13}^{n_1} x_{23}^{-n_1} \\
& \times \frac{(-)^{n_1} \Gamma(4 - D + 2j + 2n_1) \Gamma(2 - \frac{D}{2} + n_1)}{n_1! j! \Gamma(3 - \frac{D}{2} + j + 2n_1)} \\
& + \frac{i(-)^{D/2-3} (p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} x_{23}^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \xi_{m_3}^j x_{13}^{n_1} \frac{\Gamma(1 + j + n_1) \Gamma(4 - D + 2j)}{(j!)^2 n_1! \Gamma(3 - \frac{D}{2} + j + n_1)} \\
& \times \Gamma(\frac{D}{2} - 1 + n_1) \\
& + \frac{i(-)^{D/2-3} (p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} x_{23}^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \xi_{m_3}^j \frac{\Gamma(4 - D + 2j)}{j! \Gamma(3 - \frac{D}{2} + j)}. \tag{A4}
\end{aligned}$$

$$\begin{aligned}
C_{0,6}^b(p_1^2, p_2^2, p_3^2, m^2) &= \frac{i(-)^{D/2-3} (p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} x_{13}^{D/2-2} \\
& \times \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m_3}^j x_{13}^{-j+n_1} x_{23}^{n_2} \\
& \times \frac{\Gamma(1 + n_1 + n_2) \Gamma(4 - D + 2j) \Gamma(\frac{D}{2} - 1 - j + n_1 + n_2)}{j! n_1! n_2! \Gamma(3 - \frac{D}{2} + j + n_2) \Gamma(\frac{D}{2} - 1 - j + n_1)} \\
& = \frac{i(-)^{D/2-3} (p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3)} x_{13}^{D/2-2} \\
& \times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} x_{13}^{n_1} x_{23}^{n_2} \frac{\Gamma(1 + n_1 + n_2) \Gamma(\frac{D}{2} - 1 + n_1 + n_2)}{n_1! n_2! \Gamma(\frac{D}{2} - 1 + n_1) \Gamma(3 - \frac{D}{2} + n_2)} \\
& + \frac{i(-)^{D/2-3} (p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} x_{13}^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_2=0}^{\infty} \sum_{n_1=1}^{\infty} \xi_{m_3}^j x_{13}^{n_1} x_{23}^{n_2} \\
& \times \frac{\Gamma(1 + j + n_1 + n_2) \Gamma(4 - D + 2j) \Gamma(\frac{D}{2} - 1 + n_1 + n_2)}{j! n_2! (j + n_1)! \Gamma(3 - \frac{D}{2} + j + n_2) \Gamma(\frac{D}{2} - 1 + n_1)} \\
& + \frac{i(-)^{D/2-3} (p_3^2)^{D/2-3} \Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} x_{13}^{D/2-2}
\end{aligned}$$

$$\begin{aligned}
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=1}^{\infty} \xi_{m_3}^{j+n_1} x_{13}^{-j} x_{23}^{j+n_2} \Gamma(2 - \frac{D}{2} + j) \Gamma(\frac{D}{2} - 1 + n_2) \\
& \times \frac{(-)^j \Gamma(1 + j + n_1 + n_2) \Gamma(4 - D + 2j + 2n_1)}{(j + n_1)! (j + n_2)! n_1! \Gamma(3 - \frac{D}{2} + 2j + n_1 + n_2)} \\
& + \frac{i(-)^{D/2-3} (p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} x_{13}^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m_3}^{j+n_1+n_2} x_{13}^{-j-n_2} x_{23}^{n_2} \frac{\Gamma(2 - \frac{D}{2} + j + n_2)}{\Gamma(2 - \frac{D}{2} + j)} \\
& \times \frac{(-)^{n_2} \Gamma(1 + n_1 + n_2) \Gamma(4 - D + 2j + 2n_1 + 2n_2)}{(j + n_1 + n_2)! n_1! n_2! \Gamma(3 - \frac{D}{2} + j + n_1 + 2n_2)} \\
& + \frac{i(-)^{D/2-3} (p_3^2)^{D/2-3} \Gamma(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} x_{13}^{D/2-2} \\
& \times \sum_{j=0}^{\infty} \sum_{n_2=1}^{\infty} \xi_{m_3}^{j+n_2} x_{13}^{-n_2} x_{23}^{n_2} \\
& \times \frac{(-)^{n_2} \Gamma(4 - D + 2j + 2n_2) \Gamma(2 - \frac{D}{2} + n_2)}{n_2! j! \Gamma(3 - \frac{D}{2} + j + 2n_2)} \\
& + \frac{i(-)^{D/2-3} (p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} x_{13}^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_2=1}^{\infty} \xi_{m_3}^j x_{23}^{n_2} \\
& \times \frac{\Gamma(1 + j + n_2) \Gamma(4 - D + 2j) \Gamma(\frac{D}{2} - 1 + n_2)}{(j!)^2 n_2! \Gamma(3 - \frac{D}{2} + j + n_2)} \\
& + \frac{i(-)^{D/2-3} (p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} x_{13}^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \xi_{m_3}^j \frac{\Gamma(4 - D + 2j)}{j! \Gamma(3 - \frac{D}{2} + j)}. \tag{A5}
\end{aligned}$$

$$\begin{aligned}
C_{0,7}^b(p_1^2, p_2^2, p_3^2, m^2) &= - \frac{i(-)^{D/2-3} (p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3) \Gamma(4-D)} (x_{13} x_{23})^{D/2-2} \\
& \times \sum_{j=0}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m_3}^j x_{13}^{-j+n_1} x_{23}^{-j+n_2} \frac{\Gamma(4 - D + 2j)}{j! n_1! n_2!} \\
& \times \frac{\Gamma(\frac{D}{2} - 1 - j + n_1 + n_2) \Gamma(D - 3 - 2j + n_1 + n_2)}{\Gamma(\frac{D}{2} - 1 - j + n_1) \Gamma(\frac{D}{2} - 1 - j + n_2)} \\
& = - \frac{i(-)^{D/2-3} (p_3^2)^{D/2-3} \Gamma^2(2 - \frac{D}{2}) \Gamma^2(\frac{D}{2} - 1)}{(4\pi)^{D/2} \Gamma(D-3)} (x_{13} x_{23})^{D/2-2} \\
& \times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} x_{13}^{n_1} x_{23}^{n_2} \frac{\Gamma(\frac{D}{2} - 1 + n_1 + n_2) \Gamma(D - 3 + n_1 + n_2)}{n_1! n_2! \Gamma(\frac{D}{2} - 1 + n_1) \Gamma(\frac{D}{2} - 1 + n_2)}
\end{aligned}$$

$$\begin{aligned}
& - \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma(2 - \frac{D}{2})\Gamma(\frac{D}{2} - 1)}{\Gamma(D - 3)\Gamma(4 - D)} (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_2=0}^{\infty} \sum_{n_1=1}^{\infty} \xi_{m3}^{j+n_2} x_{13}^{j+n_1} x_{23}^{-j} \frac{(-)^j \Gamma(4 - D + 2j + 2n_2)}{(j + n_2)!(2j + n_1 + n_2)!n_2!} \\
& \times \frac{\Gamma(\frac{D}{2} - 1 + j + n_1 + n_2)\Gamma(D - 3 + n_1)\Gamma(2 - \frac{D}{2} + j)}{\Gamma(\frac{D}{2} - 1 + j + n_1)} \\
& - \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma^2(2 - \frac{D}{2})\Gamma^2(\frac{D}{2} - 1)}{\Gamma(D - 3)\Gamma(4 - D)} (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \xi_{m3}^j x_{13}^{n_1} x_{23}^{n_2} \Gamma(4 - D + 2j) \\
& \times \frac{\Gamma(\frac{D}{2} - 1 + j + n_1 + n_2)\Gamma(D - 3 + n_1 + n_2)}{j!(j + n_1)!(j + n_2)!\Gamma(\frac{D}{2} - 1 + n_1)\Gamma(\frac{D}{2} - 1 + n_2)} \\
& - \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma(2 - \frac{D}{2})\Gamma(\frac{D}{2} - 1)}{\Gamma(D - 3)\Gamma(4 - D)} (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=1}^{\infty} \xi_{m3}^{j+n_1} x_{13}^{-j} x_{23}^{j+n_2} \frac{(-)^j \Gamma(4 - D + 2j + 2n_1)}{(j + n_1)!n_1!(2j + n_1 + n_2)!} \\
& \times \frac{\Gamma(\frac{D}{2} - 1 + j + n_1 + n_2)\Gamma(D - 3 + n_2)\Gamma(2 - \frac{D}{2} + j)}{\Gamma(\frac{D}{2} - 1 + j + n_2)} \\
& - \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \Gamma(2 - \frac{D}{2})\Gamma(\frac{D}{2} - 1) (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=1}^{\infty} \xi_{m3}^{j+n_1+n_2} x_{13}^{-j-n_2} x_{23}^{n_2} \frac{\Gamma(\frac{D}{2} - 1 + n_1 + n_2)}{\Gamma(4 - D + j)} \\
& \times \frac{(-)^{n_2} \Gamma(4 - D + 2j + 2n_1 + 2n_2)\Gamma(2 - \frac{D}{2} + j + n_2)}{(j + n_1 + n_2)!n_1!(j + n_1 + 2n_2)!\Gamma(\frac{D}{2} - 1 + n_2)} \\
& - \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \Gamma(2 - \frac{D}{2})\Gamma(\frac{D}{2} - 1) (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m3}^{j+n_1+n_2} x_{13}^{-j-n_2} x_{23}^{-j-n_1} \frac{\Gamma(2 - \frac{D}{2} + j + n_1)}{\Gamma(2 - \frac{D}{2} + j)} \\
& \times \frac{(-)^j \Gamma(4 - D + 2j + 2n_1 + 2n_2)\Gamma(2 - \frac{D}{2} + j + n_2)}{(j + n_1 + n_2)!n_1!n_2!\Gamma(4 - D + 2j + n_1 + n_2)} \\
& - \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \xi_{m3}^{j+n_1+n_2} x_{13}^{-j} x_{23}^{-n_1} \frac{\Gamma(2 - \frac{D}{2} + j)}{\Gamma(4 - D + j + n_1)} \\
& \times \frac{\Gamma(4 - D + 2j + 2n_1 + 2n_2)\Gamma(\frac{D}{2} - 1 + n_2)\Gamma(2 - \frac{D}{2} + n_1)}{(j + n_1 + n_2)!(n_1 + n_2)!(j + n_2)!}
\end{aligned}$$

$$\begin{aligned}
& - \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1) (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_2=0}^{\infty} \sum_{n_1=1}^{\infty} \xi_{m3}^{j+n_1+n_2} x_{13}^{n_1} x_{23}^{-j-n_1} \frac{\Gamma(2 - \frac{D}{2} + j + n_1)}{\Gamma(4 - D + j)} \\
& \times \frac{(-)^{n_1} \Gamma(4 - D + 2j + 2n_1 + 2n_2) \Gamma(\frac{D}{2} - 1 + n_1 + n_2)}{(j + n_1 + n_2)! (j + 2n_1 + n_2)! n_2! \Gamma(\frac{D}{2} - 1 + n_1)} \\
& - \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \Gamma(\frac{D}{2} - 1) (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \xi_{m3}^{j+n_1} x_{13}^{-j} x_{23}^{-n_1} \frac{\Gamma(2 - \frac{D}{2} + j) \Gamma(2 - \frac{D}{2} + n_1)}{j! n_1! (j + n_1)!} \\
& \times \frac{\Gamma(4 - D + 2j + 2n_1)}{\Gamma(4 - D + j + n_1)} \\
& - \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1)}{\Gamma(4 - D)} (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \xi_{m3}^{j+n_1} x_{13}^{-j} x_{23}^j \Gamma(4 - D + 2j + 2n_1) \\
& \times \frac{(-)^j \Gamma(\frac{D}{2} - 1 + j + n_1) \Gamma(2 - \frac{D}{2} + j)}{(j + n_1)! n_1! \Gamma(1 + 2j + n_1) \Gamma(\frac{D}{2} - 1 + j)} \\
& - \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1)}{\Gamma(4 - D)} (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=0}^{\infty} \sum_{n_1=1}^{\infty} \xi_{m3}^{j+n_1} x_{13}^{n_1} x_{23}^{-n_1} \Gamma(4 - D + 2j + 2n_1) \\
& \times \frac{(-)^{n_1} \Gamma(\frac{D}{2} - 1 + j + n_1) \Gamma(2 - \frac{D}{2} + n_1)}{(j + n_1)! (j + 2n_1)! \Gamma(1 + j) \Gamma(\frac{D}{2} - 1 + n_1)} \\
& - \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \Gamma(2 - \frac{D}{2}) (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=1}^{\infty} \xi_{m3}^{j+n_1} x_{13}^{-j} \Gamma(4 - D + 2j + 2n_1) \\
& \times \frac{\Gamma(\frac{D}{2} - 1 + n_1) \Gamma(2 - \frac{D}{2} + j)}{((j + n_1)!)^2 n_1! \Gamma(4 - D + j)} \\
& - \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma^2(2 - \frac{D}{2}) \Gamma(\frac{D}{2} - 1)}{\Gamma(D - 3) \Gamma(4 - D)} (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_1=0}^{\infty} \xi_{m3}^j x_{13}^{n_1} \Gamma(4 - D + 2j) \\
& \times \frac{\Gamma(\frac{D}{2} - 1 + j + n_1) \Gamma(D - 3 + n_1)}{(j!)^2 (j + n_1)! \Gamma(\frac{D}{2} - 1 + n_1)}
\end{aligned}$$

$$\begin{aligned}
& -\frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}}\Gamma(2-\frac{D}{2})(x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_2=1}^{\infty} \xi_{m_3}^{j+n_2} x_{23}^{-j} \Gamma(4-D+2j+2n_2) \\
& \times \frac{\Gamma(\frac{D}{2}-1+n_2)\Gamma(2-\frac{D}{2}+j)}{((j+n_2)!)^2 n_2! \Gamma(4-D+j)} \\
& -\frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma^2(2-\frac{D}{2})\Gamma(\frac{D}{2}-1)}{\Gamma(D-3)\Gamma(4-D)} (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \sum_{n_2=0}^{\infty} \xi_{m_3}^j x_{23}^{n_2} \Gamma(4-D+2j) \\
& \times \frac{\Gamma(\frac{D}{2}-1+j+n_2)\Gamma(D-3+n_2)}{(j!)^2 (j+n_2)! \Gamma(\frac{D}{2}-1+n_2)} \\
& + \frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \frac{\Gamma^2(2-\frac{D}{2})}{\Gamma(4-D)} (x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \xi_{m_3}^j \frac{\Gamma(4-D+2j)\Gamma(\frac{D}{2}-1+j)}{(j!)^3} \\
& -\frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \Gamma(2-\frac{D}{2})\Gamma(\frac{D}{2}-1)(x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \xi_{m_3}^j x_{23}^{-j} \frac{\Gamma(4-D+2j)\Gamma(2-\frac{D}{2}+j)}{(j!)^2 \Gamma(4-D+j)} \\
& -\frac{i(-)^{D/2-3}(p_3^2)^{D/2-3}}{(4\pi)^{D/2}} \Gamma(2-\frac{D}{2})\Gamma(\frac{D}{2}-1)(x_{13}x_{23})^{D/2-2} \\
& \times \sum_{j=1}^{\infty} \xi_{m_3}^j x_{13}^{-j} \frac{\Gamma(4-D+2j)\Gamma(2-\frac{D}{2}+j)}{(j!)^2 \Gamma(4-D+j)}. \tag{A6}
\end{aligned}$$

Appendix B: Some coefficients in ε expansions

$$\begin{aligned}
A_{n_1, n_2}^{(a)}(x, y) &= 4\gamma_E \ln x \ln y + \ln^2 x \ln y + \ln x \ln^2 y + 8\psi^3(1+n_1+n_2) \\
& + 8\psi^2(1+n_1+n_2) \left[2\gamma_E + \ln x + \ln y - \psi(1+n_2) \right] \\
& - 8\gamma_E \ln x \psi(1+n_2) - 2\ln^2 x \psi(1+n_2) - 2\ln x \ln y \psi(1+n_2) \\
& + 8\gamma_E \psi'(1+n_1+n_2) + 4\ln x \psi'(1+n_1+n_2) + 4\ln y \psi'(1+n_1+n_2) \\
& - 4\psi(1+n_2) \psi'(1+n_1+n_2) - 2\psi(1+n_1) \left[4\gamma_E \ln y + \ln x \ln y \right. \\
& \left. + \ln^2 y + 4\psi(1+n_1+n_2)^2 + 2\psi(1+n_1+n_2) \left(4\gamma_E + \ln x + 2\ln y \right) \right]
\end{aligned}$$

$$\begin{aligned}
& -2\psi(1+n_2) \Big) - 2(4\gamma_E + \ln x + \ln y)\psi(1+n_2) + 2\psi'(1+n_1+n_2) \Big] \\
& + 2\psi(1+n_1+n_2) \Big[4\gamma_E \ln x + \ln^2 x + 4\gamma_E \ln y + 3 \ln x \ln y + \ln^2 y \\
& - 2(4\gamma_E + 2 \ln x + \ln y)\psi(1+n_2) + 6\psi'(1+n_1+n_2) \Big] \\
& + 2\psi''(1+n_1+n_2) ,
\end{aligned}$$

$$\begin{aligned}
A'_{j,n_1,n_2}^{(a)}(x,y) &= \ln x \ln y + 4\psi^2(1+j+n_1+n_2) - \ln x \psi(1+j+n_2) - \ln y \psi(1+n_1) \\
& + \psi(1+j+n_2)\psi(1+n_1) + 2\psi(1+j+n_1+n_2) \Big[\ln(xy) - \psi(1+j+n_2) \\
& - \psi(1+n_1) - \psi(1+n_2) \Big] - \ln x \psi(1+n_2) + \psi(1+n_1)\psi(1+n_2) \\
& + \psi(1+j+n_1) \Big[-\ln y - 2\psi(1+j+n_1+n_2) + \psi(1+j+n_2) \\
& + \psi(1+n_2) \Big] + 2\psi'(1+j+n_1+n_2) ,
\end{aligned}$$

$$B_{j,n_1,n_2}^{(a)} = -\ln x_{13} + \psi(1+j+n_1+n_2) + \psi(1+n_1) - 2\psi(1+n_1+n_2) ,$$

$$C_{j,n_1,n_2}^{(a)} = -\ln x_{23} + \psi(1+j+n_1+n_2) + \psi(1+n_2) - 2\psi(1+n_1+n_2) ,$$

$$\begin{aligned}
D_{j,n_1,n_2}^{(a)} &= \ln^2(x_{13}x_{23}) - \ln^2 \xi_{33} + 8\psi^2(j) + \psi^2(j+n_1) \\
& - \psi(j+n_1+n_2) \Big[4 \ln(x_{13}x_{23}) - 2 \ln \xi_{33} - 3\psi(j+n_1+n_2) \Big] \\
& + \psi(1+j+n_1+n_2) \Big[2 \ln \xi_{33} - 2\psi(j+n_1+n_2) - \psi(1+j+n_1+n_2) \Big] \\
& + 2\psi(j) \Big[3 \ln(x_{13}x_{23}) - \ln \xi_{33} - 3\psi(j+n_1) - 5\psi(j+n_1+n_2) \\
& + \psi(1+j+n_1+n_2) - 3\psi(j+n_2) \Big] - 4\psi'(j) + 3\psi'(j+n_1+n_2) \\
& - 2\psi(j+n_1) \Big[\ln(x_{13}x_{23}) - 2\psi(j+n_1+n_2) - \psi(j+n_2) \Big] \\
& - \psi(j+n_2) \Big[2 \ln(x_{13}x_{23}) - 4\psi(j+n_1+n_2) - \psi(j+n_2) \Big] \\
& + \psi'(j+n_1) + \psi'(1+j+n_1+n_2) + \psi'(j+n_2) ,
\end{aligned}$$

$$\begin{aligned}
E_{j,n_1,n_2}^{(a)} &= \ln x_{31} \ln x_{21} + \psi^2(1+j+n_1+n_2) - 2 \ln x_{21} \psi(1+n_1) \\
& + \ln x_{31} \psi(1+n_1+n_2) + \ln x_{21} \psi(1+n_1+n_2) \\
& - 2\psi(1+n_1)\psi(1+n_1+n_2) + \psi^2(1+n_1+n_2) \\
& - \psi(1+j+n_2) \Big[\ln x_{31} - 2\psi(1+n_1) + \psi(1+n_1+n_2) \Big] \\
& + \psi(1+j+n_1+n_2) \Big[\ln(x_{31}x_{21}) - \psi(1+j+n_2) - 2\psi(1+n_1) \\
& + 2\psi(1+n_1+n_2) - \psi(1+n_2) \Big] - \ln x_{31} \psi(1+n_2) \\
& + 2\psi(1+n_1)\psi(1+n_2) - \psi(1+n_1+n_2)\psi(1+n_2) \\
& + \psi'(1+j+n_1+n_2) + \psi'(1+n_1+n_2) ,
\end{aligned}$$

$$\begin{aligned}
G_{j,n_1,n_2}^{(a)} &= 2\pi^2 + \ln^2 x_{31} + 2 \ln x_{31} \ln x_{21} - \ln^2 x_{21} + \ln^2 \xi_{31} + 2\psi^2(j) - \psi^2(j+n_1) \\
&\quad - \psi(j+n_1+n_2) \left[4 \ln x_{31} - 4 \ln x_{21} + 2 \ln \xi_{31} + 3\psi(j+n_1+n_2) \right] \\
&\quad - \psi(1+j+n_1+n_2) \left[2 \ln \xi_{31} - 2\psi(j+n_1+n_2) - \psi(1+j+n_1+n_2) \right] \\
&\quad - 2\psi(1+n_1) \left[\ln x_{31} + 3 \ln x_{21} - \ln \xi_{31} - 5\psi(j+n_1+n_2) \right. \\
&\quad \left. + \psi(1+j+n_1+n_2) \right] + \psi(1+n_1+n_2) \left[2 \ln(x_{31}x_{21}) - 4\psi(j+n_1+n_2) \right. \\
&\quad \left. - 2\psi(1+n_1) + \psi(1+n_1+n_2) \right] - 2\psi(j+n_1) \left[\ln(x_{31}/x_{21}) \right. \\
&\quad \left. + 2\psi(j+n_1+n_2) - 3\psi(1+n_1) + \psi(1+n_1+n_2) \right] \\
&\quad + 4\psi(j) \left[\ln x_{31} - 2\psi(1+n_1) + \psi(1+n_1+n_2) \right] - 2\psi'(j) - \psi'(j+n_1) \\
&\quad - 3\psi'(j+n_1+n_2) - \psi'(1+j+n_1+n_2) + \psi'(1+n_1+n_2) , \\
H_{j,n_1,n_2}^{(a)} &= \ln x_{31}^2 - 2 \ln x_{31} \ln \xi_{31} - \psi(1+j+n_1+n_2) \left[2 \ln x_{31} + 2 \ln \xi_{31} - 6\psi(1+n_1) \right. \\
&\quad \left. + 3\psi(1+j+n_1+n_2) - 4\psi(1+j+n_2) + 2\psi(1+n_1+n_2) \right] \\
&\quad + \psi(1+n_1) \left[4 \ln \xi_{31} - 2 \ln x_{31} - 8\psi(1+j+n_2) \right] + 4 \ln x_{31} \psi(1+j+n_2) \\
&\quad + \psi(1+n_1+n_2) \left[2 \ln x_{31} - 2 \ln \xi_{31} + 4\psi(1+j+n_2) - 2\psi(1+n_1) \right. \\
&\quad \left. + \psi(1+n_1+n_2) \right] - 6\psi(j) \left[\ln x_{31} + \psi(1+j+n_1+n_2) - 2\psi(1+n_1) \right. \\
&\quad \left. + \psi(1+n_1+n_2) \right] + 2\psi(1+j) \left[\ln x_{31} + \psi(1+j+n_1+n_2) - 2\psi(1+n_1) \right. \\
&\quad \left. + \psi(1+n_1+n_2) \right] - 3\psi'(1+j+n_1+n_2) + \psi'(1+n_1+n_2) , \\
P_{j,n_1,n_2}^{(a)} &= \ln x_{31}^2 + 2 \ln x_{31} \ln \xi_{31} + \psi^2(1+j+n_1) + \psi^2(1+j+n_1+n_2) \\
&\quad - 2 \ln x_{31} \psi(1+n_1) - 4 \ln \xi_{31} \psi(1+n_1) - 2 \ln x_{31} \psi(n_2) + 4\psi(1+n_1)\psi(n_2) \\
&\quad + 2\psi(1+j+n_1+n_2) \left[\ln x_{31} + \ln \xi_{31} - \psi(1+n_1) - \psi(n_2) - \psi(1+n_2) \right] \\
&\quad + 2\psi(1+j+n_1) \left[\ln(x_{31}\xi_{31}) + \psi(1+j+n_1+n_2) - \psi(1+n_1) - \psi(n_2) \right. \\
&\quad \left. - \psi(1+n_2) \right] - 2 \ln x_{31} \psi(1+n_2) + 4\psi(1+n_1)\psi(1+n_2) \\
&\quad + \psi'(1+j+n_1) + \psi'(1+j+n_1+n_2) , \\
Q_{j,n_1,n_2}^{(a)} &= \ln^2 \eta_{33} + \psi^2(1+j+n_1) - \psi^2(1+j+n_1+n_2) - 4 \ln \eta_{33} \psi(2+j+n_1+n_2) \\
&\quad - 2\psi(1+j+n_1+n_2)\psi(2+j+n_1+n_2) + 3\psi^2(2+j+n_1+n_2) \\
&\quad + 2 \ln \eta_{33} \psi(1+j+n_2) - 4\psi(2+j+n_1+n_2)\psi(1+j+n_2) + \psi^2(1+j+n_2) \\
&\quad - 2\psi(1+j) \left[\ln \eta_{33} + \psi(1+j+n_1) - \psi(1+j+n_1+n_2) \right. \\
&\quad \left. - 3\psi(2+j+n_1+n_2) + \psi(1+j+n_2) \right]
\end{aligned}$$

$$\begin{aligned}
& +2\psi(1+j+n_1)\left[\ln \eta_{33} - 2\psi(2+j+n_1+n_2) + \psi(1+j+n_2)\right] \\
& +\psi'(1+j+n_1) - \psi'(1+j+n_1+n_2) - 3\psi'(2+j+n_1+n_2) \\
& +\psi'(1+j+n_2)
\end{aligned} \tag{B1}$$

$$\begin{aligned}
A_{j,n_1,n_2}^{(b)} &= \ln x_{13} \ln x_{23} + \psi^2(1+j+n_1+n_2) - \ln x_{13} \psi(1+j+n_2) \\
& - \ln x_{23} \psi(1+n_1) + \psi(1+j+n_2)\psi(1+n_1) + \ln x_{13} \psi(1+n_1+n_2) \\
& + \ln x_{23} \psi(1+n_1+n_2) - \psi(1+j+n_2)\psi(1+n_1+n_2) \\
& - \psi(1+n_1)\psi(1+n_1+n_2) + \psi^2(1+n_1+n_2) - \psi(1+j+n_1)\left[\ln x_{23} \right. \\
& \left. + \psi(1+j+n_1+n_2) - \psi(1+j+n_2) + \psi(1+n_1+n_2) - \psi(1+n_2)\right] \\
& + \psi(1+j+n_1+n_2)\left[\ln x_{13} + \ln x_{23} - \psi(1+j+n_2) - \psi(1+n_1) \right. \\
& \left. + 2\psi(1+n_1+n_2) - \psi(1+n_2)\right] - \ln x_{13} \psi(1+n_2) + \psi(1+n_1)\psi(1+n_2) \\
& - \psi(1+n_1+n_2)\psi(1+n_2) + \psi'(1+j+n_1+n_2) + \psi'(1+n_1+n_2) , \\
B_{j,n_1,n_2}^{(b)} &= 2\pi^2 + \ln^2 x_{13} + 2 \ln x_{13} \ln x_{23} - \ln^2 x_{23} + \ln^2 \xi_{m3} + 2\psi^2(j) - \psi^2(j+n_1) \\
& - 4 \ln x_{13} \psi(2j+2n_1+2n_2) + 4 \ln x_{23} \psi(2j+2n_1+2n_2) \\
& - 4\psi^2(2j+2n_1+2n_2) - 2 \ln \xi_{m3} \psi(1+j+n_1+n_2) + \psi^2(1+j+n_1+n_2) \\
& - 4 \ln x_{23} \psi(1+j+2n_1+n_2) + 8\psi(2j+2n_1+2n_2)\psi(1+j+2n_1+n_2) \\
& - 2\psi^2(1+j+2n_1+n_2) - 2 \ln x_{13} \psi(1+n_1) - 2 \ln x_{23} \psi(1+n_1) \\
& + 4\psi(2j+2n_1+2n_2)\psi(1+n_1) + \psi^2(1+n_1) + 2 \ln x_{13} \psi(1+n_1+n_2) \\
& + 2 \ln x_{23} \psi(1+n_1+n_2) - 4\psi(2j+2n_1+2n_2)\psi(1+n_1+n_2) \\
& - 2\psi(1+n_1)\psi(1+n_1+n_2) + \psi^2(1+n_1+n_2) - 2\psi(j+n_1)\left[\ln x_{13} \right. \\
& \left. - \ln x_{23} + 2\psi(2j+2n_1+2n_2) - 2\psi(1+j+2n_1+n_2) - \psi(1+n_1) \right. \\
& \left. + \psi(1+n_1+n_2)\right] + 4\psi(j)\left[\ln x_{13} - \psi(1+j+2n_1+n_2) - \psi(1+n_1) \right. \\
& \left. + \psi(1+n_1+n_2)\right] - 2\psi'(j) - \psi'(j+n_1) - 4\psi'(2j+2n_1+2n_2) \\
& - \psi'(1+j+n_1+n_2) + 2\psi'(1+j+2n_1+n_2) - \psi'(1+n_1) \\
& + \psi'(1+n_1+n_2) , \\
C_{j,n_1,n_2}^{(b)} &= B_{j,n_2,n_1}^{(b)}(x_{13} \leftrightarrow x_{23}) , \\
D_{j,n_1,n_2}^{(b)} &= \ln^2(x_{13}x_{23}) - \ln^2 x_{m3} + 4\psi^2(2j+n_1+n_2) + \psi^2(j)
\end{aligned}$$

$$\begin{aligned}
& -2 \ln x_{13} \psi(j + n_1) - 2 \ln x_{23} \psi(j + n_1) + \psi^2(j + n_1) \\
& -4 \ln x_{13} \psi(2j + 2n_1 + 2n_2) - 4 \ln x_{23} \psi(2j + 2n_1 + 2n_2) \\
& +4\psi(j + n_1)\psi(2j + 2n_1 + 2n_2) + 4\psi^2(2j + 2n_1 + 2n_2) \\
& +2 \ln x_{m3} \psi(1 + j + n_1 + n_2) - \psi^2(1 + j + n_1 + n_2) \\
& +2\psi(j) \left[\ln(x_{13}x_{23}) - \psi(j + n_1) - 2\psi(2j + 2n_1 + 2n_2) - \psi(j + n_2) \right] \\
& +4\psi(2j + n_1 + n_2) \left[\ln(x_{13}x_{23}) + \psi(j) - \psi(j + n_1) \right. \\
& \left. -2\psi(2j + 2n_1 + 2n_2) - \psi(j + n_2) \right] - 2 \ln(x_{13}x_{23})\psi(j + n_2) \\
& +2\psi(j + n_1)\psi(j + n_2) + 4\psi(2j + 2n_1 + 2n_2)\psi(j + n_2) + \psi^2(j + n_2) \\
& -4\psi'(2j + n_1 + n_2) - \psi'(j) + \psi'(j + n_1) + 4\psi'(2j + 2n_1 + 2n_2) \\
& +\psi'(1 + j + n_1 + n_2) + \psi'(j + n_2) , ,
\end{aligned}$$

$$E_{j,n}^{(b)}(x) = \gamma_E - \ln x - \psi(1 + j + n) + \psi(1 + j + 2n) + \psi(1 + n) ,$$

$$G_{j,n}^{(b)}(x) = \gamma_E - \ln x + \psi(j) - 2\psi(j + n) + 2\psi(2j + 2n) + \psi(n) . \quad (\text{B2})$$

-
- [1] CMS Collaboration, Phys. Lett. **B716**(2012)30.
- [2] ATLAS Collaboration, Phys. Lett. **B716**(2012)1.
- [3] CEPC-SPPC study group, *CEPC-SPPC preliminary conceptual design report. 1. Physics and detector*, IHEP-CEPC-DR-2015-01, 2015.
- [4] T. Behnke et al., *The International Linear Collider Technical Design Report - Volume I: Executive Summary*, arXiv:1306.6327 [physics.acc-ph].
- [5] G. Apollinari, et al., *High-Luminosity Large Hadron Collider (HL-LHC): Preliminary Design Report*, Technical Report CERN-2015-005 (2015).
- [6] V. A. Smirnov, *Analytic Tools for Feynman Integrals*, (Springer, Heidelberg 2012), and references therein.
- [7] G. t'Hooft, M. Veltman, Nucl. Phys. **B153**(1979)365.
- [8] J. Collins, *Renormalization*, Cambridge Univ. Press, 1984.
- [9] G. Passarino, M. Veltman, Nucl. Phys. **B160**(1979)151.
- [10] E. E. Boos, and A. I. Davydychev, Vestn. Mosk. Univ. **28**(1987)8.
- [11] A. I. Davydychev, and J. B. Tausk, Nucl. Phys. **B397**(1993)123.

- [12] A. I. Davydychev, Phys. Rev. **D61**(2000)087701.
- [13] E. E. Boos, and A. I. Davydychev, Theor. Math. Phys. **89**(1991)1052.
- [14] A. I. Davydychev, J. Phys. **A25**(1992)5587.
- [15] A. I. Davydychev, J. Math. Phys.**33**(1992)358.
- [16] A. I. Davydychev, J. Math. Phys.**32**(1991)1052.
- [17] A. I. Davydychev, Nucl. Instrum. Meth. **A559**(2006)293.
- [18] O. V. Tarasov, Nucl. Phys. B(Proc. Suppl)**89**(2000)237.
- [19] J. Fleischer, F. Jegerlehner, and O. V. Tarasov, Nucl. Phys. **B672**(2003)303.
- [20] M. E. Taylor, *Partial differential equations* (Springer, Heidelberg 2012).
- [21] N. I. Ussyukina, and A. I. Davydychev, Phys. Lett. **B298**(1993)363.
- [22] V. A. Smirnov, Phys. Lett. **B460**(1999)397.
- [23] J. B. Tausk, Phys. Lett. **B469**(1999)225.
- [24] M. Y. Kalmykov, and B. A. Kniehl, Phys. Lett. **B714**(2012)103.
- [25] L. J. Slater, *Generalised Hypergeometric Functions* (Cambridge University Press 1966).
- [26] J. Leray, *Le Calcul Différentiel et Intégral sur Une Variété Analytique Complexe*, Bull. Soc. Math., **87**(1959)81.
- [27] J. Horn, Math. Ann. **34**(1889)544.
- [28] T.-F. Feng, C.-H. Chang, J.-B. Chen, Z.-H. Gu, and H.-B. Zhang, Nucl. Phys. **B927**(2018)516.
- [29] V. V. Bytev, M. Y. Kalmykov, and B. A. Kniehl, Nucl. Phys. **B836**(2010)129.
- [30] V. V. Bytev, M. Y. Kalmykov, and B. A. Kniehl, Comput. Phys. Commun. **184**(2013)2332.
- [31] M. Y. Kalmykov, and B. A. Kniehl, JHEP**1707**(2017)031.
- [32] X. C. Wang, *Finite element method* (Tsinghua University Press 2003, in Chinese).