

## ASAS-SN IDENTIFICATION OF FY SCT AS A DETACHED ECLIPSING BINARY SYSTEM WITH A $\sim 2.6$ YEAR PERIOD

T. Jayasinghe,<sup>1</sup> K. Z. Stanek,<sup>1, 2</sup> C. S. Kochanek,<sup>1, 2</sup> B. J. Shappee,<sup>3</sup> T. W. -S. Holoiien,<sup>4</sup> T. A. Thompson,<sup>1, 2</sup> J. L. Prieto,<sup>5, 6</sup> Subo Dong,<sup>7</sup> M. Pawlak,<sup>8</sup> and O. Pejcha<sup>8</sup>

<sup>1</sup>Department of Astronomy, The Ohio State University, 140 West 18th Avenue, Columbus, OH 43210, USA

<sup>2</sup>Center for Cosmology and Astroparticle Physics, The Ohio State University, 191 W. Woodruff Avenue, Columbus, OH 43210

<sup>3</sup>Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

<sup>4</sup>Carnegie Fellow, The Observatories of the Carnegie Institution for Science, 813 Santa Barbara St., Pasadena, CA 91101, USA

<sup>5</sup>Núcleo de Astronomía de la Facultad de Ingeniería, Universidad Diego Portales, Av. Ejército 441, Santiago, Chile

<sup>6</sup>Millennium Institute of Astrophysics, Santiago, Chile

<sup>7</sup>Kavli Institute for Astronomy and Astrophysics, Peking University, Yi He Yuan Road 5, Hai Dian District, China

<sup>8</sup>Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles University in Prague, Czech Republic


*Keywords:* binaries: eclipsing — stars: variables: general — surveys

The All-Sky Automated Survey for SuperNovae (ASAS-SN, Shappee et al. 2014; Kochanek et al. 2017) has monitored the entire visible sky to a depth of  $\sim 17$  mag in the V-band since 2014. ASAS-SN data is well suited for the discovery and study of variable stars (Jayasinghe et al. 2018a; Shields et al. 2018). In Jayasinghe et al. (2018b), we have uniformly analyzed the ASAS-SN light curves of  $\sim 412,000$  known variables in the VSX catalog (Watson et al. 2006), providing homogeneously classified samples of variable groups for further study.

Here we report the identification of a long period detached eclipsing binary system. ASASSN-V J184258.61–105928.4 (FY Sct) was discovered by Harwood (1960) with a spectral type of M6, and is currently classified as a semi-regular (SRB) variable in the VSX catalog. FY Sct ( $l, b = 22.184, -3.135$ ) was also classified as a variable source in the INTEGRAL-OMC catalog of variable sources (Alfonso-Garzón et al. 2012). The ASAS-SN data spanning 2014–2018 captures a single, deep ( $A \sim 1.5$  mag) eclipse. The similarity of this light curve to the recently identified long period detached eclipsing binary ASASSN-V J192543.72+402619.0 (Jayasinghe et al. 2018c) piqued our interest. We supplemented the ASAS-SN light curve with V-band data from the All-Sky Automated Survey (ASAS; Pojmanski 2002), extending the light curve back to the year 2000 (Figure 1). We calculated periodograms and arrived at an orbital period of  $P_{\text{orb}} \sim 939$  d. Based on this orbital period, we highlight the expected times of eclipse using the red dotted lines in the top panel of Figure 1. The ASAS data captures three eclipses. We see clear evidence for pulsational variability when the system is out-of-eclipse, with a period of  $P_{\text{pulse}} = 78$  d. This pulsational variability resembles the variability of semi-regular variables, and likely explains the current VSX classification as a SRB variable.

Multi-band photometry from ALLWISE (Wright, et al. 2010) and 2MASS (Skrutskie et al. 2006) provides colors of  $W1 - W2 = -0.125$  and  $J - K_s = 1.292$  typical of an M-giant (Zhong, et al. 2014). The probabilistic Gaia DR2 distance of  $d \sim 6$  kpc (Bailer-Jones et al. 2018; Gaia Collaboration et al. 2018) gives  $M_{K_s} = -5.7$ , which is also typical of a M-giant. The eclipse profile is suggestive of a disk eclipsing binary (Dong et al. 2014) rather than a stellar companion. The primary eclipse lasts  $\sim 73$  days, and the next eclipse is expected to begin on  $\sim$ Sept.28, 2018. We encourage further photometric and spectroscopic observations, particularly during the eclipse, to further understand this peculiar binary system.

# ASASSN-V J184258.61-105928.4



**Figure 1.** The top panel shows the combined ASAS-SN (blue) and ASAS (black) light curve for ASASSN-V J184258.61–105928.4. The middle panel shows the combined light curve phased with the orbital period of  $P_{\text{orb}} \sim 939$  d, and the bottom panel shows the combined light curve excluding the eclipses and phased with the pulsational period of  $P_{\text{pulse}} = 78$  d.

## REFERENCES

- Alfonso-Garzón, J., Domingo, A., Mas-Hesse, J. M., et al. 2012, *A&A*, 548, A79.
- Brown, T. M., Latham, D. W., Everett, M. E., & Esquerdo, G. A. 2011, *AJ*, 142, 112
- Dong, S., Katz, B., Prieto, J. L., et al. 2014, *ApJ*, 788, 41.
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, arXiv:1804.09365
- Bailer-Jones, C. A. L., Rybizki, J., Fouesneau, M., Mantelet, G., & Andrae, R. 2018, *AJ*, 156, 58
- Harwood, M. 1960, *Annalen van de Sterrewacht te Leiden*, 21, 387
- Jayasinghe, T., Kochanek, C. S., Stanek, K. Z., et al. 2018, *MNRAS*, 477, 3145
- Jayasinghe, T., Stanek, K. Z., Kochanek, C. S., et al. 2018, arXiv:1809.07329
- Jayasinghe, T., Stanek, K. Z., Kochanek, C. S., et al. 2018, *Research Notes of the American Astronomical Society*, 2, 125
- Kochanek, C. S., Shappee, B. J., Stanek, K. Z., et al. 2017, *PASP*, 129, 104502
- Pojmanski, G. 2002, *AcA*, 52, 397
- Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, *ApJ*, 788, 48
- Shields, J. V., Jayasinghe, T., Stanek, K. Z., et al. 2018, arXiv:1809.04075
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Watson, C. L., Henden, A. A., & Price, A. 2006, *Society for Astronomical Sciences Annual Symposium*, 25, 47
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, 140, 1868.
- Zhong, J., Lépine, S., Li, J., et al. 2014, *EAS Publications Series*, 409.