GENERIC ENDS OF THE MODULI SPACE OF $SL(n, \mathbb{C})$ -HIGGS BUNDLES

LAURA FREDRICKSON

ABSTRACT. Given a generic ray of Higgs bundles $(\overline{\partial}_E, t\varphi)$, we describe the corresponding family of hermitian metrics h_t solving Hitchin's equations via gluing methods. In the process, we construct a family of approximate solutions h_t^{app} which differ from the actual harmonic metrics h_t by error terms of size $e^{-\delta t}$. Such families of explicit approximate solutions have already proved useful for answering finer questions about the asymptotic geometry of the Hitchin moduli space.

1. INTRODUCTION

In this paper, we describe solutions of Hitchin's equations near the generic ends of the SU(n)-Hitchin moduli space by constructing good approximate solutions and perturbing them to actual solutions. Our paper generalizes Mazzeo-Swoboda-Weiss-Witt's results for the SU(2)-Hitchin moduli space [MSWW16, MSWW14], which has already been useful in their more recent work of the asymptotic geometry of the Hitchin moduli space [MSWW17]. Gaiotto-Moore-Neitzke give a conjectural description of the hyperkähler metric on the SU(n)-Hitchin moduli space [GMN09, GMN10], and our finer description of solutions of SU(n)-Hitchin's equations near the ends is a first step towards proving their conjecture. In fact, a number of conjectures from mathematicians and physics about \mathcal{M} remain open because they require a finer knowledge of the ends of the moduli space than provided by traditional algebro-geometric techniques alone. As demonstrated in [MSWW17], constructive analytic techniques complement these well, so we take this approach.

1.1. **Fixed data.** Fix $C = C(I, g_C, \omega)$ a compact Kähler curve of genus ≥ 2 with metric g_C , complex structure I, and symplectic form ω_C . Let K_C be the canonical line bundle. Fix $E \rightarrow C$ a complex vector bundle of rank n and degree d. Let Det E be the determinant line bundle. The groups Aut(E) and End(E) respectively denote the automorphisms and endomorphisms of the complex vector bundle E which induce the identity map on Det E.

Additionally fix a holomorphic structure, $\overline{\partial}_{\text{Det }E}$, and a hermitian structure, $h_{\text{Det }E}$, on the complex line bundle Det *E* such that $h_{\text{Det }E}$ is Hermitian-Einstein for the holomorphic line bundle (Det *E*, $\overline{\partial}_{\text{Det }E}$). We do not normalize the Riemannian volume $\text{vol}_{g_C}(C)$ of the curve *C*. Consequently, the Hermitian-Einstein condition states that the curvature of the

associated Chern connection $D = D(\overline{\partial}_{\text{Det }E}, h_{\text{Det }E})$ satisfies

$$F_D = -\sqrt{-1} \text{deg} E \frac{2\pi\omega_C}{\text{vol}_{g_C}(C)} \text{Id}_{\text{Det}\,E}.$$
(1.1)

Given this fixed data, let \mathcal{M} be the associated Hitchin moduli space. The Hitchin moduli space consists of triples $(\overline{\partial}_E, \varphi, h)$ solving Hitchin's equations up to complex gauge equivalence, defined in (1.4). Here,

- $\overline{\partial}_E$ is a holomorphic structure on *E*,
- $\varphi \in \Omega^{1,0}(C, \operatorname{End} E)$ is the Higgs field, and
- *h* is a hermitian metric on *E*.

Additionally, the induced holomorphic and hermitian structures on Det *E* must agree with the fixed structures $\overline{\partial}_{\text{Det }E}$ and $h_{\text{Det }E}$. We say that such a triple $(\overline{\partial}_E, \varphi, h)$ is a *solution of SU*(*n*)-*Hitchin's equations* if

$$\overline{\partial}_E \varphi = 0, \qquad F_{D(\overline{\partial}_E,h)}^{\perp} + [\varphi, \varphi^{\dagger_h}] = 0, \tag{1.2}$$

where $D(\overline{\partial}_E, h)$ is the Chern connection, $\varphi^{\dagger_h} \in \Omega^{0,1}(C, \operatorname{End} E)$ is the *h*-hermitian adjoint, and F_D^{\perp} denotes the trace-free part of the the curvature of *D*, i.e.

$$F_{D(\bar{\partial},h)}^{\perp} = F_{D(\bar{\partial}_{E},h)} + \sqrt{-1} \frac{\deg\left(E\right)}{\operatorname{rank}(E)} \frac{2\pi\omega_{C}}{\operatorname{vol}_{g_{C}}(C)} \operatorname{Id}_{E}.$$
(1.3)

We call such *h* the *harmonic metric* for the Higgs bundle $(\overline{\partial}_E, \varphi)$. A Higgs bundle $(\overline{\partial}_E, \varphi)$ admits a harmonic metric if, and only if, $(\overline{\partial}_E, \varphi)$ is polystable¹.

The action of the complex gauge group Aut(E) is as follows: given $g \in Aut(E)$,

$$g \cdot (\overline{\partial}_E, \varphi, h) = (g^{-1} \circ \overline{\partial}_E \circ g, g^{-1} \varphi g, g \cdot h), \quad \text{where } (g \cdot h)(v, w) = h(gv, gw). \quad (1.4)$$

Unitary formulation of Hitchin's equations. There's an equivalent unitary formation of Hitchin's equations. In this formulation, we additionally fix a hermitian metric h_0 on the complex vector bundle $E \rightarrow C$. Now, the Hitchin moduli space consists of pairs (d_A, Φ) , where

- d_A is a h_0 -unitary connection, and
- $\Phi \in \Omega^{1,0}(\mathbb{C}, \operatorname{End} E)$,

solving $\overline{\partial}_A \Phi = 0$ and $F_A^{\perp} + [\Phi, \Phi^{\dagger_{h_0}}] = 0$ —up to h_0 -unitary gauge equivalence.

We can pass back and between these two formulations. Given the pair (d_A, Φ) , we get the associated triple $(\overline{\partial}_A, \Phi, h_0)$. Conversely, given a triple $(\overline{\partial}_E, \varphi, h)$, there is an End *E*valued h_0 -hermitian section *H* such that $h(v, w) = h_0(Hv, w)$. Take the complex gauge

¹ A Higgs bundle ($\overline{\partial}_E, \varphi$) is *stable* if for all φ -invariant subbundles $F, \mu(F) < \mu(E)$; here, $\mu(F) := \frac{\deg(F)}{\operatorname{rank}(F)}$ is the slope of the bundle. A Higgs bundle is *polystable* if it is the direct sum of stable Higgs bundles of the same slope.

transformation $g = H^{-1/2}$. Observe that in general, $(g \cdot h)(v, w) = h_0((g^{\dagger_{h_0}}Hg)v, w)$; consequently, for our choice of gauge transformation $g = H^{-1/2}$, indeed $(g \cdot h) = h_0$. Then, for the complex gauge action in (1.4), $g \cdot (\overline{\partial}_E, \varphi, h) = (H^{1/2} \circ \overline{\partial}_E \circ H^{-1/2}, H^{1/2}\varphi H^{-1/2}, h_0)$. Consequently, the associated pair (d_A, Φ) is defined by $\overline{\partial}_A = H^{1/2} \circ \overline{\partial}_E \circ H^{-1/2}$ and $\Phi = H^{1/2}\varphi H^{-1/2}$.

Remark 1.1. Locally, it will be convenient to work in both holomorphic and unitary gauges. A local basis $\{s_1, \dots, s_n\}$ of sections of *E* is holomorphic if s_i are holomorphic sections of $(E, \overline{\partial}_E)$. A local basis $\{s_1, \dots, s_n\}$ of sections of *E* is unitary if $h_0(s_i, s_j) = \delta_{ij}$.

1.2. **Summary of results.** Fix a polystable Higgs bundle $(\overline{\partial}_E, \varphi)$ in a non-degenerate fiber of the Hitchin fibration $\mathcal{M} \to \mathcal{B}$. Consider the \mathbb{R}_t^+ -family of Higgs bundles $(\overline{\partial}_E, t\varphi)$. We seek to describe the corresponding family of harmonic metric h_t for $t \gg 0$, as shown in Figure 1.1. To describe h_t ,

- first, we construct a singular hermitian metric h_{ℓ} (*A posteriori*, we find in Corollary 6.4 that $h_{\ell} = \lim_{t \to \infty} h_{t}$.) (§3.1);
- then, we construct a family of approximate solutions, h_t^{app}, built from the singular hermitian metric h_l and a family of local model solutions (§4.1);
- we prove that this family of approximate hermitian metrics h_t^{app} solves Hitchin's equations up to an exponentially-decaying error (Proposition 4.10); and
- finally, we perturb from the approximate solutions h_t^{app} to the actual solutions h_t using a contraction mapping argument. This last point is the content of the main theorem, Theorem 6.1.



FIGURE 1.1. A \mathbb{R}^+ -family of Higgs bundles approaching the $(t = \infty)$ -ends of the Hitchin moduli space.

The strategy of the proof outlined above is the same as the strategy in the n = 2 case appearing in [MSWW16, MSWW14]. We highlight some notable differences. First, in the n = 2 case, the singular hermitian metric h_{ℓ} could be desingularized using a single 2 × 2 model solution. In the rank n case, we need $K \times K$ model solutions for $K = 2, \dots, n$ to desingularize h_{ℓ} . We discuss these model solutions in §4.1.2. Secondly, the proof that the inverse of the linearized operator is bounded (Proposition 5.2) requires substantial modification from [MSWW16]. (In Remark 5.3, we make a lengthy remark about why Mazzeo-Swoboda-Weiss-Witt's method does not work if n > 2.) *Remark* 1.2. Note that in both Mazzeo-Swoboda-Weiss-Witt's proof in [MSWW16, MSWW14] and the one here, we use the fact that Hitchin's equations are conformal. After fixing a polystable Higgs bundle $(\overline{\partial}_E, \varphi)$ in a non-degenerate fiber of $\mathcal{M} \to \mathcal{B}$, we take a conformal metric g'_C on C which is flat on disks around the zeros of the discriminant section Δ_{φ} defined in (2.4). The convenience of the metric g'_C will be discussed further in §4.

1.3. Acknowledgements. Many thanks to Rafe Mazzeo and Andy Neitzke for helpful discussions.

2. Higgs bundles in \mathcal{M}'

2.1. **Review of Hitchin fibration.** The SU(n)-Hitchin moduli space \mathcal{M} is a complex integrable system with half-dimensional base \mathcal{B} . The Hitchin fibration is

$$\begin{array}{rcl}
\text{Hit}: & \mathcal{M} & \to & \mathcal{B} \\ & & (\overline{\partial}_E, \varphi, h) & \mapsto & \operatorname{char}_{\varphi}(\lambda),
\end{array} \tag{2.1}$$

where $\operatorname{char}_{\varphi}(\lambda)$ is the characteristic polynomial of $\varphi \in \Omega^{1,0}(C, \operatorname{End} E)$. The Hitchin base \mathcal{B} can be identified with the complex vector space $\bigoplus_{i=2}^{n} H^{0}(C, K_{C}^{i}) \ni \mathbf{b} = (q_{2}, \cdots, q_{n})$, under the map from $\operatorname{char}_{\varphi}(\lambda)$ to its coefficients

$$\operatorname{char}_{\varphi}(\lambda) = \lambda^{n} + q_{2}\lambda^{n-2} + \dots + q_{n-1}\lambda + q_{n}.$$
(2.2)

A point $\mathbf{b} \in \mathcal{B}$ encodes the eigenvalues of φ . We can geometrically package the eigenvalues as a ramified n: 1-cover cut out of the total space of holomorphic cotangent bundle $K_C \rightarrow C$ by the equation

$$\Sigma = \{\lambda \in K_C : \operatorname{char}_{\varphi}(\lambda) = 0\}.$$
(2.3)

Call $\Sigma \xrightarrow{\pi} C$ the *spectral cover*.

The fiber $\operatorname{Hit}^{-1}(\mathbf{b})$ is a compact abelian variety if, and only if, the spectral curve $\Sigma_{\mathbf{b}}$ is smooth. Let \mathcal{B}' be this locus where the spectral cover $\Sigma_{\mathbf{b}}$ is smooth. We restrict our attention to Higgs bundles in the *regular locus* $\mathcal{M}' = \operatorname{Hit}^{-1}(\mathcal{B}')$, and call such Higgs bundles *regular*.

Given $(\overline{\partial}_E, \varphi)$, the discriminant section Δ_{φ} is

$$\Delta_{\varphi}: C \rightarrow K_{C}^{n^{2}-n}$$

$$p \mapsto \prod_{1 \leq i < j \leq n} (\lambda_{i}(p) - \lambda_{j}(p))^{2}.$$

$$(2.4)$$

As shown in Figure 2.1, the associated spectral curve $\Sigma \xrightarrow{\pi} C$ is ramified at the zeros of the discriminant section $Z = \Delta_{\varphi}^{-1}(0) \subset C$. Given $(\overline{\partial}_E, \varphi) \in \mathcal{M}'$, the map $\pi : \Sigma \to \mathbb{C}$ restricted



FIGURE 2.1. The spectral cover Σ is an n : 1 cover of C, ramified at Z.

to a neighborhood of a point $\tilde{p} \in \tilde{Z}$ looks like

$$\pi: \widetilde{\mathbb{D}} \to \mathbb{D}$$

$$w \mapsto w^{K} = z.$$
(2.5)

(This is indeed a smooth curve by the Jacobi criterion. The curve is the zero set of $f(w,z) := w^K - z$, and ∇f does not vanish at the point (0,0).) The point $\tilde{p} \in \tilde{Z}$ contributes a zero of order K - 1 to Δ_{φ} .

Remark 2.1. In the case where n = 2, a $SL(2, \mathbb{C})$ -Higgs bundle $(\overline{\partial}_E, \varphi)$ is in \mathcal{M}' if, and only if, $\Delta_{\varphi} = -4 \det \varphi$ has only simple zeros. Note that for n > 2, the space of regular Higgs bundles is slightly larger than the space of Higgs bundles for which the discriminant section Δ_{φ} has only simple zeros.

2.2. Local model near a ramification point for a Higgs bundle $(\overline{\partial}_E, \varphi) \in \mathcal{M}'$. The next proposition gives a local model around ramification points $p \in Z$ for regular Higgs bundles.



FIGURE 2.2. In the disk around $p \in Z$, we have n = 11 and $K_1 = K_2 = 3$, $K_3 = 2$, $K_4 = K_5 = K_6 = 1$.

Proposition 2.2. (Local model for $(\overline{\partial}_E, \varphi)$ around ramification points) Let $(\overline{\partial}_E, \varphi)$ be a polystable regular Higgs bundle. Let $p \in Z \subset C$ be a ramification point. Then, there are: a partition of n as $n = K_1 + \cdots + K_{m_p}$, local coordinates z_1, \cdots, z_{m_p} centered at p, and a local holomorphic trivialization of E over a disk \mathbb{D} centered at p such that

$$\overline{\partial}_E = \overline{\partial}$$
 (2.6)

$$arphi \ = \ igoplus_{j=1}^{m_p} \left(\lambda_{(j)} \mathbf{1}_{K_j} + egin{pmatrix} 0 & 1 & & \ & 0 & \ddots & \ & & \ddots & 1 \ & & \ddots & 1 \ z_j & & & 0 \end{pmatrix}_{K_j imes K_j}
ight).$$

Here, $\{\lambda_1, \dots, \lambda_n\}$ are the eigenvalues of φ , and $\lambda_{(i)}$ is the average of the cluster of K_i eigenvalues

$$\lambda_{(j)} = \sum_{k=s_{j-1}+1}^{s_j} \lambda_k \quad \text{where } s_i = \sum_{q=1}^i K_q.$$
 (2.7)

Remark 2.3. For the $SL(2, \mathbb{C})$ case, see [MSWW16, Lemma 4.2] which is considerably simpler and features an explicit gauge transformation. It is difficult to write such an explicit gauge transformation for arbitrary rank.

Proof. Take a disk \mathbb{D} centered at p without additional ramification points. Partition the eigenvalues by the value at p, and call these distinct values $\lambda_{(j)}(p)$. Let K_j be the associated cluster size. Because Σ is smooth, there is exactly one sheet of Σ going through $\lambda_{(j)}(p)$, and the spectral curve $\pi : \Sigma \to C$ through $\lambda_{(j)}(p)$ is locally given by

$$\pi_j : \widetilde{\mathbb{D}}_j \to \mathbb{D}$$

$$w_j \mapsto w_j^{K_j} = z_j$$
(2.8)

for some local holomorphic function z_i . We can arrange that z_i satisfies

$$\prod_{k=s_{j-1}+1}^{s_j} (x + \lambda_{(j)} - \lambda_k) = x^{K_j} - z_j dz_j^{K_j}.$$
(2.9)

(In the case K = 2, this is equivalent to the standard argument (e.g. [Mas86, p. 216]) showing that there is a local holomorphic coordinate z_i centered at p such that

$$(\lambda_1 - \lambda_2)^2 = 4z_j dz_j^2.)$$
(2.10)

We can work locally with each cluster of size K_j , and for convenience we may shift the eigenvalues so that $\lambda_{(j)} = 0$; to avoid notational clutter, we drop all the indices j related to cluster number, and number the eigenvalues $\lambda_1, \dots, \lambda_K$. The associated eigenvalues $\lambda_1, \dots, \lambda_K$ —or more precisely their pullbacks $\pi^* \lambda_i$ —are single-valued on the ramified K : 1 local cover $\widetilde{\mathbb{D}}$. Order them so that $\lambda_j = e^{\frac{2\pi i (j-1)}{K}} w d(w^K)$. Define

$$\sigma: \widetilde{\mathbb{D}} \to \widetilde{\mathbb{D}}$$

$$w \mapsto e^{2\pi i/K} w.$$
(2.11)

The cyclic group $\mathbb{Z}_K = \langle \sigma \rangle$ acts on $\widetilde{\mathbb{D}}$, exchanging the sheets of $\pi : \widetilde{\mathbb{D}} \to \mathbb{D}$. Note that $\lambda_i = (\sigma^{i-1})^* \lambda_1$.

Because the spectral cover Σ is smooth, the associated rank 1, locally-free, torsion-free sheaf $\mathcal{L} \to \Sigma$ is actually a line bundle. Thus, choose s_1 a smooth non-vanishing holomorphic section of the eigenline associated to λ_1 . Define $s_i = (\sigma^{i-1})^* s_1$ and note that in the basis s_i of $\pi^* \mathcal{E}$, $\pi^* \varphi$ acts by multiplication by λ_i . The basis elements $\{s_i\}$ do not descend from $\widetilde{\mathbb{D}}$ to \mathbb{D} , but the following basis elements satisfy $\sigma^* s'_i = s'_i$, and hence descend.

$$s_{1}' = \frac{1}{K} \sum_{i=1}^{K} s_{i}$$

$$s_{i}' = \frac{1}{Kw^{K}(d(w^{K}))^{K-(i-1)}} \sum_{k=1}^{K} \lambda_{k}^{K-(i-1)} s_{k}$$

$$i = 2, \cdots, K$$

$$(2.12)$$

Note that s'_i is nonsingular and non-vanishing at w = 0. In this basis,

$$\pi^* \varphi(s_1') = w^K \mathbf{d}(w^K) \varphi(s_K'), \qquad \pi^* \varphi(s_i') = \mathbf{d}(w^K) \varphi(s_{i-1}') \quad \text{for } i = 2, \cdots, K.$$
(2.13)

Define the basis e_i by $\pi^* e_i = s'_i$. In this holomorphic basis the $K \times K$ block of φ is

$$\begin{pmatrix} 0 & 1 & & \\ & 0 & \ddots & \\ & & \ddots & 1 \\ z & & 0 \end{pmatrix} dz.$$
 (2.14)

If the average of the eigenvalues $\lambda_{(j)} \neq 0$, then we simply add $\lambda_{(j)} \mathbf{1}_K$, as claimed in (2.6). Note that this holomorphic gauge is not unique since the section s_1 can be multiplied by any non-vanishing holomorphic function f.

In a block where K = 1, the associated eigenvalue λ is not ramified, so we simply choose e to be a smooth section of the associated eigenline over the base \mathbb{D} .

Remark 2.4. The sections s'_i appearing in (2.12) accomplish something slightly subtle. In the case K = 2, Proposition 2.2 produces the following basis and sections:

$$\varphi = \begin{pmatrix} 0 & 1 \\ z & 0 \end{pmatrix} dz, \quad s_1 = \pi^* \begin{pmatrix} 1 \\ \sqrt{z} \end{pmatrix}, \lambda_1 = \pi^* (\sqrt{z} dz), \quad s_2 = \pi^* \begin{pmatrix} 1 \\ -\sqrt{z} \end{pmatrix}, \lambda_2 = \pi^* (-\sqrt{z} dz)$$

In particular, note that s_1 and s_2 become linearly dependent at w = 0. Despite this, the sections

$$s'_{1} = \pi^{*} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \qquad s'_{2} = \pi^{*} \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$
 (2.15)

are linearly independent—even at w = 0.

Remark 2.5. By shrinking \mathbb{D} , we may assume that the disks around different points of *Z* do not intersect. Shrinking \mathbb{D} further, we may assume that the difference between the eigenvalues of φ is bounded below by some positive constant $\epsilon_{\lambda} > 0$ on $C - \bigcup_{p \in Z} \mathbb{D}_p$. By possibly taking a smaller ϵ_{λ} , we may assume that *on* \mathbb{D} , the difference between the averaged-eigenvalues $\lambda_{(j)}$ are bounded below by ϵ_{λ} . By rescaling the Riemannian metric on g_C , we may assume that each disk \mathbb{D}_p centered at p has radius one.

As shown in Figure 2.3, Proposition 2.5 gives a local model only when the ramification points all lie above the same point. Deforming this, we can also give a local model when the ramification points lie above points that are nearby. This is the content of Corollary 2.6, a direct corollary of the proof of Proposition 2.2.



FIGURE 2.3. Proposition 2.2 gives a local model for local spectral covers corresponding to the left figure. Corollary 2.6 gives a similar local model for the right figure.

Corollary 2.6. Let $(\overline{\partial}_E, \varphi)$ be a polystable regular Higgs bundle. Let \mathbb{D} be an open neighborhood over which the spectral cover has m_p connected components, each containing at most one point of \widetilde{Z} . Then, there are: a partition of n as $n = K_1 + \cdots + K_{m_p}$, local coordinates z_1, \cdots, z_{m_p} with z_i centered at $\pi(\widetilde{p}_i)$, and a local holomorphic trivialization of E over a disk \mathbb{D} centered at p such that

$$\overline{\partial}_{E} = \overline{\partial} \qquad (2.16)$$

$$\varphi = \bigoplus_{j=1}^{m_{p}} \left(\lambda_{(j)} \mathbf{1}_{K_{j}} + \begin{pmatrix} 0 & 1 & & \\ & 0 & \ddots & \\ & & \ddots & 1 \\ & & \ddots & 1 \\ z_{j} & & & 0 \end{pmatrix}_{K_{j} \times K_{j}} \right).$$

2.3. **Stratification of** \mathcal{M}' . Given a Higgs bundle $(\overline{\partial}_E, \varphi) \in \mathcal{M}'$, we get a collection of partitions $n = K_{1,p} + K_{2,p} + \cdots + K_{m_p,p}$ labeled by $p \in Z$. For $K = 2, 3, \ldots, n$ define

$$N_K = \#\{(p,i) \in Z \times \mathbb{N} : K_{p,i} = K\}.$$
(2.17)

(Since Δ_{φ} is a section of $K_C^{n^2-n}$,

$$\sum_{K=2}^{n} (K-1)N_K = 2(n^2 - n)(g - 1).)$$
(2.18)

This gives us a map onto a discrete space:

$$\Xi: \mathcal{M}' \to \mathbb{N}^{n-1}$$

$$(\overline{\partial}_E, \varphi) \mapsto (N_2, \cdots, N_n).$$

$$(2.19)$$

The map Ξ gives us a stratification of \mathcal{M}' .

3. LIMITING CONFIGURATIONS

One of the salient properties of the limiting metric $h_{\infty} = \lim_{t\to\infty} h_t$ is that it solves the "decoupled SU(n)-Hitchin's equations" by [Moc15, Theorem 2.7].

Definition 3.1. Given a polystable Higgs bundle $(\overline{\partial}_E, \varphi) \in \mathcal{M}$, a hermitian metric *h* solves the *decoupled SU*(*n*)-*Hitchin's equations* if

$$[\varphi, \varphi^{\dagger_h}] = 0, \qquad F_{D(\bar{\partial}_F, h)}^{\perp} = 0. \tag{3.1}$$

and det $h = h_{\text{Det }E}$.

Fix a polystable regular Higgs field $(\overline{\partial}_E, \varphi)$. In this section, we construct a metric h_ℓ solving the decoupled SU(n)-Hitchin's equations. It is worth emphasizing that there are many solutions of the decoupled SU(n)-Hitchin's equations, and each of these solutions depends on a choice of parabolic weights (Remark 3.4). We make the "correct" choice of parabolic weights in our construction, though this is only justified *a posteriori* in Corollary 6.4 when we prove that $h_\ell = h_\infty$. The subscript ℓ is used for "limiting."

3.1. **Construction of limiting metrics.** Given a polystable regular Higgs field $(\bar{\partial}_E, \varphi)$, Construction 3.2 produces a singular hermitian metric h_ℓ , unique up to rescaling by a constant. This metric arises as the pushforward of the Hermitian-Einstein metric on the associated spectral line bundle $\mathcal{L} \to \Sigma$ equipped with a specific parabolic structure. By Proposition 3.3, the triple $(\bar{\partial}, \varphi, h_\ell)$ solves the decoupled SU(n)-Hitchin's equations— possible after some constant rescaling of h_ℓ . As mentioned above, in Corollary 6.4, we will prove that $\lim_{t\to\infty} h_t = h_\ell$. Thus, we call this particular triple which solves the decoupled SU(n)-Hitchin's equations a *limiting configuration*.



FIGURE 3.1. At a point $\tilde{p} \in \tilde{Z}$ where the spectral cover is locally K : 1, put parabolic weight $\frac{1-K}{2}$.

Construction 3.2. *Given* $(\overline{\partial}_E, \varphi)$ *a regular polystable Higgs bundle, let* $\mathcal{L} \to \Sigma$ *be the associated spectral data.*

- Equip the holomorphic line bundle $\mathcal{L} \to \Sigma$ with parabolic structure: At each point $\tilde{p}_j \in \tilde{Z}$ add the parabolic weight $\frac{1-K_j}{2}$ (as shown in Figure 3.1).
- Equip the parabolic line bundle L → Σ with a hermitian structure: For parabolic line bundles—such as L— there is a Hermitian-Einstein metric adapted to the parabolic structure[Sim90, Biq96]². The Hermitian-Einstein metric solves

$$F_{\mathcal{L}} = -2\pi\sqrt{-1}\frac{\text{pdeg }\mathcal{L}}{\text{rank }\mathcal{L}}\frac{\pi^*\omega_C}{\text{vol}_{\pi^*g_C}(\Sigma)}\text{Id}_{\mathcal{L}}.$$
(3.2)

and is unique up to rescaling by a constant.

Define h_ℓ on E|_{C-Z} from the orthogonal push-forward of the Hermitian-Einstein metric h_L on L → Σ. I.e. decompose E into eigenspaces of φ; these eigenspaces are orthogonal with respect to h_ℓ; on each eigenspace h_ℓ agrees with the metric induced by h_L.

Proposition 3.3. *Given a polystable regular Higgs bundle, Construction 3.2 produces a unique hermitian metric* h_{ℓ} *solving the* SU(n)*-decoupled Hitchin's equations.*

Proof. Construction 3.2 determines a hermitian metric h_{ℓ} on E_{C-Z} up to rescaling by a constant. Any such metric h_{ℓ} solves the decoupled Hitchin's equations. Since φ and h_{ℓ} are diagonal in the basis of eigenbundles on φ , $\left[\varphi, \varphi^{\dagger}_{h_{\ell}}\right] = 0$.

CLAIM: The parabolic degree of \mathcal{L} is equal to the degree of E.

Proof: \triangleright The statement $\text{pdeg}\mathcal{L} = \text{deg}E$ holds because of the choice of parabolic weights. A cluster of size K contributes a zero of order K - 1 to Δ_{φ} ; at such a point $\tilde{p} \in \tilde{Z}$, we assigned the parabolic weight $\frac{1-K}{2}$. Since Δ_{φ} has $2(n^2 - n)(g - 1)$ zeros (counted with multiplicity), the sum of all parabolic weights is $-\frac{1}{2} \cdot 2(n^2 - n)(g - 1)$. Consequently,

pdeg
$$\mathcal{L} = \deg \mathcal{L} + \sum_{\tilde{p} \in \tilde{Z}} \alpha_p$$
 (3.3)

$$= \left(\deg E + (n^2 - n)(g - 1) \right) + \left(-\frac{1}{2} \right) 2(n^2 - n)(g - 1)$$

$$= \deg E.$$

The condition $F_{D(\overline{\partial}_E,h_\ell)}^{\perp} = 0$ holds because pdeg $\mathcal{L} = \deg E$. The induced metric $\det(h_\ell)$ is a Hermitian-Einstein metric on Det *E*, consequently it is a constant multiple of the fixed Hermitian-Einstein metric $h_{\text{Det }E}$. Rescale h_ℓ by a constant so that these two Hermitian-Einstein metrics agree.

²Technically, the result in [Biq96] is only for parabolic line bundles of parabolic degree 0, however, it is straightforward to extend the results to arbitrary degree.

Remark 3.4. Note that in the proof we did not use the individual values of the parabolic weight. We only used the fact that the sum of all parabolic weights was $-(n^2 - n)(g - 1)$. In Construction 3.2, we could take any collection of parabolic weights summing to $-(n^2 - n)(g - 1)$ and produce a hermitian metric solving the decoupled SU(n)-Hitchin's equations. However, this hermitian metric agrees with h_{∞} only for our choice of parabolic weights (Corollary 6.4).

3.2. Local model near a ramification point for a limiting configuration in \mathcal{M}' . The next proposition gives a local model for the limiting configuration in Proposition 3.3.

Proposition 3.5. There is a holomorphic gauge satisfying the conditions of Proposition 2.2 in which

$$h_{\ell} = \bigoplus_{j=1}^{m_{p}} \begin{pmatrix} |z_{j}|^{-2\alpha_{K_{j},1}} & & \\ & \ddots & \\ & & |z_{j}|^{-2\alpha_{K_{j},K_{j}}} \end{pmatrix}_{K_{j} \times K_{j}}.$$
 (3.4)

Here, the constants $\alpha_{K,i}$ *are*

$$\alpha_{K,i} = \frac{2i - (K+1)}{2K}.$$
(3.5)

Proof. First, assume deg E = 0. We can work locally with each cluster of size K_j . As in the proof of Proposition 2.2, we assume that we are working with the first cluster and drop all subscripts relating to the cluster index. The key idea is that we use up the remaining gauge freedom in Proposition 2.2 by multiplying the section s_1 by a non-vanishing holomorphic function in order to arrange that

$$h_{\mathcal{L}}(s_1, s_1) = \sqrt{K} |w|^{1-K}.$$
(3.6)

Let $h_{\mathcal{L}}$ be the Hermitian-Einstein metric on $\mathcal{L} \to \Sigma$ which is adapted to the hermitian metric. There are two consequences of this. First, because $h_{\mathcal{L}}$ is adapted to the parabolic structure on $\mathcal{L} \to \Sigma$ at \tilde{p}_i , $h_{\mathcal{L}}(s_1, s_1) \sim |w|^{-2 \cdot \frac{1-K}{2}} = |w|^{K-1}$. Secondly, because $h_{\mathcal{L}}$ is Hermitian-Einstein and $\log h_{\mathcal{L}}(s_1, s_1)$ is harmonic. Any harmonic function on the punctured disk $\widetilde{\mathbb{D}}^{\times}$ can be written $\operatorname{Re}(f(w)) + c \log(|w|)$ where f(w) is holomorphic on $\widetilde{\mathbb{D}}^{\times}$ and c is some constant; hence

$$\log h_{\mathcal{L}}(s_1, s_1) = \operatorname{Re}(f(w)) + (K - 1)\log(|w|).$$
(3.7)

The function f is bounded because $h_{\mathcal{L}}$ is adapted, hence it extends to a holomorphic function on \mathbb{D} . We replace s_1 with the section $K^{1/2}e^{-\frac{f}{2}}s_1$ that satisfies

$$\log h_{\mathcal{L}}\left(K^{1/2} \mathrm{e}^{-\frac{f}{2}} s_1, K^{1/2} \mathrm{e}^{-\frac{f}{2}} s_1\right) = \log\left(K \mathrm{e}^{-\mathrm{Re}(f)} h_{\mathcal{L}}(s_1, s_1)\right) = (K-1)\log(|w|) + \log K.$$
(3.8)



FIGURE 4.1. The curvature $F_{D(\bar{\partial},h_{\ell})}^{\perp}$ is concentrated at $p \in Z$, illustrated by orange spikes. Approximate solutions h_t^{app} are constructed by desingularizing h_{ℓ} by gluing in smooth model solutions (shown in blue).

Consequently, $h_{\mathcal{L}}(s_i, s_i) = K|w|^{K-1}$ for $i = 1, \dots K$. Then, we see that

$$\pi^* h_{\ell}(s_i, s_k) = \begin{cases} 0 & \text{if } i \neq k \\ |w|^{K+1-2i} & \text{if } i = k \end{cases}.$$
(3.9)

Hence in the basis $\{e_i\}$, the hermitian metric is as claimed.

Note that when K = 1, the associated eigenvalue λ is not ramified. Consequently, the associated section e satisfies $\log h_{\ell}(e, e) = \operatorname{Re}(f(z))$ for f(z) harmonic on the disk \mathbb{D} —rather than its cover. Thus, by replacing e with the section $e^{-\frac{f}{2}}e$, we see that $h_{\ell}(e^{-\frac{f}{2}}e, e^{-\frac{f}{2}}e) = 1$, as desired.

Note that det $h_{\ell} = 1$ because h_{ℓ} is block diagonal and the determinant of each $K \times K$ block is 1.

If deg $E \neq 0$, then we simply note that log $h_{\mathcal{L}}(s_1, s_1)$ minus some multiple of the Kähler potential is harmonic, and repeat the argument above.

4. A FAMILY OF APPROXIMATE SOLUTIONS

Fix a regular polystable Higgs bundle $(\overline{\partial}_E, \varphi)$. Consider the \mathbb{R}_t^+ -family of Higgs bundles $(\overline{\partial}_E, t\varphi)$. Ultimately, we seek to describe the corresponding family of harmonic metric h_t for large values of t. In this section, we construct a \mathbb{R}^+ -family of approximate solutions h_t^{app} by desingularizing the limiting configuration h_ℓ in §3. As shown in Figure 4.1, the metric h_ℓ is singular at $p \in Z$, so we glue in smooth solutions of Hitchin's equations on the disks \mathbb{D} around each ramification point $p \in Z$. These smooth models are described in §4.1.3. Because these smooth models are defined on disks in \mathbb{C} with its usual flat metric, we take a conformal metric g'_C on C which is flat in each disk \mathbb{D} (Remark 1.2). The approximate solutions h_t^{app} are defined in §4.2.

4.1. **Model solutions.** For each cluster rank *K*, we describe the necessary family of model solutions of rank parameterized by $t \in \mathbb{R}^+$. All of these model solutions are on \mathbb{C} with its flat metric. We begin by reviewing the K = 2 family of model solution featured in [MSWW16] in §4.1.1 before turning to the higher rank versions in §4.1.2. We conclude

by describing the model solutions for the regular Higgs bundle $(\overline{\partial}_E, t\varphi)$ on the disk \mathbb{D} in §4.1.3.

4.1.1. *The* K = 2 *family of model solution*. The following family of model solutions is featured in [MSWW16].

Definition 4.1. The SU(2) *t-model solution* is

$$\overline{\partial}_{E}^{(2)} = \overline{\partial}$$

$$\varphi^{(2)} = \begin{pmatrix} 0 & 1 \\ z & 0 \end{pmatrix} dz$$

$$h_{t}^{(2), \text{mod}} = \begin{pmatrix} |z|^{1/2} e^{u_{t}(|z|)} & \\ & |z|^{-1/2} e^{-u_{t}(|z|)} \end{pmatrix}$$
(4.1)

where $u_t : \mathbb{R}^+ \to \mathbb{R}$ is solution of

$$\left(\frac{d^2}{d|z|^2} + \frac{1}{|z|}\frac{d}{d|z|}\right)u_t = 8t^2|z|\sinh(2u_t).$$
(4.2)

with asymptotics

$$u_t(|z|) \sim \frac{1}{\pi} K_0(\frac{8t}{3}|z|^{\frac{3}{2}}) \text{ as } |z| \to \infty$$

 $u_t(|z|) \sim -\frac{1}{2} \log(|z|) \text{ as } |z| \to 0.$

Remark 4.2. The u_t are related by

$$u_t = \rho_t^* u_1 \qquad \rho_t(z) = t^{2/3} z.$$
 (4.3)

Remark 4.3. Note that $h_t^{(2),\text{mod}}$ has a chance of being smooth at |z| = 0 because of the coefficient of $\log(|z|)$ appearing in the expansion around |z| = 0. Mazzeo-Swoboda-Weiss-Witt prove that it is smooth in [MSWW16, Corollary 3.4]. Moreover, note that the pointwise limit $\lim_{t\to\infty} h_t^{(2),\text{mod}}$ is diag $(|z|^{1/2}, |z|^{-1/2})$. This is the 2 × 2 block appearing in the limiting metric h_ℓ in (3.4).

Remark 4.4. In unitary gauge (see the discussion at the end of $\S1.1$), the SU(2) t-model solution is written

$$A_{t}^{(2),\text{mod}} = d + \left(\frac{1}{8} + \frac{|z|}{4}\frac{du_{t}}{d|z|}\right) \begin{pmatrix} 1 \\ -1 \end{pmatrix} \left(\frac{dz}{z} - \frac{d\overline{z}}{\overline{z}}\right)$$

$$\Phi_{t}^{(2),\text{mod}} = \begin{pmatrix} 0 & |z|^{1/2}e^{u_{t}(|z|)} \\ \frac{z}{|z|^{1/2}}e^{-u_{t}(|z|)} & 0 \end{pmatrix} dz$$
(4.4)

In [MSWW16], as well as in [GMN09], the model solution takes this shape.

4.1.2. The rank K family of model solutions from $\mathcal{M}_{K,1}$. For any rank K, there is a single SU(K) model solution on \mathbb{C} with its flat metric which generalizes the the SU(2) (t = 1)-model solution in §4.1.1. For each K, there is a one-point moduli space $\mathcal{M}_{K,1}$ of solutions of the SU(K)-Hitchin's equations on \mathbb{CP}^1 with an irregular singularity at $\{\infty\}$ such that the eigenvalues of the Higgs field are $\lambda_r = e^{2\pi i r/K} z^{1/K} dz$. The point $[(\overline{\partial}_E, \varphi, h)]$ is fixed by a U(1)-action, consequently the solution of Hitchin's equations can be written down relatively explicitly [FN17]:

Proposition 4.5. [FN17, Proposition 3.9 & Lemma 3.13] *The one-point point moduli space* $\mathcal{M}_{K,1} = [(\bar{\partial}_E, \varphi, h^{(K), \text{mod}})]$ where

$$\overline{\partial}_{E} = \overline{\partial}, \qquad \varphi = \begin{pmatrix} 0 & 1 & & \\ & 0 & \ddots & \\ & & \ddots & 1 \\ z & & 0 \end{pmatrix} dz, \qquad h^{(K), \text{mod}} = \begin{pmatrix} |z|^{-2\alpha_{K,1}} e^{u_{K,1}} & & \\ & & \ddots & \\ & & |z|^{-2\alpha_{K,K}} e^{u_{K,K}} \end{pmatrix}.$$
(4.5)

The constants α_i *are*

$$\alpha_{K,i} = \frac{2i - (K+1)}{2K}.$$
(4.6)

The real-valued functions $u_{K,i}(z) = u_{K,i}(|z|)$ *satisfy the symmetry* $u_{K,i} = -u_{K,K+1-i}$ *and solve*

$$\frac{1}{4} \left(\frac{\mathrm{d}^2}{\mathrm{d}|z|^2} + \frac{1}{|z|} \frac{\mathrm{d}}{\mathrm{d}|z|} \right) u_{K,i} = |z|^{\frac{2}{K}} \left(\mathrm{e}^{u_{K,i} - u_{K,i+1}} - \mathrm{e}^{u_{K,i-1} - u_{K,i}} \right)$$
(4.7)

with the following boundary conditions:

- The function $u_{K,i}$ decays to 0 as $|z| \to \infty$.
- *Near* 0, $u_{K,i} \sim 2\alpha_{K,i} \log |z|$.

Letting $\mathbf{u}(|z|) = (u_{K,1}(|z|), \dots, u_{K,K}(|z|))$, the function $\|\mathbf{u}(|z|)\|^2$ is decreasing and exhibits exponential decay at ∞ . More precisely, for $\epsilon > 0$, take $R_{\epsilon} > 0$ such that $\|\mathbf{u}(R_{\epsilon})\| < \epsilon$. Then, there is a constant c > 0 (depending explicitly on ϵ and K) such that

$$\|\mathbf{u}\|^{2}(\rho) \leq \epsilon^{2} \frac{K_{0}(c\zeta(\rho))}{K_{0}(c\zeta(R_{\epsilon}))} \qquad \text{for } \rho > R_{\epsilon},$$
(4.8)

where K_0 is the modified Bessel function of first kind and $\zeta(|z|) = \frac{2K}{K+1}|z|^{\frac{(K+1)}{K}}$.

Remark 4.6. The bound in (4.8) is not sharp. The constant $c = (2C_{\epsilon}C_{K})^{-1/2}$ where $C_{\epsilon} > 1$ with $\lim_{\epsilon \searrow 0} C_{\epsilon} = 1$; the first few values C_{K} are $C_{2} = 4$, $C_{3} = 3$, $C_{4} = 2$, $C_{5} = \frac{5-\sqrt{5}}{2}$. In the case where K = 2, $u_{1} = -u_{2} \sim K_{0}(\frac{8}{3}\rho^{3/2})$; consequently the constant $\frac{4}{3}c$ at best approaches $\frac{\sqrt{8}}{3}$ —considerably worse than the optimal constant $\frac{16}{3}$.

Remark 4.7. With the change of variables above, $u_{K,i}(\zeta)$ solve the system of equations

$$\left(\frac{d^2}{d\zeta^2} + \frac{1}{\zeta}\frac{d}{d\zeta}\right)u_{K,i} = e^{u_{K,i}-u_{K,i+1}} - e^{u_{K,i-1}-u_{K,i}}.$$
(4.9)

This is the radial version of the coupled system of PDE known as "2d cyclic affine Toda lattice with opposite sign." Because of the symmetry, this is actually a coupled system of $\lfloor \frac{K-1}{2} \rfloor$ ODEs.

The solution of Hitchin's equations in Proposition (4.5) appears earlier in the literature, where it is called a solution of the " tt^* -Toda equations." The tt^* -Toda equations are a special case of the tt^* -equations which were introduced by Cecotti and Vafa to describe certain deformations of supersymmetric quantum field theories [CV91, CV92]. (Not every solution of the tt^* -equations is a solution of Hitchin's equations on a Riemann surface, and conversely, not every solution of Hitchin's equations gives a solution of the tt^* -equations. However, these coincide here roughly because $\mathcal{M}_{K,1}$ is a one-point moduli space fixed by a circle action and a real involution.) These particular solutions were also studied in [GL10, GL13, Moc13].

We now introduce the parameter $t \in \mathbb{R}^+$. Define rescaled functions

$$u_{K,i,t} = \rho_{K,t}^* u_{K,i}, \qquad \text{where } \rho_{K,t} : r \to t^{\frac{\kappa}{K+1}} r. \tag{4.10}$$

The following triple $(\overline{\partial}_E, t\varphi, h_t^{(K), \text{mod}})$ solves Hitchin's equations:

$$\overline{\partial}_{E} = \overline{\partial}, \qquad t\varphi = t \begin{pmatrix} 0 & 1 & & \\ & 0 & \ddots & \\ & & \ddots & 1 \\ z & & 0 \end{pmatrix} dz, \qquad h_{t}^{(K), \text{mod}} = \begin{pmatrix} |z|^{-2\alpha_{1}} e^{u_{K,1,t}} & & \\ & \ddots & \\ & & |z|^{-2\alpha_{K}} e^{u_{K,K,t}} \end{pmatrix}.$$

$$(4.11)$$

4.1.3. *Family of model solutions for* $(\overline{\partial}_E, t\varphi)$. We will use the following family of model solutions to desingularize h_ℓ .

Definition 4.8. Let $(\overline{\partial}_E, \varphi, h_\ell)$ be as in (2.6) & (3.4). Define a hermitian metric

$$h_{t}^{\text{mod}} = \bigoplus_{j=1}^{m_{p}} \begin{pmatrix} |z_{j}|^{-2\alpha_{K_{j},1}} e^{u_{K_{j},1,t}(|z_{j}|)} & & \\ & \ddots & \\ & & |z_{j}|^{-2\alpha_{K_{j},K_{j}}} e^{u_{K_{j},K_{j},t}(|z_{j}|)} \end{pmatrix}_{K_{j} \times K_{j}}.$$
 (4.12)

Call the *t*-family ($\overline{\partial}_E$, $t\varphi$, h_t^{mod}) the *family of model solutions* of Hitchin's equations.

Remark 4.9. In unitary gauge, this is

$$A_{t}^{\text{mod}} = \bigoplus_{j=1}^{m_{p}} \begin{pmatrix} -\frac{\alpha_{K_{j},1}}{2} + \frac{|z_{j}|}{4} \frac{du_{K_{j},1,t}}{d|z_{j}|} & & \\ & \ddots & \\ & -\frac{\alpha_{K_{j},K_{j}}}{2} + \frac{|z_{j}|}{4} \frac{du_{K_{j},K_{j},t}}{d|z_{j}|} \end{pmatrix} \begin{pmatrix} \left(\frac{dz_{j}}{z_{j}} - \frac{d\overline{z}_{j}}{d\overline{z}_{j}}\right) & (4.13) \\ & K_{j} \times K_{j} \end{pmatrix} \\ \Phi_{t}^{\text{mod}} = \bigoplus_{j=1}^{m_{p}} \begin{pmatrix} 0 & |z_{j}|^{\frac{1}{K_{j}}} e^{\frac{u_{K_{j},K_{j},t}-u_{K_{j},2,t}}{2}} & & \\ & 0 & \ddots & \\ & & \ddots & |z_{j}|^{\frac{1}{K_{j}}} e^{\frac{u_{K_{j},K_{j}-1,t}-u_{K_{j},K_{j},t}}{2}} \end{pmatrix} dz_{j} \end{pmatrix}$$

Note that if $K_j = 1$, the 1×1 block in A_t^{mod} is (0) and the block in Φ_t^{mod} is the eigenvalue $(\lambda_{(j)})$.

4.2. **Description of approximate solutions.** The following non-linear operator measures the the failure of $(\overline{\partial}_E, \varphi, h)$ to be a solution of Hitchin's equations:

$$\mathbf{F}(\overline{\partial}_E, \varphi, h) := H^{1/2} \left(F_{D(\overline{\partial}_E, h)}^{\perp} + [\varphi, \varphi^{\dagger_h}] \right) H^{-1/2}.$$
(4.14)

Observe that we conjugate by the End(*E*)-valued section $H^{1/2}$ (discussed at the end of §1.1) which satisfies $h(v,w) = h_0(H^{1/2}v, H^{1/2}w)$. By doing this, the output $\mathbf{F}(\bar{\partial}_E, \varphi, h)$ is an h_0 -unitary section of $\Omega^{1,1}(C, \mathfrak{su}(E))$. (Equivalently in the unitary formulation of Hitchin's equations, this operator **F** is equal to $\mathbf{F}(\mathbf{d}_A, \Phi) = F_A^{\perp} + [\Phi, \Phi^{\dagger_{h_0}}]$.)

Definition/Proposition 4.10. *Choose a smooth cutoff function* $\chi : [0, \infty) \rightarrow [0, 1]$ *such that*

$$\chi \Big|_{[0,\frac{1}{2}]} = 1 \quad and \quad \chi \Big|_{[1,\infty)} = 0.$$
 (4.15)

On \mathbb{D}_p , in the local gauge of Proposition 3.5, define h_t^{app} by

$$h_{t}^{\text{app}} = \bigoplus_{j=1}^{m_{p}} \begin{pmatrix} |z_{j}|^{-2\alpha_{K_{j},1}} e^{\chi(|z_{j}|)u_{K_{j},1,t}(|z_{j}|)} & & \\ & \ddots & \\ & & |z_{j}|^{-2\alpha_{K_{j},K_{j}}} e^{\chi(|z_{j}|)u_{K_{j},K_{j},t}(|z_{j}|)} \end{pmatrix}_{K_{j} \times K_{j}}$$
(4.16)

On $C^{\text{ext}} = C - \overline{\bigcup_p \mathbb{D}_p}$, define $h_t^{\text{app}} = h_\ell$.

For $t_0 > 0$ sufficiently large, there exists positive constants c, δ such that for $t > t_0$

$$\left\|\mathbf{F}(\bar{\partial}_E, t\varphi, h_t^{\mathrm{app}})\right\|_{L^2(C)} \le ce^{-\delta t},\tag{4.17}$$

for **F** defined in (4.14). Because of the exponential decay in *t*, call the family $\{(\overline{\partial}_E, t\varphi, h_t^{app})\}_{t>t_0}$ a family of approximate solutions.

Remark 4.11. In unitary gauge, the family $(A_t^{app}, \Phi_t^{app})$ is given by inserting the cutoff function χ into the expressions $(A_t^{mod}, \Phi_t^{mod})$ in (4.13)

Proof of Proposition 4.10. On C^{ext} , $h_t^{\text{app}} = h_\ell$. Because $(\overline{\partial}_E, \varphi, h_\ell)$ solves the decoupled Hitchin's equations, $\mathbf{F}(\overline{\partial}_E, t\varphi, h_t^{\text{app}})$ vanishes on C^{ext} . The $L^2(C)$ -norm is simply the sum of the $L^2(\mathbb{D}_p)$ -norms of each of the individual $K_j \times K_j$ blocks. Dropping indices, the relevant $K \times K$ piece is

$$\overline{\partial}_{E} = \overline{\partial} \quad t\varphi = t \begin{pmatrix} \lambda & 1 & & \\ & \lambda & \ddots & \\ & & \ddots & 1 \\ z & & \lambda \end{pmatrix} dz, \quad h_{t}^{app} = \begin{pmatrix} |z|^{-2\alpha_{K,1}} e^{\chi u_{K,1,t}} & & \\ & & \ddots & \\ & & |z|^{-2\alpha_{K,K}} e^{\chi u_{K,K,t}} \end{pmatrix}.$$
(4.18)

On the *K* × *K* block, the value of $F_{D(\overline{\partial}_{E}, \varphi)} + t^{2}[\varphi, \varphi^{\dagger_{h_{t}^{app}}}] = 0$ is the diagonal matrix whose (i, i) entry is

$$\left(-\frac{1}{4}\left(\frac{d^{2}}{d|z|^{2}}+\frac{1}{|z|}\frac{d}{d|z|}\right)\chi u_{K,i,t}+t^{2}|z|^{\frac{2}{K}}\left(e^{\chi u_{K,i,t}-\chi u_{K,i+1,t}}-e^{\chi u_{K,i-1,t}-\chi u_{K,i,t}}\right)\right)dz\wedge d\overline{z}.$$
(4.19)

(Note that without the cutoff function χ , this vanishes.) From the exponential decay of $u_{K,i}(|z|)|$ in |z| like $e^{-c|z|^{\frac{K+1}{K}}}$ (see Proposition 4.5), we see that—fixing $|z| - u_{K,i,t}(|z|)$ decays in t like e^{-ct} . To see that the expression in (4.19) is exponentially decaying, we break it into pieces. For $t \gg 0$, there is a constant C_1 close to 1 such that

$$\begin{aligned} \left| e^{\chi u_{K,i,t} - \chi u_{K,i+1,t}} - 1 \right| &\leq C_1 \left(\chi u_{K,i,t} - \chi u_{K,i+1,t} \right) \leq C_1 \left(\left| u_{K,i,t} \right| + \left| u_{K,i+1,t} \right| \right) \quad (4.20) \\ \left| e^{\chi u_{K,i-1,t} - \chi u_{K,i,t}} - 1 \right| &\leq C_1 \left(\chi u_{K,i-1,t} - \chi u_{K,i,t} \right) \leq C_1 \left(\left| u_{K,i-1,t} \right| + \left| u_{K,i,t} \right| \right). \end{aligned}$$

Additionally, because $u_{K,i,t}$ and its derivatives in |z| are all exponentially decaying in t, there is a constant C_2 depending on the maximum of $|\chi'|$ and $|\chi''|$ such that

$$\left| \left(\frac{\mathrm{d}^2}{\mathrm{d}|z|^2} + \frac{1}{|z|} \frac{\mathrm{d}}{\mathrm{d}|z|} \right) \chi u_{K,i,t} \right| \leq C_2 \mathrm{e}^{-ct}.$$

$$(4.21)$$

The exponential decay of (4.19) follows.

5. PROPERTIES OF THE LINEARIZATION

In Proposition 4.10, we proved that the family h_t^{app} of approximate solutions was close to solving Hitchin's equations. The metrics h_t^{app} failed to solve Hitchin's equations only on the union of the gluing annuli around $p \in Z$ where the value of the cutoff function $\chi(|z_i|)$ differed from 0 or 1—and on those gluing annuli, the error was exponentially decaying in *t*.

Looking forward, the main theorem (Theorem 6.1) states something much stronger: for $t \gg 0$, the approximate metric h_t^{app} is close to the actual harmonic h_t solving Hitchin's equations in the sense that

$$h_t(v,w) = h_t^{\operatorname{app}}(\mathrm{e}^{-\gamma_t} v, \mathrm{e}^{-\gamma_t} w), \qquad (5.1)$$

for γ_t "small."

Following the conventions of Mazzeo-Swoboda-Weiss-Witt, we do the analysis using the unitary formulation of Hitchin's equations discussed in §1.1. We fix a hermitian metric h_0 on E. We replace the triple $(\overline{\partial}_E, t\varphi, h_t^{app})$ with a pair $(d_{A_t}, t\Phi_t)$ such that $(\overline{\partial}_E, t\varphi, h_t^{app})$ and $(\overline{\partial}_{A_t}, t\Phi_t, h_0)$ are complex gauge equivalent with respect to the action in (1.4), i.e. $[(\overline{\partial}_E, t\varphi, h_t^{app})]$ and $[(\overline{\partial}_{A_t}, t\Phi_t, h_0)]$ define the same point in the Hitchin moduli space \mathcal{M} .

There are two other interesting actions of the complex gauge group on the space of triples $(\bar{\partial}_E, \varphi, h)$. For these, the equation $\bar{\partial}_E \varphi = 0$ is preserved by the action of the complex gauge group; however, the equation $F_{D(\bar{\partial}_E,h)}^{\perp} + [\varphi, \varphi^{\dagger_h}] = 0$ is not preserved. In the first action, the complex gauge group acts transitively on the space of hermitian metrics by

$$g \cdot_1 (\overline{\partial}_E, \varphi, h) = (\overline{\partial}_E, \varphi, g \cdot h)$$
 where $(g \cdot h)(v, w) = h(gv, gw)$. (5.2)

If $(\overline{\partial}_E, \varphi)$ is polystable, then in the complex gauge orbit, there is a hermitian metric $g \cdot h$ solving Hitchin's equations. In the second action, we fix the hermitian metric and take the action

$$g \cdot_2 (\overline{\partial}_E, \varphi, h) = (g \circ \overline{\partial}_E \circ g^{-1}, g \varphi g^{-1}, h).$$
(5.3)

This second action induces a complex gauge action on the space of pairs (d_A, Φ) in the unitary formulation of Hitchin's equations:

$$g \cdot (\mathbf{d}_A, \Phi) = (D(g \circ \overline{\partial}_E \circ g^{-1}, h_0), g \Phi g^{-1}), \tag{5.4}$$

where *D* is the Chern connection associated to the pair. Note that these two actions of the complex gauge transformation satisfy

$$g \cdot \left(g \cdot_2(\overline{\partial}_E, \varphi, h)\right) = g \cdot_1(\overline{\partial}_E, \varphi, h), \tag{5.5}$$

where g is the action of the complex gauge transformation in (1.4).

We are interested in finding the complex gauge transformation g such that $g \cdot (d_{A_t}, t\Phi_t)$ (defined in (5.4)) solves Hitchin's equations. Since Hitchin's equations are invariant under h_0 -unitary gauge transformations, we take the standard slice of the complex gauge transformations modulo h_0 -unitary gauge transformations by assuming that $g = e^{-\gamma}$ is h_0 -hermitian. Define the operator

$$\mathbf{F}_{t}^{\mathrm{app}}(\gamma) := -i \star \left(F_{A_{t}^{\mathrm{exp}(-\gamma)}} + t^{2} [\mathrm{e}^{-\gamma} \Phi_{t} \mathrm{e}^{\gamma}, \mathrm{e}^{\gamma} \Phi_{t}^{\dagger_{h_{0}}} \mathrm{e}^{-\gamma}] \right).$$
(5.6)

(We add the superscript to clarify that this expression is based at the approximate solution $(\bar{\partial}_E, t\varphi, h_t^{app})$.) We are interested in the family γ_t satisfying $\mathbf{F}_t^{app}(\gamma_t) = 0$. Note that this is equivalent to finding an h_0 -unitary γ_t satisfying (5.1).

In this section, we study the linearization of $\mathbf{F}_t^{\text{app}}$ and prove bounds on its inverse (Proposition 5.2). The linearization of $\mathbf{F}_t^{\text{app}}$ at 0 is

$$L_t \gamma := D \mathbf{F}_t^{\mathrm{app}}(0)[\gamma] = \frac{d}{d\epsilon} \Big|_{\epsilon=0} \mathbf{F}_t^{\mathrm{app}}(\epsilon \gamma)$$

$$= \Delta_{A_t} \gamma - i \star t^2 M_{\Phi_t} \gamma$$
(5.7)

where

$$\Delta_{A_t} := d_{A_t}^* d_{A_t} \gamma$$

$$M_{\Phi_t} \gamma := [\Phi_t^* \wedge [\Phi_t, \gamma]] - [\Phi_t \wedge [\Phi_t^*, \gamma]].$$
(5.8)

First note that L_t is a positive operator.

Proposition 5.1. [MSWW16, Proposition 5.1] If $\gamma \in \Omega^0(\mathfrak{sl}(E))$, then

$$\langle L_t \gamma, \gamma \rangle_{L^2} = \| \mathbf{d}_A \gamma \|_{L^2}^2 + 2t^2 \| [\Phi, \gamma] \|_{L^2}^2 + 2t^2 \| [\Phi^*, \gamma] \|_{L^2}^2 \ge 0.$$
(5.9)

Consequently, restricted to $\Omega^0(\mathfrak{isu}(E))$, L_t has no kernel.

We now prove that its inverse $L_t^{-1} : L^2(i\mathfrak{su}(E)) \to H^2(i\mathfrak{su}(E))$ is bounded.

Proposition 5.2. For t_0 sufficiently large, there is are constants $\tilde{C}_1, \tilde{C}_2 > 0$ such that

(a) $\|L_t^{-1}\|_{\mathcal{L}(L^2, L^2)} \le \tilde{C}_1,$

(b)
$$||L_t^{-1}||_{\mathcal{L}(L^2, H^2)} \leq \tilde{C}_2 t^2.$$

Remark 5.3. For the SU(2) case, the analog of Proposition 5.2a is stated in [MSWW16, Lemma 6.3]. An important ingredient of their strategy is the the domain decomposition principle in [BÖ0]. They decompose *C* into disjoint pieces: neighborhoods \mathbb{D}_p around each point $p \in Z$, plus the remaining piece $C^{\text{ext}} = C - \bigcup_p \mathbb{D}_p$. On each piece, they find a lower bound for the first Neumann eigenvalue. Then, the domain decomposition principle gives a lower bound on the first global eigenvalue.

One might hope that this method of proof works for SU(n) when n > 2. However, this does not work because the Neumann boundary problem on each disk \mathbb{D} has kernel. By explicit computation of L_t in the basis of (4.16) on \mathbb{D} (see 5.55), one can compute that the Neumann kernel consists of constant traceless diagonal matrices with the shape

$$\gamma = \bigoplus_{j=1}^{m_p} \gamma_{(j)} \mathbf{1}_{K_j}$$
(5.10)

Consequently, we pursue a global strategy that does not use the domain decomposition principle.

Proof of Proposition 5.2a. (Necessary lemmata appear in §5.1.) Define

$$\mathbb{L}_t \gamma := \Delta_{A_t} \gamma - i \star M_{\Phi_t} \gamma. \tag{5.11}$$

Since M_{Φ_t} is a semi-positive operator, $L_t \geq \mathbb{L}_t$. Consider the eigenvalues $\{\lambda_\ell^t\}$ of \mathbb{L}_t . These are all positive by Proposition 5.1. We will prove that the lowest eigenvalue λ_0^t of \mathbb{L}_t is bounded below by some constant $\kappa > 0$ as $t \to \infty$. Suppose to the contrary that $\lambda_0^t \to 0$.

We define a family of weight functions $\mu_t : C \to \mathbb{R}^+$ as follows: Around each point $p \in Z$, work in the gauge from Proposition 2.2 & 3.5 and let *z* be some holomorphic coordinate centered at *p*. (The coordinate *z* need not be any of the holomorphic coordinates z_i appearing in Proposition 2.2.) Order the elements of the partition of *n* so that $K_1 \ge K_2 \ge \cdots \ge K_{m_p}$. In the unit disk $\mathbb{D}_{p_i} = \{|z_j| \le 1\}$ around p_j , define the weight



FIGURE 5.1. Weight function μ_t

function by

$$\mu_t(z) := \min\left(\left(t^{-\frac{2K_1}{K_1+1}} + |z|^2\right)^{1/2}, 1\right).$$
(5.12)

On the rest of the surface, define

$$\mu_t(x) := 1 \qquad x \in C \setminus \{\mathbb{D}_{p_i}\}_{p_i \in Z}$$
(5.13)

The weight function μ_t (shown in Figure 5.1) is continuous. (Its lack of regularity is immaterial, and we could easily introduce a smoothed version.) Note that μ_t increases in |z| with minimum $\mu_t(0) = t^{-\frac{K_1}{K_1+1}}$. The family $\{\mu_t\}$ is uniformly bounded above by 1, and the family is also bounded away from 0 on any set where $|z| > \epsilon > 0$.

Let ψ_t denote an eigensection of the first eigenvalue λ_0^t . Fix some constant $\delta > 0$. (We will choose a good value of δ later in the proof.) We normalize ψ_t —multiplying it by a constant— so that

$$\sup_{C} \mu_t^{\delta} |\psi_t| = 1. \tag{5.14}$$

In what follows, we show the supremum of $\mu_t^{\delta} |\psi_t|$ cannot be achieved at any point of *C*—contradicting our initial assumption that $\lambda_0^t \to 0$.

CLAIM: There is no value of $\epsilon \in (0, 1]$ for which there exists a constant η such that for all *t*—or rather for some unbounded subsequence $\{t_k\}$ —

$$\sup_{\substack{C-\bigcup\\p_i\in Z}\{|z_j|\leq\epsilon\}}\mu_t^{\delta}|\psi_t|\geq\eta>0.$$
(5.15)

Proof of Claim: \triangleright Since (5.14) holds, then for any choice of $\epsilon > 0 |\psi_t| \le \mu_t^{-\delta} \le \epsilon^{-\delta}$ on $C - \bigcup_{p_j \in Z} \{|z_j| \le \epsilon\}$. Because the eigensections $\{\psi_t\}$ are uniformly bounded in L^{∞} on $C - \bigcup_{p_j \in Z} \{|z_j| < \epsilon\}$, by compactness, we may obtain a subsequence of ψ_t which converges in L^{∞} on the punctured surface C - Z to a section ψ_{∞} ; moreover, by elliptic regularity, the $\{\psi_t\}$ and limiting ψ_{∞} are in C^{∞} . In the region $C - \bigcup_Z \{|z_j| < \epsilon\}$, the coefficients of \mathbb{L}_t are converging smoothly; thus ψ_{∞} satisfies

$$\mathbb{L}_{\infty}\psi_{\infty} = 0 \qquad \text{on } C - Z. \tag{5.16}$$

Furthermore, $|\psi_{\infty}|$ is non-zero from our assumption in (5.15).

Using the local conic regularity theory at $p \in Z$ (see [Maz91, MW15]), ψ_{∞} has an asymptotic expansion in powers of r = |z| with coefficients which are trigonometric functions of the angular variable θ . Because ψ_{∞} is bounded at $p \in Z$, all of the powers of r in the expansion of ψ_{∞} at p are nonnegative. Now,

$$\langle \mathbb{L}_{\infty}\psi_{\infty},\psi_{\infty}\rangle = -\star d\star \langle d_{A_{\infty}}\psi_{\infty},\psi_{\infty}\rangle + \|d_{A_{\infty}}\psi_{\infty}\|^{2} + 2\|[\Phi_{\infty},\psi_{\infty}\|^{2}.$$
(5.17)

Note that $\star \langle d_{A_{\infty}} \psi_{\infty}, \psi_{\infty} \rangle$ vanishes at each point $p \in Z$ (These reasons are elaborated in a more general setting in (5.67).); hence, doing integration by parts,

$$0 = \langle L_{\infty}\psi_{\infty}, \psi_{\infty} \rangle_{L^{2}(C)} = \| \mathbf{d}_{A_{\infty}}\psi_{\infty} \|_{L^{2}(C)}^{2} + 2 \| [\Phi_{\infty}, \psi_{\infty}] \|_{L^{2}(C)}^{2}.$$
(5.18)

From Proposition 5.1, there is no global non-zero solution satisfying both $d_{A_{\infty}}\psi_{\infty} = 0$ and $[\Phi_{\infty}, \psi_{\infty}] = 0$, hence $\psi_{\infty} = 0$.

Now, suppose the claim is false, i.e. suppose there is a choice of $\epsilon \in (0, 1]$ and $\eta > 0$ such that for all *t*

$$\sup_{\substack{C-\bigcup\\p_i\in Z}}\mu_t^{\delta}|\psi_t| \ge \eta.$$
(5.19)

Then, $\sup_{C-\bigcup_{p_j\in \mathbb{Z}}\{|z_j|\leq \epsilon\}} |\psi_t| \geq \sup_{C-\bigcup_{p_j\in \mathbb{Z}}\{|z_j|\leq \epsilon\}} \eta \mu_t^{-\delta} \geq \eta$, so ψ_{∞} is non-zero, a contradiction. \triangleleft

Let $\{q_t\} \to \overline{q}$ be a convergent sequence of points at which the supremum of $\mu_t^{\delta} |\psi_t|$ in (5.14) is achieved. From the CLAIM, we see that $\overline{q} = p \in Z$ and that $\mu_t^{\delta} |\psi_t|$ tends to zero pointwise (and in fact uniformly in compact subsets) on C - Z. Let $z : \mathbb{D} \to \mathbb{C}$ be the chosen holomorphic coordinate centered at p. Define $z_t := z(q_t)$. Note that $\{z_t\}$ converge to zero because $\{q_t\}$ converge to p. Let K_0 be the largest integer less than or equal to K_1 for which $t^{\frac{K_0}{K_0+1}}z_t$ is bounded above by some constant R. Note that K_0 is automatically nonnegative. We will now show that if we assume that $\lambda_0^t \to 0$, then the supremum of (5.14) also can't be achieved at a point $p \in Z$, by separately considering two cases, depending on whether z_t converge to zero more quickly (CASE A, $K_0 > 0$) or more slowly (CASE B, $K_0 = 0$).

CASE A. $K_0 > 0$: Define a rescaling

$$\rho_{K_0,t}: z \to t^{\frac{K_0}{K_0+1}} z =: w.$$
(5.20)

Let $w_t = \rho_{K_0,t}(z_t)$, and note that $|w_t| \leq R$. Now, pullback and rescale the eigensections ψ_t , taking

$$\Psi_t := t^{-\frac{\delta K_0}{K_0 + 1}} (\rho_{K_0, t}^{-1})^* \psi_t.$$
(5.21)

On the disk, the bound in (5.14) is

$$\left(t^{-\frac{2K_1}{K_1+1}} + |z|^2\right)^{\delta/2} |\psi_t(z)| \le 1.$$
(5.22)

This implies that

$$\left(t^{-\frac{2K_1}{K_1+1}+\frac{2K_0}{K_0+1}}+|w|^2\right)^{\delta/2}|\Psi_t(w)| \le 1,$$
(5.23)

with equality attained at w_t . Since $K_0 \leq K_1$, $-\frac{2K_1}{K_1+1} + \frac{2K_0}{K_0+1} \leq 0$. Consequently,

$$|\Psi_{\infty}(w)| \leq \begin{cases} \left(1 + |w|^2\right)^{-\delta/2} & \text{if } K_0 = K_1 \\ |w|^{-\delta} & \text{if } K_0 < K_1. \end{cases}$$
(5.24)

Since the disk $\{|w| \leq R\}$ is compact, a subsequence of w_t converges to some \overline{w} , hence $|\Psi_{\infty}(\overline{w})| \neq 0$. By Lemma 5.6,

$$\lim_{t \to \infty} t^{-\frac{2K_0}{K_0+1}} (\rho_{K_0,t}^{-1})^* \mathbb{L}_t = \Delta_{\widetilde{A}} = \left(\bigoplus_{j: \ K_j > K_0} A_\infty \right) \oplus \left(\bigoplus_{j: \ K_j = K_0} A_{\text{mod}} \right) \oplus \left(\bigoplus_{j: \ K_j < K_0} A_0 \right).$$
(5.25)

The expressions for A_{∞} , A_{mod} , and A_0 are given in (5.40). Because the coefficients of the operators in (5.25) converge smoothly to $\Delta_{\tilde{A}}$, the non-zero Ψ_{∞} satisfies the bound in (5.24) and

$$\Delta_{\widetilde{A}} \Psi_{\infty} = 0. \tag{5.26}$$

By Proposition 5.8, for $\delta > 0$ sufficiently small, there is no non-zero solution. But Ψ_{∞} is non-zero! Thus, CASE A cannot hold.

CASE B. $K_0 = 0$: Lastly, suppose that $|\rho_{1,t}(z_t)| = |t^{1/2}z_t|$ is unbounded. Define a rescaling

$$\sigma_t : z \to |z_t|^{-1} z =: \widetilde{w} \tag{5.27}$$

Let $\tilde{w}_t = \sigma_t(z_t)$, and note that $|\tilde{w}_t| = 1$. Now, pullback the eigensection ψ_t and rescale it by an aptly chosen constant

$$\widetilde{\Psi}_t := |z_t|^{\delta} (\sigma_t^{-1})^* \psi_t \tag{5.28}$$

The constant is chosen so that the bound in (5.14) implies that

$$(|z_t|^{-2}t^{-\frac{2K_1}{K_1+1}} + |\widetilde{w}|^2)^{\delta/2} |\widetilde{\Psi}_t(\widetilde{w})| \le 1.$$
(5.29)

Taking the limit of the bounds in (5.29), we see that $\widetilde{\Psi}_{\infty}$ satisfies

$$|\widetilde{\Psi}_{\infty}(\widetilde{w})| \le |\widetilde{w}|^{-\delta}.$$
(5.30)

Since $\widetilde{\Psi}_t$ achieves the bound in (5.29) at \widetilde{w}_t which has norm one, $\widetilde{\Psi}_{\infty}$ also achieves the bound in (5.30) on the unit circle; hence $\widetilde{\Psi}_{\infty}$ is non-zero.

In the rescaling limit,

$$\lim_{t \to \infty} (\sigma_t^{-1})^* M_{\Phi_t} = M_{\Phi_\infty} \qquad \lim_{t \to \infty} |z_t|^{-2} (\sigma_t^{-1})^* \Delta_{A_t} = \Delta_{A_\infty}; \tag{5.31}$$

consequently,

$$\lim_{t \to \infty} |z_t|^{-2} (\sigma_t^{-1})^* \mathbb{L}_t = \Delta_{A_\infty},$$
(5.32)

where A_{∞} is defined in (5.40). Thus, $\tilde{\Psi}_{\infty}$ is non-zero and satisfies

$$\Delta_{A_{\infty}} \widetilde{\Psi}_{\infty}(\widetilde{w}) = 0, \qquad |\widetilde{\Psi}_{\infty}(\widetilde{w})| \le |\widetilde{w}|^{-\delta}.$$
(5.33)

By Proposition 5.8, for $\delta > 0$ sufficiently small, there is no non-zero solution. But $\widetilde{\Psi}_{\infty}$ is non-zero. Thus, CASE B too is impossible.

In summary, we have shown that it is impossible that $\lambda_0^t \to 0$.

Proof of Proposition 5.2b. The proof that $||L_t^{-1}||_{\mathcal{L}(L^2, H^2)} \leq \tilde{C}t^2$ is a direct adaption of the proof of in the SU(2) case [MSWW16, Lemma 6.5]. The graph norm of $\Delta_{A_{\infty}}$ is equivalent to the standard Sobolev H^2 -norm [MSWW16, Lemma 6.5]. Consequently, we will prove that there is a constant \tilde{C}' such that

$$\sqrt{\|L_t^{-1}u\|_{L^2}^2 + \|\Delta_{A_\infty}L_t^{-1}u\|_{L^2}^2} \le \tilde{C}'t^2\|u\|_{L^2}.$$
(5.34)

Define

$$\widetilde{L}_t = \Delta_{A_\infty} - \mathbf{i} \star t^2 M_{\Phi_\infty}.$$
(5.35)

(For comparison, recall from (5.7) that $L_t = \Delta_{A_t} - i \star t^2 M_{\Phi_t}$.) Then, note that

$$\|\Delta_{A_{\infty}}L_{t}^{-1}u\|_{L^{2}} \leq \|u\|_{L^{2}} + \|(\Delta_{A_{\infty}}-L_{t})L_{t}^{-1}u\|_{L^{2}}$$
(5.36)

$$\leq \|u\|_{L^{2}} + \left(t^{2}\|M_{\Phi_{\infty}}\|_{\mathcal{L}(L^{2},L^{2})} + \|(\widetilde{L}_{t}-L_{t})\|_{\mathcal{L}(L^{2},L^{2})}\right)\|L_{t}^{-1}\|_{\mathcal{L}(L^{2},L^{2})}\|u\|_{L^{2}}.$$

The bound $||M_{\Phi_{\infty}}||_{\mathcal{L}(L^{2},L^{2})} \leq c_{M}$ follows in Lemma 5.4. The bound $||\widetilde{L}_{t} - L_{t}||_{\mathcal{L}(L^{2},L^{2})} \leq Ce^{-\delta t}$ follows because Φ_{t} converges to Φ_{∞} exponentially in t and A_{t} converges to A_{∞} exponentially in t. (See [MSWW16, Lemma 6.5] for the case of $K \times K = 2 \times 2$ blocks.) The bound $||L_{t}^{-1}||_{\mathcal{L}(L^{2},L^{2})} \leq \widetilde{C}_{1}$ is from Proposition 5.2a. Thus we obtain the desired bound in (5.34).

In the proof of Proposition 5.2b, we used the following bound on $M_{\Phi_{\infty}}$.

Lemma 5.4. There is a constant c_M such that at any point of C

$$|M_{\Phi_{\infty}}|_{g_C,h_\ell} \le c_M. \tag{5.37}$$

Proof. Over Σ , $\pi^* \mathcal{E}$ decomposes as the sum of eigenline bundles of $\pi^* \varphi$. Let \mathcal{L}_i be the line bundle corresponding to globally-defined eigenvalue $\pi^* \lambda_i$. To see the bound on $M_{\Phi_{\infty}}$, note that, pulled-back from C to Σ , π^* End $\mathcal{E} = \oplus \text{Hom}(\mathcal{L}_i, \mathcal{L}_j)$ and the (i, j)-entry of $\pi^* M_{\Phi_{\infty}}$ is

$$(\pi^* M_{\Phi_{\infty}} \gamma)_{ij} = 2|\lambda_i - \lambda_j|^2 \gamma_{ij}.$$
(5.38)

The difference between the eigenvalues of φ are bounded above, hence $|M_{\Phi_{\infty}}|$ is bounded.

5.1. Lemmata for Proposition 5.2a: Local analysis of $\Delta_{\widetilde{A}}$. Take $p \in Z$ with associated partition $n = K_1 + \cdots + K_{m_p}$ and holomorphic coordinates z_1, \cdots, z_{m_p} centered at p in Proposition 2.2. Given a holomorphic coordinate z centered at p, define biholomorphic functions $f_i = z_i \circ z^{-1}$ such that $f_i(0) = 0$. Given a choice of positive number J, define

$$\widetilde{A} := \left(\bigoplus_{j: K_j > J} A_{\infty}\right) \oplus \left(\bigoplus_{j: K_j = J} A_{\text{mod}}\right) \oplus \left(\bigoplus_{j: K_j < J} A_0\right),$$
(5.39)

where the $K \times K$ blocks are

$$A_{\infty} = \begin{pmatrix} -\frac{\alpha_{K,1}}{2} & & \\ & \ddots & \\ & & -\frac{\alpha_{K,K}}{2} \end{pmatrix} \begin{pmatrix} \frac{dz}{z} - \frac{d\overline{z}}{\overline{z}} \end{pmatrix}$$
(5.40)
$$A_{\text{mod}} = \begin{pmatrix} -\frac{\alpha_{K,1}}{2} + \frac{|z|}{4} \frac{du_{K,1,1}(|f'(0)||z|)}{d|z|} & & \\ & & \ddots & \\ & & -\frac{\alpha_{K,K}}{2} + \frac{|z|}{4} \frac{du_{K,K,1}(|f'(0)||z|)}{d|z|} \end{pmatrix} \begin{pmatrix} \frac{dz}{z} - \frac{d\overline{z}}{\overline{z}} \end{pmatrix}$$
$$K \times K$$

$$A_{0} = \mathbf{0}_{K \times K}.$$



FIGURE 5.2. The values K_1, \dots, K_{m_p} are separated into three different categories: less than, equal to, or greater than some critical integer (here, 3). The limiting \tilde{A} consequently features three types of blocks: $A_0, A_{\text{mod}}, A_{\infty}$.

Example 5.5. *For example, if* n = 2 + 1 + 1*,*

$$\widetilde{A} = \left(\frac{1}{8} + \frac{|z|}{4}\frac{du_1}{d|z|}\right) \begin{pmatrix} 1 & 0 & 0 & 0\\ 0 & -1 & 0 & 0\\ 0 & 0 & 0 & 0\\ 0 & 0 & 0 & 0 \end{pmatrix} \left(\frac{dz}{z} - \frac{d\overline{z}}{\overline{z}}\right),$$
(5.41)

where $u_1 = u_{2,1,t=1}$ is the function in (4.2).

In this section, we prove that \tilde{A} appears naturally in a rescaling limit (Lemma 5.6). This is used in the proof of Proposition 5.2a in (5.58). We then prove in Proposition 5.8 that for $\delta > 0$ sufficiently small, there are no solutions of

$$\Delta_{\widetilde{A}} \Psi = 0 \qquad |\Psi| \le |z|^{-\delta}. \tag{5.42}$$

Lemma 5.6. Define a rescaling

$$\rho_{J,t}: z \mapsto t^{\frac{J}{J+1}} z. \tag{5.43}$$

Then, for the approximate solution (A_t, Φ_t) on \mathbb{D} in the unitary gauge of Remark 4.11, we have

$$\lim_{t \to \infty} t^{-\frac{2J}{J+1}} (\rho_{J,t}^{-1})^* \Delta_{A_t} = \Delta_{\widetilde{A}}.$$
(5.44)

Proof. First note that for any holomorphic function f such that f(0) = 0, we have

$$\lim_{t \to \infty} (f \circ \rho_{J,t}^{-1})^* \left(\frac{\mathrm{d}z}{z} - \frac{\mathrm{d}\overline{z}}{\overline{z}} \right) = \left(\frac{\mathrm{d}z}{z} - \frac{\mathrm{d}\overline{z}}{\overline{z}} \right).$$
(5.45)

This follows from expanding the holomorphic—hence analytic—function f as $f(x) = \sum_{i=0}^{\infty} \frac{1}{i!} f^{(i)}(0) x^i$ in the following expression:

$$\lim_{t \to \infty} \frac{\mathrm{d}(f(t^{-\frac{J}{J+1}}w))}{f(t^{-\frac{J}{J+1}}w)} = \frac{\lim_{t \to \infty} f'(t^{-\frac{J}{J+1}}w)\mathrm{d}w}{t^{\frac{J}{J+1}}f(t^{-\frac{J}{J+1}}w)} = \frac{f'(0)\mathrm{d}w}{f'(0)w} = \frac{\mathrm{d}w}{w}.$$
(5.46)

Secondly, note that for similar reasons

$$\lim_{t \to \infty} (f \circ \rho_{J,t}^{-1})^* \left(\frac{|z|}{4} \frac{d}{d|z|} \right) = \frac{|z|}{4} \frac{d}{d|z|}.$$
(5.47)

(Here, it is convenient to use that $|z|\frac{\partial}{\partial |z|} = z\frac{\partial}{\partial z} + \overline{z}\frac{\partial}{\partial \overline{z}}$.)

We work separately in each $K \times K$ block. The computation of the limits is based on the following observation:

$$(\rho_{J,t}^{-1})^* u_{K,i,t}(r) = u_{K,i,t}\left(t^{-\frac{J}{J+1}}r\right) = u_{K,i,t}\left(t^{-\frac{K}{K+1}} t^{\frac{K}{K+1} - \frac{J}{J+1}}r\right) = u_{K,i,t-1}\left(t^{\frac{K}{K+1} - \frac{J}{J+1}}r\right)$$
(5.48)

We see that if J < K, then $\frac{K}{K+1} - \frac{J}{J+1} > 0$, so $\lim_{t\to\infty} t^{\frac{K}{K+1} - \frac{J}{J+1}} r = \infty$. Each function $u_{K,i,t}$ decays to 0 at ∞ , hence

$$\lim_{t \to \infty} (f \circ \rho_{J,t}^{-1})^* \left(\frac{|z|}{4} \frac{\mathrm{d}u_{K,i,t}(|z|)}{\mathrm{d}|z|} \right) = 0, \tag{5.49}$$

and consequently, in the limit of (5.44), the $K \times K$ block is A_{∞} .

If J = K, then $\frac{K}{K+1} - \frac{J}{J+1} = 0$, so $\lim_{t\to\infty} t^{\frac{K}{K+1} - \frac{J}{J+1}} z = z$. Additionally, note that because f is analytic

$$\left(f \circ \rho_{K,t}^{-1} \right)^* u_{K,i,t}(|z|) = u_{K,i,t} \left(\left| f(t^{-\frac{K}{K+1}}z) \right| \right)$$

$$= u_t \left(\left| \sum_{k=0}^{\infty} \frac{1}{k!} f^{(k)}(0) \left(t^{-\frac{K}{K+1}}z \right)^k \right| \right)$$

$$= u_1 \left(\left| \sum_{k=1}^{\infty} \frac{1}{k!} f^{(k)}(0) \left(t^{-\frac{K}{K+1}} \right)^{k-1} z^k \right| \right)$$

$$(5.50)$$

In the last line, we use that $u_t(t^{-\frac{K}{K+1}}r) = u_1(r)$ and f(0) = 0. Consequently, taking the limit, we obtain

$$\lim_{t \to \infty} \left(u_t \left(\left| f(t^{-\frac{K}{K+1}} z) \right| \right) = \lim_{t \to \infty} u_1 \left(\left| \sum_{k=1}^{\infty} \frac{1}{k!} f^{(k)}(0) \left(t^{-\frac{K}{K+1}} \right)^{k-1} z^k \right| \right) = u_1 \left(\left| f'(0) \right| |z| \right).$$
(5.51)

Hence,

$$\lim_{t \to \infty} (f \circ \rho_{J,t}^{-1})^* \left(\frac{|z|}{4} \frac{\mathrm{d}u_{K,i,t}(|z|)}{\mathrm{d}|z|} \right) = \left(\frac{|z|}{4} \frac{\mathrm{d}u_{K,i,t=1}(|f'(0)| |z|)}{\mathrm{d}|z|} \right); \tag{5.52}$$

in the limit of (5.44), the $K \times K$ block is A_{mod} .

Lastly, if J > K, then $\frac{K}{K+1} - \frac{J}{J+1} < 0$, so $\lim_{t\to\infty} t^{\frac{K}{K+1} - \frac{J}{J+1}} z = 0$. Since $\frac{|z|}{4} \frac{du_{K,i,1}(0)}{d|z|} = \frac{\alpha_{K,i}}{2}$, we have

$$\lim_{t \to \infty} (f \circ \rho_{J,t}^{-1})^* \left(-\frac{\alpha_{K,i}}{2} + \frac{|z|}{4} \frac{\mathrm{d}u_{K,i,t}(|z|)}{\mathrm{d}|z|} \right) = 0;$$
(5.53)
the $K \times K$ block is A_0 .

in the limit of (5.44), the $K \times K$ block is A_0 .

We now analyze the kernel of $\Delta_{\widetilde{A}}$. The operator $\Delta_{\widetilde{A}}$ on the punctured plane $\mathbb{C} \setminus \{0\}$ is an differential edge operator³, and so the theory in [Maz91, MW15] applies. We first compute the indicial roots of the operator. A number $\nu \in \mathbb{C}$ is called an indicial root for

³This is roughly because, in polar coordinates, the operator $r^2\Delta_{\widetilde{A}}$ can be written in terms of $r\partial_r$ and ∂_{θ} .

 $\Delta_{\widetilde{A}}$ if there exists some function $\zeta = \zeta(\theta)$ such that

$$\Delta_{\widetilde{A}}(r^{\nu}\zeta(\theta)) = O(r^{\nu-1}), \tag{5.54}$$

rather than the expected rate $O(r^{\nu-2})$ [MSWW16, Definition 4.2]. If $\Delta_{\widetilde{A}}\psi = 0$, we see that ψ has a inhomogeneous asymptotic development around 0 (or ∞) in terms of the indicial roots of $\Delta_{\widetilde{A}}$ at 0 (or, respectively, ∞).

Lemma 5.7. In this basis of $\mathfrak{sl}(n, \mathbb{C})$, the operator $\Delta_{\widetilde{A}}$ fully decouples. On in (i, j) block, $\Delta_{\widetilde{A}}$ acts as

$$\Delta_{\widetilde{A}}\gamma_{ij} = \left(d + (\widetilde{A}_{ii} - \widetilde{A}_{jj})\right)^* \left(d + (\widetilde{A}_{ii} - \widetilde{A}_{jj})\right)\gamma_{ij}.$$
(5.55)

The set indicial roots of

$$\Delta_{\widetilde{A}}: \Gamma(\mathfrak{isu}(n)) \to \Gamma(\mathfrak{isu}(n)) \tag{5.56}$$

at |z| = 0 is $\Gamma(\Delta_{\widetilde{A}}, 0) = \mathbb{Z} \sqcup S_0$, where S_0 is a discrete set, symmetric around the origin, with $S_0 \subset (-1, 1)$. Similarly, the indicial roots at $|z| = \infty$ is $\Gamma(\Delta_{\widetilde{A}}, \infty) = \mathbb{Z} \sqcup S_\infty$, where S_∞ is a discrete set, symmetric around the origin, with $S_\infty \subset (-1, 1)$.

Proof. The operator $\Delta_{\widetilde{A}}$ decouples as in (5.55) because \widetilde{A} is diagonal. Because $\widetilde{A}_{jj} = 2if_i(r)d\theta$, we can compute that

$$\Delta_{\widetilde{A}}\gamma_{ij} = \left(\mathbf{d} + (\widetilde{A}_{ii} - \widetilde{A}_{jj})\right)^* \left(\mathbf{d} + (\widetilde{A}_{ii} - \widetilde{A}_{jj})\right)\gamma_{ij}$$

$$= r^{-2} \left((r\partial_r)^2 + \left(\partial_\theta + 2\mathbf{i}(f_i - f_j)\right)^2\right)\gamma.$$
(5.57)

To compute the indicial roots at 0 for $\Delta_{\widetilde{A}}$ acting on $\Gamma(\mathfrak{sl}(n, \mathbb{C}))$, we only need to look at the highest order part of $\Delta_{\widetilde{A}}$ at 0. We evaluate the function $f_i(r) - f_j(r)$ appearing in (5.57) at r = 0. In a $K \times K$ block of A_{∞} , $f_i(0) = -\frac{\alpha_{K,i}}{2}$; for both A_{mod} and A_0 , this constant is 0. Then, taking $b_{ij} = f_i(0) - f_j(0)$, we see the relevant operator is

$$\left((r\partial_r)^2 + \left(\partial_\theta + 2ib_{ij} \right)^2 \right) \gamma.$$
(5.58)

Suppose ν is an indicial root at 0 in (i, j) block. Then there is some function $\zeta(\theta) = \sum_{\ell \in \mathbb{Z}} a_{\ell} e^{i\ell\theta}$ such that $\Delta_{\widetilde{A}}|_{ij} (r^{\nu}\zeta(\theta)) = O(r^{\nu-1})$. Taking $\zeta(\theta) = e^{i\ell\theta}$,

$$0 = \left((r\partial_r)^2 + (\partial_\theta + 2ib_{ij})^2 \right) (r^{\nu} e^{i\ell\theta})$$

= $r^{\nu} e^{i\ell\theta} \left(\nu^2 - (\ell + 2b_{ij})^2 \right).$ (5.59)

Consequently, since $\ell \in \mathbb{Z}$, $\nu \in \{\mathbb{Z} + 2b_{ij}\} \cup \{\mathbb{Z} - 2b_{ij}\}$.

Further restricting to $i\mathfrak{su}(n)$, we note that $\gamma_{ji} = \overline{\gamma}_{ij}$. We compute the indicial roots at 0 for the direct sum of the (i, j)-block with the (j, i)-block. Letting $\gamma_{ij} = r^{\nu} \zeta(\theta)$, and taking

 $\zeta(\theta) = a_{\ell} e^{i\ell\theta} + a_{-\ell} e^{-i\ell\theta}$ we have

$$\Delta_{\widetilde{A}}\Big|_{ij\oplus ji} \begin{pmatrix} r^{\nu}\zeta(\theta)\\ r^{\nu}\overline{\zeta}(\theta) \end{pmatrix} = \begin{pmatrix} \left(r^{2}\partial_{r}^{2} + r\partial_{r} + \left(\partial_{\theta} + 2\mathrm{i}b_{ij}\right)^{2}\right)r^{\nu}\left(a_{\ell}\mathrm{e}^{i\ell\theta} + a_{-\ell}\mathrm{e}^{-i\ell\theta}\right)\\ \left(r^{2}\partial_{r}^{2} + r\partial_{r} + \left(\partial_{\theta} - 2\mathrm{i}b_{ij}\right)^{2}\right)r^{\nu}\left(\overline{a}_{\ell}\mathrm{e}^{-i\ell\theta} + \overline{a}_{-\ell}\mathrm{e}^{i\ell\theta}\right) \end{pmatrix}$$
(5.60)
$$= \begin{pmatrix} r^{\nu}a_{\ell}\mathrm{e}^{i\ell\theta}\left(\nu^{2} - \left(\ell + 2b_{ij}\right)^{2}\right) + r^{\nu}a_{-\ell}\mathrm{e}^{-i\ell\theta}\left(\nu^{2} - \left(-\ell + 2b_{ij}\right)^{2}\right)\\ r^{\nu}\overline{a}_{\ell}\mathrm{e}^{-i\ell\theta}\left(\nu^{2} - \left(-\ell + 2b_{ij}\right)^{2}\right) + r^{\nu}\overline{a}_{-\ell}\mathrm{e}^{i\ell\theta}\left(\nu^{2} - \left(\ell + 2b_{ij}\right)^{2}\right) \end{pmatrix} \end{pmatrix}$$

Thus, we see that $\nu \in \{\ell + 2b_{ij}, -\ell - 2b_{ij}\} \cap \{-\ell + 2b_{ij}, \ell - 2b_{ij}\}$. Thus, as claimed:

- If $b_{ij} = 0$, then $\nu \in \mathbb{Z}$; the indicial root ν comes from $r^{\nu}\zeta(\theta)$ where $\zeta(\theta) = a_{\nu}e^{i\nu\theta} + a_{-\nu}e^{-i\nu\theta}$. (Note that the $\nu = 0$ indicial root comes from $a_0 + \hat{a}_0 \log r$.)
- If $b_{ij} \neq 0$, then $\nu \in \{\pm 2b_{ij}\}$; the indicial root ν comes from $r^{\nu}\zeta(\theta)$ where $\zeta(\theta)$ is constant.

It is worth noting that because $\alpha_{K,i} \in (-\frac{1}{2}, \frac{1}{2})$, we automatically have that $b_{ij} \in (-\frac{1}{2}, \frac{1}{2})$ as well. Hence, the indicial roots of $\Delta_{\widetilde{A}}$ at 0 are $\mathbb{Z} \sqcup S_0$, where S_0 is a discrete set, symmetric around the origin, with $S_0 \subset (-1, 1)$.

The computation of the indicial roots at ∞ is similar. We let $v = z^{-1}$ and introduce polar coordinates $v = se^{i\vartheta}$. In these coordinates, the operator in (5.57) is

$$r^{2}\Delta_{\widetilde{A}}\gamma_{ij} = \left((s\partial_{s})^{2} + \left(-\partial_{\vartheta} + 2\mathrm{i}(f_{i} - f_{j})\right)^{2}\right)\gamma.$$
(5.61)

(Here, we used that $r\partial_r = -s\partial_s$ and $\partial_\theta = -\partial_\theta$.) The computation of the indicial roots is similar. In a $K \times K$ block of A_∞ or A_{mod} , $f_i(\infty) = -\frac{\alpha_{K,i}}{2}$; for A_0 , this constant is 0. Define the constant $c_{ij} = f_i - f_j|_{r=\infty}$. Thus, as claimed:

• If $c_{ij} = 0$, then $\nu \in \mathbb{Z}$; the indicial root ν comes from $s^{\nu}\zeta(\vartheta)$ where $\zeta(\vartheta) = a_{\nu}e^{i\nu\vartheta} + a_{-\nu}e^{-i\nu\vartheta}$.

• If $c_{ij} \neq 0$, then $\nu \in \{\pm 2c_{ij}\}$; the indicial root ν comes from $s^{\nu}\zeta(\vartheta)$ where $\zeta(\vartheta) = a_0$. Hence, the indicial roots of $\Delta_{\widetilde{A}}$ at $r = \infty$ are $\mathbb{Z} \sqcup S_{\infty}$, where S_{∞} is a discrete set, symmetric around the origin, with $S_{\infty} \subset (-1, 1)$.

Using the computation of the indicial roots in Lemma 5.7, we now prove:

Proposition 5.8. There exists a $\delta > 0$ such that there are no non-zero solutions $\psi \in \Gamma(i\mathfrak{su}(n))$ solving

$$\Delta_{\widetilde{A}}\psi = 0 \qquad |\psi| \le (\epsilon + |z|^2)^{-\delta/2} \tag{5.62}$$

where $\epsilon = 0, 1$.

Proof. In the proof, we will integrate by parts and then conclude that there is no solution of $d_{\tilde{A}}\psi = 0$ solving the above bound. Because

$$\star \langle \Delta_{\widetilde{A}} \psi, \psi \rangle = \star \| \mathbf{d}_{\widetilde{A}} \psi \|^2 + \mathbf{d} \star \langle \mathbf{d}_{\widetilde{A}} \psi, \psi \rangle, \tag{5.63}$$

we will additionally need to show that $\lim_{r\to 0,\infty} \int_{S^1_r} \star \langle d_{\widetilde{A}} \psi, \psi \rangle = 0.$

Choose δ satisfying

$$\delta < \min\{|\nu| : \nu \in S_0 \cup S_\infty\},\tag{5.64}$$

noting that $S_0 \cup S_\infty$ is a discrete set that does not contain 0. Let $N_0 = (S_0 \cup \mathbb{Z}) \cap (-\delta, \infty)$. Similarly, let $N_\infty = (S_\infty \cup \mathbb{Z}) \cap (\delta, \infty)$.

The bound at 0 implies that ψ admits an asymptotic development around |z| = 0 like

$$\psi \sim \sum_{\nu \in N_0 + \mathbb{N}} r^{\nu} \zeta_{\nu}(\theta).$$
(5.65)

As shown in Figure 5.3, because of our choice of δ , $N_0 + \mathbb{N} \subset [0, \infty)$.



FIGURE 5.3. The indicial roots at 0 are $S_0 \sqcup \mathbb{Z}$. With the given choice of δ , the set of all powers in the expansion are nonnegative.

Furthering studying the $\star \langle \mathbf{d}_{\widetilde{A}} \psi, \psi \rangle$ term in (5.63), let $\widetilde{A} = \Xi(r) 2i d\theta$. Then, that

$$\begin{aligned} \star \langle \mathbf{d}_{\widetilde{A}} \psi, \psi \rangle &= \star \langle \partial_{r} \psi \mathrm{d}r + \partial_{\theta} \psi \mathrm{d}\theta + 2\mathrm{i}[\Xi, \psi] \mathrm{d}\theta, \psi \rangle \\ &= \langle \partial_{r} \psi, \psi \rangle r \mathrm{d}\theta - \langle \partial_{\theta} \psi + 2\mathrm{i}[\Xi, \psi], \psi \rangle r^{-1} \mathrm{d}r. \end{aligned}$$
(5.66)

To see that $\lim_{r\to 0} \int_{S_r^1} \star \langle \mathbf{d}_{\widetilde{A}} \psi, \psi \rangle = 0$, note that the $\langle \partial_r \psi, \psi \rangle r d\theta$ term tends to zero pointwise. If we integrate on S_r^1 , then the dr component does not matter. However, we want to show that it's not problematic to perturb the loop S_r^1 —even though the indicial roots $\nu \leq 1$ look problematic because of the " r^{-1} ". Since $\psi \in \Gamma(\mathfrak{isu}(n))$, it is a straightforward computation to check that

$$\langle \partial_{\theta} \psi + 2i[\Xi, \psi], \psi \rangle r^{-1} dr = \langle \partial_{\theta} \psi, \psi \rangle r^{-1} dr.$$
(5.67)

From the indicial root computation in Lemma 5.7, we see that if $\nu < 1$, then $\zeta'_{\nu}(\theta) = 0$; thus the problematic-looking terms in (5.66) vanish. It follows that $\lim_{r \to 0} \int_{S_r^1} \star \langle d_{\widetilde{A}} \psi, \psi \rangle = 0$.

Similarly, in the coordinate $v = z^{-1} = se^{i\vartheta}$, the bound on ψ is given by $\psi(v) \le |v|^{\delta}$. As shown in Figure 5.4, because of our choice of δ , $N_{\infty} + \mathbb{N} \subset (0, \infty)$. Using a similar



FIGURE 5.4. The indicial roots at ∞ are $S_{\infty} \sqcup \mathbb{Z}$. With the given choice of δ , the set N_{∞} of all permissible indicial roots are positive.

computation, we can see that $\lim_{r\to\infty}\int_{S_r^1}\star\langle \mathsf{d}_{\widetilde{A}}\psi,\psi\rangle=0.$

Consequently, by (5.63), we can equivalently show that there is no non-zero solution $\psi \in \Gamma(i\mathfrak{su}(n))$ solving

$$\mathbf{d}_{\widetilde{A}}\psi = 0 \qquad |\psi| \le (\epsilon + |z|^2)^{-\delta/2} \tag{5.68}$$

where $\epsilon = 0, 1$. We now analyze now analyze this problem. The (i, j) block of $d_{\tilde{A}}\gamma$ is

$$0 = (\mathbf{d}_{\widetilde{A}}\gamma)_{ij} = \partial_r \gamma_{ij} \mathbf{d}r + \partial_\theta \gamma_{ij} \mathbf{d}\theta + (f_i - f_j)\gamma_{ij} 2\mathbf{i}\mathbf{d}\theta.$$
(5.69)

Note the following consequence:

$$0 = \partial_{\theta}(\partial_r \gamma_{ij}) = \partial_r(\partial_{\theta} \gamma_{ij}) = \partial_r(2\mathbf{i}(f_j - f_i)\gamma_{ij}).$$
(5.70)

Consequently, there are three distinct cases.

- (1) If $f_j f_i = 0$, then γ_{ij} is constant. Imposing the asymptotic decay condition, we see that $\gamma_{ij} = 0$.
- (2) If $f_i f_i$ is not constant, it follows that $\gamma_{ij} = 0$ (using (5.70) and $\partial_r \gamma_{ij} = 0$).
- (3) If $f_j f_i$ is a non-zero constant, then we see that we'd like to take $\gamma_{ij} = ce^{2i\theta(f_j f_i)}$. However, this is a function on the punctured-plane if, and only if $f_j - f_i \in \mathbb{Z}$. However, note that for each function $f_i(r)$, $f_i(0) \in (-\frac{1}{2}, \frac{1}{2})$. Consequently, this case is not possible.

Note that in the last two cases we did not use the decay condition.

Thus, we've proved that there is no non-zero solution of (5.68) and thereby no non-zero solution of (5.62).

Remark 5.9. The proof is easier in the case where there are no blocks A_{mod} appearing in \tilde{A} . In this case, $\Delta_{\tilde{A}}$ is a dilation-covariant conic operator. Solutions are superpositions of functions of the form $r^{\nu}\zeta_{\nu}(\theta)$ where ν is an indicial root of the operator. The bounds at 0 force $\nu \geq 0$, while the bounds at ∞ force $\nu < 0$. These are incompatible, hence the zero solution is the only solution.

6. PERTURBATION TO A SOLUTION OF HITCHIN'S EQUATIONS

In this section we prove that h_t^{app} is close to the harmonic metric h_t in the space of hermitian metrics. We work in unitary formulation of Hitchin's equations, discussed at the beginning of §5.

Main Theorem 6.1. Fix a Higgs bundle $(\overline{\partial}_E, \varphi) \in \mathcal{M}'$ and let δ be the constant in Proposition 4.10. Given any $\epsilon > 0$ and choice of t_0 , there exists a constant C such that for $t > t_0$ there is a unique h_0 -hermitian γ_t satisfying $\|\gamma_t\|_{H^2(\mathfrak{isu}(E))} \leq Ce^{(-\delta+\epsilon)t}$, such that $\mathbf{F}_t^{\operatorname{app}}(\gamma_t) = 0$, i.e. $(\mathbf{d}_{A_t^{\exp(-\gamma_t)}}, \Phi_t^{\exp(-\gamma_t)}))$ solves Hitchin's equations. (Equivalently, $h_t(v, w) = h_t^{\operatorname{app}}(e^{-\gamma_t}v, e^{-\gamma_t}w)$ is harmonic.) Moreover, there exists a constant C' such that γ_t is unique in the the ball of radius $C't^{-4-\epsilon}$.



FIGURE 6.1. Theorem 6.1 describes the size of γ_t solving $\mathbf{F}_t^{\text{app}}(\gamma_t) = 0$, i.e. solving Hitchin's equations. Alternatively, it describes the relation between h_t and h_t^{app} in the space {hermitian metrics} $\times \mathbb{R}_t^+$. In this language, the dotted blue curve would be labeled h_t^{app} and the solid green curve would be labeled h_t .

Theorem 6.1 will be proved using a contraction mapping argument, as in [MSWW16]. The map F_t^{app} defined in (5.6) is naturally a map between the following Sobolev spaces

$$\mathbf{F}_t^{\mathrm{app}}: H^2(i\mathfrak{su}(E)) \to L^2(i\mathfrak{su}(E)).$$
(6.1)

Observe that $\mathbf{F}_t^{\text{app}}(\gamma_t) = 0$ if, and only if, γ_t is a fixed point of the map

$$\mathbf{T}_{t}: H^{2}(i\mathfrak{su}(E)) \rightarrow H^{2}(i\mathfrak{su}(E).$$

$$\gamma \mapsto \gamma - L_{t}^{-1} \left(\mathbf{F}_{t}^{\mathrm{app}}(\gamma) \right).$$
(6.2)

Expanding $\mathbf{F}_{t}^{\text{app}}$ into a constant, linear, and non-linear term $\mathbf{F}_{t}^{\text{app}}$

$$\mathbf{F}_{t}^{\mathrm{app}}(\gamma) = \mathbf{F}_{t}^{\mathrm{app}}(0) + L_{t}(\gamma) + Q_{t}(\gamma), \tag{6.3}$$

the map \mathbf{T}_t is

$$\mathbf{T}_{t}(\gamma) = -(L_{t})^{-1} (\mathbf{F}_{t}^{\mathrm{app}}(0) + Q_{t}(\gamma)).$$
(6.4)

To show there is some ball $B_{\rho_t} \in H^2(i\mathfrak{su}(E))$ centered at the zero section (corresponding to h_t^{app}) on which \mathbf{T}_t is a contraction mapping of B_{ρ_t} , we additionally need an estimate on the nonlinear terms in the expansion of the operator \mathbf{T}_t in (6.4).

6.1. Estimates for nonlinear terms. We prove the analog of [MSWW16, Lemma 6.8] for the case of regular $SL(n, \mathbb{C})$ -Higgs bundles.

Lemma 6.2. The approximate solution satisfies

$$\|A_t\|_{C^1} \le Ct \tag{6.5}$$

on the disk \mathbb{D} , so that for any H^{k+1} section γ , k = 0, 1,

$$\|\mathbf{d}_{A_t}\gamma\|_{H^k} \le Ct \|\gamma\|_{H^{k+1}},\tag{6.6}$$

and moreover,

$$\|L_t \gamma\|_{L^2} \le Ct^2 \|\gamma\|_{H^2}.$$
(6.7)

Proof. We first prove the bound in (6.5). From Remark 4.11, A_t is diagonal with diagonal elements like

$$-\frac{\alpha_{K,i}}{2} + \frac{|z|}{4} \frac{d(u_{K,i,t}\chi)}{d|z|}.$$
(6.8)

This has the same asymptotics as

$$f_{i,t} := -\frac{\alpha_{K,i}}{2} + \frac{|z|}{4} \frac{\mathrm{d}u_{K,i,t}}{\mathrm{d}|z|},\tag{6.9}$$

thus the diagonal elements of A_t are uniformly bounded in t. Since, we're computing the C^1 bound, we will show that $|\partial_{|z|}f_{i,t}| \leq Ct$. Recall that $u_{K,i,t} = \zeta_t^* v_{K,i}$ where $v_{K,i}$ solve the affine Toda lattice in (4.9) and $\zeta_t(|z|) = \frac{2K}{K+1}|z|^{\frac{K+1}{K}}$. Consequently,

$$f_{i,t} = \zeta_t^* g_{K,i} \qquad g_{K,i}(\zeta) := \left(-\frac{\alpha_{K,i}}{2} + \frac{(K+1)\zeta}{4K} \zeta \partial_{\zeta} v_{K,i}(\zeta) \right).$$
(6.10)

Note that $\lim_{\zeta \to 0} g'_{K,i}(\zeta) = 0$ and $g'_{K,i}(\zeta)$ asymptotically decays like $e^{-c_1\zeta}$ from (4.8). Thus there is a constant $C_1 > 0$ such that for all z,

$$\left|f_{i,t}'(|z|)\right| = \left|g_{K,i}'(\zeta_t(|z|)) \cdot 2t|z|^{\frac{1}{K}}\right| \le C_1 t.$$
(6.11)

Consequently the derivative of (6.8) obeys a similar bound. The estimate in (6.5) follows.

The bounds in (6.6) and (6.7) are immediate corollaries. For (6.6), observe that

$$\|\mathbf{d}_{A_{t}}\gamma\|_{H^{k}} \leq \|\mathbf{d}\gamma\|_{H^{k}} + \|[A_{t},\gamma]\|_{H^{k}} \leq \|\gamma\|_{H^{k+1}} + \|A_{t}\|_{C^{k}}\|\gamma\|_{H^{k}} \leq C't\|\gamma\|_{H^{k+1}}.$$
 (6.12)

The bound in (6.7) follows from (6.6) and the bound on M_{Φ_t} in (5.37).

Using this lemma, we can derive the following estimate on the nonlinear terms in (6.3).

Lemma 6.3 (Estimate on nonlinear terms). [MSWW16, Lemma 6.9] *There exists a constant* $\hat{C} > 0$ *such that*

$$\|Q_t(\gamma_1) - Q_t(\gamma_2)\|_{L^2} \le \widehat{C}\xi t^2 \|\gamma_1 - \gamma_2\|_{H^2}$$
(6.13)

for all $\xi \in (0,1]$ and γ_1, γ_2 satisfying $\|\gamma_i\| \leq \xi$.

Proof of Lemma 6.3. The proof in [MSWW16, Lemma 6.9] carries over to the case of regular $SL(n, \mathbb{C})$ -Higgs bundles using Lemma 6.2 in place of [MSWW16, Lemma 6.8].

6.2. Main Theorem.

Proof of Main Theorem 6.1. For all $\xi \in (0, 1]$ and $t > t_0$, hermitian sections γ_1, γ_2 satisfying $\|\gamma_i\| \leq \xi$,

$$\|\mathbf{T}_{t}(\gamma_{1}-\gamma_{2})\|_{H^{2}} = \|-L_{t}^{-1}\left(Q_{t}(\gamma_{1})-Q_{t}(\gamma_{2})\right)\|_{H^{2}}$$

$$\leq \|L_{t}^{-1}\|_{\mathcal{L}(L^{2},H^{2})}\|\|Q_{t}(\gamma_{1})-Q_{t}(\gamma_{2})\|_{L^{2}}$$

$$\leq \tilde{C}t^{2}\cdot\hat{C}\xi t^{2}\|\gamma_{1}-\gamma_{2}\|_{L^{2}}.$$
(6.14)

In the last line, we used the bounds in Lemma 6.3 and Proposition 5.2. Consequently setting ξ equal to $r_t = \frac{1}{\tilde{C}\tilde{C}t^4}$, \mathbf{T}_t is a contraction on the ball of radius r_t .

To see that there is a radius $\rho_t \leq r_t$ such that $\mathbf{T}_t(B_{\rho_t}) \subset B_{\rho_t}$, note that

$$\|\mathbf{T}_{t}(\gamma)\|_{H^{2}} \leq \tilde{C}t^{2} \cdot \hat{C}\xi t^{2} \|\gamma\|_{L^{2}} + \|\mathbf{T}_{t}(0)\|_{H^{2}}$$

$$\leq \tilde{C}t^{2} \cdot \hat{C}\xi t^{2} \|\gamma\|_{L^{2}} + \tilde{C}t^{2}Ce^{-\delta t}.$$
(6.15)

Note that $\|\mathbf{T}_t(0)\|$ decays exponentially in t (Proposition 4.10) like $ce^{-\delta t}$ for $t > t_0$. Consequently, given any $\epsilon > 0$, there exists a constant C such that $\mathbf{T}_t(B_{\rho_t}) \subset B_{\rho_t}$ for $\xi = \rho_t = Ce^{-(\delta + \epsilon)t}$. Moreover, there exists a constant C' such that $\mathbf{T}_t(B_{\rho_t'}) \subset B_{\rho_t'}$ for $\xi = \rho_t' = C't^{-4-\epsilon}$.

 \mathbf{T}_t is a contraction mapping on B_{ρ_t} and $B_{\rho'_t}$. Consequently, there is exists a fixed point γ_t of \mathbf{T}_t in the ball $B_{Ce^{(-\delta+\epsilon)t}}$. Moreover, it is unique in the ball $B_{C't^{-4-\epsilon}}$.

Corollary 6.4. As $t \to \infty$, the harmonic metrics h_t converge pointwise to h_{∞} .

Proof. The approximate metrics h_t^{app} converge to h_{∞} pointwise by construction. The metric $h_t(v, w) = h_t^{app}(e^{-\gamma_t}v, e^{-\gamma_t}w)$; consequently, because $\gamma_t \to 0$, h_t also converge to h_{∞} pointwise.

Remark 6.5. It's worth noting that this description of h_t in terms of h_t^{app} does not work uniformly in \mathcal{M}' . As discussed in §2.3, in our construction of h_t^{app} , we have stratified \mathcal{M}' . Our construction works uniformly provided we stay in a single piece of the stratification. However, the radius of the disks \mathbb{D}_p we use in the desingularization of h_ℓ goes to zero when we pass between strata, as shown in Figure 6.2. Ideally, our construction would



FIGURE 6.2. Local model of spectral cover as one passed from (LEFT) $N_2 = 4$ to (RIGHT) $N_5 = 1$.

work uniformly on all of \mathcal{M}' since it should not be problematic for ramification points to coalesce unless there is a collapsing cycle in the spectral cover, as shown in Figure 6.3.

REFERENCES

- [B00] C. Bär, "The Dirac operator on hyperbolic manifolds of finite volume," J. Differential Geom. 54 no. 3, (2000) 439–488, arXiv:math/0010233.
- [Biq96] O. Biquard, "Sur les fibrés paraboliques sur une surface complexe," J. London Math. Society 53 (1996) 392–216.



FIGURE 6.3. If the two indicated points of *Z* coalesce, then a cycle in Σ collapses.

- [CV91] S. Cecotti and C. Vafa, "Topological-antitopological fusion," Nucl. Phys. B367 (1991) 359–461.
- [CV92] S. Cecotti and C. Vafa, "Exact Results for Supersymmetric Sigma Models," Phys. Rev. Lett 68 (1992) 0903–906, arXiv:hep-th/9111016.
- [FN17] L. Fredrickson and A. Neitzke, "From S¹-fixed points to W-algebra representations," arXiv:1709.06142.
- [GL10] M. Guest and C.-S. Lin, "Nonlinear PDE aspects of the *tt** equations of Cecotti and Vafa," arXiv:1010.1889.
- [GL13] M. Guest and C.-S. Lin, "Isomonodromy aspects of the *tt** equations of Cecotti and Vafa II. Riemann-Hilbert problem," arXiv:1312.4825.
- [GMN09] D. Gaiotto, G. W. Moore, and A. Neitzke, "Wall-crossing, Hitchin systems, and the WKB approximation," arXiv:0907.3987.
- [GMN10] D. Gaiotto, G. W. Moore, and A. Neitzke, "Four-dimensional wall-crossing via threedimensional field theory," *Commun. Math. Phys.* **299** (2010) 163–224, arXiv:0807.4723.
- [Mas86] H. Masur, "Lower bounds for the number of saddle connections and closed trajectories of a quadratic differential," in *Holomorphic Functions and Moduli I*, pp. 215–228. 1986.
- [Maz91] R. Mazzeo, "Elliptic theory of differential edge operators, I," Comm. Par. Diff. Eqns 16 no. 10, (1991) 1615–1664.
- [Moc13] T. Mochizuki, "Harmonic bundles and Toda lattices with opposite sign," arXiv:1301.1718.
- [Moc15] T. Mochizuki, "Asymptotic behavior of certain families of harmonic bundles on Riemann surfaces," arXiv:1508.05997.
- [MSWW14] R. Mazzeo, J. Swoboda, H. Weiss, and F. Witt, "Limiting configurations for solutions of Hitchin's equation," *Semin. Theor. Spectr. Geom.* **31** (2012-2014) 91–116, arXiv:1502.01692.
- [MSWW16] R. Mazzeo, J. Swoboda, H. Weiss, and F. Witt, "Ends of the moduli space of Higgs bundles," *Duke Math. J.* 165 no. 12, (2016) 2227–2271, arXiv:1405.5765.
- [MSWW17] R. Mazzeo, J. Swoboda, H. Weiss, and F. Witt, "Asymptotic Geometry of the Hitchin Metric," arXiv:1709.03433.
- [MW15] R. Mazzeo and H. Weiss, "Teichmüller theory for conic surfaces," arXiv:1509.07608.
- [Sim90] C. T. Simpson, "Harmonic bundles on noncompact curves," *Journal of the America Mathematical Society* **3** no. 3, (1990) 713–770.