

ON CHARACTERISTICS OF HYPERFIELDS OBTAINED AS QUOTIENTS OF FINITE FIELDS

ANTONIO FRIGO, HAHN LHEEM, AND DYLAN LIU

PROPOSED BY: MATTHEW BAKER
COUNSELOR: ORON PROPP

ABSTRACT. Hyperstructures are a natural extension of regular algebraic structures in which one of the operations, known as the hyperoperation, is multivalued; a hyperfield is such an extension on a field. M. Krasner (1962) proved that the quotient \mathbb{F}_p/G , where G is a subgroup of units in \mathbb{F}_p is a hyperfield. The characteristic of a field may be explicitly determined from the order of the field, but there are no existing generalizations for determining the characteristic of a hyperfield of the form \mathbb{F}_p/G . We show that for odd primes p , there exists an explicit form for the characteristic of the hyperfield \mathbb{F}_p/G and $|G| = 1, 2, 3, 4$. Finally, we prove a general form of the characteristic for hyperfields where $|G|$ is prime.

1. INTRODUCTION

Hyperfields are a natural generalization of fields. Instead of the addition operation taking two elements and producing exactly one output, the \boxplus operation takes input of two sets and outputs a set in return. In our paper, we aim to study hyperfields constructed from $\mathbb{Z}/p\mathbb{Z}$ in the method described by Krasner [1]. We consider the motivating question:

Characterize the submonoids $S \subseteq \mathbb{N}$ for which S is the characteristic of a hyperfield.

The characteristic of a hyperfield is analogous to that of a field, except that the characteristic is a submonoid of \mathbb{N} rather than a natural number.

Our paper is structured as follows. We first provide the necessary definitions and prove basic propositions in §2. Next, we prove explicit forms of generating sets of type (p, n) -hyperfields for specific values of n , along with some computational data. We then prove a finite criterion for characteristics when n is prime in §4. Computational data for all (p, n) -hyperfields of primes p up to 200 can be found in Appendix A.

We will prove the following theorems in §3 and §4:

Theorem. *Let \mathbb{H} be an (p, n) -hyperfield with $|G| = 1$ and $p \geq 2$. Then $\text{gen}(\mathbb{H}) = \{p\}$.*

Theorem. *Let \mathbb{H} be an (p, n) -hyperfield with $|G| = 2$ and $p \geq 3$. Then $\text{gen}(\mathbb{H}) = \{2, p\}$.*

Theorem. *Let \mathbb{H} be an (p, n) -hyperfield with $|G| = 3$ and $p \geq 7$. Then there always exists positive integral solutions (a, b) to $a^2 - ab + b^2 = p$. Let m, n be the two possible values of $a + b$. Then the generating set is in the form $\{3, m, n\}$.*

Theorem. Let \mathbb{H} be an (p, n) -hyperfield with $|G| = 4$ and $p \geq 5$. Let (a, b) be the positive integral solution to $a^2 + b^2 = p$ and let $n = a + b$. Then the generating set is in the form $\{2, n\}$.

Theorem. Let q be a prime. Let ζ_q be the primitive q -th root of unity. The characteristic of the (p, q) -hyperfield is generated by the set

$$\{p, q\} \cup \left\{ \sum_{i=0}^{q-2} a_i \mid N \left(\sum_{i=0}^{q-2} a_i \zeta_q^i \right) \equiv 0 \pmod{p} \right\},$$

such that a_i are integers with $0 \leq a_i < p$.

1.1. Acknowledgements. The authors would especially like to thank Oron Propp for providing constant guidance and support throughout our research. We would also like to thank Dr. Matt Baker for proposing this project, as well as Dr. Henry Cohn and Dylan Pentland for their helpful insights. Finally, we thank the PROMYS program and the Clay Mathematics Institute for giving us this research opportunity.

2. DEFINITIONS, TERMINOLOGY, AND BASIC PROPERTIES

We now define some necessary concepts.

Definition 1. (*Hyperoperation*) Given two non-empty subsets $A, B \in S$, we define

$$A \boxplus B = \bigcup_{a \in A, b \in B} (a \boxplus b).$$

This operation is closed in S and is associative.

We can now construct analogs for groups, rings, and fields.

Definition 2. (*Canonical Hypergroup*) A hypergroup G has the hyperoperation \boxplus and a zero element. The hyperoperation must satisfy the following constraints:

- (*Additive Identity*) $0 \boxplus x = \{x\}$ for all $x \in G$.
- (*Hyperinverse*) Given $x \in G$, there exists unique $y \in G$ such that $0 \in x \boxplus y$.
- (*Reversibility*) $x \in y \boxplus z$ if and only if $z \in x \boxplus (-y)$

Since we consider only canonical hypergroups, we will refer to them as a hypergroup for conciseness.

Definition 3. (*Hyperring*) A hyperring R is a hypergroup with an additional operation \odot and element I that satisfy the following:

- The tuple (R, \odot, I) is a commutative monoid.
- The tuple $(R, \boxplus, 0)$ is a commutative hypergroup.
- (*Absorption*) For all $x \in R$, $x \odot 0 = 0$.
- (*Distributive*) $a \odot (x \boxplus y) = (a \odot x) \boxplus (a \odot y)$ for all $a, x, y, \in R$.

Proposition 1. (*Canonical*) The distributive law implies the reversibility axiom.

Proof. Without loss of generality, let $x \in y \boxplus z$. Note that $0 \in -x \boxplus x \subset -x \boxplus (y \boxplus z) = (-x \boxplus y) \boxplus z = -(x \boxplus -y) \boxplus z$, where the last equality follows because $-1 \odot (x \boxplus y) = (-1 \odot x) \boxplus (-1 \odot y)$ from distributivity. Because additive inverses are unique, this implies that $z \in x \boxplus -y$. An analogous argument proves the reverse direction. \square

Finally, we define a hyperfield, which is the focus of the paper.

Definition 4. (*Hyperfield*) A hyperfield is a hyperring F with $0 \neq 1$ and every non-zero element of F has a multiplicative inverse.

We extend a particular property of fields (its characteristic) to hyperfields.

Definition 5. (*Characteristic*) Let \mathbb{H} be a hyperfield. The characteristic of \mathbb{H} , which we will denote as $\text{char}(\mathbb{H})$, is the submonoid of \mathbb{N} consisting of all $n \in \mathbb{N}$ such that $0 \in nx$ for all $x \in F$, where nx is defined to be $\underbrace{x \boxplus x \boxplus \cdots \boxplus x}_{n \text{ } x\text{'s}}$.

Definition 6. (*Generating set*) We call the finite set $G = \{g_1, g_2, \dots, g_k\}$ of distinct natural numbers the generating set of a submonoid $S \subseteq \mathbb{N}$ if $S = \{g_1x_1 + g_2x_2 + \cdots + g_kx_k \mid g_i \in G, x_i \in \mathbb{N} \cup \{0\}\}$ and $|G|$ is minimal. For hyperfield (respectively, hyperring) \mathbb{H} , we denote the generating set of $\text{char}(\mathbb{H})$ as $\text{gen}(\mathbb{H})$.

Definition 7. (*Continuous*) We call a submonoid continuous at k if and only if every integer greater than or equal to k is contained in the submonoid.

We will mostly examine the structure of hyperfields generated in the following manner:

Proposition 2. (*Krasner*) For any arbitrary ring (resp. a field) R , let G be a subgroup of the group R^\times of units in R . Then the set of R/G units for the action of G on R by multiplication produces a hyperring (resp. a hyperfield) structure.

Proof. Proven by Krasner in [1]. \square

Remark. Note that when referring to G , we will use angle brackets to denote the set, namely $\langle \text{ and } \rangle$.

Definition 8. (*Type (p, n) -hyperfield*) We let a hyperfield of type (p, n) be of the form $(\mathbb{Z}/p\mathbb{Z})/G$ for positive prime p and subgroup $G \subseteq (\mathbb{Z}/p\mathbb{Z})^\times$. A hyperfield is the (p, n) -hyperfield if the subgroup G has size n .

Remark. This always yields a valid hyperfield by Proposition 2.

Definition 9. For hyperrings $H = R/G$ of type (p, n) , the operation \odot satisfies $[a] \odot [b] = [ab]$, where $a, b \in R$ and $[a]$ is the set generated by the multiplicative action of G on a , while the operation \boxplus is such that $[a] \boxplus [b]$ is the set of all $[c] \in H$ that are subsets of $\{x + y \mid x \in [a], y \in [b]\}$.

We now prove some useful propositions.

Proposition 3. To check whether $n \in \text{char}(F/G)$, it suffices to check only that $0 \in n[1]$

Proof. We see first that $0 \in n[1]$ is a necessary condition for $n \in \text{char}(F/G)$, as $1 \in F$. Now we show that it is sufficient: Let $a \in F$. Then, $a \odot (n[1]) = n[a]$ by distributivity. Since 0 is the absorption element, $0 \in n[a]$, we're done. Note that this proof extends to hyperrings as well. \square

Theorem 1. *If $G = \langle g_1, g_2, \dots, g_m \rangle$, then $\sum_{i=1}^m a_i g_i = 0$ in the ring implies that $\sum_{i=1}^m a_i \in \text{char}(F/G)$, where the $a_i \in \mathbb{Z}_{\geq 0}$. Additionally, $\sum_{i=1}^m a_i \in \text{char}(F/G)$ only if there exists an m -tuple (a_1, a_2, \dots, a_m) such that $\sum_{i=1}^m a_i g_i = 0$.*

Proof. It suffices to consider the case where G acts upon 1 by Proposition 3.

Since we are taking the setwise sum, we see that $0 \in n[1]$ if and only if we can add n of the elements in the group together to obtain zero. So, we add some combination of units together, with a_i instances of g_i in the sum, for a total of $n = \sum_{i=1}^m a_i$ summands. Since we assumed that adding the combination of g_i yields zero, $n = \sum_{i=1}^m a_i \in \text{char}(F/G)$. \square

3. GENERATING SETS OF (p, n) -HYPERFIELDS

3.1. Computations. Programs were written in Magma[2], C and Python to compute the generating sets of (p, n) -hyperfields. Table 1 contain the generating set for the (p, n) -hyperfield for some primes p and when $n = 1, 2, 3, 4$. Tables for all (p, n) -hyperfields for $p < 200$ is included in the Appendix, though data for all primes less than 1500 was calculated.

TABLE 1. $|G| = 1, 2, 3, 4$

$ G = 1$		$ G = 2$		$ G = 3$		$ G = 4$	
p	Generating Set	p	Generating Set	p	Generating Set	p	Generating Set
2	{2}	3	{2, 3}	7	{3, 4, 5}	5	{2, 3}
3	{3}	5	{2, 5}	13	{3, 5, 7}	13	{2, 5}
5	{5}	7	{2, 7}	19	{3, 7, 8}	17	{2, 5}
7	{7}	11	{2, 11}	31	{3, 7, 11}	29	{2, 7}
11	{11}	13	{2, 13}	37	{3, 10, 11}	37	{2, 7}
13	{13}	17	{2, 17}	43	{3, 8, 13}	41	{2, 9}
17	{17}	19	{2, 19}	61	{3, 13, 14}	53	{2, 9}
19	{19}	23	{2, 23}	67	{3, 11, 16}	61	{2, 11}
23	{23}	29	{2, 29}	73	{3, 10, 17}	73	{2, 11}
29	{29}	31	{2, 31}	79	{3, 13, 17}	89	{2, 13}
31	{31}	37	{2, 37}	97	{3, 14, 19}	97	{2, 13}
37	{37}	41	{2, 41}	103	{3, 13, 20}	101	{2, 11}
41	{41}	43	{2, 43}	109	{3, 17, 19}	109	{2, 13}
43	{43}	47	{2, 47}	127	{3, 19, 20}	113	{2, 15}
47	{47}	53	{2, 53}	139	{3, 16, 23}	137	{2, 15}
53	{53}	59	{2, 59}	151	{3, 19, 23}	149	{2, 17}

3.2. Some useful lemmas.

Lemma 2. *For any subgroup of units of $\mathbb{Z}/p\mathbb{Z}$, there is a generator of the subgroup. In particular, if the subgroup is size $|G|$, then a generator of the subgroup is $g^{\frac{p-1}{|G|}}$, where g is a generator of $(\mathbb{Z}/p\mathbb{Z})^\times$.*

Proof. Consider two elements in G , namely g^{da} and g^{db} where g is a generator in $\mathbb{Z}/p\mathbb{Z}$ and $(a, b) = 1$. Since there always exists a unique multiplicative inverse in the subgroup, by Bezout's famous lemma, g^d is also in the subgroup, which generates both g^{da} and g^{db} . We can continue this process inductively, ultimately showing that there is a generator for all elements in G . The second half of the statement follows immediately. \square

Lemma 3. *For any $\mathbb{Z}/p\mathbb{Z}$ with a subgroup $G = \langle g_0, g_1, \dots, g_m \rangle$ of $(\mathbb{Z}/p\mathbb{Z})^\times$, if $kp \geq \sum b_i g_i$ for non-negative integers b_i and a positive integer k , then $s \in S$ iff $s = kp - \sum b_i (g_i - 1)$.*

Proof. We first note that $(a_1, a_2, \dots, a_m) = (kp, 0, \dots, 0)$ for any arbitrary natural number k causes $\sum_{i=1}^m a_i g_i = kp = 0$, so $kp + 0 + \dots + 0 = kp \in S$.

We can replace any $(a_1, \dots, a_i, \dots, a_m)$ with $(a_1 - g_i, \dots, a_i + 1, \dots, a_m)$, which still yields a sum of $\sum_{i=1}^m a_i g_i = kp = 0$, but instead causes $kp - g_{x_1} + 1 - g_{x_2} + 1 - \dots - g_{x_n} + 1 \in S$ for some sequence of x_i , where $1 \leq x_i \leq m$, as long as $a_1 \geq 0$. This algorithm gives all possible solutions to $\sum_{i=1}^m a_i g_i = kp$, so the reverse direction holds. We can simplify this statement to the one stated in the lemma, where b_j represents the number of times some $x_i = j$. \square

Lemma 4. *For any $\mathbb{Z}/p\mathbb{Z}$ and a nontrivial subgroup of units G , $\text{char}((\mathbb{Z}/p\mathbb{Z})/G)$ is continuous at $p - 1$.*

Proof. Because of the nontriviality of the subgroup, there exists an element $g_i \in G$ such that $g_i - 1$ is a unit modulo p . Then, as b_i ranges from 0 to $p - 1$, $s = kp - \sum b_i (g_i - 1)$ hits every residue modulo p , and since $kp \geq \sum b_i g_i$, this implies that $s \geq p - 1 \geq b_i$, so any value greater than or equal to $p - 1$ is valid. \square

Lemma 5. *The equation $a^2 - ab + b^2$ has two distinct solutions $(a, b) \in \mathbb{N} \cup \{0\}$.*

Proof. To do this, we first prove that $a^2 - ab + b^2 = p$ has two distinct solutions up to order in the natural numbers. Consider the Eisenstein integer $a + b\omega \in \mathbb{Z}[\omega]$ and its norm, namely $N(a + b\omega) = a^2 - ab + b^2 = (a + b\omega)(a + b\omega^2)$. Because $N(a + b\omega)$ is a rational prime, $a + b\omega$ and $a + b\omega^2$ must be primes in $\mathbb{Z}[\omega]$.

Since $\mathbb{Z}[\omega]$ has unique prime factorization up to associates, we see that $a + b\omega$ and $a + b\omega^2$ and their associates are the only Eisenstein integers whose norms satisfy $a^2 - ab + b^2 = p$. If we assume without loss of generality $a > b$, then the only two solutions to $a^2 - ab + b^2 = p$ are (a, b) and $(a, a - b)$. \square

3.3. Explicit forms of generating sets. We have the following four theorems:

Theorem 6. *Let \mathbb{H} be an (p, n) -hyperfield with $|G| = 1$ and $p \geq 2$. Then $\text{gen}(\mathbb{H}) = \{p\}$.*

Proof. Using Theorem 1, we want all $n \in \mathbb{N}$ such that $n \times 1 = 0$ in $\mathbb{Z}/p\mathbb{Z}$, since $G = \{1\}$. This implies that $a = 0$, so a must be any positive multiple of p , leading to the generating set $\{p\}$. \square

Theorem 7. *Let \mathbb{H} be an (p, n) -hyperfield with $|G| = 2$ and $p \geq 3$. Then $\text{gen}(\mathbb{H}) = \{2, p\}$.*

Proof. Since $G = \langle -1, 1 \rangle$, it suffices by Theorem 1 to characterize $a + b$, where $a - b = 0$. We can then let $a = b + kp$, so $a + b = 2b + pk$. Since b and k can be any natural number, the generating set of the characteristic is $\{2, p\}$. \square

Theorem 8. *Let \mathbb{H} be an (p, n) -hyperfield with $|G| = 3$ and $p \geq 7$. Then there always exists positive integral solutions (a, b) to $a^2 - ab + b^2 = p$. Let m, n be the two possible values of $a + b$. Then the generating set is in the form $\{3, m, n\}$.*

Proof. Let $F = \mathbb{Z}/p\mathbb{Z}$, with prime $p \equiv 1 \pmod{3}$, and $G = \langle \omega, \omega^2, 1 \rangle$, where ω is a third root of unity. Since $1 \times \omega + 1 \times \omega^2 + 1 \times 1 = 0$, we see that $1 + 1 + 1 = 3$ is an element of $\text{gen}(F/G)$. To find the other generators of the generating set, it suffices to only consider the equation $a\omega + b = 0$.

Consider the equation $a_1\omega + a_2\omega^2 + a_3 = 0$. Let $m = \min(a_1, a_2, a_3)$. We know that since $\omega + \omega^2 + 1 = 0$, we have $m(\omega + \omega^2 + 1) = 0$. Hence

$$\begin{aligned} a_1\omega + a_2\omega^2 + a_3 - m(\omega + \omega^2 + 1) &= 0 \\ (a_1 - m)\omega + (a_2 - m)\omega^2 + (a_3 - m) &= 0. \end{aligned}$$

Since $m = \min(a_1, a_2, a_3)$, one of these terms is zero. We can then multiply by some power of ω to yield an equation in the form $a\omega + b = 0$.

Then, we have $a\omega \equiv -b \pmod{p}$, so cubing yields $a^3 \equiv -b^3 \pmod{p}$. So, p necessarily divides $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$ if we want $a + b \in \text{char}(F/G)$. However, we know if $p|(a + b)$, then $a + b \geq p$, so by Lemma 4, $a + b \in \text{char}(F/G)$ automatically. Hence we only need to consider $p|(a^2 - ab + b^2)$ for generators of $\text{char}(F/G)$.

In fact, we claim that the two values of $a + b$ such that $a^2 - ab + b^2 = p$, are not only in $\text{char}(F/G)$ but also the other two elements of the generating set of $\text{char}(F/G)$.

Without loss of generality, let $a > b$. Let $(a, b), (a, a - b)$ are the solutions to $x^2 - xy + y^2 = p$ (we know these are the only two by Lemma 5). It suffices to show that if all pairs of positive integers (c, d) satisfy $c^2 - cd + d^2 = kp$ for integer $k > 1$, then $\max(a + b, 2a - b) \leq c + d$.

We first show that $k \leq 3$. Suppose $k > 3$. Noting c, d are positive, we have

$$(c + d)^2 > c^2 - cd + d^2 = kp \implies c + d > \sqrt{kp}.$$

Furthermore,

$$a^2 - ab + b^2 = (a - b)^2 + ab = p \implies ab < p.$$

Thus,

$$a^2 - ab + b^2 = (a + b)^2 - 3ab = p \implies (a + b)^2 = p + 3ab < 4p \implies a + b < 2\sqrt{p}.$$

But for integer $k > 3$, we have

$$a + b < 2\sqrt{p} \leq \sqrt{kp} < c + d.$$

Hence if $k > 3$, then $\max(a + b, 2a - b) \leq c + d$.

Furthermore, $c^2 - cd + d^2 = kp = k(a^2 - ab + b^2)$, so since norms are multiplicative k must be the norm of some Eisenstein integer $f + g\omega$. We have that $k \neq 2$ since $f^2 - fg + g^2 = (f + g)^2 - 3fg \equiv 2 \pmod{3}$ has no solutions.

Now consider $k = 3$. By process of elimination from above, $c^2 - cd + d^2 = 3p$.

Let $x + y\omega \in \mathbb{Z}[\omega]$ and $N(x + y\omega) = 3$. From above, $c + d\omega = (x + y\omega)(a + b\omega)$ for positive integers a, b, c, d . Again, without loss of generality let $a > b$. (If $a < b$, then take the associate $-\omega(a + b\omega) = b + (b - a)\omega$.) As $N(1 + \omega) = 3$, $x + y\omega$ must be some associate of $1 + \omega$. Taking all products of $(a + b\omega)$ by an associate of $1 + \omega$ yield

$$(c, d) = (a + b, 2b - a), (a - 2b, 2a - b), (2a - b, b + a).$$

If $a > 2b$, then the minimum value of $c + d$ such that $c, d > 0$ is $(a - 2b) + (2a - b) = 3a - 3b$. However,

$$a > 2b \implies 2a > 4b \implies a + b < 3a - 3b = c + d.$$

If $a < 2b$, then $\min(c + d)$ becomes $(a + b) + (2b - a) = 3b$, and $a < 2b \implies a + b < 3b = c + d$.

Therefore, if $p \mid (x^2 - xy + y^2)$ for positive integers x, y , then $\min(x + y)$ is achieved when $x^2 - xy + y^2 = p$. By Lemma 5, there are exactly two solutions $(a_1, b_1), (a_2, b_2)$ to this equation, and hence $a_1 + b_1$ and $a_2 + b_2$ are generators of $\text{char}(F/G)$. \square

Theorem 9. *Let \mathbb{H} be an (p, n) -hyperfield with $|G| = 4$ and $p \geq 5$. Let (a, b) be the positive integral solution to $a^2 + b^2 = p$ and let $n = a + b$. Then the generating set of $\text{gen}(\text{char}(\mathbb{H}))$ is $\{2, n\}$.*

Proof. Let $F = \mathbb{Z}/p\mathbb{Z}$ for prime $p \equiv 1 \pmod{4}$. By Lemma 2, let g be the generator of the subgroup G . Then g is a fourth root of unity, so $G = \langle g, g^2, g^3, g^4 \rangle = \langle g, -1, -g, 1 \rangle$. If we wish to find the characteristic of \mathbb{H} , by Proposition 3, it suffices to find all positive integers n such that $0 \in n[1] = n\{g, -1, -g, 1\}$.

Because $1 \cdot 1 + 1 \cdot (-1) = 0$, $1 + 1 = 2$ must be a generator. Then, there can only be one other possible generator, which can be found by considering only $ag + b = 0$. To see why, note that we can take the equation $a_1g - a_2 - a_3g + a_4 = 0$ and subtract $0 = \min(a_1, a_3)(g - g) + \min(a_2, a_4)(-1 + 1)$ (thus $a_1 + a_2 + a_3 + a_4 - 2(\min(a_1, a_3) + \min(a_2, a_4)) \in \text{char}(F/G)$) and choose the more convenient value of g . Then, notice that $ag \equiv -b \pmod{p}$, so $-a^2 \equiv b^2 \pmod{p}$. Thus, p necessarily divides $a^2 + b^2$. In fact, we claim that the value of $a + b$ where $a^2 + b^2 = p$ is not only in $\text{char}(F/G)$ but also the other generator of $\text{char}(F/G)$.

Note that the maximum value of $a + b$ when $a^2 + b^2 = p$ occurs when $a = b$, so $\max(a + b) = \sqrt{2p}$. The minimum value of $a + b$ when $a^2 + b^2 = kp$, where $k > 1$, however, occurs when one of a or b is equal to zero, so $\min(a + b) = \sqrt{kp} \geq \sqrt{2p}$. So, the other generator is the smallest value of $a + b$ when $p \mid a^2 + b^2$, which occurs when $a^2 + b^2 = p$. It is well-known that the pair of natural numbers (a, b) is the unique solution up to ordering, so the value of n is also unique. \square

Conjecture 1. *For a given generating set $\{2, n\}$ where n is a positive odd integer, then there must exist a hyperfield \mathbb{H} such that $\text{gen}(\mathbb{H}) = \{2, n\}$.*

Remark. By Theorem 9, we have that $a^2 + b^2 = (a + bi)(a - bi) = p$. Note how $a^2 + b^2$ is the norm form in the Gaussian integers $\mathbb{Z}[i]$. Let $\pi \in \mathbb{Z}[i]$. It is a canonical fact that if $N(\pi)$ is prime, then π is a prime in the Gaussian integers. Since $a + b = n$, the conjecture reduces to showing that there always exists a Gaussian

prime that is n away from the line $y = x$ for all odd n . This is a very explicit statement regarding the distribution of the Gaussian primes, leading us to believe that this is a very difficult conjecture to prove.

We have verified this result computationally up to $n = 89441$.

4. FINITE CRITERION FOR GENERATING SET OF (p, n) -HYPERFIELD

Theorems 8 and 9 provide an explicit method for computing generators from solutions to bivariate polynomials. We aim to generalize these theorems and find, for every n such that $|G| = n$, a corresponding polynomial whose solutions can compute generators. We first give a finite criterion for when n is prime, before generalizing to all positive integers n .

We first define the norm of a cyclotomic field.

Definition 10. Let $f(\zeta_n) = f_0 + f_1\zeta_n + \cdots + f_k\zeta_n^k \in \mathbb{Z}[\zeta_n]$, where ζ_n is a primitive n -th root of unity. The norm of $f(\zeta_n)$ is the integer

$$N(f(\zeta_n)) = \prod_{i \in (\mathbb{Z}/n\mathbb{Z})^\times} f(\zeta_n^i).$$

In other words, the norm of $f \in \mathbb{Z}[\zeta_n]$ is the product of all Galois conjugates of f , including itself.

We are now prepared to construct a generating set of (p, q) -hyperfields for prime q .

Theorem 10. Let q be a prime. Let ζ_q be the primitive q -th root of unity. The characteristic of the (p, q) -hyperfield is generated by the set

$$\{p, q\} \cup \left\{ \sum_{i=0}^{q-2} a_i \mid N \left(\sum_{i=0}^{q-2} a_i \zeta_q^i \right) \equiv 0 \pmod{p} \right\},$$

such that a_i are integers with $0 \leq a_i < p$.

Proof. Let S be the set defined above and T be the minimal generating set. It suffices to show that T is a subset of S .

Let g be a generator in subgroup G . Every element in the characteristic must be of the form $a_0 + a_1g + \cdots + a_{q-1}g^{q-1}$, where a_i are non-negative integers and $a_0 + a_1g + \cdots + a_{q-1}g^{q-1} = 0$ in \mathbb{F}_p . However, we can reduce this equation. Let $\min(a_0, a_1, \dots, a_{q-1}) = a_i$. Then,

$$\begin{aligned} 0 &= g^{q-i-1} (a_0 + a_1g + \cdots + a_{q-1}g^{q-1}) \\ &= a_0g^{q-i-1} + \cdots + a_i g^{q-1} + \cdots + a_{q-1}g^{2q-i-2} \\ &= a_{i+1} + a_{i+2}g + \cdots + a_{q-1}g^{q-i-2} + a_0g^{q-i-1} + \cdots + a_i g^{q-1} \\ &= (a_{i+1} - a_i) + (a_{i+2} - a_i)g + \cdots + (a_{i-1} - a_i)g^{q-2}, \end{aligned}$$

with all $a_j - a_i$ being non-negative by minimality of a_i .

Note that this reduction yields all zero coefficients if either all the coefficients a_i were originally 1) equal, or 2) divisible by p . (In other words, all a_i are congruent modulo p .) If all coefficients were equal, then

$a_0 + a_1 + \cdots + a_{q-1} = q \cdot a_0$, so q is a generator. If all coefficients are $0 \pmod p$, then the sum of coefficients is some multiple of p , so p is also included in the generating set. Note, however, that because usually p is significantly large, p is not in the minimal generating set for any (p, n) -hyperfield. We thus assume that p is never in the minimal generating set, which we have verified computationally.

We claim that all other generators are produced from the sum of coefficients of some expression in terms of g with at most $q - 1$ terms.

Lemma 11. *Let $m \neq p, q$ be in the minimal generating set. Then*

$$m = b_0 + b_1 + \cdots + b_{q-2},$$

where b_i are elements in \mathbb{F}_p such that $b_0 + b_1g + \cdots + b_{q-2}g^{q-2} = 0$ in \mathbb{F}_p .

Proof. Suppose that $m = c_0 + \cdots + c_{q-1}$ such that $\gamma = c_0 + c_1g + \cdots + c_{q-1}g^{q-1} \equiv 0 \pmod p$, with c_i being integers. If one coefficient c_j is zero, then we can multiply γ by some power of g , shift indices, and reach the desired result. Furthermore, if all coefficients are equal, then $m = q$, contradiction. Thus, assume all c_i are positive and not all equal.

We first prove that $c_i \in \mathbb{F}_p$. Let $c_i \equiv d_i \pmod p$.

$$\begin{aligned} c_0 + c_1g + \cdots + c_{q-1}g^{q-1} &\equiv 0 \pmod p \\ \implies d_0 + d_1g + \cdots + d_{q-1}g^{q-1} &\equiv 0 \pmod p. \end{aligned}$$

Let $m' = d_0 + \cdots + d_{q-1}$. Because $c_i \equiv d_i \pmod p$, $m = m' + kp$ for some non-negative integer k . Thus, m can be generated by m' and p , so m cannot be in the minimal generating set. Thus, all coefficients c_i must be in \mathbb{F}_p .

Suppose again that $m = c_0 + \cdots + c_{q-1}$ such that $\gamma = c_0 + c_1g + \cdots + c_{q-1}g^{q-1} \equiv 0 \pmod p$, with $0 \leq c_i < p$. Assume that all c_i are positive and not all equal. Without loss of generality let $\min\{c_0, \dots, c_{q-1}\} = c_{q-1}$ (if not, then multiply γ by some power of g and shift indices such that the coefficient of g^{q-1} is minimal). We can use our reduction method shown earlier to get

$$\gamma = (c_0 - c_{q-1}) + (c_1 - c_{q-1})g + \cdots + (c_{q-2} - c_{q-1})g^{q-2} = 0.$$

Note that all coefficients of g^i are non-negative by minimality of c_{q-1} . Let $m'' = (c_0 - c_{q-1}) + \cdots + (c_{q-2} - c_{q-1})$. We see that

$$m'' + q \cdot c_{q-1} = \left(\sum_{i=0}^{q-2} c_i - c_{q-1} \right) + q \cdot c_{q-1} = \sum_{i=0}^{q-1} c_i = m,$$

so m can be generated by m'' and q . Therefore, m cannot be in the minimal generating set, and all elements in the minimal generating set are either q or expressed as the sum of at most $q - 1$ terms. \square

For notation, we let $M = T \setminus \{q\}$, or the set of all generators in the minimal generating set that are not equal to q . From our theorem, it then suffices to prove that

$$M \subseteq \left\{ \sum_{i=0}^{q-2} a_i \mid N \left(\sum_{i=0}^{q-2} a_i \zeta_q^i \right) \equiv 0 \pmod p \right\}$$

for $a_i \in \mathbb{F}_p$.

We now define a norm analogue for \mathbb{F}_p .

Definition 11. Let n, p be positive integers such that p is prime and $n|p-1$. Let $f(g) = f_0 + f_1g + \dots + f_kg^k \in \mathbb{F}_p$, where g is a primitive n -th root in \mathbb{F}_p . The p -norm of $f(g)$ is the integer

$$N'(f(g)) = \prod_{i \in (\mathbb{Z}/n\mathbb{Z})^\times} f(g^i).$$

Let

$$\begin{aligned} \alpha_1(\zeta_n) &= a_0 + a_1\zeta_n + \dots + a_k\zeta_n^k \in \mathbb{F}_p[\zeta_n] \\ \alpha_2(g) &= a_0 + a_1g + \dots + a_kg^k \in \mathbb{F}_p, \end{aligned}$$

where ζ_n is a primitive n -th root of unity and g is a primitive n -th root in \mathbb{F}_p . To compare $N(\alpha_1)$ and $N'(\alpha_2)$, we prove the following lemma:

Lemma 12. There exists a homomorphism $\rho : \mathbb{F}_p[\zeta_n] \rightarrow \mathbb{F}_p$ that maps ζ_n to g and every element in \mathbb{F}_p to itself.

Proof. To show that a homomorphism exists, it suffices to prove the following statement.

Lemma 13. Let ζ_n be a primitive n -th root of unity and g be a primitive n -th root in \mathbb{F}_p . Let V_1, V_2 be vector spaces over field \mathbb{F}_p with bases $\{1, \zeta_n, \dots, \zeta_n^{n-1}\}$ and $\{1, g, \dots, g^{n-1}\}$, respectively. Then there exists a linear map $f : V_1 \rightarrow V_2$ that maps ζ_n^k to g^k .

Proof. Let $\mathbf{v}_1, \mathbf{v}_2$ be vectors in V_1 , and c is in field \mathbb{F}_p . We will prove the following:

$$f(\mathbf{v}_1 + \mathbf{v}_2) = f(\mathbf{v}_1) + f(\mathbf{v}_2) \tag{1}$$

$$f(c\mathbf{v}_1) = cf(\mathbf{v}_1). \tag{2}$$

Let $\mathbf{v}_1 = a_0 + a_1\zeta_n + \dots + a_{n-1}\zeta_n^{n-1}$ and $\mathbf{v}_2 = b_0 + b_1\zeta_n + \dots + b_{n-1}\zeta_n^{n-1}$. The proof for (1) is straightforward:

$$\begin{aligned} f(\mathbf{v}_1 + \mathbf{v}_2) &= f((a_0 + b_0) + (a_1 + b_1)\zeta_n + \dots + (a_{n-1} + b_{n-1})\zeta_n^{n-1}) \\ &= (a_0 + b_0) + (a_1 + b_1)g + \dots + (a_{n-1} + b_{n-1})g^{n-1} \\ f(\mathbf{v}_1) + f(\mathbf{v}_2) &= (a_0 + a_1g + \dots + a_{n-1}g^{n-1}) + (b_0 + b_1g + \dots + b_{n-1}g^{n-1}) \\ &= (a_0 + b_0) + (a_1 + b_1)g + \dots + (a_{n-1} + b_{n-1})g^{n-1} \end{aligned}$$

Therefore $f(\mathbf{v}_1 + \mathbf{v}_2) = f(\mathbf{v}_1) + f(\mathbf{v}_2)$.

The proof for (2) is also straightforward:

$$\begin{aligned} f(c\mathbf{v}_1) &= f(ca_0 + ca_1\zeta_n + \dots + ca_{n-1}\zeta_n^{n-1}) \\ &= c(a_0 + a_1g + \dots + a_{n-1}g^{n-1}) \\ cf(\mathbf{v}_1) &= cf(a_0 + a_1\zeta_n + \dots + a_{n-1}\zeta_n^{n-1}) \\ &= c(a_0 + a_1g + \dots + a_{n-1}g^{n-1}), \end{aligned}$$

so $f(c\mathbf{v}_1) = cf(\mathbf{v}_1)$.

Thus, f is a linear map because we know how every vector in the basis of V_1 maps to V_2 and the two conditions for a linear map hold. \square

Because of the existence of a linear map by Lemma 13, there exists a homomorphism from $\mathbb{F}_p[\zeta_n]$ to \mathbb{F}_p that maps ζ_n to g . We can check that ρ preserves multiplication. Let

$$\begin{aligned}x &= x_0 + x_1\zeta_n + \cdots + x_{n-1}\zeta_n^{n-1} \\y &= y_0 + y_1\zeta_n + \cdots + y_{n-1}\zeta_n^{n-1},\end{aligned}$$

with $x, y \in \mathbb{F}_p[\zeta_n]$. Then

$$\begin{aligned}\rho(x \cdot y) &= \rho((x_0 + \cdots + x_{n-1}\zeta_n^{n-1})(y_0 + \cdots + y_{n-1}\zeta_n^{n-1})) \\&= \rho\left(\sum_{i=0}^{2n-2} \sum_{\substack{j+k=i \\ 0 \leq j, k < n}} x_j y_k \zeta_n^i\right) \\&= \sum_{i=0}^{2n-2} \sum_{\substack{j+k=i \\ 0 \leq j, k < n}} x_j y_k g^i \\&= (x_0 + \cdots + x_{n-1}g^{n-1})(y_0 + \cdots + y_{n-1}g^{n-1}) \\&= \rho(x) \cdot \rho(y),\end{aligned}$$

so $\rho(x \cdot y) = \rho(x) \cdot \rho(y)$ and ρ preserves multiplication. \square

Let $i \in (\mathbb{Z}/q\mathbb{Z})^\times$. By Lemma 12, if ρ is the homomorphism, then $\rho(\alpha_1(\zeta_q^i)) = \alpha_2(g^i)$. Taking the product over all units i in $\mathbb{Z}/q\mathbb{Z}$, $\rho(N(\alpha_1(\zeta_q))) = N'(\alpha_2(g))$. But $N(\alpha_1(\zeta_q))$ is an integer, so by definition of ρ in Lemma 12,

$$N(\alpha_1(\zeta_q)) \equiv N'(\alpha_2(g)) \pmod{p}. \quad (3)$$

Suppose $m = b_0 + b_1 + \cdots + b_{q-2}$ is in the generating set for $b_i \in \mathbb{F}_p$. Let $\beta = b_0 + b_1g + \cdots + b_{q-2}g^{q-2} \equiv 0 \pmod{p} \implies p|\beta$. By Definition 11,

$$\beta | N'(\beta).$$

Let $\beta(\zeta_q) = b_0 + b_1\zeta_q + \cdots + b_{q-2}\zeta_q^{q-2}$. By Equation 3,

$$\begin{aligned}N'(\beta) &\equiv N(\beta(\zeta_q)) \pmod{p} \\ \implies p|\beta | N'(\beta) &\equiv N(\beta(\zeta_q)) \pmod{p} \\ \implies N(\beta(\zeta_q)) &\equiv 0 \pmod{p}.\end{aligned}$$

This is true for all generators $m \in M$, so

$$M \subseteq \left\{ \sum_{i=0}^{q-2} a_i \mid N\left(\sum_{i=0}^{q-2} a_i \zeta_q^i\right) \equiv 0 \pmod{p} \right\}$$

for $a_i \in \mathbb{F}_p$. \square

Theorem 10 is significant in the following two ways:

- (1) For every hyperfield of type (p, q) , with q prime, there exists a homogeneous polynomial f of degree $q - 1$ whose solutions produce a generating set, and
- (2) The criterion is finite: there are at most p^{q-1} tuples (a_0, \dots, a_{q-2}) to test.

In particular, the homogeneous polynomial f whose solutions produce a generating set is the polynomial in $q - 1$ variables a_0, a_1, \dots, a_{q-2} that computes the norm of $a_0 + a_1\zeta_q + \dots + a_{q-2}\zeta_q^{q-2}$.

We can improve on the number of tuples to test in (2). First, we assume that every generator in the minimal generating set is less than p , so it suffices to check only the tuples whose elements sum to some positive value less than p . Furthermore, q is always a generator, so we may exclude all tuples whose elements sum to any multiples of q . By simple counting arguments, the total number of tuples we must test is at most

$$\binom{p+q-2}{q-1} - \sum_{k=0}^{\frac{p-q-1}{q}} \binom{q-2+kq}{q-2}.$$

REFERENCES

- [1] Marc Krasner, *A CLASS OF HYPERRINGS AND HYPERFIELDS* Internat. J. Math. & Math. Sci. Vol. 6 No. 2 (1983) 307-312
- [2] Wieb Bosma, John Cannon, and Catherine Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput., 24 (1997), 235–265.

APPENDIX A. COMPUTATIONAL DATA

p	$ G $	Gen. Set	p	$ G $	Gen. Set	p	$ G $	Gen. Set	p	$ G $	Gen. Set	p	$ G $	Gen. Set
2	1	2	43	42	2	89	2	2	127	42	2	173	4	2
3	1	3	43	21	3 5	89	11	4 6 7	127	2	2	173	2	2
3	2	2	43	14	2	89	22	2	131	1	131	173	86	2
5	1	5	47	1	47	89	44	2	131	65	3 5	173	172	2
5	2	2	47	23	3 5	89	8	2	131	2	2	173	43	3 5
5	4	2	47	2	2	89	4	2	131	10	2	179	1	179
7	1	7	47	46	2	97	4	2	131	26	2	179	178	2
7	3	3 5	53	1	53	97	12	2	131	13	4 6 7	179	2	2
7	6	2	53	52	2	97	48	2	131	5	5 9 11 12	179	89	3 5
7	2	2	53	13	3 5	97	16	2	131	130	2	181	60	2
11	1	11	53	2	2	97	6	2	137	136	2	181	180	2
11	5	3 5	53	4	2	97	96	2	137	4	2	181	36	2
11	10	2	53	26	2	97	2	2	137	1	137	181	12	2
11	2	2	59	1	59	97	1	97	137	34	2	181	18	2
13	1	13	59	2	2	97	3	3 19	137	17	4 6 7	181	30	2
13	12	2	59	58	2	97	24	2	137	2	2	181	10	2
13	2	2	59	29	3 5	97	8	2	137	8	2	181	20	2
13	3	3 7	61	30	2	97	32	2	137	68	2	181	1	181
13	4	2	61	15	3 5	101	1	101	139	1	139	181	9	3 8
13	6	2	61	3	3 14	101	50	2	139	23	4 6 7	181	6	2
17	1	17	61	5	4 7	101	25	3 5	139	46	2	181	3	3 26
17	2	2	61	1	61	101	2	2	139	3	3 23	181	45	3 5
17	16	2	61	10	2	101	20	2	139	69	3 5	181	15	3 7
17	4	2	61	60	2	101	4	2	139	138	2	181	5	5 7
17	8	2	61	6	2	101	100	2	139	2	2	181	4	2
19	1	19	61	20	2	101	10	2	139	6	2	181	90	2
19	18	2	61	12	2	101	5	5 8 9 11	149	1	149	181	2	2
19	3	3 8	61	2	2	103	1	103	149	2	2	191	38	2
19	2	2	61	4	2	103	102	2	149	4	2	191	5	5 8 9 11
19	6	2	67	1	67	103	3	3 20	149	148	2	191	19	3 5
19	9	3 5	67	33	3 5	103	6	2	149	37	3 5	191	1	191
23	1	23	67	6	2	103	17	3 5	149	74	2	191	10	2
23	2	2	67	3	3 16	103	34	2	151	150	2	191	190	2
23	22	2	67	11	4 6 7	103	2	2	151	10	2	191	95	3 5
23	11	3 5	67	2	2	103	51	3 5	151	3	3 23	191	2	2
29	1	29	67	66	2	107	1	107	151	5	5 9 11 13	193	6	2
29	7	4 6 7	67	22	2	107	106	2	151	75	3 5	193	3	3 25
29	2	2	71	1	71	107	2	2	151	25	3 5	193	2	2
29	4	2	71	10	2	107	53	3 5	151	1	151	193	192	2
29	28	2	71	70	2	109	27	3 5	151	50	2	193	8	2
29	14	2	71	2	2	109	18	2	151	30	2	193	96	2
31	1	31	71	35	3 5	109	54	2	151	15	3 5	193	64	2
31	5	5 7 8 9	71	7	4 6 7	109	4	2	151	6	2	193	1	193
31	10	2	71	14	2	109	1	109	151	2	2	193	32	2
31	3	3 11	71	5	5 7 8 9	109	12	2	157	6	2	193	48	2
31	6	2	73	3	3 17	109	2	2	157	39	3 5	193	16	2
31	2	2	73	4	2	109	6	2	157	2	2	193	12	2
31	30	2	73	18	2	109	36	2	157	52	2	193	4	2
31	15	3 5	73	6	2	109	9	3 7	157	78	2	193	24	2
37	1	37	73	36	2	109	108	2	157	1	157	197	4	2
37	36	2	73	24	2	109	3	3 19	157	13	4 6 7	197	7	5 7 8 9
37	12	2	73	1	73	113	8	2	157	26	2	197	49	3 5
37	2	2	73	72	2	113	28	2	157	3	3 25	197	28	2
37	6	2	73	8	2	113	4	2	157	12	2	197	1	197
37	9	3 5	73	2	2	113	56	2	157	156	2	197	2	2
37	3	3 11	73	12	2	113	1	113	157	4	2	197	196	2
37	18	2	73	9	3 5	113	2	2	163	9	3 7	197	14	2
37	4	2	79	1	79	113	7	5 7 8 9	163	2	2	197	98	2
41	5	5 7 8 9	79	2	2	113	16	2	163	162	2	199	9	3 7
41	10	2	79	39	3 5	113	112	2	163	18	2	199	2	2
41	40	2	79	13	4 6 7	113	14	2	163	1	163	199	99	3 5
41	1	41	79	6	2	127	6	2	163	27	3 5	199	11	6 8 9 10 11
41	8	2	79	26	2	127	21	3 5	163	81	3 5	199	6	2
41	2	2	79	78	2	127	1	127	163	6	2	199	1	199
41	20	2	79	3	3 17	127	126	2	163	54	2	199	33	3 5
41	4	2	83	1	83	127	9	3	163	3	3 25	199	198	2

p	$ G $	Gen. Set	p	$ G $	Gen. Set	p	$ G $	Gen. Set	p	$ G $	Gen. Set	p	$ G $	Gen. Set
43	1	43	83	82	2	127	63	3 5	167	1	167	199	66	2
43	3	3 13	83	2	2	127	3	3 20	167	83	3 5	199	3	3 28
43	6	2	83	41	3 5	127	7	7 9 10 11 12 13	167	2	2	199	18	2
43	7	3 7	89	1	89	127	14	2	167	166	2	199	22	2
43	2	2	89	88	2	127	18	2	173	1	173			