# FINITE GROUPS WITH A PRESCRIBED NUMBER OF CYCLIC SUBGROUPS II

RICHARD BELSHOFF, JOE DILLSTROM, AND LES REID

ABSTRACT. In [4], Tărnăuceanu described the finite groups G having exactly |G| - 1 cyclic subgroups. In [1], the authors used elementary methods to completely characterize those finite groups Ghaving exactly  $|G| - \Delta$  cyclic subgroups for  $\Delta = 2, 3, 4$  and 5. In this paper, we prove that for any  $\Delta > 0$  if G has exactly  $|G| - \Delta$ cyclic subgroups, then  $|G| \leq 8\Delta$  and therefore the number of such G is finite. We then use the computer program GAP to find all Gwith exactly  $|G| - \Delta$  cyclic subgroups for  $\Delta = 1, \ldots, 32$ .

## 1. INTRODUCTION

Throughout this paper, a cyclic group of order n is denoted  $C_n$ and the dihedral group of order 2n is denoted  $D_{2n}$ . In [4], Marius Tărnăuceanu proved the following theorem.

**Theorem 1** (Tărnăuceanu). A finite group G has exactly |G|-1 cyclic subgroups if and only if G is one of the following groups:  $C_3$ ,  $C_4$ ,  $S_3$ , or  $D_8$ .

Let G be a finite group, and let C(G) denote the poset of cyclic subgroups of G. We define  $\Delta(G)$ , or just  $\Delta$  if the group is understood, to be the difference between the order of G and the number of cyclic subgroups of G:

$$\Delta(G) = |G| - |C(G)|.$$

It is known that  $\Delta = 0$  if and only if G is an elementary abelian 2-group, i.e.,  $G = C_2^n$  for some n.

In [4] where Tărnăuceanu described all groups satisfying  $\Delta = 1$ , he also posed the following natural open problem.

**Problem.** Describe the finite groups G having  $|G| - \Delta$  cyclic subgroups, where  $2 \leq \Delta \leq |G| - 1$ .

In [1] we determined those finite groups for which  $\Delta = 2, 3, 4$ , and 5. In this paper we will show that  $|G| \leq 8\Delta(G)$  and use this bound and a GAP program to find all G such that  $\Delta(G) = 1, \ldots 32$ .

Date: October 22, 2018.

# 2. Main Result

Let G be a finite group. Denote by C(G) the poset of cyclic subgroups of G, and let  $\Delta(G) = |G| - |C(G)|$ . For any positive integer d, let  $n_d = |\{H \in C(G) : |H| = d\}|$  be the number of cyclic subgroups of order d. Since every element of G generates a cyclic subgroup and  $\phi(d)$  is the number of generators of a cyclic group of order d, it follows that

$$\sum_{d\geq 1} n_d \phi(d) = |G|.$$

Because  $|C(G)| = \sum_{d \ge 1} n_d$ , we have  $\sum n_d(\phi(d) - 1)$ 

$$\sum_{d \ge 1} n_d(\phi(d) - 1) = \Delta(G). \tag{(\star)}$$

Since  $\phi(d) \ge 2$  for d > 2, we have

$$\Delta(G) = \sum_{d \ge 1} n_d(\phi(d) - 1) \ge \sum_{d > 2} n_d = |C(G)| - i_2(G), \quad (1)$$

where  $i_2(G)$  denotes the number of elements of order 2 or less.

Now by [3], if  $G \not\cong C_2^k$ , then  $i_2(G) \leq \frac{3}{4}|G|$  with equality if and only if  $G \cong D_8 \times C_2^k$  for some  $k \geq 0$ . If  $\Delta(G) > 0$ , then  $G \not\cong C_2^k$  so we have

$$\begin{aligned} |G| &= \Delta(G) + |C(G)| \\ &\leq \Delta(G) + \Delta(G) + i_2(G) \\ &= 2\Delta(G) + i_2(G) \\ &\leq 2\Delta(G) + \frac{3}{4}|G|. \end{aligned}$$

Solving the inequality for |G| yields the following result.

**Theorem 2.** For any finite group G, if  $\delta > 0$  and  $\Delta(G) = \delta$ , then  $|G| \leq 8\delta$ . Equality only occurs when  $\delta = 2^k$  and in that case  $G \cong D_8 \times C_2^k$ .

**Corollary 1.** If  $\delta > 0$ , the number of groups such that  $\Delta(G) = \delta$  is finite.

This gives an alternate proof of Tărnăuceanu's result. If  $\Delta(G) = 1$ , then  $|G| \leq 8$  and one only need check the nine groups (up to isomorphism) of order less than 8  $[C_1, C_2, C_3, C_4, C_2 \times C_2, C_5, C_6, D_6, C_7]$  (if |G| = 8 we must have  $G \cong D_8$ ). In the remainder of this paper, we will report on the results of a GAP program that finds all groups of order less than  $8\delta$  such that  $\Delta(G) = \delta$  for  $\delta = 1, \ldots, 32$ .

# 3. Computer Results

Here we list all groups with difference  $\delta$ , for  $\delta = 1$  to 32. The GAP notation is as follows:

A x B denotes a direct product of a group A by a group B,

A:B denotes a split extension of A by B,

A.B denotes just an extension of A by B (split or nonsplit). For each group, we list both GAP's StructureDescription and GroupID, in the form

## StructureDescription = GroupID

For example, the line below that reads

D8 = [8,3]

means that the dihedral group of order 8 is the third group of order 8 in GAP's SmallGroups library.

## Four groups with difference 1 (Tărnăuceanu [4]).

```
C3 = [ 3, 1 ]
C4 = [ 4, 1 ]
S3 = [ 6, 1 ]
D8 = [ 8, 3 ]
```

```
Four groups with difference 2 (Tărnăuceanu [5], and BDR [1]).
```

C6 = [ 6, 2 ] C4 x C2 = [ 8, 2 ] D12 = [ 12, 4 ] C2 x D8 = [ 16, 11 ]

Three groups with difference 3 (BDR [1]).

C5 = [ 5, 1 ] Q8 = [ 8, 4 ] D10 = [ 10, 1 ]

Eleven groups with difference 4 (BDR [1]).

C8 = [ 8, 1 ] C3 x C3 = [ 9, 2 ] A4 = [ 12, 3 ] C6 x C2 = [ 12, 5 ] (C4 x C2) : C2 = [ 16, 3 ] D16 = [ 16, 7 ] C4 x C2 x C2 = [ 16, 10 ] (C4 x C2) : C2 = [ 16, 13 ] (C3 x C3) : C2 = [ 18, 4 ] C2 x C2 x S3 = [ 24, 14 ] C2 x C2 x D8 = [ 32, 46 ]

Three groups with difference 5 (BDR [1]).

C7 = [ 7, 1 ] C3 : C4 = [ 12, 1 ] D14 = [ 14, 1 ]

# Thirteen groups with difference 6.

C9 = [9, 1] C10 = [10, 2] C12 = [12, 2]  $C4 \times C4 = [16, 2]$  C4 : C4 = [16, 4] QD16 = [16, 8]  $C2 \times Q8 = [16, 12]$  D18 = [18, 1] D20 = [20, 4] D24 = [24, 6]  $(C2 \times C2 \times C2 \times C2) : C2 = [32, 27]$   $(C4 \times C4) : C2 = [32, 34]$   $(C2 \times D8) : C2 = [32, 49]$ 

# Three groups with difference 7.

C3 x S3 = [ 18, 3 ] (C6 x C2) : C2 = [ 24, 8 ] S4 = [ 24, 12 ]

# Fifteen groups with difference 8.

C8 x C2 = [ 16, 5 ] C8 : C2 = [ 16, 6 ] Q16 = [ 16, 9 ] C6 x C3 = [ 18, 5 ] C5 : C4 = [ 20, 3 ] C2 x A4 = [ 24, 13 ] C6 x C2 x C2 = [ 24, 15 ] C2 x ((C4 x C2) : C2) = [ 32, 22 ] (C4 x C2 x C2) : C2 = [ 32, 28 ] C2 x D16 = [ 32, 39 ] C4 x C2 x C2 x C2 = [ 32, 45 ]

C2 x ((C4 x C2) : C2) = [ 32, 48 ] C2 x ((C3 x C3) : C2) = [ 36, 13 ] C2 x C2 x C2 x S3 = [ 48, 51 ] C2 x C2 x C2 x D8 = [ 64, 261 ]

# Three groups with difference 9.

C11 = [ 11, 1 ] D22 = [ 22, 1 ] C4 x S3 = [ 24, 5 ]

# Eleven groups with difference 10.

```
C14 = [ 14, 2 ]

C2 x (C3 : C4) = [ 24, 7 ]

C3 x D8 = [ 24, 10 ]

D28 = [ 28, 3 ]

((C4 x C2) : C2) : C2 = [ 32, 6 ]

C4 x D8 = [ 32, 25 ]

(C4 x C2 x C2) : C2 = [ 32, 30 ]

(C4 x C4) : C2 = [ 32, 31 ]

(C2 x D8) : C2 = [ 32, 43 ]

(C2 x Q8) : C2 = [ 32, 50 ]

S3 x S3 = [ 36, 10 ]
```

## Eight groups with difference 11.

C13 = [ 13, 1 ] C15 = [ 15, 1 ] C16 = [ 16, 1 ] C5 : C4 = [ 20, 1 ] SL(2,3) = [ 24, 3 ] D26 = [ 26, 1 ] D30 = [ 30, 3 ] D32 = [ 32, 18 ]

Twenty-one groups with difference 12.

C18 = [ 18, 2 ] C10 x C2 = [ 20, 5 ] C7 : C3 = [ 21, 1 ] C3 : Q8 = [ 24, 4 ] C12 x C2 = [ 24, 9 ] (C4 x C2) : C4 = [ 32, 2 ] (C8 x C2) : C2 = [ 32, 9 ] C4 x C4 x C2 = [ 32, 21 ] C2 x (C4 : C4) = [ 32, 23 ]

```
(C4 x C4) : C2 = [ 32, 24 ]
(C2 x Q8) : C2 = [ 32, 29 ]
(C4 x C4) : C2 = [ 32, 33 ]
C2 x QD16 = [ 32, 40 ]
(C8 x C2) : C2 = [ 32, 42 ]
C2 x C2 x Q8 = [ 32, 47 ]
D36 = [ 36, 4 ]
C2 x C2 x D10 = [ 40, 13 ]
C2 x D24 = [ 48, 36 ]
C2 x ((C2 x C2 x C2 x C2) : C2) = [ 64, 202 ]
C2 x ((C4 x C4) : C2) = [ 64, 211 ]
C2 x ((C2 x D8) : C2) = [ 64, 264 ]
```

# Five groups with difference 13.

(C3 x C3) : C3 = [ 27, 3 ] C3 x C3 x C3 = [ 27, 5 ] (C3 x C3) : C4 = [ 36, 9 ] D8 x S3 = [ 48, 38 ] (C3 x C3 x C3) : C2 = [ 54, 14 ]

# Thirteen groups with difference 14.

C20 = [20, 2]  $C3 \times Q8 = [24, 11]$  (C8 : C2) : C2 = [32, 7]  $(C4 \times C4) : C2 = [32, 11]$   $C4 \times Q8 = [32, 26]$   $(C2 \times C2) \cdot (C2 \times C2 \times C2) = [32, 32]$  C4 : Q8 = [32, 35]  $(C2 \times Q8) : C2 = [32, 44]$   $C6 \times S3 = [36, 12]$  D40 = [40, 6]  $C2 \times ((C6 \times C2) : C2) = [48, 43]$   $C2 \times S4 = [48, 48]$   $D8 \times D8 = [64, 226]$ 

# Four groups with difference 15.

```
C17 = [ 17, 1 ]
C3 : C8 = [ 24, 1 ]
QD32 = [ 32, 19 ]
D34 = [ 34, 1 ]
```

```
Thirty groups with difference 16.
```

```
C24 = [24, 2]
C3 \times D10 = [30, 2]
(C8 \times C2) : C2 = [32, 5]
Q8 : C4 = [32, 10]
C8 : C4 = [32, 13]
C8 : C4 = [32, 14]
C8 \times C2 \times C2 = [32, 36]
C2 \times (C8 : C2) = [32, 37]
(C8 \times C2) : C2 = [32, 38]
C2 \times Q16 = [32, 41]
C3 \times A4 = [36, 11]
C6 \times C6 = [36, 14]
C2 \times (C5 : C4) = [40, 12]
D48 = [48, 7]
C2 \times C2 \times A4 = [48, 49]
(C2 \times C2 \times C2 \times C2) : C3 = [48, 50]
C6 \times C2 \times C2 \times C2 = [48, 52]
(C2 \times ((C4 \times C2) : C2)) : C2 = [64, 60]
(C2 \times C2 \times D8) : C2 = [64, 73]
C2 \times C2 \times ((C4 \times C2) : C2) = [64, 193]
C2 \times ((C4 \times C2 \times C2) : C2) = [64, 203]
(C2 \times C2 \times D8) : C2 = [64, 215]
(C2 \times C2 \times D8) : C2 = [64, 216]
C2 \times C2 \times D16 = [64, 250]
C4 \times C2 \times C2 \times C2 \times C2 = [64, 260]
C2 \times C2 \times ((C4 \times C2) : C2) = [64, 263]
(C2 \times ((C4 \times C2) : C2)) : C2 = [64, 266]
C2 \times C2 \times ((C3 \times C3) : C2) = [72, 49]
C2 \times C2 \times C2 \times C2 \times C3 = [96, 230]
C2 \times C2 \times C2 \times C2 \times D8 = [128, 2320]
```

# Eight groups with difference 17.

C19 = [ 19, 1 ] C21 = [ 21, 2 ] C7 : C4 = [ 28, 1 ] (C3 x C3) : C4 = [ 36, 7 ] D38 = [ 38, 1 ] (C10 x C2) : C2 = [ 40, 8 ] D42 = [ 42, 5 ] (C4 x S3) : C2 = [ 48, 41 ]

#### Seventeen groups with difference 18.

```
C22 = [22, 2]
C5 \times C5 = [25, 2]
C8 \times C4 = [32, 3]
C8 : C4 = [32, 4]
C2 . ((C4 x C2) : C2) = (C2 x C2) . (C4 x C2) = [ 32, 8 ]
C4 : C8 = [32, 12]
D44 = [44, 3]
(C12 \times C2) : C2 = [48, 14]
C2 \times C4 \times S3 = [48, 35]
(C12 \times C2) : C2 = [48, 37]
(C5 \times C5) : C2 = [50, 4]
(((C4 \times C2) : C2) : C2) : C2 = [64, 138]
(C8 \times C4) : C2 = [64, 174]
(C2 \times C2 \times D8) : C2 = [64, 227]
(C4 \times D8) : C2 = [64, 231]
((C4 \times C2 \times C2) : C2) : C2 = [64, 241]
((C4 \times C4) : C2) : C2 = [64, 242]
```

# Eight groups with difference 19.

C9 x C3 = [ 27, 2 ] C9 : C3 = [ 27, 4 ] Q32 = [ 32, 20 ] C4 x D10 = [ 40, 5 ] (C7 : C3) : C2 = [ 42, 1 ] (C3 x D8) : C2 = [ 48, 15 ] (C2 x (C3 : C4)) : C2 = [ 48, 39 ] (C9 x C3) : C2 = [ 54, 7 ]

# Twenty-eight groups with difference 20.

```
C14 x C2 = [ 28, 4 ]

C5 x S3 = [ 30, 1 ]

C4 . D8 = C4 . (C4 x C2) = [ 32, 15 ]

C3 x (C3 : C4) = [ 36, 6 ]

(C2 x (C3 : C4)) : C2 = [ 48, 19 ]

GL(2,3) = [ 48, 29 ]

A4 : C4 = [ 48, 30 ]

C2 x C2 x (C3 : C4) = [ 48, 42 ]

C6 x D8 = [ 48, 45 ]

C2 x C2 x D14 = [ 56, 12 ]

(C4 x C2 x C2 x C2) : C2 = [ 64, 67 ]

(C4 x C4 x C2) : C2 = [ 64, 71 ]
```

```
(C2 \times ((C4 \times C2) : C2)) : C2 = [64, 75]
C2 \times (((C4 \times C2) : C2) : C2) = [64, 90]
(C2 \times C2 \times D8) : C2 = [64, 128]
C2 \times C4 \times D8 = [64, 196]
(C4 \times D8) : C2 = [64, 199]
C2 \times ((C4 \times C2 \times C2) : C2) = [64, 205]
(C4 \times C2 \times C2 \times C2) : C2 = [64, 206]
C2 \times ((C4 \times C4) : C2) = [64, 207]
(C4 \times C4 \times C2) : C2 = [64, 213]
(C2 \times ((C4 \times C2) : C2)) : C2 = [64, 218]
(C4 \times D8) : C2 = [64, 219]
(C4 \times D8) : C2 = [64, 221]
C2 \times ((C2 \times D8) : C2) = [64, 254]
(C2 \times D16) : C2 = [64, 257]
C2 \times ((C2 \times Q8) : C2) = [64, 265]
C2 \times S3 \times S3 = [72, 46]
```

## Five groups with difference 21.

C23 = [ 23, 1 ] C9 : C4 = [ 36, 1 ] C12 x C3 = [ 36, 8 ] D46 = [ 46, 1 ] (C12 x C3) : C2 = [ 72, 33 ]

## Thirty groups with difference 22.

```
C25 = [25, 1]
C26 = [26, 2]
C28 = [28, 2]
C30 = [30, 4]
C16 \times C2 = [32, 16]
C16 : C2 = [32, 17]
C2 \times (C5 : C4) = [40, 7]
(C4 \times C4) : C3 = [48, 3]
C24 : C2 = [48, 6]
C2 \times SL(2,3) = [48, 32]
D50 = [50, 1]
D52 = [52, 4]
((C3 \times C3) : C3) : C2 = [54, 5]
((C3 \times C3) : C3) : C2 = [54, 8]
C3 \times ((C3 \times C3) : C2) = [54, 13]
D56 = [56, 5]
D60 = [60, 12]
```

```
(((C4 x C2) : C2) : C2) : C2 = [ 64, 34 ]
((C4 x C4) : C2) : C2 = [ 64, 134 ]
(((C4 x C2) : C2) : C2) : C2 = [ 64, 139 ]
(C4 x D8) : C2 = [ 64, 140 ]
(C2 x D16) : C2 = [ 64, 177 ]
C2 x D32 = [ 64, 186 ]
(C4 x D8) : C2 = [ 64, 228 ]
(C2 x C2 x Q8) : C2 = [ 64, 229 ]
(C4 x D8) : C2 = [ 64, 232 ]
(C4 x D8) : C2 = [ 64, 234 ]
(C4 x D8) : C2 = [ 64, 236 ]
(C4 x D8) : C2 = [ 64, 240 ]
((C2 x C2) . (C2 x C2 x C2)) : C2 = [ 64, 243 ]
```

## Four groups with difference 23.

C27 = [ 27, 1 ] (C3 x Q8) : C2 = [ 48, 17 ] Q8 x S3 = [ 48, 40 ] D54 = [ 54, 1 ]

#### Fifty-nine groups with difference 24.

```
(C2 \times C2) : C9 = [36, 3]
C18 \times C2 = [36, 5]
C13 : C3 = [39, 1]
C5 : Q8 = [40, 4]
C10 \times C2 \times C2 = [40, 14]
C2 \times (C7 : C3) = [42, 2]
C3 \times D14 = [42, 4]
C4 \times (C3 : C4) = [48, 11]
(C3 : C4) : C4 = [48, 12]
C12 : C4 = [48, 13]
C3 \times ((C4 \times C2) : C2) = [48, 21]
C3 \times D16 = [48, 25]
C4 \times A4 = [48, 31]
SL(2,3) : C2 = [48, 33]
C2 \times (C3 : Q8) = [48, 34]
C12 \times C2 \times C2 = [48, 44]
C3 \times ((C4 \times C2) : C2) = [48, 47]
C13 : C4 = [52, 3]
(C4 \times C2 \times C2) : C4 = [64, 23]
C2 \times ((C4 \times C2) : C4) = [64, 56]
C4 \times ((C4 \times C2) : C2) = [64, 58]
```

```
(C2 \times (C4 : C4)) : C2 = [64, 61]
((C4 \times C2) : C4) : C2 = [64, 62]
(C2 \times (C4 : C4)) : C2 = [64, 66]
(C4 \times C4 \times C2) : C2 = [64, 69]
(C2 \times C2 \times Q8) : C2 = [64, 74]
(C2 \times (C4 : C4)) : C2 = [64, 77]
(C2 \times (C4 : C4)) : C2 = [64, 78]
(C2 \times (C4 : C4)) : C2 = [64, 80]
(((C4 \times C2) : C2) : C2) : C2 = [64, 91]
C2 \times ((C8 \times C2) : C2) = [64, 95]
(C2 \times (C8 : C2)) : C2 = [64, 99]
(C2 \times D16) : C2 = [64, 130]
(C2 \times QD16) : C2 = [64, 131]
(C8 \times C2 \times C2) : C2 = [64, 147]
(C2 \times (C8 : C2)) : C2 = [64, 150]
C4 \times C4 \times C2 \times C2 = [64, 192]
C2 \times C2 \times (C4 : C4) = [64, 194]
C2 \times ((C4 \times C4) : C2) = [64, 195]
C4 \times ((C4 \times C2) : C2) = [64, 198]
(C4 \times Q8) : C2 = [64, 201]
C2 \times ((C2 \times Q8) : C2) = [64, 204]
C2 \times ((C4 \times C4) : C2) = [64, 209]
(C4 \times C4 \times C2) : C2 = [64, 210]
(C2 \times C2 \times Q8) : C2 = [64, 217]
(C4 \times D8) : C2 = [64, 220]
(C4 \times Q8) : C2 = [64, 223]
((C2 \times Q8) : C2) : C2 = [64, 224]
C2 \times C2 \times QD16 = [64, 251]
C2 \times ((C8 \times C2) : C2) = [64, 253]
(C2 \times (C8 : C2)) : C2 = [64, 256]
(C2 \times QD16) : C2 = [64, 258]
C2 \times C2 \times C2 \times Q8 = [64, 262]
C2 \times C2 \times D18 = [72, 17]
C2 \times C2 \times C2 \times D10 = [80, 51]
C2 \times C2 \times D24 = [96, 207]
C2 \times C2 \times ((C2 \times C2 \times C2 \times C2) : C2) = [128, 2163]
C2 \times C2 \times ((C4 \times C4) : C2) = [128, 2172]
C2 \times C2 \times ((C2 \times D8) : C2) = [128, 2323]
```

# Six groups with difference 25.

(C3 : C8) : C2 = [ 48, 16 ] C3 x C3 x S3 = [ 54, 12 ] S3 x D10 = [ 60, 8 ] (C6 x C6) : C2 = [ 72, 35 ] (S3 x S3) : C2 = [ 72, 40 ] (C3 x A4) : C2 = [ 72, 43 ]

# Twenty-nine groups with difference 26.

```
C32 = [32, 1]
C5 : C8 = [40, 3]
C5 \times D8 = [40, 10]
C2 . S4 = SL(2,3) . C2 = [48, 28]
C2 \times ((C3 \times C3) : C3) = [54, 10]
C6 \times C3 \times C3 = [54, 15]
((C8 : C2) : C2) : C2 = [64, 32]
(C4 \times C4) : C4 = [64, 35]
D64 = [64, 52]
C4 \times D16 = [64, 118]
(C4 \times D8) : C2 = [64, 123]
((C4 \times C4) : C2) : C2 = [64, 135]
((C4 \times C4) : C2) : C2 = [64, 136]
(C2 \times QD16) : C2 = [64, 141]
(C4 \times D8) : C2 = [64, 144]
(C8 \times C4) : C2 = [64, 167]
((C8 \times C2) : C2) : C2 = [64, 171]
(C8 \times C4) : C2 = [64, 173]
(C8 \times C4) : C2 = [64, 176]
(C2 \times D16) : C2 = [64, 190]
Q8 \times D8 = [64, 230]
(C4 \times Q8) : C2 = [64, 233]
(C4 \times Q8) : C2 = [64, 235]
(C4 \times Q8) : C2 = [64, 237]
((C4 \times C4) : C2) : C2 = [64, 244]
(C6 \times S3) : C2 = [72, 23]
C2 \times ((C3 \times C3) : C4) = [72, 45]
C2 \times D8 \times S3 = [96, 209]
C2 \times ((C3 \times C3 \times C3) : C2) = [108, 44]
```

# Five groups with difference 27.

C29 = [ 29, 1 ] C36 = [ 36, 2 ] (C14 x C2) : C2 = [ 56, 7 ] D58 = [ 58, 1 ] D72 = [ 72, 6 ]

#### Fifty-seven groups with difference 28.

```
C20 \times C2 = [40, 9]
C8 \times S3 = [48, 4]
C24 : C2 = [48, 5]
C3 : Q16 = [48, 8]
C12 \times C4 = [48, 20]
C3 \times (C4 : C4) = [48, 22]
C3 \times QD16 = [48, 26]
C6 \times Q8 = [48, 46]
C3 \times D18 = [54, 3]
(C9 : C3) : C2 = [54, 6]
A5 = [60, 5]
((C8 \times C2) : C2) : C2 = [64, 8]
C4 \times C4 \times C4 = [64, 55]
(C4 \times C4) : C4 = [64, 57]
C4 \times (C4 : C4) = [64, 59]
(C4 \times C4) : C4 = [64, 63]
(C4 \times C4) : C4 = [64, 64]
(C4 \times C4) : C4 = [64, 65]
(C4 : C4) : C4 = [64, 68]
(C4 : C4) : C4 = [64, 70]
(C2 \times Q8) : C4 = [64, 72]
(C4 \times C2) : Q8 = [64, 76]
(C2 \times C2 \times C2) . (C2 \times C2 \times C2) = [64, 79]
(C2 \times C2 \times C2) . (C2 \times C2 \times C2) = [64, 81]
(C2 \times C2 \times C2) . (C2 \times C2 \times C2) = [64, 82]
C2 \times ((C8 : C2) : C2) = [64, 92]
(C8 \times C2 \times C2) : C2 = [64, 97]
(C2 \times (C8 : C2)) : C2 = [64, 98]
C2 \times ((C4 \times C4) : C2) = [64, 101]
(C2 \times (C8 : C2)) : C2 = [64, 102]
(C2 \times C2 \times Q8) : C2 = [64, 129]
(C2 \times Q16) : C2 = [64, 133]
(C8 \times C2 \times C2) : C2 = [64, 146]
(C2 \times (C8 : C2)) : C2 = [64, 149]
(C2 \times D16) : C2 = [64, 153]
(C2 \times (C4 : C4)) : C2 = [64, 161]
(C2 \times (C4 : C4)) : C2 = [64, 162]
((C8 \times C2) : C2) : C2 = [64, 163]
C2 \times C4 \times Q8 = [64, 197]
(C4 \times Q8) : C2 = [64, 200]
```

```
C2 \times ((C2 \times C2) \cdot (C2 \times C2 \times C2)) = [64, 208]
C2 \times (C4 : Q8) = [64, 212]
(C4 \times Q8) : C2 = [64, 214]
(C4 \times Q8) : C2 = [64, 222]
(C4 : Q8) : C2 = [64, 225]
C2 \times ((C2 \times Q8) : C2) = [64, 255]
(C2 \times Q16) : C2 = [64, 259]
A4 \times S3 = [72, 44]
C2 \times C6 \times S3 = [72, 48]
C2 \times D40 = [80, 37]
(C12 \times C4) : C2 = [96, 81]
C2 \times C2 \times ((C6 \times C2) : C2) = [96, 219]
C2 \times C2 \times S4 = [96, 226]
(C2 \times C2 \times C2 \times C2 \times C2 \times C2) : C2 = [128, 1578]
(C4 \times C4 \times C4) : C2 = [128, 1599]
C2 \times D8 \times D8 = [128, 2194]
(C2 \times ((C2 \times D8) : C2)) : C2 = [128, 2326]
```

## Nine groups with difference 29.

C31 = [ 31, 1 ] C33 = [ 33, 1 ] C5 : C8 = [ 40, 1 ] C11 : C4 = [ 44, 1 ] C3 : Q16 = [ 48, 18 ] C4 x D14 = [ 56, 4 ] D62 = [ 62, 1 ] D66 = [ 66, 3 ] (C6 x S3) : C2 = [ 72, 22 ]

Twenty-seven groups with difference 30.

```
C34 = [ 34, 2 ]
C5 \times Q8 = [ 40, 11 ]
C2 \times (C3 : C8) = [ 48, 9 ]
(C3 : C8) : C2 = [ 48, 10 ]
(C4 : C8) : C2 = [ 64, 12 ]
(C4 \times C2 \times C2) : C4 = [ 64, 33 ]
(C16 \times C2) : C2 = [ 64, 38 ]
C4 \times QD16 = [ 64, 119 ]
(C4 \times Q8) : C2 = [ 64, 121 ]
((C4 \times C4) : C2) : C2 = [ 64, 137 ]
(Q8 : C4) : C2 = [ 64, 142 ]
(C2 \times Q16) : C2 = [ 64, 145 ]
```

```
(C8 : C4) : C2 = [ 64, 155 ]
(C8 : C4) : C2 = [ 64, 157 ]
(C8 : C4) : C2 = [ 64, 159 ]
(C8 x C4) : C2 = [ 64, 169 ]
(Q8 : C4) : C2 = [ 64, 170 ]
(C2 x Q16) : C2 = [ 64, 178 ]
C2 x QD32 = [ 64, 187 ]
(C16 x C2) : C2 = [ 64, 189 ]
Q8 : Q8 = [ 64, 238 ]
Q8 x Q8 = [ 64, 239 ]
(C2 x C2) . (C2 x C2 x C2 x C2) = [ 64, 245 ]
D68 = [ 68, 4 ]
C4 x ((C3 x C3) : C2) = [ 72, 32 ]
(C2 x C2 x C2 x C3 x C2 = [ 96, 89 ]
(D8 x S3) : C2 = [ 96, 216 ]
```

#### Five groups with difference 31.

C35 = [ 35, 1 ] C15 : C4 = [ 60, 7 ] D70 = [ 70, 3 ] (C3 x C3) : Q8 = [ 72, 41 ] D8 x D10 = [ 80, 39 ]

#### Sixty-four groups with difference 32.

```
C40 = [40, 2]
C7 \times S3 = [42, 3]
C24 \times C2 = [48, 23]
C3 \times (C8 : C2) = [48, 24]
C3 \times Q16 = [48, 27]
C6 \times D10 = [60, 10]
((C8 \times C2) : C2) : C2 = [64, 4]
(C2 \times Q8) : C4 = [64, 9]
(C8 \times C2) : C4 = [64, 18]
(C4 \times C4) : C4 = [64, 20]
(C8 \times C2) : C4 = [64, 21]
C2 \times ((C8 \times C2) : C2) = [64, 87]
(C2 \times (C8 : C2)) : C2 = [64, 88]
(C8 \times C2 \times C2) : C2 = [64, 89]
(C2 \times (C8 : C2)) : C2 = [64, 94]
C2 \times (Q8 : C4) = [64, 96]
(Q8 : C4) : C2 = [64, 100]
C2 \times (C8 : C4) = [64, 106]
```

```
C2 \times (C8 : C4) = [64, 107]
(C8 : C4) : C2 = [64, 108]
(C8 : C4) : C2 = [64, 109]
(C2 \times Q16) : C2 = [64, 132]
(C2 \times Q16) : C2 = [64, 148]
(C2 \times Q16) : C2 = [64, 151]
(C2 \times QD16) : C2 = [64, 152]
(Q8 : C4) : C2 = [64, 164]
(Q8 : C4) : C2 = [64, 165]
(C8 : C4) : C2 = [64, 166]
C8 \times C2 \times C2 \times C2 = [64, 246]
C2 \times C2 \times (C8 : C2) = [64, 247]
C2 \times ((C8 \times C2) : C2) = [64, 248]
(C2 \times (C8 : C2)) : C2 = [64, 249]
C2 \times C2 \times Q16 = [64, 252]
C17 : C4 = [68, 3]
(C3 \times (C3 : C4)) : C2 = [72, 21]
C6 \times A4 = [72, 47]
C6 \times C6 \times C2 = [72, 50]
D80 = [80, 7]
(C2 \times (C5 : C4)) : C2 = [80, 34]
C2 \times C2 \times (C5 : C4) = [80, 50]
C2 \times D48 = [96, 110]
(C2 \times C2 \times C2 \times S3) : C2 = [96, 144]
(C6 \times D8) : C2 = [96, 147]
(C6 \times D8) : C2 = [96, 211]
C2 \times C2 \times C2 \times A4 = [96, 228]
C2 \times ((C2 \times C2 \times C2 \times C2) : C3) = [96, 229]
C6 \times C2 \times C2 \times C2 \times C2 = [96, 231]
C2 \times ((C2 \times ((C4 \times C2) : C2)) : C2) = [128, 1009]
C2 \times ((C2 \times C2 \times D8) : C2) = [128, 1116]
(C2 \times C2 \times C2 \times D8) : C2 = [128, 1135]
(C2 \times C2 \times C2 \times D8) : C2 = [128, 1165]
C2 \times C2 \times C2 \times ((C4 \times C2) : C2) = [128, 2151]
C2 \times C2 \times ((C4 \times C2 \times C2) : C2) = [128, 2164]
C2 \times ((C2 \times C2 \times D8) : C2) = [128, 2177]
C2 \times ((C2 \times C2 \times D8) : C2) = [128, 2178]
(C2 \times C2 \times C2 \times D8) : C2 = [128, 2216]
(C2 \times ((C4 \times C4) : C2)) : C2 = [128, 2230]
C2 \times C2 \times C2 \times D16 = [128, 2306]
C4 \times C2 \times C2 \times C2 \times C2 \times C2 = [128, 2319]
C2 \times C2 \times C2 \times ((C4 \times C2) : C2) = [128, 2322]
```

C2 x ((C2 x ((C4 x C2) : C2)) : C2) = [ 128, 2325 ] C2 x C2 x C2 x ((C3 x C3) : C2) = [ 144, 196 ] C2 x C2 x C2 x C2 x C2 x C3 = [ 192, 1542 ] C2 x C2 x C2 x C2 x C2 x C2 x D8 = [ 256, 56083 ]

## 4. Acknowledgement

This paper is a continuation of [1], which was an expansion and revision of the Master's thesis of the second author, directed by the first and third authors.

## References

- [1] R. Belshoff, J. Dillstrom, and L. Reid. Finite groups with a prescribed number of cyclic subgroups. To appear in Communications in Algebra.
- [2] GAP Groups, Algorithms, and Programming, Version 4.8.5, The GAP Group (2016), http://www.gap-system.org.
- [3] G.A. Miller. Note on the possible number of operators of order 2 in a group of order 2<sup>n</sup>. Ann. of Math. (2) 7 (1906), no. 2, 55-60.
- [4] Marius Tărnăuceanu. Finite groups with a certain number of cyclic subgroups. *Amer. Math. Monthly* 122 (2015), 275–276.
- [5] Marius Tărnăuceanu, Finite groups with a certain number of cyclic subgroups II, http://arxiv.org/abs/1604.04974.

Department of Mathematics, Missouri State University, Springfield, MO 65897

E-mail address, Corresponding author: RBelshoff@MissouriState.edu

Department of Mathematics, Missouri State University, Springfield, MO 65897

E-mail address: Dillstrom008@live.missouristate.edu

DEPARTMENT OF MATHEMATICS, MISSOURI STATE UNIVERSITY, SPRINGFIELD, MO 65897

*E-mail address*: LesReid@MissouriState.edu