Discrete and Ultradiscrete Periodic Phase Soliton Equations

Hidetomo Nagai¹^{*}, Yasuhiro Ohta²[†], and Ryogo Hirota^{3‡}

¹Department of Mathematics, Tokai University, 4-1-1 Kitakaname, Hiratsuka, Kanagawa 259-1292, Japan

²Department of Mathematics, Kobe University, 1-1 Rokkodai, Nada, Kobe, Hyogo 657-8501, Japan

³Professor Emeritus, Waseda University, 3-4-1 Okubo, Shinjuku, Tokyo 169-8555, Japan

Abstract

We propose new type of discrete and ultradiscrete soliton equations, which admit extended soliton solution called periodic phase soliton solution. The discrete equation is derived from the discrete DKP equation and the ultradiscrete one is obtained by applying the ultradiscrete limit. The soliton solutions have internal freedom and change their shape periodically during propagation. In particular, the ultradiscrete solution reduces into the solution to the ultradiscrete hungry Lotka-Volterra equation in a special case.

1 Introduction

The periodic phase soliton (PPS) is a recently discovered wave for the ultradiscrete hungry Lotka-Volterra (uhLV) equation[1]. Usual soliton is a traveling wave which maintains its shape during propagation, while the PPS is the one which keeps changing its shape periodically. Since the PPS solutions have many internal freedoms, they can exhibit various types of interactions. As far as the authors know, this type of periodically oscillating solution has been given for the uhLV equation only. The PPS in Ref. [1] is the ultradiscrete solution and even its discrete analogue has not been reported yet.

In this paper the PPS solutions are constructed for the following discrete soliton equation,

$$\frac{u_m^{l+1}}{u_m^l} = \prod_{k=1}^M \frac{1 + \delta u_{m-k}^l}{1 + \delta u_{m+k}^{l+1}} \frac{1 + \delta w_{m+k}^{l+1}}{1 + \delta w_{m-k}^l},$$

$$\frac{w_m^{l+1}}{w_m^l} = \prod_{k=1}^M \frac{1 + \delta u_{m-k+1}^l}{1 + \delta u_{m+k-1}^{l+1}} \frac{1 + \delta w_{m-k+1}^{l+1}}{1 + \delta w_{m-k+1}^l},$$

$$\frac{u_{m+M-1}^{l+1}}{u_m^l} = \frac{w_{m+M}^{l+1}}{w_{m-1}^l},$$
(1)
(2)

where m and l denote discrete space and time variables, respectively, M means a range of interaction and δ is a difference interval. The time evolution of u_m^l and w_m^l is described by (1), and Eq.(2) gives a consistent constraint. We call (1) and (2) a discrete PPS (dPPS) equation. It is noted that dPPS

^{*}hdnagai@tokai-u.jp

[†]ohta@math.sci.kobe-u.ac.jp

[‡]Deceased 17 January 2015

equation reduces to the discrete hungry Lotka-Volterra equation when $w_m^l = 0$. By applying the variable transformations,

$$u_m^l = e^{U_m^l/\varepsilon}, \quad w_m^l = e^{W_m^l/\varepsilon}, \quad \delta = e^{-1/\varepsilon}$$
(3)

and taking the ultradiscrete limit[2], i.e., $\varepsilon \to +0$, (1) and (2) reduce to the ultradiscrete PPS (uPPS) equation,

$$U_m^{l+1} - U_m^l = \sum_{k=1}^M \left(\max(1, U_{m-k}^l) - \max(1, U_{m+k}^{l+1}) + \max(1, W_{m+k}^{l+1}) - \max(1, W_{m-k}^{l}) \right),$$

$$W_m^{l+1} - W_m^l = \sum_{k=1}^M \left(\max(1, U_{m-k+1}^l) - \max(1, U_{m+k-1}^{l+1}) + \max(1, W_{m+k-1}^{l+1}) - \max(1, W_{m-k+1}^{l}) \right),$$
(4)

$$U_{m+M-1}^{l+1} - U_m^l = W_{m+M}^{l+1} - W_{m-1}^l.$$
(5)

The ultradiscrete analogue of PPS solutions for the above system are also constructed. Moreover, we show that the PPS solutions for uhLV equation are derived from the ones for the above uPPS equation as a special case.

The contents of this article are as follows. In section 2, we give the discrete DKP equation and its solution. In section 3, we derive the dPPS equation and solution from the discrete DKP. In section 4, we ultradiscretize the PPS solution. In section 5, we show the relation between the obtained solution and that of the uhLV equation. Finally, the concluding remarks are given in section 6.

2 Discrete DKP Equation and Solution

Let us start from the discrete DKP equation[3, 4], which is expressed by

$$\delta(\tau(k+1,l+1,m+1)\tau(k,l,m) - \tau(k+1,l+1,m)\tau(k,l,m+1)) = \tau(k+1,l,m+1)\tau(k,l+1,m) - \tau(k+1,l,m)\tau(k,l+1,m+1),$$
(6)

where δ is a constant. We have the pfaffian solution to discrete DKP equation,

$$\tau(k,l,m) = pf(\alpha_1,\alpha_2,\dots,\alpha_N,\beta_N,\dots,\beta_2,\beta_1)_{k,l,m},$$
(7)

where the pfaffian elements are defined by

$$pf(\alpha_i, \alpha_j)_{k,l} = a_{ij}\eta_i(k, l)\eta_j(k, l), \quad pf(\alpha_i, \beta_j)_{k,l,m} = \delta_{i,j} + \eta_i(k, l)\psi_j(m), \quad pf(\beta_i, \beta_j)_m = b_{ij}\psi_{ij}(m), \quad (8)$$

$$a_{ij} = \frac{q_j - q_i}{1 - q_i q_j}, \qquad \eta_i(k, l) = q_i^k \omega_i^l c_i, \qquad \omega_i = \frac{q_i + \delta}{1 + \delta q_i},$$

$$\delta_{i,j} = \begin{cases} 1 & (i = j) \\ 0 & (i \neq j) \end{cases}, \qquad \psi_i(m) = p_i^m \phi_i(m), \qquad b_{ij} = \frac{1}{1 - p_i^M p_j^M},$$

$$\psi_{ij}(m) = \sum_{1 \le \mu \le M} (\psi_i(m + \mu)\psi_j(m + \mu - 1) - \psi_i(m + \mu - 1)\psi_j(m + \mu)),$$
(9)

where p_i , q_i and c_i are arbitrary constants, M is a positive integer, and $\phi_i(m)$ is a periodic function which satisfies $\phi_i(m+M) = \phi_i(m)$. Here and hereafter we denote k, l and m as suffix, and may omit unshifted independent variables for simplicity. Then the pfaffian (7) satisfies Eq.(6) (see Appendix B).

3 Discrete PPS Equation and Solution

We shall reduce the discrete DKP equation and solution given in the previous section into the dPPS equation and solution.

Proposition 3.1 Suppose $q_i = p_i^M$ in (9). Then the pfaffian (7) satisfies

$$\tau(k+1, l, m) = \tau(k, l, m+M).$$
(10)

(11)

Proof. Let us use the triangle matrix notation,

$$\begin{aligned} & \left| \begin{array}{cccc} \mathrm{pf}(\alpha_{1},\alpha_{2}) & \dots & \mathrm{pf}(\alpha_{1},\alpha_{N}) & \mathrm{pf}(\alpha_{1},\beta_{N}) & \mathrm{pf}(\alpha_{1},\beta_{N-1}) & \dots & \mathrm{pf}(\alpha_{1},\beta_{1}) \\ & \ddots & \vdots & & \vdots & & \vdots & & \vdots \\ & & \mathrm{pf}(\alpha_{N-1},\alpha_{N}) & \mathrm{pf}(\alpha_{N-1},\beta_{N}) & \mathrm{pf}(\alpha_{N-1},\beta_{N-1}) & \dots & \mathrm{pf}(\alpha_{N-1},\beta_{1}) \\ & & & \mathrm{pf}(\alpha_{N},\beta_{N}) & \mathrm{pf}(\alpha_{N},\beta_{N-1}) & \dots & \mathrm{pf}(\alpha_{N},\beta_{1}) \\ & & & & \mathrm{pf}(\beta_{N},\beta_{N-1}) & \dots & \mathrm{pf}(\beta_{N},\beta_{1}) \\ & & & & \ddots & \vdots \\ & & & & \mathrm{pf}(\beta_{2},\beta_{1}) \end{array} \end{aligned}$$

Then we have

$$\tau(k+1,l,m) = \left| \begin{array}{c} \operatorname{pf}(\alpha_{i},\alpha_{j})_{k+1} & \operatorname{pf}(\alpha_{i},\beta_{j})_{k+1} \\ \operatorname{pf}(\beta_{i},\beta_{j})_{k+1} \end{array} \right| \\ = \left| \begin{array}{c} p_{i}^{M} p_{j}^{M} \operatorname{pf}(\alpha_{i},\alpha_{j})_{k} & \delta_{i,j} + p_{i}^{M} \eta_{i}(k,l) \psi_{j}(m) \\ \operatorname{pf}(\beta_{i},\beta_{j})_{k} \end{array} \right| \\ = \left(\prod_{i=1}^{N} p_{i}^{M} \right) \left| \begin{array}{c} \operatorname{pf}(\alpha_{i},\alpha_{j})_{k} & \delta_{i,j} / p_{i}^{M} + \eta_{i}(k,l) \psi_{j}(m) \\ \operatorname{pf}(\beta_{i},\beta_{j})_{k} \end{array} \right| .$$

$$(12)$$

On the other hand, we also have

$$\tau(k,l,m+M) = \begin{vmatrix} \operatorname{pf}(\alpha_{i},\alpha_{j})_{m+M} & \operatorname{pf}(\alpha_{i},\beta_{j})_{m+M} \\ \operatorname{pf}(\beta_{i},\beta_{j})_{m+M} \end{vmatrix}$$

$$= \begin{vmatrix} \operatorname{pf}(\alpha_{i},\alpha_{j})_{m} & \delta_{i,j} + p_{j}^{M} \eta_{i}(k,l)\psi_{j}(m) \\ p_{i}^{M} p_{j}^{M} \operatorname{pf}(\beta_{i},\beta_{j})_{m} \end{vmatrix}$$

$$= \left(\prod_{i=1}^{N} p_{i}^{M}\right) \mid \operatorname{pf}(\alpha_{i},\alpha_{j})_{m} \quad \delta_{i,j}/p_{j}^{M} + \eta_{i}(k,l)\psi_{j}(m) \\ \operatorname{pf}(\beta_{i},\beta_{j})_{m} \end{vmatrix} ,$$

$$(13)$$

which is identical with (12). This completes the proof.

Proposition 3.1 shows the pfaffian (7) under the condition $q_i = p_i^M$ also satisfies

$$\delta(\tau(k, l+1, m+1+M)\tau(k, l, m) - \tau(k, l+1, m+M)\tau(k, l, m+1)) = \tau(k, l, m+1+M)\tau(k, l+1, m) - \tau(k, l, m+M)\tau(k, l+1, m+1).$$
(14)

Denote f_m^l as $\tau(k, l, m - Ml)$. Then Eq.(14) is reduced into

$$\delta(f_{m+M+1}^{l+1}f_{m-M}^{l} - f_{m+M}^{l+1}f_{m-M+1}^{l}) = f_{m+1}^{l}f_{m}^{l+1} - f_{m}^{l}f_{m+1}^{l+1}.$$
(15)

Thus we have the following proposition [5].

Proposition 3.2 Let f_m^l be

$$f_m^l = pf(a_1, a_2, \dots, a_N, b_N, \dots, b_2, b_1)_{l,m},$$
(16)

where

$$pf(a_i, a_j) = a_{ij}s_i(l, m)s_j(l, m), \quad pf(a_i, b_j) = \delta_{i,j} + s_i(l, m)\phi_j(m), \quad pf(b_i, b_j) = b_{ij}\phi_{ij}(m),$$
(17)

$$s_i(l,m) = p_i^{m-Ml} \omega_i^l c_i,$$

$$\phi_{ij}(m) = \sum_{1 \le \mu \le M} (p_i^{\mu} p_j^{\mu-1} \phi_i(m+\mu) \phi_j(m+\mu-1) - p_i^{\mu-1} p_j^{\mu} \phi_i(m+\mu-1) \phi_j(m+\mu)).$$
(18)

The definitions of other symbols are the same as (9) with $q_i = p_i^M$. Then, the pfaffian (16) satisfies Eq.(15).

Proof. We show $\tau(k, l, m - Ml)$ coincides with the pfaffian (16). In a similar manner to proposition 3.1, we have

$$\tau(k,l,m-Ml) = \left| \begin{array}{c} pf(\alpha_i,\alpha_j)_{m-Ml} & pf(\alpha_i,\beta_j)_{m-Ml} \\ pf(\beta_i,\beta_j)_{m-Ml} \end{array} \right| \\ = \left| \begin{array}{c} a_{ij}p_i^{kM}\omega_i^l c_i p_j^{kM}\omega_j^l c_j & \delta_{i,j} + p_i^{kM}\omega_i^l c_i p_j^{m-Ml}\phi_j(m) \\ p_i^{m-Ml}p_j^{m-Ml}b_{ij}\phi_{ij} \end{array} \right|.$$

$$(19)$$

Replacing c_i as $p_i^{-kM}c_i$, it is rewritten as

$$\left| \begin{array}{cc} a_{ij}\omega_{i}^{l}c_{i}\omega_{j}^{l}c_{j} & \delta_{i,j} + \omega_{i}^{l}c_{i}p_{j}^{m-Ml}\phi_{j}(m) \\ p_{i}^{m-Ml}p_{j}^{m-Ml}b_{ij}\phi_{ij} \end{array} \right|$$

$$= \left(\prod_{i=1}^{N} p_{i}^{m-Ml}\right) \quad \left| \begin{array}{c} a_{ij}\omega_{i}^{l}c_{i}\omega_{j}^{l}c_{j} & \delta_{i,j}/p_{j}^{m-Ml} + \omega_{i}^{l}c_{i}\phi_{j}(m) \\ b_{ij}\phi_{ij} \end{array} \right|.$$

$$(20)$$

On the other hand, we also have

$$\begin{aligned}
f_m^l &= \left| \begin{array}{c} \operatorname{pf}(a_i, a_j) & \operatorname{pf}(a_i, b_j) \\ \operatorname{pf}(b_i, b_j) \end{array} \right| \\
&= \left| \begin{array}{c} a_{ij} p_i^{m-Ml} \omega_i^l c_i p_j^{m-Ml} \omega_j^l c_j & \delta_{i,j} + p_i^{m-Ml} \omega_i^l c_i \phi_j(m) \\ & b_{ij} \phi_{ij} \end{array} \right| \\
&= \left(\prod_{i=1}^N p_i^{m-Ml} \right) \left| \begin{array}{c} a_{ij} \omega_i^l c_i \omega_j^l c_j & \delta_{i,j} / p_i^{m-Ml} + \omega_i^l c_i \phi_j(m) \\ & b_{ij} \phi_{ij} \end{array} \right|,
\end{aligned} \tag{21}$$

which is identical with (20). This completes the proof.

Now we have the following proposition.

Proposition 3.3 Eq.(15) derives (1) and (2) through transformations

$$u_m^l = \frac{f_{m-M}^l f_{m+M+1}^{l+1}}{f_m^l f_{m+1}^{l+1}}, \qquad w_m^l = \frac{f_{m+M}^{l+1} f_{m-M+1}^l}{f_{m+1}^l f_m^{l+1}}.$$
(22)




Figure 1: 2-soliton solution u_m^l .

Proof. Eq.(2) follows from (22) straightforwardly. Eq.(15) is rewritten as

$$1 + \delta u_m^l = \frac{f_{m+1}^l f_m^{l+1}}{f_m^l f_{m+1}^{l+1}} + \delta \frac{f_{m+M}^{l+1} f_{m-M+1}^l}{f_m^l f_{m+1}^{l+1}},$$

$$1 + \delta w_m^l = \frac{f_m^l f_{m+1}^{l+1}}{f_{m+1}^l f_m^{l+1}} + \delta \frac{f_{m+M+1}^{l+1} f_{m-M}^l}{f_{m+1}^l f_m^{l+1}}.$$
(23)

In particular,

$$\frac{1+\delta u_m^l}{1+\delta w_m^l} = \frac{f_{m+1}^l f_{m+1}^{l+1}}{f_m^l f_{m+1}^{l+1}} \tag{24}$$

holds. Thus, we have

$$\prod_{k=1}^{M} \frac{1+\delta u_{m-k}^{l}}{1+\delta u_{m+k}^{l+1}} \frac{1+\delta w_{m+k}^{l+1}}{1+\delta w_{m-k}^{l}} = \prod_{k=1}^{M} \frac{f_{m+1-k}^{l} f_{m-k}^{l+1}}{f_{m-k}^{l} f_{m+1-k}^{l+1}} \frac{f_{m+k}^{l+1} f_{m+1+k}^{l+2}}{f_{m+1-k}^{l+1} f_{m+1-k}^{l+2}} = \frac{u_{m}^{l+1}}{u_{m}^{l}}.$$
(25)

The other equation can be obtained similarly. This completes the proof.

Due to proposition 3.3, we may call (15) as the dPPS equation in bilinear form. Thus we have derived the pfaffian solution to the dPPS equation. Figures 1, 2, 3 and 4 show the behaviour of the dPPS solution u_m^l and w_m^l in the case of N = M = 2. Here l and m denote time and space variables respectively. In Fig.1 and Fig.2, we set $\delta = 1/3$, $(p_1, p_2, c_1, c_2) = (5, 3, 1, 1)$ and $\phi_i(m) = 1$ for i = 1, 2. They behave as the usual solution solutions. In Fig.3 and Fig.4, we set $\delta = 1/3$, $(p_1, p_2, c_1, c_2) = (5, 3, 1, 1)$ and

$$\phi_1(m) = \begin{cases} 1 & (m: \text{ even}) \\ 2 & (m: \text{ odd}) \end{cases}, \qquad \phi_2(m) = \begin{cases} 3 & (m: \text{ even}) \\ 5 & (m: \text{ odd}) \end{cases}.$$
(26)

We can observe that the high jaggy wave collides with low one and overtakes. The shape of the solitary wave changes periodically.






Figure 2: 2-soliton solution w_m^l .








Figure 3: 2-dPPS solution u_m^l .




Figure 4: 2-dPPS solution w_m^l .

4 Ultradiscretization

In this section we ultradiscretize the pfaffian solution (16). For ultradiscretizing we assume $0 < p_1 < p_2 < \cdots < p_N < 1, 0 < c_i, 0 < \phi_i(m), 0 < \delta$ and

$$\frac{p_i}{p_j} > \max_{1 \le m \le M} \frac{\phi_i(m-1)\phi_j(m)}{\phi_i(m)\phi_j(m-1)},$$

$$p_i p_j \le \min_{1 \le m \le M} \frac{\phi_i(m)\phi_j(m-1)}{\phi_i(m+1)\phi_j(m)},$$
(27)

for i > j. In the case of usual soliton, i.e., when $\phi_i(m)$ is independent of m, the above condition (27) is obviously satisfied. From (27), we have

$$p_{i}\phi_{i}(m)\phi_{j}(m-1) > p_{j}\phi_{i}(m-1)\phi_{j}(m),$$

$$\phi_{i}(m)\phi_{j}(m-1) \ge p_{i}p_{j}\phi_{i}(m+1)\phi_{j}(m),$$
(28)

for i > j and $m = 1, 2, \ldots, M$. Now we introduce reparametrizations,

$$p_i = e^{-P_i/\varepsilon}, \quad \delta = e^{-1/\varepsilon}, \quad \omega_i = e^{\Omega_i/\varepsilon}, \quad c_i = e^{C_i/\varepsilon}, \quad \phi_i(m) = e^{\Phi_i(m)/\varepsilon},$$
 (29)

and we are going to take a limit $\varepsilon \to +0$ for ultradiscretization. It is commented that P_i is positive, Ω_i is negative and $P_i < P_j$ for i > j. In the following, we show the ultradiscrete limit of PPS solutions, first for N = 2, 3 and next for general N.

4.1 Case of N = 2

In the case of $N = 2, f_m^l$ is expressed by

$$\begin{aligned}
f_m^l &= pf(a_1, a_2, b_2, b_1) \\
&= 1 + s_1\phi_1(m) + s_2\phi_2(m) + a_{12}s_1s_2b_{21}\phi_{21}(m).
\end{aligned}$$
(30)

We can ultradiscretize each variable as

$$\begin{aligned}
\omega_i &\to \Omega_i = \max(-MP_i, -1), \\
s_i(l, m) &\to S_i(l, m) = -(m - Ml)P_i + l\Omega_i + C_i, \\
a_{12} &\to -MP_2, \\
b_{21} &\to 0
\end{aligned}$$
(31)

from (18). Let us ultradiscretize $\phi_{21}(m)$, which consists of 2M terms. The greatest term among these terms is $p_2\phi_2(m+1)\phi_1(m)$ since we can derive

$$p_{i}^{\mu}p_{j}^{\mu-1}\phi_{i}(m+\mu)\phi_{j}(m+\mu-1) > p_{i}^{\mu-1}p_{j}^{\mu}\phi_{i}(m+\mu-1)\phi_{j}(m+\mu),$$

$$p_{i}^{\mu}p_{j}^{\mu-1}\phi_{i}(m+\mu)\phi_{j}(m+\mu-1) \ge p_{i}^{\mu+1}p_{j}^{\mu}\phi_{i}(m+\mu+1)\phi_{j}(m+\mu),$$
(32)

for $\mu = 1, 2, \ldots, M$ from (28). Thus we obtain

$$\phi_{21}(m) \rightarrow -P_2 + \Phi_2(m+1) + \Phi_1(m).$$
 (33)

Therefore the ultradiscrete analogue of (30) is given by

$$f_m^l \to F_m^l = \max(0, S_1 + \Phi_1(m), S_2 + \Phi_2(m), S_1 + S_2 - (M+1)P_2 + \Phi_1(m) + \Phi_2(m+1)).$$
 (34)




Figure 5: 2-soliton solution U_m^l .




Figure 6: 2-soliton solution W_m^l .

Note F_m^l leads solutions to the uPPS equation (4) and (5) through the transformations

$$U_m^l = F_{m-M}^l + F_{m+M+1}^{l+1} - F_m^l - F_{m+1}^{l+1}, \qquad W_m^l = F_{m+M}^{l+1} + F_{m-M+1}^l - F_{m+1}^l - F_m^{l+1}.$$
(35)

Figures 5, 6, 7 and 8 show behaviour of the uPPS solution U_m^l and W_m^l in the case of N = M = 2. In Fig.5 and Fig.6, we set $(P_1, P_2, C_1, C_2) = (10, 2, 0, 0)$ and $\Phi_i(m) = 0$ for i = 1, 2. In Fig.7 and Fig.8, we set $(P_1, P_2, C_1, C_2) = (10, 2, 0, 0)$ and

$$\Phi_1(m) = \begin{cases} 4 & (m: \text{ even}) \\ 3 & (m: \text{ odd}) \end{cases}, \qquad \Phi_2(m) = \begin{cases} 1 & (m: \text{ even}) \\ 6 & (m: \text{ odd}) \end{cases}.$$
(36)

They behave similarly to the discrete ones.

4.2 Case of N = 3

In the case of N = 3, f_m^l is expanded as

$$f_{m}^{l} = pf(a_{1}, a_{2}, a_{3}, b_{3}, b_{2}, b_{1})$$

$$= 1 + s_{1}\phi_{1}(m) + s_{2}\phi_{2}(m) + s_{3}\phi_{3}(m)$$

$$+ a_{12}s_{1}s_{2}b_{21}\phi_{21}(m) + a_{13}s_{1}s_{3}b_{31}\phi_{31}(m) + a_{23}s_{2}s_{3}b_{32}\phi_{32}(m)$$

$$+ s_{1}s_{2}s_{3}(a_{12} - a_{13} + a_{23})(b_{21}\phi_{3}(m)\phi_{21}(m) - b_{31}\phi_{2}(m)\phi_{31}(m) + b_{32}\phi_{1}(m)\phi_{32}(m)).$$
(37)

One can prove that $a_{12} - a_{13} + a_{23}$ is equal to $a_{12}a_{13}a_{23}$ and it is ultradiscretized as $-MP_2 - 2MP_3$. Moreover, we can determine the greatest term among the expansion of $b_{21}\phi_3\phi_{21} - b_{31}\phi_2\phi_{31} + b_{32}\phi_1\phi_{32}$




Figure 7: 2-uPPS solution U_m^l .




Figure 8: 2-uPPS solution W_m^l .

as $p_2 p_3^2 \phi_1(m) \phi_2(m+1) \phi_3(m+2)$. It is ultradiscretized as $-P_2 - 2P_3 + \Phi_1(m) + \Phi_2(m+1) + \Phi_3(m+2)$ (see appendix D). Thus the ultradiscrete analogue of (37) is obtained by

$$F_m^l = \max\left(0, S_1 + \Phi_1(m), S_2 + \Phi_2(m), S_3 + \Phi_3(m), S_1 + S_2 - (M+1)P_2 + \Phi_1(m) + \Phi_2(m+1), S_1 + S_3 - (M+1)P_3 + \Phi_1(m) + \Phi_3(m+1), S_2 + S_3 - (M+1)P_3 + \Phi_2(m) + \Phi_3(m+1), S_1 + S_2 + S_3 - (M+1)P_2 - 2(M+1)P_3 + \Phi_1(m) + \Phi_2(m+1) + \Phi_3(m+2)\right).$$
(38)

4.3 Case of general N

Let us ultradiscretize f_m^l for general N. First, we use the following relation,

$$pf(a_1, \dots, a_N, b_N, \dots, b_1) = \sum_{0 \le n \le N} \sum_{I_n \subset \{1, 2, \dots, N\}} pf(a_{i_1}, a_{i_2}, \dots, a_{i_n}) pf(b_{i_n}, \dots, b_{i_2}, b_{i_1}),$$
(39)

where $\sum_{I_n \subset \{1,2,\ldots,N\}}$ denotes the summation over all *n*-element subsets $I_n = \{i_1, i_2, \ldots, i_n\}$ chosen from $\{1, 2, \ldots, N\}$. When n = 0, we define $\sum_{I_n \subset \emptyset}$ is 1. When n is odd, we define

$$pf(a_{i_1}, a_{i_2}, \dots, a_{i_n}) = pf(a_{i_1}, a_{i_2}, \dots, a_{i_n}, \emptyset), \quad pf(b_{i_n}, \dots, b_{i_2}, b_{i_1}) = pf(\emptyset, b_{i_n}, \dots, b_{i_2}, b_{i_1})$$
(40)

with the new entries

$$pf(a_i, \emptyset) = s_i(l, m), \qquad pf(\emptyset, b_i) = \phi_i(m).$$
 (41)

For example, f_m^l in the case of N = 2, 3 are expressed by

$$f_{m}^{l} = 1 + pf(a_{1}, \emptyset)pf(\emptyset, b_{1}) + pf(a_{2}, \emptyset)pf(\emptyset, b_{2}) + pf(a_{1}, a_{2})pf(b_{2}, b_{1}),$$

$$f_{m}^{l} = 1 + pf(a_{1}, \emptyset)pf(\emptyset, b_{1}) + pf(a_{2}, \emptyset)pf(\emptyset, b_{2}) + pf(a_{3}, \emptyset)pf(\emptyset, b_{3})$$

$$+ pf(a_{1}, a_{2})pf(b_{2}, b_{1}) + pf(a_{1}, a_{3})pf(b_{3}, b_{1}) + pf(a_{2}, a_{3})pf(b_{3}, b_{2})$$

$$+ pf(a_{1}, a_{2}, a_{3}, \emptyset)pf(\emptyset, b_{3}, b_{2}, b_{1}),$$
(42)

and they correspond to (30) and (37). The proof of the relation (39) is given in appendix C.

Let us consider the expansion of $pf(a_1, a_2, ..., a_N)$ and $pf(b_N, ..., b_2, b_1)$ respectively. We assume N is even for simplicity. Then $pf(a_1, a_2, ..., a_N)$ can be expanded as follows[6].

$$pf(a_1, a_2, \dots, a_N) = \prod_{1 \le i < j \le N} a_{ij} \prod_{1 \le i \le N} s_i.$$

$$(43)$$

Hence introducing transformations (29) under the assumption $0 < p_1 < p_2 < \cdots < p_N < 1$, we ultradiscretize

$$pf(a_1, a_2, \dots, a_N) \to \sum_{1 \le i \le N} S_i - \sum_{1 \le i < j \le N} MP_j = \sum_{1 \le i \le N} S_i - M \sum_{2 \le j \le N} (j-1)P_j.$$
(44)

The expansion of $pf(b_N, \ldots, b_2, b_1)$ cannot be factorized as $pf(a_1, a_2, \ldots, a_N)$. However, the greatest term of $pf(b_N, \ldots, b_2, b_1)$ is determined as $\prod_{i=1}^N p_i^{i-1} \phi_i(m+i-1)$ under the assumption (28) (see appendix D), thus,

$$pf(b_N, \dots, b_2, b_1) \to \sum_{1 \le i \le N} \Big(-(i-1)P_i + \Phi_i(m+i-1) \Big).$$
 (45)

Therefore we obtain the ultradiscrete analogue of (39) as

$$F_m^l = \max_{\mu_i = 0, 1} \Big(\sum_{1 \le i \le N} \mu_i S_i(l, m) - (M+1) \sum_{1 \le i < j \le N} \mu_i \mu_j P_j + \sum_{1 \le j \le N} \mu_j \Phi_j(m + \sum_{1 \le i \le j - 1} \mu_i) \Big),$$
(46)

where $\max_{\mu_i=0,1} X(\mu_1, \mu_2, \dots, \mu_N)$ denotes the maximum value of X in 2^N possible cases $\{\mu_1, \mu_2, \dots, \mu_N\}$ replacing each μ_i by 0 or 1. We note F_m^l satisfies the following equation

$$\max(F_{m+M+1}^{l+1} + F_{m-M}^{l} - 1, F_{m}^{l} + F_{m+1}^{l+1}) = \max(F_{m+M}^{l+1} + F_{m-M+1}^{l} - 1, F_{m+1}^{l} + F_{m}^{l+1}), \quad (47)$$

which is obtained by ultradiscretizing (15).

5 Relation between the uPPS Equation and the uhLV Equation

The discrete hungry Lotka-Volterra equation is ultradiscretized as

$$F_{m+1}^{l} + F_{m}^{l+1} = \max(F_{m}^{l} + F_{m+1}^{l+1}, F_{m-M}^{l} + F_{m+M+1}^{l+1} - 1).$$
(48)

In addition, the uPPS solution to Eq.(48) is proposed as follows[1]:

$$F_m^l = \max_{\mu_i = 0, 1} \left(\sum_{1 \le i \le N} \mu_i S_i(l, m) - (M+1) \sum_{1 \le i < j \le N} \mu_i \mu_j P_j + \sum_{1 \le j \le N} \mu_j \Phi_j(m + \sum_{1 \le i \le j - 1} \mu_i) \right),$$
(49)

where $S_i(l,m) = (MP_i - 1)l - P_im + C_i$. Here P_i and C_i are arbitrary parameters satisfying $1/M \le P_N \le \cdots \le P_2 \le P_1$. Function $\Phi_i(m)$ is a periodic function satisfying $\Phi(m) = \Phi(m+M)$ and the following relations,

$$P_i \ge |\Phi_i(m) - \Phi_i(m+1)|, |P_i - P_j| \ge |\Phi_i(m) - \Phi_i(m+1) - (\Phi_j(m) - \Phi_j(m+1))|.$$
(50)

We show (46) is reduced into (49). Assume $P_i \ge 1/M$ for i = 1, 2, ..., N in (46), then S_i is rewritten as $l \max(0, MP_i - 1) - mP_i + C_i = (MP_i - 1)l - mP_i + C_i$. Moreover ultradiscretization of (28) derives

$$P_{j} - P_{i} > \Phi_{j}(m) - \Phi_{j}(m-1) - (\Phi_{i}(m) - \Phi_{i}(m-1)),$$

$$P_{j} + P_{i} \ge \Phi_{j}(m) - \Phi_{j}(m-1) + \Phi_{i}(m+1) - \Phi_{i}(m),$$
(51)

for i > j and m = 1, 2, ..., M. Thus, if we set distinct wave numbers P_i and periodic functions $\Phi_i(m)$ so that they satisfy (50) and (51), the solution (46) is reduced into (49). In other words, the solution (46) under the assumptions satisfies the uhLV equation (48).

6 Concluding Remarks

We derived the dPPS equation from the discrete DKP equation. It has the dPPS solution, which presents internal oscillatory motion. Moreover, the dPPS solution under the assumptions (28) is ultradiscretized, and we obtained the uPPS solution which is also periodically oscillating and satisfies the uPPS equation (47). We showed that the solution in this paper reduces to the uPPS solution to the uhLV equation in a special case. It suggests that there may be a direct connection between DKP and KP in ultradiscrete case. It might be interesting to clarify the relation between integrable hierarchies and construct various soliton equations admitting periodic phase soliton solutions for both discrete and ultradiscrete limit. There are other choices of parameters which derive solutions for the uPPS equation. Investigating such solutions may be a future work.

Acknowledgments

This work was supported by JSPS KAKENHI Grand Numbers JP24340029, JP26610029 and JP18H01130.

A A Formula for Pfaffian

We first show the following theorem[7].

Theorem A.1 Let n, m be any positive integers. Suppose

$$pf(i,j) = pf(d_1, d_2, \dots, d_{2m}, \alpha_i, \alpha_j) / pf(d_1, d_2, \dots, d_{2m}),$$
(52)

for $i, j = 1, 2, \ldots, 2n$. Then we have

$$pf(1, 2, \dots, 2n) = pf(d_1, d_2, \dots, d_{2m}, \alpha_1, \alpha_2, \dots, \alpha_{2n}) / pf(d_1, d_2, \dots, d_{2m}).$$
(53)

Proof. We prove the theorem by using mathematical induction on n. Assume

$$pf(1, 2, \dots, 2n - 2) = \frac{pf(d_1, d_2, \dots, d_{2m}, \alpha_1, \alpha_2, \dots, \alpha_{2n-2})}{pf(d_1, d_2, \dots, d_{2m})}.$$
(54)

The pfaffian pf(1, 2, ..., 2n) is obtained by the identity

$$pf(1, 2, \dots, 2n) = \sum_{j=2}^{2n} (-1)^j pf(1, j) pf(2, 3, \dots, \hat{j}, \dots, 2n),$$
(55)

where the symbol \hat{j} means the deletion of j. By using (52) and (54), the right-hand side is rewritten as

$$\sum_{j=2}^{2n} (-1)^j \frac{\mathrm{pf}(d_1, d_2, \dots, d_{2m}, \alpha_1, \alpha_j)}{\mathrm{pf}(d_1, d_2, \dots, d_{2m})} \frac{\mathrm{pf}(d_1, d_2, \dots, d_{2m}, \alpha_2, \alpha_3, \dots, \widehat{\alpha_j}, \dots, \alpha_{2n})}{\mathrm{pf}(d_1, d_2, \dots, d_{2m})},$$
(56)

and moreover we have

$$\sum_{j=2}^{2n} (-1)^{j} \mathrm{pf}(d_{1}, d_{2}, \dots, d_{2m}, \alpha_{1}, \alpha_{j}) \mathrm{pf}(d_{1}, d_{2}, \dots, d_{2m}, \alpha_{2}, \alpha_{3}, \dots, \widehat{\alpha_{j}}, \dots, \alpha_{2n})$$

$$= \mathrm{pf}(d_{1}, d_{2}, \dots, d_{2m}) \mathrm{pf}(d_{1}, d_{2}, \dots, d_{2m}, \alpha_{1}, \alpha_{2}, \dots, \alpha_{2n}),$$
(57)

due to the Plücker relation of pfaffian type[8]. Therefore we obtain (53).

B Proof of the Discrete DKP Equation (6)

We show the pfaffian (7) satisfies (6) by using Theorem A.1. In this appendix, we denote (k, l, m) as (k_1, k_2, k_3) , and $\tau(k) = \tau(k_1, k_2, k_3) = pf(\alpha_1, \ldots, \alpha_N, \beta_N, \ldots, \beta_1)_{k_1, k_2, k_3}$. Moreover we use the following notations,

$$\tau(k_{1}) = \tau(k_{1} + 1, k_{2}, k_{3}) = pf(\alpha_{1}, \alpha_{2}, \dots, \alpha_{N}, \beta_{N}, \dots, \beta_{2}, \beta_{1})_{k_{1} + 1, k_{2}, k_{3}},$$

$$\tau(\widetilde{k_{1}}) = \tau(k_{1}, k_{2} + 1, k_{3} + 1) = pf(\alpha_{1}, \alpha_{2}, \dots, \alpha_{N}, \beta_{N}, \dots, \beta_{2}, \beta_{1})_{k_{1}, k_{2} + 1, k_{3} + 1},$$

$$\tau(\overline{k}) = \tau(k_{1} + 1, k_{2} + 1, k_{3} + 1) = pf(\alpha_{1}, \alpha_{2}, \dots, \alpha_{N}, \beta_{N}, \dots, \beta_{2}, \beta_{1})_{k_{1} + 1, k_{2} + 1, k_{3} + 1},$$
(58)

and $\tau(k_2)$, $\tau(k_3)$, $\tau(\tilde{k_2})$, $\tau(\tilde{k_3})$ are similarly defined. Let us introduce new indices, d_1 , d_2 , d_3 and d_4 , which are defined by

$$pf(d_0, d_1) = pf(d_0, d_2) = 1, \quad pf(d_1, d_2) = -\delta, pf(d_0, d_3) = pf(d_1, d_3) = pf(d_2, d_3) = -1.$$
(59)

In these notations, the discrete DKP equation (6) is expressed by

$$pf(d_0, d_1, d_2, d_3)\tau(\vec{k})\tau(k) = pf(d_0, d_1)pf(d_2, d_3)\tau(k_1)\tau(\vec{k_1}) - pf(d_0, d_2)pf(d_1, d_3)\tau(k_2)\tau(\vec{k_2}) + pf(d_0, d_3)pf(d_1, d_2)\tau(k_3)\tau(\vec{k_3}).$$
(60)

Now let us denote the (i, j)-elements of pfaffians $\tau(k_l)$, $\tau(\tilde{k}_l)$, $\tau(\bar{k})$ as $\tau_{ij}(k_l)$, $\tau_{ij}(\tilde{k}_l)$, $\tau_{ij}(\bar{k})$, respectively, where l = 1, 2, 3 and $i, j = \alpha_1, \alpha_2, \ldots, \alpha_N, \beta_1, \beta_2, \ldots, \beta_N$. Then by defining pfaffian elements,

$$pf(d_0, \alpha_h) = pf(d_3, \alpha_h) = \eta_h(k_1, k_2), \quad pf(d_1, \alpha_h) = q_h \eta_h(k_1, k_2), \quad pf(d_2, \alpha_h) = \omega_h \eta_h(k_1, k_2), \\ pf(d_0, \beta_h) = pf(d_1, \beta_h) = pf(d_2, \beta_h) = \psi_h(k_3), \quad pf(d_3, \beta_h) = \psi_h(k_3 + 1), \quad (1 \le h \le N),$$
(61)

we obtain
$$p_{h}(a_{0}, p_{h}) = p_{h}$$

$$\tau_{ij}(k_l) = pf(d_0, d_l, i, j) / pf(d_0, d_l),$$

$$\tau_{ij}(\tilde{k}_l) = pf(d_1, \hat{d}_l, d_3, i, j) / pf(d_1, \hat{d}_l, d_3),$$

$$\tau_{ij}(\bar{k}) = pf(d_0, d_1, d_2, d_3, i, j) / pf(d_0, d_1, d_2, d_3),$$

(62)

for l = 1, 2, 3 and $i, j = \alpha_1, \alpha_2, \ldots, \alpha_N, \beta_1, \beta_2, \ldots, \beta_N$. For example, in the case of $l = 1, i = \alpha_i, j = \alpha_j$, we have

$$\tau_{\alpha_{i}\alpha_{j}}(k_{1}) = pf(\alpha_{i}, \alpha_{j})_{k_{1}+1, k_{2}, k_{3}} = a_{ij}q_{i}q_{j}\eta_{i}(k_{1}, k_{2})\eta_{j}(k_{1}, k_{2}) = pf(d_{0}, d_{1}, \alpha_{i}, \alpha_{j})/pf(d_{0}, d_{1}),$$

$$\tau_{\alpha_{i}\alpha_{j}}(\tilde{k}_{1}) = pf(\alpha_{i}, \alpha_{j})_{k_{1}, k_{2}+1, k_{3}+1} = a_{ij}\omega_{i}\omega_{j}\eta_{i}(k_{1}, k_{2})\eta_{j}(k_{1}, k_{2}) = pf(d_{2}, d_{3}, \alpha_{i}, \alpha_{j})/pf(d_{2}, d_{3}),$$

$$\tau_{\alpha_{i}\alpha_{j}}(\bar{k}) = pf(\alpha_{i}, \alpha_{j})_{k_{1}+1, k_{2}+1, k_{3}+1} = a_{ij}q_{i}q_{j}\omega_{i}\omega_{j}\eta_{i}(k_{1}, k_{2})\eta_{j}(k_{1}, k_{2})$$

$$= pf(d_{0}, d_{1}, d_{2}, d_{3}, \alpha_{i}, \alpha_{j})/pf(d_{0}, d_{1}, d_{2}, d_{3}).$$
(63)

The other cases of (62) can be proved in a similar way. Then due to Theorem A.1, the pfaffians are given by

$$\tau(k_l) = pf(d_0, d_l, \alpha_1, \alpha_2, \dots, \alpha_N, \beta_N, \dots, \beta_2, \beta_1) / pf(d_0, d_l),$$

$$\tau(\widetilde{k}_l) = pf(d_1, \widehat{d}_l, d_3, \alpha_1, \alpha_2, \dots, \alpha_N, \beta_N, \dots, \beta_2, \beta_1) / pf(d_1, \widehat{d}_l, d_3),$$

$$\tau(\overline{k}) = pf(d_0, d_1, d_2, d_3, \alpha_1, \alpha_2, \dots, \alpha_N, \beta_N, \dots, \beta_2, \beta_1) / pf(d_0, d_1, d_2, d_3),$$

(64)

and (60) is rewritten as

$$pf(d_0, d_1, d_2, d_3, \alpha_1, \alpha_2, \dots, \alpha_N, \beta_N, \dots, \beta_2, \beta_1)pf(\alpha_1, \alpha_2, \dots, \alpha_N, \beta_N, \dots, \beta_2, \beta_1)$$

$$=pf(d_0, d_1, \alpha_1, \alpha_2, \dots, \alpha_N, \beta_N, \dots, \beta_2, \beta_1)pf(d_2, d_3, \alpha_1, \alpha_2, \dots, \alpha_N, \beta_N, \dots, \beta_2, \beta_1)$$

$$-pf(d_0, d_2, \alpha_1, \alpha_2, \dots, \alpha_N, \beta_N, \dots, \beta_2, \beta_1)pf(d_1, d_3, \alpha_1, \alpha_2, \dots, \alpha_N, \beta_N, \dots, \beta_2, \beta_1)$$

$$+pf(d_0, d_3, \alpha_1, \alpha_2, \dots, \alpha_N, \beta_N, \dots, \beta_2, \beta_1)pf(d_1, d_2, \alpha_1, \alpha_2, \dots, \alpha_N, \beta_N, \dots, \beta_2, \beta_1),$$
(65)

which is nothing but the pfaffian's identity[9]. This completes the proof.

C Proof of (39)

In this appendix, we prove the relation (39) by showing the correspondence of the terms which have $s_1s_2...s_n$ in the both sides. When n is even, the terms on the right hand side of (39) are given by $pf(a_1, a_2, ..., a_n)pf(b_n, ..., b_2, b_1)$ since

$$pf(a_1, a_2, \dots, a_n) = \sum' sgn \begin{pmatrix} 1 & 2 & \dots & n \\ i_1 & i_2 & \dots & i_n \end{pmatrix} pf(a_{i_1}, a_{i_2}) pf(a_{i_3}, a_{i_4}) \dots pf(a_{i_{n-1}}, a_{i_n}) = \sum' sgn \begin{pmatrix} 1 & 2 & \dots & n \\ i_1 & i_2 & \dots & i_n \end{pmatrix} a_{i_1 i_2} a_{i_3 i_4} \dots a_{i_{n-1} i_n} s_1 s_2 \dots s_n,$$
(66)

where \sum' denotes the summation over all permutations $\{i_1, i_2, \ldots, i_n\}$ of $\{1, 2, \ldots, n\}$ which satisfy the inequalities

$$i_1 < i_2, \ i_3 < i_4, \ \dots, \ i_{n-1} < i_n,$$
 and $i_1 < i_3 < \dots < i_{n-1}.$ (67)

Let us consider the terms on the left hand side of (39). We can find that all of the terms which have $s_1s_2...s_n$ belong to

$$pf(a_1, a_2, \dots, a_n, b_n, \dots, b_2, b_1) pf(a_{N-n}, b_{N-n}) pf(a_{N-n+1}, b_{N-n+1}) \dots pf(a_N, b_N)$$
(68)

since $pf(a_i, b_j)$ or $pf(a_i, a_j)$ has $s_i\phi_j$ or $a_{ij}s_is_j$ for $i \neq j$. Then we consider

$$pf(a_1, a_2, \dots, a_n, b_n, \dots, b_2, b_1).$$
 (69)

Expansion of (69) has the terms

$$(-1)^{i_1+i_2+j_1+j_2} \operatorname{pf}(a_{i_1}, b_{j_2}) \operatorname{pf}(a_{i_2}, b_{j_1}) \operatorname{pf}(\bullet), (-1)^{i_1+i_2+j_1+j_2+1} \operatorname{pf}(a_{i_1}, b_{j_1}) \operatorname{pf}(a_{i_2}, b_{j_2}) \operatorname{pf}(\bullet),$$
(70)

where $1 \le i_1 < i_2 \le n$, $1 \le j_1 < j_2 \le n$, and $pf(\bullet)$ denotes

$$pf(a_1,\ldots,\widehat{a_{i_1}},\ldots,\widehat{a_{i_2}},\ldots,a_n,b_n,\ldots,\widehat{b_{j_2}},\ldots,\widehat{b_{j_1}},\ldots,b_1).$$
(71)

Each of (70) has $s_{i_1}s_{i_2}$, however, the sum of them does not since

$$pf(a_{i_1}, b_{j_2})pf(a_{i_2}, b_{j_1})pf(\bullet) - pf(a_{i_1}, b_{j_1})pf(a_{i_2}, b_{j_2})pf(\bullet) = (\delta_{i_1, j_2}\delta_{i_2, j_1} + \delta_{i_1, j_2}s_{i_2}\phi_{j_1} + \delta_{i_2, j_1}s_{i_1}\phi_{j_2} - \delta_{i_1, j_1}\delta_{i_2, j_2} - \delta_{i_1, j_1}s_{i_2}\phi_{j_2} - \delta_{i_2, j_2}s_{i_1}\phi_{j_1})pf(\bullet).$$
(72)

Thus, all the terms of $s_1 s_2 \ldots s_n$ among the expansion of (69) are expressed by

$$\sum' \operatorname{sgn} \begin{pmatrix} 1 & 2 & \dots & n \\ i_1 & i_2 & \dots & i_n \end{pmatrix} \operatorname{pf}(a_{i_1}, a_{i_2}) \operatorname{pf}(a_{i_3}, a_{i_4}) \dots \operatorname{pf}(a_{i_{n-1}}, a_{i_n}) \\ \times \sum' \operatorname{sgn} \begin{pmatrix} 1 & 2 & \dots & n \\ j_n & j_{n-1} & \dots & j_1 \end{pmatrix} \operatorname{pf}(b_{j_n}, b_{j_{n-1}}) \dots \operatorname{pf}(b_{j_4}, b_{j_3}) \operatorname{pf}(b_{j_2}, b_{j_1}),$$
(73)

and it corresponds to $pf(a_1, a_2, \ldots, a_n)pf(b_n, \ldots, b_2, b_1)$.

When n is odd, the terms of $s_1 s_2 \ldots s_n$ on the right hand side of (39) are given by $pf(a_1, a_2, \ldots, a_n, \emptyset) pf(\emptyset, b_n, \ldots, b_2, b_1)$. On the other hand, the terms on the left hand side are obtained from

$$\sum_{1 \le i \le n} \sum_{1 \le j \le n} (-1)^{i+j} \operatorname{pf}(a_i, b_j) \operatorname{pf}(a_1, \dots, \widehat{a_i}, \dots, a_n, b_n, \dots, \widehat{b_j}, \dots, b_1),$$
(74)

among which the terms containing $s_1 s_2 \dots s_n$ appear in

$$\sum_{1 \le i \le n} \sum_{1 \le j \le n} (-1)^{i+j} s_i \phi_j(m) \operatorname{pf}(a_1, \dots, \widehat{a_i}, \dots, a_n) \operatorname{pf}(b_n, \dots, \widehat{b_j}, \dots, b_1)$$

=
$$\sum_{1 \le i \le n} \sum_{1 \le j \le n} (-1)^{i+j} \operatorname{pf}(a_i, \emptyset) \operatorname{pf}(\emptyset, b_j) \operatorname{pf}(a_1, \dots, \widehat{a_i}, \dots, a_n) \operatorname{pf}(b_n, \dots, \widehat{b_j}, \dots, b_1)$$

=
$$\operatorname{pf}(a_1, a_2, \dots, a_n, \emptyset) \operatorname{pf}(\emptyset, b_n, \dots, b_2, b_1).$$
 (75)

Therefore, the proof is done.

D Expansion of $pf(b_N, \ldots, b_2, b_1)$

We consider the expansion of $pf(b_N, \ldots, b_2, b_1)$ and its greatest term. We use a notation

$$\Gamma^{\mu}_{ij} = p^{\mu}_i p^{\mu-1}_j \phi_i(m+\mu) \phi_j(m+\mu-1).$$
(76)

Then, we get $\Gamma^{\mu}_{ij} > \Gamma^{\mu}_{ji}$ for i > j and $\Gamma^{\mu}_{ij} \ge \Gamma^{\nu}_{ij}$ for $\mu < \nu$ from (28). Moreover the followings also hold.

$$\Gamma_{i_{1}i_{2}}^{\mu}\Gamma_{i_{3}i_{4}}^{\mu} - \Gamma_{i_{1}i_{4}}^{\mu}\Gamma_{i_{3}i_{2}}^{\mu} = 0,
\Gamma_{i_{1}i_{2}}^{\mu+1}\Gamma_{i_{3}i_{4}}^{\mu} - \Gamma_{i_{1}i_{3}}^{\mu+1}\Gamma_{i_{2}i_{4}}^{\mu} = 0,
\phi_{i_{1}}(m)\Gamma_{i_{2}i_{3}}^{1} - \phi_{i_{3}}(m)\Gamma_{i_{2}i_{1}}^{1} = 0,$$
(77)

for $i_1, i_2, i_3, i_4 = 1, 2, ..., N$ and $\mu = 1, 2, ...$ Using this notation, $pf(b_i, b_j)$ can be expressed by

$$pf(b_{i},b_{j}) = \frac{1}{1 - p_{i}^{M} p_{j}^{M}} \sum_{\mu=1}^{M} (p_{i}^{\mu} p_{j}^{\mu-1} \phi_{i}(m+\mu) \phi_{j}(m+\mu-1) - p_{j}^{\mu} p_{i}^{\mu-1} \phi_{j}(m+\mu) \phi_{i}(m+\mu-1))$$

$$= \sum_{\nu=0}^{\infty} p_{i}^{\nu M} p_{j}^{\nu M} \sum_{\mu=1}^{M} (p_{i}^{\mu} p_{j}^{\mu-1} \phi_{i}(m+\mu) \phi_{j}(m+\mu-1) - p_{j}^{\mu} p_{i}^{\mu-1} \phi_{j}(m+\mu) \phi_{i}(m+\mu-1))$$

$$= \sum_{\mu=1}^{\infty} (p_{i}^{\mu} p_{j}^{\mu-1} \phi_{i}(m+\mu) \phi_{j}(m+\mu-1) - p_{j}^{\mu} p_{i}^{\mu-1} \phi_{j}(m+\mu) \phi_{i}(m+\mu-1))$$

$$= \sum_{\mu=1}^{\infty} (\Gamma_{ij}^{\mu} - \Gamma_{ji}^{\mu})$$
(78)

since $p_i < 1$ and $\phi_i(m+M) = \phi_i(m)$. When N is even, we obtain

$$\begin{aligned} & pf(b_{N}, b_{N-1}, \dots, b_{2}, b_{1}) \\ &= (-1)^{N/2} pf(b_{1}, b_{2}, \dots, b_{N}) \\ &= (-1)^{N/2} \sum' \operatorname{sgn} \begin{pmatrix} 1 & 2 & \dots & N \\ i_{1} & i_{2} & \dots & i_{N} \end{pmatrix} \sum_{\mu_{1}=1}^{\infty} (\Gamma_{i_{1}i_{2}}^{\mu_{1}} - \Gamma_{i_{2}i_{1}}^{\mu_{1}}) \sum_{\mu_{3}=1}^{\infty} (\Gamma_{i_{3}i_{4}}^{\mu_{3}} - \Gamma_{i_{4}i_{3}}^{\mu_{3}}) \\ &\cdots \sum_{\mu_{N-1}=1}^{\infty} (\Gamma_{i_{N-1}i_{N}}^{\mu_{N-1}} - \Gamma_{i_{N}i_{N-1}}^{\mu_{N-1}}) \\ &= (-1)^{N/2} \sum_{\mu_{1},\mu_{3},\dots,\mu_{N-1}=1}^{\infty} \sum_{\mu_{1},\mu_{3},\dots,\mu_{N-1}=1}^{\infty} \sum_{\mu_{1},\mu_{3},\dots,\mu_{N-1}=1}^{\mu_{N}} \sum_{\mu_{1},\mu_{3},\dots,\mu_{N-1}=1}^{\infty} \sum_{\mu_{1},\mu_{3},\dots,\mu_{N-1}=1}^{\mu_{N}} \sum_{\mu_{1},\mu_{2},\dots,\mu_{N-1}=1}^{\mu_{N}} \sum_{\mu_{1},\mu_{2},\dots,\mu_{N-1}=1}^{\mu_{1}} \sum_{\mu_{1},\mu_{2},\dots,\mu_{N-1}=1}^{\mu_{1}} \sum_{\mu_{1},\mu_{2},\dots,\mu_{N-1}=1}^{\mu_{1}} \sum_{\mu_{1},\mu_{2},\dots,\mu_{N-1}=1}^{\mu_{N}} \sum_{\mu_{1},\mu_{2},\dots,\mu_{N-1}=1}^{\mu_{N}} \sum_{\mu_{1},\mu_{2},\dots,\mu_{N-1}=1}^{\mu_{N}} \sum_{\mu_{1},\mu_{2},\dots,\mu_{N-1}=1}^{\mu_{N}} \sum_{\mu_{1},\mu_{2},\dots,\mu_{N-1}=1}^{\mu_{N}} \sum_{\mu_{1},\mu_{2},\dots,\mu_{N-1}=1}^{\mu_{N}} \sum_{\mu_{1},\mu_{2},\dots,\mu_{N-1}=1}^{\mu_{N}} \sum_{\mu_{1},\mu_{2},\dots,\mu_{N-1}=1}^{\mu_{N}} \sum_{\mu_{1},\mu_{2},\dots$$

where \sum'' denotes the summation over all permutations $\{i_1, i_2, \ldots, i_N\}$ of $\{1, 2, \ldots, N\}$ which satisfy the inequalities $i_1 < i_3 < \cdots < i_{N-1}$. From (77), the terms which have $\Gamma^{\mu}_{i_1i_2}\Gamma^{\mu}_{i_3i_4}$ or $\Gamma^{\mu+1}_{i_1i_2}\Gamma^{\mu}_{i_3i_4}$ in (79) vanish for $\mu = 1, 2, \ldots$ and $i_1, i_2, i_3, i_4 = 1, 2, \ldots, N$. In addition, $\Gamma^{\mu}_{i_2i_1}\Gamma^{\mu+2}_{i_4i_3} > \Gamma^{\mu}_{i_4i_3}\Gamma^{\mu+2}_{i_2i_1}$ holds for $i_1 < i_2 < i_3 < i_4$ and $\mu = 1, 2, 3, \ldots$ from (28). Due to these facts, the greatest term in (79) is given as $\Gamma^{1}_{21}\Gamma^{3}_{43}\ldots\Gamma^{N-1}_{N,N-1} = \prod_{i=1}^{N} p_i^{i-1}\phi_i(m+i-1)$. When N is odd,

$$pf(b_N, \dots, b_2, b_1) = (-1)^{(N-1)/2} pf(\emptyset, b_1, b_2, \dots, b_N)$$

$$= (-1)^{(N-1)/2} \sum_{\mu_2, \mu_4, \dots, \mu_{N-1}=1}^{\infty} \sum_{j=1}^{N} sgn \begin{pmatrix} 1 & 2 & \dots & N \\ i_1 & i_2 & \dots & i_N \end{pmatrix} \phi_{i_1} \Gamma_{i_2 i_3}^{\mu_2} \dots \Gamma_{i_{N-1} i_N}^{\mu_{N-1}}.$$
(80)

From (77), the terms which have $\phi_{i_1}\Gamma_{i_2i_3}^1$ also vanish for $i_1, i_2, i_3 = 1, 2, \ldots, N$. Thus the greatest term is given as $\phi_1(m)\Gamma_{32}^2\Gamma_{54}^4\ldots\Gamma_{N,N-1}^{N-1} = \prod_{i=1}^N p_i^{i-1}\phi_i(m+i-1)$.

References

- [1] S. Nakamura, J. Phys. A: Math. Theor. 42, 495204 (2009).
- [2] T. Tokihiro, D. Takahashi, J. Matsukidaira, and J. Satsuma, Phys. Rev. Lett. 76, 3247 (1996).
- [3] E. Date, M. Jimbo, M. Kashiwara, and T. Miwa, Publ. RIMS, Kyoto Univ. 18, 1077 (1982).
- [4] M. Jimbo and T. Miwa, Publ. RIMS, Kyoto Univ. 19, 943 (1983).
- [5] R. Hirota, Y. Ohta, and H. Nagai, Reports of RIAM Symposium No.23AO-S7, 90 (2012) [in Japanese].
- [6] S. Tsujimoto and R. Hirota, J. Phys. Soc. Jpn. 65, 2797 (1996).
- [7] R. Hirota, RIMS Kôkyûroku Bessatsu **B41**, 1 (2013) [in Japanese].
- [8] Y. Ohta, RIMS Kôkyûroku 822, 197 (1993) [in Japanese].
- [9] R. Hirota, The Direct Method in Soliton Theory (Cambridge University Press, Cambridge, 2004).