Path Category For Free* Open Morphisms From Coalgebras With Non-Deterministic Branching

Thorsten Wißmann¹[0000-0001-8993-6486], Jérémy Dubut^{2,3}, Shin-ya Katsumata², and Ichiro Hasuo^{2,4}

Friedrich-Alexander-Universität Erlangen-Nürnberg, Germany thorsten.wissmann@fau.de
National Institute of Informatics, Tokyo, Japan {dubut,s-katsumata,hasuo}@nii.ac.jp
Japanese-French Laboratory for Informatics, Tokyo, Japan
SOKENDAI, Kanagawa, Japan

Abstract. There are different categorical approaches to variations of transition systems and their bisimulations. One is coalgebra for a functor G, where a bisimulation is defined as a span of G-coalgebra homomorphism. Another one is in terms of path categories and open morphisms, where a bisimulation is defined as a span of open morphisms. This similarity is no coincidence: given a functor G, fulfilling certain conditions, we derive a path-category for pointed G-coalgebras and lax homomorphisms, such that the open morphisms turn out to be precisely the G-coalgebra homomorphisms. The above construction provides path-categories and trace semantics for free for different flavours of transition systems: (1) non-deterministic tree automata (2) regular nondeterministic nominal automata (RNNA), an expressive automata notion living in nominal sets (3) multisorted transition systems. This last instance relates to Lasota's construction, which is in the converse direction.

Keywords: Coalgebra · Open maps · Categories · Nominal Sets

1 Introduction

Coalgebras [27] and open maps [16] are two main categorical approaches to transition systems and bisimulations. The former describes the branching type of systems as an endofunctor, a system becoming a coalgebra and bisimulations being spans of coalgebra homomorphisms. Coalgebra theory makes it easy to consider state space types in different settings, e.g. nominal sets [17,18] or algebraic categories [5,11,21]. The latter, open maps, describes systems as objects of

^{*} This research was supported by ERATO HASUO Metamathematics for Systems Design Project (No. JPMJER1603), JST. The first author was supported by the DFG project MI 717/5-1. He expresses his gratitude for having been invited to Tokyo, which initiated the present work.

this paper	worlds	data	systems	func. sim.	func. bisim.	(bi)simulation	•
	open maps	$J \colon \mathbb{P} \longrightarrow \mathbb{M}$ Def. 2.4	$\mathbf{obj}\left(\mathbb{M}\right)$	$\mathbf{mor}\left(\mathbb{M}\right)$	open maps Def. 2.5	- 33/ /36	Lasota's
	coalgebra	$G \colon \mathbb{C} \longrightarrow \mathbb{C}$ Def. 2.7	pointed G-coalg. Sec. 2.2	lax hom. Def. 2.8	coalg. hom. Def. 2.6		

Table 1. Two approaches to categorical (bi)simulations

a category and the execution types as particular objects called paths. In this case, bisimulations are spans of open morphisms. Open maps are particularly adapted to extend bisimilarity to history dependent behaviors, e.g. true concurrency [8,7], timed systems [24] and weak (bi)similarity [9]. Coalgebra homomorphisms and open maps are then key concepts to describe bisimilarity categorically. They intuitively correspond to functional bisimulations, that is, those maps between states whose graph is a bisimulation.

We are naturally interested in the relationship between those two categorical approaches to transition systems and bisimulations. A reduction of open maps situations to coalgebra was given by Lasota using multi-sorted transition systems [20]. In this paper, we give the reduction in the other direction: from the category $\mathsf{Coalg}_l(TF)$ of pointed TF-coalgebras and lax homomorphisms, we construct the path-category Path and a functor $J:\mathsf{Path} \longrightarrow \mathsf{Coalg}_l(TF)$ such that Path -open morphisms coincide with strict homomorphisms, hence functional bisimulations. Here, T is a functor describing the branching behaviour and F describes the input type, i.e. the type of data that is processed (e.g. words or trees). This development is carried out with the case where T is a powerset-like functor, and covers transition systems allowing non-deterministic branching.

The key concept in the construction of Path are F-precise maps. Roughly speaking in set, a map $f \colon X \longrightarrow FY$ is F-precise if every $y \in Y$ is used precisely once in f, i.e. there is a unique x such that y appears in f(x) and additionally y appears precisely once in f(x). Such an F-precise map represents one deterministic step (of shape F). Then a path $P \in P$ ath is a finite sequence of deterministic steps, i.e. finitely many precise maps. J converts such a data into a pointed TF-coalgebra. There are many existing notions of paths and traces in coalgebra [4,12,13,22], which lack the notion of precise map, which is crucial for the present work.

Once we set up the situation J: Path \longrightarrow Coalg $_l(TF)$, we are on the framework of open map bisimulations. Our construction of Path using precise maps is justified by the characterisation theorem: Path-open morphisms and strict coalgebra homomorphisms coincide (Theorem 3.20 and Theorem 3.24). This coincidence relies on the concept of path-reachable coalgebras, namely, coalgebras such that every state can be reached by a path. Under mild conditions, path-reachability is equivalent to an existing notion in coalgebra, defined as the non-existence of a proper sub-coalgebra (Section 3.5). Additionally, this characterization produces a canonical trace semantics for free, given in terms of paths (Section 3.6).

We illustrate our reduction with several concrete situations: different classes of non-deterministic top-down tree automata using analytic functors (Section 4.1), Regular Nondeterministic Nominal Automata (RNNA), an expressive automata notion living in nominal sets (Section 4.2), multisorted transition systems, used in Lasota's work to construct a coalgebra situation from an open map situation (Section 4.3).

Notation. We assume basic categorical knowledge and notation (see e.g. [1,3]). The cotupling of morphisms $f: A \to C$, $g: B \to C$ is denoted by $[f,g]: A+B \to C$, and the unique morphism to the terminal object is $!: X \to 1$ for every X.

2 Two categorical approaches for bisimulations

We introduce the two formalisms involved in the present paper: the open maps (Section 2.1) and the coalgebras (Section 2.2). Those formalisms will be illustrated on the classic example of Labelled Transition Systems (LTSs).

Definition 2.1. Fix a set A, called the alphabet. A labelled transition system is a triple (S, i, Δ) with S a set of states, $i \in S$ the initial state, and $\Delta \subseteq S \times A \times S$ the transition relation. When Δ is obvious from the context, we write $s \stackrel{a}{\longrightarrow} s'$ to mean $(s, a, s') \in \Delta$.

For instance, the tuple $(\{0, \dots, n\}, 0, \{(k-1, a_k, k) \mid 1 \le k \le n\})$ is an LTS, and called the *linear system* over the word $a_1 \cdots a_n \in A^*$. To relate LTSs, one considers functions that preserves the structure of LTSs:

Definition 2.2. A morphism of LTSs from (S, i, Δ) to (S', i', Δ') is a function $f: S \longrightarrow S'$ such that f(i) = i' and for every $(s, a, s') \in \Delta$, $(f(s), a, f(s')) \in \Delta'$. LTSs and morphisms of LTSs form a category, which we denote by LTS_A.

Some authors choose other notions of morphisms (e.g. [16]), allowing them to operate between LTSs with different alphabets for example. The usual way of comparing LTSs is by using simulations and bisimulations [25]. The former describes what it means for a system to have at least the behaviours of another, the latter describes that two systems have exactly the same behaviours. Concretely:

Definition 2.3. A simulation from (S, i, Δ) to (S', i', Δ') is a relation $R \subseteq S \times S'$ such that (1) $(i, i') \in R$, and (2) for every $s \stackrel{a}{\longrightarrow} t$ and $(s, s') \in R$, there is $t' \in S'$ such that $s' \stackrel{a}{\longrightarrow} t'$ and $(t, t') \in R$. Such a relation R is a bisimulation if $R^{-1} = \{(s', s) \mid (s, s') \in R\}$ is also a simulation.

Morphisms of LTSs are functional simulations, i.e. functions between states whose graph is a simulation. So how to model (1) systems, (2) functional simulations and (3) functional bisimulations categorically? In the next two sections, we will describe known answers to this question, with open maps and coalgebra. In both cases, it is possible to capture similarity and bisimilarity of two LTSs T and T'. Generally, a simulation is a (jointly monic) span of a functional bisimulation

and a functional simulation, and a bisimulation is a simulation whose converse is also a simulation, as depicted in Table 1. Consequently, to understand similarity and bisimilarity on a general level, it is enough to understand functional simulations and bisimulations.

2.1 Open maps

The categorical framework of open maps [16] assumes functional simulations to be already modeled as a category \mathbb{M} . For example, for $\mathbb{M} := \mathsf{LTS}_A$, objects are LTSs, and morphisms are functional simulations. Furthermore, the open maps framework assumes another category \mathbb{P} of 'paths' or 'linear systems', together with a functor J that tells how a 'path' is to be understood as a system:

Definition 2.4 [16]. An open map situation is given by categories \mathbb{M} ('systems' with 'functional simulations') and \mathbb{P} ('paths') together with a functor $J : \mathbb{P} \to \mathbb{M}$.

For example with $\mathbb{M} := \mathsf{LTS}_A$, we pick $\mathbb{P} := (A^*, \leq)$ to be the poset of words over A with prefix order. Here, the functor J maps a word $w \in A^*$ to the linear system over w, and $w \leq v$ to the evident functional simulation $J(w \leq v) : Jw \longrightarrow Jv$.

In an open map situation $J \colon \mathbb{P} \longrightarrow \mathbb{M}$, we can abstractly represent the concept of a run in a system. A run of a path $w \in \mathbb{P}$ in a system $T \in \mathbb{M}$ is simply defined to be an \mathbb{M} -morphism of type $Jw \longrightarrow T$. With this definition, each \mathbb{M} -morphism $h \colon T \longrightarrow T'$ (i.e. functional simulation) inherently transfers runs: given a run $x \colon Jw \longrightarrow T$, the morphism $h \cdot x \colon Jw \longrightarrow T'$ is a run of w in T'. In the example open map situation $J \colon (A^*, \leq) \longrightarrow \mathsf{LTS}_A$, a run of a path $w = a_1 \cdots a_n \in A^*$ in an LTS $T = (S, i, \Delta)$ is nothing but a sequence of states $x_0, \ldots, x_n \in S$ such that $x_0 = i$ and $x_{k-1} \xrightarrow{a_k} x_k$ holds for all $1 \le k \le n$.

We introduce the concept of open map [16]. This is an abstraction of the property posessed by functional bisimulations. For LTSs $T=(S,i,\Delta)$ and $T'=(S',i',\Delta')$, an LTS_A-morphism $h\colon T\longrightarrow T'$ is a functional bisimulation if the graph of h is a bisimulation. This implies the following relationship between runs in T and runs in T'. Suppose that $w\leq w'$ holds in A^* , and a run x of w in T is given as in (1); here n,m are lengths of w,w' respectively. Then for any run y' of w' in T' extending $h\cdot x$ as in (2), there is a run x' of w' extending x, and moreover its image by h coincides with y' (that is, $h\cdot x'=y'$). Such x' is obtained by repetitively applying the condition of functional bisimulation.

$$\rightarrow \underbrace{i \xrightarrow{w_1} x_1 \xrightarrow{w_2} \cdots \xrightarrow{w_n} x_n}^{x} \xrightarrow{w'_{n+1}} x'_{n+1} \xrightarrow{w'_{n+2}} \cdots \xrightarrow{w'_m} x'_m \quad (\text{in } T) \quad (1)$$

$$\rightarrow \underbrace{i' \xrightarrow{w_1} h(x_1) \xrightarrow{w_2} \cdots \xrightarrow{w_n} h(x_n) \xrightarrow{w'_{n+1}} y'_{n+1} \xrightarrow{w_{n+2}} \cdots \xrightarrow{w'_m} y'_m}_{i'} \quad (\text{in } T') \quad (2)$$

Observe that y' extending $h \cdot x$ can be represented as $y' \cdot J(w \leq w') = h \cdot x$, and x' extending x as $x' \cdot J(w \leq w') = x$. From these, we conclude that if an LTS_A-morphism $h \colon T \longrightarrow T'$ is a functional bisimulation, then for any $w \leq w'$

in A^* and run $x: Jw \longrightarrow T$ and $y': Jw' \longrightarrow T'$ such that $y' \cdot J(w \leq w') = h \cdot x$, there is a run $x': Jw' \longrightarrow T$ such that $x' \cdot J(w \leq w') = x$ and $h \cdot x' = y'$ (the converse also holds if all states of T are reachable). This necessary condition of functional bisimulation can be rephrased in any open map situation, leading us to the definition of open map.

Definition 2.5 [16]. Let $J: \mathbb{P} \longrightarrow \mathbb{M}$ be an open map situation. An \mathbb{M} -morphism $h: T \longrightarrow T'$ is said to be open if for every morphism $\Phi: w \longrightarrow w' \in \mathbb{P}$ making the square on the right commute, there is x' making the two triangles commute. $Jw \xrightarrow{x} T$ $\downarrow h$ $Jw' \xrightarrow{y'} T'$

Open maps are closed under composition and stable under pullback [16].

2.2 Coalgebras

The theory of G-coalgebras is another categorical framework to study bimulations. The type of systems is modelled using an endofunctor $G \colon \mathbb{C} \longrightarrow \mathbb{C}$ and a system is then a coalgebra for this functor, that is, a pair of an object S of \mathbb{C} (modeling the state space), and of a morphism of type $S \longrightarrow GS$ (modeling the transitions). For example for LTSs, the transition relation is of type $\Delta \subseteq S \times A \times S$. Equivalently, this can be defined as a function $\Delta \colon S \longrightarrow \mathcal{P}(A \times S)$, where \mathcal{P} is the powerset. In other words, the transition relation is a coalgebra for the Set-functor $\mathcal{P}(A \times _)$. Intuitively, this coalgebra gives the one-step behaviour of an LTS: S describes the state space of the system, \mathcal{P} describes the 'branching type' as being non-deterministic, $A \times S$ describe the 'computation type' as being linear, and the function itself lists all possible futures after one-step of computation of the system. Now, changing the underlying category or the endofunctor allows to model different types of systems. This is the usual framework of coalgebra, as described for example in [27].

Initial states are modelled coalgebraically by a pointing to the carrier $i: I \longrightarrow S$ for a fixed object I in \mathbb{C} , describing the 'type of initial states' (see e.g. [2, Sec. 3B]). For example, an initial state of an LTS is the same as a function from the singleton set $I := \{*\}$ to the state space S. This object I will often be the final object of \mathbb{C} , but we will see other examples later. In total, an I-pointed G-coalgebra is a \mathbb{C} -object S together with morphisms $\alpha: S \longrightarrow GS$ and $i: I \longrightarrow S$. E.g. an LTS is an I-pointed G-coalgebra for $I = \{*\}$ and $GX = \mathcal{P}(A \times X)$.

In coalgebra, functional bisimulations are the first class citizens to be modelled as homomorphisms. The intuition is that those preserve the initial state, and preserve and reflect the one-step relation.

Definition 2.6. An I-pointed G-coalgebra homomorphism $I \xrightarrow{i} S \xrightarrow{\alpha} GS$ from $I \xrightarrow{i} S \xrightarrow{\alpha} GS$ to $I \xrightarrow{i'} S' \xrightarrow{\alpha'} GS'$ is a morphism $f: S \longrightarrow S'$ making the right-hand diagram commute. $I \xrightarrow{i'} S \xrightarrow{\alpha'} GS'$

For instance, when $G = \mathcal{P}(A \times _)$, one can easily see that a function f is a G-coalgebra homomorphism iff it is a functional bisimulation. Thus, if we want to capture functional simulations in LTSs, we need to weaken the condition of homomorphism to the inequality $Gf(\alpha(s)) \subseteq \alpha'(f(s))$ (instead of equality). To

express this condition for general G-coalgebras, we introduce a partial order $\sqsubseteq_{X,Y}$ on each homset $\mathbb{C}(X,GY)$ in a functorial manner.

Definition 2.7. A partial order on G-homsets is a functor $\sqsubseteq : \mathbb{C}^{\mathsf{op}} \times \mathbb{C} \longrightarrow \mathsf{Pos}$ such that $U \cdot \sqsubseteq = \mathbb{C}(_, G_-)$; here, $U \colon \mathsf{Pos} \longrightarrow \mathsf{Set}$ is the forgetful functor from the category Pos of posets and monotone functions.

The functoriality of \sqsubseteq amounts to that $f_1 \sqsubseteq f_2$ implies $Gh \cdot f_1 \cdot g \sqsubseteq Gh \cdot f_2 \cdot g$.

Definition 2.8. Given a partial order on G-homsets, an I-pointed lax G-coalgebra homomorphism $f: (S, \alpha, i) \longrightarrow I \xrightarrow{i} S \xrightarrow{\alpha} GS$ (S', α', i') is a morphism $f: S \longrightarrow S'$ making the right-hand diagram commute. The I-pointed G-coalgebras and lax homomorphisms form a category, denoted by $Coalg_I(I, G)$.

Conclusion 2.9. In Set, with $I = \{*\}$, $G = \mathcal{P}(A \times _)$, define the order $f \sqsubseteq g$ in $\mathsf{Set}(X, \mathcal{P}(A \times Y))$ iff for every $x \in X$, $f(x) \subseteq g(x)$. Then $\mathsf{Coalg}_l(\{*\}, \mathcal{P}(A \times _)) = \mathsf{LTS}_A$. In particular, we have an open map situation

$$\mathbb{P} = (A^{\star}, \leq) \quad \overset{J}{\longrightarrow} \quad \mathbb{M} = \mathsf{LTS}_A = \mathsf{Coalg}_l(\{*\}, \mathcal{P}(A \times _))$$

and the open maps are precisely the coalgebra homomorphisms (for reachable LTSs). In this paper, we will construct a path category \mathbb{P} for more general I and G, such that the open morphisms are precisely the coalgebra homomorphisms.

3 The open map situation in coalgebras


Lasota's construction [20] transforms an open map situation $J \colon \mathbb{P} \longrightarrow \mathbb{M}$ into a functor G (with a partial order on G-homsets), together with a functor $\operatorname{Beh} \colon \mathbb{M} \longrightarrow \operatorname{Coalg}_l(I,G)$ that sends open maps to G-coalgebra homomorphisms (see section 4.3 for details). In this paper, we provide a construction in the converse direction for functors G of a certain shape.

As exemplified by LTSs, it is a common pattern that G is the composition G = TF of two functors [12], where T is the branching type (e.g. partial, or non-deterministic) and F is the data type, or the 'linear behaviour' (words, trees, words modulo α -equivalence). If we instantiate our path-construction to $T = \mathcal{P}$ and $F = A \times$, we obtain the known open map situation for LTSs (Conclusion 2.9).

Fix a category $\mathbb C$ with pullbacks, functors $T,F:\mathbb C\longrightarrow\mathbb C$, an object $I\in\mathbb C$ and a partial order \sqsubseteq^T on T-homsets. They determine a coalgebra situation $(\mathbb C,I,TF,\sqsubseteq)$ where \sqsubseteq is the partial order on TF-homsets defined by $\sqsubseteq_{X,Y}=\sqsubseteq_{X,FY}^T$. Under some conditions on T and F, we construct a path-category $\mathsf{Path}(I,F+1)$ and an open map situation $\mathsf{Path}(I,F+1)\hookrightarrow \mathsf{Coalg}_l(I,TF)$ where TF-coalgebra homomorphisms and $\mathsf{Path}(I,F+1)$ -open morphisms coincide.

3.1 Precise morphisms

While the path category is intuitively clear for $FX = A \times X$, it is not for inner functors F that model tree languages. For example for $FX = A + X \times X$, a $\mathcal{P}F$ -coalgebra models transition systems over binary trees with leaves labelled in A,






Fig. 1. A non-precise map f that factors through the F-precise $f': X \longrightarrow Y' \times Y' + \{\bot\}$

instead of over words. Hence, the paths should be these kind of binary trees. We capture the notion of tree like shape ("every node in a tree has precisely one route to the root") by the following abstract definition:

Definition 3.1. For a functor $F: \mathbb{C} \longrightarrow \mathbb{C}$, a morphism $s: S \longrightarrow FR$ is called F-precise if for all f, g, h the following implication holds:

Remark 3.2. If F preserves weak pullbacks, then a morphism s is F-precise iff it fulfils the above definition for q = id.

Example 3.3. Intuitively speaking, for a polynomial Set-functor F, a map $s: S \to \mathbb{R}$ FR is F-precise iff every element of R is mentioned precisely once in the definition of the map f. For example, for $FX = A \times X + \{\bot\}$, the case needed later for LTSs, a map $f: X \longrightarrow FY$ is precise iff for every $y \in Y$, there is a unique pair $(x,a) \in X \times A$ such that f(x) = (a,y). For $FX = X \times X + \{\bot\}$ on Set, the map $f: X \longrightarrow FY$ in Figure 1 is not F-precise, because y_2 is used three times (once in $f(x_2)$ and twice in $f(x_3)$, and y_3 and y_4 do not occur in f at all. However, $f': X \longrightarrow FY'$ is F-precise because every element of Y' is used precisely once in f', and we have that $Fh \cdot f' = f$. Also note that f' defines a forest where X is the set of roots, which is closely connected to the intuition that, in the F-precise map f', from every element of Y', there is precisely one edge up to a root in X.

So when transforming a non-precise map into a precise map, one duplicates elements that are used multiple times and drops elements that are not used. We will cover functors F for which this factorization pattern provides F-precise maps. If F involves unordered structure, this factorization needs to make choices, and so we restrict the factorization to a class \mathcal{S} of objects that have that choiceprinciple (see Example 4.5 later):

Definition 3.4. Fix a class of objects $S \subseteq \text{obj} \mathbb{C}$ closed un- $S \xrightarrow{\exists f'} FY'$ der isomorphism. We say that F admits precise factorizations w.r.t. S if for every $f: S \to FY$ with $S \in S$, there exist $Y' \in S$, $h: Y' \to Y$ and $f': S \to FY'$ F-precise with $Fh \cdot f' = f$.

$$S \xrightarrow{\exists f'} FY'$$

$$\forall f \qquad \downarrow_{Fh}$$

$$FY$$




Fig. 2. A path of length 4 for $FX = \{a\} \times X + X \times X + \{\bot\}$ with $I = \{*\}$.

For $\mathbb{C} = \mathsf{Set}$, \mathcal{S} contains all sets. However for the category of nominal sets, \mathcal{S} will only contain the strong nominal sets (see details in subsection 4.2).

Remark 3.5. Precise morphisms are essentially unique. If $f_1: X \longrightarrow FY_1$ and $f_2: X \longrightarrow FY_2$ are F-precise and if there is some $h: Y_1 \longrightarrow Y_2$ with $Fh \cdot f_1 = f_2$, then h is an isomorphism. Consequently, if $f: S \longrightarrow FY$ with $S \in \mathcal{S}$ is F-precise and F-admits precise factorizations, then $Y \in \mathcal{S}$.

Functors admitting precise factorizations are closed under basic constructions:

Proposition 3.6. The following functors admit precise factorizations w.r.t. S:

- 1. Constant functors, if \mathbb{C} has an initial object 0 and $0 \in \mathcal{S}$.
- 2. $F \cdot F'$ if $F : \mathbb{C} \longrightarrow \mathbb{C}$ and $F' : \mathbb{C} \longrightarrow \mathbb{C}$ do so.
- 3. $\prod_{i \in I} F_i$, if all $(F_i)_{i \in I}$ do so and S is closed under I-coproducts.
- 4. $\coprod_{i\in I} F_i$, if all $(F_i)_{i\in I}$ do so, $\mathbb C$ is I-extensive and $\mathcal S$ is closed under I-coproducts.
- 5. Right-adjoint functors, if and only if its left-adjoint preserves $\mathcal{S}\text{-objects}$.

Example 3.7. When $\mathbb C$ is infinitary extensive and $\mathcal S$ is closed under coproducts, every polynomial endofunctor $F\colon\mathbb C\longrightarrow\mathbb C$ admits precise factorizations w.r.t. $\mathcal S$. This is in particular the case for $\mathbb C=\mathcal S=\mathsf{Set}$. In this case, we shall see later (Section 4.1) that many other Set-functors, e.g. the bag functor $\mathcal B$, where $\mathcal B(X)$ is the set of finite multisets, have precise factorizations. In contrast, $F=\mathcal P$ does not admit precise factorizations, and if $f\colon X\longrightarrow \mathcal PY$ is $\mathcal P$ -precise, then $f(x)=\emptyset$ for all $x\in X$.

3.2 Path categories in pointed coalgebras

We define a path for I-pointed TF-coalgebras as a tree according to F. Following the observation in Example 3.3, one layer of the tree is modelled by a F-precise morphism and hence a path in a TF-coalgebra is defined to be a finite sequence of (F+1)-precise maps, where the $_++1$ comes from the dead states w.r.t. T; the argument is given later in Remark 3.23 when reachability is discussed. Since the $_++1$ is not relevant yet, we define $\mathsf{Path}(I,F)$ in the following and will use $\mathsf{Path}(I,F+1)$ later. For simplicity, we write X_n for finite families $(X_k)_{0 \le k \le n}$.

Definition 3.8. The category Path(I,F) consists of the following. An object is $(\mathbf{P}_{n+1},\mathbf{p}_n)$ for an $n \in \mathbb{N}$ with $P_0 = I$ and \mathbf{p}_n a family of F-precise maps $(p_k \colon P_k \longrightarrow FP_{k+1})_{k < n}$. We say that $(\mathbf{P}_{n+1},\mathbf{p}_n)$ is a path of length n. A morphism $\phi_{n+1} \colon (\mathbf{P}_{n+1},\mathbf{p}_n) \longrightarrow (\mathbf{Q}_{m+1},\mathbf{q}_m)$, $m \ge n$, is a family $(\phi_k \colon P_k \longrightarrow Q_k)_{k \le n}$ with $\phi_0 = \operatorname{id}_I$ and $q_k \cdot \phi_k = F\phi_{k+1} \cdot p_k$ for all $0 \le k \le n$.

Example 3.9. Paths for $FX = A \times X + 1$ and $I = \{*\}$ singleton are as follows. First, a map $f : I \longrightarrow FX$ is precise iff (up-to isomorphism) either X = I and f(*) = (a, *) for some $a \in A$; or $X = \emptyset$ and $f(*) = \bot$. Then a path is isomorphic to an object of the form: $P_i = I$ for $i \le k$, $P_i = \emptyset$ for i > k, $p_i(*) = (a_i, *)$ for i < k, and $p_k(*) = \bot$. A path is the same as a word, plus some "junk", concretely, a word in $A^*.\bot^*$. For LTSs, an object in Path(I,F) with $FX = A \times X$ is simply a word in A^* . For a more complicated functor, Figure 2 depicts a path of length 4, which is a tree for the signature with one unary, one binary symbol, and a constant. The layers of the tree are the sets P_4 . Also note that since every p_i is F-precise, there is precisely one route to go from every element of a P_k to *.

Remark 3.10. The inductive continuation of Remark 3.5 is as follows. Given a morphism ϕ_{n+1} in Path(I, F), since ϕ_0 is an isomorphism, then ϕ_k is an isomorphism for all $0 \le k \le n$. If F admits precise factorizations and if $I \in \mathcal{S}$, then for every path (P_{n+1}, p_n) , all P_k , $0 \le k \le n$, are in \mathcal{S} .

Remark 3.11. If in Definition 3.4, the connecting morphism $h: Y' \longrightarrow Y$ uniquely exists, then it follows by induction that the hom-sets of Path(I, F) are at most singleton. This is the case for all polynomial functors, but not the case for the bag functor on sets (discussed in subsection 4.1).

Definition 3.12. The path poset PathOrd(I, F) is the set $F^nF^{m-n}1$ $\coprod_{0 \le n} \mathbb{C}(I, F^n1)$ equipped with the order: for $u: I \longrightarrow F^n1$ and $v: I \longrightarrow F^m1$, we define $u \le v$ if $n \le m$ and $F^n(!) \cdot v = u$. $I \xrightarrow{u} F^n1$

So $u \le v$ if u is the truncation of v to n levels. This matches the morphisms in $\mathsf{Path}(I,F)$ that witnesses that one path is prefix of another:

Proposition 3.13. 1. The functor Comp: Path $(I, F) \longrightarrow$ PathOrd(I, F) defined by $I = P_0 \stackrel{p_0}{\longrightarrow} FP_1 \cdots \longrightarrow F^nP_n \stackrel{F^n!}{\longrightarrow} F^n1$ on $(\mathbf{P}_{n+1}, \mathbf{p}_n)$ is full, and reflects isos. 2. If F admits precise factorizations w.r.t. S and $I \in S$, then Comp is sujective. 3. If additionally h in Def. 3.4 is unique, then Comp has a right-inverse.

In particular, $\mathsf{PathOrd}(I,F)$ is $\mathsf{Path}(I,F)$ up to isomorphism. In the instances, it is often easier to characterize $\mathsf{PathOrd}(I,F)$. This also shows that $\mathsf{Path}(I,F)$ contains the elements – understood as morphisms from I – of the finite start of the final chain of $F\colon 1\stackrel{!}{\leftarrow} F1 \stackrel{F!}{\leftarrow} F^21 \stackrel{F^2!}{\leftarrow} F^31 \leftarrow \cdots$.

Example 3.14. When $FX = A \times X + 1$, $F^n 1$ is isomorphic to the set of words in $A^*.\bot^*$ of length n. Consequently, $\mathsf{PathOrd}(I,F)$ is the set of words in $A^*.\bot^*$, equipped with the prefix order. In this case, Comp is an equivalence of categories.

3.3 Embedding paths into pointed coalgebras

The paths (P_{n+1}, p_n) embed into $Coalg_l(I, TF)$ as one expects it for examples like Figure 2: one takes the disjoint union of the P_k , one has the pointing $I = P_0$ and the linear structure of F is embedded into the branching type T.

During the presentation of the results, we require T, F, and I to have certain properties, which will be introduced one after the other. The full list of assumptions is summarized in Table 2:

 $(\operatorname{Ax1})$ – The main theorem will show that coalgebra homomorphisms in $\operatorname{\mathsf{Coalg}}_l(I,TF)$ are the open maps for the path category $\operatorname{\mathsf{Path}}(I,F+1)$. So from now on, we assume that $\mathbb C$ has finite coproducts and to use the results from the previous sections, we fix a class $\mathcal S\subseteq\operatorname{\mathbf{obj}}\mathbb C$ such that F+1 admits precise factorizations w.r.t. $\mathcal S$ and that $I\in\mathcal S$.

 $(\operatorname{Ax2})$ – Recall, that a family of morphisms $(e_i\colon X_i\longrightarrow Y)_{i\in I}$ with common codomain is called jointly epic if for $f,g\colon Y\longrightarrow Z$ we have that $f\cdot e_i=g\cdot e_i\ \forall i\in I$ implies f=g. For Set, this means, that every element $g\in Y$ is in the image of some e_i . Since we work with partial orders on T-homsets, we also need the generalization of this property if $f\sqsubseteq g$ are of the form $Y\longrightarrow TZ'$.

(Ax3) – In this section, we encode paths as a pointed coalgebra by constructing a functor $J: \mathsf{Path}(I, F+1) \hookrightarrow \mathsf{Coalg}_l(I, TF)$. For that we need to embed the linear behaviour FX+1 into TFX. This is done by a natural transformation $[\eta, \bot]: \mathrm{Id} + 1 \longrightarrow T$, and we require that $\bot: 1 \longrightarrow T$ is a bottom element for \sqsubseteq .

Example 3.15. For the case where T is the powerset functor \mathcal{P} , η is given by the unit $\eta_X(x) = \{x\}$, and \bot is given by empty sets $\bot_X(*) = \emptyset$.

Definition 3.16. We have an inclusion functor J: Path $(I, F+1) \hookrightarrow \mathsf{Coalg}_l(I, TF)$ that maps a path $(\mathbf{P}_{n+1}, \mathbf{p}_n)$ to an I-pointed TF-coalgebra on $\coprod \mathbf{P}_{n+1} := \coprod_{0 \le k \le n} P_k$. The pointing is given by $\mathsf{in}_0 \colon I = P_0 \longrightarrow \coprod \mathbf{P}_{n+1}$ and the structure by:

$$\coprod_{0 \le h \le n} P_k + P_n \xrightarrow{[(F \mathsf{in}_{k+1} + 1) \cdot p_k]_{0 \le k < n} + !} F \coprod \mathbf{P}_{n+1} + 1 \xrightarrow{[\eta, \bot]} TF \coprod \mathbf{P}_{n+1}.$$

Example 3.17. In the case of LTSs, a path, or equivalently a word $a_1...a_k.\perp...\perp \in A^*.\perp^*$, is mapped to the finite linear system over $a_1...a_k$ (see Section 2.1), seen as a coalgebra (see Section 2.2).

Proposition 3.18. Given a morphism $[x_k]_{k \leq n} : \coprod P_{n+1} \longrightarrow X$ for some system (X, ξ, x_0) and a path (P_{n+1}, p_n) , we have

$$J(\boldsymbol{P}_{n+1},\boldsymbol{p}_n) \xrightarrow{[x_k]_{k \leq n}} (X,\xi,x_0) \iff \forall k < n \colon \Pr_{k\downarrow} \xrightarrow{Fx_{k+1}+1} \sqsubseteq \underset{[\eta,\bot]_X}{\sqsubseteq} \downarrow \xi \\ FP_{k+1}+1 \longrightarrow FX+1 \longrightarrow TFX.$$

Also note that the pointing x_0 of the coalgebra is necessarily the first component of any run in it. In a run $[x_k]_{k < n}$, p_k corresponds to an edge from x_k to x_{k+1} .

Example 3.19. For LTSs, since the P_k are singletons, x_k just picks the kth state of the run. The right-hand side of this lemma describes that this is a run iff there is a transition from the kth state and the (k+1)—th state.

3.4 Open morphisms are exactly coalgebra homomorphisms

In this section, we prove our main contribution, namely that $\mathsf{Path}(I, F+1)$ open maps in $\mathsf{Coalg}_l(I, TF)$ are exactly coalgebra homomorphisms. For the first




Table 2. Main assumptions on $F, T: \mathbb{C} \longrightarrow \mathbb{C}, \sqsubseteq^T, S \subseteq \mathbf{obj} \mathbb{C}$

direction of the main theorem, that is, that coalgebra homomorphisms are open, we need two extra axioms:

(Ax4) – describing that the order on $\mathbb{C}(X,TY)$ is point-wise. This holds for the powerset because every set is the union of its singleton subsets.

(Ax5) – describing that $\mathbb{C}(X,TY)$ admits a choice-principle. This holds for the powerset because whenever $y \in h[x]$ for a map $h: X \longrightarrow Y$ and $x \subseteq X$, then there is some $\{x'\} \subseteq x$ with h(x') = y.

Theorem 3.20. Under the assumptions of Table 2, a coalgebra homomorphism in $Coalg_l(I, TF)$ is Path(I, F + 1)-open.

The converse is not true in general, because intuitively, open maps reflect runs, and thus only reflect edges of reachable states, as we have seen in Section 2.1. The notion of a state being reached by a path is the following:

Definition 3.21. A system (X, ξ, x_0) is path-reachable if the family of runs $[x_k]_{k \le n} : J(\mathbf{P}_{n+1}, \mathbf{p}_n) \longrightarrow (X, \xi, x_0)$ (of paths from $\mathsf{Path}(I, F+1)$) is jointly epic.

Example 3.22. For LTSs, this means that every state in X is reached by a run, that is, there is a path from the initial state to every state of X.

Remark 3.23. In Definition 3.21, it is crucial that we consider $\mathsf{Path}(I, F+1)$ and not $\mathsf{Path}(I, F)$ for functors incorporating 'arities ≥ 2 '. This does not affect the example of LTSs, but for I=1, $FX=X\times X$ and $T=\mathcal{P}$ in Set, the coalgebra (X,ξ,x_0) on $X=\{x_0,y_1,y_2,z_1,z_2\}$ given by $\xi(x_0)=\{(y_1,y_2)\},\ \xi(y_1)=\{(z_1,z_2)\},\ \xi(y_2)=\xi(z_1)=\xi(z_2)=\emptyset$ is path-reachable for $\mathsf{Path}(I,F+1)$. There is no run of a length 2 path from $\mathsf{Path}(I,F)$, because y_2 has no successors, and so there is no path to z_1 or to z_2 .

Theorem 3.24. Under the assumptions of Table 2, if (X, ξ, x_0) is path-reachable, then an open morphism $h: (X, \xi, x_0) \longrightarrow (Y, \zeta, y_0)$ is a coalgebra homomorphism.

3.5 Connection to other notions of reachability

There is another concise notion for reachability in the coalgebraic literature [2].

Definition 3.25. A subcoalgebra of (X, ξ, x_0) is a coalgebra homomorphism $h: (Y, \zeta, y_0) \longrightarrow (X, \xi, x_0)$ that is carried by a monomorphism $h: X \rightarrowtail Y$. Furthermore (X, ξ, x_0) is called reachable if it has no proper subcoalgebra, i.e. if any subcoalgebra h is an isomorphism.

Under the following assumptions, this notion coincides with the path-based definition of reachability (Definition 3.21).

Assumption 3.26. For the present subsection 3.5, let \mathbb{C} be cocomplete, have (epi,mono)-factorizations and wide pullbacks of monomorphisms.

The first direction follows directly from Theorem 3.20:

Proposition 3.27. Every path-reachable (X, ξ, x_0) has no proper subcoalgebra.

For the other direction it is needed that TF preserves arbitrary intersections, that is, wide pullbacks of monomorphisms. In Set, this means that for a family $(X_i \subseteq Y)_{i \in I}$ of subsets we have $\bigcap_{i \in I} TFX_i = TF \bigcap_{i \in I} X_i$ as subsets of TFY.

Proposition 3.28. If, furthermore, for every monomorphism $m: Y \longrightarrow Z$, the function $\mathbb{C}(-,Tm)\colon \mathbb{C}(X,TY) \longrightarrow \mathbb{C}(X,TZ)$ reflects joins and if TF preserves arbitrary intersections, then a reachable coalgebra (X,ξ,x_0) is also path-reachable.

All those technical assumptions are satisfied in the case of LTSs, and will also be satisfied in all our instances in section 4.

3.6 Trace semantics for pointed coalgebras

The characterization from Theorem 3.20 and Theorem 3.24 points out a natural way of defining a trace semantics for pointed coalgebras. Indeed, the paths category $\mathsf{Path}(I,F+1)$ provides a natural way of defining the runs of a system. A possible way to go from runs to trace semantics is to describe accepting runs as the subcategory $J' \colon \mathsf{Path}(I,F) \hookrightarrow \mathsf{Path}(I,F+1)$. We can define the *trace semantics* of a system (X,ξ,x_o) as the set:

$$\operatorname{tr}(X,\xi,x_0) = \{\operatorname{\mathsf{Comp}}(\boldsymbol{P}_{n+1},\boldsymbol{p}_n) \mid \exists \operatorname{run} [x_k]_{k \leq n} \colon JJ'(\boldsymbol{P}_{n+1},\boldsymbol{p}_n) \longrightarrow (X,\xi,x_0)$$
 with $(\boldsymbol{P}_{n+1},\boldsymbol{p}_n) \in \operatorname{\mathsf{Path}}(I,F)\}$

Since Path(I, F)-open maps preserve and reflect runs, we have the following:

Let us look at two LTS-related examples (we will describe some others in the next section). First, for $FX = A \times X$. The usual trace semantics is given by all the words in A^* that are labelled of a run of a system. This trace semantics is obtained because $\mathsf{PathOrd}(I,F) = \coprod_{n \geq 0} A^n$ and because Comp maps every path to its underlying word. Another example is given for $FX = A \times X + \{ \checkmark \}$, where \checkmark marks final states. In this case, a path in $\mathsf{Path}(I,F)$ of length n is either a path that can still be extended or encodes less than n steps to an accepting state \checkmark . This obtains the trace semantics containing the set of accepted words, as in automata theory, plus the set of possibly infinite runs.

4 Instances

4.1 Analytic functors and tree automata

In Example 3.7, we have seen that every polynomial Set-functors, in particular the functor $X \mapsto A \times X$, has precise factorizations with respect to all sets. This allowed us to see LTSs, modelled as $\{*\}$ -pointed $\mathcal{P}(A \times _)$ -coalgebra, as an instance of our theory. This allowed us in particular to describe their trace semantics using our path category in Section 3.6. This can be extended to tree automata as follows. Assume given a signature Σ , that is, a collection $(\Sigma_n)_{n\in\mathbb{N}}$ of disjoint sets. When σ belongs to Σ_n , we say that n is the arity of σ or that σ is a symbol of arity n. A top-down non-deterministic tree automata as defined in [6] is then the same as a $\{*\}$ -pointed $\mathcal{P}F$ -coalgebra where F is the polynomial functor $X\mapsto\coprod_{\sigma\in\Sigma_n}X^n$. For this functor, $F^n(1)$ is the set of trees over $\Sigma\sqcup\{*(0)\}$ of depth at most n+1 such that a leaf is labelled by * if and only if it is at depth n+1. Intuitively, elements of $F^n(1)$ are partial runs of length n that can possibly be extended. Then, the trace semantics of a tree automata, seen as a pointed coalgebra, is given by the set of partial runs of the automata. In particular, this contains the set of accepted finite trees as those partial runs without any *, and the set of accepted infinite trees, encoded as the sequence of their truncations of depth n, for every n.

In the following, we would like to extend this to other kinds of tree automata by allowing some symmetries. For example, in a tree, we may not care about the order of the children. This boils down to quotient the set X^n of n-tuples, by some permutations of the indices. This can be done generally given a subgroup G of the permutation group \mathfrak{S}_n on n elements by defining X^n/G as the quotient of X^n under the equivalence relation: $(x_1,\ldots,x_n)\equiv_G (y_1,\ldots,y_n)$ iff there is $\pi\in G$ such that for all $i, x_i=y_{\pi(i)}$. Concretely, this means that we replace the polynomial functor F by a so-called analytic functor:

Definition 4.1 [14,15]. An analytic Set-functor is a functor of the form $FX = \coprod_{\sigma \in \Sigma_n} X^n/G_{\sigma}$ where for every $\sigma \in \Sigma_n$, we have a subgroup G_{σ} of the permutation group \mathfrak{S}_n on n elements.

Example 4.2. Every polynomial functor is analytic. The bag-functor is analytic, with $\Sigma = (\{*\})_{n \in \mathbb{N}}$ has one operation symbol per arity and $G_{\sigma} = \mathfrak{S}_{\mathsf{ar}(\sigma)}$ is the full permutation group on $\mathsf{ar}(\sigma)$ elements. It is the archetype of an analytic functor, in the sense that for every analytic functor $F \colon \mathsf{Set} \longrightarrow \mathsf{Set}$, there is a natural transformation into the bag functor $\alpha \colon F \longrightarrow \mathcal{B}$. If F is given by Σ and G_{σ} as above, then α_X is given by

$$FX = \coprod_{\sigma \in \Sigma_n} X^n / G_{\sigma} \ \twoheadrightarrow \ \coprod_{\sigma \in \Sigma_n} X^n / \mathfrak{S}_n \ \to \ \coprod_{n \in \mathbb{N}} X^n / \mathfrak{S}_n = \mathcal{B}X.$$

Proposition 4.3. For an analytic Set-functor F, the following are equivalent (1) a map $f: X \longrightarrow FY$ is F-precise, (2) $\alpha_Y \cdot f$ is \mathcal{B} -precise, (3) every element of Y appears precisely once in the definition of f, i.e. for every $y \in Y$, there is exactly one x in X, such that f(x) is the equivalence class of a tuple (y_1, \ldots, y_n) where there is an index i, such that $y_i = y$; and furthermore this index is unique. So every analytic functor has precise factorizations w.r.t. Set.

4.2 Nominal Sets: Regular Nondeterministic Nominal Automata

We derive an open map situation from the coalgebraic situation for regular nondeterministic nominal automata (RNNAs) [28]. They are an extension of automata to accept words with binders, consisting of literals $a \in \mathbb{A}$ and binders |a| for $a \in \mathbb{A}$; the latter is counted as length 1. An example of such a word of length 4 is $a|_cbc$, where the last c is bound by |c. The order of binders makes difference: $|a|_bab \neq |a|_bba$. RNNAs are coalgebraically represented in the category of nominal sets [10], a formalism about atoms (e.g. variables) that sit in more complex structures (e.g. lambda terms), and gives a notion of binding. Because the choice principles (Ax4) and (Ax5) are not satisfied by every nominal sets, we instead use the class of strong nominal sets for the precise factorization (Definition 3.4).

Definition 4.4 [10,26]. Fix a countably infinite set \mathbb{A} , called the set of atoms. For the group $\mathfrak{S}_{\mathsf{f}}(\mathbb{A})$ of finite permutations on the set \mathbb{A} , a group action (X,\cdot) is a set X together with a group homomorphism \cdot : $\mathfrak{S}_{\mathsf{f}}(\mathbb{A}) \longrightarrow \mathfrak{S}_{\mathsf{f}}(X)$, written in infix notation. An element $x \in X$ is supported by $S \subseteq \mathbb{A}$, if for all $\pi \in \mathfrak{S}_{\mathsf{f}}(\mathbb{A})$ with $\pi(a) = a \ \forall a \in S$ we have $\pi \cdot x = x$. A nominal set is a group action for $\mathfrak{S}_{\mathsf{f}}(\mathbb{A})$ such that every $x \in X$ is finitely supported, i.e. supported by a finite $S \subseteq \mathbb{A}$. A map $f: (X, \cdot) \longrightarrow (Y, \star)$ is equivariant if for all $x \in X$ and $\pi \in \mathfrak{S}_{\mathsf{f}}(\mathbb{A})$ we have $f(\pi \cdot x) = \pi \star f(x)$. The category of nominal sets and equivariant maps is denoted by Nom. A nominal set (X, \cdot) is called strong if for all $x \in X$ and $\pi \in \mathfrak{S}_{\mathsf{f}}(\mathbb{A})$ with $\pi \cdot x = x$ we have $\pi(a) = a$ for all $a \in \mathsf{supp}(x)$.

Intuitively, the support of an element is the set of free literals. An equivariant map can forget some of the support of an element, but can never introduce new atoms, i.e. $\mathsf{supp}(f(x)) \subseteq \mathsf{supp}(x)$. The intuition behind strong nominal sets is that all atoms appear in a fixed order, that is, \mathbb{A}^n is strong, but $\mathcal{P}_{\mathsf{f}}(\mathbb{A})$ (the finite powerset) is not. We set \mathcal{S} to be the class of strong nominal sets:

Example 4.5. The Nom-functor of unordered pairs admits precise factorizations w.r.t. strong nominal sets, but not w.r.t. all nominal sets.

In the application, we fix the set $I = \mathbb{A}^{\# n}$ of distinct *n*-tuples of atoms $(n \ge 0)$ as the pointing. The hom-sets $\mathsf{Nom}(X, \mathcal{P}_{\mathsf{ufs}}Y)$ are ordered point-wise.

Proposition 4.6. Uniformly finitely supported powerset $\mathcal{P}_{\mathsf{ufs}}(X) = \{Y \subseteq X \mid \bigcup_{y \in Y} \mathsf{supp}(y) \text{ finite}\}\$ satisfies (Ax2-5) w.r.t. \mathcal{S} the class of strong nominal sets.⁵

As for F, we study an LTS-like functor, extended with the binding functor [10]:

Definition 4.7. For a nominal set X, define the α -equivalence relation \sim_{α} on $\mathbb{A} \times X$ by: $(a,x) \sim_{\alpha} (b,y) \Leftrightarrow \exists c \in \mathbb{A} \setminus \operatorname{supp}(x) \setminus \operatorname{supp}(y)$ with $(a c) \cdot x = (b c) \cdot y$. Denote the quotient by $[\mathbb{A}]X := \mathbb{A} \times X/\sim_{\alpha}$. The assignment $X \mapsto [\mathbb{A}]X$ extends to a functor, called the binding functor $[\mathbb{A}]$: Nom \longrightarrow Nom.

RNNA are precisely $\mathcal{P}_{\mathsf{ufs}}F$ -coalgebras for $FX = \{\checkmark\} + [\mathbb{A}]X + \mathbb{A} \times X$ [28]. In this paper we additionally consider initial states for RNNAs.

⁵ There are two variants of powersets discussed in [28]. The finite powerset \mathcal{P}_f also fulfils the axioms. However, *finitely supported* powerset \mathcal{P}_{fs} does not fulfil (Ax5).

Proposition 4.8. The binding functor $[\mathbb{A}]$ admits precise factorizations w.r.t. strong nominal sets and so does $FX = \{\checkmark\} + [\mathbb{A}]X + \mathbb{A} \times X$.

An element in PathOrd($\mathbb{A}^{\# n}, F$) may be regarded as a word with binders under a context $\mathbf{a} \vdash w$, where $\mathbf{a} \in \mathbb{A}^{\# n}$, all literals in w are bound or in \mathbf{a} , and w may end with \checkmark . Moreover, two word-in-contexts $\mathbf{a} \vdash w$ and $\mathbf{a}' \vdash w'$ are identified if their closures are α -equivalent, that is, $|a_1 \cdots a_n w| = |a_1 \cdots a_n w'$. The trace semantics of a RNNA T contains all the word-in-contexts corresponding to runs in T. This trace semantics distinguishes whether words are concluded by \checkmark .

4.3 Subsuming arbitrary open morphism situations

Lasota [20] provides a translation of a small path-category $\mathbb{P} \hookrightarrow \mathbb{M}$ into a functor $\mathbb{F} \colon \mathsf{Set}^{\mathbf{obj}\,\mathbb{P}} \longrightarrow \mathsf{Set}^{\mathbf{obj}\,\mathbb{P}}$ defined by $\mathbb{F}(X_P)_P = (\prod_{Q \in \mathbb{P}} (\mathcal{P}(X_Q))^{\mathbb{P}(P,Q)})_{P \in \mathbb{P}}$. So the hom-sets $\mathsf{Set}^{\mathbf{obj}\,\mathbb{P}}(X,\mathbb{F}Y)$ have a canonical order, namely the point-wise inclusion. This admits a functor Beh from \mathbb{M} to \mathbb{F} -coalgebras and lax coalgebra homomorphisms, and Lasota shows that $f \in \mathbb{M}(X,Y)$ is \mathbb{P} -open iff $\mathsf{Beh}(f)$ is a coalgebra homomorphism. In the following, we show that we can apply our framework to \mathbb{F} by a suitable decomposition $\mathbb{F} = TF$ and a suitable object I for the initial state pointing. As usual in open map papers, we require that \mathbb{P} and \mathbb{M} have a common initial object $0_{\mathbb{P}}$. Observe that we have $\mathbb{F} = T \cdot F$ where

$$T(X_P)_{P\in\mathbb{P}} = (\mathcal{P}(X_P))_{P\in\mathbb{P}}$$
 and $F(X_P)_{P\in\mathbb{P}} = (\coprod_{Q\in\mathbb{P}} \mathbb{P}(P,Q) \times X_Q)_{P\in\mathbb{P}}$.

Lasota considers coalgebras without pointing, but one indeed has a canonical pointing as follows. For $P \in \mathbb{P}$, define the characteristic family $\chi^P \in \mathsf{Set}^{\mathsf{obj}\,\mathbb{P}}$ by $\chi^P_Q = 1$ if P = Q and $\chi^P_Q = \emptyset$ if $P \neq Q$. With this, we fix the pointing $I = \chi^{0_{\mathbb{P}}}$.

Proposition 4.9. T, F and I satisfy the axioms from Table 2, with $S = \mathsf{Set}^{\mathbf{obj}\,\mathbb{P}}$.

The path category in $\mathsf{Coalg}_l(I, TF)$ from our theory can be described as follows.

Proposition 4.10. An object of Path(I, F) is a sequence of composable \mathbb{P} -morphisms $0_{\mathbb{P}} \xrightarrow{m_1} P_1 \xrightarrow{m_2} P_2 \cdots \xrightarrow{m_n} P_n$.

5 Conclusions and Further work

We proved that coalgebra homomorphisms for systems with non-deterministic branching can be seen as open maps for a canonical path-category, constructed from the computation type F. This limitation to non-deterministic systems is unsurprising: as we have proved in Section 4.3 on Lasota's work [20], every open map situation can been encoded as a coalgebra situation with a powerset-like functor, so with non-deterministic branching. As a future work, we would like to extend this theory of path-categories to coalgebras for further kinds of branching, especially probabilistic and weighted. This will require (1) to adapt open maps to allow those kinds of branching (2) adapt the axioms from Table 2, by replacing the "+1" part of (Ax1) to something depending on the branching type.

References

- Adámek, J., Herrlich, H., Strecker, G.E.: Abstract and concrete categories: The joy of cats. online and enhanced edition of the book published in 1990 by John Wiley and Sons (http://katmat.math.uni-bremen.de/acc/acc.pdf) (2004)
- 2. Adámek, J., Milius, S., Moss, L.S., Sousa, L.: Well-pointed coalgebras. Logical Methods in Computer Science 9(3), 1–51 (2013)
- Awodey, S.: Category Theory. Oxford University Press, Inc., New York, NY, USA, 2nd edn. (2010)
- Beohar, H., Küpper, S.: On path-based coalgebras and weak notions of bisimulation. In: 7th Conference on Algebra and Coalgebra in Computer Science, CALCO 2017, June 12-16, 2017, Ljubljana, Slovenia. pp. 6:1– 6:17 (2017). https://doi.org/10.4230/LIPIcs.CALCO.2017.6, https://doi.org/ 10.4230/LIPIcs.CALCO.2017.6
- Bonchi, F., Silva, A., Sokolova, A.: The Power of Convex Algebras. In: Meyer, R., Nestmann, U. (eds.) 28th International Conference on Concurrency Theory (CONCUR 2017). vol. 85, pp. 23:1–23:18. Dagstuhl, Germany (2017). https://doi.org/10.4230/LIPIcs.CONCUR.2017.23
- Comon, H., Dauchet, M., Gilleron, R., Jacqemard, F., Lugiez, D., Löding, C., Tison, S., Tommasi, M.: Tree Automata Techniques and Applications. Online http://tata.gforge.inria.fr (2007)
- 7. Dubut, J., Goubault, E., Goubault-Larrecq, J.: Natural homology. In: Proceedings of the 42nd International Colloquium on Automata, Languages and Programming (ICALP'15) Part II. Lecture Notes in Computer Scienc, vol. 9135, pp. 171–183. Springer (July 2015)
- Fahrenberg, U., Legay, A.: History-Preserving Bisimilarity for Higher-Dimensional Automata via Open Maps. Electronic Notes in Theoretical Computer Science 298, 165–178 (November 2013)
- 9. Fiore, M.P., Cattani, G.L., Winskel, G.: Weak bisimulation and open maps. In: 14th Annual IEEE Symposium on Logic in Computer Science (LICS'99). pp. 67–76 (1999)
- Gabbay, M., Pitts, A.M.: A new approach to abstract syntax involving binders. In: Longo, G. (ed.) Proceedings of the Fourteenth Annual IEEE Symp. on Logic in Computer Science, LICS 1999. pp. 214–224. IEEE Computer Society Press (July 1999)
- Hansen, H.H., Klin, B.: Pointwise extensions of GSOS-defined operations. Math. Struct. Comp. Sci. 21(1), 321–361 (2011)
- 12. Hasuo, I., Jacobs, B., Sokolova, A.: Generic trace semantics via coinduction. Logical Methods in Computer Science **3**(4) (2007)
- 13. Jacobs, B., Sokolova, A.: Traces, executions and schedulers, coalgebraically. In: CALCO. Lecture Notes in Computer Science, vol. 5728, pp. 206–220. Springer (2009)
- Joyal, A.: Une théorie combinatoire des séries formelles. Advances in Mathematics 42(1), 1–82 (1981)
- Joyal, A.: Foncteurs analytiques et espèces de structures. In: Proceedings of the "Colloque de combinatoire énumérative". pp. 126–159. Lecture Notes in Mathematics, Springer-Verlag (1986)
- Joyal, A., Nielsen, M., Winskel, G.: Bisimulation from Open Maps. Information and Computation 127, 164–185 (1996)


- Kozen, D., Mamouras, K., Petrisan, D., Silva, A.: Nominal kleene coalgebra. In: Automata, Languages, and Programming - 42nd International Colloquium, ICALP 2015, Kyoto, Japan, July 6-10, 2015, Proceedings, Part II. pp. 286– 298 (2015). https://doi.org/10.1007/978-3-662-47666-6_23, https://doi.org/10. 1007/978-3-662-47666-6_23
- Kurz, A., Petrisan, D., Severi, P., de Vries, F.: Nominal coalgebraic data types with applications to lambda calculus. Logical Methods in Computer Science 9(4) (2013). https://doi.org/10.2168/LMCS-9(4:20)2013, https://doi.org/10.2168/LMCS-9(4:20)2013
- Kurz, A., Petrisan, D., Velebil, J.: Algebraic theories over nominal sets. CoRR abs/1006.3027 (2010)
- 20. Lasota, S.: Coalgebra morphisms subsume open maps. Theoretical Computer Science 280(1), 123 135 (2002)
- 21. Milius, S.: A sound and complete calculus for finite stream circuits. In: Proc. 25th Annual Symposium on Logic in Computer Science (LICS'10). pp. 449–458 (2010)
- Milius, S., Pattinson, D., Schröder, L.: Generic trace semantics and graded monads. In: Moss, L.S., Sobocinski, P. (eds.) Proc. 6th Conference on Algebra and Coalgebra in Computer Science, CALCO 2015. Leibniz International Proceedings in Informatics, vol. 35, pp. 253–269 (2015), http://www8.cs.fau.de/_media/research: papers:traces-gm.pdf
- 23. Milius, S., Schröder, L., Wißmann, T.: Regular behaviours with names. Applied Categorical Structures **24**(5), 663–701 (2016)
- 24. Nielsen, M., Hune, T.: Bisimulation and Open Maps for Timed Transition Systems. Fundamenta Informaticae 38, 61–77 (1999)
- Park, D.: Concurrency and Automata on Infinite Sequences. Theoretical Computer Science 104, 167–183 (1981)
- Pitts, A.M.: Nominal Sets: Names and Symmetry in Computer Science, Cambridge Tracts in Theoretical Computer Science, vol. 57. Cambridge University Press (2013)
- 27. Rutten, J.: Universal coalgebra: a theory of systems. Theoretical Computer Science **249**(1), 3 80 (2000)
- Schröder, L., Kozen, D., Milius, S., Wißmann, T.: Nominal automata with name binding. In: Esparza, J., Murawski, A. (eds.) Proc. 20th International Conference on Foundations of Software Science and Computation Structures (FoSSaCS 2017). Lecture Notes in Computer Science, vol. 10203, pp. 124–142. Springer (2017)

A Omitted Proofs

Proof of Remark 3.2

Lemma A.1. Assuming that F preserves weak pullbacks, a morphism $s: S \longrightarrow FR$ is F-precise iff for all f, h the following implication holds:

Proof. Sufficiency is clear, because one can fix D = R and $g = \mathrm{id}_R$ in the definition of F-precise. For necessity, consider s, f, g, h as in the definition of F-precise. The pullback of g along h is weakly preserved by F, and so we have the following commuting diagram:



Hence, s induces some $d: R \longrightarrow P$ with $Fd \cdot s = f'$ and $\pi_2 \cdot d = \mathrm{id}_R$. The witness that s is F-precise is $\pi_1 \cdot d: R \longrightarrow C$, because $F(\pi_1 \cdot d) \cdot s = f$ and $h \cdot \pi_1 \cdot d = g$. \square

Proof of Remark 3.5

Apply that f_2 is F-precise and one obtains a lifting $d: Y_2 \longrightarrow Y_1$ with $Fd \cdot f_2 = f_1$ and $h \cdot d = \mathrm{id}_{Y_2}$, i.e. d has a left inverse, h. Additionally, since f_1 is F-precise, $Fd \cdot f_2 = f_1$ induces some $d': Y_1 \longrightarrow Y_2$ with $f_2 = Fd' \cdot f_1$ and $d \cdot d' = \mathrm{id}_{Y_1}$. Since d has both a left and right inverse, it is an isomorphism, and so is its left-inverse h.

For the second part, we a have a precise factorisation of f, that is, a F-precise morphism $f' \colon S \longrightarrow Y'$ with $Y' \in \mathcal{S}$, and a morphism $h \colon Y' \longrightarrow Y$ such that $f = Fh \cdot f'$. By the previous point, h is an isomorphism, and since \mathcal{S} is closed under isomorphisms, $Y \in \mathcal{S}$.

Proof of Proposition 3.6

- 1. Fix Z an object of \mathbb{C} , and assume given a morphism $f: S \longrightarrow Z = \mathbb{C}_Z Y$, with $S \in \mathcal{S}$. Since a morphism of the form $f': S \longrightarrow F0$ is always F-precise, then $f' = f: S \longrightarrow Z = \mathbb{C}_Z 0$ is \mathbb{C}_Z -precise. Taking h as the unique morphism from 0 to Y, $Fh \cdot f' = \mathrm{id}_Z \cdot f = f$.
- 2. Let us start with the following lemma:

Lemma A.2. If $f: S \longrightarrow FR$ is F-precise and $f': R \longrightarrow F'Q$ is F'-precise, then $Ff' \cdot f$ is $F \cdot F'$ -precise.

Proof. Assume given the following situation:

$$S \xrightarrow{k} FF'C$$

$$f \downarrow \qquad \qquad \downarrow FF'u$$

$$FF' \downarrow \qquad \qquad \downarrow FF'u$$

$$FF'Q \xrightarrow{FF'v} FF'W$$

In particular, we have $(FF'v \cdot Ff') \cdot f = FF'u \cdot k$. Since f is F-precise, there is $d \colon R \longrightarrow F'C$ such that

$$k = Fd \cdot f$$
$$F'v \cdot f' = F'u \cdot d$$

Since f' is F'-precise, there is $d': Q \longrightarrow C$ such that

$$d = F'd' \cdot f'$$
$$v = u \cdot d'$$

This implies that $k = FF'd' \cdot (Ff' \cdot f)$.

Let us prove this point now. We start with a morphism $f\colon S\longrightarrow FF'Y$, with $S\in\mathcal{S}$. Since F admits precise factorisations w.r.t. \mathcal{S} , there are $f'\colon S\longrightarrow FY'$ F-precise and $h\colon Y'\longrightarrow F'Y$ with $Y'\in\mathcal{S}$, and $Fh\cdot f'=f$. Now, since F' admits precise factorisations w.r.t. \mathcal{S} , there are $f''\colon Y'\longrightarrow F'Y''$ F'-precise and $h'\colon Y''\longrightarrow Y$ such that $F'h'\cdot f''=h$. Consequently, $f=FF'h'\cdot (Ff''\cdot f')$ and $Ff''\cdot f'$ is FF'-precise by the previous lemma.

3. Let us start with the following lemma:

Lemma A.3. For a family of functors $F_i : \mathbb{C} \longrightarrow \mathbb{C}$, $i \in I$, and F_i -precise morphisms $f_i : X \longrightarrow F_iY_I$, $g : X \longrightarrow GY_G$, then we have a $\prod_{i \in I} F_i$ -precise morphism

$$\langle F_i \mathsf{in}_i \cdot f_i \rangle \colon X \longrightarrow \prod_{i \in I} F_i (\prod_{j \in I} Y_j).$$

Proof. Consider a square

$$X \xrightarrow{\langle v_i \rangle_{i \in I}} \prod_{i \in I} F_i W \qquad X \xrightarrow{v_i} FW$$

$$\prod_{i \in I} F_i Y_i \qquad \left| F_i u \right|_{i \in I} \implies F_i Y_i \qquad F_i u$$

$$\prod_{i \in I} F_i (\coprod_{j \in I} Y_j) \xrightarrow{[F_i h]_{i \in I}} \prod_{i \in I} F_i Z \qquad F_i (\coprod_{j \in I} Y_j) \xrightarrow{F_i h} F_i Z$$

Since f_i is F_i -precise, we obtain some $d_i \colon Y_i \longrightarrow W$ with $F_i d_i \cdot f_i = v_i$ and $u \cdot d_i = h \cdot \mathsf{in}_i$. We have $[d_j]_{j \in I} \colon \coprod_{j \in I} Y_j \longrightarrow W$ with

$$(\prod_{i \in I} F_i[d_j]_{j \in I}) \cdot \langle F_i \mathsf{in}_i \cdot f_i \rangle_{i \in I} = \langle Fd_i \cdot f_i \rangle_{i \in I} = \langle v_i \rangle_{i \in I}$$

and

$$u \cdot [d_i]_{i \in I} = [u \cdot d_i]_{i \in I} = [h \cdot \mathsf{in}_i]_{i \in I} = h.$$

Let us prove this point now. Consider $\langle f_i \rangle_{i \in I} \colon X \longrightarrow \prod_{i \in I} F_i Y$ and consider the F_i -precise factorizations:

$$X \xrightarrow{f_i} F_i Y$$

$$f'_i \xrightarrow{} f_i h_i$$

$$F_i Y_i$$

By Lemma A.3 we have that $\langle F_i \mathsf{in}_i \cdot f_i' \rangle \colon X \longrightarrow \prod_{i \in I} F_i (\coprod_{j \in I} Y_j)$ is $\prod_{i \in I} F_i$ -precise and it is the $\prod_{i \in I} F_i$ -precise factorization of $\langle f_i \rangle_{i \in I}$:

$$(\prod_{i \in I} F_i)[h_i]_{i \in I} \cdot \langle F_i \mathsf{in}_i \cdot f_i' \rangle_{i \in I} = \langle F_i h_i \cdot f_i' \rangle_{i \in I} = \langle f_i \rangle_{i \in I}.$$

4. By *I*-extensive we mean 'extensive' if *I* is finite and 'infinitary extensive' if *I* is infinite. In any case, \mathbb{C} is *I*-extensive if for for all families $(X_i)_{i\in I}, (Y_i)_{i\in I}, (g_i)_{i\in I}, (x_i)_{i\in I}$ and h with

$$\begin{array}{ccc} X_i & \xrightarrow{x_i} & P \\ g_i \downarrow & & \downarrow h \\ Y_i & \xrightarrow{\operatorname{in}_i} & \coprod_{i \in I} Y_i \end{array} \quad \forall i \in I$$

the following equivalence holds:


$$(x_i \colon X_i \longrightarrow P)_{i \in I}$$
 is a coproduct \iff

for all $i \in I$ the above square is a pullback

Let us start with the following lemma:

Lemma A.4. Given functors $F_i : \mathbb{C} \longrightarrow \mathbb{C}$, $i \in I$, on an I-extensive category, and morphisms $f_i : A_i \longrightarrow F_i X_i$, $i \in I$. Then $\coprod_{i \in I} (F_i \text{in}_i \cdot f_i) : \coprod_{i \in I} A_i \longrightarrow (\coprod_{i \in I} F_i) (\coprod_{i \in I} X_i)$ is $\coprod_{i \in I} F_i$ -precise if f_i is F_i -precise for every $i \in I$.

Proof. Consider a commutative square



Note that by the extensivity of \mathbb{C} we can assume that the top morphism $\coprod_{i\in I} A_i \longrightarrow \coprod_{i\in I} F_i C$ is indeed a coproduct of morphisms. Hence we have $F_i g_i \cdot f_i = F_i c \cdot a_i$, for every $i \in I$. Since f_i is F_i -precise, this induces some $d_i \colon X_i \longrightarrow C$ with $F_i d_i \cdot f_i = a_i$ and $c \cdot g_i = d_i$. In total, we have $[d_i]_{i\in I} \colon \coprod_{i\in I} X_i \longrightarrow C$ with $c \cdot [d_i]_{i\in I} = [g_i]_{i\in I}$ and

$$\left(\prod_{i\in I} F_i\right)[d_i]_{i\in I} \cdot \prod_{i\in I} (F_i \mathsf{in}_i \cdot f_i) = \prod_{i\in I} (F_i d_i \cdot f_i) = \prod_{i\in I} a_i \qquad \Box$$

Let us prove this point now. Given a morphism $h: X \longrightarrow (\coprod_{i \in I} F_i)(Y)$, construct the pullbacks in the *I*-extensive category \mathbb{C} :

$$X_{i} \xrightarrow{\operatorname{in}_{i}} X$$

$$f_{i} \downarrow \xrightarrow{\operatorname{in}_{i}} (\coprod_{i \in I} F_{i})(Y)$$
 for every $i \in I$.

So we have $X = \coprod_{i \in I} X_i$ with the coproduct injections as in the top row of the above pullback diagram, and in particular $h = [\inf_i \cdot f_i]_{i \in I} = \coprod_{i \in I} f_i$. Let f'_i be the F_i -precise morphisms through which f_i factors for every $i \in I$:

$$\forall i \in I \qquad X_i \xrightarrow{\exists f_i'} FY_i \qquad \coprod_{i \in I} X_i \xrightarrow{\coprod_{i \in I} F \text{in}_i \cdot f_i'} (\coprod_{i \in I} F_i)(\coprod_{i \in I} Y_i)$$

$$\downarrow_{Fy_i} \implies \downarrow_{h = \coprod_{i \in I} f_i} (\coprod_{i \in I} F_i)(\coprod_{i \in I} Y_i)$$

$$(\coprod_{i \in I} F_i)(Y)$$

By Lemma A.4, $\coprod_{i \in I} Fin_i \cdot f_i'$ is $\coprod_{i \in I} F_i$ -precise.

5. We first show that $\eta_X \colon X \longrightarrow R(LX)$ is R-precise for every $X \in \mathbb{C}$. Since right adjoints preserve limits, R preserves (weak) pullbacks and it suffices

to check Lemma $A.1/Remark\ 3.2.$ For any suitable commutative triangle we have

$$X \xrightarrow{g} RZ$$

$$\eta_X \downarrow \qquad Rh$$

$$RLX$$

Since $L \dashv R$, there exists a unique g' with

$$X \xrightarrow{g} RZ \qquad LX \xrightarrow{g'} Z$$

$$\uparrow_{\eta_X} \downarrow \qquad \text{and } id_{LX} \downarrow \qquad h$$

$$RLX \qquad LX$$

by the universal mapping property of $L\dashv R$ and by the naturality of the isomorphism $\mathbb{C}(LX,Z)\cong\mathbb{C}(X,RZ)$ respectively. Now, we can prove this point:

- For necessity and $X \in \mathcal{S}$, we have that $\eta_X : X \longrightarrow RLX$ is R-precise by the previous argument, and hence $LX \in \mathcal{S}$ by Remark 3.5.
- For sufficiency, consider a morphism $f: X \longrightarrow RY$. The adjunction induces a unique $f': LX \longrightarrow Y$ with $Rf' \cdot \eta_X = f$, where $\eta_X: X \longrightarrow RLX$ is R-precise as shown above. Since L preserves objects in S, we have that $LX \in S$.

Proof of Example 3.7

Given an \mathcal{P} -precise $f: X \longrightarrow \mathcal{P}Y$, define $f': X \longrightarrow \mathcal{P}(Y + Y)$ by $f'(x) = \{\operatorname{inl}(y), \operatorname{inr}(y) \mid y \in f(x)\}$, and so $\mathcal{P}[\operatorname{id}_Y, \operatorname{id}_Y] \cdot f' = f$. Hence, we have some $d: Y \longrightarrow Y + Y$ with $[\operatorname{id}_Y, \operatorname{id}_Y] \cdot d = \operatorname{id}_Y$ and $f' = \mathcal{P}d \cdot f$. The first equation implies that for every $y \in Y$, d(y) is $\operatorname{inl}(y)$ or $\operatorname{inr}(y)$. So for $x \in X$ and for every $y \in Y$, either $\operatorname{inl}(y)$ or $\operatorname{inr}(y)$ is in $d[f(x)] = (\mathcal{P}d \cdot f)(x) = f'(x)$, which is a contradiction unless X or Y is empty. Hence, $f(x) = \emptyset$ for all $x \in X$.

Proof of Remark 3.10

The proof is by induction on the length. Since $P_0 = I = Q_0$, $\phi_0 : P_0 \longrightarrow Q_0$ is an isomorphism. Assume that $\phi_k : P_k \longrightarrow Q_k$ is isomorphic for k < n. Since $q_k \cdot \phi_k$ and p_k are F + 1-precise, we have that ϕ_{k+1} is isomorphic by Remark 3.5.

The second part of the remark is a consequence of the second part of Remark 3.5.

Proof of Proposition 3.13

1. - Functoriality. consider a path morphism ϕ_{n+1} : $(P_{n+1}, p_n) \longrightarrow (Q_{m+1}, q_m)$ and use that all components are isos (Remark 3.5)

- Hence $\mathsf{Comp}(\boldsymbol{P}_{n+1},\boldsymbol{p}_n) \leq \mathsf{Comp}(\boldsymbol{Q}_{m+1},\boldsymbol{q}_m).$
- Fullness. Since PathOrd(I, F) is partially ordered, it sufficies to show that whenever PathOrd(I, F)(Comp (P_{n+1}, p_n) , Comp (Q_{m+1}, q_m)) is non-empty, then so is Path $(I, F)(P_{n+1}, p_n)$, (Q_{m+1}, q_m) . We have $n \leq m$ and construct a morphism $\phi_{n+1} \colon (P_{n+1}, p_n) \longrightarrow (Q_{m+1}, q_m)$ by induction. There is nothing to do in the base case $\phi_0 := \operatorname{id}_I$. For the step, assume some $\phi_k \colon P_k \longrightarrow Q_k$ making the following diagram commute:

$$P_{k} \xrightarrow{\phi_{k}} Q_{k} \xrightarrow{q_{k}} FQ_{k+1} \xrightarrow{\cdots} F^{n-k}Q_{n}$$

$$\downarrow^{p_{k}} \qquad \qquad \downarrow^{F^{n-k}!}$$

$$FP_{k+1} \xrightarrow{\cdots} F^{n-k}P_{n} \xrightarrow{F^{n-k}!} F^{n-k}1$$

Since p_k is F-precise, we obtain a morphism $\phi_{k+1}: P_{k+1} \longrightarrow Q_{k+1}$ with

$$P_{k} \xrightarrow{\phi_{k}} Q_{k} \qquad P_{k+1} \xrightarrow{\phi_{k+1}} Q_{k+1} \longrightarrow F^{n-k-1}Q_{n}$$

$$\downarrow^{p_{k}} \qquad \downarrow^{q_{k}} \text{ and } \qquad \downarrow^{F^{n-k-1}!} \qquad \downarrow^{F^{n-k-1}!} F^{n-k-1}1.$$

So, ϕ_{k+1} is indeed a morphism in Path(I, F).

2. For every $u: I \longrightarrow F^n 1$ we can define a path $(\boldsymbol{P}_{n+1}, \boldsymbol{p}_n)$ inductively, starting with $P_0 := I \in \mathcal{S}$. Given some \mathbb{C} -morphism $u': P_k \longrightarrow F^{n-k} 1$, $k \leq n$, $P_k \in \mathcal{S}$, we are done with $r = \mathrm{id}_1$ if k = n. If k < n consider the F-precise factorization

$$P_k \xrightarrow{r} FP_{k+1}$$

$$\downarrow^{Fr'}$$


$$FF^{n-k-1}$$

providing $P_{k+1} \in \mathcal{S}$ and $p_k \colon P_k \longrightarrow FP_{k+1}$ and some $u'' \colon P_{k+1} \longrightarrow F^{n-k-1}1$. This defines the families \mathbf{P}_{n+1} and \mathbf{p}_n with $\mathsf{Comp}(\mathbf{P}_{n+1}, \mathbf{p}_n) = u$.

3. By Remark 3.11.

Proof of Proposition 3.18

For all k < n, the following diagram commutes:




For sufficiency, assume that $[x_k]_{k \leq n}$ is a morphism in $\mathsf{Coalg}_l(I, TF)$. Hence for all k < n:

$$[\eta, \bot] \cdot (Fx_{k+1} + 1) \cdot p_k = TF[x_j]_{j \le n} \cdot \rho \cdot \mathsf{in}_k \sqsubseteq \xi \cdot [x_j]_{j \le n} \cdot \mathsf{in}_k = \xi \cdot x_k.$$

For necessity, we have for all k < n:

$$\begin{array}{c|c} P_k & \xrightarrow{\operatorname{in}_k} & \coprod_{j \leq n} P_j & \xrightarrow{[x_j]_{j \leq n}} X \\ \downarrow p_k & \downarrow & \downarrow & \downarrow \\ \downarrow p_k & \downarrow & \downarrow & \downarrow \\ \downarrow & \downarrow & \downarrow & \downarrow \\ FP_{k+1} + 1 \xrightarrow{Fp_{k+1}+1} FX + 1 & \xrightarrow{[\eta,\bot]} TFX \\ \downarrow & \uparrow & \uparrow \\ \hline TF[x_j]_{j \leq n} \cdot \rho \cdot \operatorname{in}_k \\ \end{array}$$

and



Since the family of coproduct injections in_k is jointly epic we have by (Ax2) that $[x_k]_{k < n}$ is indeed a weak homomorphism.

Proof of Theorem 3.20

As usual in open map proofs, this direction of the characterization theorem is shown by considering only Path-morphisms of length difference 1. Indeed every Path-morphism ϕ from a path of length n to a path of length $m \geq n+2$ can be expressed as a composition of Path-morphisms of smaller length difference $\phi = \psi \cdot \theta$. When applying the openness of a morphism h to θ , we obtain a diagonal $d : \operatorname{dom}(\psi) \longrightarrow \operatorname{dom}(h)$ which allows applying the openness of h to ψ , yielding the desired lifting for ϕ .




Consider a $\mathsf{Path}(I, F+1)$ -morphism $\phi_{n+1} \colon (P_{n+1}, p_n) \longrightarrow (Q_{n+2}, q_{n+1})$ and a commuting square in $\mathsf{Coalg}_l(I, TF)$ where h is a homomorphism:

$$J(\boldsymbol{P}_{n+1}, \boldsymbol{p}_n) \xrightarrow{[x_k]_{k \le n}} (X, \xi, x_0)$$


$$\downarrow J\phi_{n+1} \downarrow \qquad \qquad \downarrow h$$

$$J(\boldsymbol{Q}_{n+2}, \boldsymbol{q}_{n+1}) \xrightarrow{[y_k]_{k \le n+1}} (Y, \zeta, y_0)$$

By precomposition with in_n we obtain the following commuting diagram:



Applying (Ax5) to the outer part of the diagram yields some $f: P_n \longrightarrow FX + 1$ and the commuting diagram:



Since q_n is F + 1-precise, we obtain the following:

$$P_{n} \xrightarrow{f} FX + 1 \qquad X$$

$$q_{n} \cdot \phi_{n} \downarrow \qquad \qquad \qquad \downarrow h$$

$$FQ_{n+1} + 1 \qquad \qquad Q_{n+1} \xrightarrow{y_{n+1}} Y$$

Define $d_k : Q_k \longrightarrow X$ by $x_k \cdot \phi_k^{-1}$ for $k \leq n$. It remains to show that

$$[d_k]_{k \le n+1} \colon J(\boldsymbol{Q}_{n+1}, \boldsymbol{Q}_n) \longrightarrow (X, \xi, x_0)$$

is indeed a morphism in $\mathsf{Coalg}_l(I,TF)$ and that it makes the desired diagram in $\mathsf{Coalg}_l(I,TF)$ commute.

1. That $[d_k]_{k \le n+1}$ is a weak homomorphism follows from Proposition 3.18 because we have:

$$Q_{n} \xrightarrow{\phi_{n}^{-1}} P_{n} \xrightarrow{x_{n}} X$$

$$Q_{n} \downarrow \text{ Def. } d_{n+1} \downarrow f \sqsubseteq \downarrow \xi$$

$$FQ_{n+1} + 1 \xrightarrow{Fd_{n+1}+1} FX + 1 \xrightarrow{[\eta, \bot]} TFX$$

and for all k < n:

$$Q_{k} \xrightarrow{\phi_{k}^{-1}} P_{k} \xrightarrow{x_{k}} X$$

$$\downarrow q_{n} \downarrow \qquad \downarrow f \qquad \text{by Proposition 3.18} \qquad \downarrow \xi$$

$$FQ_{k+1} + 1 \xrightarrow{F\phi_{n+1}^{-1} + 1} FP_{k+1} + 1 \xrightarrow{Fx_{k+1}} FX + 1 \xrightarrow{[\eta, \bot]} TFX$$

$$\downarrow Fd_{k+1}$$

2. The first desired commutativity $[d_k]_{k \le n+1} \cdot J\phi_{n+1} = [x_k]_{k \le n}$ is clear by the definition of d_k . The second commutativity $[y_k]_{k \le n+1} = h \cdot [d_k]_{k \le n+1}$ is proven because the coproduct injections are jointly epic. For $k \le n$ we clearly have $y_k = h \cdot x_k \cdot \phi_k^{-1} = h \cdot d_k$. For k = n+1 we have $y_{n+1} = h \cdot d_{n+1}$ by the definition of d_{n+1} .

Proof of Theorem 3.24

Having $\zeta \cdot h \supseteq TFh \cdot \xi$ already, we only need to show:

$$\begin{array}{c|c} X & \xrightarrow{\xi} TFX \\ h \downarrow & \swarrow & \downarrow^{TFh} \\ Y & \xrightarrow{\zeta} TFY \end{array}$$

By path-reachability, it suffices to show that for all runs $[x_k]_{k \le n}$: $J(\boldsymbol{P}_{n+1}, \boldsymbol{p}_n) \longrightarrow (X, \xi, x_0)$ we have:

$$\coprod_{k \leq n} P_k \xrightarrow{[x_k]_{k \leq n}} X \xrightarrow{\xi} TFX
\downarrow [x_k]_{k \leq n} \qquad \qquad \downarrow TFh
X \xrightarrow{h} Y \xrightarrow{\zeta} TFY$$

By (Ax2), coproduct injections are jointly epic in an ordered-enriched sense, so by induction, it suffices to prove that:

$$\begin{array}{c|c} P_n & \xrightarrow{x_n} & X & \xrightarrow{\xi} & TFX \\ x_n \downarrow & & & \downarrow TFh \\ X & \xrightarrow{h} & Y & \xrightarrow{\zeta} & TFY \end{array}$$

By (Ax4), we can prove this by showing that for all $p'_n: P_n \longrightarrow TFY$ we have. We will prove the implication

In the following, this implication is proved for all p'_n . For such a $p'_n: P_n \longrightarrow FY+1$, since $P_n \in \mathcal{S}$, (Ax1) yields an object $P_{n+1} \in \mathcal{S}$ and morphisms $p_n: P_n \longrightarrow FP_{n+1}+1$ and $y_{n+1}: P_{n+1} \longrightarrow Y$ with $(Fy_{n+1}+1) \cdot p_n = p'_n$. So we have a new path (P_{n+2}, p_{n+1}) and a morphism defined by

$$[y_k]_{k \le n+1} : \coprod_{k \le n+1} P_k \longrightarrow Y$$
 with $y_k = x_k$ for $k \le n$, and y_{n+1} as above.

This is indeed a lax homomorphism $J(\boldsymbol{P}_{n+2}, \boldsymbol{p}_{n+1}) \longrightarrow (Y, \zeta, y_0)$ because of Proposition 3.18 and because for all k < n:

$$\begin{array}{c|c} P_k & \xrightarrow{x_k} X & \xrightarrow{h} Y \\ & & & & & \downarrow \zeta \\ \hline FX + 1 & \xrightarrow{[\eta, \bot]} TFX & \xrightarrow{TFh} TFY \\ \hline Fx_{k+1} + 1 & & & & \downarrow \uparrow \\ \hline FP_{k+1} + 1_{Fy_{k+1} + 1} FY + 1 & & & \downarrow \uparrow \end{array}$$

and

$$P_{n} \xrightarrow{x_{n}} X \xrightarrow{h} Y$$

$$\downarrow p_{n} \downarrow \qquad \qquad \downarrow \zeta$$

$$FP_{n+1} + 1 \xrightarrow{Fy_{n+1} + 1} FY + 1 \xrightarrow{[\eta_{FY}, \bot_{FY}]} TFY$$

Since $y_k = x_k$, for $k \le n$, the constructed lax homomorphism makes the following square commute

$$\begin{split} J(\boldsymbol{P}_{n+1},\boldsymbol{p}_n) & \xrightarrow{[x_k]_{k \leq n}} (X,\xi,x_0) \\ J(\mathrm{id}_{P_k})_{k \leq n+1} \!\!\!\! \int_{h} & \text{in } \mathsf{Coalg}_l(I,TF). \\ J(\boldsymbol{P}_{n+2},\boldsymbol{p}_{n+1}) & \xrightarrow{[y_k]_{k \leq n+1}} (Y,\zeta) \end{split}$$

Since h is open, we obtain a diagonal lifting, that is, a lax homomorphism $[d_k]_{k \leq n+1} : J\mathbf{P}_{n+1} \longrightarrow (X, \xi, x_0)$ with $h \cdot d_k = y_k$ for $k \leq n + 1$ and $d_k = x_k$ for $k \leq n$. This finally proves the desired implication:

$$P_{n} \xrightarrow{d_{n} = x_{n}} X$$

$$\downarrow p_{n} \qquad \sqsubseteq \text{ by Proposition 3.18} \qquad \downarrow \xi$$

$$FP_{n+1} + 1 \xrightarrow{Fd_{n+1}+1} FX + 1 \xrightarrow{[\eta, \bot]} TFX$$

$$\downarrow FY_{n+1}+1 \qquad \downarrow TFY$$

$$FY + 1 \xrightarrow{[\eta, \bot]} TFY$$

Remarks on Assumption 3.26

The category of sets and the category of nominal sets have (epi,mono)-factorizations. However, the assumption is rather unusual, because in most categories, a morphism that is both epi and mono is not necessarily an isomorphism. By the assumption of (epi,mono)-factorizations together with the cocompleteness of \mathbb{C} , the following are equivalent for any family $(e_i \colon X_i \longrightarrow Y)_{i \in I}$:

- 1. $(e_i)_{i \in I}$ are jointly epic.
- 2. $[e_i]_{i \in I} : \coprod_{i \in I} X_i \longrightarrow Y$ is an epimorphism.
- 3. $[e_i]_{i \in I} : \coprod_{i \in I} X_i \longrightarrow Y$ is a strong epimorphism.
- 4. $(e_i)_{i \in I}$ are jointly strong epic.

All the results in subsection 3.5 work when replacing 'jointly epic' by 'jointly strong epic' and by replacing '(epi,mono)-factorizations' by the very common assumption of '(strong epi,mono)-factorizations'. Since in all our instances (possibly sorted Sets, Nominal Sets), all epimorphisms are strong, we phrase the results and proofs in terms of (jointly) epic families for the sake of simplicity.

Proof of Proposition 3.27

Consider a subcoalgebra $h: (Y, \zeta, y_0) \longrightarrow (X, \xi, x_0)$. For every run $[x_k]_{k \leq n}: J(\boldsymbol{P}_{n+1}, \boldsymbol{p}_n) \longrightarrow (X, \xi, x_0)$, we have the commuting square:

$$J0 = (I, \perp_{FI} \cdot !, \mathrm{id}_I) \xrightarrow{!} (Y, \zeta, y_0)$$

$$J! \downarrow \qquad \qquad \downarrow h$$

$$J(\boldsymbol{P}_{n+1}, \boldsymbol{p}_n) \xrightarrow{[x_k]_{k \leq n}} (X, \xi, x_0)$$

and since h is open by Theorem 3.20, we have a run $[d_k]_{k \le n}$ of $(\boldsymbol{P}_{n+1}, \boldsymbol{p}_n)$ in (Y, ζ, y_0) . Since $[x_k]_{k \le n}$ is epic and $h \cdot d_k = x_k$ for all $0 \le k \le n$, we have that the mono h is an epimorphism and hence an isomorphism.

Proof of Proposition 3.28


For functors $H: \mathbb{C} \longrightarrow \mathbb{C}$ preserving intersections, we have breadth-first-search:

Lemma A.5. Let $H: \mathbb{C} \longrightarrow \mathbb{C}$ preserve arbitrary intersections and let \mathbb{C} have $(\mathcal{E}, mono)$ -factorizations, countable coproducts, and arbitrary intersections. For an I-pointed H-coalgebra $I \xrightarrow{i} X \xrightarrow{\xi} HX$, there are monomorphisms $(m_k: X_k \rightarrowtail X)$ with the property that:


- 1. m_0 is the mono-part of the $(\mathcal{E}, mono)$ -factorization of i.
- 2. m_{k+1} is the least subobject of X such that $\xi \cdot m_k$ factors through Hm_{k+1} .
- 3. The union of the m_k , i.e. the image of $[m_k]_{k\geq 0}$: $\coprod_{k\geq 0} X_k \longrightarrow X$ is the carrier of a reachable subcoalgebra of (X, ξ, x_0) . In particular, if (X, ξ, x_0) is reachable, then $[m_k]_{k\geq 0} \in \mathcal{E}$.

Intuitively, $X_k \rightarrow X$ contains precisely the states that are k steps away from the initial state X_0 , and so the union $\bigcup_{k>0} X_k$ contains all reachable states.

Proof. 1. and 2. Let $I \xrightarrow{i'} X_0 \xrightarrow{m_0} X$ be the $(\mathcal{E}, \text{mono})$ -factorization of $i: I \longrightarrow X$. For the inductive step, assume $m_k: X_k \rightarrowtail X$. Consider the family of subobjects $m': X' \rightarrowtail X$ with the property that there exists some $\xi': X_k \longrightarrow HX'$ with $Hm' \cdot \xi' = \xi \cdot m_k$. Denote the intersection of these subobjects by $m_{k+1}: X_{k+1} \rightarrowtail X$, i.e. we have the wide pullback:




Note that since H preserves intersections, i.e. pullbacks of monomorphisms, it preserves monomorphisms, because a morphism m is mono. So the above wide pullback of monos is mapped again to a pullback of monos, and we have the commutative diagram:



By the universal property of the wide pullback, there is a unique map $\xi_k \colon X_k \longrightarrow HX_{k+1}$ with $Hm_{k+1} \cdot \xi_k = \xi \cdot m_k$. Indeed, $m_{k+1} \colon X_{k+1} \rightarrowtail X$ is the least subobject of X with this property, because any other $m' \colon X' \rightarrowtail X$ with this property is itself included in the diagram for the intersection and hence $X_{k+1} \rightarrowtail X'$.

3. Consider the $(\mathcal{E}, \text{mono})$ -factorization $\coprod_{k\geq 0} X_k \stackrel{e}{\to} Y \stackrel{h}{\to} X$ of $[m_k]_{k\geq 0}$, then with the ξ_k as above, we have the commutative diagram:




and since $e \in \mathcal{E}$ is left-orthogonal to the mono Hh, there is a unique morphism $y \colon Y \longrightarrow HY$ making e and h coalgebra homomorphisms. It remains to show that $(Y, y, e \cdot \mathsf{in}_0 \cdot i')$ does not have a proper subcoalgebra. Assume a pointed subcoalgebra $f \colon (Z, \zeta, z_0) \longrightarrow (Y, y, e \cdot i')$, with $f \colon Z \rightarrowtail Y$. We show that e factors through f, by constructing maps $d_k \colon X_k \longrightarrow Z$ with $e \cdot \mathsf{in}_k = f \cdot d_k$ inductively. For k = 0, we have the following commuting square which uniquely induces d_0 :

$$I \xrightarrow{i'} X_0$$

$$z_0 \downarrow \xrightarrow{\exists ! d_0} \downarrow e \cdot \mathsf{in}_0$$

$$Z \xrightarrow{\iota} f Y$$

Given $d_k : X_0 \longrightarrow Z$ with $f \cdot d_k = e \cdot \mathsf{in}_k$, we have the commutative diagram:



So $h \cdot f$ is a subobject with the property that $\xi \cdot m_k$ factors through $H(h \cdot f)$. Since by item (2.), X_{k+1} is the least subobject with this property, there is some $d_{k+1} \colon X_{k+1} \rightarrowtail Z$ with $(h \cdot f) \cdot d_{k+1} = m_{k+1}$ and in particular $f \cdot d_{k+1} = e \cdot \mathsf{in}_{k+1}$ since h is monic.


In total, we have $f \cdot [d_k]_{k \geq 0} = [e \cdot \mathsf{in}_k]_{k \geq 0} = e \in \mathcal{E}$, and so $f \in \mathcal{E}$ (by \mathcal{E} -laws in factorizations) and hence the mono f must be an isomorphism.

We now can continue with the proof of the main statement, and in the following we instantiate Lemma A.5 with H = TF, $\mathcal{E} = \text{epi}$.

Proof (of Proposition 3.28). Let $\xi_k \colon X_k \longrightarrow TFX_{k+1}$ the witness of item 2 of Lemma A.5, and so $TFm_{k+1} \cdot \xi_k = \xi \cdot m_k$ and m_{k+1} is the least subobject with

this property. In the following, we choose sets \mathcal{F}_k , $k \geq 0$, containing morphisms $(E, e) \in \mathcal{F}_k$, $e: E \longrightarrow X_k$, $E \in \mathcal{S}$, inductively:


- 1. \mathcal{F}_0 contains only $x_0' : I \to X_0$, where $x_0 = m_0 \cdot x_0'$ is the pointing $I \to X$, as provided by item 1 of Lemma A.5.
- 2. For every $(E, e) \in \mathcal{F}_k$, let $G_{(E, e)} \subseteq \mathbb{C}(E, FX_{k+1} + 1)$ be the set of morphisms $g: E \longrightarrow FX_{k+1} + 1$ with:



For each $g \in G_{(E,e)}$, choose some object $Y_g \in \mathcal{S}$, and morphisms $y_g \colon Y_g \longrightarrow X_{k+1}$ and a F+1-precise $p_g \colon E \longrightarrow FY_g+1$ with $g = (Fy_g+1) \cdot p_g$, according to (Ax1). Define $\mathcal{F}_{k+1} := \{(Y_g,y_g) \mid (E,e) \in \mathcal{F}_k, g \in G_{(E,e)}\}.$

We need to prove some properties about the \mathcal{F}_k

- By construction, for every $(E', e') \in \mathcal{F}_{k+1}$, there is some $(E, e) \in \mathcal{F}_k$ and an F + 1-precise map $p_q \colon E \longrightarrow FE' + 1$ with



Consequently by Proposition 3.18, for every $(E, e) \in \mathcal{F}_k$, $k \geq 0$, there is a run $[x_j]_{j \leq k} : J(\mathbf{P}_{k+1}, \mathbf{p}_k) \longrightarrow (X, \xi, x_0)$ with $P_k = E$, $x_k = \operatorname{in}_k \cdot e$.

- For every family \mathcal{F}_k , $k \geq 0$, the morphism $[e]_{(E,e)\in\mathcal{F}_k}$ is an isomorphism; this means that for the factorization:

$$[e]_{(E,e)\in\mathcal{F}_k} \equiv \left(\prod_{(E,e)\in\mathcal{F}_k} E \stackrel{q_k}{\twoheadrightarrow} \bar{X}_k \stackrel{s_k}{\rightarrowtail} X_k\right)$$

we have that s_k is an isomorphism for every $k \geq 0$.

$$\xi_k \cdot e \overset{(\text{Ax4})}{=} \bigsqcup_{g \in G_{(E,e)}} [\eta_{FX_{k+1}}, \bot_{FX_{k+1}}] \cdot g$$

$$\overset{\text{Def. } p_g, y_g}{=} \bigsqcup_{g \in G_{(E,e)}} [\eta_{FX_{k+1}}, \bot_{FX_{k+1}}] \cdot (Fy_g + 1) \cdot p_g$$

$$\overset{(Y_G, y_g) \in \mathcal{F}_{k+1}}{=} \bigsqcup_{g \in G_{(E,e)}} [\eta_{FX_{k+1}}, \bot_{FX_{k+1}}] \cdot (F(s_{k+1} \cdot q_{k+1} \cdot \mathsf{in}_{y_g}) + 1) \cdot p_g$$

$$\overset{\text{Naturality}}{=} \underbrace{\bigsqcup_{g \in G_{(E,e)}} TFs_{k+1} \cdot \left[\eta_{FX'_{k+1}}, \bot_{FX'_{k+1}}\right] \cdot \left(F\left(q_{k+1} \cdot \mathsf{in}_{y_g}\right) + 1\right) \cdot p_g }_{\text{-reflection}} TFs \cdot \underbrace{\bigsqcup_{g \in G_{(E,e)}} \left[\eta_{FX'_{k+1}}, \bot_{FX'_{k+1}}\right] \cdot \left(F\left(q_{k+1} \cdot \mathsf{in}_{y_g}\right) + 1\right) \cdot p_g }_{\text{-short hand } v_{(E,e)} \colon E \longrightarrow TF\bar{X}_{k+1}}$$

Hence, $TFs \cdot v_{(E,e)} = \xi_k \cdot e$ for all $(E,e) \in \mathcal{F}_k$. Since the $e \in \mathcal{F}_k$ are jointly epic by the induction hypothesis, we have a unique diagonal ξ'_k in the following square:

Since ξ_k was constructed to be the least morphism, s_{k+1} is necessarily an isomorphism.

Now we have that $f_k := [e]_{(E,e) \in \mathcal{F}_k} : \coprod_{(E,e) \in \mathcal{F}_k} E \longrightarrow X_k$ is epic for every $k \geq 0$. Since (X, ξ, x_0) is reachable, $[m_k]_{k \geq 0} : \coprod_{k \geq 0} X_k \longrightarrow X$ (cf. item 3 of Lemma A.5) is an epimorphism. Hence, the family $(m_k \cdot f_k)_{k \geq 0}$ is jointly epic; this family is contained in the family of runs in (X, ξ, x_0) , and so the family of runs is jointly epic.

Proof of Proposition 4.3

The first step is to describe when a map is \mathcal{B} -precise:

Lemma A.6. A map $s: X \longrightarrow \mathcal{B}Y$ is \mathcal{B} -precise iff for all $y \in Y$,

$$\sum_{x \in S} s(x)(y) = 1.$$

Proof. Sufficiency is proven by the two inequalities:

 (≤ 1) Let $t_{X,Y} : \mathcal{B}Y \times X \longrightarrow \mathcal{B}(Y \times X)$ be the functorial strength defined as for all Set-functors by $t_{X,Y}(b,x) = \mathcal{B}(y \mapsto (y,x))(b)$. We thus have the commuting diagram

$$X \xrightarrow{\langle s, \mathrm{id}_X \rangle} \mathcal{B}Y \times X \xrightarrow{t_{X,Y}} \mathcal{B}(Y \times X)$$

$$\downarrow \mathcal{B}\pi_1$$

$$\mathcal{B}Y$$

Since s is \mathcal{B} -precise, we obtain a map $d: Y \longrightarrow Y \times X$ with $\pi_1 \cdot d = \operatorname{id}_Y$ and $t_{X,Y} \cdot \langle s, \operatorname{id}_X \rangle = \mathcal{B}d \cdot s$, that means $\mathcal{B}(y \mapsto (y,x))(s(x)) = \mathcal{B}(y \mapsto (y,d'(y)))(s(x))$, with $d' = \pi_2 \cdot d$. Hence, every $y \in Y$ appears in at most one s(x), namely only in s(d'(y)) or not at all; this means for every $y \in Y$, only one summand of $\sum_{x \in S} s(x)(y)$ is non-zero.

To see that every summand of $\sum_{x \in S} s(x)(y)$ is at most 1, define $f: X \longrightarrow \mathcal{B}(\mathbb{N} \times Y)$ by

$$f(x)(n,y) = \begin{cases} 1 & \text{if } n < s(x)(y) \\ 0 & \text{otherwise} \end{cases}$$

and so

$$\mathcal{B}(\pi_2)(f(x))(y) = \sum_{\substack{(n,y) \in \mathbb{N} \times Y \\ n < s(x)(y)}} f(x)(n,y) = \sum_{\substack{(n,y) \in \mathbb{N} \times Y \\ n < s(x)(y)}} 1 = s(x)(y)$$

Hence, we have some map $d: Y \longrightarrow \mathbb{N} \times Y$ with $f = \mathcal{B}(d) \cdot s$ and d(y) = (d'(y), y) for some $d': Y \longrightarrow \mathbb{N}$. The first equality expands to

$$f(x)(n,y) = \mathcal{B}(d)(s(x))(n,y) = \sum_{\substack{y' \in Y \\ d(y') = (n,y)}} s(x)(y') = \begin{cases} s(x)(y) & \text{if } n = d'(y) \\ 0 & \text{otherwise} \end{cases}$$

This implies that for every x and y there is at most one $n \in N$ such that f(x)(n,y) > 0, hence $s(x)(y) \le 1$.

(≥ 1) Assume that $\sum_{x \in S} s(x)(y) = 0$ for some $y \in Y$. Define $s' : X \longrightarrow \mathcal{B}(Y \setminus \{y\})$ by s'(x)(y') = s(x)(y'). With the obvious inclusion $i : Y \setminus \{y\} \hookrightarrow Y$ we have $s = \mathcal{B}i \cdot s'$ and so the \mathcal{B} -precise s induces a map $d : Y \longrightarrow Y \setminus \{y\}$ with $i \cdot d = \mathrm{id}_Y$, a contradiction.

For necessity, consider f and h with

$$\begin{array}{ccc}
X & \xrightarrow{f} & \mathcal{B}C \\
\downarrow s & & \downarrow & \\
\mathcal{B}Y & & & \\
\end{array} \tag{3}$$

By assumption on s we have for all $y \in Y$:

$$1 = \sum_{x \in X} s(x)(y) \stackrel{\text{(3)}}{=} \sum_{x \in X} \sum_{\substack{c \in C \\ h(c) = y}} f(x)(c) = \sum_{\substack{x \in X, c \in C \\ h(c) = y}} f(x)(c)$$

Since all summands of the right-hand sum is are non-negative integers, there exists precisely one $x_y \in X, c_y \in C$ with $h(c_y) = y$ such that $f(x)(c) \neq 0$. Hence, define $d: Y \longrightarrow C$ by this witness: $d(y) = c_y$. This implies directly that h(d(y)) = y and that s(x)(y) = f(x)(d(y)) for all $x \in X, y \in Y$.

where the last equality holds because d is injective.

Now, we want to transfer preciseness from the bag functor to any analytic functor using the natural transformation α . This is done using the following:

Lemma A.7. Let F preserve weak pullbacks and let $\alpha \colon F \longrightarrow G$ be a natural transformation whose naturality squares are weak pullbacks. Then a morphism $f \colon X \longrightarrow FY$ is F-precise iff $\alpha_Y \cdot f$ is G-precise. If G admits precise factorizations w.r.t. $S \subseteq \mathbf{obj} \mathbb{C}$, then so does F.


Proof. For **sufficiency**, let $f: X \longrightarrow FY$ be F-precise and consider a commuting diagram

$$X \xrightarrow{g} GW$$

$$\alpha_Y \cdot f \downarrow \qquad \qquad \downarrow Gw$$

$$GY \xrightarrow{Gy} GZ$$

Since the naturality square for w is a weak pullback, a morphism $g' \colon X \longrightarrow FZ$ exists, making the following diagram commute:



Since f is F-precise, we obtain a morphism $d: Y \longrightarrow W$ with $w \cdot d = y$ and $Fd \cdot f = g'$, and thus also $g = \alpha_W \cdot g' = \alpha_W \cdot Fd \cdot f = Gd \cdot \alpha_Y \cdot f$ as desired.

For **necessity**, we use the simplified version of F-precise because F preserves weak pullbacks. So let $\alpha_Y \cdot f$ be G-precise and consider a commuting diagram

$$X \xrightarrow{g} FZ$$

$$f \downarrow \qquad \qquad Fz$$

$$FY$$


Hence, $Gz \cdot \alpha_Z \cdot g = \alpha_Y \cdot f$ and since $\alpha_Y \cdot f$ is G-precise we obtain a morphism $d\colon Y \longrightarrow Z$ with

$$X \xrightarrow{g} FZ \xrightarrow{\alpha_Z} GZ$$

$$f \downarrow \qquad \qquad \downarrow Gd \qquad \& \quad z \cdot d = id_Y$$


$$FY \xrightarrow{\alpha_Y} GY$$

The naturality square for d is a weak pullback, and so we have some $f': X \longrightarrow FY$ making the following diagrams commute:



So d is indeed a diagonal lifting: $Fd \cdot f = Fd \cdot f' = g$.

For **precise factorizations**, consider $f: X \longrightarrow FY$ with $X \in \mathcal{S}$. For the G-precise factorization $Gy' \cdot g'$ of $\alpha_Y \cdot f$ we have:



Since the naturality square is a weak pullback, we obtain some $f' \colon X \longrightarrow FY'$ with $Fy' \cdot f' = f$ and $Y' \in \mathcal{S}$ as desired.

To conclude, since analytic functors preserve weak pullbacks, it is enough to prove the following:

Lemma A.8. The naturality squares of $\alpha \colon F \longrightarrow \mathcal{B}$ are weak pullback squares.

Proof. Consider a map $f: X \longrightarrow Y$ and elements $t \in FY$, $s \in \mathcal{B}X$ with $\alpha_Y(t) = \mathcal{B}f(s)$. We can write t as an equivalence class $t = [\sigma(y_1, \ldots, y_n)]$. Every y_i is in the image of f, and so any $t' = [\sigma(x_1, \ldots, x_n)] \in FX$ with $f(x_i) = y_i$ for all $1 \le i \le n$ has the property that $\alpha_X(t') = s$ and Ff(t') = t.

So we obtain Proposition 4.3 as a corollary of Lemma A.7.

Auxiliary Lemma in Nominal Sets

In this section, we will use the following standard lemma on strong nominal sets (see e.g. [23, Prop. 5.10] or [19, p. 3]):

Lemma A.9. Given a strong nominal set X, a subset $O \subseteq X$ such that for every $x' \in X$ there is precisely one $x \in X$ with a $\pi \in \mathfrak{S}_{\mathsf{f}}(\mathbb{A})$ fulfilling $x' = \pi \cdot x$, and a map $f \colon O \longrightarrow Y$ into a nominal set Y with $\mathsf{supp}(f(x)) \subseteq \mathsf{supp}(x) \, \forall x \in O$. Then there is a unique equivariant map $f' \colon X \longrightarrow Y$ that agrees with f on O. Such a subset $O \subseteq X$ exists for every nominal set X.

Details for Example 4.5

Factorization w.r.t. strong nominal sets. Let $F: \mathsf{Nom} \longrightarrow \mathsf{Nom}$ be the functor of unordered pairs:

$$FX = \{ p \subseteq X \mid 1 \le |p| \le 2 \}.$$

with element-wise nominal structure. Given a strong nominal set X and an equivariant map $f: X \longrightarrow FY$, define $s: X \longrightarrow F(X + X)$ with

$$s(x) = {\inf(x), \inf(x)}.$$

To see that s is F-precise, consider a commutative square

$$X \xrightarrow{g} FW$$

$$\downarrow s \qquad \downarrow Fu$$

$$F(X+X) \xrightarrow{F[h_1,h_2]} FZ.$$

Spelling out the commutativity, we have:

$$\{h_1(x), h_2(x)\} = F[h_1, h_2](s(x)) = Fu(g(x)) \text{ for all } x \in X.$$

And so $\{h_1(x), h_2(x)\} = Fu(g(x)) = \{u(w_1), u(w_2)\}$ for $g(x) = \{w_1, w_2\}$. For every orbit of X choose some $x \in X$, and let $d_1(x), d_2(x) \in W$ such that $u(d_1(x)) = a_1(x)$ and $u(d_2(x)) = a_2(x)$ and $\{d_1(x), d_2(x)\} = g(x)$ (this involves a finite choice if |g(x)| = 2 and |Fu(g(x))| = 1 but is unique otherwise). Because X is a strong nominal set and by Lemma A.9, d_1 and d_2 extend to equivariant maps $d_1, d_2 \colon X \longrightarrow W$ with $\{d_1(x), d_2(x)\} = g(x)$ and $u(d_i(x)) = h_i(x)$, $i \in \{1, 2\}$. So in total $g = Fd \cdot s$ and $[h_1, h_2] = u \cdot [d_1, d_2]$.

To see that f factors through s, consider g := f, Z = 1, $u = ! : W \longrightarrow Z$, and $h_1 = h_2 = ! : X \longrightarrow 1$, using that $F1 \cong 1$, and so the above square commutes trivially and the diagonal fill in proves that f factors through s as desired.

No factorization w.r.t. all nominal sets. In order to prove that F does not admit precise factorizations w.r.t. all nominal sets, consider the (unique) equivariant map

$$f \colon P \longrightarrow F1 \cong 1 \qquad \text{ with } P = \big\{ \{a,b\} \mid a,b \in \mathbb{A}, a \neq b \big\}$$

Let $s: P \longrightarrow FX$ be its precise factorization, so necessarily $f = F! \cdot s$, with $!: X \longrightarrow 1$. In the following, we derive information about s and X.

As a first aspect, consider the constant equivariant map $t: P \longrightarrow F2$ with $t(\{a,b\}) = \{0,1\}$. Hence,

$$P \xrightarrow{t} F2$$

$$\downarrow s \qquad \downarrow F! \implies \exists d \colon X \longrightarrow 2 \colon Fd \cdot s = t$$

$$FX \xrightarrow{F!} F1$$

For every $\{a,b\} \in P$, $s(\{a,b\}) = \{x_1,x_2\}$ for some $x_1,x_2 \in X$, and $\{d(x_1),d(x_2)\} = \{0,1\}$. Since $0,1 \in 2$ are in different orbits, x_1,x_2 are in different orbits in X, i.e. so for all $\pi \in \mathfrak{S}_{\mathsf{f}}(\mathbb{A})$, $x_1 \neq \pi \cdot x_2$.

As a second aspect, consider $p: P \longrightarrow F(\mathbb{A}^2)$ defined by

$$p(\{a,b\}) = \{(a,b), (b,a)\}.$$

Again, we can use the commutativity:

$$P \xrightarrow{p} F \mathbb{A}^{2}$$

$$\downarrow s \qquad \downarrow f \qquad \downarrow_{F!} \implies \exists u \colon X \longrightarrow \mathbb{A}^{2} \colon Fu \cdot s = p$$

$$FX \xrightarrow{F!} F1$$

For $\{a,b\} \in P$ and $s(\{a,b\}) = \{x_1,x_2\}$ as before, $\{u(x_1),u(x_2)\} = \{(a,b),(b,a)\}$ and the equivariance and functionality of u implies

$$u(x_1) \neq u(x_2) = (a b) \cdot u(x_1) = u((a b) \cdot x_1) \Longrightarrow (a b) \cdot x_1 \neq x_1.$$

Together with $\sigma \cdot x_1 \neq x_2$ from above, we have that $(a b) \cdot x_1 \notin \{x_1, x_2\}$. This leads to a contradiction to the equivariance of s:

$$\{x_1, x_2\} = s(\{a, b\}) = s((ab) \cdot \{a, b\}) = (ab) \cdot s(\{a, b\})$$

$$= (ab) \cdot \{x_1, x_2\} = \{(ab) \cdot x_1, (ab) \cdot x_2\} \neq \{x_1, x_2\}.$$

Proof of Proposition 4.6

Let us check the axioms:

- (Ax2): obtain as in Set.
- (Ax3): We indeed have natural transformations $\eta: X \longrightarrow \mathcal{P}_{\mathsf{ufs}}(X)$ for singleton sets and $\bot: 1 \longrightarrow \mathcal{P}_{\mathsf{ufs}}(X)$ for the empty set.
- (Ax4): consider a equivariant $A \longrightarrow \mathcal{P}_{\mathsf{ufs}}B$ with A strong. Since f is by definition an upper bound for the $\mathcal{F} = \{ [\eta_B, \bot_B] \cdot f' \sqsubseteq f \mid f' \colon A \longrightarrow B+1 \}$, we need to prove that it is the least upper bound. Let g be an upper bound for \mathcal{F} . In order to have $f \sqsubseteq g$ we need to show for every $a \in A$ and $b \in f(a)$ that $b \in g(a)$. Since $b \in f(a)$ and f(a) is ufs, we have $\mathsf{supp}(b) \subseteq \mathsf{supp}(f(a)) \subseteq \mathsf{supp}(a)$. Define $f' \colon A \longrightarrow B+1$ via Lemma A.9, where $O \subseteq A$ is chosen with $a \in O$ and $f'' \colon O \longrightarrow B+1$ is defined by $f''(a) = \mathsf{inl}(b)$ and $f''(x) = \mathsf{inr}(*)$ for $x \neq a$. Now, f' is the unique equivariant map extending f''. Since $[\eta_B, \bot] \cdot f''(x) \subseteq f(x)$ for all $x \in O$, we have that $[\eta_B, \bot] \cdot f' \sqsubseteq f$, and so $[\eta_B, \bot] \cdot f' \in \mathcal{F}$, and thus $[\eta_B, \bot] \cdot f' \sqsubseteq g$, hence $b \in g(a)$.
- (Ax5): consider a weakly commuting square:

$$A \xrightarrow{x} \mathcal{P}_{\mathsf{ufs}} X$$

$$y \downarrow \qquad \qquad \downarrow \mathcal{P}_{\mathsf{ufs}} h$$

$$Y \xrightarrow{\eta_{Y}} \mathcal{P}_{\mathsf{ufs}} Y$$

where A is strong. Pick any subset $O \subseteq A$ fulfilling the assumption of Lemma A.9. For each $a \in A$ we have that $y(a) \in h[x(a)]$, so for each $a \in O$, there exists some $a' \in x(a)$ with h(a') = y(a). For each $a \in O$, denote this witness by x'(a) := a'. By Lemma A.9, x' extendes to an equivariant map $c : A \longrightarrow X$ with $c(a) = x'(a) \in x(a)$ for all $a \in O$. For every $b \in A$, there is some $\pi \in \mathfrak{S}_{\mathbf{f}}(\mathbb{A})$ and $a \in A$ with $b = \pi \cdot a$, and hence $x'(b) = x'(\pi \cdot a) = \pi \cdot x'(a) \in \pi \cdot x(a) = x(b)$, i.e. $\eta_X \cdot x' \subseteq x$. By construction of c, we have that $h \cdot x' = y$.

Proof of Proposition 4.8

To prove this, by Proposition 3.6.4, it is enough to prove that the binding functor [A] admits precise factorizations w.r.t. strong nominal sets.

Recall from [26] that $[\mathbb{A}]$ has the left adjoint $\mathbb{A}\#_-$, sending X to the nominal set $\mathbb{A}\#X := \{(a,x) \in \mathbb{A} \times X \mid a \not\in \mathsf{supp}(x)\}$. If X is a strong nominal set, then so is $\mathbb{A}\#X$, hence $[\mathbb{A}]$ admits precise factorizations w.r.t. strong nominal sets by Proposition 3.6.5.

Proof of Proposition 4.9

All the axioms are proved as in Set, component-wise. The only interesting axiom is to prove that $F\{X_P\}_{P\in\mathbb{P}}=\{\coprod_{Q\in\mathbb{P}}\mathbb{P}(P,Q)\times X_Q\}_{P\in\mathbb{P}}$ admits precise factorisations. Since $F=\coprod_{Q\in\mathbb{P}}H_Q\times\pi_Q$ with:

- H_Q being the constant functor on $(\mathbb{P}(P,Q))_{P\in\mathbb{P}}$,
- π_Q being the functor mapping $\{X_P\}_{P\in\mathbb{P}}$ to $\{X_Q\}_{P\in\mathbb{P}}$,

by Proposition 3.6.(1,3,4), it is enough to prove that π_Q admits precise factorisations.

For that purpose, given a morphism, i.e. family of maps, $(f_P)_{P\in\mathbb{P}}: \{X_P\}_{P\in\mathbb{P}} \to \pi_Q(\{Y_P\}_{P\in\mathbb{P}}) = \{Y_Q\}_{P\in\mathbb{P}}$, define

$$(Z_P)_{P\in\mathbb{P}}$$
 with $Z_Q=\coprod_{P\in\mathbb{P}}X_P$ and $Z_P=\emptyset$ for $P\neq Q$.

and $(\mathsf{in}_P)_{P\in\mathbb{P}} \colon \{X_P\}_{P\in\mathbb{P}} \longrightarrow \pi_Q(Z_P)_{P\in\mathbb{P}}$

$$\operatorname{in}_P\colon X_P \longrightarrow (\pi_Q(Z_P)_{P\in\mathbb{P}})_P = Z_Q = \coprod_{P\in\mathbb{P}} X_P$$

and $(h_P)_{P\in\mathbb{P}}: (Z_P)_{P\in\mathbb{P}} \longrightarrow (Y_P)_{P\in\mathbb{P}}$ with

$$h_Q = [f_P]_{P \in \mathbb{P}} \colon \coprod_{P \in \mathbb{P}} X_P \longrightarrow Y_Q, \quad h_P = ! \colon \emptyset \longrightarrow Y_Q \text{ for } P \neq Q.$$

Obviously, $(\pi_Q h)_P \cdot \mathsf{in}_P = f_P$ for all $P \in \mathbb{P}$. It remains to show that $(\mathsf{in}_P)_{P \in \mathbb{P}}$ is π_Q -precise. Consider a commutative square:

$$\begin{split} &(X_P)_{P\in\mathbb{P}} \xrightarrow{\quad (v_P)_{P\in\mathbb{P}} \quad} \pi_Q((V)_{P\in\mathbb{P}}) \\ &(\operatorname{in}_P)_{P\in\mathbb{P}} \downarrow \qquad \qquad \downarrow^{\pi_Q((w_P)_{P\in\mathbb{P}}) = w_Q} \\ &\pi_Q((Z)_{P\in\mathbb{P}}) \xrightarrow{\pi_Q((z_P)_{P\in\mathbb{P}}) \quad} \pi_Q((W)_{P\in\mathbb{P}}) \end{split} \tag{*}$$

Define a diagonal $(d_P)_{P\in\mathbb{P}}: (Z_P)_{P\in\mathbb{P}} \longrightarrow (V_P)_{P\in\mathbb{P}}$ by

$$d_Q = [v_P]_{P \in \mathbb{P}} \colon \coprod_{P \in \mathbb{P}} X_P \longrightarrow V_Q \quad d_P = ! \colon Z_P \longrightarrow V_P \text{ for } P \neq Q.$$

Obviously, we have the first triangle $v_P = d_Q \cdot \text{in}_P$ for all $P \in \mathbb{P}$. The other triangle follows by case distinction: for $P \neq Q$, $z_P = w_P \cdot d_P$ holds by initiallity of $Z_P = \emptyset$; we have

$$z_Q \cdot \operatorname{in}_P \stackrel{(*)}{=} w_Q \cdot v_P = w_Q \cdot d_Q \cdot \operatorname{in}_P \quad \forall P \in \mathbb{P}$$

and since the $\operatorname{in}_P \colon X_P \longrightarrow Z_Q$ are jointly epic, $z_Q = w_Q \cdot d_Q$. So $(d_P)_{P \in \mathbb{P}}$ is a diagonal lifting and so $(\operatorname{in}_P)_{P \in \mathbb{P}}$ is π_Q -precise.

Proof of Proposition 4.10

To prove this proposition, it is enough to prove the following:

Lemma A.10. A morphism $f: \chi^P \longrightarrow FY$ is F-precise iff $Y = \chi^Q$ for some $Q \in \mathbb{P}$.

Proof. Necessity. Consider $f_P \colon 1 \longrightarrow \coprod_{Q \in \mathbb{P}} \mathbb{P}(P,Q) \times Y_Q$ and let $f_P = \operatorname{in}_Q(m,x)$. Define the injective $h \colon \chi^Q \longrightarrow (Y_R)_{R \in \mathbb{P}}$ by $h_R = ! \colon \emptyset \longrightarrow Y_R$ for $R \neq Q$ and $h_Q = x \colon 1 \longrightarrow X_Q$. Furthermore, define by $g \colon \chi^P \longrightarrow F\chi^Q$ $g_P = \operatorname{in}_Q(m,*)$, and so $Fh \cdot g = f$. Since f is F-precise, there is a $d \colon Y \longrightarrow \chi^Q$ with $h \cdot d = \operatorname{id}_Y$, so in total h is an isomorphism.

Sufficiency. For $f: \chi^P \longrightarrow F\chi^Q$ let $m: P \longrightarrow Q$ with $f_P = \operatorname{in}_Q(m, *)$ and consider a commuting diagram:

$$\chi^{P} \xrightarrow{w} FW$$

$$f \downarrow \qquad \qquad \downarrow_{Fh} \text{ in } \mathsf{Set}^{\mathbf{obj}\,\mathbb{P}}$$

$$F\chi^{Q} \xrightarrow{Fz} FZ$$


And so in component P we have:

$$1 \xrightarrow{w_P} \coprod_{R \in \mathbb{P}} \mathbb{P}(P,R) \times W_R \\ \downarrow^{F_h} \qquad \qquad \downarrow^{F_h} \qquad \text{in Set} \\ \mathbb{P}(P,Q) \times \chi_Q^Q \xrightarrow{\text{in}_Q} \coprod_{R \in \mathbb{P}} \mathbb{P}(P,R) \times \chi_R^Q \xrightarrow{F_Z} \coprod_{R \in \mathbb{P}} \mathbb{P}(P,R) \times Z_R$$

Hence, (formally by the extensivity of Set) w_P is necessarily $w_P = \operatorname{in}_Q(m, x)$ for some $x \in W_Q$ with $h_Q(x) = z_Q(*)$. Define $d: \chi^Q \longrightarrow W$ by $d_Q: 1 \longrightarrow W_Q$, $d_Q = x$, and $d_R = !: \emptyset \longrightarrow W_R$ for $R \neq Q$. Obviously d fulfils both $Fd \cdot f = w$ and $h \cdot d = z$.

Notes on section 5

In probabilistic systems, it is not clear how paths should look like. Consider the probabilistic system A:



There are different possibilities how to model a path in such a system:

- 1. Simply $\bullet \longrightarrow \bullet \longrightarrow \bullet$ as in LTS for a singleton input alphabet. Of course, this makes the probabilities in the system entirely meaningless for runs.
- 2. A path and a probability $(p, \bullet \longrightarrow \bullet \longrightarrow \bullet)$ meaning that there is a path whose weights multiply to at least p. So the above system has a run for such a path if $p \le 0.08 = 0.2 \cdot 0.4$.
- 3. A path is a probabilistic system in which each state has at most one successor, and so the notion of run of path coincides with a functional simulation from that path to the system. E.g. the above system has a run for \bullet $\xrightarrow{0.1} \bullet$ $\xrightarrow{0.25} \bullet$. Clearly, there is a functional bisimulation $f: A \longrightarrow X$ from the above system to the system X given by

$$\longrightarrow x \xrightarrow{0.5} y \xrightarrow{0.7} z.$$

While A has no run for the path $\bullet \xrightarrow{0.5} \bullet$, the system X has. So $f: A \longrightarrow X$ is not an open map, even though it is a functional bisimulation, i.e. coalgebra homomorphism for the subdistribution functor.