

SCHUR'S COLOURING THEOREM FOR NON-COMMUTING PAIRS

TOM SANDERS

ABSTRACT. For G a finite non-Abelian group we write $c(G)$ for the probability that two randomly chosen elements commute and $k(G)$ for the largest integer such that any $k(G)$ -colouring of G is guaranteed to contain a monochromatic quadruple (x, y, xy, yx) with $xy \neq yx$. We show that $c(G) \rightarrow 0$ if and only if $k(G) \rightarrow \infty$.

1. INTRODUCTION

Our starting point is Schur's theorem [Sch16, Hilfssatz], the proof of which adapts to give the following.

Theorem 1.1. *Suppose that G is a finite group and \mathcal{C} is a cover of G of size k . Then there is a set $A \in \mathcal{C}$ with at least $c_k |G|^2$ triples $(x, y, xy) \in A^3$ where c_k is a constant depending only on k .*

The proof is a routine adaptation, but we shall not give it as the result as stated also follows from our next theorem.

If G is non-Abelian then we might like to ask for quadruples $(x, y, xy, yx) \in A^4$ instead of triples and establishing the following such result (which we do in §2) is the main purpose of the paper.

Theorem 1.2. *Suppose that G is a finite group and \mathcal{C} is a cover of G of size k . Then there is a set $A \in \mathcal{C}$ with $c_k |G|^2$ quadruples $(x, y, xy, yx) \in A^4$ where c_k is a constant depending only on k .*

When G is non-Abelian we should like to ensure that at least one of the quadruples found in Theorem 1.2 has $xy \neq yx$, and to this end we define the **commuting probability** of a finite group G to be

$$(1.1) \quad c(G) := \frac{1}{|G|^2} \sum_{x, y \in G} 1_{[xy=yx]}$$

in words it is the probability that a pair $(x, y) \in G^2$ chosen uniformly at random has $xy = yx$. There are many nice results about the commuting probability – see the introduction to [Heg13] for details – and it is an instructive exercise (see [Gus73]) to check that if $c(G) < 1$ then $c(G) \leq \frac{5}{8}$, so that if a group is non-Abelian there are ‘many’ pairs that do not commute. Despite this we have the following which we prove in §3.

Proposition 1.3. *Suppose that G is a finite group and $c(G) \geq \epsilon$. Then there is a cover \mathcal{C} of G of size $\exp((2 + o_{\epsilon \rightarrow 0}(1))\epsilon^{-1} \log \epsilon^{-1})$ such that if $A \in \mathcal{C}$ and $(x, y, xy, yx) \in A^4$ then $xy = yx$.*

If G is non-Abelian we write $k(G)$ for the **non-commuting Schur number** of G , that is the largest natural number such that for any cover \mathcal{C} of G of size $k(G)$ there is some $A \in \mathcal{C}$ and $(x, y, xy, yx) \in A^4$ with $xy \neq yx$. (Note that since G is assumed non-Abelian we certainly have $k(G) \geq 1$.)

The number $k(G)$ has been studied for a range of specific groups by McCutcheon in [McC] and we direct the interested reader there for examples and further questions.

Theorem 1.4. *Suppose that $(G_n)_n$ is a sequence of non-Abelian groups. Then $c(G_n) \rightarrow 0$ if and only if $k(G_n) \rightarrow \infty$.*

Proof. The right to left implication follows immediately from Proposition 1.3. We can assume that c_k is monotonically decreasing. Suppose that $c(G_n) \rightarrow 0$ and there is a k_0 and an infinite set S of naturals such that $k(G_n) < k_0$ for all $n \in S$. Let $n \in S$ be such that $c(G_n) < c_{k_0}$ which can be done since $c(G_n) \rightarrow 0$ and $c_{k_0} > 0$.

Since $k(G_n) < k_0$ there is a cover \mathcal{C} of G_n of size k_0 such that if $A \in \mathcal{C}$ and $(x, y, xy, yx) \in A^4$ then $xy = yx$. By Theorem 1.2 there is $A \in \mathcal{C}$ such that $(x, y, xy, yx) \in A^4$ for at least $c_{k_0}|G_n|^2$ quadruples. But then by design $xy = yx$ for all these pairs and so $c(G_n) \geq c_{k_0}$, a contradiction which proves the result. \square

Before closing the section we need to acknowledge our debt to previous work. In [McC12] McCutcheon proves that $k(S_n) \rightarrow \infty$ as $n \rightarrow \infty$. A short calculation shows that $c(S_n) \rightarrow 0$ as $n \rightarrow \infty$, and the possibility of showing that $k(G_n) \rightarrow \infty$ as $c(G_n) \rightarrow 0$ is identified by Bergelson and Tao in the remarks after [BT14, Theorem 11]. Earlier, in [BT14, Footnote 4], they also observe the significance of Neumann's work [Neu89] which is the main idea behind the proof of Proposition 1.3.

Write $D(G)$ for the smallest dimension of a non-trivial unitary representation of G .¹ In [BT14, Corollary 8] the authors show that $k(G_n) \rightarrow \infty$ as $D(G_n) \rightarrow \infty$, and in fact go further proving a density result. For general finite groups there can be no density result; we refer the reader to the discussion after [BT14, Theorem 11] for more details.

2. PROOF OF THEOREM 1.2

The proof of Theorem 1.2 is inspired by an attempt to translate the proof of [BM98, Theorem 3.4] into a combinatorial setting. There the authors use a recurrence theorem [BMZ97, Theorem 5.2]; in its place we use a version of the Ajtai-Szemerédi Corners Theorem [AS74] for finite groups. This was proved by Solymosi [Sol13, Theorem 2.1] using the triangle removal lemma.

Theorem 2.1. *There is a function $f_\Delta : (0, 1] \rightarrow (0, 1]$ such that if G is a finite group and $\mathcal{A} \subset G^2$ has size at least $\alpha|G|^2$ then*

$$S(\mathcal{A}) := \frac{1}{|G|^3} \sum_{x, y, z \in G} 1_{\mathcal{A}}(x, y) 1_{\mathcal{A}}(zx, y) 1_{\mathcal{A}}(x, yz) \geq f_\Delta(\alpha).$$

¹This is called the quasirandomness of G in [BT14, Definition 1] following the work of Gowers [Gow08]

Proof. Following the proof of [Sol13, Theorem 2.1], form a tripartite graph with three copies of G as the vertex sets (call them V_1, V_2, V_3) and joining $(x, y) \in V_1 \times V_2$ if and only if $(x, y) \in \mathcal{A}$; $(y, w) \in V_2 \times V_3$ if and only if $(y^{-1}w, y) \in \mathcal{A}$; and $(x, w) \in V_1 \times V_3$ if and only if $(x, wx^{-1}) \in \mathcal{A}$. The map $G^3 \rightarrow G^3; (x, y, w) \mapsto (x, y, y^{-1}wx^{-1})$ is a bijection and (x, y, w) is a triangle in this graph if and only if $(x, y), (zx, y), (x, yz) \in \mathcal{A}$ where $z = y^{-1}wx^{-1}$.

It follows from [Tao06, Theorem 1.1] that one can remove at most $3 \cdot o_{S(\mathcal{A}) \rightarrow 0}(|G|^2) = o_{S(\mathcal{A}) \rightarrow 0}(|G|^2)$ elements from \mathcal{A} to make the graph triangle-free. On the other hand if $(x, y) \in \mathcal{A}$ then (x, y, xy) is a triangle in the above graph, hence we must have removed all elements from \mathcal{A} and $\alpha|G|^2 \leq o_{S(\mathcal{A}) \rightarrow 0}(|G|^2)$ from which the result follows. \square

There are a number of subtleties around the extent to which one can replace, say, (zx, y) with (xz, y) , and we refer the reader to the papers of Solymosi [Sol13] and Austin [Aus16] for some discussion.

We take the convention, as we can, that the function f_Δ is monotonically increasing and $f_\Delta(x) \leq x$ for all $x \in (0, 1]$. Even with Fox's work [Fox11] we only have $f_\Delta(\alpha)^{-1} \leq T(O(\log \alpha^{-1}))$ in general. However, when G is Abelian much better bounds are known as a result of the beautiful arguments of Shkredov [Shk06b, Shk06a, Shk09]. It seems likely these could be adapted to give a bound with a tower of bounded height if the Fourier analysis is adapted to the non-Abelian setting in the same way as it is for Roth's theorem in [San17]. Doing so would give a quantitative version of [BT14, Theorem 10] (see [BT14, Remark 44]), but the improvement to Theorem 1.2 would only be to replace a wowzer-type function with a tower as we shall see shortly.

We shall prove the following proposition from which Theorem 1.2 follows immediately on inserting the bound for f_Δ given by Theorem 2.1.

Proposition 2.2. *Suppose G is a finite group and \mathcal{C} is a cover of G of size k . Then there is a set $A \in \mathcal{C}$ with $(g^{(k+1)}(1))^2|G|^2$ quadruples $(x, y, xy, yx) \in A^4$ where $g^{(k+1)}$ is the $(k+1)$ -fold composition of g with itself, and $g : (0, 1] \rightarrow (0, 1]; \alpha \mapsto (3k)^{-1}f_\Delta(\alpha^k)$.*

Proof. Write A_1, \dots, A_k for the sets in \mathcal{C} ordered so that they have density $\alpha_1 \geq \dots \geq \alpha_k$ respectively; since \mathcal{C} is a cover we have $\alpha_1 \geq \frac{1}{k}$. Let $r \in \{1, \dots, k\}$ be minimal such that

$$(2.1) \quad \frac{1}{3}f_\Delta(\alpha_1 \cdots \alpha_r) \geq \alpha_{r+1} + \cdots + \alpha_k,$$

which is possible since the sum on the right is empty and so 0 when $r = k$. From minimality and the order of the α_i s we have

$$\alpha_{i+1} > \frac{1}{3k}f_\Delta(\alpha_1 \cdots \alpha_i) \text{ for all } 1 \leq i \leq r-1.$$

The function f_Δ is monotonically increasing and $f_\Delta(x) \leq x$ for all $x \in (0, 1]$ so it follows from the above that $\alpha_r \geq g^{(r)}(1) \geq g^{(k)}(1)$.

Now, suppose that $s_1, \dots, s_r \in G$ and write

$$\mathcal{A}_i := \{(x, y) \in G^2 : xs_iy \in A_i\} \text{ for } 1 \leq i \leq r.$$

Then

$$\mathbb{E}_{s_i \in G} 1_{\mathcal{A}_i}(x, y) = \alpha_i \text{ for all } x, y \in G \text{ and } 1 \leq i \leq r,$$

and so

$$\mathbb{E}_{s \in G^r} \left| \bigcap_{i=1}^r \mathcal{A}_i \right| = \sum_{x, y \in G} \mathbb{E}_{s \in G^r} \prod_{i=1}^r 1_{\mathcal{A}_i}(x, y) = \alpha_1 \cdots \alpha_r |G|^2.$$

By averaging we can pick some $s \in G^r$ such that $\mathcal{A} := \bigcap_{i=1}^r \mathcal{A}_i$ has $|\mathcal{A}| \geq \alpha_1 \cdots \alpha_r |G|^2$.

By the definition of f_Δ (from Theorem 2.1) we have

$$\mathbb{E}_{x, y, z \in G} 1_{\mathcal{A}}(x, y) 1_{\mathcal{A}}(zx, y) 1_{\mathcal{A}}(x, yz) = S(\mathcal{A}) \geq f_\Delta(\alpha_1 \cdots \alpha_r);$$

write

$$Z := \left\{ z \in G : \mathbb{E}_{x, y \in G} 1_{\mathcal{A}}(x, y) 1_{\mathcal{A}}(zx, y) 1_{\mathcal{A}}(x, yz) \geq \frac{1}{3} f_\Delta(\alpha_1 \cdots \alpha_r) \right\}.$$

Then

$$\begin{aligned} \mathbb{P}(Z) + \frac{1}{3} f_\Delta(\alpha_1 \cdots \alpha_r) &\geq \mathbb{E}_{x, y, z \in G} 1_{Z \sqcup (G \setminus Z)}(z) 1_{\mathcal{A}}(x, y) 1_{\mathcal{A}}(zx, y) 1_{\mathcal{A}}(x, yz) \\ &= S(\mathcal{A}) \geq f_\Delta(\alpha_1 \cdots \alpha_r), \end{aligned}$$

and hence $\mathbb{P}(Z) \geq \frac{2}{3} f_\Delta(\alpha_1 \cdots \alpha_r)$. But then

$$\begin{aligned} \mathbb{P}(Z \setminus (A_{r+1} \cup \cdots \cup A_k)) &\geq \frac{2}{3} f_\Delta(\alpha_1 \cdots \alpha_r) - (\alpha_{r+1} + \cdots + \alpha_k) \\ &\geq \frac{1}{3} f_\Delta(\alpha_1 \cdots \alpha_r) \end{aligned}$$

by (2.1). Since $\bigcup_{i=1}^k A_i = G$ we conclude that there is some $1 \leq i \leq k$ such that

$$\mathbb{P}((Z \setminus (A_{r+1} \cup \cdots \cup A_k)) \cap A_i) \geq \frac{1}{3r} f_\Delta(\alpha_1 \cdots \alpha_r).$$

If course $(Z \setminus (A_{r+1} \cup \cdots \cup A_k)) \cap A_j = \emptyset$ for $r < j \leq k$ and so we may assume $i \leq r$.

Write $Z' := (Z \setminus (A_{r+1} \cup \cdots \cup A_k)) \cap A_i$. Since $Z' \subset Z$ we have

$$\mathbb{E}_{x, y} 1_{\mathcal{A}_i}(x, y) 1_{\mathcal{A}_i}(zx, y) 1_{\mathcal{A}_i}(x, yz) \geq \mathbb{E}_{x, y} 1_{\mathcal{A}}(x, y) 1_{\mathcal{A}}(zx, y) 1_{\mathcal{A}}(x, yz) \geq \frac{1}{3} f_\Delta(\alpha_1 \cdots \alpha_r)$$

for all $z \in Z'$. On the other hand every $z \in Z'$ has $z \in A_i$ and so we conclude that there are at least

$$\frac{1}{3} f_\Delta(\alpha_1 \cdots \alpha_r) |G|^2 \cdot \frac{1}{3r} f_\Delta(\alpha_1 \cdots \alpha_r) |G|$$

triples $(x, y, z) \in G^3$ such that

$$z \in A_i, xs_i y \in A_i, zxs_i y \in A_i, \text{ and } xs_i yz \in A_i.$$

The map $(x, y, z) \mapsto (xs_i y, z)$ has all fibres of size $|G|$ and so there are at least

$$\frac{1}{9r} f_\Delta(\alpha_1 \cdots \alpha_r)^2 |G|^2 \geq (g(\alpha_r))^2 |G|^2$$

pairs $(a, b) \in G^2$ such that $a, b, ab, ba \in A_i$. This gives the result. \square

3. PROOF OF PROPOSITION 1.3

The key idea comes from Neumann's theorem [Neu89, Theorem 1] which is already identified in [BT14, Footnote 4]. Neumann's theorem describes the structure of groups G for which $c(G) \geq \epsilon$ – they are the groups containing normal subgroups $K \leq H \leq G$ such that K and G/H have size $O_\epsilon(1)$ and H/K is Abelian. Neumann's theorem was further developed in [Ebe15, Theorem 2.4], but both arguments provide a more detailed structure than we require.

We have made some effort to control the exponent; results such as [Ebe15, Lemma 2.1] or [Ols76, Theorem 2.2] could be used in place of Kemperman's Theorem in what follows at the possible expense of the 2 becoming slightly larger. Moving the $2 + o_{\epsilon \rightarrow 0}(1)$ below 1 would require a slightly different approach as we normalise a subgroup of index around ϵ^{-1} at a certain point which costs us a term of size ϵ^{-1} .

Proposition (Proposition 1.3). *Suppose that G is a finite group and $c(G) \geq \epsilon$. Then there is a cover \mathcal{C} of G of size $\exp((2 + o_{\epsilon \rightarrow 0}(1))\epsilon^{-1} \log \epsilon^{-1})$ such that if $A \in \mathcal{C}$ and $(x, y, xy, yx) \in A^4$ then $xy = yx$.*

Proof. We work with the conjugation action of G on itself (*i.e.* $(g, x) \mapsto g^{-1}xg$), and write x^G for the conjugacy class of x (the orbit of x under this action), and $C_G(x)$ for the centre of x in G (the stabiliser of x under this action).

Let $\eta, \nu \in (0, 1]$ be parameters (we shall take $\nu = \frac{1}{2}$ and $\eta = \epsilon / \log \epsilon^{-1}$) to be optimised later and put

$$X := \{x \in G : |x^G| \leq \eta^{-1}\}.$$

Then

$$\epsilon|G|^2 \leq |G|^2 \mathbb{P}(xy = yx) = \sum_x |C_G(x)| = |G| \sum_x \frac{1}{|x^G|} \leq \sum_{x \in X} |G| + \sum_{x \notin X} \eta |G|.$$

Writing $\kappa := |X|/|G|$ we can rearrange the above to see that $\kappa \geq (\epsilon - \eta)/(1 - \eta)$.

Suppose that $s \in \mathbb{N}$ is maximal such that

$$|\overbrace{X \cdots X}^{s \text{ times}}| \geq (1 + (1 - \nu)(s - 1)) |X|.$$

Certainly there is some $s \in \mathbb{N}$ since the inequality certainly holds for $s = 1$, and there is a maximal such with $s \leq \frac{\kappa^{-1} - \nu}{1 - \nu}$ since $|X| \geq \kappa |G|$.

Since $1_G^G = \{1_G\}$ we have $1_G \in X$ and $1_G \in X \cdots X$ for any s -fold product. By Kemperman's Theorem [Kem56, Theorem 5] (also recorded on [Ols84, p111], and which despite the additive notation does not assume commutativity) it follows that there is some $H \leq G$ such that

$$|\overbrace{X \cdots X}^{s+1 \text{ times}}| \geq |\overbrace{X \cdots X}^{s \text{ times}}| + |X| - |H| \text{ and } H \subset \overbrace{X \cdots X}^{s+1 \text{ times}}.$$

By maximality of s we know that

$$(1 + (1 - \nu)s) |X| > |\overbrace{X \cdots X}^{s+1 \text{ times}}| \geq (1 + (1 - \nu)(s - 1)) |X| + |X| - |H|,$$

so that $|H| > \nu|X|$, and so $|G/H| < \nu^{-1}\kappa^{-1}$.

Let K be the kernel of the action of left multiplication by G on G/H *i.e.* $K := \{x \in G : xgH = gH \text{ for all } g \in G\}$. The action induces a homomorphism from G to $\text{Sym}(G/H)$ so that by the First Isomorphism Theorem

$$K \triangleleft G \text{ and } |G/K| \leq |\text{Sym}(G/H)| \leq |G/H|!.$$

Each $x \in H$ (and hence each $x \in K$ since $xH = H$ for such x) can be written as a product of $s+1$ elements of X . Moreover, the function $x \mapsto |x^G|$ is sub-multiplicative *i.e.* $|(xy)^G| \leq |x^G||y^G|$ and so it follows that

$$|x^G| \leq \eta^{-(s+1)} \leq R := \left\lfloor \eta^{-\frac{\kappa^{-1}+1-2\nu}{1-\nu}} \right\rfloor \text{ for all } x \in X^{s+1},$$

and in particular for all $x \in K$. Thus for each $x \in K$ there is an injection $\phi_{x^G} : x^G \rightarrow \{1, \dots, R\}$. With this notation we can define our covering: let

$$\mathcal{S} := \{\{x \in K : \phi_{x^G}(x) = i\} : 1 \leq i \leq R\} \text{ and } \mathcal{C} := ((G/K) \setminus \{K\}) \cup \mathcal{S},$$

so that \mathcal{S} is a cover of K and \mathcal{C} is a cover of G . Now

$$\begin{aligned} |\mathcal{C}| &\leq |G/K| - 1 + |\mathcal{S}| \leq \lfloor \nu^{-1}\kappa^{-1} \rfloor! - 1 + R \\ &\leq \exp \left(\max \left\{ \nu^{-1}\kappa^{-1} \log \nu^{-1}\kappa^{-1}, \frac{\kappa^{-1} + 1 - 2\nu}{1 - \nu} \log \eta^{-1} \right\} + O(1) \right). \end{aligned}$$

Optimise this by taking $\nu = \frac{1}{2}$ and $\eta = \epsilon / \log \epsilon^{-1}$ as mentioned before so that $\kappa \geq \epsilon(1 - o_{\epsilon \rightarrow 0}(1))$ and $\log \eta^{-1} = (1 + o_{\epsilon \rightarrow 0}(1)) \log \epsilon^{-1}$.

Suppose that $A \in \mathcal{C}$ and $x, y, xy, yx \in A$. If $A \in (G/K) \setminus \{K\}$ then $xK = yK = xyK = yxK = A$. Since $K \triangleleft G$ we have $xK = xyK = (xK)(yK)$ and so $yK = K$ which is a contradiction. It follows that $A \in \mathcal{S}$ and hence $x, y, xy, yx \in K$. We conclude $\phi_{(xy)^G}(xy) = \phi_{(yx)^G}(yx)$ but $xy = y^{-1}(yx)y$ and so $(xy)^G = (yx)^G$. Since $\phi_{(xy)^G}$ is an injection we conclude that $xy = yx$ as required.

The result is proved. \square

REFERENCES

- [AS74] M. Ajtai and E. Szemerédi. Sets of lattice points that form no squares. *Stud. Sci. Math. Hungar.*, 9:9–11 (1975), 1974. URL <https://dcm.epfl.ch/wp-content/uploads/2018/10/AjtaiSzemeredi-1974-SetsOfLatticePointsThatFormNo>
- [Aus16] T. Austin. Ajtai-Szemerédi theorems over quasirandom groups. In *Recent trends in combinatorics*, volume 159 of *IMA Vol. Math. Appl.*, pages 453–484. Springer, [Cham], 2016, arXiv:1503.08746. doi:10.1007/978-3-319-24298-9_19.
- [BM98] V. Bergelson and R. McCutcheon. Recurrence for semigroup actions and a non-commutative Schur theorem. In *Topological dynamics and applications (Minneapolis, MN, 1995)*, volume 215 of *Contemp. Math.*, pages 205–222. Amer. Math. Soc., Providence, RI, 1998. doi:10.1090/conm/215/02942.
- [BMZ97] V. Bergelson, R. McCutcheon, and Q. Zhang. A Roth theorem for amenable groups. *Amer. J. Math.*, 119(6):1173–1211, 1997. doi:10.1353/ajm.1997.0035.
- [BT14] V. Bergelson and T. C. Tao. Multiple recurrence in quasirandom groups. *Geometric and Functional Analysis*, 24(1):1–48, Feb 2014, arXiv:1211.6372. doi:10.1007/s00039-014-0252-0.

- [Ebe15] S. Eberhard. Commuting probabilities of finite groups. *Bulletin of the London Mathematical Society*, 47(5):796–808, 2015, arXiv:1411.0848. doi:10.1112/blms/bdv050.
- [Fox11] J. Fox. A new proof of the graph removal lemma. *Ann. of Math. (2)*, 174(1):561–579, 2011, arXiv:1006.1300. doi:10.4007/annals.2011.174.1.17.
- [Gow08] W. T. Gowers. Quasirandom groups. *Comb. Probab. Comput.*, 17(3):363–387, 2008, arXiv:0710.3877. doi:10.1017/S0963548307008826.
- [Gus73] W. H. Gustafson. What is the probability that two group elements commute? *The American Mathematical Monthly*, 80(9):1031–1034, 1973. doi:10.2307/2318778.
- [Heg13] P. Hegarty. Limit points in the range of the commuting probability function on finite groups. *J. Group Theory*, 16(2):235–247, 2013, arXiv:1207.0760. doi:10.1515/jgt-2012-0040.
- [Kem56] J. H. B. Kemperman. On complexes in a semigroup. *Nederl. Akad. Wetensch. Proc. Ser. A.*, 59:247–254, 1956. URL <http://www.math.haifa.ac.il/~seva/MathOverflow/OnComplexes.pdf>.
- [McC] R. McCutcheon. Non-commutative Schur configurations in finite groups. URL <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.538.5511&rep=rep1&type=pdf>.
- [McC12] R. McCutcheon. Monochromatic permutation quadruples—a Schur thing in S_n . *The American Mathematical Monthly*, 119(4):342–343, 2012. doi:10.4169/amer.math.monthly.119.04.342.
- [Neu89] P. M. Neuman. Two combinatorial problems in group theory. *Bulletin of the London Mathematical Society*, 21(5):456–458, 1989. doi:10.1112/blms/21.5.456.
- [Ols84] J. E. Olson. On the sum of two sets in a group. *Journal of Number Theory*, 18(1):110 – 120, 1984. doi:10.1016/0022-314X(84)90047-7.
- [Ols76] J. E. Olson. Sums of sets of group elements. *Acta Arith.*, 28(2):147–156, 1975/76. doi:10.4064/aa-28-2-147-156.
- [San17] T. Sanders. Solving $xz = y^2$ in certain subsets of finite groups. *Q. J. Math.*, 68(1):243–273, 2017, arXiv:1611.05632. doi:10.1093/qmath/haw056.
- [Sch16] I. Schur. Über die Kongruenz $x^m + y^m \equiv z^m \pmod{p}$. *Jahresber. Dtsch. Math.-Ver.*, 25:114–117, 1916. URL <http://eudml.org/doc/145475>.
- [Shk06a] I. D. Shkredov. On a generalization of Szemerédi’s theorem. *Proc. London Math. Soc. (3)*, 93(3):723–760, 2006, arXiv:math/0503639. doi:10.1017/S0024611506015991.
- [Shk06b] I. D. Shkredov. On a problem of Gowers. *Izv. Ross. Akad. Nauk Ser. Mat.*, 70(2):179–221, 2006, arXiv:math/0405406. doi:10.1070/IM2006v070n02ABEH002316.
- [Shk09] I. D. Shkredov. On a two-dimensional analogue of Szemerédi’s theorem in abelian groups. *Izv. Ross. Akad. Nauk Ser. Mat.*, 73(5):181–224, 2009, arXiv:0705.0451. doi:10.1070/IM2009v073n05ABEH002472.
- [Sol13] J. Solymosi. Roth-type theorems in finite groups. *European J. Combin.*, 34(8):1454–1458, 2013, arXiv:1201.3670. doi:10.1016/j.ejc.2013.05.027.
- [Tao06] T. C. Tao. A variant of the hypergraph removal lemma. *J. Comb. Theory Ser. A*, 113(7):1257–1280, 2006, arXiv:math/0503572. doi:10.1016/j.jcta.2005.11.006.

MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, RADCLIFFE OBSERVATORY QUARTER, WOODSTOCK ROAD, OXFORD OX2 6GG, UNITED KINGDOM

E-mail address: tom.sanders@maths.ox.ac.uk