THE NITROGEN CARRIER IN INNER PROTOPLANETARY DISKS

Klaus M. Pontoppidan

Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

COLETTE SALYK

Vassar College Physics and Astronomy Department, 124 Raymond Avenue, Poughkeepsie, NY 12604, USA

Andrea Banzatti

Lunar and Planetary Laboratory, The University of Arizona, Tucson, AZ, 85721, USA

Geoffrey A. Blake

Division of Geological and Planetary Sciences, California Institute of Technology, MC 150-21, 1200 E California Blvd., Pasadena, CA 91125, USA

CATHERINE WALSH

School of Physics and Astronomy, University of Leeds, Leeds, LS2 9JT, UK

JOHN H. LACY

Department of Astronomy, The University of Texas at Austin, 2515 Speedway, Stop C1400, Austin, TX 78712, USA

Matthew J. Richter

Department of Physics, University of California Davis, 1 Shields Avenue, Davis, CA 95616, USA

Draft version February 12, 2019

ABSTRACT

The dominant reservoirs of elemental nitrogen in protoplanetary disks have not yet been observationally identified. Likely candidates are HCN, NH₃ and N₂. The relative abundances of these carriers determine the composition of planetesimals as a function of disk radius due to strong differences in their volatility. A significant sequestration of nitrogen in carriers less volatile than N₂ is likely required to deliver even small amounts of nitrogen to the Earth and potentially habitable exo-planets. While HCN has been detected in small amounts in inner disks (< 10 au), so far only relatively insensitive upper limits on inner disk NH₃ have been obtained. We present new Gemini-TEXES high resolution spectroscopy of the 10.75 μ m band of warm NH₃, and use 2-dimensional radiative transfer modeling to improve previous upper limits by an order of magnitude to [NH₃/H_{nuc}] < 10⁻⁷ at 1 au. These NH₃ abundances are significantly lower than those typical for ices in circumstellar envelopes ([NH₃/H_{nuc}] ~ 3 × 10⁻⁶). We also consistently retrieve the inner disk HCN gas abundances using archival Spitzer spectra, and derive upper limits on the HCN ice abundance in protostellar envelopes using archival ground-based 4.7 μ m spectroscopy ([HCN_{ice}]/[H₂O_{ice}]< 1.5 – 9%). We identify the NH₃/HCN ratio as an indicator of chemical evolution in the disk, and use this ratio to suggest that inner disk nitrogen is efficiently converted from NH₃ to N₂, significantly increasing the volatility of nitrogen in planet-forming regions.

1. INTRODUCTION

Understanding the pathways of volatiles from the interstellar medium to planets is the subject of intense debate, with the Earth and the Solar system at its center, but with a growing generalization to exoplanetary systems (Moriarty et al. 2014; Ciesla et al. 2015). It remains unclear how the Earth obtained its primary volatiles (carbon, nitrogen and hydrogen/water), and the answer may include multiple sources, including a combination of cometary (Greenwood et al. 2011; Hartogh et al. 2011) and chondritic late impactors (Morbidelli et al. 2000;

Tartèse & Anand 2013). More broadly, both in-situ accretion, and the later delivery, of volatile elements to exoplanets is likely intimately linked to the chemistry of protoplanetary disks. The bulk molecular carriers of a given element during the formation of planetesimals determine its volatility; that is, how much of the element can be found in a solid form as a function of temperature, and therefore how much can be incorporated into planetesimals at a given location in the disk. Once sequestered into planetesimals, the volatiles may be delivered to forming terrestrial planets, either as a local process, or through later delivery during dynamical scattering events (Morbidelli et al. 2000).

The molecular environment in the inner regions of pro-

toplanetary disks (within a few au) is likely fundamentally different from that of the outer, icy regions (10s of AU or more). As there is recent evidence, e.g., from submillimeter dust imaging (Zhang et al. 2018), that some planets exist at large radii of 10-100 AU, we denote the inner $\sim 10\,\mathrm{AU}$ as the "inner planet-forming region", to indicate that this is where the Solar system formed its planets, and where almost all known exoplanets likely formed.

1.1. Nitrogen in the solar system

Delivery of volatiles to terrestrial planets is an inefficient process. Indeed, carbon, nitrogen and oxygen (CNO) are highly abundant in volatiles in molecular clouds, prior to, and during, the star-formation process, with 10-35% of elemental nitrogen accounted for in observations of ices (Öberg et al. 2011a; Pontoppidan et al. 2014). The main carriers of nitrogen in molecular clouds and circumstellar envelopes are N₂, NH₃, HCN and possibly a carbon-dominated refractory dust component. In ices a minor amount of what is likely the OCN⁻ ion is also sometimes seen.

Conversely, nitrogen is highly depleted in the bulk Earth relative to its cosmic abundance, likely by as much as 5–6 orders of magnitude, and by 1–2 orders of magnitude relative to the chondritic N/H₂O abundance (Marty 2012). While it is uncertain how much nitrogen might be sequestered in the deep mantle, this uncertainty is unlikely to increase the terrestrial nitrogen abundance by more than a factor of two (Halliday 2013). In solar system comets, elemental nitrogen is also depleted, albeit by a somewhat smaller value of roughly two orders of magnitude (Mumma & Charnley 2011). Further, the relative depletion of nitrogen in the Earth is much higher than that of carbon and hydrogen (as carried by water). Together, relative abundances of nitrogen in various solar system bodies suggest that the bulk molecular carriers of nitrogen in the Earth-forming disk material were significantly more volatile than water. N₂, for instance, is extremely volatile and generally only available for inclusion into solids below $\sim 15\,\mathrm{K}$ for pure ice, and $\sim 25\,\mathrm{K}$ in the case of N_2 frozen on water (Bisschop et al. 2006; Fayolle et al. 2016), corresponding to distances beyond 20-50 au. Therefore, nitrogen present in the Earth and other terrestrial planets is likely delivered in the form of a less volatile carrier.

1.2. Nitrogen in the inner planet-forming region

The dust temperatures in the inner regions of protoplanetary disks are sufficiently high that no ices will exist at any depth. In this region, the dominant CNO carriers will either be in the gas-phase or in a refractory component, stable to temperatures of at least a few 100 K. Consequently, infrared observations of gas-phase lines from the most abundant nitrogen-bearing molecules will directly constrain major carriers within a few au. The most likely bulk carriers of nitrogen anywhere are NI and $\rm N_2$. However, these species have no electric dipole or quadropole transitions, nor magnetic dipole transitions, and the atomic ground state has no hyperfine splittings, such that they are unobservable in the gas-phase. Their abundance can be crudely estimated as the fraction of the elemental abundance not accounted for by observation of

secondary species, such as HCN and NH_3 (Pontoppidan et al. 2014).

Warm NH₃ in disks has not yet been detected. Mandell et al. (2012) reported upper limits on warm NH₃ in inner disks of $[NH_3/H_2O] < 0.16 - 0.2$, based on highresolution VLT-CRIRES spectra of transitions from the ν_1 band around $3 \, \mu \text{m}$. Assuming an absolute water abundance of $[H_2O/H_{nuc}] = 5 \times 10^{-5}$, this corresponds to a relatively weak upper limit on the absolute NH₃ abundance of $[NH_3/H] < 10^{-5}$ (where $n(H_{\text{nuc}}) = n(H) + 2n(H_2)$). For instance, this limit would not have detected NH₃ in concentrations similar to those in interstellar ices. NH₃ has some of its strongest warm bands in 8–13 μ m regions, of which some were observable with the R = 600 highresolution mode of Spitzer-IRS. Using these bands, Salyk et al. (2011) reported stronger, but uncertain, upper limits of $[NH_3/H_2O] < 0.01$, based on a simple slab model and assuming an NH₃ gas temperature of 400 K. If accurate, these limits suggest that NH₃ has been destroyed in the inner disk, relative to the primordial ice reservoir.

1.3. Nitrogen in outer disks and the cold interstellar medium

In contrast to the inner disk, the outer disk is characterized by the sequestration of bulk volatiles into ices, with only a small fraction reaching the gas-phase due to the action of non-thermal desorption mechanisms. Recently, gas-phase NH₃ was detected in the outer disk of TW Hya (50–100 au, Salinas et al. 2016). The location of the detected NH₃ gas within the TW Hya disk is not well constrained, and it may not be cospatial with the detected cold water vapor (Hogerheijde et al. 2011). The detected gas has a low local abundance, but is thought to be a tracer of photo-evaporation, indicating the presence of a much more massive reservoir of ice. The inferred NH₃ ice abundance relative to water, although with significant uncertainty, could be higher than that of solar system comets by as much as an order of magnitude, thus potentially accounting for most of the elemental nitrogen. While the conversion between gas-phase abundances and the abundances of the underlying ice reservoir is uncertain, the current best evidence nevertheless suggests that NH₃ is an abundant carrier of nitrogen in the cold outer parts of protoplanetary disks.

In interstellar ices, NH_3 is the primary detected nitrogen carrier ($[NH_3/H_2O] \sim 0.1$) (Bottinelli et al. 2010; Öberg et al. 2011a). The abundance of NH_3 in interstellar ices is even somewhat higher than in comets (Öberg et al. 2011a). While HCN is commonly detected in inner disks from mid-infrared spectroscopy (Salyk et al. 2011; Mandell et al. 2012; Najita et al. 2013), and in outer disks through its rotational lines (Öberg et al. 2010), it has yet to be detected in interstellar ices.

1.4. This paper

With this paper, we aim is to derive a more comprehensive picture of the evolution of dominant nitrogen carriers toward the formation of planets. To this end, we employ new and archival infrared spectroscopy of warm gas in inner protoplanetary disks to constrain the major reservoirs of nitrogen near 1 au. Specifically, we report a deep $10.7\,\mu\mathrm{m}$ search, at high spectral resolution, for warm NH₃ at ~ 1 au in protoplanetary disks

around solar-mass young stars. Using two-dimensional radiative transfer models to retrieve the amount of warm NH_3 in the disk surfaces, we derive robust upper limits on the NH_3 abundance, improving on previous limits on warm NH_3 gas in disks by up to an order of magnitude. We use the upper limits on inner disk NH_3 to investigate whether inner disk chemistry increases the average volatility of bulk nitrogen by destroying NH_3 , and potentially re-partitioning it into highly volatile N_2 . We also report new upper limits on the abundance of HCN ice in prestellar dust using archival M-band spectroscopy. Finally, we discuss the implications this may have on our understanding of delivery of nitrogen to the Earth and terrestrial exoplanets.

2. OBSERVATIONS AND DATA

To compare the chemical composition of carriers of bulk nitrogen in inner protoplanetary disks to outer disks, protostars and molecular clouds, we obtained highresolution spectroscopy of a section of the strong $10 \,\mu\mathrm{m}$ $NH_3 \nu_2$ band ("umbrella mode") for a small sample of protoplanetary disks using the TEXES spectrometer (Lacy et al. 2002) mounted on the Gemini North telescope on Mauna Kea, Hawai'i. This mode is one of the strongest NH₃ bands, and is located in a clear part of the Earth's transmission spectrum without strong telluric absorption lines. It has previously been detected in absorption from the ground, using TEXES, toward a number of massive protostars (Knez et al. 2009; Barentine & Lacy 2012), with inferred abundances of $[NH_3/H_{nuc}] \sim 5 \times 10^{-7}$, as well as in Jupiter's atmosphere (Fletcher et al. 2014). The strongest lines in the covered spectral range (10.715–10.77 μ m) trace energies of 1400–1800 K above the ground state. These are similar to the upper level energies of the rotational water lines in the 15-30 μ m range traced by Spitzer, which are known to originate in $\sim 500\,\mathrm{K}$ gas (Carr & Najita 2011; Salyk et al. 2011). That is, the mid-infrared NH₃ and water lines will trace the same gas reservoir, and the observations will provide a robust measurement of the [NH₃/H₂O] ratio, with the option to infer a local [NH₃/H_{nuc}] abundance using a canonical $[H_2O/H_{nuc}]$ abundance of 5×10^{-5} . In Figure 1 the TEXES setting is shown in the context of the wider NH_3 spectrum.

We targeted three bright protoplanetary disks around solar-mass young stars, AS 205N, DR Tau and RNO 90, which were selected based on the presence of strong mid-infrared rotational water line emission and the rovibrational HCN ν_3 band at $14 \,\mu\mathrm{m}$ (Salyk et al. 2011) as observed with the Spitzer Space Telescope InfraRed Spectrograph (IRS, Houck et al. 2004). We also retrieved Spitzer high-resolution spectra of the same sample of disks from Pontoppidan et al. (2010b) to derive consistent HCN abundances using the strong $14 \,\mu \text{m} \, \nu_2$ bending mode. All three disks have velocity-resolved ground-based 12.4 μ m spectroscopy of a few of the water lines (Pontoppidan et al. 2010a; Banzatti et al. 2014), demonstrating that the water emission originates from the inner disks. The presence of water emission allows for the derivation of meaningful relative abundances of NH₃ and HCN. Further, all three disks orbit stars of $0.8 - 1.0 M_{\odot}$, and of ages 1-2 Myr, implying that the three objects trace general properties of this specific demographic.

Finally, we use archival ice spectroscopy of a sample of unrelated protostars to estimate new upper limits on the general HCN abundance in star-forming ices prior to the formation of protoplanetary disks. While the NH₃ abundance in circumstellar ices was measured by Spitzer (Bottinelli et al. 2010), HCN has not yet been detected in the solid phase, and few or no upper limits exist in the literature. Specifically, we estimate upper limits on the HCN ice abundance in a sample of young stars with Spitzerbased NH₃ ice detections using the 4.757 μ m (2102 cm⁻¹) HC-N stretch in spectra obtained with the ISAAC instrument on the ESO Very Large Telescope, as part of the large program 164.I-0605 (van Dishoeck et al. 2003).

2.1. Target properties

RNO 90 is a solar-mass $(1.5 M_{\odot}; Pontoppidan et al.$ 2011) star surrounded by a classical protoplanetary disk. It is also one of the brightest disks known to be rich in molecular emission lines in the infrared, with doublepeaked line profiles and spectro-astrometry demonstrating an origin in a Keplerian disk (Pontoppidan et al. 2008). AS 205N is an optical component in a 1.3" binary. It is somewhat more massive than the Sun, and is still accreting at a relatively high rate of $8 \times 10^{-8} M_{\odot} \,\mathrm{yr}^{-1}$ (Prato et al. 2003; Salyk et al. 2013). It has rovibrational CO line profiles that have been interpreted as evidence of a disk-wind flow (Pontoppidan et al. 2011; Salvk et al. 2014). Thus, AS 205N represents a more active, possibly younger system. DR Tau is a well-known variable, high-accretion T Tauri disk, and the brightest known molecule-rich disk in the Taurus star-forming region (Carr & Najita 2011). Similar to AS 205N, the CO profiles of DR Tau indicate a partial origin in a disk wind.

2.2. TEXES spectroscopy of NH₃

We obtained spectra of the $^{14}{\rm NH_3}$ Q-branch of the ν_2 band near $10.75\,\mu{\rm m}$ (930 cm $^{-1}$) toward the three protoplanetary disks around the solar-mass young stars AS 205N, RNO 90 and DR Tau. We used the cross-dispersed high-medium configuration, which results in instantaneous coverage between 10.715 and 10.775 $\mu{\rm m}$ at a spectral resolving power of $\lambda/\Delta\lambda\sim85,000$. A two-dimensional radiative transfer model of the RNO 90 disk (Blevins et al. 2016) predicts that this spectral region contains some of the strongest NH₃ lines visible from the ground (see Figure 1).

The TEXES data were reduced using standard procedures, including division by a flat field, linearization, registration and co-addition of individual nod-cycles. The spectra were calibrated using equivalent observations of bright asteroids, which are used to remove the combined signature of the system spectral response function and telluric absorption. The wavelength solution was determined using an atmospheric model of the sky emission spectrum. Based on the fit of the observed sky background and the sky emission model, we estimate that the wavelength solution is accurate to $\sim 3\,\mathrm{km\,s^{-1}}$.

The reduced TEXES spectra are presented in Figure 2. They show no detection of NH₃ lines in any of the disks down to the 5-10% line-to-continuum level. Some residual low-frequency noise is apparent, which may be due to residual fringing not removed by the telluric calibrators. This leads to spurious features with widths similar to those of individual spectral orders ($\sim 0.01 \, \mu \text{m}$,




Fig. 1.— Model spectrum showing the NH₃ ν_2 band region at different spectral resolving powers, relevant for TEXES (top curve; $R=85\,000$), JWST-MIRI (middle curve; $R=2\,000$) and Spitzer-IRS (bottom curve; R=600). The shaded region indicates the coverage of the TEXES NH₃ spectral setting. The model shown assumes an inner disk NH₃ abundance of 2.5×10^{-7} per H. Note that in actual disk spectra, other molecular species contribute additional strong lines, some of which will be blended with some of the NH₃ lines.

TABLE 1 SOURCE PROPERTIES AND TEXES OBSERVING LOG

Disk	M_* $[M_{\odot}]$	Luminosity ^a $[L_{\odot}]$	Distance [pc]	Incl. [degrees]	Date	Calibrator	Int. time [Minutes]
AS 205N	$1.0^{\rm b}$	$4.1^{\rm b}$	128^{d}	$20^{\rm e}$	2014 Aug 11	Eunomia	19
DR Tau	0.8^{e}	2.2^{c}	196^{d}	9^{e}	2014 Aug 14	Europa	26
RNO 90	$1.5^{\rm e}$	2.9^{c}	117^{d}	$37^{\rm e}$	$2014~\mathrm{Aug}~13$	Eunomia	45

Note. —

- ^a Total (stellar+accretion) system luminosity.
- ^b Andrews et al. (2010), corrected to the GAIA distance.
- ^c Blevins et al. (2016), corrected to the GAIA distance.
- ^d Gaia Collaboration et al. (2016, 2018)
- ^e Pontoppidan et al. (2011)

or $200-300\,\mathrm{km\,s^{-1}}$). However, based on the observed widths of the (energetically similar) CO fundamental rovibrational lines of $37-92\,\mathrm{km\,s^{-1}}$ (Banzatti & Pontoppidan 2015), the low-frequency noise is not likely to be confused with intrinsic NH₃ line emission.

$2.3. \ \ Archival \ observations \ of \ HCN \ ice \ in \ circumstellar \\ envelopes$

In addition to NH₃, HCN is another potential carrier of significant amounts of nitrogen. While there are many detections of HCN gas in protoplanetary disks (Dutrey et al. 1997; Öberg et al. 2011b), as well as in protostellar envelopes (Jørgensen et al. 2004), there is, to our knowledge, no direct detection of a solid counterpart in interstellar ices, nor have any upper limits been reported. Indirect tracers of solid HCN exist, such as observations of warm gas in hot core regions, thought to trace recently evaporated ices. Based on inference from observations of warm gas-phase HCN, HCN ice is expected to have relatively low abundance compared to major ice species; approximately 2.5×10^{-7} relative to H, or 0.3-0.5% relative to water (Lahuis & van Dishoeck 2000).

In order to better estimate the primordial HCN content of primordial ices, for comparison with the observed abundances of HCN in disks, we report estimated upper limits on HCN ice on grain mantles present during the cold phases of star formation. We use archival M-

band spectra from Keck-NIRSPEC and VLT-ISAAC of the strong, narrow 4.75 $\mu m~(2100\,{\rm cm^{-1}})$ C-N stretch of HCN toward a number of circumstellar envelopes around low-mass young stars. This is the most promising band for HCN detection, as the stronger C-H stretch around 3.2 μm is blended with the 3.1 μm water ice band, and other bands are weaker (Gerakines et al. 2004). The ISAAC spectra were previously published in Pontoppidan et al. (2003), whereas the Keck-NIRSpec spectra of RNO 91 and EC 90 have not previously been published. We include sources from Pontoppidan et al. (2003) for which the water ice band has an optical depth $\tau>1$, and with a signal-to-noise ratio of $\sim\!\!50$ or more.

A challenge to using the CN stretch to detect HCN ice is that it is often affected by absorption and/or emission from warm gas-phase CO, requiring high resolution spectroscopy ($R \gtrsim 10\,000$) to clearly separate any broad ice absorption from multiple CO gas lines. We define the continuum as a linear function, fit to regions between $4.71-4.735\,\mu\mathrm{m}$ and $4.79-4.82\,\mu\mathrm{m}$. The optical depth spectra are then calculated as $\tau = -\ln(F/C)$, where F and C are the flux density and continuum spectra, respectively. The upper limits are derived as 3σ values using the pure HCN ice spectrum and band strength $(5.1\times10^{-18}\,\mathrm{cm\,molec^{-1}})$ from Gerakines et al. (2004). The shape of the HCN band is somewhat sensitive to composition of the ice mantles, with mixtures




Fig. 2.— High resolution TEXES spectra of part of the NH₃ ν_2 mode, compared to the RNO 90 reference disk model (Blevins et al. 2016), assuming different inner disk NH₃ abundances relative to H at ~1 au. Some response function and/or fringe residuals are seen, but these are not coincident with the expected NH₃ lines, and tend to be much broader. The derived upper limits fall between the models with NH₃ abundances of 5×10^{-8} and 5×10^{-7} per H.

with other polar molecules, such as water, leading to signifant broadening and reddening (Noble et al. 2013). The resulting upper limits on the HCN ice column densities are given in Table 2. Compared to the water ice column densities toward the same sources, the upper limits on HCN generally correspond to maximum [HCN/H₂O] abundances of 2-5%, consistent with the relative abundance of gas-phase HCN observed in hot cores. It follows that improvements in sensitivity to solid HCN by at least an order of magnitude are likely necessary to detect it in protostellar sightlines.

3. ANALYSIS

3.1. Radiative transfer modeling and abundance retrieval

In order to retrieve upper limits of the local abundance of NH₃ gas at ~ 1 au, we use the modeling framework described in Zhang et al. (2013) and Blevins et al. (2016). This framework uses the two-dimensional line radiative transfer code RADLite (Pontoppidan et al. 2009) to render predicted line spectra given a disk density structure and molecular abundance structure. The model was fitted in detail to the broad-band SED and 10-180 μm Spitzer-IRS and Herschel-PACS water spectra for RNO 90. It was used to retrieve inner disk H₂O and CO abundances (see Pontoppidan et al. 2014; Blevins et al. 2016, for details on the modeling approach).

The model includes both dust and gas. The dust density structure is fitted to the observed continuum Spectral Energy Distribution (SED). The dust model includes dust grains up to $40~\mu \mathrm{m}$ in size, with a power law size index of -2.5, relevant for the disk surface traced by the infrared molecular lines; larger grains are assumed to have settled to the midplane, where they no longer contribute to the infrared properties of the disk. The molecular

abundances are degenerate with respect to the assumed gas-to-dust ratio, such that lower absolute abundances tend to be needed to produce a given observed line flux for higher gas-to-dust ratios. This is due to a larger gas column being visible for lower opacity disks. Following the procedure in Blevins et al. (2016), we assume a constant gas-to-dust mass ratio of 100 everywhere in the disk. While this is unlikely to be universally true, the infrared molecular spectra trace a relatively small part of the disk (the surface at ~ 1 au), so the assumed gasto-dust ratio is relevant for a relatively restricted region. There is significant uncertainty in the surface gas-to-dust ratio as competing processes are thought to be active. For instance, dust settling and growth acts to increase the gas-to-dust ratio (Meijerink et al. 2009; Horne et al. 2012; Carmona et al. 2014), disk dispersal mechanisms such as photoevaporation tend to decrease it (Bruderer et al. 2014; Ansdell et al. 2016) and chemical effects may mask the true ratio (Kama et al. 2016; Miotello et al. 2017).

The gas temperature is estimated using the thermochemical calculation of Najita et al. (2011), and scaled to the specific RADLite dust model using the vertical column density at each disk radius (see Figure 4 in Blevins et al. (2016)). The level populations are assumed to be in local thermodynamic equilibrium. While the critical densities of infrared molecular transitions for collisions with $\rm H_2$ tend to be high ($n \sim 10^{10}-10^{12}\,\rm cm^{-3}$), a combination of infrared pumping from the local warm dust and with collisions with atomic H provides an efficient mechanism leading to near-thermalization. This has been observed both for rovibrational CO lines (Blake & Boogert 2004; Thi et al. 2013), as well as for the mid-infrared rotational $\rm H_2O$ lines (with high rotational quantum numbers) (Meijerink et al. 2009), and has been supported by models for

Disk	$N(\mathrm{NH_3})$	$N(\mathrm{H_2O})$	N(HCN)	Instrument	Reference
IRAS 08242-5050	4.77 ± 0.46	77.9 ± 7.7	< 2.7	VLT-ISAAC	(Pontoppidan et al. 2003)
SVS 4-5	2.4:	56.5 ± 11.3	< 4.0	VLT-ISAAC	(Pontoppidan et al. 2003)
R CrA IRS5	0.91 ± 0.23	35.8 ± 2.6	< 2.6	VLT-ISAAC	(Pontoppidan et al. 2003)
L1489 IRS	2.31 ± 0.3	42.6 ± 5.1	< 1.7	Keck-NIRSPEC	(Boogert et al. 2002)
RNO 91	2.03 ± 0.3	42.5 ± 3.6	< 1.5:	Keck-NIRSPEC	This paper
EC 90	0.67 ± 0.2	16.9 ± 1.6	< 0.8	Keck-NIRSPEC	This paper
CrA IRS 7B	3:	110.1 ± 19.7	< 1.9	VLT-ISAAC	(Pontoppidan et al. 2003)
WL 6	1.2093 ± 0.24	41.7 ± 6	< 2.9	VLT-ISAAC	(Pontoppidan et al. 2003)
IRS 44	1.258 ± 0.25	34 ± 4	< 1.2	VLT-ISAAC	(Pontoppidan et al. 2003)
IRS 46	0.65 ± 0.13	12.8 ± 2	< 3.1	VLT-ISAAC	(Pontoppidan et al. 2003)
IRS 51	0.53 ± 0.11	22.1 ± 3	< 0.3	VLT-ISAAC	(Pontoppidan et al. 2003)
IRS 63	1.16 ± 0.23	20.4 ± 3	< 0.3	VLT-ISAAC	(Pontoppidan et al. 2003)

TABLE 2
NITROGEN-BEARING ICE COLUMN DENSITIES IN PROTOSTELLAR ENVELOPES

Note. — All column densities are provided in units of $10^{17} \, \mathrm{cm}^{-2}$. Colons indicate high uncertainty.

HCN (Bruderer et al. 2015). For disks around early-type stars, UV fluorescence can in some cases also be important (Brittain et al. 2007), but this is expected to be a small effect for disks around low-mass and solar-mass stars where the local UV fields are less intense. While the expectation is that non-LTE effects are relatively small, in particular for lower-mass stars with low UV fields, non-LTE models for $\rm NH_3$ and HCN are ultimately needed to confirm the results presented in this work.

The warm water and CO ro-vibrational line-tocontinuum ratios in all three disks fall within a factor of 2-3 of each other, and the previous applications of this retrieval approach indicate water and CO abundances that are consistent with canonical abundances of $[H_2O/H_{nuc}] \sim [CO/H_{nuc}] \sim 5 \times 10^{-5}$ (Pontoppidan & Blevins 2014). The three sources also have similar spectral energy distributions, and their luminosities all fall within a factor 2 (see Table 1). Consequently, we consider the three disks to differ only to first order, which means we can use the model spectra for RNO 90, scaling by constant factors to the observed 10.7 μ m continua of AS 205N and DR Tau. The implicit assumption is that, all other model parameters (abundance structure) being equal, infrared line fluxes scale linearly with source luminosity within the narrow range of luminosity of the sample (as does the continuum). Even if there are inherent uncertainties in the absolute abundance relative to H, the abundances relative to water and CO will be self-consistent.

The molecular abundances are implemented as "jump models", in which the abundance is high inside a critical radius, $R_{\rm crit}$, and low outside. Further, the gas abundance is assumed to be low where the dust temperature is below the relevant freeze-out temperature. If $R_{\rm crit}$ is very large, the disk abundance structure follows the location of the snow line. Conversely, if $R_{\rm crit}$ is small, the abundance jump may be interpreted as being due to a chemical effect, rather than freeze-out.

3.2. Inner disk NH₃ upper limits

Using the RNO 90 structure model, we calculate a grid of models by varying the NH₃ gas abundance between [NH₃/H_{nuc}] = 5×10^{-8} and 10^{-6} , as models outside of this range are clearly inconsistent with the data (see Figure 2). Since more face-on configurations will increase the line-to-continuum ratios, and therefore the detectability for the same abundance, we calculate a sep-

arate model grid for the known disk inclinations (see Table 1). The non-detections offer no constraints on a critical radius, so we assume the NH₃ abundance has a large $R_{\rm crit} = 11 \, {\rm au}$, which is the measured location of the surface water snowline in RNO 90 by Blevins et al. (2016). The mid-plane water snowline in the same disk is at 1 au, but this is not likely to be traced by our data. For each model in the grid we then identify spectral channels of the model spectrum with significant potential line emission (> 10\% of the peak line flux) and calculate the leastsquares (χ^2) statistic relative to the observed spectrum on those channels. The model with the lowest abundance that can be excluded with 99% confidence is identified and adopted as the appropriate upper limit. A caveat to upper limits is that we have no measurement of line ratios and opacities, and we therefore cannot rule out higher local NH₃ abundances with a small filling factor. That is, it is possible that high abundance NH₃ is concentrated in a small area ($\ll 10 \,\mathrm{AU}$); we can only say that the average abundance in the inner disk surface is low.

3.3. New inner disk HCN abundances

Since a key comparative nitrogen reservoir in the inner disk is warm HCN, we use the same radiative transfer framework to retrieve the HCN abundance by fitting to the Spitzer spectra presented in Pontoppidan et al. (2010b). Salyk et al. (2011) estimated abundances for the same data set using a simple slab model, however a more self-consistent 2-dimensional RADLite model is expected to significantly improve the estimate. The HCN abundance is constrained not only by the fundamental $14\,\mu\mathrm{m}\ v = 01^10 \to 00^00$ intensity, but also by the shape of the band and the presence of the $v = 02^{0}0 \rightarrow 01^{1}0$ band near $14.3 \,\mu\text{m}$. These bands have different opacities and excitation temperatures, and allow for an independent estimate of the critical radius of a jump model. For HCN, we find that a small critical radius of 1 ± 0.2 au is needed to fit the shape of both the 14 and 14.3 μ m bands. This is illustrated in Figure 5, where a large radius leads to HCN with too low opacity, as seen by a narrow $14 \,\mu \mathrm{m}$ band and relatively weak $14.3 \,\mu\mathrm{m}$ band.

Bruderer et al. (2015) presented a non-LTE model for HCN in the AS 205N disk. They found that the primary excitation mechanism is radiative pumping, and that retrieved abundances using the $14 \,\mu\mathrm{m}$ band are essentially unaffected by an assumption of LTE. Their model also




Fig. 3.— Fits of the $4.75\,\mu\mathrm{m}$ ($2100\,\mathrm{cm}^{-1}$) band of solid HCN to archival M-band spectra of nearby embedded young stars. The curves show the 1 and 5σ upper limits on the strengths of the HCN band. The vertical rectangles indicate the regions of the spectrum used to define the continuum. The deep absorption feature at $4.67\,\mu\mathrm{m}$ is due to CO ice. Also visible in most spectra are emission and absorption lines from ro-vibrational CO gas-phase transitions.

provides a benchmark to the RADLite model, as they estimate the HCN for the same Spitzer spectrum of the AS 205N disk. For AS205N, we estimate an absolute abundance of [HCN/H_{nuc}] = 2×10^{-6} for a gas-to-dust ratio of 100. In comparison, Bruderer et al. (2015) also find that a sharp jump is needed around 1 au and estimate an inner abundance of [HCN/H_{nuc}] = 3×10^{-7} for an assumed gas-to-dust ratio of 1000. These values are consistent given the degeneracy of the assumed gas-to-dust ratio (if Bruderer et al. had assumed a smaller gas-to-dust ratio, the retrieved abundance would increase in proportion).

The NH_3 limits and HCN gas abundances are shown in Table 3.

4. DISCUSSION

4.1. Comparison to NH₃ in the outer disk

A detection of high abundances of cold NH₃ gas was reported by Salinas et al. (2016) using Herschel-HIFI observations of the ortho-NH₃ $1_0 - 0_0$ line at 572.5 GHz. The line was reported to have a spectral FWHM of $0.9 \,\mathrm{km}\,\mathrm{s}^{-1}$. which corresponds to disk radii $\gtrsim 30\,\mathrm{AU}$, assuming a stellar mass of $0.7 M_{\odot}$ (Herczeg & Hillenbrand 2014) and an inclination of the outer disk of 7° (Qi et al. 2004). While the retrieved gas-phase NH₃ abundance is low relative to the total nitrogen content, its presence is interpreted as being the photo-desorption product of a much larger reservoir of NH₃ ice. The inferred nebular ice abundance, while uncertain, could be as high as $[NH_3/H] \sim 10^{-5}$. even higher than that inferred by solar system comets (up to $[{\rm NH_3/H}] \sim 10^{-6}$; see references in Figure 6). The inferred total NH₃ abundance in the outer disk of TW Hya is comparable to that of the most enriched interstellar ices observed in young stellar envelopes (Bottinelli et al. 2010).

4.2. Constraints on the nitrogen budget

In combination with existing detections of warm HCN from the same region of the disk, the presented NH₃ observations constrain the budget of elemental nitrogen, not sequestered in N₂, in the inner disk surface. We find that at radii inside of 10 au, the average NH₃ abundance is at least an order of magnitude lower than that of interstellar ice, and at least marginally lower than that of comets. We rule out the possibility that a significant amount of nitrogen is sequestered in NH₃ in the inner disk surface. In comparison, the inner disk HCN abundance is higher than that of comets at 1 au, but likely significantly lower at radii between 1 and 10 au. This indicates that the chemistry of disk volatiles in the terrestrial region has been significantly altered from its primordial state, possibly with a greater fraction of the nitrogen driven into N_2 .

In Figure 6, we compare the observed abundances, relative to water, of HCN and NH₃ in inner regions of disks to those of comets and interstellar ices. The figure demonstrates that the new upper limits on the inner disk NH₃ abundance are significantly lower than those observed in reservoirs tracing colder material. Although the NH₃ abundance in comets is already known to be depleted relative to interstellar ices, the inner disk abundance appears to be depleted by at least another order of magnitude. At the same time, a self-consistent abundance




Fig. 4.— Left: the gas temperature structure. Some temperature contours are indicated in units of K. Right: The NH₃ abundance of a model close to the upper limit. The solid white curves show the $\tau=1$ surface at $10.7\,\mu\mathrm{m}$, whereas the dashed curves indicate the surface corresponding to a vertical column density of $10^{22}\,\mathrm{cm}^{-2}$.

 ${\bf TABLE~3} \\ {\bf Observed~gas\text{-}phase~nitrogen~carriers~in~inner~disks}$

Disk	$[\mathrm{HCN}/\mathrm{H}_{\mathrm{nuc}}]$	$[{\rm NH_3/H_{\rm nuc}}]^{\rm a}$	$[{\rm H_2O/H}]$	% Missing N ^b
AS 205N	$(2 \pm 0.5) \times 10^{-6}$	$< 1.1 \times 10^{-7}$	5×10^{-5}	97.0
DR Tau	$(1 \pm 0.2) \times 10^{-6}$	$< 1.4 \times 10^{-7}$	5×10^{-5}	95.5
RNO 90	$(3 \pm 1.0) \times 10^{-6}$	$< 2.5 \times 10^{-7}$	5×10^{-5}	98.5

Note. — All absolute abundances are referenced to an assumed gas-todust ratio of 100.

retrieval suggests that HCN abundances are somewhat enhanced in the inner disk. The upper limits on HCN in interstellar ices are not constraining, as they are consistent with both comets and the inner disk abundances. The implication is that some of the nitrogen lost from NH₃ could be driven into HCN in the innermost disk, inside of 1 au. Outside of this region, the HCN abundance remains low. The presented data do not reveal where the liberated nitrogen went in the $\gtrsim 1\,\mathrm{AU}$ region, but chemical models suggest N₂ is a likely candidate (see Section 4.3).

Ån important caveat is that the observations only probe the disk surface, down to vertically integrated column densities of a few $\times 10^{22}\,\mathrm{cm^{-2}}$ (see Figure 4). Thus, the mid-plane region, where the bulk of the molecular mass resides is not directly constrained. Nevertheless, given that all disk material is inherited from a protostellar envelope, the fact that the observed upper limits on the NH₃ abundance in the inner disk are ~ 50 times lower than that observed in the outer disk of TW Hya is strong evidence for the efficient destruction of NH₃ in inner disk surfaces.

4.3. Comparison to chemical models

In recent years, a number of studies of the observable chemistry of inner disks have become available to better understand the observed infrared tracers of warm molecular gas in disks (Heinzeller et al. 2011; Schwarz & Bergin 2014; Walsh et al. 2015; Agúndez et al. 2018). Models

including full nitrogen-bearing chemical networks predict that the bulk carriers of nitrogen in protoplanetary disks consist of $\rm N_2$, HCN and NH₃, with only relatively minor contributions from other simple species. Disk models by Schwarz & Bergin (2014) predict large abundances of gasphase NH₃ at a few au in the midplane exceeding 10^{-5} in models that initialized with most nitrogen in NH₃, and an order of magnitude less in models initialized with most nitrogen in N or N₂. Disk models also generally predict large abundances of NH₃ ice throughout the disk midplane at radii where the dust temperature remains below $\sim 60\,\rm K$, and that there is a tendency to form more NH₃ ice with time (Furuya & Aikawa 2014).

If photodissociation is the main destruction mechanism of all three species, which is an appropriate assumption within a column density of $10^{22} \,\mathrm{cm}^{-2}$, the following destructive reactions apply:

$$N_2 + h\nu \to N + N \tag{1}$$


$$HCN + h\nu \rightarrow H + CN$$
 (2)


$$NH_3 + h\nu \rightarrow NH_2 + H \text{ or } NH + H_2$$
 (3)


For a 4000 K blackbody, the unshielded photodissociation rates for the above three reactions are: $3.2 \times 10^{-16}\,\mathrm{s^{-1}}$, $5.7 \times 10^{-12}\,\mathrm{s^{-1}}$, and $3.6 \times 10^{-9}\,\mathrm{s^{-1}}$, respectively (Heays et al. 2017). The balance of N between these three species will be governed by the rates of destruction versus reformation from constituent

 $^{^{\}rm a}$ Upper limits are 3σ

 $^{^{\}rm b}$ Assuming a solar nitrogen abundance of [N/H_{nuc}] = 6.76 \times 10⁻⁵ (Asplund et al. 2009).










FIG. 5.— Fits of the inner disk HCN abundance to archival Spitzer spectra using the RADLite grid. The water abundance is held constant at $[{\rm H_2O/H_{nuc}}]{=}5\times10^{-5},$ while the HCN abundance and radial size of the region where the HCN abundance is high are varied. The RNO90 fit is consistent with that presented in Pontoppidan et al. (2014). Top: Best fits assuming a large critical radius. Bottom three: Best fits assuming $R_{\rm crit}\leq 1~{\rm au}$.

atoms/radicals. The unshielded destruction rates already indicate a preferential depletion of NH₃ and HCN relative to N_2 in the inner disk surface. Furthermore, we have recently learned that N₂ self-shields efficiently (Li et al. 2013; Heavs et al. 2014). N₂ is therefore even more robust against photodissociation than both HCN and NH₃ in the disk surface. Together, this provides theoretical support for the preferential sequestration of N from NH₃ into N₂. Reformation of HCN will occur through $CN + H_2$ which has a small barrier easily overcome in warm-to-hot gas. Reformation of NH₃ from N, NH, or NH₂, on the other hand, has several barriers, which will slow this reaction sequence down, even in warm-to-hot gas. Any available atomic N is easily converted to N₂ via barrierless reactions, e.g., $N + NO \rightarrow N_2 + O$ or $N + CN \rightarrow N_2 + N$, provided there is adequate supply of small radicals in the gas-phase.

The discrepancy between the low NH₃ abundance found in the inner disk and the potentially large reservoir of NH₃ ice in the outer disk is that, once the temperature is sufficiently cold to retain both N, NH, NH₂, and atomic H on dust grain surfaces, then the formation of NH₃ ice via sequential hydrogenation is rapid. Once

formed, the high binding energy of NH₃ protects it in the ice until the dust grains are warmed to the sublimation temperature, at which point gas-phase chemistry takes over. Indeed, HCN, has an experimental binding energy on astrophysical surfaces in the range 3400-3800 K with some uncertainty (Noble et al. 2013; Rice et al. 2018), whereas NH₃ has a binding energy of 4000 K on compact water ice. When mixed with water NH₃ desorbs at the same temperature as the water, or $\sim 145 \, K$ at laboratory pressures (He et al. 2016). Chemical models therefore predict a reservoir of abundant gas-phase NH₃ within its snowline in the disk midplane due to a lack of efficient destruction mechanisms deeper in the disk. In this regime. NH₃ is mostly chemically inert, although cosmic rays can modify the composition and elemental partitioning over long timescales (> 1 Myr; see Eistrup et al. 2016, 2018). Over time, cosmic ray processing also has the effect of driving NH_3 into N_2 (Eistrup et al. 2018).

Ultimately, the ability of the inner disk to retain abundant NH₃ likely depends on the efficiency of vertical mixing to expose shielded NH₃ from the midplane to the surface chemistry, and mix the processed results back down. Both vertical and radial transport are known to link disparate chemical regions in the disk (Ilgner et al. 2004; Semenov & Wiebe 2011). More specifically, the theoretical expectation is that vertical mixing will transport chemically rich material upwards, and increase the abundances of many species in the inner disk surface, including NH₃ (Heinzeller et al. 2011). The fact that we do not see abundant NH₃ in the surface could mean that either vertical mixing is weak, or that the surface destruction of NH₃ is fast. An argument against weak vertical mixing is that the water abundance is known to be high in all three observed disks, suggesting that we are indeed witnessing the active destruction of NH₃ and transfer of nitrogen into N_2 .

In Figure 6, we compare the observed and predicted NH₃ and HCN abundances in the inner disk surface for a $0.5\,M_\odot$ star from Walsh et al. (2015) as a function of radius. Note that this model is static, and midplane abundances may be affected by the presence of vertical mixing. It is expected that the inner disk surface has reached chemical steady state, whereas the midplane at the same radii evolves slower, resulting in significant chemical decoupling. Further, the inner disk midplane chemistry, because of the longer chemical time scale, is dependent on its initial condition.

It is seen that the observed upper limits on the average NH₃ abundance are consistent with the model just outside of 0.3 au. At radii between 0.3 and several au, the model HCN abundances are not consistent with the observations, while the NH₃ abundances are very low, consistent with the upper limits. The combination of the HCN and NH₃ data therefore supports a scenario in which we are observing relatively abundant HCN in the innermost disk within 1 au, and that the NH₃ abundance is low throughout the inner disk surface, except perhaps for a very small region in the inner few tenths of an au. It also predicts that modest improvements in our sensitivity to warm NH₃ should lead to detections from the same gas giving rise to the observed warm HCN.

4.4. Potential impact on planetary composition




FIG. 6.— Observed evolution of the nitrogen chemistry, as traced by the HCN and NH₃ abundances, from interstellar ices to inner protoplanetary disk surfaces. Inner disks are defined at regions within 10 au from the central star. Also included for reference are the observed abundances in Solar System comets. The comet data points from Smyth et al. (1995); Schloerb et al. (1987); Meier et al. (1994); Biver et al. (1999); Kawakita & Watanabe (1998); Bird et al. (1997); Biver et al. (2012); Kawakita & Mumma (2011); Kobayashi & Kawakita (2009); Dello Russo et al. (2014). The elliptical regions are added as a guide to indicate the approximate regions that may be populated for ices, comets and inner disks. Finally, the data points are compared to the model abundances of the inner disk surface of a T Tauri disk from Walsh et al. (2015). Note that since this figure displays ratios of observed molecular species, it is insensitive to absolute abundances and the assumed gas-to-dust ratio. The range of the outer disk NH₃ abundance in TW Hya from Salinas et al. (2016) is also indicated.

This strong chemical evolution in nitrogen and other common elements has been used to trace the path of biogenic material from the interstellar medium to planetary systems. The degree of volatile depletion in planetesimals and terrestrial (exo)planets can be constrained by models (Lee et al. 2010; Furuya & Aikawa 2014; Schwarz & Bergin 2014), or by directly measuring the abundances of the main carbon, nitrogen and hydrogen carriers in protoplanetary disks (e.g., Hogerheijde et al. 2011; Pontoppidan et al. 2014; McClure et al. 2015; Min et al. 2016). For instance, the ratio of carbon to nitrogen (C/N) in comets, chondrites and the Earth was used by Bergin et al. (2015) to constrain the delivery and loss of volatiles from terrestrial planets, generally finding an increasing C/N ratio as material was transported toward the disk radius forming the Earth. In this picture, the solar system exhibits evidence for increasing nitrogen volatility relative to water and carbon at smaller distances to the Sun. If a significant fraction of the inner disk nitrogen is redistributed from the less volatile NH_3 to HCN and N_2 , as suggested by the observations in this work, this leads to an increase of the C/N ratio in condensible species, at least inside of a few au. This is consistent with the measured C/N ratio in chondrites and the Earth. Other mechanisms than just nitrogen volatility may also act to change the C/N ratio, such as differential loss of volatiles from parent bodies at later stages in the evolution of the planetary system (Kerridge

1999; Marty 2012). Nevertheless, the primordial branching ratio of nitrogen carriers is likely to play a key role in the volatile composition of planet-forming material.

5. CONCLUSIONS

We have presented new, sensitive upper limits on the amount of warm NH₃ gas in the inner regions of protoplanetary disks known to be rich in water. We also presented upper limits on the abundance of HCN in interstellar ices, as well as new estimates of the inner disk surface HCN abundance. We compared the relative NH₃ and HCN abundances in different reservoirs, tracing different evolutionary stages toward the formation of planets. We found that the observed NH₃ abundance relative to H₂O is lower than predicted by static chemical models. The observed highly depleted NH₃ abundances in inner disk surfaces at $\sim 1\,\mathrm{au}$ suggests that a global route for NH₃ destruction in the inner disk surface is efficient. However, we are not constraining the midplane abundances nor to which degree these may be affected by mixing processes. If vertical mixing is efficient, then the inner disk nitrogen reservoir is likely globally depleted in NH₃, leading to a higher mean volatility of the planetforming nitrogen reservoir. This could have important consequences for the ability of the disk to deliver nitrogen to the surfaces of terrestrial planets. Future sensitive observations of warm NH₃ with the James Webb Space Telescope will be critical for constraining the nitrogen chemistry of planet-forming material.

We are grateful to the referee for a constructive report that helped to clarify the paper. K.M.P. and A.B. acknowledges financial support by a NASA Origins of the Solar System grant No. OSS 11-OSS11-0120, a NASA Planetary Geology and Geophysics Program under grant NAG 5-10201. This work is based in part on observations obtained at the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the National Research Council (Canada), CONICYT (Chile), Ministerio de Ciencia, Tecnologa e Innovacin Productiva (Argentina), and Ministrio da Cincia, Tecnologia e Inovao (Brazil). This work is also based in part on observations obtained at the European Southern Observatory,

Paranal, Chile, within the observing program 164.I-0605. This work is also based in part on observations made with the Spitzer Space Telescope, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. C.W. acknowledges support from the University of Leeds and the Science and Technology Facilities Council (grant number ST/R000549/1). This research made use of Astropy, a community-developed core Python package for Astronomy (Astropy Collaboration, 2018). This work has made use of data from the European Space Agency (ESA) mission Gaia (https://www.cosmos.esa.int/gaia), processed by the Gaia Data Processing and Analysis Consortium (DPAC, https://www.cosmos.esa.int/web/ gaia/dpac/consortium). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the Gaia Multilateral Agreement. Facilities: Gemini, VLT, Spitzer.

REFERENCES

Agúndez, M., Roueff, E., Le Petit, F., & Le Bourlot, J. 2018, A&A, 616, A19

Andrews, S. M., Wilner, D. J., Hughes, A. M., Qi, C., & Dullemond, C. P. 2010, ApJ, 723, 1241

Ansdell, M., Williams, J. P., van der Marel, N., et al. 2016, ApJ, 828, 46

Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, ARA&A, 47, 481

Banzatti, A., Meyer, M. R., Manara, C. F., Pontoppidan, K. M., & Testi, L. 2014, ApJ, 780, 26

Banzatti, A., & Pontoppidan, K. M. 2015, ApJ, 809, 167 Barentine, J. C., & Lacy, J. H. 2012, ApJ, 757, 111

Bergin, E. A., Blake, G. A., Ciesla, F., Hirschmann, M. M., & Li, J. 2015, Proceedings of the National Academy of Science, 112, 8065

Bird, M. K., Janardhan, P., Wilson, T. L., et al. 1997, Earth Moon and Planets, 78, 21

Bisschop, S. E., Fraser, H. J., Öberg, K. I., van Dishoeck, E. F., & Schlemmer, S. 2006, A&A, 449, 1297

Biver, N., Bockelée-Morvan, D., Crovisier, J., et al. 1999, AJ, 118, 1850

Biver, N., Crovisier, J., Bockelée-Morvan, D., et al. 2012, A&A, 539, A68

Blake, G. A., & Boogert, A. C. A. 2004, ApJ, 606, L73

Blevins, S. M., Pontoppidan, K. M., Banzatti, A., et al. 2016, ApJ, 818, 22

Boogert, A. C. A., Hogerheijde, M. R., & Blake, G. A. 2002, ApJ, 568, 761

Bottinelli, S., Boogert, A. C. A., Bouwman, J., et al. 2010, ApJ, 718, 1100

Brittain, S. D., Simon, T., Najita, J. R., & Rettig, T. W. 2007, ApJ, 659, 685

Bruderer, S., Harsono, D., & van Dishoeck, E. F. 2015, A&A, 575, A94

Bruderer, S., van der Marel, N., van Dishoeck, E. F., & van Kempen, T. A. 2014, A&A, 562, A26

Carmona, A., Pinte, C., Thi, W. F., et al. 2014, A&A, 567, A51 Carr, J. S., & Najita, J. R. 2011, ApJ, 733, 102

Ciesla, F. J., Mulders, G. D., Pascucci, I., & Apai, D. 2015, ApJ, 804, 9

Dello Russo, N., Vervack, R. J., Kawakita, H., et al. 2014, Icarus, 238, 125

Dutrey, A., Guilloteau, S., & Guelin, M. 1997, A&A, 317, L55
 Eistrup, C., Walsh, C., & van Dishoeck, E. F. 2016, A&A, 595, A83

—. 2018, A&A, 613, A14

Fayolle, E. C., Balfe, J., Loomis, R., et al. 2016, ApJ, 816, L28
 Fletcher, L. N., Greathouse, T. K., Orton, G. S., et al. 2014,
 Icarus, 238, 170

Furuya, K., & Aikawa, Y. 2014, ApJ, 790, 97

Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, A&A, 595, A1

Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, A&A, 616, A1

Gerakines, P. A., Moore, M. H., & Hudson, R. L. 2004, Icarus, 170, 202

Greenwood, J. P., Itoh, S., Sakamoto, N., et al. 2011, Nature Geoscience, 4, 79

Halliday, A. N. 2013, Geochim. Cosmochim. Acta, 105, 146
Hartogh, P., Lis, D. C., Bockelée-Morvan, D., et al. 2011, Nature, 478, 218

He, J., Acharyya, K., & Vidali, G. 2016, ApJ, 823, 56

Heays, A. N., Bosman, A. D., & van Dishoeck, E. F. 2017, A&A, 602, A105

Heays, A. N., Visser, R., Gredel, R., et al. 2014, A&A, 562, A61Heinzeller, D., Nomura, H., Walsh, C., & Millar, T. J. 2011, ApJ, 731, 115

Herczeg, G. J., & Hillenbrand, L. A. 2014, ApJ, 786, 97
 Hogerheijde, M. R., Bergin, E. A., Brinch, C., et al. 2011,
 Science, 334, 338

Horne, D., Gibb, E., Rettig, T. W., et al. 2012, ApJ, 754, 64
Houck, J. R., Roellig, T. L., Van Cleve, J., et al. 2004, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 5487, Optical, Infrared, and Millimeter Space Telescopes, ed. J. C. Mather, 62–76

Ilgner, M., Henning, T., Markwick, A. J., & Millar, T. J. 2004, A&A, 415, 643

Jørgensen, J. K., Schöier, F. L., & van Dishoeck, E. F. 2004, A&A, 416, 603

Kama, M., Bruderer, S., van Dishoeck, E. F., et al. 2016, A&A, 592, A83

Kawakita, H., & Mumma, M. J. 2011, ApJ, 727, 91

Kawakita, H., & Watanabe, J.-i. 1998, ApJ, 495, 946

Kerridge, J. F. 1999, Space Sci. Rev., 90, 275

Knez, C., Lacy, J. H., Evans, II, N. J., van Dishoeck, E. F., & Richter, M. J. 2009, ApJ, 696, 471

Kobayashi, H., & Kawakita, H. 2009, ApJ, 703, 121

Lacy, J. H., Richter, M. J., Greathouse, T. K., Jaffe, D. T., & Zhu, Q. 2002, PASP, 114, 153

Lahuis, F., & van Dishoeck, E. F. 2000, A&A, 355, 699

Lee, J.-E., Bergin, E. A., & Nomura, H. 2010, ApJ, 710, L21

Li, X., Heays, A. N., Visser, R., et al. 2013, A&A, 555, A14 Mandell, A. M., Bast, J., van Dishoeck, E. F., et al. 2012, ApJ, 747, 92

Marty, B. 2012, Earth and Planetary Science Letters, 313, 56 McClure, M. K., Espaillat, C., Calvet, N., et al. 2015, ApJ, 799, 162

Meier, R., Eberhardt, P., Krankowsky, D., & Hodges, R. R. 1994, A&A, 287, 268

Meijerink, R., Pontoppidan, K. M., Blake, G. A., Poelman, D. R., & Dullemond, C. P. 2009, ApJ, 704, 1471

Min, M., Bouwman, J., Dominik, C., et al. 2016, A&A, 593, A11

- Miotello, A., van Dishoeck, E. F., Williams, J. P., et al. 2017, A&A, 599, A113
- Morbidelli, A., Chambers, J., Lunine, J. I., et al. 2000, Meteoritics and Planetary Science, 35, 1309
- Moriarty, J., Madhusudhan, N., & Fischer, D. 2014, ApJ, 787, 81 Mumma, M. J., & Charnley, S. B. 2011, ARA&A, 49, 471
- Najita, J. R., Ádámkovics, M., & Glassgold, A. E. 2011, ApJ, 743, 147
- Najita, J. R., Carr, J. S., Pontoppidan, K. M., et al. 2013, ApJ, 766, 134
- Noble, J. A., Theule, P., Borget, F., et al. 2013, MNRAS, 428, 3262
- Öberg, K. I., Boogert, A. C. A., Pontoppidan, K. M., et al. 2011a, ApJ, 740, 109
- Öberg, K. I., Qi, C., Fogel, J. K. J., et al. 2010, ApJ, 720, 480 —. 2011b, ApJ, 734, 98
- Pontoppidan, K. M., Blake, G. A., & Smette, A. 2011, ApJ, 733, 84
- Pontoppidan, K. M., Blake, G. A., van Dishoeck, E. F., et al. 2008, ApJ, 684, 1323
- Pontoppidan, K. M., & Blevins, S. M. 2014, Faraday Discussions, 169, 49
- Pontoppidan, K. M., Meijerink, R., Dullemond, C. P., & Blake, G. A. 2009, ApJ, 704, 1482
- Pontoppidan, K. M., Salyk, C., Bergin, E. A., et al. 2014, Protostars and Planets VI, 363
- Pontoppidan, K. M., Salyk, C., Blake, G. A., & Käufl, H. U. 2010a, ApJ, 722, L173
- Pontoppidan, K. M., Salyk, C., Blake, G. A., et al. 2010b, ApJ, 720, 887

- Pontoppidan, K. M., Fraser, H. J., Dartois, E., et al. 2003, A&A, 408, 981
- Prato, L., Greene, T. P., & Simon, M. 2003, ApJ, 584, 853 Qi, C., Ho, P. T. P., Wilner, D. J., et al. 2004, ApJ, 616, L11
- Rice, T. S., Bergin, E. A., Jørgensen, J. K., & Wampfler, S. F. 2018, ApJ, 866, 156
- Salinas, V. N., Hogerheijde, M. R., Bergin, E. A., et al. 2016, A&A, 591, A122
- Salyk, C., Herczeg, G. J., Brown, J. M., et al. 2013, ApJ, 769, 21 Salyk, C., Pontoppidan, K., Corder, S., et al. 2014, ApJ, 792, 68
- Salyk, C., Pontoppidan, K. M., Blake, G. A., Najita, J. R., & Carr, J. S. 2011, ApJ, 731, 130
- Schloerb, F. P., Kinzel, W. M., Swade, D. A., & Irvine, W. M. 1987, A&A, 187, 475
- Schwarz, K. R., & Bergin, E. A. 2014, ApJ, 797, 113
- Semenov, D., & Wiebe, D. 2011, ApJS, 196, 25
- Smyth, W. H., Combi, M. R., Roesler, F. L., & Scherb, F. 1995, ApJ, 440, 349
- Tartèse, R., & Anand, M. 2013, Earth and Planetary Science Letters, 361, 480
- Thi, W. F., Kamp, I., Woitke, P., et al. 2013, A&A, 551, A49 van Dishoeck, E. F., Dartois, E., Pontoppidan, K. M., et al. 2003, The Messenger, 113, 49
- Walsh, C., Nomura, H., & van Dishoeck, E. 2015, A&A, 582, A88
 Zhang, K., Pontoppidan, K. M., Salyk, C., & Blake, G. A. 2013, ApJ, 766, 82
- Zhang, S., Zhu, Z., Huang, J., et al. 2018, ApJ, 869, L47