Primitive prime divisors in the critical orbits of one-parameter families of rational polynomials

By RUFEI REN

Department of Mathematics, Fudan University, 220 Handan Rd., Yangpu District, Shanghai 200433, China e-mail: rufeir@fudan.edu.cn

(Received 27 October 2020)

Abstract

For a polynomial $f(x) \in \mathbb{Q}[x]$ and rational numbers c, u, we put $f_c(x) := f(x) + c$, and consider the Zsigmondy set $\mathcal{Z}(f_c, u)$ associated to the sequence $\{f_c^n(u) - u\}_{n \geq 1}$, see Definition [1](#page-0-0).1, where f_c^n is the *n*-st iteration of f_c . In this paper, we prove that if u is a rational critical point of f, then there exists an $M_f > 0$ such that $M_f \geq 0$ $\max_{c\in\mathbb{Q}}\{\#\mathcal{Z}(f_c, u)\}.$

Contents

1. Introduction

For every polynomial $f(x) \in \mathbb{Q}[x]$ and $\alpha \in \mathbb{Q}$ we put $f_{\alpha}(x) := f(x) + \alpha$. Therefore, f_{α} can be considered as a one-parameter family of polynomials. For every $u \in \mathbb{Q}$ we write

$$
\mathbb{S}_{f,u} \coloneqq \{c \in \mathbb{Q} \mid \{f_c^n(u) - u\}_{n \geq 1} \text{ is infinite}\},\
$$

where f_c^n is the *n*-st iteration of f_c . In particular, if $u = 0$, we put $\mathbb{S}_f := \mathbb{S}_{f,0}$.

We denote by $val_p(-)$ the p-adic valuation of Q normalized by $val_p(p) = 1$. In keeping with the terminology of [[5](#page-14-1)], for every polynomial $f(x) \in \mathbb{Q}[x]$, $u \in \mathbb{Q}$ and $n \geq 1$ we say that p is a **primitive prime divisor** of $f^{n}(u) - u$ if $\text{val}_{p}(f^{n}(u) - u) > 0$ and $\operatorname{val}_p(f^k(u) - u) \leq 0$ for all $1 \leq k < n$.

Definition 1.1. The Zsigmondy set of the sequence $\{f^{(n)}(u) - u\}_{n \geq 1}$ is defined by

 $\mathcal{Z}(f, u) \coloneqq \{n \geq 1 \mid f^{n}(u) - u \text{ has no primitive prime divisor}\}.$

The primary application of bounds on the Zsigmondy set is towards understanding

arboreal Galois representations associated to iteration of rational maps over number fields. It is first studied by Bang [[1](#page-14-2)] and Zsigmondy [[9](#page-14-3)]. Since then, there have been quite a few research papers on characterizing/bounding Zsigmondy sets of various sequences in various settings, e.g., Carmichael [[2](#page-14-4)], Schinzel [[8](#page-14-5)], Rice [[7](#page-14-6)], Ingram–Silverman [[5](#page-14-1)], Doerksen–Haensch [[3](#page-14-7)], Gratton–Nguyen–Tucker[[4](#page-14-8)], Krieger [[6](#page-14-9)], etc.

In this work, we are interested in the size of the Zsigmondy set of a sequence obtained from the critical orbit of polynomials with rational coefficients of degree $d \geq 2$. We denote by Σ the set of finite primes of $\mathbb Z$ and reserve p for a prime number.

We first state our main theorem.

Theorem 1.2. For every polynomial $f \in \mathbb{Q}[x]$ of degree $d \geq 2$ with a critical point $u \in \mathbb{Q}$ there is a constant $\mathbf{M}_f > 0$, depending only on f (independent of $c \in \mathbb{Q}$), such that

$$
\#\mathcal{Z}(f_c, u) \leqslant \mathbf{M}_f
$$

for every $c \in \mathbb{S}_{f,u}$.

It is worth mentioning that Rice [[7](#page-14-6)] was the first to prove the finiteness of $\mathcal{Z}(f, 0)$ for each individual polynomial $f(x) \neq x^d$. In [[3](#page-14-7)], Doerksen–Haensch prove Theorem [1](#page-1-1):2 for the case that $f(x) = x^d$, $u = 0$ and $c \in \mathbb{Z}$, which is generalized by Krieger in [[6](#page-14-9)] to every $c \in \mathbb{Q}$, see [[6](#page-14-9), Theorem [1](#page-1-1).1]. Our contribution is to prove Theorem 1.2 for general polynomials which are not necessary to be monic nor integer. As we consider polynomials f that are more complicated than x^d , we did not aim to get the sharpest uniform bound M_f .

Definition 1.3. A polynomial $g(x) \in \mathbb{Q}[x]$ is called x^2 -divisible if it has degree $d \ge 2$ and is of the form

$$
g(x) = u_d x^d + \dots + u_2 x^2 \in \mathbb{Q}[x].
$$

At the last section we will prove that the following theorem implies Theorem [1](#page-1-1).2. **Theorem 1.4.** Given an x^2 -divisible $g(x) \in \mathbb{Z}[x]$ of degree $d \geq 3$ there is a constant $\mathbf{M}_q > 0$, depending only on g, such that

$$
\#\mathcal{Z}(g_c,0)\leqslant \mathbf{M}_g
$$

for every $c \in \mathbb{S}_q$.

Note that one can give an explicit expression of the lower bound \mathbf{M}_g when combining the decomposition of \mathbb{S}_q in Proposition [2](#page-2-0) \cdot 2 with Propositions [3](#page-11-1) \cdot 3, 3 \cdot 4, 3 \cdot 7 and 3 \cdot 8.

This paper is inspired by Krieger's work in [[6](#page-14-9)]. We generalize her result from the special polynomial $f(x) = x^d$ to arbitrary polynomials in Q[x]. We first address the difficulties on this generalization as follows.

The first difficulty is from dealing with the non-monic case, in which the denominator $f_c^n(c)$ is no longer always equal to d^n 's power of the denominator of c. To conquer it, we introduce a factorization of an integer with respect to the leading term u_d of f, see [\(2.9\)](#page-4-0), which allows us to focus on the major factor of the denominator of $f_c^n(c)$.

The second difficulty is from the critical points of large multiplicities. Due to this reason, some arguments in [[6](#page-14-9)] do not work for our case. For example, Krieger uses Mahler's theorem to control $|f_c^n(c)|$ by $|f_c^{n-1}(c) - f_c^{-1}(0)|$. However, this estimation might not be enough when $f_c^{n-1}(c)$ is very close to a critical point with large multiplicity. It forces us to control $|f_c^n(c)|$ in Proposition 2·[11](#page-7-0) by $|f_c^{n-N}(c) - f_c^{-N}(0)|$ for some relatively large $N > 1$.

Acknowledgment

The author would like to thank Tom Tucker and Shenhui Liu for their valuable discussions.

2. Introduction of Proposition [2](#page-2-0):2 and some estimates

We split this section into two parts. In the first part, we introduce our main technical result Proposition 2.2 whose proof will be given in §[3,](#page-12-0) and prove that it implies Theorem [1](#page-1-0).4. In the second part, we focus on estimating $\ln |A_n|$ which appears in Proposition $2·2$.

Let us first set conventions and introduce some notations.

- (1) We set $\mathbb{N} := \{1, 2, \ldots\}$ to be the set of natural numbers and for every $n \in \mathbb{N}$ we denote by $[n]$ the finite set $\{1, 2, \ldots, n\}.$
- (2) We denote by Σ the set of finite primes of $\mathbb Z$ and reserve p for a prime number. For every $n \in \mathbb{Z}\backslash\{0\}$ the sum $\sum_{p|n}$ and the product $\prod_{p|n}$ are taken over all its distinct prime factors whose number is denoted by $\omega(n)$.

We will always write an x^2 -divisible $g(x) \in \mathbb{Z}[x]$ by $g(x) = u_d x^d + \cdots + u_2 x^2 \in \mathbb{Z}[x]$, and define its length by

$$
L_g := 1 + \sum_{i=2}^{d-1} |u_i|/|u_d|.
$$

For every x^2 -divisible $g(x) \in \mathbb{Z}[x]$, $c \in \mathbb{Q}$ and $n \geq 0$ we write the $(n + 1)$ -st iteration $g_c^{n+1}(0)$ as

$$
g_c^{n+1}(0) = g_c^n(c) := \frac{A_n}{B_n},
$$

where $B_n > 0$, A_n are coprime and both depend on g and c. Clearly, we have $c = g_c(0)$ A_0 .

B₀.
Definition 2.1. For an x^2 -divisible $g(x) \in \mathbb{Z}[x]$ and a set S in \mathbb{S}_g we call that g has rapidly increasing numerators on S if there exists an integer $N > 0$ such that for every $c \in S$ there is a finite set J_c with $\#J_c \le N$ such that for every $n \notin J_c$ we have

$$
\ln|A_n| > \sum_{p|n} \ln|A_{n/p}| \,. \tag{2.1}
$$

We now state our main proposition, which is followed by the proof of Theorem [1](#page-1-0).4. **Proposition 2.2** (Main Proposition). Every x^2 -divisible $g(x) \in \mathbb{Z}[x]$ has rapidly increasing numerators on \mathbb{S}_q .

Proof of Theorem 1.[4](#page-1-0) in assuming proposition [2](#page-2-0).2. By [[6](#page-14-9), Lemma 2.3 and Corollary 2.4], if $n \in \mathcal{Z}(g_c, 0)$, then $A_n | \prod_{p|n} A_{n/p}$ and hence

$$
\ln |A_n| \leqslant \sum_{p|n} \ln |A_{n/p}|.
$$

Together with Proposition [2](#page-2-0) \cdot 2, this finishes the proof.

The naive idea of proving Proposition [2](#page-2-0) \cdot 2 is to give a lower bound for $\ln |A_n|$ and an upper bound for $\prod_{p|n} |A_{n/p}|$ such that the lower bound is always greater than the upper bound when n is large enough. Consider that

$$
\ln |A_n| = \ln B_n + \ln |g_c^n(c)|. \tag{2.2}
$$

口

It is sufficient for us to control $\ln B_n$ and $\ln |g_c^n(c)|$.

2.1. Upper bounds for $\ln B_n$ and $\ln |g_c^n(c)|$

Lemma 2.3. Given any x^2 -divisible $g(x) \in \mathbb{Z}[x]$ and $c \in \mathbb{Q}$, for every $n \geq 0$ we have (1) $\ln B_n \leq d^n \ln B_0;$

(2) $\ln |g_c^n(c)| \leq d^n \ln (2|u_d| \max\{|c|, 4L_g\}).$

Proof. (1) Since $g_c^n(c)$ can be written as $A'_n/B_0^{d^n}$ for some $A'_n \in \mathbb{Z}$, we have $B_n|B_0^{d^n}$. (2) It is enough to prove

$$
|g_c^n(c)| \leq (2|u_d|)^{\frac{d^n-1}{d-1}} \left(\max\{|c|, 4L_g\} \right)^{d^n}
$$
 for all $n \geq 0$.

For $n = 0$, we have $|g_c^0(c)| = |c| \le \max\{|c|, 4L_g\}$.

Assume that the desired inequality holds for some $n \geq 0$ and temporarily denote its right side by T_n . Then we have

$$
|g_c^{n+1}(c)| \le |u_d||g_c^n(c)|^d + \sum_{i=2}^{d-1} |u_i||g_c^n(c)|^i + |c| \le |u_d|(T_n + L_g)T_n^{d-1}
$$

$$
\le |u_d| \cdot 2T_n \cdot T_n^{d-1} = (2|u_d|)^{\frac{d^{n+1}-1}{d-1}} \left(\max\{|c|, 4L_g\}\right)^{d^{n+1}}.
$$

the proof follows by induction.

The proof follows by induction.

2.2. A lower bound for $\ln B_n$

Consider that

$$
\frac{A_{n+1}}{B_{n+1}} = g_c^{n+1}(c) = g_c\left(\frac{A_n}{B_n}\right) = \sum_{i=2}^d u_i \frac{A_n^i}{B_n^i} + \frac{A_0}{B_0}.
$$
\n(2.3)

Lemma 2.4. Given any x^2 -divisible $g(x) \in \mathbb{Z}[x]$ and $c \in \mathbb{Q}$, for every $n \geq 0$ if $p \in I(B_n)$, then we have

- (i) $p \in I(B_{n+1})$ and
- (ii) $\operatorname{val}_p(B_{n+1}) = d\operatorname{val}_p(B_n) \operatorname{val}_p(u_d).$

Proof. Note that for every $n \geq 0$ we have

$$
\text{val}_p\left(\frac{u_d A_n^d}{B_n^d}\right) = \text{val}_p(u_d) + d\text{val}_p(A_n) - d\text{val}_p(B_n),\tag{2.4}
$$

$$
\operatorname{val}_p\left(\frac{u_i A_n^i}{B_n^i}\right) \geqslant (1-d)\operatorname{val}_p(B_n), \quad i = 2, \dots, d-1,
$$
\n
$$
(2.5)
$$

$$
\text{val}_p(\frac{A_0}{B_0}) \geqslant -\text{val}_p(B_0). \tag{2.6}
$$

Therefore, if $p \in I(B_n)$, then we have val_p $(A_n) = 0$ and

$$
\operatorname{val}_p\left(\frac{u_d A_n^d}{B_n^d}\right) < (1-d)\operatorname{val}_p(B_n) \leqslant \operatorname{val}_p\left(\frac{u_i A_n^i}{B_n^i}\right) \tag{2.7}
$$

for every $2 \leq i \leq d - 1$. Note that the term on the right hand side of this inequality does not exist for the case $d = 2$.

We now prove this lemma by induction.

Primitive prime divisors in the critical orbits of rational polynomials 5

For $n = 0$ we have $val_p(B_0) > val_p(u_d)$. Combined with [\(2.3\)](#page-3-2) and [\(2.7\)](#page-3-3), this implies

$$
\operatorname{val}_p\left(\frac{A_1}{B_1}\right) = \operatorname{val}_p\left(\frac{u_d A_0^d}{B_0^d}\right) = \operatorname{val}_p(u_d) - d \operatorname{val}_p(B_0) < -\operatorname{val}_p(u_d)
$$

and hence $p \in I(B_1)$.

Now we assume that this lemma holds for every $0 \le k \le n$.

(1) If $p \notin I(B_0)$, we have

$$
\text{val}_p(B_0) \leqslant \text{val}_p(u_d) < \text{val}_p(B_n).
$$

Combined with (2.3) and (2.7) , this implies

$$
\operatorname{val}_p\left(\frac{A_{n+1}}{B_{n+1}}\right) = \operatorname{val}_p\left(\frac{u_d A_n^d}{B_n^d}\right) = \operatorname{val}_p(u_d) - d \operatorname{val}_p(B_n) < -\operatorname{val}_p(u_d) \tag{2.8}
$$

and hence $p \in I(B_{n+1}).$

(2) If $p \in I(B_0)$, then by induction, we have $p \in I(B_n)$ and

$$
\operatorname{val}_p(B_n) = d^n \operatorname{val}_p(B_0) - \operatorname{val}_p(u_d) \sum_{i=0}^{n-1} d^i \geqslant \operatorname{val}_p(B_0).
$$

Combining (2.3) with (2.7) , we also obtain (2.8) . This completes the induction. \Box

For every $a \in \mathbb{Z}$ we denote

$$
I(a) \coloneqq \{ p \in \Sigma \mid \operatorname{val}_p(a) > \operatorname{val}_p(u_d) \},\
$$

and put

$$
\hat{a} := \prod_{p \in I(a)} p^{\text{val}_p(a)}.
$$
\n(2.9)

When $I(a)$ is empty, we put $\hat{a} := 1$. Note that we always have

$$
|a| \le |u_d \hat{a}|. \tag{2.10}
$$

Lemma 2.5. Given any x^2 -divisible $g(x) \in \mathbb{Z}[x]$ of degree $d \geq 3$ and $c \in \mathbb{Q}$, for every $0 \leq n' \leq n$ we have

$$
\ln B_n \geqslant \frac{d^{n-n'}}{3} \ln \widehat{B_{n'}}.
$$

Proof. If $\widehat{B_{n'}} = 1$, it is trivial.

Now we assume that $\widehat{B_{n'}} \ge 2$. It is enough to show that every prime $p \in I(B_{n'})$ satisfies

$$
\operatorname{val}_p(B_n) \geq \frac{d^{n-n'}}{3} \operatorname{val}_p(B_{n'}).
$$
\n(2.11)

Using Lemma [2](#page-3-4).4 inductively, we have

$$
\text{val}_p(B_n) = d^{n-n'} \text{val}_p(B_{n'}) - \text{val}_p(u_d) \sum_{i=0}^{n-n'-1} d^i \geq \left(d^{n-n'} - \sum_{i=0}^{n-n'-1} d^i \right) \text{val}_p(B_{n'}).
$$

From our assumption that $d \geq 3$, we have

$$
d^{n-n'} - \sum_{i=0}^{n-n'-1} d^i \ge d^{n-n'} - 2d^{n-n'-1} \ge \frac{1}{3}d^{n-n'},
$$

which completes the proof.

2.3. The lower bound for $\ln |g_c^n(c)|$

Lemma 2·6. Given any x^2 -divisible $g(x) \in \mathbb{Z}[x]$ and $c \in \mathbb{Q}$, if $|g_c^{n'}(c)| \ge \max\{4L_g, |c|\}$ for some $n' \geqslant 0$, then for every $n \geqslant n'$ we have

$$
|g_c^n(c)| \geq 2^{-\frac{d^{n-n'}-1}{d-1}} \cdot |g_c^{n'}(c)|^{d^{n-n'}}.
$$

Clearly, in this case c is in the basin of infinity for g_c .

Proof. The proof follows from induction. For $n = n'$ this lemma is trivial.

Assume that this lemma holds for some $n \geq n'$. Then we have

$$
|g_c^n(c)|\geqslant 2^{-\frac{d^{n-n'}-1}{d-1}}\cdot |g_c^{n'}(c)|^{d^{n-n'}}\geqslant \max\{4L_g,|c|\}\cdot |4L_g/2|^{\frac{d^{n-n'}-1}{d-1}}\geqslant \max\{4L_g,|c|\},
$$

and hence

$$
|g_c^{n+1}(c)| \ge |u_d||g_c^n(c)|^d - \sum_{i=2}^d |u_i||g_c^n(c)|^i - |c|
$$

\n
$$
\ge |u_d| \left(|g_c^n(c)|^d - L_g|g_c^n(c)|^{d-1}\right) = |g_c^n(c)|^{d-1}|u_d| \left(|g_c^n(c)| - L_g\right)
$$

\n
$$
\ge |g_c^n(c)|^d/2 \ge 2^{-\frac{d^{n-n'+1}-1}{d-1}} \cdot |g_c^{n'}(c)|^{d^{n-n'+1}}.\quad \Box \quad (2.12)
$$

Corollary 2.7. Given any x^2 -divisible $g(x) \in \mathbb{Z}[x]$ and $c \in \mathbb{Q}$, if $|g_c^{n'}(c)| \ge \max\{4L_g, |c|\}$ for some $n' \geqslant 0$, then for every $n \geqslant n'$ we have

$$
\ln |g_c^n(c)| \geq d^{n-n'} \ln |g_c^{n'}(c)/2|.
$$

Proof. It follows directly from Lemma [2](#page-5-1) \cdot 6.

Given an algebraic number $\gamma \in \mathbb{C}$ of degree ℓ with conjugates $\gamma_1 := \gamma, \gamma_2, \ldots, \gamma_\ell$ over $\mathbb{Q},$ let a_0 be an integer such that the coefficients of the polynomial $g(X) = a_0 \prod_{i=1}^{\ell} (X - \gamma_i)$ are integers of gcd 1, then we define the Mahler measure of γ by

$$
M(\gamma) := |a_0| \prod_{i=1}^{\ell} \max(1, |\gamma_i|).
$$

Notation 2.8. For every $r \ge 1$ and $\delta > 0$ we put

$$
W(r,\delta) := 2 \times 10^7 \delta^{-4} \cdot \ln 4r \cdot \ln \ln 4r.
$$

Theorem 2.9 ([[10](#page-14-10)], Theorem 1). Let $0 < \delta < 1$. Then for every algebraic number γ of degree $r \geq 1$, there are at most $W(r, \delta)$ solutions $a/b \in \mathbb{Q}$ to

$$
|a/b - \gamma| < M(a/b)^{-2-\delta} \tag{2.13}
$$

 \Box

with $M(a/b) \ge \max\{4^{2/\delta}, M(\gamma)\}.$

Theorem [2](#page-5-2) \cdot 9 implies the following result.

Corollary 2.10. Given any x^2 -divisible $g(x) \in \mathbb{Z}[x]$, for every $L > 0$, $N \in \mathbb{N}$, $c \in$ $[-L, L] \cap \mathbb{Q}$ and $\gamma \in \mathbb{C}$ such that $g_c^N(\gamma) = 0$, there is an integer $D > 0$, independent of c, such that

$$
|a/b - \gamma| < \frac{1}{(2bD)^3} \tag{2.14}
$$

has at most $W(d^N, 1/10)$ rational solutions a/b with $b \ge \max\left\{4^{20}, (\vert u_d \vert B_0 D)^{d^N}\right\}$.

Primitive prime divisors in the critical orbits of rational polynomials 7

Proof. Let $\gamma_{c,1}, \ldots, \gamma_{c,d^N}$ be the roots of $g_c^N(x) = 0$ in $\mathbb C$ which are not necessary to be distinct. Since $g_c^N(x)$ is continuous as a function of x and c, there exists an integer $D > 1$ such that for every $c \in [-L, L] \cap \mathbb{Q}$ and $1 \leq i \leq d^N$ we have

$$
|\gamma_{c,i}| < D. \tag{2.15}
$$

Without loss of generality, we put $\gamma := \gamma_{c,1}$ and $h(x) := a_0 \prod_{i=1}^{\ell} (x - \gamma_{c,i})$ to be the minimal polynomial of γ with integer coefficients of gcd 1.

Since $B_0^{dN} g_c^N(\gamma) = 0$ and $B_0^{dN} g_c^N(x)$ is a polynomial with integer coefficients, we have $h(x)|B_0^{d^N} g_c^N(x)$. Combined with Gauss's lemma, this implies

$$
a_0|(u_d B_0)^{d^N}.\tag{2.16}
$$

Combining (2.15) with (2.16) , we have

$$
M(\gamma) < (|u_d|B_0)^{d^N} D^\ell. \tag{2.17}
$$

On the other hand, for every rational number a/b in the lowest terms such that $|a/b| \leq$ 2D we have

$$
b \le M(a/b) \le 2bD. \tag{2.18}
$$

Note that Theorem [2](#page-5-2) \cdot 9 still holds when we do the following modifications.

- (1) Restricting this theorem to a set of algebraic numbers and changing $M(\gamma)$ to a function of γ which is larger than $M(\gamma)$ for every γ in this set.
- (2) Changing the right hand side of (2.13) to a function of a/b which is less than $M(a/b)^{-2-\delta}$ for every rational number a/b .
- (3) Changing the second $M(a/b)$ in Theorem [2](#page-5-2).9 to a function of a/b which is less than $M(a/b)$ for every rational number a/b .

Therefore, combined with (2.17) and (2.18) , Theorem [2](#page-5-2) \cdot 9 implies that there are at most $W(\ell, \delta)$ rational solutions a/b to

$$
|a/b - \gamma| < (2bD)^{-2-\delta} \tag{2.19}
$$

such that $|a/b| \leq 2D$ and $b \geq \max\{4^{2/\delta}, (|u_d|B_0)^{d^N}D^{\ell}\}.$

For rational number a/b such that $|a/b| \ge 2D$ we have

$$
|a/b - \gamma| \ge D > 1 > (2bD)^{-2-\delta}.
$$

Together with [\(2.19\)](#page-6-4), this shows that there at most $W(\ell, \delta)$ rational solutions $a/b \in \mathbb{Q}$ to

$$
|a/b - \gamma| < (2bD)^{-2-\delta} \tag{2.20}
$$

with $b \ge \max\{4^{2/\delta}, (|u_d|B_0)^{d^N}D^{\ell}\}.$

Take $\delta := 1/10$. Combining $W(\ell, 1/10) \leq W(d^N, 1/10)$ with the modification(3) above, we can replace ℓ by d^N and $-2 - 1/10$ by -3 , which completes the proof. □

Now we consider $|c| \leq 4L_g$. Recall that \mathbb{S}_g is the set that contains all rational number c such that $\{g_c^n(0)\}\$ is infinite. By Corollary [2](#page-5-4)·7 with $n' = 0$, for every $c \in (-\infty, -4L_g]$ $[4L_g, \infty)$ we have $\lim_{n\to\infty} \ln |g_c^n(c)| = \infty$ and hence $\mathbb{S}_g \supset (-\infty, -4L_g] \cup [4L_g, \infty)$.

We denote by \mathbb{U}_g^0 the finite subset of $\mathbb{S}_g \cap \left[-4L_g, 4L_g\right]$ consisting of all the rational numbers with denominator dividing u_d and put $\mathbb{U}_g := \left[-4L_g, 4L_g \right] \cap (\mathbb{S}_g \backslash \mathbb{U}_g^0)$. The following proposition aims at dealing the case $c \in \mathbb{U}_g$. It is worth noting that $\widehat{B_0} \geq 2$ for all $c \in \mathbb{U}_q$.

Proposition 2.11. For an x^2 -divisible $g(x) \in \mathbb{Z}[x]$ and a real number $\alpha \in [-4L_g, 4L_g]$ such that $g(x) \neq u_d x^d$ or $\alpha \neq 0$, there exists $0 < \delta < L_g$, $C > 0$ and an integer $N \geq 0$ such that for every $c \in (\alpha - \delta, \alpha + \delta) \cap \mathbb{S}_g$ if $\widehat{B_{n'}} \geq 2$ for some $n' \geq 0$, then there is a finite set $S_c \subset \mathbb{N}$ of bounded cardinality $N + n'$ such that for every $n \notin S_c$ we have

$$
\ln |g^n_c(c)| \geqslant \min \left\{(-1 + 1/d) \ln B_n + \ln C, \ \ln \delta \right\}.
$$

Proof. Let

$$
N_0:=\left\lceil\frac{2\ln 3}{\ln(d/(d-1))}\right\rceil+1,
$$

which satisfies

$$
9(d-1)^{N_0-1} \leq d^{N_0-1}.\tag{2.21}
$$

Let $\gamma_1,\ldots,\gamma_r \in \mathbb{C}$ be the distinct roots of $g_{\alpha}^{N_0}(x) = 0$ of multiplicity m_1,\ldots,m_r , respectively. Choose an $0 < \epsilon < 1$ small enough such that for any two distinct $i, j \in [r]$ we have $|\gamma_i - \gamma_j| > 3\epsilon$.

By continuity of $g_c^N(x)$ as a function of x and c, there exists $0 < \delta < L_g$ such that for every $1 \leq i \leq r$ and $\alpha', \beta \in \mathbb{R}$ with $|\alpha' - \alpha| < \delta$ and $|\beta| < \delta$ there are exactly m_i roots of $g_{\alpha'}^{N_0}(x) - \beta = 0$ in the disk $O(\gamma_i, \epsilon) \subset \mathbb{C}$.

Now we consider an arbitrary $c \in (\alpha - \delta, \alpha + \delta) \cap \mathbb{S}_q$.

Let Γ be the multiset consisting of all the roots of $g_c^{N_0}(x) = 0$, i.e. two elements in Γ could be the same. From the argument above, for every $n \ge N_0$ if $|g_c^n(c)| < \delta$, then there exists $1 \leq i_0 \leq r$ such that $g_c^{n-N_0}(c) \in O(\gamma_{i_0}, \epsilon)$. We put

$$
\Gamma_1 := \Gamma \backslash O(\gamma_{i_0}, \epsilon)
$$
 and $\Gamma_2 := \Gamma \cap O(\gamma_{i_0}, \epsilon)$.

Note that we have

$$
\Gamma = \Gamma_1 \cup \Gamma_2, \ \#(\Gamma) = d^{N_0} \text{ and } \#(\Gamma_2) = m_{i_0}.
$$
 (2.22)

Now we count the distance between $g_c^{n-N_0}(c)$ and the points in Γ . For every $\xi \in \Gamma_1$, from our choice of ϵ , we have

$$
\left|\xi - g_c^{n - N_0}(c)\right| > 3\epsilon - 2\epsilon = \epsilon. \tag{2.23}
$$

For every $\xi \in \Gamma_2$, by Corollary 2:[10](#page-5-5) with $L := 4L_g$ and $N := N_0$, there is an integer $D > 0$, independent of c, such that

$$
|a/b - \xi| < \frac{1}{(2bD)^3} \tag{2.24}
$$

has at most $W(d^{N_0}, 1/10)$ rational solutions a/b with $b \ge \max\{4^{20}, (\vert u_d \vert B_0 D)^{d^{N_0}}\}.$ Put

 $N_1 := \left[\log_d 120 \right] + N_0$ and $N_2 := \left[3 \log_d \log_2(u_d^2 D + 1) \right] + 2N_0$.

Then for every $n \geq N_1$, by Lemma [2](#page-4-2).5, we have

$$
\ln B_{n+n'-N_0} \geq \frac{d^{n-N_0}}{3} \ln \widehat{B_{n'}} \geq \frac{d^{n-N_0}}{3} \ln 2 \geq 20 \ln 4.
$$

If $\widehat{B_0} = 1$, by Lemma [2](#page-4-2).5, [\(2.10\)](#page-4-3) and the choice of N_2 , for every $n \ge N_2$ we have

$$
\ln B_{n+n'-N_0} \geqslant \frac{d^{n-N_0}}{3} \ln \widehat{B_{n'}} \geqslant \frac{d^{n-N_0}}{3} \ln 2 \geqslant d^{N_0} \ln (u_d^2 D) \geqslant d^{N_0} \ln (|u_d| B_0 D).
$$

Primitive prime divisors in the critical orbits of rational polynomials 9 If $\widehat{B}_0 \ge 2$ $\widehat{B}_0 \ge 2$, by Lemma 2.5 and [\(2.10\)](#page-4-3) again, for every $n \ge N_2$ we have

$$
\ln B_{n-N_0} \geq \frac{d^{n-N_0}}{3} \ln \widehat{B_0} \geq \left(\frac{d^{n-N_0}}{3} - d^{N_0}\right) \ln 2 + d^{N_0} \ln \widehat{B_0}
$$

$$
\geq d^{N_0} \ln(u_d^2 \widehat{B_0} D) \geq d^{N_0} \ln(|u_d| B_0 D).
$$

Therefore, there are most $W(d^{N_0}, 1/10)$ many integers $n \ge \max\{N_1 + n', N_2 + n'\}$ such that $g_c^{n-N_0}(c)$ is a rational solution to [\(2.24\)](#page-7-1). Combined with

$$
|g_c^n(c)| = |g_c^{N_0}(g_c^{n-N_0}(c))| = \prod_{\xi \in \Gamma} |g_c^{n-N_0}(c) - \xi| = \prod_{\xi \in \Gamma_1} |g_c^{n-N_0}(c) - \xi| \cdot \prod_{\xi \in \Gamma_2} |g_c^{n-N_0}(c) - \xi|,
$$

this implies that for all but at most $\#\Gamma_2 \cdot W(d^{N_0}, 1/10)$ many $n \ge \max\{N_1 + n', N_2 + n'\}$ we have

$$
\ln|g_c^n(c)| \ge \#\Gamma_1 \cdot \ln \epsilon - 3 \cdot \#\Gamma_2 \ln(2DB_{n-N_0}). \tag{2.25}
$$

Combining [\(2.22\)](#page-7-2) and our assumption $0 < \epsilon < 1$, we have

$$
\#\Gamma_1 \cdot \ln \epsilon - 3 \cdot \#\Gamma_2 \ln (2DB_{n-N_0}) \geq d^{N_0} \ln \left(\frac{\epsilon}{8D^3} \right) - 3\#\Gamma_2 \ln B_{n-N_0}.
$$

From our assumption that $g(x) \neq u_d x^d$ or $\alpha \neq 0$, we have

$$
\#\Gamma_2 = m_{i_0} \leq (d-1)^{N_0}.
$$

Therefore, the previous statement implies that for all but at most $(d-1)^{N_0}W(d^{N_0}, 1/10)$ many $n \ge \max\{N_1 + n', N_2 + n'\}$ we have

$$
\ln|g_c^n(c)| \ge d^{N_0} \ln\left(\frac{\epsilon}{8D^3}\right) - 3(d-1)^{N_0} \ln B_{n-N_0}.\tag{2.26}
$$

 \Box

By (2.10) and (2.21) , we have

$$
3(d-1)^{N_0}\ln B_{n-N_0}\leqslant \frac{d^{N_0-1}(d-1)}{3}\ln B_{n-N_0}\leqslant \frac{d^{N_0-1}(d-1)}{3}\ln(|u_d|\widehat{B_{n-N_0}}).
$$

By Lemma $2·5$, we obtain

$$
\frac{d^{N_0-1}(d-1)}{3}\ln(|u_d|\widehat{B_{n-N_0}}) \le \frac{d^{N_0-1}(d-1)}{3}\ln|u_d| + \frac{d-1}{d}\ln B_n
$$

 $\le d^{N_0}\ln|u_d| + \frac{d-1}{d}\ln B_n$

The two inequality above implies that

Right Hand Side of (2.26)
$$
\geq d^{N_0} \ln \left(\frac{\epsilon}{8|u_d|D^3} \right) + (-1 + 1/d) \ln B_n
$$
,

which completes the proof.

3. Proof of Proposition [2](#page-2-0).2

The basic idea of proving Proposition [2](#page-2-0).2 is to show that for each x^2 -divisible $g(x) \in$ $\mathbbm{Z}[x]$ there exists a finite cover of \mathbb{S}_g as follows:

- (1) $(-\infty, -4L_g] \cup [4L_g, \infty);$
- (2) $(\alpha \delta_{g,\alpha}, \alpha + \delta_{g,\alpha}) \cap \mathbb{U}_g$ for finitely many α in $[-4L_g, 4L_g]$ with $0 < \delta_{g,\alpha} < L_g$;
- (3) the finite set \mathbb{U}_g^0 ,

such that g has rapidly increasing numerators on every set in this cover.

Recall that for every $n \in \mathbb{N}$ we denote by $\omega(n)$ the number of its distinct prime divisors. For convenience, we put $s_d(n) := \sum_{p|n} d^{\frac{n}{p}}$. Then we have the following estimation. **Lemma 3.1.** For every $d \ge 2$ and every $n \ge 30$, we have $s_d(n) \le d^{\frac{3n}{5}}$.

Proof. For every integer $n \ge 2$ we have $n \ge 2^{\omega(n)}$ and hence

$$
\omega(n) \leqslant \log_2 n. \tag{3.1}
$$

Since for every prime divisor p of n we have $n/p \le n/2$. Combined with [\(3.1\)](#page-9-2), we have

$$
s_d(n) \leq d^{\frac{n}{2}}\omega(n) \leq d^{\frac{n}{2}}\log_2 n. \tag{3.2}
$$

On the other hand, for every $n \geq 30$ we have

$$
\log_2 n < 5 < 2^3 \leq d^{\frac{n}{10}}.
$$

Together with [\(3.2\)](#page-9-3), this finishes the proof.

Lemma 3.2. Given any x^2 -divisible $g(x) \in \mathbb{Z}[x]$ and $c \in \mathbb{Q}$, for every $n \geq 30$ we have

$$
\sum_{p|n}\ln \left| A_{n/p}\right|\leqslant d^{\frac{3n}{5}}\ln \Big(2|u_d|^2\widehat{B_0}\max\left\{|c|,4L_g\right\}\Big).
$$

Proof. By Lemma [2](#page-3-5).3, for every $n \geq 0$ we have

$$
\sum_{p|n} \ln |A_{n/p}| \leq \sum_{p|n} d^{n/p} \ln (2|u_d| B_0 \max\{|c|, 4L_g\}).
$$

Together with Lemma 3.1 and (2.10) , this finishes the proof.

Proposition 3.3. Every x^2 -divisible $g(x) \in \mathbb{Z}[x]$ of degree $d \geq 3$ has rapidly increasing numerators on $(-\infty, -4L_g] \cup [4L_g, \infty)$.

Proof. Let c be an arbitrary rational number in $(-\infty, -4L_q] \cup [4L_q, \infty)$.

By Lemma [2](#page-5-4).5, Corollary 2.7 with $n' = 0$ and $|c| \ge 4L_g \ge 4$, for every $n \ge 0$ we have

$$
\ln|A_n| \geq d^n \ln|c/2| + \frac{d^n}{3} \ln \widehat{B_0} \geq \frac{d^n}{3} \ln|c\widehat{B_0}|. \tag{3.3}
$$

Combined with Lemma [3](#page-9-5).2 and $|c| \ge 4L_g$, this implies that for every $n \ge 30$ we have

$$
\ln |A_n| - \sum_{p|n} \ln |A_{n/p}| \geqslant -d^{\frac{3n}{5}} \ln(2|u_d|^2) + \left(d^n/3 - d^{\frac{3n}{5}}\right) \ln \left|c\widehat{B_0}\right|.
$$

Therefore, there exists an integer $N > \max\left\{\frac{5}{2}\log_d 3, 30\right\}$, which only depends on g, such that for every $n \ge N$ and every rational number $c \in (-\infty, -4L_g] \cup [4L_g, \infty)$ we have

$$
\ln |A_n| > \sum_{p|n} \ln |A_{n/p}|.
$$

Thus we prove this proposition.

We next prove the following.

Proposition 3.4. Every polynomial $g(x) = u_d x^d \in \mathbb{Z}[x]$ of degree $d \geq 3$ has rapidly increasing numerators on $\left(-\frac{1}{|4u_d|}, \frac{1}{|4u_d|}\right)$ $\big) \cap \mathbb{S}_g.$

Since $0\notin\mathbb S_q,$ it is sufficient to show the following two lemmas.

 \Box

□

 \Box

Primitive prime divisors in the critical orbits of rational polynomials 11 **Lemma 3.5.** Every polynomial $g(x) = u_d x^d \in \mathbb{Z}[x]$ of degree $d \geq 3$ has rapidly increasing numerators on $\left(0, \frac{1}{4|u_d|}\right)$ $\Big) \cap \mathbb{S}_g.$

Proof. For every $c \in (0, \frac{1}{4|u_d|}) \cap \mathbb{S}_g$ and $n \geq 0$ we have

$$
|c| \le |g_c^n(c)| \le (|u_d| + 1)^{\frac{d^n - 1}{d - 1}} |c| \le (|u_d| + 1)^{d^n} |c| \tag{3.4}
$$

and $\widehat{B_0} \geq 2$.

Combining Lemmas [2](#page-4-2).3(1), 2.5 for $n' = 0$ with [\(3.4\)](#page-10-0), for every $n \ge 0$ we have

$$
\ln |A_n| - \sum_{p|n} \ln |A_{n/p}| \ge (1 - \omega(n)) \ln |c| + \frac{d^n}{3} \ln \widehat{B_0} - s_d(n) \ln \big((|u_d| + 1)B_0 \big).
$$

Together with [\(2.10\)](#page-4-3) and lemma [3](#page-9-4).1, this implies that for every $n \geq 30$ we have

$$
\ln |A_n| - \sum_{p|n} \ln |A_{n/p}| \ge -d^{\frac{3n}{5}} \ln ((|u_d| + 1)|u_d|) + \left(\frac{d^n}{3} - d^{\frac{3n}{5}}\right) \ln \widehat{B_0}.
$$

From $\widehat{B_0} \ge 2$, there exists an integer $N \ge 30$ such that for every $n \ge N$ we have

$$
\ln |A_n| - \sum_{p|n} \ln |A_{n/p}| > 0,
$$

which completes the proof.

Lemma 3 \cdot **6.** Every polynomial $g(x) = u_d x^d \in \mathbb{Z}[x]$ of degree $d \geq 3$ has rapidly increasing numerators on $\left(-\frac{1}{4|u_d|},0\right) \cap \mathbb{S}_g$.

Proof. Note that when d is odd, we may replace c with $-c$ and the forward orbit of 0 will be unchanged, modulo sign. By Lemma [3](#page-10-1) $:5$, we immediately prove this case.

Therefore, it is sufficient to study the case that d is even. We first show that for every $c \in \left(-\frac{1}{4|u_d|}, 0\right)$ and every $n \geq 0$ we have

$$
|c|(1-|u_d||c|^{d-1}) \leq |g_c^n(c)| \leq |c|. \tag{3.5}
$$

 \Box

For $n = 0$, we have $|g_c^0(c)| = |c|$. Assume that [\(3.5\)](#page-10-2) holds for some $n \ge 0$. Since $g_c(x)$ is negative and decreasing on $\left(-\frac{1}{4|u_d|}, 0\right)$, we have

$$
|c|(1-|u_d||c|^{d-1}) = |g_c(c)| \leq |g_c(g_c^n(c))| \leq |f(0)| = |c|,
$$

which proves (3.5) by induction.

Combining Lemmas $2·3(1)$, $2·5$ with (3.5) , we have

$$
\ln |A_n| - \sum_{p|n} \ln |A_{n/p}| \ge \ln (|c| \cdot (1 - |u_d||c|^{d-1})) + \frac{d^n}{3} \ln \widehat{B}_0 - s_d(n) \ln B_0 - \omega(n) \ln |c|
$$

$$
\ge \ln(1 - |u_d||c|^{d-1}) + \frac{d^n}{3} \ln \widehat{B}_0 - s_d(n) \ln B_0.
$$

Together with [\(2.10\)](#page-4-3), Lemma [3](#page-9-4).1 and $|c| < \frac{1}{4|u_d|}$, this implies that for every $n \ge 30$ we have

$$
\ln|A_n| - \sum_{p|n} \ln|A_{n/p}| \ge \ln(3/4) + d^{\frac{3n}{5}} \ln|u_d| + \left(\frac{d^n}{3} - d^{\frac{3n}{5}}\right) \ln \widehat{B}_0.
$$
 (3.6)

On the other hand, for every $c \in \left(-\frac{1}{4|u_d|}, 0\right)$, we have $\widehat{B}_0 \geq 2$. Combined with (3.6) , this proves that there exists an integer $N \ge 30$ such that for every $n \ge N$ we have

$$
\ln |A_n| - \sum_{p|n} \ln |A_{n/p}| > 0,
$$

which completes the proof.

Proposition 3.7. Given any x^2 -divisible $g(x) \in \mathbb{Z}[x]$ of degree $d \geq 3$ and $\alpha \in [-4L_g, 4L_g]$ such that $g(x) \neq u_d x^d$ or $\alpha \neq 0$, there is an $0 < \delta < L_g$ such that g has rapidly increasing numerators on $c \in (\alpha - \delta, \alpha + \delta) \cap \mathbb{U}_q$.

Proof. Note that for every $c \in \mathbb{U}_g$ we have $\widehat{B_0} \ge 2$. By Proposition 2·[11](#page-7-0) with $n' = 0$, there is a 3-tuple $0 < \delta < L_g$, $C > 0$ and $N_1 > 0$ such that for every $c \in (\alpha - \delta, \alpha + \delta) \cap \mathbb{U}_g$ there is a finite set $S_c \subset \mathbb{N}$ of bounded cardinality N_1 such that for every $n \notin S_c$ we have

$$
\ln |g_c^n(c)| \geqslant \min \left\{(-1+1/d) \ln B_n + \ln C, \ln \delta \right\},\
$$

and therefore

$$
\ln|A_n| \geqslant \min\left\{\frac{1}{d}\ln B_n + \ln C, \ \ln B_n + \ln \delta\right\}.
$$
 (3.7)

On the other hand, by Lemma [2](#page-4-2) \cdot 5, we have

$$
\ln(B_n) + \ln \delta - d^{\frac{3n}{5}} \ln \left| 2cu_d^2 \widehat{B}_0 \right| \ge \ln \delta + \left(\frac{d^n}{3} - d^{\frac{3n}{5}}\right) \ln \widehat{B}_0 - d^{\frac{3n}{5}} \ln |2cu_d^2|,
$$

$$
\frac{1}{d} \ln(B_n) + \ln C - d^{\frac{3n}{5}} \ln \left| 2cu_d^2 \widehat{B}_0 \right| \ge \ln C + \left(\frac{d^{n-1}}{3} - d^{\frac{3n}{5}}\right) \ln \widehat{B}_0 - d^{\frac{3n}{5}} \ln |2cu_d^2|.
$$

Combined with Lemma [3](#page-9-5) $:2, (3.7)$ $:2, (3.7)$ and $|c| \leq 5L_q$, this implies that there exists an integer $N_2 \geq 30$ such that for every rational number $c \in (\alpha - \delta, \alpha + \delta)$ and $n \in \{N_2, N_2+1, \dots, \} \setminus S_c$ we have

$$
\ln |A_n| > \sum_{p|n} \ln |A_{n/p}|.
$$

Taking $N \coloneqq N_1 + N_2$, we prove this proposition.

Now we turn our attention to the finite set \mathbb{U}_g^0 . **Proposition 3.8.** Every x^2 -divisible $g(x) \in \mathbb{Z}[x]$ of degree $d \geq 3$ has rapidly increasing numerators on the finite set \mathbb{U}_g^0 .

Proof. It is sufficient to show that for each individual rational number in \mathbb{U}_g^0 there are finite many $n \in \mathbb{N}$ satisfying (2.1) .

Let c be an arbitrary rational number in \mathbb{U}_g^0 . We first show that there must exist an integer n' such that either $|g_c^{n'}(c)| > 4L_g$ or $\widehat{B_{n'}} \ge 2$. Suppose that for every $n \ge 0$ we have $\widehat{B_n} = 1$, i.e., $B_n|u_d$. Since $c \in \mathbb{S}_g$ and there are only finitely many integers in $[-4L_g, 4L_g]$ with denominator dividing u_d , we know that there must exist an $n' \geq 0$ such that $|g_c^{n'}(c)| > 4L_g$.

(1) When $|c| \leq 4L_g$, $\widehat{B_0} = 1$ and there exists an integer $n' \geq 1$ such that $|g_c^{n'}(c)| > 4L_g$. Combining these conditions with Corollary [2](#page-5-4).7 and Lemma [3](#page-9-5).2, for every $n \ge \max\{30, n'\}$ we have

$$
\ln |A_n| - \sum_{p|n} \ln |A_{n/p}| \geqslant -d^{\frac{3n}{5}} \ln(8L_g |u_d|^2) + d^{n-n'} \left| g_c^{n'}(c)/2 \right|.
$$

 \Box

 \Box

Primitive prime divisors in the critical orbits of rational polynomials 13 Clearly, there exists an integer $N > \max\{30, n'\}$ such that for every $n \ge N$ we have

$$
\ln |A_n| > \sum_{p|n} \ln |A_{n/p}|.
$$

(2) When $|c| \le 4L_g$, $\widehat{B_0} = 1$ and there is an integer $n' \ge 1$ such that $\widehat{B_{n'}} \ge 2$. Similar to Proposition 3.7 , we combine Lemma 2.5 with Proposition 2.11 , and obtain a finite set $S_c \subset \mathbb{N}$, an integer N, $\delta_c > 0$ and $C_c > 0$ such that for every $n \geq n'$ if $n \notin S_c$, then

$$
\ln |g_c^n(c)| \geqslant \min \left\{ (-1 + 1/d) \ln B_n + \ln C_c, \ln \delta_c \right\},\
$$

and therefore

$$
\ln|A_n| \geqslant \min\left\{\frac{1}{d}\ln B_n + \ln C_c, \ln B_n + \ln \delta_c\right\}.
$$
 (3.8)

On the other hand, by Lemma [2](#page-4-2).5, we have

$$
\ln B_n + \ln \delta_c - d^{\frac{3n}{5}} \ln \left| 2cu_d^2 \widehat{B}_0 \right| \ge \ln \delta_c + \frac{d^{n-n'}}{3} \ln \widehat{B}_{n'} - d^{\frac{3n}{5}} \ln |2cu_d^2|,
$$

$$
\frac{1}{d} \ln B_n + \ln C_c - d^{\frac{3n}{5}} \ln \left| 2cu_d^2 \widehat{B}_0 \right| \ge \ln C_c + \frac{d^{n-n'-1}}{3} \ln \widehat{B}_{n'} - d^{\frac{3n}{5}} \ln |2cu_d^2|.
$$

Combined with Lemma [3](#page-9-5) $:2, (3.8)$ $:2, (3.8)$ and $|c| \leq 4L_g$, this implies that there exists an integer $N_1 > \max\{n', 30\}$ such that for every $n \in \{N_1, N_1 + 1, \dots\} \backslash S_c$ we have

$$
\ln |A_n| > \sum_{p|n} \ln |A_{n/p}|.
$$

Put $N := \#S_c + N_1$. Then we complete the proof.

Proof of Proposition [2](#page-2-0).2. By Proposition [3](#page-11-0).7, for every $g(x) \neq u_d x^d$ or $\alpha \neq 0$ and every real number $\alpha \in [-4L_g, 4L_g]$, there is an $0 < \delta_{g,\alpha} < L_g$ such that g has rapidly increasing numerators on $c \in (\alpha - \delta_{g,\alpha}, \alpha + \delta_{g,\alpha}) \cap \mathbb{U}_g$.

For $g(x) = u_d x^d$ and $\alpha = 0$, if we put $\delta_{g,\alpha} := \frac{1}{4|u_d|}$, then we proved in Proposition [3](#page-9-1).4 that g has rapidly increasing numerators on $c \in (-\delta_{g,\alpha}, \delta_{g,\alpha}) \cap \mathbb{U}_g$.

Now for every x^2 -divisible $g(x) \in \mathbb{Z}[x]$ we obtain a cover of \mathbb{S}_g as

$$
(-\infty, -4L_g] \cup [4L_g, \infty) \cup \mathbb{U}_g^0 \cup \bigcup_{\alpha \in [-4L_g, 4L_g]} \left((\alpha - \delta_{g,\alpha}, \alpha + \delta_{g,\alpha}) \cap \mathbb{U}_g \right).
$$
 (3.9)

Note that

$$
\bigcup_{\alpha \in [-4L_g, 4L_g]} (\alpha - \delta_{g,\alpha}, \alpha + \delta_{g,\alpha})
$$

is an open cover of the closed interval $[-4L_g, 4L_g]$, which has a finite cover. We use the center α to represent the interval $(\alpha - \delta_{g,\alpha}, \alpha + \delta_{g,\alpha})$ in this finite cover, and put T to be the index set of α .

Therefore, we obtain a finite cover of \mathbb{S}_q as follows:

$$
(-\infty, -4L_g] \cup [4L_g, \infty) \cup \mathbb{U}_g^0 \cup \bigcup_{\alpha \in T} \Big((\alpha - \delta_{g,\alpha}, \alpha + \delta_{g,\alpha}) \cap \mathbb{U}_g \Big).
$$

By Propositions [3](#page-11-1) \cdot 3, 3 \cdot 4, 3 \cdot 7 and 3 \cdot 8, we know that g has rapidly increasing numerators on each set in this cover. It implies that g also has rapidly increasing numerators on \mathbb{S}_q , which completes the proof. 囗

$$
\Box
$$

[4](#page-1-0). Theorem $1·4$ implies Theorem $1·2$

At the end of this section, we will prove that Theorem 1.4 implies the following proposition which leads to Theorem 1.2 .

Proposition 4.1. For every x^2 -divisible polynomial $g(x) \in \mathbb{Q}[x]$ there is a constant $\mathbf{M}_g > 0$, depending only on g, such that

$$
\#\mathcal{Z}(g_c,0)\leqslant \mathbf{M}_g
$$

for every $c \in \mathbb{S}_q$.

For every $n \in \mathbb{Z}\backslash\{0\}$, we denote by $\omega(n)$ the number of distinct prime factors of n. **Lemma 4.2.** Let $g(x) \in \mathbb{Q}[x]$ be x^2 -divisible and $a \in \mathbb{Z}\backslash\{0\}$. Then with $h(x) := \frac{1}{a}$ $rac{1}{a}g(ax)$ we have

$$
|\#\mathcal{Z}(g_c,0) - \#\mathcal{Z}(h_{c/a},0)| \leq \omega(a) \quad \text{for every } c \in \mathbb{Q}.
$$

Proof. Consider that

$$
g_c^n(ax) = uh_{c/a}^n(x)
$$
 and $g_c^n(0) = ah_{c/a}^n(0)$.

For every $p \nmid u$ we have p is a primitive prime divisor of $g_c^n(0)$ if and only if it is a primitive prime divisor of $h_{c/a}^n(0)$. Therefore, the difference between $\#\mathcal{Z}(g_c, 0)$ and $\#\mathcal{Z}(h_{c/a}, 0)$ can not exceed the number of prime factors of a, which completes the proof. □

Lemma 4.3. For any x^2 -divisible $g(x) \in \mathbb{Q}[x]$ and $t \in \mathbb{Q} \setminus \{0\}$, Proposition 4.[1](#page-13-1) holds for g if and only if it holds for $\frac{1}{t}g(tx)$.

Proof. Let $t = \frac{a}{b}$ be an arbitrary rational number. Put

$$
h(x) := \frac{1}{t}g(tx)
$$
 and $h_1(x) := \frac{1}{a}g(ax)$.

Note that $h_1(x) = \frac{1}{b}h(bx)$.

By Lemma [4](#page-13-2).2 for every $c \in \mathbb{Q}$ we have

$$
|\# \mathcal{Z}(g_c, 0) - \# \mathcal{Z}((h_1)_{c/a}, 0)| \leq \omega(a),
$$

$$
|\# \mathcal{Z}(h_{c/t}, 0) - \# \mathcal{Z}((h_1)_{c/a}, 0)| \leq \omega(b),
$$

which implies

$$
|\#\mathcal{Z}(g_c,0) - \#\mathcal{Z}(h_{c/t},0)| \leq \omega(a) + \omega(b).
$$

Since $\omega(a)$ and $\omega(b)$ are both independent of c, we complete the proof.

 \Box

Proposition [4](#page-1-0).4. Theorem 1.4 implies Theorem 1.2 .

Proof. Step I: We first prove that Theorem [1](#page-1-0).4 implies Proposition [4](#page-13-1).1.

Given any x²-divisible polynomial $g(x) \in \mathbb{Q}[x]$ of degree $d \geq 2$, if $d = 2$, then we can find $t \in \mathbb{Q}$ such that $\frac{1}{t}g(tx) = x^2$. Combining Lemma [4](#page-13-3).3 with [[6](#page-14-9), Theorem 1.1], we prove Proposition [4](#page-13-1).1 for this case.

If $d \geq 3$, there is $t \in \mathbb{Z}$ such that $\frac{1}{t}g(tx) \in \mathbb{Z}[x]$. Combined with Lemma [4](#page-13-3).3, this proves Proposition [4](#page-13-1).1 for this case.

Step II: Now we prove that Proposition [4](#page-13-1).1 implies Theorem [1](#page-1-1).2.

For every polynomial $g(x) \in \mathbb{Q}[x]$ with a critical point $u \in \mathbb{Q}$, if we put $f(x) :=$ $g(x + u) - u$, then we know that 0 is a critical point of f and for every $c \in \mathbb{Q}$ we have $f_c^n(0) = g_c^n(u) - u$ and hence

$$
\mathcal{Z}(g_c, u) = \mathcal{Z}(f_c, 0).
$$

Primitive prime divisors in the critical orbits of rational polynomials 15

 \Box

This completes the proof.

REFERENCES

- [1] A. S. Bang. Talteoretiske undersøgelser. Tidsskrift Mat. (5) 4 (1886), 70–80; 130–137.
- [2] R. D. CARMICHAEL. On the numerical factors of the arithmetic forms $\alpha^n \pm \beta^n$. Ann. of Math. (1/4) 15 (1913), 49–70.
- [3] K. Doerksen and A. Haensch. Primitive prime divisors in zero orbits of polynomials. Integers (3) 12 (2011), 465–473.
- [4] C. Gratton, K. Nguyen, and T. J. Tucker. ABC implies primitive prime divisors in arithmetic dynamic. Bull. Lond. Math. Soc. (6) 45 (2013), 1194–1208.
- [5] P. Ingram and J. Silverman. Primitive divisors in arithmetic dynamics. Math. Proc. Cambridge Philos. Soc. (2) 146 (2009), 289–302.
- [6] H. KRIEGER. Primitive prime divisors in the critical orbit of $z^d + c$. Int. Math. Res. Not. IMRN 23 (2013), 5498–5525.
- [7] B. Rice. Primitive prime divisors in polynomial arithmetic dynamics. Integers (1) 7 (2007), A26, 1–16.
- [8] A. SCHINZEL. Primitive divisors of the expression $a^n - b^n$ in algebraic number fields. J. Reine Angew. Math. 268/269 (1974), 27–33.
- [9] K. Zsigmondy. Zur Theorie der Potenzreste. Monatsh. Math. Phys. (1) 3 (1892), 265–284.
- [10] J. H. Evertse. The number of algebraic numbers of given degree approximating a given algebraic number. Analytic Number Theory, Y. Motohashi (ed.), London Math. Soc. Lecture Notes Ser. 247, Cambridge Univ. Press (1998), 53–83.