

Coexistence of Non-Periodic Attractors

Liviana Palmisano

May 19, 2020

Abstract

In the space of polynomial maps of \mathbb{R}^2 of degree at least two, there are codimension 3 laminations of maps with at least 3 period doubling Cantor attractors. The leaves of the laminations are real-analytic and they have uniform diameter. The closure of each lamination contains the codimension one tangency locus of a saddle point. Asymptotically, the leaves of each lamination align with the leaves of the eigenvalue foliation. This is an example of general coexistence theorems valid for higher dimensional real-analytic unfoldings of two dimensional homoclinic tangencies.

1 Introduction

In order to understand the long term behavior of a dynamical system, one possible approach is to study the set where a lot of orbits spend most of the time. This set is called the attractor of the system. Moreover, as soon as an attractor is detected, one would like to know for which other parameters a similar attractor occurs in a family of systems of the same type. This says how much and in which form an attractor is stable. Attractive periodic orbits and hyperbolic attractors, for example, persist by changing parameters in an open set. They have the strongest form of stability.

We study here real-analytic two-dimensional unfoldings of maps with a strong homoclinic tangency, see Definition 2.7. What makes such families special is that, like in most systems with frictions, the first return maps to a neighborhood near the homoclinic tangency, are close to Hénon maps, see [6, 17]. The set of parameters where this holds, consists of the so-called Hénon strips. The main theorems rely on a refinement of this fact. In the usual real setting, Hénon-like maps are defined as smooth perturbation of a certain form of the Hénon maps. In our real-analytic context, we refer to *Hénon-like maps* as holomorphic perturbations, defined on a bi-disk in \mathbb{C}^2 , of quadratic maps, i.e.

$$HF \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x^2 + \nu + \epsilon(x, y) \\ x \end{pmatrix}.$$

The precise definition is given in (5.4). In particular, given a real-analytic unfolding, the first return maps are Hénon-like maps and they have the following properties, see Theorem 3.61, Theorem 4.1, and Theorem 4.46.

Theorem A. *Let $F_{t,a}$ be a real-analytic two-dimensional unfolding of the homoclinic tangency at $a = 0$. Then, for any n large enough, there exists a Hénon strip \mathcal{H}_n in parameter space, such that, for all $(t, a) \in \mathcal{H}_n$, the first return map $F_{t,a}^n$, to an appropriate domain, becomes a Hénon-like map $HF_{t,a}$, after a holomorphic coordinate change. Furthermore, the parameter dependence is given by*

$$\begin{aligned}\frac{\partial HF_{t,a}}{\partial a} &= \left(L(t, a) + O\left(\frac{n}{\mu^n(t, a)}\right) \right) \mu^{2n}(t, a), \\ \frac{\partial HF_{t,a}}{\partial t} &= \left(M(t, a) + O\left(\frac{1}{\mu^n(t, a)}\right) \right) n\mu^n(t, a),\end{aligned}$$

where $\mu(t, a)$ is the unstable eigenvalue of the saddle point and the constants $L(t, a)$ and $M(t, a)$ are real-analytic functions of the parameters.

Theorem A shows a universal aspect of the parameter dependence of the return maps, namely a parameter change results in essentially a translation. Theorem A will play a crucial role in finding maps in real-analytic unfoldings with period doubling Cantor attractors¹. In fact, the manifold structure of the period doubling locus is only known in the Hénon-like setting, see [11].

The study of part of the local dynamics in unfoldings is reduced to the study of the dynamics of Hénon-like maps. In the Hénon family and in other two-dimensional Hénon-like families the following attractors has been detected:

- there are maps, for an open set of parameters, having one attractive periodic orbit, a sink,
- there are maps, for a positive Lebesgue set of parameters, having a strange attractor², see [3, 15],
- there are maps having a period doubling Cantor attractor and they form a smooth curve in the parameter space, see [11].

A natural question is to ask if there are maps in the Hénon family having two or more of these attractors simultaneously and in which form they are "observable". A strategy to find them is to start with a map having a periodic attracting orbit and try to find parameters in the open set where the periodic orbit persists which have also another attractor. This approach has been introduced originally by Newhouse in [16]. One difference from this classical construction is that we need to make a selection of parameters. This can be a very sophisticated procedure, which has been carried out in for example [4, 5, 6, 8, 10, 18] using different strategies. In [4, 18], the authors find parameters in the Hénon family corresponding to maps having either coexistence of periodic attractors (sinks), or periodic attractors and one non-periodic attractor. Here, using the method introduced in [4], we solve the more delicate problem to find parameters in finite dimensional unfoldings whose corresponding maps have multiple non-periodic attractors combined with sinks. Non-periodic attractors are much less stable than the periodic ones, they can be easily destroyed by changing parameters. Despite this we can prove that multiple period doubling Cantor attractors coexist along real-analytic leaves of a lamination. We summarize in the following statement Proposition 5.3, Theorem 6.1 and Theorem 7.3.

¹See Definition 5.1

²See Definition 7.13

Theorem B. *Let M , \mathcal{P} and \mathcal{T} be real-analytic manifolds and $F : (\mathcal{P} \times \mathcal{T}) \times M \rightarrow M$ be a real-analytic family with $\dim(\mathcal{P}) = 2$. If there exists $\tau_0 \in \mathcal{T}$ such that $F_0 : (\mathcal{P} \times \{\tau_0\}) \times M \rightarrow M$ is an unfolding of a map f_{τ_0} with a strong homoclinic tangency, then for $k = 1, 2$, there exists a codimension k lamination of maps with at least k period doubling Cantor attractors which persist along the leafs. The homoclinic tangency persists along a global codimension one manifold in $\mathcal{P} \times \mathcal{T}$ and this tangency locus is contained in the closure of the lamination. Moreover, the leafs of the lamination are real-analytic and they have a uniform positive diameter when $\dim(\mathcal{T}) \geq 1$.*

Theorem B states that, when $\dim(\mathcal{T}) = 0$, there is a set of maps with two period doubling Cantor attractors which start to move creating a real-analytic lamination when $\dim(\mathcal{T}) \geq 1$. Our method combined with the method in [4], allows also to find parameters in which finitely many sinks and multiple period doubling Cantor attractors coexist, see Theorem 6.2 and Theorem 7.11.

Furthermore, we find other coexistence phenomena persisting along codimension three laminations. In these cases we are able to give a description of the asymptotic direction of the leafs of the laminations. This occurs in the so called *saddle deforming* unfoldings, see Definition 7.4. In saddle deforming unfoldings the level sets of the eigenvalue pair of a saddle point define the codimension two *eigenvalue foliation* of the tangency locus. A saddle deforming unfolding is such that it contains a three dimensional subfamily transversal to this foliation. In these unfoldings, the leafs of each codimension three coexistence lamination align with the leafs of the eigenvalue foliation associated to a saddle point. The coexistence phenomena and their stability stated in Theorem 7.5 and Theorem 7.12 are summarized in the following.

Theorem C. *Let M , \mathcal{P} and \mathcal{T} be real-analytic manifolds and $F : (\mathcal{P} \times \mathcal{T}) \times M \rightarrow M$ be a saddle deforming unfolding, then the following holds:*

- *there exists a codimension 3 lamination 3PD of maps with at least 3 period doubling Cantor attractors which persist along the leafs,*
- *there exists a codimension 3 lamination NHPD of maps with infinitely many sinks and at least 1 period doubling Cantor attractor which persist along the leafs.*

The leafs of the laminations are real-analytic and they have a uniform positive diameter when their dimension is at least one. Moreover, the laminations align with the eigenvalue foliation. In particular, for each leaf of the eigenvalue foliation, there is a sequence of leafs of 3PD and a sequence of leafs of NHPD which accumulate at this eigenvalue leaf.

Laminations of coexisting attractors have been found already in [4]. Here we are able to find laminations of non-periodic attractors and to describe the asymptotic direction of the leafs. This reveals further universal and global aspects of the bifurcation pattern.

We would like to stress that using Theorem A it is possible to find coexistence of 3 coexisting period doubling Cantor attractors near points with a triple homoclinic tangency. The existence of triple homoclinic points is shown in [12]. Such a construction would give a rather small set of this parameters. An integral part of our method is the simultaneous creation of new attractors together with new tangencies. The process is given by an explicit algorithm. This allows for a precise geometrical description of the laminations. In particular, the asymptotic of the laminations is controlled by the eigenvalue foliation.

Our method allows also to find coexistence of period doubling Cantor attractors and a strange one. In this case, we do not have laminations because the stability of the strange

attractors in families with at least three parameters is not yet understood. Theorem 7.14 states the following.

Theorem D. *Let M , \mathcal{P} and \mathcal{T} be real-analytic manifolds and $F : (\mathcal{P} \times \mathcal{T}) \times M \rightarrow M$ be a saddle deforming unfolding, then the set of maps with at least 2 period doubling Cantor attractors and one strange attractor has Hausdorff dimension at least $\dim(\mathcal{P} \times \mathcal{T}) - 2$.*

The previous theorems and the method of the proofs apply in particular to the Hénon family or any family of polynomial maps of \mathbb{R}^2 of degree at least 2, see Theorem 8.2 and Theorem 8.3.

Acknowledgements The author was supported by the Trygger foundation, Project CTS 17:50, and partially by the NSF grant 1600503.

2 Preliminaries

In this section we collect definitions and facts needed in the sequel. A more elaborate exposition can be found in [4]. The following well-known linearization result is due to Sternberg.

Theorem 2.1. *Given $(\lambda, \mu) \in \mathbb{R}^2$, there exists $\kappa \in \mathbb{N}$ such that the following holds. Let M be a two-dimensional real-analytic manifold and let $f : M \rightarrow M$ be a diffeomorphism with saddle point $p \in M$ having unstable eigenvalue $|\mu| > 1$ and stable eigenvalue λ . If*

$$\lambda \neq \mu^{k_1} \text{ and } \mu \neq \lambda^{k_2} \tag{2.2}$$

for $k = (k_1, k_2) \in \mathbb{N}^2$ and $2 \leq |k| = k_1 + k_2 \leq \kappa$, then f is \mathcal{C}^4 linearizable.

Definition 2.3. *Let M be a two-dimensional real-analytic manifold and let $f : M \rightarrow M$ be a diffeomorphism with a saddle point $p \in M$. We say that p satisfies the \mathcal{C}^4 non-resonance condition if (2.2) holds.*

Theorem 2.4. *Let M be a two-dimensional real-analytic manifold and let $f : M \rightarrow M$ be a diffeomorphism with a saddle point $p \in M$ which satisfies the \mathcal{C}^4 non-resonance condition. Let $0 \in \mathcal{P} \subset \mathbb{R}^n$ and let $F : M \times \mathcal{P} \rightarrow M$ be a real-analytic family with $F_0 = f$. Then, there exists a neighborhood U of p and a neighborhood V of 0 such that, for every $t \in V$, F_t has a saddle point $p_t \in U$ satisfying the \mathcal{C}^4 non-resonance condition. Moreover p_t is \mathcal{C}^4 linearizable in the neighborhood U and the linearization depends \mathcal{C}^4 on the parameters.*

The proofs of Theorem 2.1 and Theorem 2.4 can be found in [9, 13]. In the sequel we introduce the concept of a map with a strong homoclinic tangency which appears already in [4]. This is a map on a two-dimensional manifold with a saddle point, whose eigenvalues satisfy a contraction condition as in (f2). Moreover, a map with a strong homoclinic tangency also has a non-degenerate homoclinic tangency and a transversal homoclinic intersection satisfying (f6), (f7) and (f8), see Figure 1. All conditions defining a map with a strong homoclinic tangency are open in the space of maps with an homoclinic and transversal tangency. Also, except for (f2), all conditions are dense. An example is the Hénon family which contains maps with a strong homoclinic tangency. A map with these properties will then be "unfolded" to create a two-dimensional family.




Figure 1: A map with a strong homoclinic tangency

Definition 2.5. Let M be a two-dimensional real-analytic manifold and let $f : M \rightarrow M$ be a local diffeomorphism satisfying the following conditions:

- (f1) f has a saddle point p with unstable eigenvalue $|\mu| > 1$ and stable eigenvalue λ ,
- (f2) $|\lambda||\mu|^3 < 1$,
- (f3) p satisfies the \mathcal{C}^4 non-resonance condition,
- (f4) f has a non degenerate homoclinic tangency, $q_1 \in W^u(p) \cap W^s(p)$,
- (f5) f has a transversal homoclinic intersection, $q_2 \in W^u(p) \pitchfork W^s(p)$,
- (f6) let $[p, q_2]^u \subset W^u(p)$ be the arc connecting p to q_2 , then there exist arcs $W_{\text{loc},n}^u(q_2) = [q_2, u_n]^u \subset W^u(q_2)$ such that $[p, q_2]^u \cap [q_2, u_n]^u = \{q_2\}$ and

$$\lim_{n \rightarrow \infty} f^n(W_{\text{loc},n}^u(q_2)) = [p, q_2]^u,$$

- (f7) there exist neighborhoods $W_{\text{loc},n}^u(q_1) \subset W^u(q_1)$ such that

$$\lim_{n \rightarrow \infty} f^n(W_{\text{loc},n}^u(q_1)) = [p, q_2]^u,$$

- (f8) there exists $N \in \mathbb{N}$ such that

$$f^{-N}(q_1) \in [p, q_2]^u.$$

A map f with these properties is called a map with a strong homoclinic tangency, see Figure 1.

Remark 2.6. *If the unstable eigenvalue is negative, $\mu < -1$, then (f6), (f7), and (f8) are redundant.*

Following [17], we define an unfolding of a map f with a strong homoclinic tangency. This is a family created by "adding" to f two parameters. Details follow. Let $\mathcal{P} = [-r, r]^2$ with $r > 0$. Given a map f with a strong homoclinic tangency, we consider a real-analytic family $F : \mathcal{P} \times M \rightarrow M$ through f with the following properties:

(F1) $F_{0,0} = f$,

(F2) $F_{t,a}$ has a saddle point $p(t, a)$ with unstable eigenvalue $|\mu(t, a)| > 1$, with stable one $\lambda(t, a)$ and

$$\frac{\partial \mu}{\partial t} \neq 0,$$

(F3) let $\mu_{\max} = \max_{(t,a)} |\mu(t, a)|$, $\lambda_{\max} = \max_{(t,a)} |\lambda(t, a)|$ and assume

$$\lambda_{\max} \mu_{\max}^3 < 1,$$

(F4) there exists a real-analytic function $[-r, r] \ni t \mapsto q_1(t) \in W^u(p(t, 0)) \cap W^s(p(t, 0))$ such that $q_1(t)$ is a non degenerate homoclinic tangency of $F_{t,0}$.

(F5) there exists a real-analytic function $[-r, r]^2 \ni (t, a) \mapsto q_2(t, a) \in W^u(p(t, a)) \cap W^s(p(t, a))$ such that $q_2(t, a)$ is a transversal homoclinic intersection of $F_{t,a}$.

According to Theorem 2.4 we may make a change of coordinates to ensure that the family F is \mathcal{C}^4 and that, for all $(t, a) \in [-r_0, r_0]^2$ with $0 < r_0 < r$, $F_{t,a}$ is linear on the ball $[-2, 2]^2$, namely

$$F_{t,a} = \begin{pmatrix} \lambda(t, a) & 0 \\ 0 & \mu(t, a) \end{pmatrix}.$$

Moreover, the saddle point $p(t, a) = (0, 0)$ and the local stable and unstable manifolds satisfy:

- $W_{\text{loc}}^s(0) = [-2, 2] \times \{0\}$,
- $W_{\text{loc}}^u(0) = \{0\} \times [-2, 2]$,
- $q_1(t) \subset W_{\text{loc}}^s(0)$,
- $q_2(t, a) \in \{0\} \times \left(\frac{1}{\mu}, 1\right) \subset W_{\text{loc}}^u(0)$.

Consider the function $y \mapsto (F_{t,a}^N(0, y))_y$. Let $q_3(t, a) \in \{0\} \times (0, 2) \subset W_{\text{loc}}^u(0)$ be the solution of

$$\frac{\partial (F_{t,a}^N(0, y))_y}{\partial y} = 0$$

with $F_{t,a}^N(q_3(t, 0)) = q_1(t)$. Observe that q_3 depends in a \mathcal{C}^3 manner on the parameters. Without loss of generality we may assume that $q_3(t, a) = (0, 1)$. We define $q_1(t, a) = F_{t,a}^N(q_3(t, a))$ and it points in the positive y direction.

Definition 2.7. A family $F_{t,a}$ is called an unfolding of f if it can be reparametrized such that

$$\frac{\partial(q_1(t,a))_y}{\partial a} \neq 0.$$

Remark 2.8. A generic two-dimensional family through f can locally be reparametrized to become an unfolding.

3 Hénon-like normalization

In the sequel we prove that, given a real-analytic unfolding F , then the first return maps to an appropriately chosen domain can be *straightened* to obtain a family consisting of *Hénon-like maps*, see (5.4). The straightened first return maps are holomorphic maps defined on a uniform domain in \mathbb{C}^2 and they are arbitrarily close to degenerate Hénon maps. The idea of the proof is inspired by the classical construction contained for example in [6, 17]. However, in order to apply the main theorem in [11], we need first return maps with uniform holomorphic extensions. For this reason and for the study of the parameter dependence we include the following self-contained construction.

Consider a real-analytic unfolding $F_{t,a}$ of a strong homoclinic tangency, say with $(t,a) \in (-1,1) \times (-1,1)$. We may assume that this family extends to an holomorphic family of the form

$$\mathbb{D} \times \mathbb{D} \ni (t,a) \mapsto F_{t,a} : U \times U \rightarrow \mathbb{C}^2,$$

where U is a domain in \mathbb{C} . There is a local holomorphic change of coordinates such that the saddle point becomes $(0,0)$ and the local stable manifold contains, in the x -axis, the disc of radius 4 centered around 0, denoted by \mathbb{D}_4 . Similarly, the local unstable manifold contains the disc \mathbb{D}_4 in the y -axis. Moreover, the restriction of each map to the invariant manifolds is linearized, that is

$$F_{t,a}(x,0) = (\lambda x, 0) \text{ and } F_{t,a}(0,y) = (0, \mu y).$$

From now on we will study the map $F_{t,a}^{n+N}$ in this new coordinates. The following lemma is stated and proved in [4], see Lemma 17.

Lemma 3.1. *If $(x,y) \in \mathbb{D}_4 \times \mathbb{D}_4$ and $F^i(x,y) \in \mathbb{D}_4 \times \mathbb{D}_4$, for $i \leq n$, then*

$$DF^n(x,y) = \begin{pmatrix} a_{11}\lambda_1^n \mu^n & a_{12}\lambda_1^n \mu^n \\ a_{21} & a_{22}\mu^n \end{pmatrix},$$

where a_{kl} are uniformly bounded holomorphic functions and $a_{22} \neq 0$ and uniformly away from zero.

From (6.17) in [4], there exists an holomorphic function $\mathbb{D} \ni t \mapsto sa_n(t)$ such that, $|sa_n(t)| = O(1/\mu^n)$ and in the parameter $(t, sa_n(t))$ the periodic point p_t , called the strong sink of period $n+N$, has trace zero. Choose $E > 0$ and define, for n large enough, the n^{th} Hénon strip by

$$\mathcal{H}_n = \left\{ (t,a) \in [-t_0, t_0] \times [-a_0, a_0] \mid |a - sa_n(t)| \leq \frac{E}{|\mu(t, sa_n(t))|^{2n}} \right\}. \quad (3.2)$$

The precise choice of E will be made in (5.6). For $(t,a) \in \mathcal{H}_n$ the coordinates of the strong sink are given by $p_t = (p_x(t), p_y(t))$. Without loss of generality we may assume that $p_x(t) = 2$ for all $t \in \mathbb{D}$.

3.1 The straightened map

Choose $(t, a) \in \mathcal{H}_n$. In this section we apply a coordinate change to the restriction of $F_{t,a}^{n+N}$ to a relevant domain such that, the resulting map resembles a Hénon map in the sense that vertical lines go to horizontal lines, see (3.11). This map is called the *straightened* map. First we construct the coordinate change $\sigma_{t,a}$, called the *straightening* map, defined on the domain $HB(t, a)$, called the *straightening box*. Let $\mathbb{D}_1 \subset \mathbb{C}$ be the disk in the complex plane centered around $p_x(t) = 2$. Consider the vertical foliation in $\mathbb{D}_1 \times \mathbb{C}$. The domain of the straightening map

$$\sigma_{t,a} : HB(t, a) \rightarrow \mathbb{D}_1 \times \mathbb{D}_1$$

is the straightening box

$$HB(t, a) = (F_{t,a}^{n+N})^{-1}(\mathbb{D}_1 \times \mathbb{C}) \cap (\mathbb{D}_1 \times \mathbb{C})$$

and it is defined by

$$\sigma_{t,a}(x, y) = \left((F_{t,a}^{n+N}(x, y))_x, x \right)$$

where $(F_{t,a}^{n+N}(x, y))_x$ denotes the x component of $F_{t,a}^{n+N}(x, y)$. Refer to Figure 2

Remark 3.3. *Observe that the box $HB(t, a)$ has two natural foliations. Namely, one given by $\mathbb{D}_1 \times \mathbb{C}$, called the vertical foliation and the other given by $(F_{t,a}^{n+N})^{-1}(\mathbb{D}_1 \times \mathbb{C})$, called the almost horizontal foliation. The straightening map turns the vertical foliation into the horizontal foliation and the almost horizontal foliation into the vertical foliation.*




Figure 2: Straightening box and straightening map

Lemma 3.4. *For every $x \in \mathbb{D}_1$,*

$$HB_x(t, a) = HB(t, a) \cap (\{x\} \times \mathbb{C})$$

is a simply connected domain of diameter proportional to $1/\mu^n(t, a)$. If $y \in HB(t, a)$ then y is proportional to $1/\mu^n(t, a)$. Moreover the straightening map $\sigma_{t,a}$ is a biholomorphism.

Proof. By adjusting the radius of the disc \mathbb{D}_1 , we may assume that there exists a simply connected domain D in the y -axis, containing $q_3(t, 0)$, such that $(F_{t,0}^N(D))_x$ contains strictly \mathbb{D}_1 . Moreover the projection $\pi_x : F_{t,0}^N(D) \rightarrow \mathbb{C}$ is univalent. Let $x \in \mathbb{D}_1$. Then $F_{t,a}^n(\{x\} \times \mathbb{C})$ contains a simply connected domain exponentially close to D in the y -axis. In particular $HB_x(t, a)$ is a simply connected domain with the projection

$$\pi_x : F_{t,a}^{n+N}(HB_x(t, a)) \rightarrow \mathbb{D}_1$$

is univalent and onto. The diameter estimate for $HB_x(t, a)$ follows from Lemma 3.1. Observe that

$$\sigma_{t,a}^{-1}(x, y) = \left(y, \left((F_{t,a}^{n+N})^{-1}(\{x\} \times \mathbb{C}) \cap (\{y\} \times \mathbb{C}) \right)_y \right).$$

In particular the straightening map $\sigma_{t,a}$ is a biholomorphism. \square

Remark 3.5. *The construction of the straightening box implies that for every $t \in \mathbb{D}$, $p(t) \in HB(t, sa_n(t))$.*

Let π_x and π_y be the orthogonal projections to the axes. The following holds.

Lemma 3.6. *Let $\gamma_1 : \mathbb{D} \rightarrow \mathbb{C}^2$ and $\gamma_2 : \mathbb{D} \rightarrow \mathbb{C}^2$ be holomorphic discs such that*

- $\pi_x(\gamma_1(\mathbb{D})) \subset \mathbb{D}$,
- $\pi_y \circ \gamma_1 : \mathbb{D} \rightarrow \mathbb{D}$ is a 2-to-1 covering map,
- $\pi_x \circ \gamma_2 : \mathbb{D} \rightarrow \mathbb{D}$ is univalent

then for $i = 1, 2$,

$$\#(\gamma_i^{-1}(\gamma_1(\mathbb{D}) \cap \gamma_2(\mathbb{D}))) \in \{1, 2\}.$$

In particular, the intersection number is 1 if and only if the two discs have a tangency.

Proof. Observe that the intersection number is an homotopy invariant. Without loss of generality we may assume that $\pi_x(\gamma_1(\mathbb{D})) = \{0\}$ and $\pi_y(\gamma_2(\mathbb{D})) = \{0\}$. The lemma follows. \square

Lemma 3.7. *For every $(t, a) \in \mathcal{H}_n$ there exists a simply connected domain $D_{t,a}$ in the y -axis such that the map*

$$\pi_y \circ F_{t,a}^N : D_{t,a} \rightarrow \mathbb{D}$$

is a 2-to-1 covering map.

Proof. When $a = 0$, the fact that $F_{t,0}^N(q_3(t, 0)) = q_1(t, 0)$ where $q_1(t, 0)$ is the homoclinic tangency, the existence of a disc $D_{t,0}$ is assured. The parameter continuity implies the existence of the disc $D_{t,a}$ when n is large enough. \square

Lemma 3.8. *For every $(t, a) \in \mathcal{H}_n$ and $x \in \mathbb{D}_1$,*

$$\pi_x \circ \sigma_{t,a} \circ F_{t,a}^{n+N} : D_x = (F_{t,a}^{n+N})^{-1}(HB(t, a)) \cap HB_x(t, a) \rightarrow \mathbb{D}_1$$

is a 2-to-1 covering map. In particular, either D_x is simply connected or $D_x = D_x^- \cup D_x^+$ is the union of two simply connected domains.

Proof. Let $x \in \mathbb{D}_1$ and consider $F_{t,a}^n : \{x\} \times \mathbb{D} \rightarrow \mathbb{C}^2$. Because of Lemma 3.1 and Lemma 3.7, there exists a disc $D \subset \{x\} \times \mathbb{D}$ such that $F_{t,a}^n(D)$ is $(\lambda\mu)^n$ close to $D_{t,a}$ and the projection

$$\pi_y \circ F_{t,a}^{n+N} : D \rightarrow \mathbb{D}$$

is a 2-to-1 covering map. Let $x_1 \in \mathbb{D}_1$. Observe that each point $(x, y) \in D$ with

$$(\pi_x \circ \sigma_{t,a} \circ F_{t,a}^{n+N})(x, y) = x_1$$

corresponds to an intersection point of $F_{t,a}^{n+N}(D)$ with $F_{t,a}^{-(n+N)}(\{x_1\} \times \mathbb{C})$. Hence, from Lemma 3.6, we get

$$\#\{(x, y) \in D \mid (\pi_x \circ \sigma_{t,a} \circ F_{t,a}^{n+N})(x, y) = x_1\} \in \{1, 2\}.$$

Moreover, observe that

$$HB_x(t, a) = \{(x, y) \in D \mid (\pi_x \circ \sigma_{t,a} \circ F_{t,a}^{n+N})(x, y) \in \mathbb{D}_1\}.$$

The lemma follows. □

Lemma 3.9. *For every $(t, a) \in \mathcal{H}_n$ and $x \in \mathbb{D}_1$, the domain D_x satisfies*

$$\text{mod}(D_x, HB_x(t, a)) \geq \frac{n}{4\pi} \log \mu(t, a) + O(1).$$

Proof. Choose $x \in \mathbb{D}_1$ and consider the map

$$\pi_y \circ F_{t,a}^{n+N} : HB_x(t, a) \rightarrow \mathbb{C}.$$

The image of this map contains a disk of definite radius centered around zero. We may assume that it contains the disc of radius 2. Moreover, the map has a unique critical point and the corresponding critical value is at distance $O(1/\mu^n)$ to zero, using that $(t, a) \in \mathcal{H}_n$. Hence, there is a disc $D \subset HB_x(t, a)$ such that

$$\pi_y \circ F_{t,a}^{n+N} : D \rightarrow \mathbb{D} \tag{3.10}$$

is a 2-to-1 covering map. By Lemma 3.4, there exists a positive constant K , such that

$$\pi_y(HB(t, a)) \subset K_n$$

where K_n is the disc of radius K/μ^n . Observe that

$$D_x \subset (\pi_y \circ F_{t,a}^{n+N})^{-1}(K_n) \cap HB_x(t, a).$$

The lemma follows from (3.10). □

We define now the domain of the straightened map $\tilde{F}_{t,a}$ as

$$\text{Dom}(\tilde{F}_{t,a}) = \sigma_{t,a}(\cup_{x \in \mathbb{D}_1} D_x) = \sigma_{t,a}\left(\left(F_{t,a}^{n+N}\right)^{-1}(HB(t, a)) \cap HB(t, a)\right) \subset \mathbb{D}_1 \times \mathbb{D}_1$$

where

$$\tilde{F}_{t,a} = \sigma_{t,a} \circ F_{t,a}^{n+N} \circ \sigma_{t,a}^{-1} : \text{Dom}(\tilde{F}_{t,a}) \rightarrow \mathbb{D}_1 \times \mathbb{D}_1. \tag{3.11}$$

By construction

$$\tilde{F}_{t,a} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} g_{t,a}(x, y) \\ x \end{pmatrix}$$

where, for each $y \in \mathbb{D}_1$, the map $x \rightarrow g_{t,a}(x, y)$ is 2-to-1 onto \mathbb{D}_1 .

3.2 Local existence of the critical point

We introduce now the critical point of the return map $F_{t,a}^{n+N} : HB(t, a) \rightarrow \mathbb{R}^2$. This point has the same meaning as the critical point of the one dimensional part of the Hénon map, i.e. $x \mapsto a + x^2$, see (8.1) In this subsection we prove the local existence of the critical point.

Definition 3.12. *A point $c = (c_x, c_y) \in HB(t, a)$ is a critical point of the map $F_{t,a}^{n+N}$ with critical value $v = (v_x, v_y) = F_{t,a}^{n+N}(c) \in HB(t, a)$ if $v_x = c_x$ and $F_{t,a}^{n+N}(HB_{c_x}(t, a))$ is tangent to the leaf of the almost horizontal foliation passing through v . Namely,*

1. $\pi_x \circ F_{t,a}^{n+N}(c_x, c_y) = c_x$,
2. $\frac{\partial}{\partial y} (\pi_x \circ \sigma_{t,a} \circ F_{t,a}^{n+N})(c_x, c_y) = 0$.

Observe that, for $a = sa_n(t)$, the strong sink $p(t)$ is a critical point. In the next proposition we prove that the strong sink persists as critical point in a neighborhood of the strong sink locus in \mathcal{H}_n .

Proposition 3.13. *For n large enough, there exist a maximal neighborhood $\mathcal{C}_n \subset \mathcal{H}_n$ of the real part of the graph of the function sa_n and a real analytic function $c : \mathcal{C}_n \rightarrow \mathbb{R}^2$ such that, for all $(t, a) \in \mathcal{C}_n$, the point $c(t, a)$ is a critical point of $F_{t,a}^{n+N}$.*

The proof of this proposition needs some preparation. Fix $(t, a) \in \mathcal{H}_n$. Let $h_{t,a}$ be a \mathcal{C}^4 diffeomorphism which locally linearizes the saddle point $(0, 0)$ of $F_{t,a}$ and let $\hat{F}_{t,a} = h_{t,a} \circ F_{t,a} \circ h_{t,a}^{-1}$ be the corresponding smooth unfolding of a strong homoclinic tangency.

Lemma 3.14. *The coordinate change $h_{t,a}$ restricted to $HB(t, a) \cap \mathbb{R}^2$ satisfies*

$$Dh_{t,a}(x, y) = \begin{pmatrix} 1 + \alpha/\mu^n(t, a) & \beta \\ \gamma/\mu^n(t, a) & 1 + \delta_x(x - 2) + \delta_y/\mu^n(t, a) \end{pmatrix} \quad (3.15)$$

where $(x, y) \in HB(t, a) \cap \mathbb{R}^2$ and $\alpha, \beta, \gamma, \delta_x, \delta_y$ are \mathcal{C}^3 functions.

Proof. Because h is \mathcal{C}^4 and preserves the x and y axis we have that $h(x, 0)$ and $h(0, y)$ are linear. Hence, without loss of generality we may assume that $h(x, 0) = x$. By rescaling in the y direction we may assume that that $\partial(h)_y/\partial y(2, 0) = 1$. Observe that, using that h preserves the y -axis

$$(h)_x = x + y(x\varphi_{1,x} + y\varphi_{1,y}),$$

where $\varphi_{1,x}$ and $\varphi_{1,y}$ are \mathcal{C}^1 functions. It follows that

$$\frac{\partial(h)_x}{\partial x} = 1 + y \left(\varphi_{1,x} + x \frac{\partial\varphi_{1,x}}{\partial x} + y \frac{\partial\varphi_{1,y}}{\partial x} \right) = 1 + \frac{\alpha}{\mu^n},$$

where we used that, because $(x, y) \in HB(t, a)$, y is proportional to $1/\mu^n$. Moreover,

$$\frac{\partial(h)_x}{\partial y} = x\varphi_{1,x} + y\varphi_{1,y} + y \left(x \frac{\partial\varphi_{1,x}}{\partial y} + \varphi_{1,y} + y \frac{\partial\varphi_{1,y}}{\partial y} \right) = \beta.$$

Write now the Taylor expansion of $(h)_y$ centered around the point $(2, 0)$. Then

$$(h)_y = y(1 + x\varphi_{2,x} + y\varphi_{2,y}),$$

where $\varphi_{2,x}$ and $\varphi_{2,y}$ are \mathcal{C}^1 functions. It follows that,

$$\frac{\partial (h)_y}{\partial x} = y \left(\varphi_{2,x} + x \frac{\partial \varphi_{2,x}}{\partial x} + y \frac{\partial \varphi_{2,y}}{\partial x} \right) = \frac{\gamma}{\mu^n}.$$

Finally,

$$\frac{\partial (h)_y}{\partial y} = 1 + x\varphi_{2,x} + y\varphi_{2,y} + y \left(x \frac{\partial \varphi_{2,x}}{\partial y} + \varphi_{2,y} + y \frac{\partial \varphi_{2,y}}{\partial y} \right) = 1 + \delta_x x + \frac{\delta_y}{\mu^n},$$

where we used again that y is proportional to $1/\mu^n$. \square

Remark 3.16. *Because $Dh(0) = id$, by shrinking the domain of linearization we may assume that $\alpha, \beta, \gamma, \delta_x, \delta_y$ are sufficiently small.*

To describe the critical point incorporating the linearized map \hat{F} we consider the function $\Phi_{t,a} : HB(t,a) \cap \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$\Phi_{t,a}(x, y) = \left(\left(\pi_x \circ h^{-1} \circ \hat{F}_{t,a}^{n+N} \circ h \right) (x, y) - x, \frac{\partial}{\partial y} \left(\pi_x \circ h^{-1} \circ \hat{F}_{t,a}^{n+N} \circ \hat{F}_{t,a}^{n+N} \circ h \right) (x, y) \right).$$

Observe that c is a critical point of $F_{t,a}$ if and only if $\Phi_{t,a}(c) = 0$. Observe that if $p(t)$ is the strong sink of $F_{t,sa_n(t)}$, then $p(t)$ is a critical point and in particular $\Phi_{t,sa_n(t)}(p(t)) = 0$. By applying the implicit function theorem we will prove that the solution $p(t)$ can be extended over a domain of the form \mathcal{C}_n as stated in Proposition 3.13. In order to apply the implicit function theorem, we need to prove that $D\Phi_{t,a}$ is non singular in any critical point. The following notation and lemmas are required.

Let $(x, y) \in HB(t,a) \cap \mathbb{R}^2$ and

- $\underline{x} = (x, y)$ with $Dh_{t,a}(\underline{x}) = \begin{pmatrix} 1 + \alpha/\mu^n & \beta \\ \gamma/\mu^n & \delta \end{pmatrix}$,
- $\underline{x}_1 = h_{t,a}(\underline{x})$,
- $\underline{x}_2 = \hat{F}_{t,a}^n(\underline{x}_1)$ with $D\hat{F}_{t,a}^N(\underline{x}_2) = \begin{pmatrix} A_2 & B_2 \\ C_2 & D_2 \end{pmatrix}$,
- $\underline{x}_3 = \hat{F}_{t,a}^N(\underline{x}_2)$ with $Dh_{t,a}^{-1}(\underline{x}_3) = \begin{pmatrix} 1 + \alpha_3/\mu^n & \beta_3 \\ \gamma_3/\mu^n & \delta_3 \end{pmatrix}$,
- $\underline{x}_4 = \hat{F}_{t,a}^n(\underline{x}_3)$ with $D\hat{F}_{t,a}^N(\underline{x}_4) = \begin{pmatrix} A_4 & B_4 \\ C_4 & D_4 \end{pmatrix}$,
- $\underline{x}_5 = \hat{F}_{t,a}^N(\underline{x}_4)$ with $Dh_{t,a}^{-1}(\underline{x}_5) = \begin{pmatrix} 1 + \alpha_5/\mu^n & \beta_5 \\ \gamma_5/\mu^n & \delta_5 \end{pmatrix}$.

Remark 3.17. *By Lemma 3.14 and Remark 3.3, δ, δ_3 and δ_5 are non zero and close to one. Observe that, by the definition of unfolding*

$$D\hat{F}_{t,0}^N(q_3(t, 0)) = \begin{pmatrix} A & B \\ C & 0 \end{pmatrix}$$

with B away from zero. Because \underline{x}_2 and \underline{x}_4 are near q_3 , without loss of generality we may assume that B_2 and B_4 are away from zero.

Observe that all coefficients introduced, $\alpha, \beta, \dots, A_2, B_2, \dots, \alpha_3, \beta_3, \dots, A_4, B_4, \dots, \alpha_5, \dots$ are functions of the point $(x, y) \in HB(t, a) \cap \mathbb{R}^2$. Moreover we will use the notation

$$D\hat{F}_{t,a}^N(\underline{x}) = \begin{pmatrix} A(\underline{x}, t, a) & B(\underline{x}, t, a) \\ C(\underline{x}, t, a) & D(\underline{x}, t, a) \end{pmatrix}.$$

The following lemmas will give estimates on these coefficients and their partial derivatives.

Lemma 3.18. *The partial derivatives at $\underline{x} = (x, y) \in HB(t, a)$ satisfy the following.*

- $\partial\alpha/\partial x, \partial\beta/\partial x, \partial\gamma/\partial x, \partial\delta/\partial x = O(1),$
- $\partial A_2/\partial x, \partial B_2/\partial x, \partial C_2/\partial x = O(1),$
- $\partial A_4/\partial x, \partial B_4/\partial x, \partial C_4/\partial x, \partial D_4/\partial x = O(\mu^n(t, a)),$
- $\partial\alpha_5/\partial x, \partial\beta_5/\partial x, \partial\gamma_5/\partial x, \partial\delta_5/\partial x = O(\mu^n(t, a)).$

Proof. The first set of estimates follows by the smoothness of $h_{t,a}$. For the second set of estimates, observe that

$$\frac{\partial \underline{x}_2}{\partial x} = \begin{pmatrix} \lambda^n & 0 \\ 0 & \mu^n \end{pmatrix} Dh_{t,a}(\underline{x}) \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} \lambda^n + \alpha(\lambda/\mu)^n \\ \gamma \end{pmatrix} = O(1).$$

Moreover,

$$\frac{\partial \underline{x}_3}{\partial x} = D\hat{F}_{t,a}^N(\underline{x}_2) \frac{\partial \underline{x}_2}{\partial x} = \begin{pmatrix} B_2\gamma + A_2\lambda^n + A_2\alpha(\lambda/\mu)^n \\ D_2\gamma + C_2\lambda^n + C_2\alpha(\lambda/\mu)^n \end{pmatrix} = O(1). \quad (3.19)$$

The third set of estimates follows from the fact that,

$$\frac{\partial \underline{x}_4}{\partial x} = \begin{pmatrix} \lambda^n & 0 \\ 0 & \mu^n \end{pmatrix} \frac{\partial \underline{x}_3}{\partial x} = \begin{pmatrix} B_2\gamma\lambda^n + A_2\lambda^{2n} + A_2\alpha(\lambda^2/\mu)^n \\ D_2\gamma\mu^n + C_2(\lambda\mu)^n + C_2\alpha\lambda^n \end{pmatrix} = O(\mu^n).$$

Finally,

$$\begin{aligned} \frac{\partial \underline{x}_5}{\partial x} &= D\hat{F}_{t,a}^N(\underline{x}_4) \frac{\partial \underline{x}_4}{\partial x} \\ &= \begin{pmatrix} A_4B_2\gamma\lambda^n + A_4A_2\lambda^{2n} + A_4A_2\alpha(\lambda^2/\mu)^n \\ C_4B_2\gamma\lambda^n + C_4A_2\lambda^{2n} + C_4A_2\alpha(\lambda^2/\mu)^n \end{pmatrix} \\ &\quad + \begin{pmatrix} B_4D_2\gamma\mu^n + B_4C_2(\lambda\mu)^n + B_4C_2\alpha\lambda^n \\ D_4D_2\gamma\mu^n + D_4C_2(\lambda\mu)^n + D_4C_2\alpha\lambda^n \end{pmatrix} = O(\mu^n). \end{aligned} \quad (3.20)$$

The last set follows. □

Lemma 3.21. *The partial derivatives at $\underline{x} = (x, y) \in HB(t, a)$ satisfy the following.*

- $\partial\alpha/\partial y, \partial\beta/\partial y, \partial\gamma/\partial y, \partial\delta/\partial y = O(1),$
- $\partial A_2/\partial y, \partial B_2/\partial y, \partial C_2/\partial y = O(\mu(t, a)^n),$
- $\partial A_4/\partial y, \partial B_4/\partial y, \partial C_4/\partial y, \partial D_4/\partial y = O(\mu(t, a)^{2n}),$
- $\partial\alpha_5/\partial y, \partial\beta_5/\partial y, \partial\gamma_5/\partial y, \partial\delta_5/\partial y = O(\mu(t, a)^{2n}).$

Proof. The first set of estimates follows by the smoothness of $h_{t,a}$. For the second set of estimates, observe that,

$$\frac{\partial \underline{x}_2}{\partial y} = \begin{pmatrix} \lambda^n & 0 \\ 0 & \mu^n \end{pmatrix} Dh_{t,a}(\underline{x}) \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} \beta \lambda^n \\ \delta \mu^n \end{pmatrix} = O(\mu^n). \quad (3.22)$$

Moreover,

$$\frac{\partial \underline{x}_3}{\partial y} = D\hat{F}_{t,a}^N(\underline{x}_2) \frac{\partial \underline{x}_2}{\partial y} = \begin{pmatrix} B_2 \delta \mu^n + A_2 \beta \lambda^n \\ D_2 \delta \mu^n + C_2 \beta \lambda^n \end{pmatrix} = O(\mu^n). \quad (3.23)$$

The third set of estimates follows from the fact that,

$$\frac{\partial \underline{x}_4}{\partial y} = \begin{pmatrix} \lambda^n & 0 \\ 0 & \mu^n \end{pmatrix} \frac{\partial \underline{x}_3}{\partial y} = \begin{pmatrix} B_2 \delta (\lambda \mu)^n + A_2 \beta \lambda^{2n} \\ D_2 \delta \mu^{2n} + C_2 \beta (\lambda \mu)^n \end{pmatrix} = O(\mu^{2n}). \quad (3.24)$$

Finally,

$$\begin{aligned} \frac{\partial \underline{x}_5}{\partial y} &= D\hat{F}_{t,a}^N(\underline{x}_4) \frac{\partial \underline{x}_4}{\partial y} \\ &= \begin{pmatrix} A_4 B_2 \delta (\lambda \mu)^n + A_4 A_2 \beta \lambda^{2n} \\ C_4 B_2 \delta (\lambda \mu)^n + C_4 A_2 \beta \lambda^{2n} \end{pmatrix} \\ &\quad + \begin{pmatrix} B_4 D_2 \delta \mu^{2n} + B_4 C_2 \beta (\lambda \mu)^n \\ D_4 D_2 \delta \mu^{2n} + D_4 C_2 \beta (\lambda \mu)^n \end{pmatrix} = O(\mu^{2n}). \end{aligned} \quad (3.25)$$

The last set follows. \square

Lemma 3.26. *The partial derivatives at the critical point c of $F_{t,a}$ satisfy the following,*

- $D_2 = D(\underline{x}_2(c), t, a) = O((\lambda(t, a)/\mu(t, a))^n)$,
- $\partial D_2 / \partial x = \gamma \partial D / \partial y = O(\gamma)$,
- $\partial D_2 / \partial y = \delta \mu^n(t, a) \partial D / \partial y (1 + O((\lambda(t, a)/\mu(t, a))^n))$

where $\delta \partial D / \partial y$ is uniformly bounded and uniformly away from zero and γ can be taken arbitrarily small by shrinking the domain of linearization.

Proof. Observe that

$$\frac{\partial \underline{x}_2}{\partial x} = \begin{pmatrix} \lambda^n & 0 \\ 0 & \mu^n \end{pmatrix} Dh_{t,a}(\underline{x}) \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} \lambda^n \left(1 + \frac{\alpha}{\mu^n}\right) \\ \gamma \end{pmatrix}.$$

As a consequence, if $\underline{x}_2 = (x_2, y_2)$,

$$\frac{\partial D_2}{\partial x} = \frac{\partial D}{\partial x} \frac{\partial x_2}{\partial x} + \frac{\partial D}{\partial y} \frac{\partial y_2}{\partial x} = O(\gamma).$$

The second estimate follows by applying Remark 3.16. Using (3.22), we get

$$\frac{\partial D_2}{\partial y} = \frac{\partial D}{\partial x} \frac{\partial x_2}{\partial y} + \frac{\partial D}{\partial y} \frac{\partial y_2}{\partial y} = O(\lambda^n) + \frac{\partial D}{\partial y} \delta \mu^n, \quad (3.27)$$

where, by Remark 3.17, δ is uniformly away from zero. Moreover $\partial D / \partial y \neq 0$ is also uniformly away from zero because \underline{x}_2 is close to $q_3(t, a)$. In fact $\partial D / \partial y(q_3(t, 0))$ is bounded

away from zero since $q_1(t, 0)$ is a non degenerate homoclinic tangency. The third estimate follows. It is left to prove the first estimate. Let c be a critical point of $F_{t,a}$ with critical value v . Let L be the almost horizontal leaf passing through v and let $w = (1, w_2)$ be the tangent vector to $h(L)$ at $h(v)$. Then

$$w_2 = O\left(\left(\frac{\lambda}{\mu}\right)^n\right).$$

Observe that, because $\underline{x}_3 = h(v)$, we have $\partial \underline{x}_3 / \partial y = \theta w$ with $\theta \neq 0$. In particular, using (3.23), we get

$$\frac{C_2\beta\lambda^n + D_2\delta\mu^n}{A_2\beta\lambda^n + B_2\delta\mu^n} = O\left(\left(\frac{\lambda}{\mu}\right)^n\right).$$

Using the fact that B_2 is away from zero, see Remark 3.17, the first estimate follows. \square

Proof of Proposition 3.13. We will show that, if \underline{x} is a critical point of $F_{t,a}$,

$$D\Phi_{t,a}(\underline{x}) = \begin{pmatrix} \phi_{11} & \phi_{12} \\ \phi_{21} & \phi_{22} \end{pmatrix} = \begin{pmatrix} -1 + O(\gamma) & B_\phi\mu(t, a)^n \\ C_\phi\mu(t, a)^{2n} & D_\phi\mu(t, a)^{3n} \end{pmatrix} \quad (3.28)$$

where B_ϕ and D_ϕ are uniformly bounded and uniformly away from zero and C_ϕ can be taken arbitrarily small by shrinking the domain of linearization. Moreover, according to Remark 3.16, γ can also be taken small. The proposition is proved by applying the implicit function theorem.

The top row of the matrix in (3.28) is given by

$$\begin{aligned} \phi_{11} &= \left[Dh_{t,a}^{-1}(\underline{x}_3) DF_{t,a}^N(\underline{x}_2) \begin{pmatrix} \lambda^n & 0 \\ 0 & \mu^n \end{pmatrix} Dh_{t,a}(\underline{x}) \right]_{11} - 1, \\ \phi_{12} &= \left[Dh_{t,a}^{-1}(\underline{x}_3) DF_{t,a}^N(\underline{x}_2), \begin{pmatrix} \lambda^n & 0 \\ 0 & \mu^n \end{pmatrix} Dh_{t,a}(\underline{x}) \right]_{12}. \end{aligned}$$

Using the notation introduced above, a calculation shows that

$$\begin{aligned} \phi_{11} &= \left(1 + \frac{\alpha_3}{\mu^n}\right) \left(A_2\lambda^n + A_2\alpha \left(\frac{\lambda}{\mu}\right)^n + B_2\gamma\right) + \beta_3 \left(C_2\lambda^n + C_2\alpha \left(\frac{\lambda}{\mu}\right)^n + D_2\gamma\right) - 1 \\ &= -1 + O(\gamma), \end{aligned} \quad (3.29)$$

where we also used that $D_2 = O(\lambda^n/\mu^n)$, see Lemma 3.26. Similarly,

$$\begin{aligned} \phi_{12} &= \left(1 + \frac{\alpha_3}{\mu^n}\right) (A_2\beta\lambda^n + B_2\delta\mu^n) + \beta_3 (C_2\beta\lambda^n + D_2\delta\mu^n) \\ &= B_2\delta\mu^n + O(1) = B_\phi\mu^n, \end{aligned} \quad (3.30)$$

where B_ϕ is uniformly bounded and uniformly away from zero, see Remark 3.17. Observe that, the second component of the function $\Phi_{t,a}(\underline{x})$ is

$$\begin{aligned} &\left[Dh_{t,a}^{-1}(\underline{x}_5) D\hat{F}_{t,a}^N(\underline{x}_4) \begin{pmatrix} \lambda^n & 0 \\ 0 & \mu^n \end{pmatrix} D\hat{F}_{t,a}^N(\underline{x}_2) \begin{pmatrix} \lambda^n & 0 \\ 0 & \mu^n \end{pmatrix} Dh_{t,a}(\underline{x}) \right]_{12} \\ &= \delta D_2 B_4 \mu^{2n} + \delta D_2 D_4 \beta_5 \mu^{2n} + \delta D_2 B_4 \alpha_5 \mu^n + \delta B_2 A_4 (\lambda\mu)^n + \delta B_2 C_4 \beta_5 (\lambda\mu)^n \\ &+ \delta B_2 A_4 \alpha_5 \lambda^n + \beta C_2 B_4 (\lambda\mu)^n + \beta C_2 D_4 \beta_5 (\lambda\mu)^n + \beta C_2 B_4 \alpha_5 \lambda^n + \beta A_2 A_4 \lambda^{2n} \\ &+ \beta A_2 C_4 \beta_5 \lambda^{2n} + \beta A_2 A_4 \alpha_5 \left(\frac{\lambda^2}{\mu}\right)^n. \end{aligned} \quad (3.31)$$

In the following calculation we often use that $\lambda\mu^3 < 1$. Using Lemma 3.18 and Lemma 3.26 we have

$$\begin{aligned}
\phi_{21} &= \frac{\partial}{\partial x} [\delta D_2 B_4 \mu^{2n} + \delta D_2 D_4 \beta_5 \mu^{2n} + \delta D_2 B_4 \alpha_5 \mu^n] + O((\lambda\mu^2)^n) \\
&= \frac{\partial D_2}{\partial x} B_4 \delta \mu^{2n} + \frac{\partial D_2}{\partial x} D_4 \delta \beta_5 \mu^{2n} + O(\mu^n) \\
&= C_\phi \mu^{2n}.
\end{aligned} \tag{3.32}$$

Observe that $C_\phi = O(\partial D_2 / \partial x) = O(\gamma)$, see Lemma 3.26. Similarly, by using Lemma 3.21 and Lemma 3.26 we have

$$\begin{aligned}
\phi_{22} &= \frac{\partial}{\partial y} [\delta D_2 B_4 \mu^{2n} + \delta D_2 D_4 \beta_5 \mu^{2n} + \delta D_2 B_4 \alpha_5 \mu^n] + O((\lambda\mu^3)^n) \\
&= \frac{\partial D_2}{\partial y} B_4 \delta \mu^{2n} + \frac{\partial D_2}{\partial y} D_4 \delta \beta_5 \mu^{2n} + O(\mu^{2n}) \\
&= D_\phi \mu^{3n}.
\end{aligned} \tag{3.33}$$

By Remark 3.16, Remark 3.17 and Lemma 3.26, D_ϕ is uniformly bounded away from zero. From (3.29), (3.30), (3.32) and (3.33) we have that $D\Phi_{t,a}(\underline{x})$ is non singular. The proposition follows by applying the implicit function theorem. \square

3.3 Global existence of the critical point

In this subsection we prove that all maps in \mathcal{H}_n have a critical point, namely $\mathcal{C}_n = \mathcal{H}_n$. To do that we need to study the a -dependence of the critical point and critical value. From propositions 3.13 and 3.47, for all $(t, a) \in \mathcal{C}_n$, the map $F_{t,a}$ has a critical point $(c_x(t, a), c_y(t, a))$ with critical value $(v_x(t, a), v_y(t, a))$, where $v_x(t, a) = c_x(t, a)$.

Proposition 3.34. *For every $(t, a) \in \mathcal{C}_n$,*

$$\begin{aligned}
\frac{\partial c_x}{\partial a} &= \frac{\partial v_x}{\partial a} = O(n), \\
\frac{\partial c_y}{\partial a} &= O\left(\frac{n}{\mu^n(t, a)}\right), \\
\frac{\partial v_y}{\partial a} &= \delta_3 + O\left(\frac{n}{\mu^n(t, a)}\right).
\end{aligned}$$

In particular,

$$|v_y - c_y| = O\left(\frac{1}{\mu^{2n}(t, a)}\right).$$

The proof of the previous proposition needs some preparation. Fix $(t, a) \in \mathcal{C}_n$. Let $(x, y) \in HB(t, a) \cap \mathbb{R}^2$ and $\underline{x} = (x, y)$, \underline{x}_1 , \underline{x}_2 , \underline{x}_3 , \underline{x}_4 , \underline{x}_5 as introduced before. Observe that all coefficients introduced, $\alpha, \beta, \dots, A_2, B_2, \dots, \alpha_3, \beta_3, \dots, A_4, B_4, \dots, \alpha_5, \dots$ are functions of the point $(x, y) \in HB(t, a) \cap \mathbb{R}^2$ and a .

Lemma 3.35. *The partial derivatives at the critical point c of $F_{t,a}$ satisfy the following.*

- $\partial\alpha/\partial a, \partial\beta/\partial a, \partial\gamma/\partial a, \partial\delta/\partial a = O(1)$,
- $\partial A_2/\partial a, \partial B_2/\partial a, \partial C_2/\partial a, \partial D_2/\partial a = O(n)$,

- $\partial\alpha_3/\partial a, \partial\beta_3/\partial a, \partial\gamma_3/\partial a, \partial\delta_3/\partial a = O(n)$,
- $\partial A_4/\partial a, \partial B_4/\partial a, \partial C_4/\partial a, \partial D_4/\partial a = O(\mu^n(t, a))$,
- $\partial\alpha_5/\partial a, \partial\beta_5/\partial a, \partial\gamma_5/\partial a, \partial\delta_5/\partial a = O(\mu^n(t, a))$.

Proof. The first set of estimates follows by the smoothness of the family of maps $h_{t,a}$. For the second set of estimates, observe that, because $h_{t,a}(x, 0) = (x, 0)$ and $y = O(1/\mu^n)$, we have

$$\frac{\partial \underline{x}_1}{\partial a} = \frac{\partial h}{\partial a} = O\left(\frac{1}{\mu^n}\right).$$

Recall that $\underline{x}_2 = \hat{F}_{t,a}^n(\underline{x}_1, t, a)$. It follows that

$$\frac{\partial x_2}{\partial a} = n\lambda^{n-1} \frac{\partial \lambda}{\partial a} x_1 + \lambda^n \frac{\partial x_1}{\partial a} = O(n\lambda^n), \quad (3.36)$$

and

$$\frac{\partial y_2}{\partial a} = n\mu^{n-1} \frac{\partial \mu}{\partial a} y_1 + \mu^n \frac{\partial y_1}{\partial a} = O(n), \quad (3.37)$$

where we used that $y_1 = O(1/\mu^n)$. As a consequence,

$$\frac{\partial A_2}{\partial a} = \frac{\partial A}{\partial x} \frac{\partial x_2}{\partial a} + \frac{\partial A}{\partial y} \frac{\partial y_2}{\partial a} + \frac{\partial A}{\partial a} = O(n).$$

Similarly the other estimates in the second statement follows. Recall that $\underline{x}_3 = \hat{F}_{t,a}^N(\underline{x}_2, t, a)$. It follows that

$$\frac{\partial x_3}{\partial a} = A_2 \frac{\partial x_2}{\partial a} + B_2 \frac{\partial y_2}{\partial a} + O(1) = O(n), \quad (3.38)$$

and

$$\frac{\partial y_3}{\partial a} = C_2 \frac{\partial x_2}{\partial a} + D_2 \frac{\partial y_2}{\partial a} + O(1) = O(1), \quad (3.39)$$

where we used (3.36), (3.37) and that $D_2 = O(\lambda^n/\mu^n)$, see Lemma 3.26. The third set of estimates follow. Recall now that $\underline{x}_4 = \hat{F}_{t,a}^n(\underline{x}_3, t, a)$. It follows that,

$$\frac{\partial x_4}{\partial a} = n\lambda^{n-1} \frac{\partial \lambda}{\partial a} x_3 + \lambda^n \frac{\partial x_3}{\partial a} = O(n\lambda^n), \quad (3.40)$$

and

$$\frac{\partial y_4}{\partial a} = n\mu^{n-1} \frac{\partial \mu}{\partial a} y_3 + \mu^n \frac{\partial y_3}{\partial a} = O(\mu^n), \quad (3.41)$$

where we used (3.38), (3.39) and that $y_3 = O(1/\mu^n)$. As a consequence,

$$\frac{\partial A_4}{\partial a} = \frac{\partial A}{\partial x} \frac{\partial x_4}{\partial a} + \frac{\partial A}{\partial y} \frac{\partial y_4}{\partial a} + \frac{\partial A}{\partial a} = O(\mu^n).$$

Similarly the other estimates in the fourth statement follows. Finally, recall that $\underline{x}_5 = \hat{F}_{t,a}^N(\underline{x}_4, t, a)$. It follows that,

$$\frac{\partial x_5}{\partial a} = A_4 \frac{\partial x_4}{\partial a} + B_4 \frac{\partial y_4}{\partial a} + O(1) = O(\mu^n), \quad (3.42)$$

and

$$\frac{\partial y_5}{\partial a} = C_4 \frac{\partial x_4}{\partial a} + D_4 \frac{\partial y_4}{\partial a} + O(1) = O(\mu^n), \quad (3.43)$$

where we used (3.40), (3.41). As before the last set follows. \square

Lemma 3.44. *The partial derivative of y_3 at the critical point c of $F_{t,a}$ satisfy the following:*

$$\frac{\partial y_3}{\partial a} = 1 + O\left(\frac{1}{\mu^n(t, a)}\right).$$

Proof. For $a = 0$, let $q_3(t, 0)$ be the point where $D(q_3, t, 0) = 0$. Recall that $\hat{F}_{t,0}^N(q_3) = q_1$ where q_1 is the tangency at $a = 0$. Because $\partial D(q_3, t, 0)/\partial y \neq 0$, there is a curve through q_3 transversal to the y -axis where $D(\underline{x}, t, 0) = 0$ for all points \underline{x} in the curve. The transversality implies that there is a curve $D(\underline{x}, t, a) = 0$, transversally intersecting the y -axis in the point $q_3(t, a)$ at the distance of $O(1/\mu^n)$ to the curve $D(\underline{x}, t, 0) = 0$, because $(t, a) \in \mathcal{C}_n$. From Lemma 3.26 we know that $D_2 = O(\lambda^n/\mu^n)$. This implies that the distance from \underline{x}_2 to this curve is of $O(\lambda^n/\mu^n)$. Moreover $x_2 = O(\lambda^n)$. Hence,

$$\text{dist}(\underline{x}_2, q_3(t, 0)) = O\left(\frac{1}{\mu^n}\right).$$

Because $\partial\left(\hat{F}_{t,0}^N(q_3(t, 0))\right)_y/\partial a = 1$, we get

$$\frac{\partial\left(\hat{F}_{t,a}^N(\underline{x}_2)\right)_y}{\partial a} = 1 + O\left(\frac{1}{\mu^n}\right).$$

Now (3.39), becomes

$$\frac{\partial y_3}{\partial a} = C_2 \frac{\partial x_2}{\partial a} + D_2 \frac{\partial y_2}{\partial a} + 1 + O\left(\frac{1}{\mu^n}\right) = 1 + O\left(\frac{1}{\mu^n}\right),$$

where we also used (3.36), (3.37) and Lemma 3.26. □

Proof of Proposition 3.34. Observe that

$$\frac{\partial c}{\partial a} = (D\Phi)^{-1} \frac{\partial \Phi}{\partial a}.$$

Let $\Phi = (\Phi_1, \Phi_2)$. We start by calculating $\partial\Phi_1/\partial a$. Observe that $\Phi_1 = (h_{t,a}^{-1}(\underline{x}_3) - x)_x$. As a consequence,

$$\frac{\partial\Phi_1}{\partial a} = \left(1 + \frac{\alpha_3}{\mu^n}\right) \frac{\partial x_3}{\partial a} + \beta_3 \frac{\partial y_3}{\partial a} + \frac{\partial h^{-1}}{\partial a} = O(n), \quad (3.45)$$

where we used (3.38) and (3.39). In order to calculate $\partial\Phi_2/\partial a$ we take the a -derivative of (3.31). Using Lemma 3.35, the partial derivatives of all terms in (3.31) with a factor λ^n will give a contribution of $O((\lambda\mu^2)^n)$. We get

$$\begin{aligned} \frac{\partial\Phi_2}{\partial a} &= \frac{\partial}{\partial a} [\delta D_2 B_4 \mu^{2n} + \delta D_2 D_4 \beta_5 \mu^{2n} + \delta D_2 B_4 \alpha_5 \mu^n] \\ &+ O((\lambda\mu^2)^n). \end{aligned}$$

Using Lemma 3.35 and the fact that $D_2 = O(\lambda^n/\mu^n)$, see Lemma 3.26, we get

$$\frac{\partial\Phi_2}{\partial a} = O(n\mu^{2n}). \quad (3.46)$$

Because the determinant of $D\Phi$ is proportional to μ^{3n} , see (3.28), we have,

$$\frac{\partial c}{\partial a} = (D\Phi)^{-1} \frac{\partial \Phi}{\partial a} = \begin{pmatrix} O(1) & O(1/\mu^{2n}) \\ O(1/\mu^n) & O(1/\mu^{3n}) \end{pmatrix} \begin{pmatrix} O(n) \\ O(n\mu^{2n}) \end{pmatrix} = \begin{pmatrix} O(n) \\ O(n/\mu^n) \end{pmatrix},$$

where we used (3.45) and (3.46). The first two equations follow. Observe that $v_y = (h_{t,a}^{-1}(\underline{x}_3))_y$. Hence,

$$\frac{\partial v_y}{\partial a} = \frac{\gamma_3}{\mu^n} \frac{\partial x_3}{\partial a} + \delta_3 \frac{\partial y_3}{\partial a} + O\left(\frac{1}{\mu^n}\right) = \delta_3 + O\left(\frac{n}{\mu^n}\right),$$

where we used (3.38) and Lemma 3.44. The last statement is an immediate consequence of the fact that $(t, a) \in \mathcal{C}_n \subset \mathcal{H}_n$. \square

Lemma 3.47. *For n large enough, $\mathcal{H}_n = \mathcal{C}_n$.*

Proof. Let $(t, sa_n(t)) \in \mathcal{C}_n$. Then $p(t) = (2, p_y(t)) \in \{2\} \times [\delta^-, \delta^+] = HB_2(t, sa_n(t)) \cap \mathbb{R}^2$. Observe that

$$\left| \left(F_{(t, sa_n(t))}^{n+N} (2, \delta^\pm) \right)_x - 2 \right| = 1.$$

As a consequence, by Lemma 3.1, there exists a constant $K > 0$ such that

$$1 = \left| \left(F_{(t, sa_n(t))}^{n+N} (p(t)) - F_{(t, sa_n(t))}^{n+N} (\delta^\pm) \right)_x \right| \leq \frac{1}{K} |p_y(t) - \delta^\pm| \mu^n.$$

Hence,

$$|p_y(t) - \delta^\pm| \geq \frac{K}{\mu^n}. \quad (3.48)$$

For $(t, a) \in \mathcal{H}_n$ let $\partial^\pm HB(t, a)$ be the two almost horizontal curves in the boundary of $HB(t, a) \cap \mathbb{R}^2$. Notice that $\partial^\pm HB$ are graphs of functions satisfying, by Lemma 12 in [4],

$$\frac{\partial}{\partial a} (\partial^\pm HB) = O\left(\frac{n}{\mu^n}\right), \quad (3.49)$$

and by Lemma 3.1,

$$\frac{\partial}{\partial x} (\partial^\pm HB) = O\left(\frac{1}{\mu^n}\right). \quad (3.50)$$

Choose $(t, a) \in \partial \mathcal{C}_n$ and suppose $(t, a) \notin \partial \mathcal{H}_n$. Then either $c = c(t, a)$ or $v = v(t, a)$ are in the boundary of $HB(t, a)$. Because of Proposition 3.34,

$$|c_x - 2| = |v_x - 2| = O\left(\frac{n}{\mu^{2n}}\right). \quad (3.51)$$

Hence,

$$\{c, v\} \cap \{\partial^+ HB \cup \partial^- HB\} \neq \emptyset. \quad (3.52)$$

Observe that $c(t, sa_n(t)) = v(t, sa_n(t)) = p(t)$ and by Proposition 3.34,

$$|c - p(t)|, |v - p(t)| = O\left(\frac{n}{\mu^{2n}}\right). \quad (3.53)$$

Consider the case $c \in \partial^+ HB$. The other cases described in (3.52) can be treated in exactly the same way. Because of (3.49), (3.50), and (3.51), we get,

$$|\delta^+ - c| = O\left(\frac{n}{\mu^{2n}}\right).$$

This last equation together with (3.48) and (3.53) lead to a contradiction. Namely,

$$\frac{K}{\mu^n} \leq |p_y(t) - \delta^+| \leq |p(t) - c| + |c - (2, \delta^+)| = O\left(\frac{n}{\mu^{2n}}\right).$$

Hence $(t, a) \in \partial\mathcal{H}_n$ and $\mathcal{C}_n = \mathcal{H}_n$. □

3.4 The normalized map

In this subsection we rescale the domain of the straightened map to unit size. Fix $(t, a) \in \mathcal{H}_n$. Recall that the return map after straightening became

$$\tilde{F}_{t,a} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} g_{t,a}(x, y) \\ x \end{pmatrix}.$$

In particular, the map $x \mapsto g_{t,a}(x, c_x(t, a))$ has a critical point at $c_{t,a} = c_x(t, a)$. Similarly as in Definition 3.12, the point $(c_{t,a}, c_{t,a})$ is a critical point of $\tilde{F}_{t,a}$ in the sense that

$$\tilde{F}_{t,a} \begin{pmatrix} c_{t,a} \\ c_{t,a} \end{pmatrix} = \begin{pmatrix} v_{t,a} \\ c_{t,a} \end{pmatrix}$$

and the image of the horizontal slice at $y = c_{t,a}$ is tangent to the vertical slice at $x = v_{t,a}$.

Lemma 3.54. *For all $(t, a) \in \mathcal{H}_n$,*

$$\frac{\partial v_{t,a}}{\partial a} = (B_4 + D_4\beta_5) \mu^n(t, a) + O(n).$$

In particular,

$$|v_{t,a} - c_{t,a}| = O\left(\frac{1}{\mu^n(t, a)}\right).$$

Proof. Observe that $v_{t,a} = (h_{t,a}^{-1}(\underline{x}_5))_x$. To calculate its a -derivative we improve the derivatives in the proof of Lemma 3.35. By Lemma 3.44, we get

$$\frac{\partial y_4}{\partial a} = n\mu^{n-1} \frac{\partial \mu}{\partial a} y_3 + \mu^n \frac{\partial y_3}{\partial a} = \mu^n + O(n).$$

By this, by (3.40) and by (3.41) we have,

$$\frac{\partial x_5}{\partial a} = A_4 \frac{\partial x_4}{\partial a} + B_4 \frac{\partial y_4}{\partial a} + O(1) = B_4 \mu^n + O(n),$$

and

$$\frac{\partial y_5}{\partial a} = C_4 \frac{\partial x_4}{\partial a} + D_4 \frac{\partial y_4}{\partial a} + O(1) = D_4 \mu^n + O(n).$$

Finally,

$$\frac{\partial v_{t,a}}{\partial a} = \left(1 + \frac{\alpha_5}{\mu^n}\right) \frac{\partial x_5}{\partial a} + \beta_5 \frac{\partial y_5}{\partial a} + O\left(\frac{1}{\mu^n}\right) = (B_4 + \beta_5 D_4) \mu^n + O(n).$$

The last equality follows from the fact that $(t, a) \in \mathcal{H}_n$. □

We rewrite

$$\tilde{F}_{t,a} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} q_{t,a}(x) + \tilde{\epsilon}_{t,a}(x, y) \\ x \end{pmatrix}$$

where $q_{t,a}(x) = g_{t,a}(x, c_{t,a})$. Observe that, the domain of $q_{t,a}$ is the simply connected domain $\pi_x(\sigma_{t,a}(D_{c_{t,a}})) = \tilde{D} \subset \mathbb{D}_1$. Moreover, because $p_x(t) = c_x(t, sa_n(t)) = 2$ and because of Proposition 3.34, we have

$$|c_{t,a} - 2| = O\left(\frac{n}{\mu^{2n}(t, a)}\right).$$

In particular, $c_{t,a}$ is exponentially close to the center of \mathbb{D}_1 . Because $|v_y| = O(1/\mu^n(t, a))$ and for every $(x, y) \in HB(t, a)$, by Lemma 3.4, $|y| = O(1/\mu^n(t, a))$ we get

$$\text{diam}(F_{t,a}^n(D_{c_{t,a}})) = O\left(\frac{1}{\mu^{n/2}(t, a)}\right). \quad (3.55)$$

In fact, $D_{c_{t,a}} \subset HB(t, a)$ with $F_{t,a}^{n+N}(D_{c_{t,a}}) \subset HB(t, a)$. Because $\pi_y(F_{t,a}^N(F_{t,a}^n(D_{c_{t,a}})))$ is contained in a disc of radius $1/\mu^n(t, a)$ centered in zero and $\pi_y(F_{t,a}^N(F_{t,a}^n(HB(t, a))))$ contains a disc of uniform size centered in zero, the estimate in (3.55) follows. Since $\tilde{D} = \pi_x(F_{t,a}^N(F_{t,a}^n(D_{c_{t,a}})))$ we have

$$\text{diam}(\tilde{D}) = O\left(\frac{1}{\mu^{n/2}(t, a)}\right). \quad (3.56)$$




Figure 3: The factorization

We will need to introduce a rescaling to bring this domain back to unit size. The following construction is illustrated in Figure 3. We factorize the map $q_{t,a}$ as

$$q_{t,a} = \varphi((z - c_{t,a})^2) \quad (3.57)$$

where $\varphi : D_0 \rightarrow \mathbb{D}_1$ is univalent and onto and its domain D_0 is the image under the map $z \mapsto (z - c_{t,a})^2$ of \tilde{D} . Let $V_1 \subset \mathbb{D}_1$ be the disc of radius $1/2$ centered at 2 . Consider the map

$$q_{t,a} : q_{t,a}^{-1}(V_1) \rightarrow V_1.$$

Let $D_1 \subset D_0$ be the disc which is the image under the map $z \mapsto (z - c_{t,a})^2$ of $q_{t,a}^{-1}(V_1)$. Then the restriction $\varphi : D_1 \rightarrow V_1$ has uniformly bounded distortion, because it has univalent extension up to the unit disc \mathbb{D}_1 .

Observe that $q_{t,a}^{-1}(V_1) \subset \tilde{D}$ has diameter proportional to the diameter of \tilde{D} which is proportional to $1/\mu^{n/2}(t, a)$, see (3.56). Hence, D_1 has diameter proportional to $1/\mu^n(t, a)$. Because φ has bounded distortion, this implies that $D\varphi(0)$ is proportional to $\mu^n(t, a)$. Define $K_{t,a}$ by

$$v_{t,a} - c_{t,a} = \frac{K_{t,a}}{D\varphi(0)}. \quad (3.58)$$

and $K = \max_{t,a} K_{t,a}$. Because of Lemma 3.54, we have that K is uniformly bounded.

Let $s > 0$ be the unique solution of $s + 1 = s^2 K/2$ and

$$U = \left\{ z \mid |z - c_{t,a}| \leq \frac{sK}{D\varphi(0)} \right\} \subset q_{t,a}^{-1}(V_1).$$

The image of U under the map $z \rightarrow (z - c_{t,a})^2$ is denoted by U^2 . Observe that U^2 is a round disc, centered around zero, of radius $s^2 K^2 / D\varphi(0)^2$. In particular the diameter of U^2 is proportional to $1/\mu^{2n}(t, a)$ and the diameter of $\varphi(U^2)$ is proportional to $1/\mu^n(t, a)$. This implies that, if $z \in U^2$, then using the Koebe Lemma as stated in [1], [2] or [7], we have

$$\frac{D\varphi(z)}{D\varphi(0)} = 1 + O\left(\frac{1}{\mu^n(t, a)}\right). \quad (3.59)$$

This implies that, for all $z \in U^2$,

$$\varphi(z) = \varphi(0) + D\varphi(0) \int_0^z \frac{D\varphi(z)}{D\varphi(0)} dz = v_{t,a} + D\varphi(0)z + O\left(\frac{1}{\mu^{2n}(t, a)}\right). \quad (3.60)$$

In particular $\varphi(U^2)$ contains a round disc V centered around $v_{t,a}$ of radius $3s^2 K^2 / 4D\varphi(0)$. Moreover, if $z \in U$,

$$|z - v_{t,a}| \leq |z - c_{t,a}| + |c_{t,a} - v_{t,a}| \leq \frac{sK}{D\varphi(0)} + \frac{K}{D\varphi(0)} = \frac{s^2 K^2}{2D\varphi(0)} < \frac{3s^2 K^2}{4D\varphi(0)}.$$

Hence,

$$U \subset V \subset q_{t,a}(U).$$

Because $sK = 1 + \sqrt{1 + 2K}$, $\text{diam}(U)D\varphi(0) = sK \geq 2$ and $\text{mod}(U, V) \geq \log(3/2)$. In particular the map $q_{t,a} : U \rightarrow V$ is a quadratic-like map with uniform modulus.

The last step is to rescale the domain U to a finite size. Let $\alpha : x \rightarrow c_{t,a} + x/D\varphi(0)$, $U_{t,a} = \alpha^{-1}(U)$ and $V_{t,a} = \alpha^{-1}(V)$. Moreover, let $Z_{t,a} : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ defined as

$$Z_{t,a} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \alpha(x) \\ \alpha(y) \end{pmatrix}.$$

We define now the domain of the Hénon-like map $HF_{t,a}$ as

$$\text{Dom}(HF_{t,a}) = U_{t,a} \times U_{t,a}$$

and

$$HF_{t,a} = Z_{t,a}^{-1} \circ \tilde{F}_{t,a} \circ Z_{t,a} : \text{Dom}(HF_{t,a}) \rightarrow \mathbb{C} \times \mathbb{C}.$$

By construction

$$HF_{t,a} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} f_{t,a}(x, y) \\ x \end{pmatrix}.$$

Theorem 3.61. *For every $(t, a) \in \mathcal{H}_n$, there exists $\nu_{t,a} \in \mathbb{R}$ such that*

$$HF_{t,a} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} f_{t,a}(x, y) \\ x \end{pmatrix} = \begin{pmatrix} x^2 + \nu_{t,a} + \epsilon_{t,a}(x, y) \\ x \end{pmatrix},$$

with $\epsilon_{t,a}(0, 0) = 0$ and

$$|\epsilon_{t,a}| = O\left(\frac{1}{\mu(t, a)^n}\right).$$

Moreover, for every $y \in U_{t,a}$, the map $f_{t,a}(\cdot, y) : U_{t,a} \rightarrow \mathbb{C}$ is quadratic-like with

$$\begin{aligned} V_{t,a} &\subset f_{t,a}(U_{t,a}, y), \\ \text{mod}(U_{t,a}, V_{t,a}) &\geq \log\left(\frac{3}{2}\right), \end{aligned}$$

and

$$\text{diam}(U_{t,a}) \geq 2.$$

Proof. The construction of $HF_{t,a}$ gives the formula for the y -coordinate. Observe that, by using (3.60),

$$\begin{aligned} f_{t,a}(x, 0) &= \alpha^{-1} \circ q_{t,a} \circ \alpha = D\varphi(0) \left[\varphi \left(\left[\frac{x}{D\varphi(0)} \right]^2 \right) - c_{t,a} \right] \\ &= D\varphi(0) \left[v_{t,a} + \frac{x^2}{D\varphi(0)} + O\left(\frac{1}{\mu^{2n}}\right) - c_{t,a} \right] \\ &= x^2 + \nu_{t,a} + O\left(\frac{1}{\mu^n}\right), \end{aligned}$$

where, by (3.58),

$$\nu_{t,a} = D\varphi(0)(v_{t,a} - c_{t,a}) = O(1). \quad (3.62)$$

By construction, for $y = 0$, we have $V_{t,a} \subset f_{t,a}(U_{t,a}, 0)$. Take $(x_0, y_0) \in U_{t,a} \times U_{t,a}$ and compare its image with the image of $(x_0, 0)$. Let

$$Z_{t,a} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} \tilde{x} \\ \tilde{y} \end{pmatrix} \quad \text{and} \quad Z_{t,a} \begin{pmatrix} x_0 \\ 0 \end{pmatrix} = \begin{pmatrix} \tilde{x} \\ c_{t,a} \end{pmatrix},$$

then

$$\tilde{y} = \frac{y_0}{D\varphi(0)} + c_{t,a}.$$

Let

$$\begin{pmatrix} x \\ y \end{pmatrix} = \sigma_{t,a}^{-1} \begin{pmatrix} \tilde{x} \\ \tilde{y} \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} x^{(0)} \\ y^{(0)} \end{pmatrix} = \sigma_{t,a}^{-1} \begin{pmatrix} \tilde{x} \\ c_{t,a} \end{pmatrix},$$

then (x, y) and $(x^{(0)}, y^{(0)})$ are on the same leaf of the almost horizontal foliation and

$$|x - x^{(0)}| = \frac{y_0}{D\varphi(0)}. \quad (3.63)$$

Let

$$\begin{pmatrix} x_2 \\ y_2 \end{pmatrix} = F_{t,a}^n \begin{pmatrix} x \\ y \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} x_2^{(0)} \\ y_2^{(0)} \end{pmatrix} = F_{t,a}^n \begin{pmatrix} x^{(0)} \\ y^{(0)} \end{pmatrix},$$

then, because of Lemma 3.1, the fact that (x_2, y_2) and $(x_2^{(0)}, y_2^{(0)})$ are on the same leaf of the foliation and (3.63), we have

$$\text{dist} \left(\begin{pmatrix} x_2 \\ y_2 \end{pmatrix}, \begin{pmatrix} x_2^{(0)} \\ y_2^{(0)} \end{pmatrix} \right) = O(\lambda^n),$$

where we used that $D\varphi(0)$ is proportional to μ^n . Let

$$\begin{pmatrix} x_3 \\ y_3 \end{pmatrix} = F_{t,a}^N \begin{pmatrix} x_2 \\ y_2 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} x_3^{(0)} \\ y_3^{(0)} \end{pmatrix} = F_{t,a}^N \begin{pmatrix} x_2^{(0)} \\ y_2^{(0)} \end{pmatrix},$$

then

$$x_3 = x_3^{(0)} \quad \text{and} \quad y_3 - y_3^{(0)} = O(\lambda^n). \quad (3.64)$$

The next step is to obtain an estimate for the difference of the x coordinates of $\sigma_{t,a}(x_3, y_3)$ and $\sigma_{t,a}(x_3^{(0)}, y_3^{(0)})$. This can be estimated by iterating by $F_{t,a}^{n+N}$. Lemma 3.1 says that this map extends distance at most by μ^n . Let $x_5 = (\sigma_{t,a}(x_3, y_3))_x$ and $x_5^{(0)} = (\sigma_{t,a}(x_3^{(0)}, y_3^{(0)}))_x$ and we obtain

$$|x_5 - x_5^{(0)}| = O((\lambda\mu)^n),$$

where we used (3.64). Finally, by applying α^{-1} one obtains

$$|f_{t,a}(x_0, y_0) - f_{t,a}(x_0, 0)| = O((\lambda\mu^2)^n),$$

where we used that $D\varphi(0)$ is proportional to μ^n . The proposition follows by recalling that $\lambda\mu^3 < 1$. \square

4 Monotonicity of the normalized family

In the previous section, we constructed families of normalized maps $HF_{t,a}$ with $(t, a) \in \mathcal{H}_n$. In this section we study the dependence of the normalized maps on the parameters.

4.1 Monotonicity in the a direction

Fix $t \in [-t_0, t_0]$ and consider the real-analytic one-parameter family

$$[-E, E] \ni \beta \mapsto HF_\beta = HF_{t, sa_n(t) + \beta / |\mu(t, sa_n(t))|^{2n}}.$$

From Theorem 3.61, we know that

$$HF_\beta \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x^2 + \nu + \epsilon_\beta(x, y) \\ x \end{pmatrix} = \begin{pmatrix} f_\beta(x, y) \\ x \end{pmatrix},$$

where ν is a function of β .

Theorem 4.1. Fix $t \in [-t_0, t_0]$. There exists a real-analytic function $[-E, E] \ni \beta \mapsto L(\beta) = D\varphi(0)B_4(c)/\mu^n \neq 0$ such that

$$\frac{\partial f_\beta}{\partial \beta} = L(\beta) + O\left(\frac{n}{\mu^n(t, sa_n(t))}\right).$$

The proof of Theorem 4.1 requires the following lemmas. We use the notation of the previous section.

Lemma 4.2. The mixed partial derivatives at the critical point c of $F_{t,a}$ satisfy the following.

- $\partial^2 \alpha / \partial a \partial y, \partial^2 \beta / \partial a \partial y, \partial^2 \gamma / \partial a \partial y, \partial^2 \delta / \partial a \partial y = O(1),$
- $\partial^2 A_2 / \partial a \partial y, \partial^2 B_2 / \partial a \partial y, \partial^2 C_2 / \partial a \partial y, \partial^2 D_2 / \partial a \partial y = O(n\mu^n),$
- $\partial^2 A_4 / \partial a \partial y, \partial^2 B_4 / \partial a \partial y, \partial^2 C_4 / \partial a \partial y, \partial^2 D_4 / \partial a \partial y = O(n\mu^{2n}),$
- $\partial^2 \alpha_5 / \partial a \partial y, \partial^2 \beta_5 / \partial a \partial y, \partial^2 \gamma_5 / \partial a \partial y, \partial^2 \delta_5 / \partial a \partial y = O(n\mu^{2n}).$

Proof. The first set of estimates follows by the smoothness of the family of maps $h_{t,a}$. For the second set of estimates, observe that, by (3.22) and Lemma 3.35,

$$\frac{\partial^2 \underline{x}_2}{\partial a \partial y} = O(n\mu^n).$$

As a consequence,

$$\begin{aligned} \frac{\partial^2 A_2}{\partial a \partial y} &= \frac{\partial}{\partial a} \left[\frac{\partial A}{\partial x} \frac{\partial x_2}{\partial y} + \frac{\partial A}{\partial y} \frac{\partial y_2}{\partial y} \right] \\ &= \frac{\partial x_2}{\partial y} \left[\frac{\partial^2 A}{\partial x^2} \frac{\partial x_2}{\partial a} + \frac{\partial^2 A}{\partial y \partial x} \frac{\partial y_2}{\partial a} + \frac{\partial^2 A}{\partial a \partial x} \right] + \frac{\partial A}{\partial x} \frac{\partial^2 x_2}{\partial a \partial y} \\ &+ \frac{\partial y_2}{\partial y} \left[\frac{\partial^2 A}{\partial x \partial y} \frac{\partial x_2}{\partial a} + \frac{\partial^2 A}{\partial y^2} \frac{\partial y_2}{\partial a} + \frac{\partial^2 A}{\partial a \partial y} \right] + \frac{\partial A}{\partial y} \frac{\partial^2 y_2}{\partial a \partial y} \\ &= O(n\mu^n), \end{aligned}$$

where we also used (3.22), (3.36) and (3.37). Similarly the other estimates in the second statement follows. For the third set of estimates, observe that, by (3.24) and Lemma 3.35

$$\frac{\partial^2 \underline{x}_4}{\partial a \partial y} = O(n\mu^{2n}),$$

and by (3.24), the fact that $D_2 = O((\lambda/\mu)^n)$, see Lemma 3.26, we get,

$$\frac{\partial \underline{x}_4}{\partial y} = O((\lambda\mu)^n). \quad (4.3)$$

As a consequence,

$$\begin{aligned} \frac{\partial^2 A_4}{\partial a \partial y} &= \frac{\partial x_4}{\partial y} \left[\frac{\partial^2 A}{\partial x^2} \frac{\partial x_4}{\partial a} + \frac{\partial^2 A}{\partial y \partial x} \frac{\partial y_4}{\partial a} + \frac{\partial^2 A}{\partial a \partial x} \right] + \frac{\partial A}{\partial x} \frac{\partial^2 x_4}{\partial a \partial y} \\ &+ \frac{\partial y_4}{\partial y} \left[\frac{\partial^2 A}{\partial x \partial y} \frac{\partial x_4}{\partial a} + \frac{\partial^2 A}{\partial y^2} \frac{\partial y_4}{\partial a} + \frac{\partial^2 A}{\partial a \partial y} \right] + \frac{\partial A}{\partial y} \frac{\partial^2 y_4}{\partial a \partial y} \\ &= O(n\mu^{2n}), \end{aligned}$$

where we also used (3.40) and (3.41). Similarly the other estimates in the third statement follows. Finally, by (3.25), Lemma 3.26 and Lemma 3.35,

$$\frac{\partial^2 \underline{x}_5}{\partial a \partial y} = O(n\mu^{2n}),$$

and by (3.25), the fact that $D_2 = O((\lambda/\mu)^n)$, see Lemma 3.26, we get,

$$\frac{\partial \underline{x}_5}{\partial y} = O((\lambda\mu)^n). \quad (4.4)$$

As a consequence,

$$\frac{\partial^2 \alpha_5}{\partial a \partial y} = O(n\mu^{2n}),$$

where we also used (3.42) and (3.43). Similarly the other estimates in the last statement follows. \square

The following lemma refers to the function Φ . Recall that Φ_2 is the second component of Φ and $\phi_{22} = \partial\Phi_2/\partial y$, see (3.28).

Lemma 4.5. *Let $(t, a) \in \mathcal{H}_n$. Then,*

$$\frac{d(\phi_{22})}{da} = O(\mu^{4n}).$$

Proof. We calculate $d(\phi_{22})/da = \partial^2\Phi_2/\partial a\partial y$ using (3.31). Observe that, by Lemma 3.35, Lemma 3.21 and Lemma 4.2, the terms in (3.31) containing a λ^n will contribute to the estimate of the a -derivative of ϕ_{22} with an order of at most $(\lambda\mu^4)^n$. Hence,

$$\frac{d(\phi_{22})}{da} = \frac{\partial^2}{\partial a \partial y} [\delta D_2 B_4 \mu^{2n} + \delta D_2 D_4 \beta_5 \mu^{2n} + \delta D_2 B_4 \alpha_5 \mu^n] + O((\lambda\mu^4)^n).$$

Observe that the mixed partial derivatives of the coefficients of μ^{2n} , μ^n are of the same order. Hence,

$$\frac{d(\phi_{22})}{da} = O\left(\frac{\partial^2}{\partial a \partial y} [\delta D_2 B_4 \mu^{2n} + \delta D_2 D_4 \beta_5 \mu^{2n}]\right) + O((\lambda\mu^4)^n).$$

Because $D_2 = O((\lambda/\mu)^n)$, see Lemma 3.26, Lemma 3.35, Lemma 3.21 and Lemma 4.2 all the terms which do not involve a partial derivative of D_2 , give a contribution of order at most $(n\lambda\mu^3)^n$. Hence

$$\begin{aligned} \frac{d(\phi_{22})}{da} &= O\left(\frac{\partial}{\partial a} \left[\delta \frac{\partial D_2}{\partial y} B_4 \mu^{2n} + \delta \frac{\partial D_2}{\partial y} D_4 \beta_5 \mu^{2n} \right]\right) \\ &+ O\left(\frac{\partial}{\partial y} \left[\delta \frac{\partial D_2}{\partial a} B_4 \mu^{2n} + \delta \frac{\partial D_2}{\partial a} D_4 \beta_5 \mu^{2n} \right]\right) + O((\lambda\mu^4)^n). \end{aligned}$$

Observe that, by (4.3) and (4.4), we have $\partial B_4/\partial y, \partial D_4/\partial y, \partial \beta_5/\partial y = O((\lambda\mu)^n)$. Using Lemma 3.35, Lemma 3.21 and Lemma 4.2 we reduce to

$$\frac{d(\phi_{22})}{da} = O\left(\frac{\partial}{\partial a} \left[\delta \frac{\partial D_2}{\partial y} B_4 \mu^{2n} + \delta \frac{\partial D_2}{\partial y} D_4 \beta_5 \mu^{2n} \right]\right) + O(n\mu^{3n}).$$

The lemma follows by applying again Lemma 3.35, Lemma 3.21 and Lemma 4.2. \square

The following lemma refers to the univalent function φ introduced in (3.57).

Lemma 4.6. *Let $(t, a) \in \mathcal{H}_n$. Then*

$$\frac{d(D\varphi(0))}{d\beta} = O(1).$$

Proof. Let $(t, a) \in \mathcal{H}_n$ and observe that the map $\tilde{F}_{t,a}$ takes the horizontal segment $y = c_{t,a}$ to a curve which is tangent to the vertical segment $x = v_{t,a}$. The tangency occurs in the point $(v_{t,a}, c_{t,a})$. The curve near the tangency point is given by

$$x = \tilde{Q}y^2 + O(y^3),$$

with coordinates centered at the tangency point and $\tilde{Q} \neq 0$. Recall that $\tilde{F}_{t,a}$ restricted to $y = c_{t,a}$ is given by $q(x) = \varphi(x - c_{t,a})^2$, see (3.57). We describe now this map with coordinates in a domain centered around $c_{t,a}$ and in the image around $v_{t,a}$. Then,

$$\tilde{Q}x^2 + O(x^3) = \varphi(x^2) = D\varphi(0)x^2 + O(x^4),$$

where we used (3.60). Hence,

$$D\varphi(0) = \tilde{Q}. \quad (4.7)$$

In order to determine \tilde{Q} , consider the point $\underline{x} = (c_x, c_y + \Delta y)$ where (c_x, c_y) is the critical point of $F_{t,a}$ and Δy is small. Then,

$$\sigma_{t,a}(\underline{x}) = (c_{t,a} + \Delta s, c_x),$$

and

$$\sigma_{t,a}(F_{t,a}^{n+N}(\underline{x})) = (v_{t,a} + X(\Delta y), c_{t,a} + \Delta s),$$

where $(v_{t,a} + X(\Delta y), c_{t,a} + \Delta s)$ is a point on the graph of $\tilde{F}_{t,a}$ restricted to the line $y = c_{t,a}$. As a consequence,

$$\begin{aligned} \tilde{Q}(\Delta s)^2 + O((\Delta s)^3) = X(\Delta y) &= \frac{\partial X}{\partial y}(c)\Delta y + \frac{\partial^2 X}{\partial y^2}(c)(\Delta y)^2 + O((\Delta y)^3) \\ &= \Phi_2(c)\Delta y + \frac{\partial \Phi_2}{\partial y}(c)(\Delta y)^2 + O((\Delta y)^3) \\ &= \phi_{22}(\Delta y)^2 + O((\Delta y)^3), \end{aligned}$$

where we used that c is the critical point, Φ_2 is the second component of Φ and $\phi_{22} = \partial \Phi_2 / \partial y$, see (3.28). From this and (4.7) we get

$$D\varphi(0) = \phi_{22} \left(\frac{dy}{ds} \right)^2. \quad (4.8)$$

We are now ready to estimate the a -derivative of $D\varphi(0)$. Observe that, using the notation from the previous section, we have $(c_{t,a} + \Delta s) = (h_{t,a}^{-1}(\underline{x}_3))_x$ and by (3.23),

$$\frac{ds}{dy} = \left(1 + \frac{\alpha_3}{\mu^n} \right) (B_2 \delta \mu^n + A_2 \beta \lambda^n) + \beta_3 (D_2 \delta \mu^n + C_2 \beta \lambda^n).$$

It follows, using Lemma 3.26, that

$$\frac{dy}{ds} = \frac{1}{B_2 \delta \mu^n} + O\left(\frac{1}{\mu^{2n}}\right). \quad (4.9)$$

Using the above expression for ds/dy and Lemma 3.35,

$$\frac{\partial}{\partial a} \left(\frac{ds}{dy} \right) = O(n\mu^n). \quad (4.10)$$

Using

$$0 = \frac{\partial}{\partial a} \left(\frac{ds}{dy} \frac{dy}{ds} \right) = \frac{dy}{ds} \frac{\partial}{\partial a} \left(\frac{ds}{dy} \right) + \frac{ds}{dy} \frac{\partial}{\partial a} \left(\frac{dy}{ds} \right),$$

(4.9) and (4.10) we have

$$\frac{\partial}{\partial a} \left(\frac{dy}{ds} \right) = O\left(\frac{n}{\mu^n}\right). \quad (4.11)$$

By taking the a -derivative of (4.8) and using Lemma 4.5, (3.33), (4.9), (4.11), the fact that $da/d\beta = 1/\mu^{2n}(t, sa_n(t))$ and $\mu^{2n}(t, a)/\mu^{2n}(t, sa_n(t)) = 1 + O(n/\mu^n(t, sa_n(t)))$, the lemma follows. \square

Lemma 4.12. For all $\underline{x} \in \sigma^{-1} \circ Z(\text{Dom}(HF_\beta))$,

$$\begin{aligned} |\underline{x} - c| &= O(1/\mu^n), \\ D_2(\underline{x}) &= O(1/\mu^n), \\ D_4(\underline{x}) &= O(1/\mu^n), \\ B_4(\underline{x}) &= B_4(c) + O(1/\mu^n). \end{aligned}$$

Proof. Take $\underline{x}_0 \in \text{Dom}(HF_\beta)$ and compare its image with the image of $(0, 0)$. Let $Z(\underline{x}_0) = \tilde{\underline{x}}$ and $Z(0, 0) = \tilde{\underline{x}}^{(0)}$. Because Z contracts distances by $O(1/\mu^n)$ we get

$$|\tilde{\underline{x}} - \tilde{\underline{x}}^{(0)}|, \left| \tilde{F}(\tilde{\underline{x}}) - \tilde{F}(\tilde{\underline{x}}^{(0)}) \right|, \left| \tilde{F}^2(\tilde{\underline{x}}) - \tilde{F}^2(\tilde{\underline{x}}^{(0)}) \right| = O\left(\frac{1}{\mu^n}\right). \quad (4.13)$$

Let $\underline{x} = (x, y) = \sigma^{-1}(\tilde{\underline{x}})$, $\underline{x}^{(0)} = \sigma^{-1}(\tilde{\underline{x}}^{(0)}) = (c_x, c_y)$. Because $\underline{x} \in HB$, we have that y is proportional to $1/\mu^n$. Hence, by (4.13) and the definition of σ , the first estimate follows. Let $\underline{x}_2 = (x_2, y_2) = F^n(\underline{x})$, $\underline{x}_2^{(0)} = (x_2^{(0)}, y_2^{(0)}) = F^n(\underline{x}^{(0)})$, $\underline{x}_3 = (x_3, y_3) = F^N(\underline{x}_2)$, $\underline{x}_3^{(0)} = F^N(\underline{x}_2^{(0)})$, $\underline{x}_4 = (x_4, y_4) = F^n(\underline{x}_3)$ and $\underline{x}_5 = (x_5, y_5) = F^N(\underline{x}_4)$. By applying F^n to \underline{x} , \underline{x}_3 and $\underline{x}^{(0)}$, (4.13) and Lemma 3.1 imply that

$$\left| x_2 - x_2^{(0)} \right|, \left| x_4 - x_2^{(0)} \right| = O(\lambda^n). \quad (4.14)$$

Because of (4.13) we have

$$|x_3 - c_x|, |x_5 - c_x| = O\left(\frac{1}{\mu^n}\right). \quad (4.15)$$

By (4.14), (4.15) and by taking the preimages under F^N of the points \underline{x}_3 , \underline{x}_5 and $\underline{x}_3^{(0)}$ we obtain

$$\left| y_2 - y_2^{(0)} \right|, \left| y_4 - y_2^{(0)} \right| = O\left(\frac{1}{\mu^n}\right). \quad (4.16)$$

From (4.14) and (4.16) we get

$$\left| \underline{x}_2 - \underline{x}_2^{(0)} \right|, \left| \underline{x}_4 - \underline{x}_2^{(0)} \right| = O\left(\frac{1}{\mu^n}\right).$$

Observe that the image of the points \underline{x}_2 and \underline{x}_4 under the linearization h coincides with the points $\underline{x}_2, \underline{x}_4$ as introduced before in the real domain. Because h is a diffeomorphism, the distances are also of the order $1/\mu^n$. Finally, since $D_2(\underline{x}) = D(\underline{x}_2)$, $D_4(\underline{x}) = D(\underline{x}_4)$, the fact that $D_2(c) = O(\lambda^n/\mu^n)$, see Lemma 3.26, and $B_4(\underline{x}) = B(\underline{x}_4)$ the lemma follows. \square

Lemma 4.17. *For all $\underline{x} \in \sigma^{-1} \circ Z(\text{Dom}(HF_\beta))$,*

$$\frac{\partial x_5}{\partial a}(\underline{x}) = B_4(\underline{x})\mu^n + O(n).$$

Proof. Recall that $q_3 = q_3(t, 0)$ is the point where $D(q_3, t, 0) = 0$ and that $F_{t,0}^N(q_3) = q_1$ where q_1 is the tangency at $a = 0$. Because $\partial D(q_3, t, 0)/\partial y \neq 0$, there is a curve through q_3 transversal to the y -axis where $D(\underline{x}, t, 0) = 0$ for all points \underline{x} in the curve. The transversality implies that there is a curve $D(\underline{x}, t, a) = 0$, transversally intersecting the y -axis in the point $q_3(t, a)$ at the distance of $O(1/\mu^n)$, because $(t, a) \in \mathcal{H}_n$. From Lemma 4.12 we know that $D_2(\underline{x}) = O(1/\mu^n)$. This implies that the distance from \underline{x}_2 to this curve is of the order $O(1/\mu^n)$. Moreover $x_2 = O(\lambda^n)$. Hence,

$$\text{dist}(\underline{x}_2, q_3) = O\left(\frac{1}{\mu^n}\right). \quad (4.18)$$

Because $\partial(F_{t,0}^N(q_3))_y/\partial a = 1$, we get

$$\frac{\partial(F_{t,a}^N(\underline{x}_2))_y}{\partial a} = 1 + O\left(\frac{1}{\mu^n}\right).$$

Now (3.39), becomes

$$\frac{\partial y_3}{\partial a}(\underline{x}) = C_2 \frac{\partial x_2}{\partial a} + D_2 \frac{\partial y_2}{\partial a} + 1 + O\left(\frac{1}{\mu^n}\right) = 1 + O\left(\frac{n}{\mu^n}\right),$$

where we also used (3.36), (3.37) and Lemma 4.12. Now (3.41), becomes

$$\frac{\partial y_4}{\partial a}(\underline{x}) = \mu^n + O(n),$$

and (3.42), becomes

$$\frac{\partial x_5}{\partial a}(\underline{x}) = B_4(\underline{x})\mu^n + O(n),$$

where we used (3.40). \square

Proof of Theorem 4.1. Let $\underline{x}_0 = (x_0, y_0) \in \text{Dom}(HF_\beta)$, $\tilde{\underline{x}} = (\tilde{x}, \tilde{y}) = Z(\underline{x}_0)$ and $\underline{x} = (x, y) = \sigma^{-1}(\tilde{\underline{x}}) \in \sigma^{-1} \circ Z(\text{Dom}(HF_\beta))$. Because $\tilde{\underline{x}} = (c_{t,a}, c_{t,a}) + \underline{x}_0/D\varphi(0)$, using Proposition 3.34, the fact that $D\varphi(0)$ is proportional to μ^n , Lemma 4.6 and the fact that $da/d\beta = O(1/\mu^{2n})$ we have

$$\frac{d\tilde{\underline{x}}}{d\beta} = O\left(\frac{n}{\mu^{2n}}\right), \quad (4.19)$$

and because $x = \tilde{y}$,

$$\frac{dx}{d\beta} = O\left(\frac{n}{\mu^{2n}}\right). \quad (4.20)$$

The estimate for $dy/d\beta$ needs some preparation. By (3.19), (4.20), (3.23), (3.38) and the fact that $da/d\beta = O(1/\mu^{2n})$ we have

$$\frac{dx_3}{d\beta} = \frac{\partial x_3}{\partial x} \frac{dx}{d\beta} + \frac{\partial x_3}{\partial y} \frac{dy}{d\beta} + \frac{\partial x_3}{\partial a} \frac{da}{d\beta} = (B_2\delta\mu^n + O(\lambda^n)) \frac{dy}{d\beta} + O\left(\frac{n}{\mu^{2n}}\right), \quad (4.21)$$

where, by Remark 3.17, B_2 and δ are uniformly away from zero. Similarly, by (3.19), (4.20), (3.23), (3.39) and Lemma 4.12 we have

$$\frac{dy_3}{d\beta} = O(1) \frac{dy}{d\beta} + O\left(\frac{1}{\mu^{2n}}\right). \quad (4.22)$$

Observe now that $\tilde{x} = (h^{-1}(\underline{x}_3))_x$ which implies, using (4.21) and (4.22),

$$\begin{aligned} \frac{d\tilde{x}}{d\beta} &= \left(1 + \frac{\alpha_3}{\mu^n}\right) \left[(B_2\delta\mu^n + O(\lambda^n)) \frac{dy}{d\beta} + O\left(\frac{n}{\mu^{2n}}\right) \right] + \beta_3 O(1) \frac{dy}{d\beta} + O\left(\frac{1}{\mu^{2n}}\right) \\ &= (B_2\delta\mu^n + O(1)) \frac{dy}{d\beta} + O\left(\frac{n}{\mu^{2n}}\right). \end{aligned}$$

By (4.19), we get

$$\frac{dy}{d\beta} = O\left(\frac{n}{\mu^{3n}}\right). \quad (4.23)$$

Observe now that, by (3.20), Lemma 4.12, (4.20), (3.25), again Lemma 4.12, (4.23) and Lemma 4.17,

$$\frac{dx_5}{d\beta} = \frac{\partial x_5}{\partial x} \frac{dx}{d\beta} + \frac{\partial x_5}{\partial y} \frac{dy}{d\beta} + \frac{\partial x_5}{\partial a} \frac{da}{d\beta} = \frac{B_4(\underline{x})}{\mu^n} + O\left(\frac{n}{\mu^{2n}}\right). \quad (4.24)$$

Similarly, by (3.20), Lemma 4.12, (4.20), (3.25), again Lemma 4.12, (4.23) and (3.43)

$$\frac{dy_5}{d\beta} = O\left(\frac{n}{\mu^{2n}}\right). \quad (4.25)$$

Because $(\tilde{F}(\tilde{x}))_x = (h^{-1}(\underline{x}_5))_x$, using (4.24) and (4.25)

$$\begin{aligned} \frac{d(\tilde{F}(\tilde{x}))_x}{d\beta} &= \left(1 + \frac{\alpha_5}{\mu^n}\right) \left[\frac{B_4(\underline{x})}{\mu^n} + O\left(\frac{n}{\mu^{2n}}\right) \right] + \beta_5 O\left(\frac{n}{\mu^{2n}}\right) + O\left(\frac{\partial h^{-1}}{\partial a}\right) \frac{1}{\mu^{2n}} \\ &= \frac{B_4(\underline{x})}{\mu^n} + O\left(\frac{n}{\mu^{2n}}\right). \end{aligned} \quad (4.26)$$

Finally observe that

$$f_\beta(\underline{x}_0) = \alpha^{-1} \left((\tilde{F}(\tilde{x}))_x \right) = \left((\tilde{F}(\tilde{x}))_x - c_{t,a} \right) D\varphi(0).$$

As a consequence,

$$\begin{aligned}\frac{df_\beta(\underline{x}_0)}{d\beta} &= D\varphi(0)\frac{d\left(\tilde{F}(\tilde{\underline{x}})\right)_x}{d\beta} - D\varphi(0)\frac{\partial c_{t,a}}{\partial a}\frac{da}{d\beta} + \left(\left(\tilde{F}(\tilde{\underline{x}})\right)_x - c_{t,a}\right)\frac{dD\varphi(0)}{d\beta} \\ &= B_4(\underline{x})\frac{D\varphi(0)}{\mu^n} + O\left(\frac{n}{\mu^n}\right),\end{aligned}$$

where we used (4.26), the fact that $D\varphi(0)$ is proportional to μ^n , Proposition 3.34, Lemma 4.6 and the fact that $\left(\tilde{F}(\tilde{\underline{x}})\right)_x \in \alpha(U_{t,a})$ which has diameter proportional to $1/\mu^n$. In conclusion, using Lemma 4.12,

$$\frac{df_\beta}{d\beta} = B_4(c)\frac{D\varphi(0)}{\mu^n} + O\left(\frac{n}{\mu^n}\right),$$

where $B_4(c)D\varphi(0)/\mu^n$ is uniformly bounded away from zero. \square

4.2 Monotonicity in the t direction

The aim of this subsection is to study the dependence of the normalized map on the t -parameter. In order to achieve that, we start by calculating the movement of the critical point while changing the parameter t .

Proposition 4.27. *For every $(t, a) \in \mathcal{H}_n$,*

$$\begin{aligned}\frac{\partial c_{t,a}}{\partial t} &= \frac{\partial c_x}{\partial t} = \frac{\partial v_x}{\partial t} = O\left(\frac{n}{\mu^n(t, a)}\right), \\ \frac{\partial v_{t,a}}{\partial t} &= B_4 n \mu^{n-1} \frac{\partial \mu}{\partial t} y_3 + O\left(\frac{n}{\mu^n(t, a)}\right).\end{aligned}$$

The proof of this proposition needs some preparation. Fix $(t, a) \in \mathcal{H}_n$. Let $(x, y) \in HB(t, a) \cap \mathbb{R}^2$ and $\underline{x} = (x, y)$, $\underline{x}_1, \underline{x}_2, \underline{x}_3, \underline{x}_4, \underline{x}_5$ as introduced before. Observe that all coefficients introduced, $\alpha, \beta, \dots, A_2, B_2, \dots, \alpha_3, \beta_3, \dots, A_4, B_4, \dots, \alpha_5, \dots$ are all functions of the point $(x, y) \in HB(t, a) \cap \mathbb{R}^2$ and t . We will vary the parameter t .

Lemma 4.28. *The coordinate change $h_{t,a}$ restricted to $\sigma_{t,a}^{-1} \circ Z_{t,a}(\text{Dom}(HF_{t,a}))$ satisfies*

$$\frac{\partial (h_{t,a})_y}{\partial t} = O\left(\frac{1}{\mu^{2n}(t, a)}\right).$$

Proof. As in the proof of Lemma 3.14, write the Taylor expansion of $(h_{t,a})_y$ centered around the point $(2, 0)$. Then

$$(h_{t,a})_y = y(1 + x\varphi_x + y\varphi_y).$$

It follows that

$$\frac{\partial (h_{t,a})_y}{\partial t} = y \left(x \frac{\partial \varphi_x}{\partial t} + y \frac{\partial \varphi_y}{\partial t} \right) = O\left(\frac{1}{\mu^{2n}}\right)$$

where we used the first equality in Lemma 4.12 with $c_x = 2$. \square

Lemma 4.29. For all $\underline{x} = (x, y) \in F_{t,a}^n \circ \sigma_{t,a}^{-1} \circ Z_{t,a} (\text{Dom}(HF_{t,a}))$,

$$\begin{aligned}\frac{\partial D(\underline{x})}{\partial t} &= O\left(\frac{1}{\mu^n(t, a)}\right), \\ \frac{\partial (F_{t,a}^N(\underline{x}))_x}{\partial t} &= O\left(\frac{1}{\mu^n(t, a)}\right), \\ \frac{\partial (F_{t,a}^N(\underline{x}))_y}{\partial t} &= O\left(\frac{1}{\mu^{2n}(t, a)}\right).\end{aligned}$$

Proof. From the first statement of Lemma 4.12 and the fact that $a = O(1/\mu^n)$, we have

$$|q_3(t) - \underline{x}| = O\left(\frac{1}{\mu^n}\right), \quad (4.30)$$

and by the hypothesis $q_3(t, 0) = (0, 1)$

$$|1 - y| = O\left(\frac{1}{\mu^n}\right). \quad (4.31)$$

Moreover, by Lemma 3.14,

$$|x| = O((\lambda\mu)^n). \quad (4.32)$$

Observe that $D(q_3(t, 0), t, 0) = 0$ when $a = 0$. Hence, for every t , $\partial D(q_3(t, 0), t, 0)/\partial t = 0$. The first statement follows by applying (4.30). Write $(F_{t,a}^N(\underline{x}))_x$ in coordinates centered in the domain around q_3 and in the image around q_1 . One gets

$$(F_{t,a}^N(\underline{x}))_x = Ax + By + aH,$$

where A, B, H are \mathcal{C}^2 functions. Hence,

$$\frac{\partial (F_{t,a}^N(\underline{x}))_x}{\partial t} = O(|x| + |y| + |a|).$$

The second statement follows. Similarly, write $(F_{t,a}^N(\underline{x}))_y$ in coordinates centered in the domain around q_3 and in the image around q_1 . One gets

$$(F_{t,a}^N(\underline{x}))_y = Cx + Q_{11}x^2 + Q_{12}xy + Q_{22}y^2 + a(1 + V_1x + V_2y)$$

where $C, Q_{11}, Q_{12}, Q_{22}, V_1, V_2$ are \mathcal{C}^2 functions. Hence

$$\frac{\partial (F_{t,a}^N(\underline{x}))_y}{\partial t} = O(|x| + |xy| + |y^2| + |ax| + |ay|).$$

The last statement follows by using (4.30), (4.32) and the fact that $\lambda\mu^3 < 1$. \square

Lemma 4.33. The partial derivatives at the critical point c of $F_{t,a}$ satisfy the following.

- $\partial\alpha/\partial t, \partial\beta/\partial t, \partial\gamma/\partial t, \partial\delta/\partial t = O(1)$,
- $\partial A_2/\partial t, \partial B_2/\partial t, \partial C_2/\partial t, \partial D_2/\partial t = O(n)$,
- $\partial\alpha_3/\partial t, \partial\beta_3/\partial t, \partial\gamma_3/\partial t, \partial\delta_3/\partial t = O(n)$,

- $\partial A_4/\partial t, \partial B_4/\partial t, \partial C_4/\partial t, \partial D_4/\partial t = O(n)$,
- $\partial \alpha_5/\partial t, \partial \beta_5/\partial t, \partial \gamma_5/\partial t, \partial \delta_5/\partial t = O(n)$.

Proof. The first set of estimates follows by the smoothness of the family of maps $h_{t,a}$. For the second set of estimates, observe that, because $h_{t,a}(x, 0) = (x, 0)$ and $y = O(1/\mu^n)$, we have

$$\frac{\partial x_1}{\partial t} = O\left(\frac{1}{\mu^n}\right),$$

and by Lemma 4.28,

$$\frac{\partial y_1}{\partial t} = O\left(\frac{1}{\mu^{2n}}\right).$$

Recall that $\underline{x}_2 = \hat{F}_{t,a}^n(\underline{x}_1, t, a)$. It follows that,

$$\frac{\partial x_2}{\partial t} = n\lambda^{n-1}\frac{\partial \lambda}{\partial t}x_1 + \lambda^n\frac{\partial x_1}{\partial t} = O(n\lambda^n), \quad (4.34)$$

and

$$\frac{\partial y_2}{\partial t} = n\mu^{n-1}\frac{\partial \mu}{\partial t}y_1 + \mu^n\frac{\partial y_1}{\partial t} = n\mu^{n-1}\frac{\partial \mu}{\partial t}y_1 + O\left(\frac{1}{\mu^n}\right). \quad (4.35)$$

Observe that, by our initial hypotheses $\partial\mu/\partial t \neq 0$ and that y_1 is proportional to $1/\mu^n$. As a consequence, $\partial y_2/\partial t$ is proportional to n . Moreover

$$\frac{\partial A_2}{\partial t} = \frac{\partial A}{\partial x}\frac{\partial x_2}{\partial t} + \frac{\partial A}{\partial y}\frac{\partial y_2}{\partial t} + \frac{\partial A}{\partial t} = O(n).$$

Similarly the other estimates in the second statement follows. Observe that, using Lemma 4.29,

$$\frac{\partial D_2}{\partial t} = \frac{\partial D}{\partial y}n\mu^{n-1}\frac{\partial \mu}{\partial t}y_1 + O\left(\frac{1}{\mu^n}\right), \quad (4.36)$$

and $\partial D/\partial y \neq 0$. Recall that $\underline{x}_3 = \hat{F}_{t,a}^N(\underline{x}_2, t, a)$. By Lemma 4.29,

$$\frac{\partial x_3}{\partial t} = A_2\frac{\partial x_2}{\partial t} + B_2\frac{\partial y_2}{\partial t} + O\left(\frac{1}{\mu^n}\right) = B_2n\mu^{n-1}\frac{\partial \mu}{\partial t}y_1 + O\left(\frac{1}{\mu^n}\right), \quad (4.37)$$

and

$$\begin{aligned} \frac{\partial y_3}{\partial t} &= C_2\frac{\partial x_2}{\partial t} + D_2\frac{\partial y_2}{\partial t} + O\left(\frac{1}{\mu^{2n}}\right) \\ &= C_2\frac{\partial x_2}{\partial t} + D_2n\mu^{n-1}\frac{\partial \mu}{\partial t}y_1 + D_2O\left(\frac{1}{\mu^n}\right) + O\left(\frac{1}{\mu^{2n}}\right) \end{aligned} \quad (4.38)$$

$$= O\left(\frac{1}{\mu^{2n}}\right), \quad (4.39)$$

where we used (4.34), (4.35) and that $D_2 = O(\lambda^n/\mu^n)$, see Lemma 3.26. The third set of estimates follow. Recall now that $\underline{x}_4 = \hat{F}_{t,a}^n(\underline{x}_3, t, a)$. Hence,

$$\frac{\partial x_4}{\partial t} = n\lambda^{n-1}\frac{\partial \lambda}{\partial t}x_3 + \lambda^n\frac{\partial x_3}{\partial t} = O(n\lambda^n), \quad (4.40)$$

and

$$\frac{\partial y_4}{\partial t} = n\mu^{n-1}\frac{\partial\mu}{\partial t}y_3 + \mu^n\frac{\partial y_3}{\partial t} = n\mu^{n-1}\frac{\partial\mu}{\partial t}y_3 + O\left(\frac{1}{\mu^n}\right), \quad (4.41)$$

where we used (4.37), (4.39). Observe that, by our initial hypotheses $\partial\mu/\partial t \neq 0$ and that y_3 is proportional to $1/\mu^n$. As a consequence $\partial y_4/\partial t$ is proportional to n . Hence,

$$\frac{\partial A_4}{\partial t} = \frac{\partial A}{\partial x}\frac{\partial x_4}{\partial t} + \frac{\partial A}{\partial y}\frac{\partial y_4}{\partial t} + \frac{\partial A}{\partial t} = O(n).$$

Similarly the other estimates in the fourth statement follows. Finally, by Lemma 4.29,

$$\frac{\partial x_5}{\partial t} = A_4\frac{\partial x_4}{\partial t} + B_4\frac{\partial y_4}{\partial t} + O\left(\frac{1}{\mu^n}\right) = B_4n\mu^{n-1}\frac{\partial\mu}{\partial t}y_3 + O\left(\frac{1}{\mu^n}\right), \quad (4.42)$$

and

$$\frac{\partial y_5}{\partial t} = C_4\frac{\partial x_4}{\partial t} + D_4\frac{\partial y_4}{\partial t} + O\left(\frac{1}{\mu^{2n}}\right) = O\left(\frac{n}{\mu^n}\right), \quad (4.43)$$

where we used (4.40), (4.41) and Lemma 4.12. As before the last set follows. \square

Proof of Proposition 4.27. Observe that

$$\frac{\partial c}{\partial t} = (D\Phi)^{-1}\frac{\partial\Phi}{\partial t}.$$

Let $\Phi = (\Phi_1, \Phi_2)$. We start by calculating $\partial\Phi_1/\partial t$. Observe that $\Phi_1 = (h_{t,a}^{-1}(\underline{x}_3))_x$. As a consequence

$$\frac{\partial\Phi_1}{\partial t} = \left(1 + \frac{\alpha_3}{\mu^n}\right)\frac{\partial x_3}{\partial t} + \beta_3\frac{\partial y_3}{\partial t} + \frac{\partial(h_{t,a}^{-1})_x}{\partial t} = B_2n\mu^{n-1}\frac{\partial\mu}{\partial t}y_1 + O\left(\frac{n}{\mu^n}\right). \quad (4.44)$$

where we used (4.37) and (4.39). In order to calculate $\partial\Phi_2/\partial t$ we take the t -derivatives of (3.31). Using Lemma 4.33, the partial derivatives of all terms in (3.31) with a factor λ^n will give a contribution of $O(n(\lambda\mu)^n)$. We get

$$\begin{aligned} \frac{\partial\Phi_2}{\partial t} &= \frac{\partial}{\partial t} [\delta D_2 B_4 \mu^{2n} + \delta D_2 D_4 \beta_5 \mu^{2n} + \delta D_2 B_4 \alpha_5 \mu^n] \\ &+ O(n(\lambda\mu)^n). \end{aligned}$$

Using Lemma 4.33, (4.36), the fact that $D_2 = O(\lambda^n/\mu^n)$, see Lemma 3.26 and Lemma 4.12, we get

$$\frac{\partial\Phi_2}{\partial t} = \delta B_4 \frac{\partial D}{\partial y} n\mu^{n-1} \frac{\partial\mu}{\partial t} y_1 \mu^{2n} + O(n\mu^n). \quad (4.45)$$

By (3.29), (3.30), (3.32), (3.33), using Lemma 4.12 and (3.27) we have

$$D\Phi = \begin{pmatrix} -1 + O(\gamma) & \delta B_2 \mu^n + O(1) \\ \delta B_4 \frac{\partial D}{\partial x} \mu^{2n} + O(\mu^n) & \delta^2 B_4 \frac{\partial D}{\partial y} \mu^{3n} + O(\mu^{2n}) \end{pmatrix},$$

and

$$\det(D\Phi) = -\delta^2 B_4 \frac{\partial D}{\partial y} \mu^{3n} \left[1 + \frac{B_2 \partial D_2 / \partial x}{\partial D / \partial y} - O(\gamma)\right] = -\delta^2 B_4 \frac{\partial D}{\partial y} \mu^{3n} [1 + \chi],$$

where χ is close to zero, see Lemma 4.12 and Remark 3.16. Moreover

$$\begin{aligned} ((D\Phi)^{-1})_{11} &= - \left[\frac{1 + O(1/\mu^n)}{1 + \chi} \right] \\ ((D\Phi)^{-1})_{12} &= \frac{B_2}{\delta B_4 \partial D / \partial y} \frac{1}{\mu^{2n}} \left[\frac{1 + O(1/\mu^n)}{1 + \chi} \right]. \end{aligned}$$

As a consequence, using (4.44) and (4.45) a cancellation happens and leads to

$$\frac{\partial c_x}{\partial t} = \frac{\partial c_{t,a}}{\partial t} = \frac{\partial v_x}{\partial t} = ((D\Phi)^{-1})_{11} \frac{\partial \Phi_1}{\partial t} + ((D\Phi)^{-1})_{12} \frac{\partial \Phi_2}{\partial t} = O\left(\frac{n}{\mu^n}\right).$$

The first equation follows. Observe that $v_{t,a} = (h_{t,a}^{-1}(x_5))_x$. Hence

$$\frac{\partial v_{t,a}}{\partial t} = \left(1 + \frac{\alpha_5}{\mu^n}\right) \frac{\partial x_5}{\partial t} + \beta_5 \frac{\partial y_5}{\partial t} + O\left(\frac{1}{\mu^n}\right) = B_4 n \mu^{n-1} \frac{\partial \mu}{\partial t} y_3 + O\left(\frac{n}{\mu^n}\right),$$

where we used (4.42) and (4.43). □

We are now ready to state the main theorem on the monotonicity of the normalized family in the t direction.

Theorem 4.46. *Fix $(t, a) \in \mathcal{H}_n$. There exists a real-analytic function $M : \mathcal{H}_n \rightarrow \mathbb{R}$ which is bounded away from zero,*

$$M(t, a) = \frac{D\varphi(0)B_4(c)y_3}{\mu} \frac{\partial \mu}{\partial t},$$

such that

$$\frac{\partial f_{t,a}}{\partial t} = M(t, a)n\mu^n(t, a) + O(n).$$

The proof of Theorem 4.46 requires the following lemmas. We use the notation of the previous section.

Lemma 4.47. *The mixed partial derivatives at the critical point c of $F_{t,a}$ satisfy the following.*

- $\partial^2 \alpha / \partial t \partial y, \partial^2 \beta / \partial t \partial y, \partial^2 \gamma / \partial t \partial y, \partial^2 \delta / \partial t \partial y = O(1),$
- $\partial^2 A_2 / \partial t \partial y, \partial^2 B_2 / \partial t \partial y, \partial^2 C_2 / \partial t \partial y, \partial^2 D_2 / \partial t \partial y = O(n\mu^n(t, a)),$
- $\partial^2 A_4 / \partial t \partial y, \partial^2 B_4 / \partial t \partial y, \partial^2 C_4 / \partial t \partial y, \partial^2 D_4 / \partial t \partial y = O(n\mu^{2n}(t, a)),$
- $\partial^2 \alpha_5 / \partial a \partial y, \partial^2 \beta_5 / \partial a \partial y, \partial^2 \gamma_5 / \partial a \partial y, \partial^2 \delta_5 / \partial a \partial y = O(n\mu^{2n}(t, a)).$

Proof. The first set of estimates follows by the smoothness of the family of maps $h_{t,a}$. For the second set of estimates, observe that, by (3.22) and Lemma 4.33

$$\frac{\partial^2 x_2}{\partial t \partial y} = O(n\mu^n).$$

As a consequence,

$$\begin{aligned}
\frac{\partial^2 A_2}{\partial t \partial y} &= \frac{\partial}{\partial t} \left[\frac{\partial A}{\partial x} \frac{\partial x_2}{\partial y} + \frac{\partial A}{\partial y} \frac{\partial y_2}{\partial y} \right] \\
&= \frac{\partial x_2}{\partial y} \left[\frac{\partial^2 A}{\partial x^2} \frac{\partial x_2}{\partial t} + \frac{\partial^2 A}{\partial y \partial x} \frac{\partial y_2}{\partial t} + \frac{\partial^2 A}{\partial t \partial x} \right] + \frac{\partial A}{\partial x} \frac{\partial^2 x_2}{\partial t \partial y} \\
&+ \frac{\partial y_2}{\partial y} \left[\frac{\partial^2 A}{\partial x \partial y} \frac{\partial x_2}{\partial t} + \frac{\partial^2 A}{\partial y^2} \frac{\partial y_2}{\partial t} + \frac{\partial^2 A}{\partial t \partial y} \right] + \frac{\partial A}{\partial y} \frac{\partial^2 y_2}{\partial t \partial y} \\
&= O(n\mu^n),
\end{aligned}$$

where we also used (3.22), (4.34) and (4.35). Similarly the other estimates in the second statement follows. For the third set of estimates, observe that, by (3.24) and Lemma 4.33

$$\frac{\partial^2 \underline{x}_4}{\partial t \partial y} = O(n\mu^{2n}),$$

and by (3.24), the fact that $D_2 = O((\lambda/\mu)^n)$, see Lemma 3.26, we get

$$\frac{\partial \underline{x}_4}{\partial y} = O((\lambda\mu)^n).$$

As a consequence,

$$\begin{aligned}
\frac{\partial^2 A_4}{\partial t \partial y} &= \frac{\partial x_4}{\partial y} \left[\frac{\partial^2 A}{\partial x^2} \frac{\partial x_4}{\partial t} + \frac{\partial^2 A}{\partial y \partial x} \frac{\partial y_4}{\partial t} + \frac{\partial^2 A}{\partial t \partial x} \right] + \frac{\partial A}{\partial x} \frac{\partial^2 x_4}{\partial t \partial y} \\
&+ \frac{\partial y_4}{\partial y} \left[\frac{\partial^2 A}{\partial x \partial y} \frac{\partial x_4}{\partial t} + \frac{\partial^2 A}{\partial y^2} \frac{\partial y_4}{\partial t} + \frac{\partial^2 A}{\partial t \partial y} \right] + \frac{\partial A}{\partial y} \frac{\partial^2 y_4}{\partial t \partial y} \\
&= O(n\mu^{2n}),
\end{aligned}$$

where we also used (4.40) and (4.41). Similarly the other estimates in the third statement follows. Finally, by (3.25), Lemma 3.26 and Lemma 4.33

$$\frac{\partial^2 \underline{x}_5}{\partial t \partial y} = O(n\mu^{2n}),$$

and by (3.25), the fact that $D_2 = O((\lambda/\mu)^n)$, see Lemma 3.26, we get

$$\frac{\partial \underline{x}_5}{\partial y} = O((\lambda\mu)^n).$$

As a consequence,

$$\frac{\partial^2 \alpha_5}{\partial t \partial y} = O(n\mu^{2n}),$$

where we also used (4.42) and (4.43). Similarly the other estimates in the last statement follows. \square

The following lemma refers to the function Φ . Recall that Φ_2 is the second component of Φ and $\phi_{22} = \partial\Phi_2/\partial y$, see (3.28).

Lemma 4.48. *Let $(t, a) \in \mathcal{H}_n$. Then*

$$\frac{d(\phi_{22})}{dt} = O(n\mu^{3n}(t, a)).$$

Proof. We calculate $d(\phi_{22})/dt = \partial^2 \Phi_2 / \partial t \partial y$ using (3.31). Observe that, by Lemma 4.33, Lemma 3.21 and Lemma 4.47, the terms in (3.31) containing a λ^n will contribute to the estimate of the t -derivative of ϕ_{22} with an order of at most $(\lambda\mu^3)^n$. Hence,

$$\frac{d(\phi_{22})}{dt} = \frac{\partial^2}{\partial t \partial y} [\delta D_2 B_4 \mu^{2n} + \delta D_2 D_4 \beta_5 \mu^{2n} + \delta D_2 B_4 \alpha_5 \mu^n] + O(n(\lambda\mu^3)^n).$$

Observe that the mixed partial derivatives of the coefficients of μ^{2n} , μ^n are of the same order. Hence,

$$\frac{d(\phi_{22})}{dt} = O\left(\frac{\partial^2}{\partial t \partial y} [\delta D_2 B_4 \mu^{2n} + \delta D_2 D_4 \beta_5 \mu^{2n}]\right) + O(n(\lambda\mu^3)^n).$$

Because $D_2 = O((\lambda/\mu)^n)$, see Lemma 3.26, Lemma 4.33, Lemma 3.21 and Lemma 4.47 all the terms which do not involve a partial derivative of D_2 , give a contribution of order at most $n(\lambda\mu^3)^n$. Hence,

$$\begin{aligned} \frac{d(\phi_{22})}{dt} &= O\left(\frac{\partial}{\partial t} \left[\delta \frac{\partial D_2}{\partial y} B_4 \mu^{2n} + \delta \frac{\partial D_2}{\partial y} D_4 \beta_5 \mu^{2n} \right]\right) \\ &+ O\left(\frac{\partial}{\partial y} \left[\delta \frac{\partial D_2}{\partial t} B_4 \mu^{2n} + \delta \frac{\partial D_2}{\partial t} D_4 \beta_5 \mu^{2n} \right]\right) + O(n(\lambda\mu^3)^n). \end{aligned}$$

Observe that, by (4.3) and (4.4), we have $\partial B_4 / \partial y, \partial D_4 / \partial y, \partial \beta_5 / \partial y = O((\lambda\mu)^n)$. Using Lemma 4.33, Lemma 3.21 and Lemma 4.47 we reduce to

$$\frac{d(\phi_{22})}{dt} = O\left(\frac{\partial}{\partial t} \left[\delta \frac{\partial D_2}{\partial y} B_4 \mu^{2n} + \delta \frac{\partial D_2}{\partial y} D_4 \beta_5 \mu^{2n} \right]\right) + O(n\mu^{3n}).$$

The lemma follows by applying again Lemma 4.33, Lemma 3.21 and Lemma 4.47. \square

The following lemma refers to the univalent function φ introduced in (3.57).

Lemma 4.49. *Let $(t, a) \in \mathcal{H}_n$. Then*

$$\frac{d(D\varphi(0))}{dt} = O(n\mu^n(t, a)).$$

Proof. Let $(t, a) \in \mathcal{H}_n$. Following the notation as in the proof of Lemma 4.6, by (4.8) we have

$$D\varphi(0) = \phi_{22} \left(\frac{dy}{ds} \right)^2. \quad (4.50)$$

We are now ready to estimate the t -derivative of $D\varphi(0)$. Recall that $(c_{t,a} + \Delta s) = (h_{t,a}^{-1}(\underline{x}_3))_x$ and by (3.23),

$$\frac{ds}{dy} = \left(1 + \frac{\alpha_3}{\mu^n} \right) (\delta B_2 \mu^n + A_2 \beta \lambda^n) + \beta_3 (D_2 \delta \mu^n + C_2 \beta \lambda^n).$$

Using the above expression for ds/dy and Lemma 4.33,

$$\frac{\partial}{\partial t} \left(\frac{ds}{dy} \right) = O(n\mu^n). \quad (4.51)$$

Using

$$0 = \frac{\partial}{\partial t} \left(\frac{ds}{dy} \frac{dy}{ds} \right) = \frac{dy}{ds} \frac{\partial}{\partial t} \left(\frac{ds}{dy} \right) + \frac{ds}{dy} \frac{\partial}{\partial t} \left(\frac{dy}{ds} \right),$$

(4.9) and (4.51) we have

$$\frac{\partial}{\partial t} \left(\frac{dy}{ds} \right) = O\left(\frac{n}{\mu^n}\right). \quad (4.52)$$

By taking the t -derivative of (4.8) and using Lemma 4.48, (3.33), (4.9), and (4.52) the lemma follows. \square

Lemma 4.53. For all $\underline{x} \in \sigma_{t,a}^{-1} \circ Z_{t,a}(\text{Dom}(HF_{t,a}))$,

$$\frac{\partial x_5}{\partial t}(\underline{x}) = B_4(\underline{x})n\mu^{n-1}(t,a) \frac{\partial \mu}{\partial t}(y_3 + \mu^n(t,a)D_2(\underline{x})y_1) + O\left(\frac{1}{\mu^n(t,a)}\right).$$

Proof. By (4.38),

$$\frac{\partial y_3}{\partial t}(\underline{x}) = C_2 \frac{\partial x_2}{\partial t} + D_2 n \mu^{n-1} \frac{\partial \mu}{\partial t} y_1 + D_2 O\left(\frac{1}{\mu^n}\right) + O\left(\frac{1}{\mu^{2n}}\right) = D_2(\underline{x})n\mu^{n-1} \frac{\partial \mu}{\partial t} y_1 + O\left(\frac{1}{\mu^{2n}}\right),$$

where we also used (4.34), and Lemma 4.12. Now (4.41), becomes

$$\frac{\partial y_4}{\partial t}(\underline{x}) = n\mu^{n-1} \frac{\partial \mu}{\partial t}(y_3 + \mu^n D_2(\underline{x})y_1) + O\left(\frac{1}{\mu^n}\right),$$

and (4.42), becomes

$$\frac{\partial x_5}{\partial t}(\underline{x}) = B_4(\underline{x})n\mu^{n-1} \frac{\partial \mu}{\partial t}(y_3 + \mu^n D_2(\underline{x})y_1) + O\left(\frac{1}{\mu^n}\right),$$

where we used (4.40). \square

Proof of Theorem 4.46. Let $\underline{x}_0 = (x_0, y_0) \in \text{Dom}(HF_{t,a})$, $\tilde{\underline{x}} = (\tilde{x}, \tilde{y}) = Z_{t,a}(\underline{x}_0)$ and $\underline{x} = (x, y) = \sigma_{t,a}^{-1}(\tilde{\underline{x}}) \in \sigma_{t,a}^{-1} \circ Z_{t,a}(\text{Dom}(HF_{t,a}))$. Because $\tilde{\underline{x}} = (c_{t,a}, c_{t,a}) + \underline{x}_0/D\varphi(0)$, using Proposition 4.27, the fact that $D\varphi(0)$ is proportional to μ^n and Lemma 4.49 we have

$$\frac{d\tilde{\underline{x}}}{dt} = O\left(\frac{n}{\mu^n}\right), \quad (4.54)$$

and because $x = \tilde{y}$,

$$\frac{dx}{dt} = O\left(\frac{n}{\mu^n}\right). \quad (4.55)$$

The estimate for dy/dt needs some preparation. By (3.19), (4.55), (3.23) and (4.37) we have

$$\begin{aligned} \frac{dx_3}{dt} &= \frac{\partial x_3}{\partial x} \frac{dx}{dt} + \frac{\partial x_3}{\partial y} \frac{dy}{dt} + \frac{\partial x_3}{\partial t} \\ &= (\delta B_2 \mu^n + O(\lambda^n)) \frac{dy}{dt} + B_2 n \mu^{n-1} \frac{\partial \mu}{\partial t} y_1 + O\left(\frac{n}{\mu^n}\right), \end{aligned} \quad (4.56)$$

where B_2 and δ as uniformly away from zero, see Remark 3.17. Similarly, by (3.19), (4.55), (3.23), (4.38) we have

$$\frac{dy_3}{dt} = O(1) \frac{dy}{dt} + O\left(\frac{n}{\mu^{2n}}\right), \quad (4.57)$$

where we also used Lemma 4.12. Observe now that $\tilde{x} = (h^{-1}(\underline{x}_3))_x$ which implies, using (4.56) and (4.57),

$$\begin{aligned} \frac{d\tilde{x}}{dt} &= \left(1 + \frac{\alpha_3}{\mu^n}\right) \left((\delta B_2 \mu^n + O(\lambda^n)) \frac{dy}{dt} + B_2 n \mu^{n-1} \frac{\partial \mu}{\partial t} y_1 + O\left(\frac{n}{\mu^n}\right) \right) + O(1) \frac{dy}{dt} \\ &= (\delta B_2 \mu^n + O(1)) \frac{dy}{dt} + B_2 n \mu^{n-1} \frac{\partial \mu}{\partial t} y_1 + O\left(\frac{n}{\mu^n}\right). \end{aligned}$$

By (4.54), we get

$$\frac{dy}{dt} = -\frac{ny_1}{\delta \mu} \frac{\partial \mu}{\partial t} \left(1 + O\left(\frac{1}{\mu^n}\right)\right). \quad (4.58)$$

Observe now that, by (3.20), Lemma 4.12, (4.55), (3.25), again Lemma 4.12, (4.58) and Lemma 4.53,

$$\begin{aligned} \frac{dx_5}{dt} &= \frac{\partial x_5}{\partial x} \frac{dx}{dt} + \frac{\partial x_5}{\partial y} \frac{dy}{dt} + \frac{\partial x_5}{\partial t} \\ &= B_4 \mu^{2n} \left(\delta D_2 + O\left(\left(\frac{\lambda}{\mu}\right)^n\right) \right) \left(-\frac{ny_1}{\delta \mu} \frac{\partial \mu}{\partial t} \left(1 + O\left(\frac{1}{\mu^n}\right)\right) \right) \\ &\quad + B_4 n \mu^{n-1} \frac{\partial \mu}{\partial t} (y_3 + \mu^n D_2 y_1) + O\left(\frac{n}{\mu^n}\right) \\ &= B_4 n \mu^{n-1} \frac{\partial \mu}{\partial t} y_3 + O\left(\frac{n}{\mu^n}\right). \end{aligned} \quad (4.59)$$

Similarly, by (3.20), Lemma 4.12, (4.55), (3.25), again Lemma 4.12, (4.58) and (4.43)

$$\frac{dy_5}{dt} = O\left(\frac{n}{\mu^n}\right). \quad (4.60)$$

Because $\left(\tilde{F}_{t,a}(\tilde{\underline{x}})\right)_x = (h_{t,a}^{-1}(\underline{x}_5))_x$, using (4.59) and (4.60)

$$\begin{aligned} \frac{d\left(\tilde{F}_{t,a}(\tilde{\underline{x}})\right)_x}{dt} &= \left(1 + \frac{\alpha_5}{\mu^n}\right) \left[B_4 n \mu^{n-1} \frac{\partial \mu}{\partial t} y_3 + O\left(\frac{n}{\mu^n}\right) \right] + O\left(\frac{n}{\mu^n}\right) \\ &= B_4 n \mu^{n-1} \frac{\partial \mu}{\partial t} y_3 + O\left(\frac{n}{\mu^n}\right). \end{aligned} \quad (4.61)$$

Finally observe that

$$f_{t,a}(\underline{x}_0) = \alpha^{-1} \left(\left(\tilde{F}(\tilde{\underline{x}}) \right)_x \right) = \left(\left(\tilde{F}(\tilde{\underline{x}}) \right)_x - c_{t,a} \right) D\varphi(0).$$

As a consequence,

$$\begin{aligned} \frac{df_{t,a}(\underline{x}_0)}{dt} &= D\varphi(0) \frac{d\left(\tilde{F}(\tilde{\underline{x}})\right)_x}{dt} - D\varphi(0) \frac{\partial c_{t,a}}{\partial t} + \left(\left(\tilde{F}(\tilde{\underline{x}}) \right)_x - c_{t,a} \right) \frac{dD\varphi(0)}{dt} \\ &= D\varphi(0) B_4(\underline{x}) n \mu^{n-1} \frac{\partial \mu}{\partial t} y_3 + O(n), \end{aligned}$$

where we used (4.61), the fact that $D\varphi(0)$ is proportional to μ^n , Proposition 4.27, Lemma 4.49 and the fact that $\left(\tilde{F}(\tilde{x})\right)_x \in \alpha(U_{t,a})$ which has diameter proportional to $1/\mu^n$. In conclusion, using Lemma 4.12,

$$\frac{df_{t,a}}{dt} = D\varphi(0)B_4(c)n\mu^{n-1}\frac{\partial\mu}{\partial t}y_3(c) + O(n),$$

where $B_4(c)D\varphi(0)y_3(c)\partial\mu/\partial t$ is uniformly bounded away from zero. \square

5 The period doubling curve

In this section we prove the existence of a curve PD_n contained in the strip \mathcal{H}_n such that, each map corresponding to a parameter point in PD_n has a period doubling Cantor attractor. We start by recalling the following definition.

Definition 5.1. *Let M be a manifold and $f : M \rightarrow M$. An invariant Cantor set $A \subset M$ is called a period doubling Cantor attractor of f if $f|_A$ is conjugated to a 2-adic adding machine and there is a neighborhood $M \supset U \supset A$ such that the orbit of almost every point in U accumulates at A .*

Remark 5.2. *A period doubling Cantor attractor has zero topological entropy. It carries a unique invariant probability measure. Strongly dissipative Hénon-like maps at the boundary of chaos have period doubling Cantor attractors, see [11].*

We define the period doubling locus PD_n as,

$$PD_n = \{(t, a) \in \mathcal{H}_n \mid HF_{t,a} \text{ has a period doubling Cantor attractor}\}.$$

Proposition 5.3. *Let $F : \mathcal{P} \times M \rightarrow M$ be a real-analytic two dimensional unfolding of a map f with a strong homoclinic tangency, then the period doubling locus PD_n contains the graph of a real-analytic function $[-t_0, t_0] \ni t \mapsto PD_n(t) \in \mathbb{R}$. Moreover*

$$\frac{dPD_n}{dt} = -\frac{n}{\mu^{n+1}(t, a)}\frac{\partial\mu}{\partial t} + O\left(\frac{n}{\mu^{2n}(t, a)}\right).$$

The proof of Proposition 5.3 needs some preparation. Denote the disk of radius 2 centered at 0 by \mathbb{D}_2 . Let Y be the vector space of holomorphic and real symmetric functions $\epsilon : \mathbb{D}_2 \times \mathbb{D}_2 \rightarrow \mathbb{D}$ with $\epsilon(0, 0) = 0$. Y is equipped with the \mathcal{C}^0 norm.

A Hénon-like map is a map $HF : [-2, 2]^2 \rightarrow \mathbb{R}^2$ of the form

$$HF \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x^2 + \nu + \epsilon(x, y) \\ x \end{pmatrix}, \quad (5.4)$$

where $\nu \in \mathbb{R}$ and $\epsilon \in Y$. The space of Hénon-like maps is $\mathcal{H} = \mathbb{R} \times Y$. Observe that the maps defined in Theorem 3.61 are in \mathcal{H} . In particular the set \mathcal{H}_n defines a two-dimensional family in \mathcal{H} .

In the space of Hénon-like maps, there exists a codimension one manifold PD of maps which have a period doubling Cantor attractor, see [11]. In particular the degenerate

Hénon family is transversal to the PD manifold. This manifold is locally the graph of a real-analytic function

$$Ch : Y_\rho \rightarrow \mathbb{R}$$

where $Y_\rho = \{\epsilon \in Y \mid |\epsilon| \leq \rho\}$ with ρ sufficiently small. In particular,

$$HF = (\nu, \epsilon) \in PD \cap (\mathbb{R} \times Y_\rho) \iff \nu - Ch(\epsilon) = 0.$$

Observe that the maps defined in Theorem 3.61 are in $\mathcal{H}_\rho = \mathbb{R} \times Y_\rho$. In particular the set \mathcal{H}_n defines a two-dimensional family in \mathcal{H}_ρ .

Lemma 5.5. *There exists an $E > 0$ such that, for n large enough and every $t \in [-t_0, t_0]$, the family*

$$[-E, E] \ni \beta \mapsto HF_\beta = HF_{t, \text{san}(t) + \beta / |\mu(t, a_n(t))|^{2n}}$$

contains HF_0 with entropy 0 and HF_{-E} with entropy at least $\log 2$.

Proof. The degenerate Hénon map $(x, y) \mapsto (x^2, x)$ has a sink at $(0, 0)$ and entropy zero. Because this is an open condition, for n large, HF_0 being arbitrarily close to this degenerate Hénon map has also entropy zero.

The degenerate Hénon map $(x, y) \mapsto (x^2 + \nu, x)$, with $\nu \leq -3$ has a horseshoe and in particular it has entropy $\log 2$. Let

$$E = \frac{4}{\min_{\cup_{n>0} \mathcal{H}_n} (B_4(c)D\varphi(0)/\mu^n)}. \quad (5.6)$$

Observe that E is a positive finite number because $B_4(c)$ is uniformly away from zero, see Remark 3.17 and $D\varphi(0)$ is proportional to μ^n . From (5.8), $\nu_{-E} \leq -3$ and in particular, for n large enough, HF_{-E} has entropy at least $\log 2$. \square

Lemma 5.7. *The family*

$$\mathcal{H}_n \ni (t, a) \mapsto HF_{t,a} \in \mathcal{H}_\rho$$

is transversal to PD and intersects PD in a single curve.

Proof. Fix $(t, a) \in \mathcal{H}_n$. From (3.62), we get

$$\frac{\partial \nu_{t,a}}{\partial a} = \frac{\partial D\varphi(0)}{\partial a} (v_{t,a} - c_{t,a}) + D\varphi(0) \left(\frac{\partial v_{t,a}}{\partial a} - \frac{\partial c_{t,a}}{\partial a} \right),$$

and by Lemma 4.6, Proposition 3.34, Lemma 3.54, we get

$$\frac{\partial \nu_{t,a}}{\partial a} = D\varphi(0)B_4(c)\mu^n + O(\mu^n). \quad (5.8)$$

As a consequence,

$$\frac{\partial \epsilon_{t,a}}{\partial a} = \frac{\partial f_{t,a}}{\partial \beta} \frac{\partial \beta}{\partial a} - \frac{\partial \nu_{t,a}}{\partial a} = O(n\mu^n), \quad (5.9)$$

where we used Theorem 4.1. Fix $t \in [-t_0, t_0]$ and consider the family $[-E, E] \ni \beta \mapsto HF_\beta = (\nu_\beta, \epsilon_\beta) \in \mathcal{H}_\rho$. The function

$$[-E, E] \ni \beta \mapsto \nu_\beta - Ch(\epsilon_\beta) \in \mathbb{R}$$

is strictly monotone, because

$$\frac{\partial}{\partial \beta} (\nu_\beta - Ch(\epsilon_\beta)) = \frac{D\varphi(0)B_4(c)}{\mu^n} + O\left(\frac{n}{\mu^n}\right) \neq 0,$$

where we used (5.8), (5.9) and the fact that $\partial a/\partial \beta$ is proportional to $1/\mu^{2n}$. This means that the curve $\beta \mapsto HF_\beta$ is transversal to the level set of $\nu - Ch(\epsilon)$. In particular, the preimage of a level set of $\nu - Ch(\epsilon)$ is the graph of a function over the t -axes.

From Lemma 5.5, we know that HF_0 has entropy zero and HF_{-E} has entropy at least $\log 2$. The PD manifold in \mathcal{H}_ρ is the graph of the function Ch . Hence, $\mathcal{H}_\rho \setminus PD$ has two connected components, one containing HF_0 and the other HF_{-E} . In particular there exists a unique β_∞ such that $HF_{\beta_\infty} \in PD$. In particular, the two dimensional family $HF_{t,a}$ intersects PD transversally in a single curve. \square

Proof of Proposition 5.3. Consider the preimage of PD under the family $\mathcal{H}_n \ni (t, a) \mapsto HF_{t,a}$. By Lemma 5.7 and its proof, this preimage is the graph of a real-analytic function and it is contained in PD_n . Abusing the notation we denote this function by $[-t_0, t_0] \mapsto PD_n(t)$. Moreover by (3.62), we get

$$\frac{\partial \nu_{t,a}}{\partial t} = \frac{\partial D\varphi(0)}{\partial t} (v_{t,a} - c_{t,a}) + D\varphi(0) \left(\frac{\partial v_{t,a}}{\partial t} - \frac{\partial c_{t,a}}{\partial t} \right),$$

and by Lemma 4.49, Proposition 4.27, Lemma 3.54, we get,

$$\frac{\partial \nu_{t,a}}{\partial t} = D\varphi(0)B_4 n \mu^{n-1} \frac{\partial \mu}{\partial t} y_3 + O(n). \quad (5.10)$$

As a consequence,

$$\frac{\partial \epsilon_{t,a}}{\partial t} = \frac{\partial f_{t,a}}{\partial t} - \frac{\partial \nu_{t,a}}{\partial t} = O(n), \quad (5.11)$$

where we used Theorem 4.46. For every $t \in [-t_0, t_0]$, one has

$$\nu_{t,PD_n(t)} - Ch(\epsilon_{t,PD_n(t)}) = 0.$$

Hence,

$$0 = \frac{\partial \nu}{\partial t} + \frac{\partial \nu}{\partial a} \frac{dPD_n}{dt} - DCh \left(\frac{\partial \epsilon}{\partial t} + \frac{\partial \epsilon}{\partial a} \frac{dPD_n}{dt} \right),$$

and by using (5.8), (5.9), (5.10), (5.11) we have

$$\frac{dPD_n}{dt} = -\frac{n}{\mu} \frac{\partial \mu}{\partial t} y_3 + O\left(\frac{n}{\mu^{2n}(t,a)}\right).$$

Observe that, using (4.31), $y_3 = 1/\mu^n(y_4) = 1/\mu^n(1 + O(1/\mu^n))$. The proposition follows. \square

6 Coexistence of non-periodic attractors

In this section we prove that each real-analytic two-dimensional unfolding contains countably many maps with two period doubling Cantor attractors as well as countably many maps with finitely many sinks and two period doubling Cantor attractors.




Figure 4: Intersections of curves of period doubling attractors

Theorem 6.1. *Let $F : \mathcal{P} \times M \rightarrow M$ be a real-analytic two dimensional unfolding of a map f with a strong homoclinic tangency, then there exists a countable set $2PD \subset \mathcal{P}$, such that, each map in $2PD$ has two period doubling Cantor attractors. The homoclinic tangency persists along a curve in \mathcal{P} and it is contained in the closure of $2PD$.*

Proof. By Proposition 4 in [4], each \mathcal{H}_n contains finitely many curves which are graphs of functions $b_{n,n_0} : [t_{n,n_0}^-, t_{n,n_0}^+] \mapsto \mathbb{R}$. The maps corresponding to points in these curves have a secondary homoclinic tangency. Moreover, for all $E > 0$ and for n large enough,

$$b_{n,n_0}(t_{n,n_0}^+) = sa_n(t) + \frac{E}{|\mu(t, sa_n(t))|^{2n}} \text{ and } b_{n,n_0}(t_{n,n_0}^-) = sa_n(t) - \frac{E}{|\mu(t, sa_n(t))|^{2n}}.$$

As consequence, each curve b_{n,n_0} crosses the strip \mathcal{H}_n and in particular the curve PD_n . Let

$$\mathcal{P}_{n,n_0} = \left\{ (t, a) \in [-t_0, t_0] \times [-a_0, a_0] \mid t \in [t_{n,n_0}^-, t_{n,n_0}^+], |a - sa_n(t)| \leq \frac{E}{|\mu(t, sa_n(t))|^{2n}} \right\}.$$

Proposition 5 in [4] says that $F : \mathcal{P}_{n,n_0} \times M \mapsto M$ can be reparametrized to become an unfolding. We apply now the previous sections to the restricted unfoldings and we get new curves $PD_m^{(n,n_0)}$, corresponding to maps with a period doubling Cantor attractor which, for m large enough, accumulate to the curves b_{n,n_0} , see Proposition 5.3. Because the curves b_{n,n_0} intersect the curve PD_n , then also the curves $PD_m^{(n,n_0)}$ intersect PD_n and the intersection points correspond to maps with 2 period doubling Cantor attractors, see Figure 4. Finally, by Proposition 3 in [4], the distance between \mathcal{P}_{n,n_0} and \mathcal{P}_{n,n_0+1} is of the order $1/n$. Hence, the tangency curve at $a = 0$ is contained in the closure of the set of maps with two period doubling Cantor attractors. \square

Theorem 6.2. *Fix $S \in \mathbb{N}$. Let $F : \mathcal{P} \times M \rightarrow M$ be a real-analytic two dimensional unfolding of a map f with a strong homoclinic tangency, then there exists a countable set $S2PD \subset \mathcal{P}$, such that, each map in $S2PD$ has at least S sinks and two period doubling Cantor attractors. The homoclinic tangency persists along a curve in \mathcal{P} and it is contained in the closure of $S2PD$.*

Proof. This is a consequence of Theorem A in [4] and Theorem 6.1. It is enough to stop the inductive procedure in the proof of Theorem A in [4] at the step S . At this moment

there are boxes $\mathcal{P}_{n,n_0}^S \subset \mathcal{P}$ which are crossed diagonally by curves of secondary homoclinic tangencies, b_{n,n_0}^S . The family restricted to each of these boxes, \mathcal{P}_{n,n_0}^S , is an unfolding of a map with a strong homoclinic tangency given by the curve b_{n,n_0}^S and all maps in \mathcal{P}_{n,n_0}^S have at least S sinks, see Proposition 5 in [4]. By applying Theorem 6.1 to each of these restricted families, we find countable sets $S2PD_{n,n_0} \subset \mathcal{P}_{n,n_0}^S$ consisting of maps with at least S sinks and two period doubling Cantor attractors. The closure of each $S2PD_{n,n_0}$ contains the curve of secondary tangencies b_{n,n_0}^S . Because, by Proposition 3 in [4], the distance between \mathcal{P}_{n,n_0}^S and \mathcal{P}_{n,n_0+1}^S is of the order $1/n$ the set $S2PD = \cup_{n,n_0} S2PD_{n,n_0}$ contains in its closure the tangency curve at $a = 0$. \square

7 Laminations of multiple attractors

We are now ready to prove the coexistence of multiple attractors and study their stability. In families with at least three parameters, we construct maps with several period doubling Cantor attractors as well as maps with sinks and period doubling Cantor attractors forming laminations. We split the discussion in two subsections in which the main theorems and their meanings are carefully presented.

7.1 Laminations in general unfoldings

In higher dimensional families with a two dimensional section which is an unfolding of a homoclinic tangency, the sinks and the two period doubling attractors constructed in Theorem 6.1, they start to move simultaneously creating codimension two laminations, see Theorem 7.3. The leafs of the laminations are real-analytic and they have uniform positive diameter. The same phenomenon holds for the finitely many sinks and the two period doubling Cantor attractors constructed in Theorem 6.2, see Theorem 7.11.

Given a two-dimensional family F which is an unfolding of a map with a strong homoclinic tangency, recall that there exist curves PD_n , in parameter space, corresponding to maps with one period doubling Cantor attractor. Furthermore, from the proof of Theorem 6.1, for every n large enough, there are finitely many curves b_{n,n_0} consisting of maps with secondary tangencies and they cross the curve PD_n . The curves b_{n,n_0} are accumulated by new period doubling curves $PD_m^{(n,n_0)}$. In the next proposition we compare the angle of the curves b_{n,n_0} and $PD_m^{(n,n_0)}$ with the angle of the curve PD_n .

Proposition 7.1. *The curves b_{n,n_0} and $PD_m^{(n,n_0)}$ cross the curve PD_n transversally and the angle is larger than $Vn\mu_{min}^{-3n/2}$ where V is a uniform constant and $\mu_{min} = \min_{(t,a)} |\mu(t,a)|$.*

Proof. By Proposition 4 in [4],

$$\frac{db_{n,n_0}}{dt} = -\frac{n}{\mu^{n+1}} \frac{\partial \mu}{\partial t} + V_{t,a} n \lambda^{\theta n} + O(|\lambda|^{\theta n}),$$

where $V_{t,a}$ is uniformly away from zero, $0 < \theta < 1/2$ and $|\lambda|^{2\theta} |\mu|^3 > 1$. In particular, by Proposition 5.3, the curve b_{n,n_0} crosses the curve PD_n transversally and the angle is larger than, $Vn\mu^{-3n/2}$ where V is a uniform constant. Because, for m large, the curve $PD_m^{(n,n_0)}$ is \mathcal{C}^1 close to b_{n,n_0} , see Proposition 5.3, the same angle estimate holds for $PD_m^{(n,n_0)}$ and the transversality with PD_n follows. \square

Remark 7.2. Observe that, by Proposition 7.1, the angle formed by the intersection of $PD_m^{(n,n_0)}$ with the curve PD_n is larger than $Vn\mu_{\min}^{-3n/2}$, for $n \geq n_0$. By Remark 10 in [4], n_0 and V are locally constant, i.e. they depend continuously on the family.

Theorem 7.3. Let M , \mathcal{P} and \mathcal{T} be real-analytic manifolds and $F : (\mathcal{P} \times \mathcal{T}) \times M \rightarrow M$ a real-analytic family with $\dim(\mathcal{P}) = 2$ and $\dim(\mathcal{T}) \geq 1$. If there exists $\tau_0 \in \mathcal{T}$ such that $F_0 : (\mathcal{P} \times \{\tau_0\}) \times M \rightarrow M$ is an unfolding of a map f_{τ_0} with a strong homoclinic tangency, then for $k = 1, 2$, there exists a codimension k lamination of maps with at least k period doubling Cantor attractors which persist along the leaves. The homoclinic tangency persists along a global codimension one manifold in $\mathcal{P} \times \mathcal{T}$ and this tangency locus is contained in the closure of the lamination. Moreover, the leaves of the lamination are real-analytic and they have a uniform positive diameter.

Proof. Without loss of generality we may assume that $\mathcal{P} \times \mathcal{T} = [-1, 1]^2 \times [-1, 1]^{r-2}$ where $r = \dim(\mathcal{P} \times \mathcal{T})$. A point in parameter space is given by (t, a, τ) and $a = 0$ corresponds to the tangency locus. Moreover, we may also assume that, for all $\tau \in \mathcal{T}$, the family restricted to $[-1, 1]^2 \times \{\tau\}$, F_τ , is an unfolding of a strong homoclinic tangency of the map f_τ . Given τ , denote the n^{th} Hénon strip of the family restricted to $[-1, 1]^2 \times \{\tau\}$ by $\mathcal{H}_n(\tau)$.

For $k = 1$, Theorem 7.5 is a reformulation of Lemma 5.7. Namely, for every n , there is a real-analytic function $PD_n : (t, \tau) \mapsto PD_n(t, \tau)$ such that the graph of PD_n in $\mathcal{P} \times \mathcal{T}$, i.e. points of the form $(t, PD_n(t, \tau), \tau)$, consists of maps with at least one period doubling Cantor attractor. Because, for a fixed τ , these graphs are contained in $\mathcal{H}_n(\tau)$ and the distance of $\mathcal{H}_n(\tau)$ to the tangency locus at $a = 0$ is of the order $1/\mu^n$, the graphs of PD_n contain the tangency locus in their closure.

Let us consider now the case $k = 2$. As in the proof of Theorem 6.1, from Proposition 7.1, we get that in the unfolding F_τ , there are period doubling curves $PD_m^{(n,n_0)}(\tau)$ and $PD_n(\tau)$ which intersect transversally in the point $(t_m^{(n,n_0)}(\tau), a_m^{(n,n_0)}(\tau), \tau)$, see Figure 4. From Proposition 7.1 we get a lower bound for the angle between these curves which is independent of the parameter τ , see Remark 7.2. This transversality implies that this intersection persists for all τ as the graph of a real-analytic function. In particular the two period doubling attractors at $(t_m^{(n,n_0)}(\tau), a_m^{(n,n_0)}(\tau), \tau)$ have their continuation in all unfoldings F_τ , creating a codimension two real-analytic leaf of the lamination. Let

$$2PD(\tau) = \{PD_m^{(n,n_0)}(\tau) \cap PD_n(\tau) \mid n, n_0, m > 0\}.$$

According to Theorem 6.1, the set $2PD(\tau)$ consists of countably many maps in the family F_τ which have two period doubling attractors. Moreover, it accumulates at the tangency curve in $\mathcal{P} \times \{\tau\}$. The set $2PD = \cup_\tau 2PD(\tau)$ is the required lamination. In particular for a given τ , the set $2PD(\tau)$ moves along the leaves of $2PD$ while varying τ and all leaves project onto $[-1, 1]^{r-2}$, they have uniform diameter. \square

7.2 Laminations in saddle deforming unfoldings

We analyze here coexisting phenomena in saddle deforming unfoldings. In these families, whose definition and meaning is explained in the following, we find codimension three laminations of coexisting attractors and we give a precise description of the asymptotic direction of the leaves of the laminations. We start with some basic definition and explanation.

Two crucial invariants associate to a saddle point are indeed its eigenvalues. One of the properties of an unfolding is that the unstable eigenvalue changes with one parameter, see Definition 2.7. We introduce here a notion of unfolding for families with at least three parameters. We require that the unfolding of the homoclinic tangency is also able to change both eigenvalues independently. In these so called *saddle deforming unfoldings*, the level sets of the eigenvalue pair define a codimension two foliation of the tangency locus, called the *eigenvalue foliation*. A saddle deforming unfolding contains a three dimensional subfamily transversal to the codimension three leafs of the eigenvalue foliation.

Definition 7.4. *Let M , \mathcal{P} and \mathcal{T} be real-analytic manifolds and $F : (\mathcal{P} \times \mathcal{T}) \times M \rightarrow M$ a real-analytic family with $\dim(\mathcal{P}) = 2$ and $\dim(\mathcal{T}) \geq 1$. If there exists $\tau_0 \in \mathcal{T}$ such that $F_0 : (\mathcal{P} \times \{\tau_0\}) \times M \rightarrow M$ is an unfolding of a map with a strong homoclinic tangency and the eigenvalues at the saddle point, after a reparametrization, satisfy*

$$\frac{\partial \mu}{\partial \tau} = 0, \quad \frac{\partial \lambda}{\partial \tau} \neq 0,$$

then the family $(t, a, \tau) \mapsto F_{t,a,\tau}$ is called a saddle deforming unfolding.

Theorem 7.5. *Let M , \mathcal{P} and \mathcal{T} be real-analytic manifolds and $F : (\mathcal{P} \times \mathcal{T}) \times M \rightarrow M$ be a saddle deforming unfolding then there exists a codimension 3 lamination 3PD of maps with at least 3 period doubling Cantor attractors which persist along the leafs. The leafs of the lamination are real-analytic and when $\dim(\mathcal{T}) \geq 2$ they have a uniform positive diameter. The homoclinic tangency persists along a global codimension one manifold in $\mathcal{P} \times \mathcal{T}$ and for each leaf of the eigenvalue foliation, there is a sequence of leafs of 3PD which accumulate at this eigenvalue leaf.*

Proof. Without loss of generality we may assume that $\mathcal{P} \times \mathcal{T} = [-1, 1]^2 \times [-1, 1]^{r-2}$ where $r = \dim(\mathcal{P} \times \mathcal{T})$. A point in parameter space is given by $(t, a, \underline{\tau})$ and $a = 0$ corresponds to the tangency locus. Moreover, we may also assume that, for all $\underline{\tau} \in \mathcal{T}$, the family restricted to $[-1, 1]^2 \times \{\underline{\tau}\}$, $F_{\underline{\tau}}$, is an unfolding of a strong homoclinic tangency. Given $\underline{\tau}$, there are period doubling curves $PD_n(\underline{\tau}) \subset [-1, 1]^2 \times \{\underline{\tau}\}$ contained in the n^{th} Hénon strip of the family restricted to $[-1, 1]^2 \times \{\underline{\tau}\}$ which we denote here by $\mathcal{H}_n(\underline{\tau})$. By Proposition 4 in [4], each $\mathcal{H}_n(\underline{\tau})$ contains finitely many curves which are graphs of functions $b_{n,n_0}(\underline{\tau}) : [t_{n,n_0}^-(\underline{\tau}), t_{n,n_0}^+(\underline{\tau})] \mapsto \mathbb{R}$. The maps corresponding to points in these curves have a strong homoclinic tangency. Moreover, by Proposition 7.1, each curve $b_{n,n_0}(\underline{\tau})$ crosses transversally the curve $PD_n(\underline{\tau})$, in a point denoted by $\mathcal{B}_{n,n_0}(\underline{\tau})$. Because of the transversality, the points $\mathcal{B}_{n,n_0}(\underline{\tau})$ define a real-analytic codimension two manifold $\mathcal{B}_{n,n_0} \subset \mathcal{P} \times \mathcal{T}$. As in the proof of Theorem 7.3, let $PD_n = \cup_{\underline{\tau}} PD_n(\underline{\tau}) = \mathcal{P}_n \subset \mathcal{P} \times \mathcal{T}$ be a codimension one manifold. Observe that, maps corresponding to points in \mathcal{P}_n have a period doubling Cantor attractor and $\mathcal{B}_{n,n_0} \subset \mathcal{P}_n$ is a codimension one manifold consisting of maps with also a strong homoclinic tangency.

In the following we prove that the restriction of the initial family $F : \mathcal{P}_n \times M \rightarrow M$ contains an unfolding of the strong homoclinic tangency of the maps in \mathcal{B}_{n,n_0} . The only condition which is not straightforward is that $d\mu$ is non zero along \mathcal{B}_{n,n_0} . After a reparametrization of the coordinates $\underline{\tau} = (\tau, \tau_2, \dots, \tau_{r-2})$, we may assume that $\partial \lambda / \partial \tau = 1$ and $\partial \lambda / \partial \tau_j = 0$ for all $2 \leq j \leq r-2$. This is possible because, by hypothesis, our family is a saddle deforming unfolding. It suffices to prove that the two dimensional restriction of $F : \mathcal{P}_n \times M \rightarrow M$ to the coordinates $(t, a(t, \tau, 0, \dots, 0), \tau, 0, \dots, 0)$ is an unfolding. In

this two dimensional family, the tangency locus \mathcal{B}_{n,n_0} is a curve parametrized by τ , say the points in this curve are of the form $(t(\tau), a(\tau), \tau)$. The aim is to prove that $d\mu/d\tau \neq 0$ along this curve. Observe that

$$\frac{d\mu}{d\tau} = \frac{\partial\mu}{\partial t} \frac{dt}{d\tau} + \frac{\partial\mu}{\partial a} \frac{da}{d\tau} + \frac{\partial\mu}{\partial\tau}$$

where $\partial\mu/\partial t \neq 0$ and $\partial\mu/\partial\tau = 0$ because our initial family is a saddle deforming unfolding. Moreover, using Proposition 5.3 and the fact that $\partial\mu/\partial\tau = 0$, one has $da/d\tau = O(n/\mu^{2n})$. Hence,

$$\frac{d\mu}{d\tau} = \frac{\partial\mu}{\partial t} \frac{dt}{d\tau} + O\left(\frac{n}{\mu^{2n}}\right). \quad (7.6)$$

Let $(\Delta t, \Delta a, \Delta\tau)$ be a tangent vector to the curve \mathcal{B}_{n,n_0} . From the proof of Theorem B in [4], we have

$$\left(1 + O\left(\frac{1}{n}\right)\right) (V\Delta t + V_1\Delta\tau) = 0.$$

Hence,

$$\frac{dt}{d\tau} = \frac{\Delta t}{\Delta\tau} = -\frac{V_1}{V} \left(1 + O\left(\frac{1}{n}\right)\right), \quad (7.7)$$

with V a function uniformly away from zero and by (4.47) in [4], there are functions, K and K' uniformly bounded away from zero, such that

$$V_1 = K \frac{\partial\lambda}{\partial\tau} + K' \frac{\partial\mu}{\partial\tau} = K \frac{\partial\lambda}{\partial\tau} \neq 0, \quad (7.8)$$

where we used the property $\partial\mu/\partial\tau = 0$ of saddle deforming unfolding. Combining (7.6), (7.7) and (7.8) we get that $d\mu/d\tau \neq 0$ and our family is an unfolding. By Theorem 7.3, \mathcal{P}_n contains a codimension 2 lamination, denoted by $3PD_{n,n_0}$, of maps with 2 period doubling Cantor attractors which persist along the real-analytic leafs of the lamination. Moreover, the lamination contains \mathcal{B}_{n,n_0} in its closure. Observe that, all maps in \mathcal{P}_n have at least one period doubling Cantor attractor of a fixed combinatorial type. Hence, every map in $3PD_{n,n_0}$ actually has at least 3 period doubling Cantor attractors.

By Proposition 3 in [4], the distance between $b_{n,n_0}(\underline{\tau})$ and $b_{n,n_0+1}(\underline{\tau})$ is of the order $1/n$. Hence, the tangency locus at $a = 0$ is contained in the closure of the lamination $3PD = \cup_{n,n_0} 3PD_{n,n_0}$ formed by maps with at least 3 period doubling Cantor attractors.

For understanding the diameter of the leafs, write $\mathcal{P} \times \mathcal{T} = [-1, 1]^2 \times [-1, 1] \times [-1, 1]^{r-3}$, the parameters as (t, a, τ, τ') and denote the set of parameters of the form $(\cdot, \cdot, \cdot, \tau')$ by $\mathcal{P}(\tau')$. For a given τ' the lamination $3PD$ intersects $\mathcal{P}(\tau')$ in countably many points which move along the leafs of $3PD$ while varying τ' . In particular, all leafs project onto $[-1, 1]^{r-3}$, they have uniform diameter.

In order to complete the proof, observe that each family \mathcal{P}_n is the graph of a function $[-1, 1]^2 \times [-1, 1]^{r-3} \ni (t, \tau, \tau') \mapsto \mathcal{P}_n(t, \tau, \tau') \in [-1, 1]$. In particular, we can reparametrize \mathcal{P}_n by $[-1, 1]^2 \times [-1, 1]^{r-3}$ using the parameters (t, τ, τ') . For every $\tau' \in [-1, 1]^{r-3}$, we can identify the two dimensional family $\mathcal{P}_n(\tau') = \mathcal{P}_n \cap \mathcal{P}(\tau')$ with $[-1, 1]^2$. This family contains the curve $\mathcal{P}_n(\tau') \cap \mathcal{B}_{n,n_0}$ of strong homoclinic tangencies and, as shown before, it is an unfolding of these tangencies. For each $\tau' \in [-1, 1]^{r-3}$ we can choose a reparametrization of $\mathcal{P}_n(\tau')$, depending real-analytically on τ' , obtaining coordinates

$(t', a', \tau') \in [-1, 1]^2 \times [-1, 1]^{r-3}$ such that $a' = 0$ corresponds to the tangency locus. Moreover we can assume

$$\frac{\partial \mu}{\partial t'} = 1, \frac{\partial \mu}{\partial \tau'} = 0, \frac{\partial \lambda}{\partial \tau'} = 0. \quad (7.9)$$

We apply now the previous sections to the restricted unfolding and we get new curves $PD'_m(\tau') \subset \mathcal{P}_n(\tau')$, corresponding to maps with a period doubling Cantor attractor, see Proposition 5.3. Using Proposition 4 in [4] and Proposition 7.1, there are finitely many curves $b'_{m,m_0}(\tau') \subset \mathcal{P}_n(\tau')$ which cross $PD'_m(\tau')$ transversally. Because this intersection depends real-analytically on τ' , the intersection points form a codimension two manifold $\mathcal{B}'_{m,m_0} \subset \mathcal{P}_n$.

Let $(\Delta t', \Delta a', \Delta \tau')$ be a tangent vector to the manifold \mathcal{B}'_{m,m_0} . From the proof of Theorem B in [4], we have

$$\left(V' + O\left(\frac{1}{m}\right) \right) \Delta t' + \sum_{i=1}^{r-3} \left(V'_i + O\left(\frac{1}{m}\right) \right) \Delta \tau'_i = 0,$$

and

$$O\left(\frac{m}{\mu^m} (\Delta t' + \Delta \tau')\right) = \Delta a',$$

with V' a function uniformly away from zero and by (4.47) in [4], there are functions, K_i and K'_i uniformly bounded away from zero, such that

$$V'_i = K_i \frac{\partial \lambda}{\partial \tau'} + K'_i \frac{\partial \mu}{\partial \tau'} = 0,$$

where we used (7.9). Hence,

$$\Delta t' = O\left(\frac{1}{m} \Delta \tau'\right) \text{ and } \Delta a' = O\left(\frac{m}{\mu^m} \Delta \tau'\right). \quad (7.10)$$

Choose a point $(t'_0, a'_0, \tau'_0) \in \mathcal{B}'_{m,m_0}$. Observe that $a'_0 = O(1/\mu^m)$. As a consequence of (7.10), the manifold \mathcal{B}'_{m,m_0} is at the distance of the order $1/m$ to a codimension two subspace defined by $t' = t'_0$ and $a' = a'_0$. By (7.9) this subspace is a level set for the eigenvalues.

As final remark, according to Proposition 7.1 for every $\tau' \in [-1, 1]^{r-3}$ there exists a sequence of curves $PD_k^{(m,m_0)}(\tau')$ accumulating in \mathcal{C}^1 at $\mathcal{B}'_{m,m_0}(\tau')$. The transversal intersection points with $PD_m(\tau')$ varies real-analytically with τ' forming a leaf of the lamination $3PD$. Hence, there are leaves of $3PD$ which accumulates at \mathcal{B}'_{m,m_0} which by them self accumulate at the leaves of the eigenvalue foliation. \square

As before one also gets the coexistence of finitely many sinks and finitely many period doubling Cantor attractor. The proof is similar to the proof of Theorem 6.2.

Theorem 7.11. *Let M, \mathcal{P} and \mathcal{T} be real-analytic manifolds and $F : (\mathcal{P} \times \mathcal{T}) \times M \rightarrow M$ be a saddle deforming unfolding, then for every $1 \leq k \leq 3$ and $S \in \mathbb{N}$, there exists a codimension k lamination $SkPD$ of maps with at least S sinks and k period doubling Cantor attractors which persist along the leaves. The homoclinic tangency persists along a global codimension one manifold in $\mathcal{P} \times \mathcal{T}$ and this tangency locus is contained in the closure of the lamination. The leaves of the lamination are real-analytic and when $1 \leq k < \dim(\mathcal{P} \times \mathcal{T})$ they have a uniform positive diameter. Moreover, for each leaf of the eigenvalue foliation, there is a sequence of leaves of $S3PD$ which accumulate at this eigenvalue leaf.*

Proof. Without loss of generality we may assume that $\mathcal{P} \times \mathcal{T} = [-1, 1]^2 \times [-1, 1]^{r-2}$ where $r = \dim(\mathcal{P} \times \mathcal{T})$. A point in parameter space is given by (t, a, τ) and $a = 0$ corresponds to the tangency locus. Moreover, we may also assume that, for all $\tau \in \mathcal{T}$, the family restricted to $[-1, 1]^2 \times \{\tau\}$, F_τ , is an unfolding of a strong homoclinic tangency of the map f_τ . We apply the inductive procedure in the proof of Theorem A in [4] S times. At this moment there are boxes $\mathcal{P}_{n,n_0}^S(\tau) \subset \mathcal{P} \times \{\tau\}$ which are crossed diagonally by curves of secondary homoclinic tangencies, $b_{n,n_0}^S(\tau)$. The family F_τ restricted to each of these boxes $\mathcal{P}_{n,n_0}^S(\tau)$ is an unfolding of a map with a strong homoclinic tangency given by the curve $b_{n,n_0}^S(\tau)$ and all maps in the box have at least S sinks, see Proposition 5 in [4]. Because the boxes $\mathcal{P}_{n,n_0}^S(\tau)$ depends analytically on τ , we can consider the union $\mathcal{P}_{n,n_0}^S = \cup_{\tau} \mathcal{P}_{n,n_0}^S(\tau)$. Since the definition of saddle deforming unfolding only involves the dependence of λ on the parameters, the restriction of the initial family to \mathcal{P}_{n,n_0}^S is still a saddle deforming unfolding and each map in this restriction has S sinks. We apply Theorem 7.3 and Theorem 7.5 to the restricted family.

We consider first the cases $k = 1, 2$. For each \mathcal{P}_{n,n_0}^S we get a codimension k lamination $SkPD_{n,n_0}$ of maps with at least S sinks and k period doubling Cantor attractors. The leaves of this lamination project onto \mathcal{T} and they contain the tangency locus of the restricted family in their closure. Observe that, by Proposition 3 in [4], the distance between $\mathcal{P}_{n,n_0}^S(\tau)$ and $\mathcal{P}_{n,n_0+1}^S(\tau)$ is of the order $1/n$ and the collection of boxes $\mathcal{P}_{n,n_0}^S(\tau)$ contains in its closure the tangency curve at $a = 0$. Hence, the set $SkPD = \cup_{n,n_0} SkPD_{n,n_0}$ contains in its closure the tangency locus at $a = 0$.

Consider the case $k = 3$. For each \mathcal{P}_{n,n_0}^S we get a codimension 3 lamination $S3PD_{n,n_0}$ of maps with at least S sinks and 3 period doubling Cantor attractors. If the dimension of the family is three, then the set $S3PD_{n,n_0}$ is countable and it accumulates at the tangency locus of \mathcal{P}_{n,n_0}^S . For the same reason as for the case $k = 1, 2$ it contains the tangency locus of the original family in its closure. If the dimension of the family is strictly larger than three, after a reparametrization of the coordinates $\tau = (\tau_1, \dots, \tau_{r-2})$, we may assume that $\partial\lambda/\partial\tau_1 = 1$ and $\partial\lambda/\partial\tau_j = 0$ for all $2 \leq j \leq r-2$. This is possible because, by hypothesis, our family is a saddle deforming unfolding. The leaves of the lamination $S3PD_{n,n_0}$ project onto $\{0\} \times \{0\} \times \{0\} \times [-1, 1]^{r-3}$ and they contain the tangency locus of the restricted family in their closure. As before, the set $S3PD = \cup_{n,n_0} S3PD_{n,n_0}$ contains in its closure the tangency locus at $a = 0$. \square

A similar phenomenon holds for laminations of infinitely many sinks and one period doubling Cantor attractor. This is stated precisely in the following.

Theorem 7.12. *Let M, \mathcal{P} and \mathcal{T} be real-analytic manifolds and $F : (\mathcal{P} \times \mathcal{T}) \times M \rightarrow M$ be a saddle deforming unfolding, then there exists a codimension 3 lamination NHPD of maps with infinitely many sinks and at least 1 period doubling Cantor attractor which persist along the leaves. The leaves of the lamination are real-analytic and when the dimension of \mathcal{T} is at least 2, they have a uniform positive diameter. Moreover, the homoclinic tangency persists along a global codimension one manifold in $\mathcal{P} \times \mathcal{T}$ and for each leaf of the eigenvalue foliation, there is a sequence of leaves of NHPD which accumulate at this eigenvalue leaf.*

Proof. We use the notation as in the proof of Theorem 7.5. Write the parameter space as $\mathcal{P} \times \mathcal{T} = [-1, 1]^2 \times [-1, 1] \times [-1, 1]^{r-3}$, the parameters as (t, a, τ, τ') and denote the set of parameters of the form $(\cdot, \cdot, \cdot, \tau')$ by $\mathcal{P}(\tau')$. Recall that $\mathcal{P}_n \subset \mathcal{P} \times \mathcal{T}$ is a codimension one manifold formed of maps with one period doubling Cantor attractor. For a given τ' ,

write $\mathcal{P}_n(\tau') = \mathcal{P}_n \cap \mathcal{P}(\tau')$. In the proof of Theorem 7.5 we proved that the restriction of our initial family, $F_{\tau'}$, to $\mathcal{P}_n(\tau')$ is a two dimensional unfolding of a strong homoclinic tangency. We apply Theorem A in [4] to $F_{\tau'}$ and we get a set $NHPD_n(\tau')$ of maps with infinitely many sinks. Each point in $NHPD_n(\tau')$ depends real-analytically on τ' and it forms a real-analytic codimension two manifold in \mathcal{P}_n . All points together form a codimension two lamination $NHPD_n$ of maps with infinitely many sinks and one period doubling Cantor attractor coming from being in \mathcal{P}_n . By construction the lamination contains the tangency loci \mathcal{B}_{n,n_0} in its closure. Let $NHPD = \cup_n NHPD_n$. As in the previous proofs, the lamination $NHPD$ contains its closure the tangency locus at $a = 0$.

Choose $\tau' \in [-1, 1]^{r-3}$ and let $f \in NHPD_n(\tau')$. From the proof of Theorem A in [4], there are nested boxes $\mathcal{P}^g(\tau')$ with $\{f\} = \cap^g \mathcal{P}^g(\tau')$. Each $\mathcal{P}^g(\tau')$ contains a real-analytic curve $b^g(\tau')$ consisting of maps with a strong homoclinic tangency associated to the original saddle point and the family restricted to $\mathcal{P}^g(\tau')$ is an unfolding of the maps in $b^g(\tau')$. Moreover, the box $\mathcal{P}^g(\tau')$ contains a curve $PD_m^g(\tau')$ consisting of maps with one period doubling Cantor attractor and the curve $b_{m,m_0}^g(\tau') \subset \mathcal{P}^{g+1}(\tau')$ of maps with a secondary homoclinic tangency. Similarly as in the proof of Theorem 7.5, by Proposition 7.1, the curve $b_{m,m_0}^g(\tau')$ intersects transversally the curve $PD_m^g(\tau')$ in a point $\mathcal{B}_{m,m_0}^g(\tau')$. Denote this intersection point by $\mathcal{B}^{g+1}(\tau') \in \mathcal{P}^{g+1}(\tau')$ and notice that it depends real-analytically on τ' forming a manifold \mathcal{B}^{g+1} . Observe that the leaf of the lamination $NHPD$ through f is the limit of the leaves \mathcal{B}^g . As was shown in the proof of Theorem 7.5, the leaves \mathcal{B}^g asymptotically align with the leaves of the eigenvalue foliation, see (7.10).

Any point in the tangency locus is in the closure of the set $NHPD$. In particular it is in the closure of a sequence of leaves of the lamination $NHPD$. In turn these leaves are in the closure of leaves of the form \mathcal{B}^g which are asymptotically align with the leaves of the eigenvalue foliation. Hence, there is a sequence of leaves of the lamination $NHPD$ which accumulates at any chosen level set of the eigenvalue foliation. This concludes the proof. \square

Our method allows also to find coexistence of period doubling Cantor attractors and a strange one. In this case, we do not have laminations because the stability of the strange attractors in families with at least three parameters is not yet understood. We start by recalling the formal definition of a *strange attractor*.

Definition 7.13. *Let M be a manifold and $f : M \rightarrow M$. An open set $U \subset M$ is called a trapping region if $\overline{f(U)} \subset U$. An attractor in the sense of Conley is*

$$\Lambda = \bigcap_{j \geq 0} f^j(U).$$

The attractor Λ is called topologically transitive if it contains a dense orbit. If Λ contains a dense orbit which satisfies the Collet-Eckmann conditions, i.e. there exist a point z , a vector $v \in T_z M$ and a constant $\kappa > 0$ such that

$$|Df^n(z)v| \geq e^{\kappa n} \text{ for all } n > 0,$$

then Λ is called a strange attractor.

Theorem 7.14. *Let M , \mathcal{P} and \mathcal{T} be real-analytic manifolds and $F : (\mathcal{P} \times \mathcal{T}) \times M \rightarrow M$ a saddle deforming unfolding, then the set of maps with at least 2 period doubling Cantor attractors and one strange attractor has Hausdorff dimension at least $\dim(\mathcal{P} \times \mathcal{T}) - 2$.*

Proof. We use the notation as in the proof of Theorem 7.5. Write the parameter space as $\mathcal{P} \times \mathcal{T} = [-1, 1]^2 \times [-1, 1] \times [-1, 1]^{r-3}$, the parameters as (t, a, τ, τ') and denote the set of parameters of the form $(\cdot, \cdot, \cdot, \tau')$ by $\mathcal{P}(\tau')$. Recall that $\mathcal{P}_n \subset \mathcal{P} \times \mathcal{T}$ is a codimension one manifold formed of maps with one period doubling Cantor attractor. For a given τ' , write $\mathcal{P}_n(\tau') = \mathcal{P}_n \cap \mathcal{P}(\tau')$. In the proof of Theorem 7.5 we proved that the restriction of our initial family, $F_{\tau'}$, to $\mathcal{P}_n(\tau')$ is an unfolding of a strong homoclinic tangency at $\mathcal{B}_{n,n_0}(\tau')$. We apply now the previous sections to the restricted unfolding and we get new curves $PD_m^{(n,n_0)}(\tau')$, corresponding to maps with a period doubling Cantor attractor, see Proposition 5.3. Using Proposition 4 in [4] and Proposition 7.1, there are finitely many curves $b_{m,m_0}^{(n,n_0)}(\tau')$ which cross $PD_m^{(n,n_0)}(\tau')$ transversally. The maps in the curves $b_{m,m_0}^{(n,n_0)}(\tau')$ have a strong homoclinic tangency and $PD_m^{(n,n_0)}(\tau')$ is a one-dimensional unfolding of the homoclinic tangency of the map at the intersection of the two curves. By Theorem A in [15], the curve $PD_m^{(n,n_0)}(\tau')$ contains a set of positive Lebesgue measure with a strange attractor. In particular we found a set, $2PDSA(\tau')$, of positive Lebesgue measure of maps with one strange attractor and at least two period doubling attractors, one from being in $\mathcal{P}_n(\tau')$ and the other from being in $PD_m^{(n,n_0)}(\tau')$. The proof is concluded by varying τ' and taking $2PDSA = \cup_{\tau'} 2PDSA(\tau')$. \square

8 The Hénon family and other polynomial families

In this section we apply our theorems to the Hénon family and other polynomial families. We start by recalling the definition of Hénon family. The two parameter family $F : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$,

$$F_{a,b} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} a + x^2 - by \\ x \end{pmatrix}, \quad (8.1)$$

is called the *Hénon family*. It is well known that the Hénon family is an unfolding of a map with a strong homoclinic tangency. A proof can be found for example in [4], or [6].

Theorem 8.2. *Fix $S \in \mathbb{N}$. The Hénon family contains a countable set $S2PD$ of maps with at least S sinks and two period doubling Cantor attractors. Moreover the set of Hénon maps with S sinks, one period doubling Cantor attractor and one strange attractor has Hausdorff dimension at least one.*

Proof. The first statement is an application of Theorem 6.2. For the second statement we stop the inductive procedure in the proof of Theorem A in [4] at the step S . At this moment there are boxes $\mathcal{P}_{n,n_0}^S \subset \mathcal{P}$ which are crossed diagonally by curves of secondary homoclinic tangencies, b_{n,n_0}^S . The family restricted to each of these boxes \mathcal{P}_{n,n_0}^S is an unfolding of a map with a strong homoclinic tangency at the curve b_{n,n_0}^S and all maps in the box have at least S sinks, see Proposition 5 in [4]. We apply now the previous sections to the restricted unfolding and we get new curves $PD_m^{(n,n_0,S)}$, corresponding to maps with a period doubling Cantor attractor, see Proposition 5.3. Using Proposition 4 in [4] and Proposition 7.1, there are finitely many curves $b_{m,m_0}^{(n,n_0,S)}$ which cross $PD_m^{(n,n_0,S)}$ transversally. The maps in the curves $b_{m,m_0}^{(n,n_0,S)}$ have a strong homoclinic tangency and $PD_m^{(n,n_0,S)}$ is a one-dimensional unfolding of the homoclinic tangency of the map at the intersection of the two curves. By Theorem A in [15], the curve $PD_m^{(n,n_0,S)}$ contains a set of positive Lebesgue measure of maps with a strange attractor. \square

Theorem 7.11, Theorem 7.12 and Theorem 7.14 give the following.

Theorem 8.3. *Let $\text{Poly}_d(\mathbb{R}^2)$ be the space of real polynomials of \mathbb{R}^2 of degree at most d , with $d \geq 2$. Given a map f in $\text{Poly}_d(\mathbb{R}^2)$ which has a strong homoclinic tangency, there is a global codimension one manifold along which this tangency persists. The following holds:*

- *for every $1 \leq k \leq 3$ and $S \in \mathbb{N}$ there exists a codimension k lamination $SkPD$ of maps with at least S sinks and k period doubling Cantor attractors which persist along the leafs,*
- *there exists a codimension 3 lamination $NHPD$ of maps with infinitely many sinks and at least 1 period doubling Cantor attractors which persist along the leafs,*
- *the set of maps with at least 2 period doubling Cantor attractors and one strange attractor has Hausdorff dimension at least $\dim(\text{Poly}_d(\mathbb{R}^2)) - 2$.*

The leafs of the laminations are real-analytic and they have uniform diameter. Moreover, for each leaf of the eigenvalue foliation, there is a sequence of leafs of $3PD$ and a sequence of leafs of $NHPD$ which accumulate at this eigenvalue leaf.

References

- [1] Ahlfors, L., *Complex Analysis*, McGraw-Hill, New York (1979).
- [2] Ahlfors, L., *Conformal Invariants*, Topics in Geometric Function Theory, McGraw-Hill, New York (1973).
- [3] Benedicks, Michael and Carleson, Lennart, *The dynamics of the Hénon map*, Ann. of Math. (2), 133, 1991, 1, 73–169.
- [4] Benedicks, Michael, Martens, Marco and Palmisano, Liviana, *Newhouse laminations*, Preprint, arXiv:1811.00617.
- [5] Benedicks, Michael and Palmisano, Liviana, *Coexistence phenomena in the Hénon family*, Preprint, arXiv:1811.00517.
- [6] Berger, P., *Zoology in the Hénon family: twin babies and Milnor’s swallows*, Preprint, arXiv:1801.05628 .
- [7] Bieberbach, L., *Conformal mapping*, Chelsea publishing, New York (1964)
- [8] Biebler, Sébastien, *Persistent homoclinic tangencies and infinitely many sinks for residual sets of automorphisms of low degree in \mathbb{C}^3* arXiv:1611.02011.
- [9] Bronstein, I. U. and Kopanskiĭ, A. Ya., *Smooth invariant manifolds and normal forms*, World Scientific Series on Nonlinear Science. Series A: Monographs and Treatises, 7, World Scientific Publishing Co., Inc., River Edge, NJ, 1994, xii+384.
- [10] Buzzard, Gregory T., *Infinitely many periodic attractors for holomorphic maps of 2 variables*, Ann. of Math. (2), 145, 1997, 2, 389-417.
- [11] De Carvalho, Andres, Lyubich, Michael and Martens, Marco, *Renormalization in the Hénon family. I. Universality but non-rigidity*, J. Stat. Phys., 121, 2005, 5-6, 611–669.

- [12] Gonchenko, Sergey, Turaev, Dmitry and Shilnikov, Leonid, *Homoclinic tangencies of arbitrarily high orders in conservative and dissipative two-dimensional maps*, Nonlinearity, 20, 2007, 241-275.
- [13] Ilyashenko, Yu. S. and Yakovenko, S. Yu., *Finitely smooth normal forms of local families of diffeomorphisms and vector fields*, Uspekhi Mat. Nauk, 46, 1991, 1(277), 3–39, 240.
- [14] Marmi, Stefano, *An Introduction To Small Divisors*, Preprint, arXiv:math/0009232 .
- [15] Mora, Leonardo and Viana, Marcelo, *Abundance of strange attractors*, Acta Math., 171, 1993,1, 1–71.
- [16] Newhouse, Sheldon E., *Diffeomorphisms with infinitely many sinks*, Topology, 13, 1974, 9–18.
- [17] Palis, Jacob and Takens, Floris, *Hyperbolicity and sensitive chaotic dynamics at homoclinic bifurcations*, Cambridge Studies in Advanced Mathematics, 35, Fractal dimensions and infinitely many attractors, Cambridge University Press, Cambridge, 1993, x+234.
- [18] Robinson, Clark, *Bifurcation to infinitely many sinks*, Comm. Math. Phys., 90, 1983, 3, 433–459.