

Astro2020 Science White Paper

Completing the Hydrogen Census in the Circumgalactic Medium at $z \sim 0$

- Thematic Areas:**
- Planetary Systems
 - Star and Planet Formation
 - Formation and Evolution of Compact Objects
 - Cosmology and Fundamental Physics
 - Stars and Stellar Evolution
 - Resolved Stellar Populations and their Environments
 - Galaxy Evolution
 - Multi-Messenger Astronomy and Astrophysics

Principal Author:

Name: D.J. Pisano

Institution: Dept. of Physics & Astronomy and the Gravitational Wave and Cosmology Center, West Virginia University

Email: djpisano@mail.wvu.edu

Phone: +1-304-293-4886

Co-authors: A. Fox, D. French (STScI), J.C. Howk, N. Lehner (University of Notre Dame), F.J. Lockman (Green Bank Observatory), K. Jones (NAIC)

Abstract:

Over the past decade, Lyman- α and metal line absorption observations have established the ubiquity of a gas-rich circumgalactic medium (CGM) around star-forming galaxies at $z \sim 0.2$ potentially tracing half of the missing baryonic mass within galaxy halos. Unfortunately, these observations only provide a statistical measure of the gas in the CGM and do not constrain the spatial distribution and kinematics of the gas. Furthermore, we have limited sensitivity to Lyman- α at $z \sim 0$ with existing instruments. As such, we remain ignorant of how this gas may flow from the CGM onto the disks of galaxies where it can fuel ongoing star-formation in the present day. Fortunately, 21-cm HI observations with radio telescopes can map HI emission providing both spatial and kinematic information for the CGM in galaxies at $z=0$. Observations with phased array feeds, radio cameras, on single-dish telescopes yield unmatched surface brightness sensitivity and survey speed. These observations can complete the census of HI in the CGM below $N_{HI} \lesssim 10^{17} \text{cm}^{-2}$ and constrain how gas accretion is proceeding in the local universe, particularly when used in concert with UV absorption line data.

1 Background

Great strides have been made in understanding the nature and evolution of galaxies over the past fifty years. We know that galaxies assemble their mass in a hierarchical manner by accreting smaller galaxies with their associated stars and dark matter and merging with other galaxies in a process that continues to the present day [1]. It is still unknown, however, how dark matter halos, and the galaxies contained therein, accrete the gas that they need to continue to form stars to the present day. Current theories suggest that there are three ways a galaxy can accrete gas. The most straight-forward is through the accretion of a satellite galaxy as part of the hierarchical assembly of a galaxy, but such gas-rich satellites are neither abundant enough nor contain enough gas to sustain star formation[2]. Alternatively, gas can flow onto galaxies in either a hot ($T \sim 10^6 \text{K}$) or cold ($T \lesssim 10^5 \text{K}$) phase [3, 4]. The hot mode involves gas falling onto galaxies in a quasi-spherical mode and is expected to be dominant for high mass galaxies in higher density environments in the present day. In contrast, the cold mode is more filamentary in nature and should dominate at high redshift and for low mass galaxies in lower density environments. While simulations disagree on the amount and exact phase of this accretion [5, 6], they all agree that accretion from the intergalactic medium through the circumgalactic medium (CGM) and onto galaxy disks should still be occurring today.

There is certainly evidence for ongoing accretion onto galaxies in the form of discrete, cold HI clouds [7, 8], which is likely tracing a larger, warm-hot ionized reservoir of gas [9]. Further evidence for cold accretion comes from Lyman-limit absorption systems with low metallicities associated with nearby galaxies [10, 11]. Absorption line studies, however, can only provide line-of-sight information and do not yield a complete picture of the gas through the CGM of individual galaxies. *In order to understand how gas is accreted onto galaxies in the present day, comprehensive surveys in both emission and absorption of the CGM is required.*

2 Current Observations of the CGM

To date, most of the exploration of the CGM has come through UV absorption line studies. The COS-HALOS project [12, 13] has used background quasars to study the Lyman- α absorption in the halos of low redshift galaxies. The project has found that HI absorption at $N_{HI} \gtrsim 10^{14} \text{cm}^{-2}$ is ubiquitous out to 150 kpc for star-forming galaxies and present in 75% of passive galaxies as well [12]. This cool CGM gas represents 25%-45% of the total baryon mass within the virial radius of the galaxy [13, 14]. Unfortunately, above $N_{HI} \sim 10^{16} \text{cm}^{-2}$ saturation of absorption lines makes it difficult to get an accurate measure of N_{HI} ; these are the Lyman Limit Systems. While below $N_{HI} \sim 10^{16-17} \text{cm}^{-2}$, HI absorption is common, particularly in the intergalactic medium, it has been impossible to image in 21-cm HI emission to date. Obtaining deeper HI emission observations is the only way forward: while Lyman- α absorption observations can reach very low column densities, their pencil-beam nature make it extremely difficult to reconstruct the full gas distribution and its kinematics. Furthermore, such observations are needed to measure N_{HI} at these column densities so that metallicities can be accurately determined. Such metallicity measurements are key to understanding if the gas in the CGM is infalling, pristine gas or enriched outflows.

Over the past decade, 21-cm HI emission observations have yielded great insights into the nature

of the CGM around nearby galaxies. Single-dish and interferometric observations have revealed both discrete HI clouds as well as diffuse HI structures that are related to previously unknown dwarf galaxies, tidal interactions or accretion events [15, 16, 17, 18, 19, 20, 21]. While such observations detect less than 10% more M_{HI} , this emission is tracing the more massive, dominant, ionized gas reservoir in the CGM of these galaxies [9]. A prime example of this are the extensive HI surveys of the M 31's CGM. [22] discovered a HI bridge between M 31 and M 33 with $N_{HI} \gtrsim 10^{17} \text{cm}^{-2}$ that they attributed to the cosmic web (Figure 1). Higher resolution observations with the Green Bank Telescope (GBT) have shown that this diffuse structure is actually comprised of discrete, higher N_{HI} clouds [23, 24]. At these HI column densities the gas clouds are mostly ionized [25], but they allow us to trace the morphology and kinematics of this reservoir. Furthermore, the CGM of M 31 itself is quite clumpy with HI covering fractions below 5% at $N_{HI} \sim 4 \times 10^{17} \text{cm}^{-2}$ [26]. As seen in Figure 2, these results are consistent with simulations, but are significantly lower than what was found from COS-HALOS [27]. This could be due to the unique properties of M 31 or represent the evolution of the CGM since $z \sim 0.2$, where most COS-HALOS galaxies reside. These results demonstrate the need for high spatial angular resolution HI surveys with excellent surface brightness sensitivity that can only be provided by large single-dish telescopes in concert with interferometers.

3 The role of 21-cm HI observations in the next decade

As can be seen from Figures 1 and 2, 21-cm HI observations of the CGM of even a single galaxy provide direct measurements of N_{HI} , independent of the optical depth of the gas, as well as the detailed morphology and kinematics of that gas. When combined with UV absorption line data, we can determine the metallicity of the CGM, which provides a strong constraint on the origin of the gas. To date, however, such deep ($N_{HI} \sim 10^{17} \text{cm}^{-2}$) HI emission observations have been limited to M 31. The filaments of gas associated with cold accretion are expected to have widths up to $\lesssim 25$ kpc [28], so spatial resolution is needed. M 31 is close enough that single-dish telescopes, like the GBT, can spatially resolve HI structures down to ~ 2 kpc in its CGM mitigating the effects of beam dilution. The GBT should be capable of resolving such filaments out to $D \sim 10$ Mpc, while Arecibo could do so out to ~ 30 Mpc. Interferometers provide better resolution and have excellent M_{HI} sensitivity, but lack the surface brightness sensitivity needed to detect such low- N_{HI} emission. These observations are valuable for detecting HI clouds around galaxies as demonstrated by HALOGAS [29]. Still to recover HI emission at low N_{HI} , as well as from compact sources, we need both single-dish and interferometric observations. If we wish to study how the properties of the CGM vary with galaxy mass and environment, we need to extend these observations to larger samples of galaxies beyond M 31.

In the next decade it will be possible to use existing single-dish telescopes, such as the GBT and Arecibo, outfitted with new phased array feeds (PAFs), or radio cameras, FLAG [30] and ALPACA, to make sensitive ($N_{HI} \lesssim 10^{17} \text{cm}^{-2}$) surveys covering the entire dark matter halo of ~ 100 galaxies within ~ 20 Mpc spanning a range of masses and environments. Due to the dramatic improvements in survey speed from PAFs, astronomers will be able to probe more diffuse gas around more galaxies. These data will be capable of resolving filamentary structures associated with cold accretion and will provide a complete census of the HI content of the CGM of these galaxies. By measuring the metallicity of these features and modeling their kinematics,

we will be able to identify ongoing accretion events. When combined with interferometer data, from the ngVLA for example, we will be able to trace accretion from the CGM directly on to the disks of galaxies.

The insight provided by such a survey will not be achievable without single-dish radio telescopes, as even the Square Kilometer Array will not achieve such excellent N_{HI} sensitivity. While FAST in China will have better sensitivity and resolution than Arecibo or the GBT, the large field of view achieved with PAFs on these telescopes will result in faster survey speeds.

Future UV studies with large-aperture space telescopes also have an important role to play in CGM studies in the next decade and beyond. UV facilities with multiplexing ability and higher sensitivity than Hubble/COS could be used to measure Lyman- α in multiple QSO sightlines in a given galaxy halo. This would allow studies of spatial variation, kinematic structure, and covering fraction of HI within individual halos. These observations would reach very low HI column densities (10^{13}cm^{-2}) and hence complement the 21-cm radio observations that probe higher HI column densities




Figure 1: Top: A map of the total HI emission associated with M 33 (lower left) and M 31 (upper right) from [22]. The contours are at $\log N_{HI} = 17.0, 17.3, 17.7, 18.0, 18.3, 18.7, 19.0, 19.3, 19.7, 20.0, 20.3,$ and 20.7 [cm^{-2}]. The beamsize is shown in the lower right of the panel. The box shows the region mapped by [24] with the GBT. Bottom: The GBT HI map from [24]. The contours are at $-1, 1, 2, 4, 6,$ and 10 times $5 \times 10^{17} \text{cm}^{-2}$. The beam size is the circle in the lower left of the image. Note that the HI structures detected by [22] are revealed to be much smaller, higher- N_{HI} features by [24], illustrating the critical importance of resolution and sensitivity.




Figure 2: Top: Locations of the AMIGA [26] GBT pointings relative to M31 and M33, where the axes are labeled with the impact parameter from the center of M31. Aside from two filled circles, representing detections of HI associated with the Magellanic Stream, all AMIGA observations yielded non-detections. Bottom: The cumulative covering fraction as a function of impact parameter of HI emission with $\log(N_{HI}) \geq 17.6$ as compared to absorption line data from COS-HALOS [27] (left), and high-velocity clouds [31] and simulations from [32]. While the AMIGA data [26] are consistent with simulations and the Milky Way high-velocity clouds, they are inconsistent with the COS-HALOS data. HI and UV absorption line observations of more nearby galaxies will help shed light on this discrepancy.

References

- [1] W. H. Press and P. Schechter. Formation of Galaxies and Clusters of Galaxies by Self-Similar Gravitational Condensation. *ApJ*, 187:425–438, February 1974.
- [2] E. M. Di Teodoro and F. Fraternali. Gas accretion from minor mergers in local spiral galaxies. *A&A*, 567:A68, July 2014.
- [3] D. Kereš, N. Katz, D. H. Weinberg, and R. Davé. How do galaxies get their gas? *MNRAS*, 363:2–28, October 2005.
- [4] D. Kereš, N. Katz, M. Fardal, R. Davé, and D. H. Weinberg. Galaxies in a simulated Λ CDM Universe - I. Cold mode and hot cores. *MNRAS*, 395:160–179, May 2009.
- [5] M. R. Joung, M. E. Putman, G. L. Bryan, X. Fernández, and J. E. G. Peek. Gas Accretion is Dominated by Warm Ionized Gas in Milky Way Mass Galaxies at $z \sim 0$. *ApJ*, 759:137, November 2012.
- [6] D. Nelson, M. Vogelsberger, S. Genel, D. Sijacki, D. Kereš, V. Springel, and L. Hernquist. Moving mesh cosmology: tracing cosmological gas accretion. *MNRAS*, 429:3353–3370, March 2013.
- [7] R. Sancisi, F. Fraternali, T. Oosterloo, and T. van der Hulst. Cold gas accretion in galaxies. *A&ARv*, 15:189–223, June 2008.
- [8] A. Fox and R. Davé, editors. *Gas Accretion onto Galaxies*, volume 430 of *Astrophysics and Space Science Library*, 2017.
- [9] N. Lehner and J. C. Howk. A Reservoir of Ionized Gas in the Galactic Halo to Sustain Star Formation in the Milky Way. *Science*, 334:955, November 2011.
- [10] J. T. Stocke, B. A. Keeney, and C. W. Danforth. A Direct Detection of Gas Accretion: The Lyman Limit System in 3C 232. *PASA*, 27:256–264, June 2010.
- [11] J. Ribaudó, N. Lehner, J. C. Howk, J. K. Werk, T. M. Tripp, J. X. Prochaska, J. D. Meiring, and J. Tumlinson. Evidence for Cold Accretion: Primitive Gas Flowing onto a Galaxy at $z \sim 0.274$. *ApJ*, 743:207, December 2011.
- [12] J. Tumlinson, C. Thom, J. K. Werk, J. X. Prochaska, T. M. Tripp, N. Katz, R. Davé, B. D. Oppenheimer, J. D. Meiring, A. B. Ford, J. M. O’Meara, M. S. Peeples, K. R. Sembach, and D. H. Weinberg. The COS-Halos Survey: Rationale, Design, and a Census of Circumgalactic Neutral Hydrogen. *ApJ*, 777:59, November 2013.
- [13] J. K. Werk, J. X. Prochaska, J. Tumlinson, M. S. Peeples, T. M. Tripp, A. J. Fox, N. Lehner, C. Thom, J. M. O’Meara, A. B. Ford, R. Bordoloi, N. Katz, N. Tejos, B. D. Oppenheimer, R. Davé, and D. H. Weinberg. The COS-Halos Survey: Physical Conditions and Baryonic Mass in the Low-redshift Circumgalactic Medium. *ApJ*, 792:8, September 2014.

- [14] J. Tumlinson, M. S. Peeples, and J. K. Werk. The Circumgalactic Medium. *ARA&A*, 55:389–432, August 2017.
- [15] K. M. Hess, D. J. Pisano, E. M. Wilcots, and J. N. Chengalur. Anomalous H I in NGC 2997. *ApJ*, 699:76–88, July 2009.
- [16] D. J. Pisano, D. G. Barnes, L. Staveley-Smith, B. K. Gibson, V. A. Kilborn, and K. C. Freeman. An H I Survey of Six Local Group Analogs. II. H I Properties of Group Galaxies. *ApJS*, 197:28, December 2011.
- [17] J. C. Mihos, K. M. Keating, K. Holley-Bockelmann, D. J. Pisano, and N. E. Kassim. The H I Environment of the M101 Group. *ApJ*, 761:186, December 2012.
- [18] D. J. Pisano. Green Bank Telescope Observations of Low Column Density H I around NGC 2997 and NGC 6946. *AJ*, 147:48, March 2014.
- [19] W. J. G. de Blok, K. M. Keating, D. J. Pisano, F. Fraternali, F. Walter, T. Oosterloo, E. Brinks, F. Bigiel, and A. Leroy. A low H I column density filament in NGC 2403: signature of interaction or accretion. *A&A*, 569:A68, September 2014.
- [20] N. M. Pingel, D. J. Pisano, G. Heald, T. H. Jarrett, W. J. G. de Blok, G. I. G. Józsa, E. Jütte, R. J. Rand, T. Oosterloo, and B. Winkel. A GBT Survey of the HALOGAS Galaxies and Their Environments. I. Revealing the Full Extent of H I around NGC 891, NGC 925, NGC 4414, and NGC 4565. *ApJ*, 865:36, September 2018.
- [21] A. Sorgho, C. Carignan, D. J. Pisano, T. Oosterloo, W. J. G. de Blok, M. Korsaga, N. M. Pingel, A. Sardone, S. Goedhart, S. Passmoor, A. Dikgale, and S. K. Sirothia. Early observations of the MHONGOOSE galaxies: getting ready for MeerKAT. *MNRAS*, 482:1248–1269, January 2019.
- [22] R. Braun and D. A. Thilker. The WSRT wide-field H I survey. II. Local Group features. *A&A*, 417:421–435, April 2004.
- [23] S. A. Wolfe, D. J. Pisano, F. J. Lockman, S. S. McGaugh, and E. J. Shaya. Discrete clouds of neutral gas between the galaxies M31 and M33. *Nature*, 497:224–226, May 2013.
- [24] S. A. Wolfe, F. J. Lockman, and D. J. Pisano. Sensitive 21cm Observations of Neutral Hydrogen in the Local Group near M31. *ApJ*, 816:81, January 2016.
- [25] P. Maloney. Sharp edges to neutral hydrogen disks in galaxies and the extragalactic radiation field. *ApJ*, 414:41–56, September 1993.
- [26] J. C. Howk, C. B. Wotta, M. A. Berg, N. Lehner, F. J. Lockman, Z. Hafen, D. J. Pisano, C.-A. Faucher-Giguère, B. P. Wakker, J. X. Prochaska, S. A. Wolfe, J. Ribaudó, K. A. Barger, L. Corlies, A. J. Fox, P. Guhathakurta, E. B. Jenkins, J. Kalirai, J. M. O’Meara, M. S. Peeples, K. R. Stewart, and J. Strader. Project AMIGA: A Minimal Covering Factor for Optically Thick Circumgalactic Gas around the Andromeda Galaxy. *ApJ*, 846:141, September 2017.

- [27] J. X. Prochaska, J. K. Werk, G. Worseck, T. M. Tripp, J. Tumlinson, J. N. Burchett, A. J. Fox, M. Fumagalli, N. Lehner, M. S. Peebles, and N. Tejos. The COS-Halos Survey: Metallicities in the Low-redshift Circumgalactic Medium. *ApJ*, 837:169, March 2017.
- [28] A. Popping, R. Davé, R. Braun, and B. D. Oppenheimer. The simulated H I sky at low redshift. *A&A*, 504:15–32, September 2009.
- [29] G. Heald, G. Józsa, P. Serra, L. Zschaechner, R. Rand, F. Fraternali, T. Oosterloo, R. Walterbos, E. Jütte, and G. Gentile. The Westerbork Hydrogen Accretion in Local GALaxieS (HALOGAS) survey. I. Survey description and pilot observations. *A&A*, 526:A118, February 2011.
- [30] D. A. Roshi, W. Shillue, B. Simon, K. F. Warnick, B. Jeffs, D. J. Pisano, R. Prestage, S. White, J. R. Fisher, M. Morgan, R. Black, M. Burnett, J. Diao, M. Ruzindana, V. van Tonder, L. Hawkins, P. Marganian, T. Chamberlin, J. Ray, N. M. Pingel, K. Rajwade, D. R. Lorimer, A. Rane, J. Castro, W. Groves, L. Jensen, J. D. Nelson, T. Boyd, and A. J. Beasley. Performance of a Highly Sensitive, 19-element, Dual-polarization, Cryogenic L-band Phased-array Feed on the Green Bank Telescope. *AJ*, 155:202, May 2018.
- [31] P. Richter. Cold Gas Accretion by High-velocity Clouds and Their Connection to QSO Absorption-line Systems. *ApJ*, 750:165, May 2012.
- [32] Z. Hafen, C.-A. Faucher-Giguère, D. Anglés-Alcázar, D. Kereš, R. Feldmann, T. K. Chan, E. Quataert, N. Murray, and P. F. Hopkins. Low-redshift Lyman limit systems as diagnostics of cosmological inflows and outflows. *MNRAS*, 469:2292–2304, August 2017.