

Computational Astrocyence: Astrocytes encode inhibitory activity into the frequency and spatial extent of their calcium elevations

Ioannis E. Polykretis
Computational Brain Lab
Rutgers University
Piscataway, NJ, USA

Vladimir A. Ivanov
Computational Brain Lab
Rutgers University
Piscataway, NJ, USA

Konstantinos P. Michmizos
Computational Brain Lab
Rutgers University
Piscataway, NJ, USA
konstantinos.michmizos@cs.rutgers.edu

Abstract—Deciphering the complex interactions between neurotransmission and astrocytic Ca^{2+} elevations is a target promising a comprehensive understanding of brain function. While the astrocytic response to excitatory synaptic activity has been extensively studied, how inhibitory activity results to intracellular Ca^{2+} waves remains elusive. In this study, we developed a compartmental astrocytic model that exhibits distinct levels of responsiveness to inhibitory activity. Our model suggested that the astrocytic coverage of inhibitory terminals defines the spatial and temporal scale of their Ca^{2+} elevations. Understanding the interplay between the synaptic pathways and the astrocytic responses will help us identify how astrocytes work independently and cooperatively with neurons, in health and disease.

Index Terms—Astrocytes, Computational Modeling, Ca^{2+} elevations, $GABA_B$ Receptors, Inositol Triphosphate (IP_3)

I. INTRODUCTION

Breaking up the neuronal monopoly of studying the brain, the implication of glia in learning and cognition is now well established [1]. While the contribution of astrocytes, the most abundant type of glial cells, to synaptic modulation of excitatory connections has been extensively studied [2], their association with inhibitory synaptic transmission still remains elusive. Inhibition is crucial for the function of neural networks, as it has been involved in generating feedforward and feedback loops [3] and preventing hyper-excitability [4]. Interestingly, astrocytes are now known to detect firing activity of inhibitory neurons, which release gamma aminobutyric acid (GABA), and respond to it through intracellular Ca^{2+} elevations of distinct temporal (repetitive short or prolonged) and spatial (highly localized or expanded) scales [5]–[7]. These Ca^{2+} waves have recently been strongly related to mental dysfunctions [8]–[11]. In fact, most brain diseases are now considered as conditions where both neurons and astrocytes are affected [12]. In support of this notion, numerous animal model studies [13]–[16] report alterations of astrocytic activity in a diseased brain state. However, the exact astrocytic mechanisms for relating the inhibitory synaptic transmission

to the temporal and spatial characteristics of Ca^{2+} elevations are not known.

In our attempt to decipher the role of inhibitory synaptic transmission in healthy and impaired astrocytic functions, we modified our recent biologically plausible compartmental model of an astrocyte [17] to include $GABA_B$ receptors, in alignment with experimental results [5], [6]. Our modeling results suggested that astrocytes exhibit intensified responsiveness to short trains of pulses, increasing the frequency of their Ca^{2+} elevations. Our results suggested that the extent of the astrocytic coverage of inhibitory axonal terminals is a decisive factor for the distinct spatial and temporal extent of astrocytic Ca^{2+} waves. Specifically, limited coverage of active inhibitory terminals induced short and repetitive Ca^{2+} elevations, which remain spatially constrained. As the coverage of terminals increased, an isolated Ca^{2+} increase superimposed to Ca^{2+} elevations from neighboring activity and triggered less frequent, but spatially more expanded Ca^{2+} elevations.

II. METHODS

We modified our 2-dimensional astrocytic model replicating the stellate structure of astrocytes and emulating the spatial allocation of the subcellular organelles in different compartments [17], [18]. Briefly, our model comprised of a somatic compartment, which extended to a number of thick branches (Fig. 1). These two compartments included both types of Endoplasmatic Reticulum (ER) reported in astrocytes; one store had receptors sensitive to Inositol Triphosphate (IP_3) and the other to Ryanodine [19]. The thick branches ramified into thinner ones, which had only IP_3 -sensitive ER [20] and finally gave rise to thin tendrils, void of organelles, the perisynaptic astrocytic processes (PAPs). The assemblies of several PAPs form the now known as astrocytic microdomains (Fig. 1, inset). In the PAPs astrocytes contacted neuronal synapses, sensed the neurotransmitter spillover outside the synaptic cleft through their receptors and triggered intracellular procedures.

The neurotransmitter receptors of astrocytes in this study are the GABA receptors (GABARs). Astrocytes have both

This work is partially funded by the Rutgers Brain Health Institute (FP00014150). Ioannis E. Polykretis is partially funded by the Onassis Foundation Scholarship.




Fig. 1. Geometry, compartments and their respective organelles in our 2-dimensional model of an astrocyte. $GABA_B$ receptors are added to our previous model [17].

metabotropic ($GABA_BRs$) and ionotropic ($GABA_ARs$) receptors [21]. Although there is some controversy about which type of receptors gives rise to Ca^{2+} elevations in astrocytes [22], an increasing amount of evidence suggests that the slower and spatially expanded Ca^{2+} waves are induced by $GABA_BRs$ [6]. That is why we focused on this type of G-protein-coupled receptors. This coupling of $GABA_BRs$ triggers the intracellular enzymatic production of IP_3 [6], which then induces Ca^{2+} release from the intracellular stores. We used the mathematical model initially presented in [23] and further modified in [17] to simulate this pathway. We used a parameter v_{GABA} to model the rate of agonist-dependent IP_3 production due to GABA. The numerical value that most accurately replicates the experimental findings [5] is $6.7 \mu M/s$.

We used the Leaky Integrate and Fire (LIF) neuronal model to simulate the spiking activity of the interneurons. We used a membrane resistance $R = 600M\Omega$ and a time constant $\tau = 60ms$. We adjusted the stimulation current to reproduce a widely accepted stimulation pattern that has been found to induce Ca^{2+} elevations in astrocytes [5]. In this pattern the

interneurons are driven to fire a constant total number of spikes grouped in 200, 100, 66 and 33 trains of 1, 2, 3, and 6 spikes, respectively. The inter-train interval is kept constant at 2s. Specifically, we used 0.5, 0.7, 1 and 1.7 nA to drive the interneurons and force them to fire the trains of 1, 2, 3 and 6 spikes respectively.

To translate the spiking activity of the interneurons to the amount of neurotransmitter sensed by the astrocytes, we modified the model presented in [24]. Given a spiking pattern, the model predicts the amount of the neurotransmitter released inside the synaptic cleft. However, the PAP senses only a percentage of this amount that spills out of the cleft. To account for that spillover, we used only a fraction of that amount of neurotransmitter that the model predicts. This amount of stimulant was subsequently fed into the astrocytic model to trigger IP_3 production and, consequently, Ca^{2+} release from the ER that can be seen as Ca^{2+} elevations.

The 33-6 protocol [5] was used for simulation brevity in all experiments, where the reproduction of another protocol was not imperative.

III. RESULTS

Our astrocytic model exhibited the GABA-induced Ca^{2+} waves that have been reported in astrocytes [5], [7]. We observed a wide range of Ca^{2+} waves emerging through the intracellular pathway described above.

First, we examined the responsiveness of the cell to different types of stimulus. We used different stimulation protocols to drive our model and the results are shown in Figure 2. We stimulated all the PAPs of a single microdomain, assuming that they all contact the axonal terminals of a single spiking interneuron or multiple distinct interneurons driven simultaneously (Fig.2, left, bottom inset). Our results suggested that the 66-3 protocol (Fig.2, right, bottom) produced a more prominent response than the respective response to the 33-6 protocol (Fig.2, left, bottom). The 66-3 protocol gave rise to multiple Ca^{2+} elevations, while the 33-6 protocol gave rise to a single elevation during the stimulation period. As seen in Fig.2 (top, insets), every spike of a single train released an amount of the available neurotransmitters. When the trains included more than three spikes, the amount of released neurotransmitter after the third spike was inconsequential for astrocytes, resulting in weaker Ca^{2+} responses. We also tested the 200-1 and 100-2 stimulation protocols and the response of the cell was always less prominent (results not shown). Therefore, our model followed closely the experimental findings that show maximal astrocytic responsiveness to the 66-3 protocol ([5], Fig.6, c).

Elevations of different spatial extent and temporal duration in neighboring cells are described ([5], Fig.6, b). We simultaneously stimulated multiple microdomains and examined whether such elevations can be observed. As seen in Fig.3, stimulation of multiple inhibitory terminals induced Ca^{2+} elevations that were temporally extended and spatially expanded (Fig.3 bottom, green curve).




Fig. 2. Astrocytic activity in response to two stimulation patterns. The GABA stimulation patterns (top) induce IP₃ production (middle), which triggers Ca²⁺ release (bottom). Maximal response in terms of number of Ca²⁺ elevations and their frequency is achieved when the spikes are grouped in 66 trains of 3 (right column) and not in 33 trains of 6 (left column).

IV. DISCUSSION

Our compartmental astrocytic model suggested possible relationships in space and time between the inhibitory synaptic transmission and intracellular Ca²⁺ elevations. Specifically, the level and the frequency of astrocytic Ca²⁺ activity depended on the stimulation pattern of the neurons. Moreover, the spatial extent and temporal duration of Ca²⁺ waves depended on the number of inhibitory terminals that an astrocyte covered.

First, our cell model aligned with experimental results [5], as it exhibited intensified responsiveness to short trains of inhibitory spikes by increasing the frequency of its Ca²⁺ elevations. Sparse distribution of the spikes or grouping in longer trains did not induce significant Ca²⁺ elevations. We propose that this strengthened response was due to the maximal exploitation of synaptic resources by the short spike trains. Conversely, sparse spikes could not fully activate the synaptic resources and longer trains depleted them, inducing weaker responses. A limited number of active synapses induced short and repetitive, spatially compartmentalized Ca²⁺ elevations. The spatial extension of these elevations was confined to the stimulated microdomain (Fig.2, bottom, red and purple curves), since the thicker branches were not affected by

the wave (Fig.2, bottom, green curve). However, the spatial resolution of the experimental study was not high enough, so that we could test this observation ([5], Fig.6, a).

Our results exhibited spatially expanded and temporally extended Ca²⁺ elevations, similar to the experimental findings ([5], Fig.6, b). Our model suggested that the number of inhibitory terminals that an astrocyte contacts is a decisive factor for the spatial extent and the temporal duration of astrocytic Ca²⁺ elevations. Limited coverage of terminals induced brief, repetitive and localized Ca²⁺ elevations. However, increasing the number of active synapses that an astrocyte ensheathes, induced simultaneous, isolated Ca²⁺ elevations in multiple microdomains. Their superposition was sufficient to trigger the Ryanodine-sensitive ER in the thicker branches and give rise to elevations that were prolonged in time, spatially expanded and had lower frequency (Fig.3 bottom, green curve).

In our previous work, we showed that astrocytes, when stimulated by excitatory neurotransmission, can induce neuronal synchronization [17] and spatial clustering of synaptic plasticity [18]. This study complements our previous results by showing that inhibitory neurotransmission may also shape Ca²⁺ elevations in astrocytes. Taken together, our approaches may help further our understanding on the emergence of astrocytic Ca²⁺ responses to the most important synaptic




Fig. 3. Prolonged and spatially expanded Ca^{2+} elevations observed with extensive astrocytic coverage of synapses. Astrocytic coverage of multiple spiking inhibitory terminals induces temporally extended and spatially expanded Ca^{2+} elevations (bottom, green).

pathways, as well as their interaction.

Overall, our computational work aims to study how astrocytes may work independently and cooperatively with neurons in health and disease.

REFERENCES

- [1] Fields, R.D. et al, 2014. Glial biology in learning and cognition. *The neuroscientist*, 20(5), pp.426-431.
- [2] Araque, A., Parpura, V., Sanzgiri, R.P. and Haydon, P.G., 1998. Glutamate-dependent astrocyte modulation of synaptic transmission between cultured hippocampal neurons. *European Journal of Neuroscience*, 10(6), pp.2129-2142.
- [3] Buzsáki, G., 1984. Feed-forward inhibition in the hippocampal formation. *Progress in neurobiology*, 22(2), pp.131-153.
- [4] Cui, Y. et al, 2018. Astroglial Kir4.1 in the lateral habenula drives neuronal bursts in depression. *Nature*, 554(7692), p.323.
- [5] Kang, J., Jiang, L., Goldman, S.A. and Nedergaard, M., 1998. Astrocyte-mediated potentiation of inhibitory synaptic transmission. *Nature neuroscience*, 1(8), p.683.
- [6] Meier, S.D., Kafitz, K.W. and Rose, C.R., 2008. Developmental profile and mechanisms of GABA-induced calcium signaling in hippocampal astrocytes. *Glia*, 56(10), pp.1127-1137.
- [7] Nilsson, M., Eriksson, P.S., Rönnbäck, L. and Hansson, E., 1993. GABA induces Ca^{2+} transients in astrocytes. *Neuroscience*, 54(3), pp.605-614.
- [8] Tian, G.F. et al, 2005. An astrocytic basis of epilepsy. *Nature medicine*, 11(9), p.973.
- [9] Wetherington, J., Serrano, G. and Dingledine, R., 2008. Astrocytes in the epileptic brain. *Neuron*, 58(2), pp.168-178.

- [10] Verkhratsky, A., Olabarria, M., Noristani, H.N., Yeh, C.Y. and Rodriguez, J.J., 2010. Astrocytes in Alzheimer's disease. *Neurotherapeutics*, 7(4), pp.399-412.
- [11] Del Zoppo, G.J., 2006. Stroke and neurovascular protection. *n engl j med*, 354(6), pp.553-555.
- [12] Barres, B.A., 2008. The mystery and magic of glia: a perspective on their roles in health and disease. *Neuron*, 60(3), pp.430-440.
- [13] Seifert, G., Carmignoto, G. and Steinhuser, C., 2010. Astrocyte dysfunction in epilepsy. *Brain research reviews*, 63(1-2), pp.212-221.
- [14] Jabs, R., Seifert, G. and Steinhuser, C., 2008. Astrocytic function and its alteration in the epileptic brain. *Epilepsia*, 49, pp.3-12.
- [15] Oberheim, N.A. et al, 2008. Loss of astrocytic domain organization in the epileptic brain. *Journal of Neuroscience*, 28(13), pp.3264-3276.
- [16] Jo, S. et al, 2014. GABA from reactive astrocytes impairs memory in mouse models of Alzheimer's disease. *Nature medicine*, 20(8), p.886.
- [17] Polykretis, I., Ivanov, V. and Michmizos, K.P., 2018, July. A Neural-Astrocytic Network Architecture: Astrocytic calcium waves modulate synchronous neuronal activity. In *Proceedings of the International Conference on Neuromorphic Systems* (p. 6). ACM.
- [18] Polykretis, I., Ivanov, V. and Michmizos, K.P., 2018, December. The Astrocytic Microdomain as a Generative Mechanism for Local Plasticity. In *International Conference on Brain Informatics* (pp. 153-162). Springer, Cham.
- [19] Golovina, V.A. and Blaustein, M.P., 1997. Spatially and functionally distinct Ca^{2+} stores in sarcoplasmic and endoplasmic reticulum. *Science*, 275(5306), pp.1643-1648.
- [20] Straub, S.V., Bonev, A.D., Wilkerson, M.K. and Nelson, M.T., 2006. Dynamic inositol trisphosphate-mediated calcium signals within astrocytic endfeet underlie vasodilation of cerebral arterioles. *The Journal of general physiology*, 128(6), pp.659-669.
- [21] Losi, G., Mariotti, L. and Carmignoto, G., 2014. GABAergic interneuron to astrocyte signalling: a neglected form of cell communication in the brain. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 369(1654), p.20130609.
- [22] De Pittá, M., Ben-Jacob, E. and Berry, H., 2019. G protein-coupled receptor-mediated calcium signaling in astrocytes. In *Computational Glioscience* (pp. 115-150). Springer, Cham.
- [23] De Pittá, M., Goldberg, M., Volman, V., Berry, H. and Ben-Jacob, E., 2009. Glutamate regulation of calcium and IP_3 oscillating and pulsating dynamics in astrocytes. *Journal of biological physics*, 35(4), pp.383-411.
- [24] Tsodyks, M.V. and Markram, H., 1997. The neural code between neocortical pyramidal neurons depends on neurotransmitter release probability. *Proceedings of the National Academy of Sciences*, 94(2), pp.719-723.