A More General Theory of Static Approximations for Conjunctive Queries

Pablo Barceló \cdot Miguel Romero \cdot Thomas Zeume

Received: date / Accepted: date

Abstract Conjunctive query (CQ) evaluation is NP-complete, but becomes tractable for fragments of bounded hypertreewidth. Approximating a hard CQ by a query from such a fragment can thus allow for an efficient approximate evaluation. While underapproximations (i.e., approximations that return correct answers only) are well-understood, the dual notion of overapproximations (i.e., approximations that return complete – but not necessarily sound – answers), and also a more general notion of approximation based on the symmetric difference of query results, are almost unexplored. In fact, the decidability of the basic problems of evaluation, identification, and existence of those approximations has been open.

This article establishes a connection between overapproximations and existential pebble games that allows for studying such problems systematically. Building on this connection, it is shown that the evaluation and identification problem for overapproximations can be solved in polynomial time. While the

Barceló is funded by Millennium Institute for Foundational Research on Data and Fondecyt Grant 1170109. Zeume acknowledges the financial support by the European Research Council (ERC), grant agreement No 683080. Romero and Zeume thank the Simons Institute for the Theory of Computing for hosting them. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 714532). The paper reflects only the authors' views and not the views of the ERC or the European Commission. The European Union is not liable for any use that may be made of the information contained therein.

P. Barceló

DCC, University of Chile & IMFD Chile

Tel.: +56229784813

E-mail: pbarcelo@dcc.uchile.cl

M. Romero

Department of Computer Science, University of Oxford

 $\hbox{E-mail: miguel.romero@cs.ox.ac.uk}$

T. Zeume TU Dortmund

 $\hbox{E-mail: thomas.zeume@cs.tu-dortmund.de}\\$

general existence problem remains open, the problem is shown to be decidable in 2EXPTIME over the class of acyclic CQs and in PTIME for Boolean CQs over binary schemata. Additionally we propose a more liberal notion of overapproximations to remedy the known shortcoming that queries might not have an overapproximation, and study how queries can be overapproximated in the presence of tuple generating and equality generating dependencies.

The techniques are then extended to symmetric difference approximations and used to provide several complexity results for the identification, existence, and evaluation problem for this type of approximations.

Keywords conjunctive queries \cdot hypertreewidth \cdot approximations \cdot existential pebble game

1 Introduction

Due to the growing number of scenarios in which exact query evaluation is infeasible – e.g., when the volume of the data being queried is very large, or when queries are inherently complex – approximate query answering has become an important area of study in databases (see, e.g. [23,32,36,19,20]). Here we focus on approximate query answering for the fundamental class of conjunctive queries (CQs).

Exact query evaluation for CQs, that is, determining whether a tuple \bar{a} is contained in the result of a query q on a database \mathcal{D} , is NP-complete. It is known that the complexity of evaluation of a CQ depends on its degree of acyclicity, which can be formalized using different notions. One of the most general and well-studied such notions corresponds to generalized hypertreewidth [25]. Notably, the classes of CQs of bounded generalized hypertreewidth can be evaluated in polynomial time (see [24] for a survey).

Following recent work on approximate query answering for CQs and some related query languages [7,8], we study approximation of CQs by queries of bounded generalized hypertreewidth. If a CQ can be approximated by such a restricted query, this provides a certificate for an efficient approximation of the evaluation problem. It is worth noticing that the approximations studied here are *static* in the sense that they depend only on the CQ q and not on the underlying database \mathcal{D} . This has clear benefits in terms of the cost of the approximation process, as q is often orders of magnitude smaller than \mathcal{D} and an approximation that has been computed once can be used for all databases. Moreover, it allows us to construct a principled approach to CQ approximation based on the well-studied notion of CQ containment [11]. Recall that a CQ q is contained in a CQ q', written $q \subseteq q'$, if $q(\mathcal{D}) \subseteq q'(\mathcal{D})$ over each database \mathcal{D} . This notion constitutes the theoretical basis for the study of several CQ optimization problems [1].

Before stating our contributions, we recall the precise notions of approximation under consideration in this article, as well as the algorithmic problems of interest.

As mentioned above, we are interested in approximating CQs by queries from the class $\mathsf{GHW}(k)$ of CQs of generalized hypertreewidth with width at most k, for some $k \geq 1$. Intuitively, an approximation of a CQ q is a query $q' \in \mathsf{GHW}(k)$ whose result when evaluated on a database \mathcal{D} is so close to the result of q, that no result of another query from $\mathsf{GHW}(k)$ is closer. A formalization of this notion was first introduced in [6], based on the following partial order \sqsubseteq_q over the CQs in $\mathsf{GHW}(k)$: if $q', q'' \in \mathsf{GHW}(k)$, then $q' \sqsubseteq_q q''$ iff over every database \mathcal{D} the symmetric difference¹ between $q(\mathcal{D})$ and $q''(\mathcal{D})$ is contained in the symmetric difference between $q(\mathcal{D})$ and $q'(\mathcal{D})$.Intuitively, $q' \sqsubseteq_q q''$ if q'' is a better $\mathsf{GHW}(k)$ -approximation of q than q'. The $\mathsf{GHW}(k)$ -approximations of q then correspond to maximal elements with respect to \sqsubseteq_q among a distinguished class of CQs in $\mathsf{GHW}(k)$.

Three notions of approximation were introduced in [6], by imposing different "reasonable" conditions on such a class. These are:

- Underapproximations: In this case we look for approximations in the set of CQs q' in GHW(k) that are contained in q, i.e., $q' \subseteq q$. This ensures that the evaluation of such approximations always produces correct (but not necessarily complete) answers to q. A GHW(k)-underapproximation of q is then a CQ q' amongst these CQs that is maximal with respect to the partial order defined by \sqsubseteq_q . Noticeably, the latter coincides with being maximal with respect to the containment partial order \subseteq among the CQs in GHW(k) that are contained in q; i.e., no other CQ in such a set strictly contains q'.
- Overapproximations: This is the dual notion of underapproximations, in which we look for minimal elements in the class of CQs q' in $\mathsf{GHW}(k)$ that contain q, i.e., $q \subseteq q'$. Hence, $\mathsf{GHW}(k)$ -overapproximations produce complete (but not necessarily correct) answers to q.
- Symmetric difference approximations: While underapproximations must be contained in the original query, and overapproximations must contain it, symmetric difference approximations impose no constraints on approximations with respect to the partial order \subseteq . Thus, the symmetric difference $\mathsf{GHW}(k)$ -approximations of q (or $\mathsf{GHW}(k)$ - Δ -approximations from now on) are the maximal CQs in $\mathsf{GHW}(k)$ with respect to \sqsubseteq_q .

The approximations presented above provide "qualitative" guarantees for evaluation, as they are as close as possible to q among all CQs in $\mathsf{GHW}(k)$ of a certain kind. In particular, under- and overapproximations are dual notions which provide lower and upper bounds for the exact evaluation of a CQ, while Δ -approximations can give us useful information that complements the one provided by under- and overapproximations. Then, in order to develop a robust theory of bounded generalized hypertreewidth static approximations for CQs, it is necessary to have a good understanding of all three notions.

The notion of underapproximation is by now well-understood [7]. Indeed, it is known that for each $k \geq 1$ the $\mathsf{GHW}(k)$ -underapproximations have good

¹ Recall that the symmetric difference between sets A and B is $(A \setminus B) \cup (B \setminus A)$.

properties that justify their application: (a) they always exist, and (b) evaluating all GHW(k)-underapproximations of a CQ q over a database \mathcal{D} is fixed-parameter tractable with the size of q as parameter. This is an improvement over general CQ evaluation for which the latter is believed not to hold [39].

In turn, the notions of overapproximations and Δ -approximations, while introduced in [6], are much less understood. In fact, no general tools have been identified so far for studying the decidability of basic problems such as:

- Existence: Does CQ q have a $\mathsf{GHW}(k)$ -overapproximation (or $\mathsf{GHW}(k)$ - Δ -approximation)?
- Identification: Is it the case that q' is a $\mathsf{GHW}(k)$ -overapproximation (resp., $\mathsf{GHW}(k)$ - Δ -approximation) of q?
- Evaluation: Given a CQ q, a database \mathcal{D} , and a tuple \bar{a} in \mathcal{D} , is it the case that $\bar{a} \in q'(\mathcal{D})$, for some $\mathsf{GHW}(k)$ -overapproximation (resp., $\mathsf{GHW}(k)$ - Δ -approximation) q' of q?

Partial results were obtained in [6], but based on ad-hoc tools. It has also been observed that some CQs have no GHW(k)-overapproximations (in contrast to underapproximations, that always exist), which was seen as a negative result.

Contributions. We develop tools for the study of overapproximations and Δ -approximations. While we mainly focus on the former, we provide a detailed account of how our techniques can be extended to deal with the latter. In the context of $\mathsf{GHW}(k)$ -overapproximations, we apply our tools to pinpoint the complexity of evaluation and identification, and make progress in the problem of existence. We also study when overapproximations do not exist and suggest how this can be alleviated. Finally, we study overapproximations in the presence of integrity constraints. Our contributions are as follows:

- 1. Link to existential pebble games. We establish a link between $\mathsf{GHW}(k)$ overapproximations and existential pebble games [34]. Such games have
 been used to show that several classes of CQs of bounded width can be
 evaluated efficiently [15,13]. Using the fact that the existence of winning
 conditions in the existential pebble game can be checked in polynomial
 time [13], we show that the identification and evaluation problems for $\mathsf{GHW}(k)$ -overapproximations are tractable.
- 2. A more liberal notion of overapproximation. We observe that non-existence of overapproximations is due to the fact that in some cases overapproximations require expressing conjunctions of infinitely many atoms. By relaxing our notion, we get that each CQ q has a (potentially infinite) GHW(k)-overapproximation q'. This q' is unique (up to equivalence). Further, it can be evaluated efficiently in spite of being potentially infinite by checking a winning condition for the existential k-pebble game on q and \mathcal{D} .
- 3. Existence of overapproximations. It is still useful to check if a CQ q has a finite GHW(k)-overapproximation q', and compute it if possible. This might allow us to optimize q' before evaluating it. There is also a difference in complexity, as existential pebble game techniques are PTIME-complete

	Existence?	Unique?	Evaluation	Existence check
GHW(k)-underapp.	always	not always	NP-hard	N/A
GHW(k)-overapp.	not always	always	PTIME*	For $k = 1$:
				2Exptime*
				PTIME* on binary schemas
				For $k > 1$: Open

Fig. 1 Summary of results on under- and overapproximations of bounded generalized hypertreewidth (in the absence of constraints). The complexity of identification coincides with that of evaluation in both cases. New results are marked with (*). All remaining results follow from [6,7].

in general [33], and thus inherently sequential, while evaluation of CQs in GHW(k) is highly parallelizable (Gottlob et al. [25]).

By exploiting automata techniques, we show that checking if a CQ q has a (finite) GHW(1)-overapproximation q' is in 2Exptime. Also, when such q' exists it can be computed in 3Exptime. This is important since GHW(1) coincides with the well-known class of acyclic CQs [40]. If the arity of the schema is fixed, these bounds drop to Exptime and 2Exptime, respectively. Also, we look at the case of binary schemas, which are for instance used in graph databases [4] and description logics [2]. In this case, we show that for Boolean CQs, GHW(1)-overapproximations can be computed efficiently via a greedy algorithm. This is optimal, as over ternary schemas we prove an exponential lower bound for the size of GHW(1)-overapproximations.

We do not know if the existence problem is decidable for k > 1. However, we show that it can be recast as an unexplored boundedness condition for the existential pebble game. Understanding the decidability boundary for such conditions is often difficult [38,9].

4. Overapproximations under constraints. It has been observed that semantic information about the data, in the form of integrity constraints, enriches the quality of approximations [5]. This is based on the fact that approximations are now defined over a restricted set of databases; namely, those that satisfy the constraints.

We study $\mathsf{GHW}(k)$ -overapproximations for the practical classes of equality-generating dependencies (egds), which subsume functional dependencies, and tuple-generating dependencies (tgds), which subsume inclusion dependencies. By extending our previously derived techniques, we show that each $\mathsf{CQ}\ q$ admits an infinite $\mathsf{GHW}(k)$ -overapproximation under a set of constraints \varSigma , in any of the following cases: \varSigma consists exclusively of egds, or \varSigma is a set of guarded tgds [10]. Recall that the latter corresponds to a well-studied extension of the class of inclusion dependencies. Such an infinite $\mathsf{GHW}(k)$ -overapproximation can be evaluated in polynomial time for the case of functional dependencies and guarded tgds (and so, for inclusion dependencies), and by a fixed-parameter tractable algorithm for egds.

Figure 1 shows a summary of these results in comparison with previously known results about underapproximations (in the absence of constraints).

Our contributions for $\mathsf{GHW}(k)$ - Δ -approximations are as follows. As a preliminary step, we show that $\mathsf{GHW}(k)$ -under- and $\mathsf{GHW}(k)$ -overapproximations are particular cases of $\mathsf{GHW}(k)$ - Δ -approximations, but not vice versa. Afterwards, as for $\mathsf{GHW}(k)$ -overapproximations, we provide a link between $\mathsf{GHW}(k)$ - Δ -overapproximations and the existential pebble game, and use it to characterize when a $\mathsf{CQ}\ q$ has at least one $\mathsf{GHW}(k)$ - Δ -approximation that is neither a $\mathsf{GHW}(k)$ -underapproximation nor a $\mathsf{GHW}(k)$ -overapproximation (a so-called incomparable $\mathsf{GHW}(k)$ - Δ -approximation). This allows us to show that the identification problem for such Δ -approximations is coNP -complete. As for the problem of checking for the existence of incomparable $\mathsf{GHW}(k)$ - Δ -approximations, we extend our automata techniques to prove that it is in $\mathsf{2EXPTIME}\$ for k=1 (and in $\mathsf{EXPTIME}\$ for fixed-arity schemas). In case such a $\mathsf{GHW}(1)$ - Δ -approximation exists, we can evaluate it using a fixed-parameter tractable algorithm. We also provide results on existence and evaluation of infinite incomparable $\mathsf{GHW}(1)$ - Δ -approximations.

Organization. Section 2 contains background notions and results, while Section 3 introduce approximations. Basic properties of overapproximations are presented in Section 4, while the existence of overapproximations is studied in Section 5. In Section 6 we study overapproximations under constraints, while in Section 7 we deal with Δ -approximations. We conclude in Section 8 with final remarks.

2 Preliminaries

2.1 Databases, homomorphisms, and conjunctive queries

Databases and homomorphisms. A relational schema σ is a finite set of relation symbols, each one of which has an arity n>0. A database $\mathcal D$ over σ is a finite set of facts of the form $R(\bar a)$, where R is a relation symbol in σ of arity n and $\bar a$ is an n-tuple of constants. We often abuse notation and write $\mathcal D$ also for the set of constants mentioned in the facts of $\mathcal D$.

Let \mathcal{D} and \mathcal{D}' be databases over σ . A homomorphism from \mathcal{D} to \mathcal{D}' is a mapping h from \mathcal{D} to \mathcal{D}' such that for every atom $R(\bar{a})$ in \mathcal{D} it is the case that $R(h(\bar{a})) \in \mathcal{D}'$. Here, we use the convention that if $\bar{a} = (a_1, \ldots, a_n)$ then $h(\bar{a}) = (h(a_1), \ldots, h(a_n))$. If \bar{a} and \bar{b} are n-ary tuples (for $n \geq 0$) in \mathcal{D} and \mathcal{D}' , respectively, we write $(\mathcal{D}, \bar{a}) \to (\mathcal{D}', \bar{b})$ if there is a homomorphism h from \mathcal{D} to \mathcal{D}' such that $h(\bar{a}) = \bar{b}$. Checking if $(\mathcal{D}, \bar{a}) \to (\mathcal{D}', \bar{b})$ is a well-known NP-complete problem.

Conjunctive queries. A conjunctive query (CQ) over schema σ is a formula of the form $q(\bar{x}) = \exists \bar{y} \bigwedge_{1 \leq i \leq m} R_i(\bar{x}_i)$, where each $R_i(\bar{x}_i)$ is an atom over σ for each i with $1 \leq i \leq m$, and the tuple \bar{x} contains precisely the *free variables*, i.e., the variables that do not appear existentially quantified in \bar{y} . Note that

our definition allows for repetitions of variables in \bar{x} . We often refer implicitly to the free variables \bar{x} and write q for $q(\bar{x})$. If \bar{x} is empty, then q is *Boolean*.

As customary, we define the evaluation of CQs in terms of homomorphisms. Recall that the canonical database \mathcal{D}_q of a CQ $q(\bar{x}) = \exists \bar{y} \bigwedge_{1 \leq i \leq m} R_i(\bar{x}_i)$ consists precisely of the atoms $R_i(\bar{x}_i)$, for $1 \leq i \leq m$. We then define the result of q over \mathcal{D} , denoted $q(\mathcal{D})$, as the set of all tuples \bar{a} of elements (i.e., constants) in \mathcal{D} such that $(\mathcal{D}_q, \bar{x}) \to (\mathcal{D}, \bar{a})$. We often do not distinguish between a CQ q and its canonical database \mathcal{D}_q , i.e., we write q for \mathcal{D}_q . If q is Boolean, then its evaluation over a database \mathcal{D} correspond to the Boolean values true or false depending on whether $q(\mathcal{D}) = \{()\}$ or $q(\mathcal{D}) = \emptyset$, respectively.

Evaluation and tractable classes of CQs. The evaluation problem for CQs is as follows: Given a CQ q, a database \mathcal{D} , and a tuple \bar{a} of elements in \mathcal{D} , is $\bar{a} \in q(\mathcal{D})$? Since this problem corresponds to checking if $(q, \bar{x}) \to (\mathcal{D}, \bar{a})$, it is NP-complete [11]. This led to a flurry of activity for finding classes of CQs for which evaluation is tractable (see, e.g., [40,12,25,28]).

Here we deal with one of the most studied such classes: CQs of bounded generalized hypertreewidth [25], also called coverwidth [13]. We adopt the definition of [13] which is better suited for working with non-Boolean queries. A tree decomposition of a CQ $q = \exists \bar{y} \bigwedge_{1 \leq i \leq m} R_i(\bar{x}_i)$ is a pair (T, χ) , where T is a tree and χ is a mapping that assigns a subset of the existentially quantified variables in \bar{y} to each node $t \in T$, such that:

- 1. For each $1 \leq i \leq m$, the variables in $\bar{x}_i \cap \bar{y}$ are contained in $\chi(t)$, for some $t \in T$.
- 2. For each variable y in \bar{y} , the set of nodes $t \in T$ for which y occurs in $\chi(t)$ is connected.

The width of node t in (T, χ) is the minimal size of an $I \subseteq \{1, \ldots, m\}$ such that $\bigcup_{i \in I} \bar{x}_i$ covers $\chi(t)$ (where we slightly abuse notation and write \bar{x}_i also for the set of variables mentioned in the tuple \bar{x}_i). The width of (T, χ) is the maximum width of the nodes of T. The generalized hypertreewidth of q is the minimum width of its tree decompositions.

For a fixed $k \geq 1$, we denote by $\mathsf{GHW}(k)$ the class of CQs of generalized hypertreewidth at most k. The CQs in $\mathsf{GHW}(k)$ can be evaluated in polynomial time; see [24].

Containment of CQs. A CQ q is contained in a CQ q', written as $q \subseteq q'$, if $q(\mathcal{D}) \subseteq q'(\mathcal{D})$ over every database \mathcal{D} . Two CQs q and q' are equivalent, denoted $q \equiv q'$, if $q \subseteq q'$ and $q' \subseteq q$.

It is known that CQ containment and CQ evaluation are, essentially, the same problem [11]. In particular, let $q(\bar{x})$ and $q'(\bar{x}')$ be CQs. Then

$$q \subseteq q' \iff \bar{x} \in q'(\mathcal{D}_q) \iff (\mathcal{D}_{q'}, \bar{x}') \to (\mathcal{D}_q, \bar{x}).$$
 (1)

Thus, $q \subseteq q'$ and $(q', \bar{x}') \to (q, \bar{x})$ – i.e., $(\mathcal{D}_{q'}, \bar{x}') \to (\mathcal{D}_q, \bar{x})$ – are used interchangeably.

Cores of CQs. A CQ q is a core [11,29] if there is no CQ q' with fewer atoms than q such that $q \equiv q'$. Given CQs q and q', we say that q' is a core of q if q' is a core and $q \equiv q'$. In other words, a core of q is a minimal CQ (in terms of number of atoms) that is equivalent to q. The following result summarizes some important properties of cores.

Proposition 1 [29] The following statements hold:

- 1. Each CQ q has a core. Moreover, there is a unique core of q up to renaming of variables. Therefore, we can talk about the core of q.
- 2. Each core q' of q is a retract of q. That is, each atom of q' is also an atom of q and we can choose the homomorphism h witnessing $(q, \bar{x}) \to (q', \bar{x}')$ to be a retraction, i.e., to be the identity on the variables of q'.

2.2 The existential cover game

Several results in the paper require applying techniques based on existential pebble games. We use a version of the *existential cover game*, that is tailored for CQs of bounded generalized hypertreewidth [13].

Let $k \geq 1$. The existential k-cover game is played by Spoiler and Duplicator on pairs (\mathcal{D}, \bar{a}) and (\mathcal{D}', \bar{b}) , where \mathcal{D} and \mathcal{D}' are databases and \bar{a} and \bar{b} are n-ary (for $n \geq 0$) tuples over the elements (i.e., the constants) in \mathcal{D} and \mathcal{D}' , respectively. The game proceeds in rounds. In each round, Spoiler places (resp., removes) a pebble on (resp., from) an element in \mathcal{D} , and Duplicator responds by placing (resp., removing) its corresponding pebble on an element in (resp., from) \mathcal{D}' . The number of pebbles is not bounded, but Spoiler is constrained as follows: At any round p of the game, if c_1, \ldots, c_ℓ ($\ell \leq p$) are the elements marked by Spoiler's pebbles in \mathcal{D} , there must be a set of at most k atoms in \mathcal{D} that contain all such elements (this is why the game is called k-cover game, as pebbled elements are covered by no more than k atoms).

Duplicator wins if she has a winning strategy, i.e., if she can indefinitely continue playing the game in such a way that after each round, if c_1, \ldots, c_ℓ are the elements that are marked by Spoiler's pebbles in \mathcal{D} and d_1, \ldots, d_ℓ are the elements marked by the corresponding pebbles of Duplicator in \mathcal{D}' , then

$$((c_1,\ldots,c_\ell,\bar{a}),(d_1,\ldots,d_\ell,\bar{b}))$$

is a partial homomorphism from \mathcal{D} to \mathcal{D}' . That is, for every atom $R(\bar{c}) \in \mathcal{D}$, where each element c of \bar{c} appears in $(c_1, \ldots, c_\ell, \bar{a})$, it is the case that $R(\bar{d}) \in \mathcal{D}'$, where \bar{d} is the tuple obtained from \bar{c} by replacing each element c of \bar{c} by its corresponding element d in $(d_1, \ldots, d_\ell, \bar{b})$. We write $(\mathcal{D}, \bar{a}) \to_k (\mathcal{D}', \bar{b})$ if Duplicator has a winning strategy.

Notice that if $(\mathcal{D}, \bar{a}) \to (\mathcal{D}', \bar{b})$ via some homomorphism h then h witnesses that Duplicator can win the k-pebble game for each k. Hence, \to_k can be seen as an "approximation" of \to :

$$\rightarrow \subset \cdots \subset \rightarrow_{k+1} \subset \rightarrow_k \subset \cdots \subset \rightarrow_1$$
.

These approximations are convenient complexity-wise: Checking if $(\mathcal{D}, \bar{a}) \to (\mathcal{D}', \bar{b})$ is NP-complete, while $(\mathcal{D}, \bar{a}) \to_k (\mathcal{D}', \bar{b})$ can be solved efficiently.

Proposition 2 [13] Fix $k \geq 1$. Checking whether $(\mathcal{D}, \bar{a}) \rightarrow_k (\mathcal{D}', \bar{b})$ is in polynomial time.

Moreover, there is a connection between \rightarrow_k and the evaluation of CQs in $\mathsf{GHW}(k)$ that we heavily exploit in our work.

Proposition 3 [13] Fix $k \geq 1$. Then $(\mathcal{D}, \bar{a}) \rightarrow_k (\mathcal{D}', \bar{b})$ iff for each CQ $q(\bar{x})$ in GHW(k) we have that if $(q, \bar{x}) \rightarrow (\mathcal{D}, \bar{a})$ then $(q, \bar{x}) \rightarrow (\mathcal{D}', \bar{b})$.

In particular, if $q(\bar{x}) \in \mathsf{GHW}(k)$ then for every \mathcal{D} and \bar{a} it is the case that

$$\bar{a} \in q(\mathcal{D}) \iff (q, \bar{x}) \to (\mathcal{D}, \bar{a}) \iff (q, \bar{x}) \to_k (\mathcal{D}, \bar{a}).$$
 (2)

That is, the "approximation" of \to provided by \to_k is sufficient for evaluating CQs in $\mathsf{GHW}(k)$. Together with Proposition 2, this proves that CQs in $\mathsf{GHW}(k)$ can be evaluated efficiently.

As a matter of fact, the equivalences established in Equation (2) hold even if q itself is not in $\mathsf{GHW}(k)$, but its core q' is in $\mathsf{GHW}(k)$. That is, the evaluation problem for the class of CQs whose core is in $\mathsf{GHW}(k)$ can be solved efficiently via the existential k-cover game. As established by Greco and Scarcello, this is precisely the boundary for when this good property holds. Indeed, for any Boolean $\mathsf{CQ}\ q$ whose core is not in $\mathsf{GHW}(k)$ it is possible to find a database $\mathcal D$ such that $q \to_k \mathcal D$ but $q \not\to \mathcal D$ [27].

2.3 Expressing the existential cover game as a CQ in GHW(k)

An instrumental tool in several of our results is that for any given pair (\mathcal{D}, \bar{a}) , where \mathcal{D} is a database and \bar{a} is a tuple of elements (i.e., constants) over \mathcal{D} , we can construct a $\operatorname{CQ} q(\bar{x})$ in $\operatorname{GHW}(k)$ that represents the possible moves of Spoiler in the existential k-cover game played from \mathcal{D} , but only up to certain number of rounds.

For simplicity, we consider a compact version of the existential k-cover game as in [13]. For a database \mathcal{D} , we say that a set S of elements of \mathcal{D} is a k-union if there exist p atoms $R_1(\bar{a}_1), \ldots R_p(\bar{a}_p) \in \mathcal{D}$ with $p \leq k$, such that $S = \bigcup_{1 \leq i \leq p} \bar{a}_i$. That is, the k-unions of \mathcal{D} are the sets that appear in a union of at most k atoms of \mathcal{D} . In the compact existential k-cover game, Spoiler is allowed, in each round, to remove and place as many pebbles as desired, as long as the resulting pebbled elements form a k-union. (Notice the difference with the standard existential k-cover game, in which in each round Spoiler is allowed to either remove or place exactly one pebble). As before, Duplicator wins the compact existential k-cover game on (\mathcal{D}, \bar{a}) and (\mathcal{D}', \bar{b}) iff she has a winning strategy, i.e., she can indefinitely continue playing the game in such a way that after each round, if c_1, \ldots, c_ℓ are the elements marked by Spoiler's pebbles in \mathcal{D} and d_1, \ldots, d_ℓ are the elements marked by the corresponding Duplicator's

pebbles in \mathcal{D}' , then $((c_1,\ldots,c_\ell,\bar{a}),(d_1,\ldots,d_\ell,\bar{b}))$ is a partial homomorphism from \mathcal{D} to \mathcal{D}' .

As can be easily seen, the compact existential k-cover game is not more powerful than the standard one. That is, for each $k \geq 1$ Duplicator wins the existential k-cover game on pairs (\mathcal{D}, \bar{a}) and (\mathcal{D}', \bar{b}) iff she wins the compact existential k-cover game on such pairs [13].

We then write $(\mathcal{D}, \bar{a}) \to_k^c (\mathcal{D}', \bar{b})$, for $k \geq 1$ and $c \geq 1$, if Duplicator has a winning strategy in the first c rounds of the (compact) existential k-cover game on (\mathcal{D}, \bar{a}) and (\mathcal{D}', \bar{b}) . That is, $(\mathcal{D}, \bar{a}) \to_k (\mathcal{D}', \bar{b})$ if and only if $(\mathcal{D}, \bar{a}) \to_k^c (\mathcal{D}', \bar{b})$ for every $c \geq 1$. The following is our desired technical result, which establishes that there is a CQ in GHW(k) that represents the possible moves of Spoiler – up to c rounds – in the existential k-cover game played from (\mathcal{D}, \bar{a}) .

Lemma 1 Let \mathcal{D} be a database and \bar{a} an n-ary tuple of elements in \mathcal{D} , for $n \geq 0$. For each $c \geq 1$ there is a CQ $q_c(\bar{x}_c)$ in $\mathsf{GHW}(k)$ such that for each database \mathcal{D}' and tuple \bar{b} of the same arity as \bar{a} we have that

$$\bar{b} \in q_c(\mathcal{D}') \iff (q_c, \bar{x}_c) \to (\mathcal{D}', \bar{b}) \iff (\mathcal{D}, \bar{a}) \to_k^c (\mathcal{D}', \bar{b}).$$

Results of a similar kind have been obtained in [34], and actually the proof of Lemma 1 can easily be obtained by adapting techniques in such paper. We provide a proof, nevertheless, as we use it in several results along the article.

Proof (Lemma 1) Fix $c \geq 1$. We shall define a database \mathcal{D}_c such that $(\mathcal{D}_c, \bar{a}) \to (\mathcal{D}', \bar{b})$ iff $(\mathcal{D}, \bar{a}) \to_k^c (\mathcal{D}', \bar{b})$, for every pair (\mathcal{D}', \bar{b}) . The lemma will follow by choosing $q_c(\bar{x}_c)$ such that $(\mathcal{D}_{q_c}, \bar{x}_c) = (\mathcal{D}_c, \bar{a})$.

Let S_1, \ldots, S_N be an enumeration of all the k-unions of \mathcal{D} , where $N \geq 1$. In order to define \mathcal{D}_c , we first consider a pair (T_c, λ_c) where T_c is an ordered rooted tree (i.e., the children of a node are ordered) with rank N and height c, and λ_c is a labelling that maps each $t \in T_c$ to a k-union of \mathcal{D} such that (i) $\lambda_c(r) = \emptyset$, for the root r, and (ii) if t_i is the i-th child of t, then $\lambda_c(t_i) = S_i$. The pair (T_c, λ_c) is simply a representation of all the possible moves of Spoiler on \mathcal{D} in the compact existential k-cover game, up to c rounds.

Now we define a pair (T_c, β_c) as follows. The intuition is that (T_c, β_c) is obtained from (T_c, λ_c) via renaming elements in the $\lambda_c(t)$'s in such a way that (T_c, λ_c) satisfies the connectivity condition of tree decompositions. An occurrence of an element $d \in \mathcal{D}$ in (T_c, λ_c) is a maximal subtree T of T_c such that $d \in \lambda_c(t)$, for all $t \in T$. For each $d \in \mathcal{D} \setminus \bar{a}$ and each occurrence T of d in (T_c, λ_c) , we introduce a new element $e_{d,T}$, and modify each $\lambda_c(t)$ with $t \in T$ by replacing d with $e_{d,T}$. The resulting labelling is β_c . Note that, for each $t \in T_c$, there is a bijection Φ_t from $\lambda_c(t)$ to $\beta_c(t)$. By construction, $\Phi_t(a) = a$, for all $a \in \lambda_c(t) \cap \bar{a}$, and $\Phi_t(d) = \Phi_{t'}(d)$, where t' is the parent of t and $d \in \lambda_c(t) \cap \lambda_c(t')$.

The database \mathcal{D}_c is obtained as follows. For each atom $R(\bar{d})$ in \mathcal{D} and node $t \in T_c$ such that $\bar{d} \subseteq \lambda_c(t)$, we add to \mathcal{D}_c the atom $R(\Phi_t(\bar{d}))$. Note that the same atom $R(\bar{d})$ can produce several atoms in \mathcal{D}_c . By definition, $\Phi_t : \lambda_c(t) \to \beta_c(t)$ and $\Phi_t^{-1} : \beta_c(t) \to \lambda_c(t)$ are partial homomorphisms from \mathcal{D} to \mathcal{D}_c and \mathcal{D}_c

to \mathcal{D} , respectively, for all $t \in T_c$. To see that the resulting CQ $q_c(\bar{x}_c)$ is in GHW(k), we consider the pair (T_c, χ_c) , where for each $t \in T_c$, $\chi_c(t) = \beta_c(t) \setminus \bar{a}$. By construction, (T_c, χ_c) is a tree decomposition of $q_c(\bar{x}_c)$ of width k.

Let us conclude by stressing that $(\mathcal{D}_c, \bar{a}) \to (\mathcal{D}', \bar{b})$ iff $(\mathcal{D}, \bar{a}) \to_k^c (\mathcal{D}', \bar{b})$, for every pair (\mathcal{D}', \bar{b}) . Suppose that $(\mathcal{D}_c, \bar{a}) \to (\mathcal{D}', \bar{b})$ via a homomorphism h. If the first move of Spoiler is the k-union S_i of \mathcal{D} , for some $i \in \{1, \ldots, N\}$, then the Duplicator responds with the partial homomorphism $h \circ \Phi_{t_i}$, where t_i is the i-th child of the root r (and hence, the domain of Φ_{t_i} is $\lambda_c(t_i) = S_i$). If in the next round Spoiler plays S_j for some $j \in \{1, \ldots, N\}$, then the Duplicator responds with $h \circ \Phi_{t_j}$, where t_j is the j-th child of t_i . This is a valid move for Duplicator since, as mentioned above, Φ_{t_i} and Φ_{t_j} are consistent. In this way, Duplicator can win the first c rounds of the game as the height of T_c is c.

Conversely, assume that $(\mathcal{D}, \bar{a}) \to_k^c (\mathcal{D}', \bar{b})$. We can define a homomorphism h from \mathcal{D}_c to \mathcal{D}' such that $h(\bar{a}) = \bar{b}$ in a top-down fashion over T_c . We start with the root r, and make Spoiler play $\Phi_r^{-1}(\beta_c(r))$ as his first move. Suppose Duplicator responds with a partial homomorphism $g_0 : \lambda_c(r) \to \mathcal{D}'$. Then $h_0 = g_0 \circ \Phi_r^{-1}$ is a partial homomorphism from $\beta_c(r)$ to \mathcal{D}' . Following this argument, and making Spoiler play accordingly, we can extend h_0 to partial homomorphisms h_1, \ldots, h_c , where the domain of h_p , with $p \in \{1, \ldots, c\}$, are the variables in some $\beta_c(t)$ such that the distance from t to root r is at most p. We can define $h = h_c$ to be our desired homomorphism from \mathcal{D}_c to \mathcal{D}' . \square

3 Approximations of CQs

Fix $k \geq 1$. Let q be a CQ. The approximations of q in $\mathsf{GHW}(k)$ are defined with respect to a partial order \sqsubseteq_q over the set of CQs in $\mathsf{GHW}(k)$. Formally, for any two CQs q', q'' in $\mathsf{GHW}(k)$ we have

$$q' \sqsubseteq_q q'' \iff \Delta(q(\mathcal{D}), q''(\mathcal{D})) \subseteq \Delta(q(\mathcal{D}), q'(\mathcal{D})), \text{ for every database } \mathcal{D},$$

where $\Delta(A, B)$ denotes the symmetric difference between sets A and B. Thus, $q' \sqsubseteq_q q''$, whenever the "error" of q'' with respect to q – measured in terms of the symmetric difference between $q''(\mathcal{D})$ and $q(\mathcal{D})$ – is contained in that of q' for each database \mathcal{D} . As usual, we write $q' \sqsubseteq_q q''$ if $q' \sqsubseteq_q q''$ but $q'' \not\sqsubseteq_q q'$.

The approximations of q in $\mathsf{GHW}(k)$ always correspond to maximal elements, with respect to the partial order \sqsubseteq_q , over a class of CQs in $\mathsf{GHW}(k)$ that satisfies certain conditions. The following three basic notions of approximation were identified in [6]:

- Underapproximations: Let q, q' be CQs such that $q' \in \mathsf{GHW}(k)$. Then q' is a $\mathsf{GHW}(k)$ -underapproximation of q if it is maximal, with respect to \sqsubseteq_q , among all CQs in $\mathsf{GHW}(k)$ that are contained in q. That is, it holds that

$$q' \subseteq q$$
, and there is no CQ $q'' \in \mathsf{GHW}(k)$ such that $q'' \subseteq q$ and $q' \sqsubseteq_q q''$.

In particular, the $\mathsf{GHW}(k)$ -underapproximations of q produce correct (but not necessarily complete) answers with respect to q over every database \mathcal{D} .

- Overapproximations: Analogously, q' is a $\mathsf{GHW}(k)$ -overapproximation of q if it is maximal, with respect to \sqsubseteq_q , among all CQs in $\mathsf{GHW}(k)$ that contain q. That is, it holds that

 $q \subseteq q'$, and there is no CQ $q'' \in \mathsf{GHW}(k)$ such that $q \subseteq q''$ and $q' \sqsubset_q q''$.

Hence, $\mathsf{GHW}(k)$ -overapproximations of q produce complete (but not necessarily correct) answers with respect to q over every database \mathcal{D} .

- Δ -approximations: In this case we impose no restriction on q'. That is, $\overline{q'}$ is a $\mathsf{GHW}(k)$ - Δ -approximation of q if it is maximal with respect to the partial order \sqsubseteq_q , i.e., there is no $q'' \in \mathsf{GHW}(k)$ such that $q' \sqsubseteq_q q''$.

Underapproximations and overapproximations admit an equivalent, but arguably simpler characterization as maximal (resp., minimal) elements, with respect to the containment partial order \subseteq , among all CQs in $\mathsf{GHW}(k)$ that are contained in q (resp., contain q). We state this characterization next.

Proposition 4 [6] Fix $k \ge 1$. Let q, q' be CQs such that $q' \in GHW(k)$. Then:

- -q' is a $\mathsf{GHW}(k)$ -underapproximation of q iff $q' \subseteq q$ and there is no CQ $q'' \in \mathsf{GHW}(k)$ such that $q' \subset q'' \subseteq q$.
- -q' is a $\mathsf{GHW}(k)$ -overapproximation of q iff $q \subseteq q'$ and there is no CQ $q'' \in \mathsf{GHW}(k)$ such that $q \subseteq q'' \subset q'$.

The basic theoretical properties of $\mathsf{GHW}(k)$ -underapproximations are by now well-understood [7]. We concentrate on $\mathsf{GHW}(k)$ -overapproximations and $\mathsf{GHW}(k)$ - Δ -approximations in this paper. We start by studying the former.

4 Overapproximations

Recall that $\mathsf{GHW}(k)$ -overapproximations are minimal elements (in terms of \subseteq) in the set of CQs in $\mathsf{GHW}(k)$ that contain q. We show an example of a $\mathsf{GHW}(1)$ -overapproximation below.

Example 1 Figure 2 shows a CQ q and its GHW(1)-overapproximation q'. The schema consists of binary symbols P_a and P_b . Nodes represent variables, and an edge labeled P_a between x and y represents the presence of atoms $P_a(x,y)$ and $P_a(y,x)$. (Same for P_b). All variables are existentially quantified. Clearly, $q \subseteq q'$ (as $q' \to q$). In addition, there is no CQ $q'' \in \mathsf{GHW}(1)$ such that $q \subseteq q'' \subset q'$. We provide an explanation for this later.

We start in Section 4.1 by stating some basic properties on existence and uniqueness of $\mathsf{GHW}(k)$ -overapproximations. Later in Section 4.2 we establish a connection between $\mathsf{GHW}(k)$ -overapproximations and the existential pebble game, which allows us to show that both the identification and evaluation problems for $\mathsf{GHW}(k)$ -overapproximations are tractable. Finally, in Section 4.4 we look at the case when $\mathsf{GHW}(k)$ -overapproximations do not exist, and suggest how this can be alleviated by allowing infinite overapproximations.




Fig. 2 The CQ q and its GHW(1)-overapproximation q' from Example 1.

4.1 Existence and uniqueness of overapproximations

As shown in [6], existence of overapproximations is not a general phenomenon. In fact, for every k > 1 there is a Boolean CQ q in GHW(k) that has no GHW(1)-overapproximation. Using the characterization given later in Theorem 7, we can strengthen this further.

Theorem 1 For each k > 1, there is a Boolean $CQ \ q \in GHW(k)$ without $\mathsf{GHW}(\ell)$ -overapproximations for any $1 \leq \ell < k$.

Figure 3 depicts examples of CQs in GHW(k), for k = 2 and k = 3, respectively, without $\mathsf{GHW}(\ell)$ -overapproximations for any $1 \leq \ell < k$. The proof of Theorem 1 is long and quite technical, and for such a reason we relegate it to the appendix.

Interestingly, when GHW(k)-overapproximations do exist, they are unique (up to equivalence). This is because, in this case, GHW(k)-overapproximations are not only the minimal elements, but also the lower bounds of the set of CQs in GHW(k) that contain q.

In order to show uniqueness, we need to introduce a simple construction that also will be useful when studying Δ -approximations (Section 7). Let $q(\bar{x})$ and $q'(\bar{x}')$ be two CQs such that $|\bar{x}| = |\bar{x}'| = n$. The disjoint conjunction of q and q' is the CQ $(q \wedge q')(\bar{z})$, with $|\bar{z}| = n$ defined as follows. First we rename each existentially quantified variable in q and q' with a different fresh variable, and then take the conjunction of the atoms in q and q'. Finally, if $\bar{x} = (x_1, \dots, x_n)$ and $\bar{x}' = (x'_1, \dots, x'_n)$, we identify x_i and x'_i for every $1 \leq i \leq n$. The *i*-th variable of \bar{z} is the variable obtained after identifying x_i and x_i' . By construction, the following hold:

- 1. $(q, \bar{x}) \to (q \land q', \bar{z})$ and $(q', \bar{x}') \to (q \land q', \bar{z})$, 2. if $(q, \bar{x}) \to (p, \bar{w})$ and $(q', \bar{x}') \to (p, \bar{w})$, for some CQ $p(\bar{w})$, then $(q \land q', \bar{z}) \to (q, \bar{w})$ (p, \bar{w}) , and
- 3. if $q, q' \in \mathsf{GHW}(k)$ for $k \ge 1$, then $q \wedge q' \in \mathsf{GHW}(k)$.

Note that property (1) and (2) tell us that $q \wedge q'$ is the least upper bound of q and q' with respect to the order \rightarrow . We have the following result:

Proposition 5 Let $q(\bar{x})$ and $q'(\bar{x}')$ be CQs such that $q' \in GHW(k)$. The following are equivalent:

1. $q'(\bar{x}')$ is a GHW(k)-overapproximation of $q(\bar{x})$.




Fig. 3 The CQ q is in GHW(2) but has no GHW(1)-overapproximations, while q' is in GHW(3) but has no GHW(ℓ)-overapproximations for $\ell \in \{1,2\}$.

2. (i) $q \subseteq q'$, and (ii) for every $CQ \ q''(\bar{x}'') \in \mathsf{GHW}(k)$, it is the case that $q \subseteq q''$ implies $q' \subseteq q''$.

Proof We only prove the nontrivial direction $(1) \Rightarrow (2)$. By contradiction, suppose that there is a CQ $q'' \in \mathsf{GHW}(k)$ such that $q \subseteq q''$ but $q' \not\subseteq q''$. Let $(q' \land q'')(\bar{z})$ be the disjoint conjunction of q' and q''. By definition, we have that $q' \land q''$ is in $\mathsf{GHW}(k)$, and $q \subseteq (q' \land q'') \subseteq q'$. But q' is a $\mathsf{GHW}(k)$ -overapproximation of q, and thus $q' \subseteq (q' \land q'')$. Again by construction of $q' \land q''$, we have that $(q' \land q'') \subseteq q''$, and then $q' \subseteq q''$. This is a contradiction.

As a corollary, we immediately obtain the following.

Corollary 1 Consider a CQ q with GHW(k)-overapproximations q_1 and q_2 . Then it is the case that $q_1 \equiv q_2$.

This result shows a stark difference between $\mathsf{GHW}(k)$ -overapproximations and $\mathsf{GHW}(k)$ -underapproximations: $\mathsf{GHW}(k)$ -overapproximations do not necessarily exist, but when they do they are unique; $\mathsf{GHW}(k)$ -underapproximations always exist but there can be exponentially many incomparable ones [7].

4.2 A link with the existential pebble game

Existential cover games can be applied to obtain a semantic characterization of GHW(k)-overapproximations as follows.

Theorem 2 Fix $k \ge 1$. Let $q(\bar{x})$ and $q'(\bar{x}')$ be CQs with $q' \in GHW(k)$. The following are equivalent:

1. $q'(\bar{x}')$ is the GHW(k)-overapproximation of $q(\bar{x})$.

2. $(q', \bar{x}') \rightarrow_k (q, \bar{x})$ and $(q, \bar{x}) \rightarrow_k (q', \bar{x}')$.

Proof Assume that $q'(\bar{x}')$ is the $\mathsf{GHW}(k)$ -overapproximation of $q(\bar{x})$. Then $(q', \bar{x}') \to (q, \bar{x})$, and therefore $(q', \bar{x}') \to_k (q, \bar{x})$. It remains to prove that $(q, \bar{x}) \to_k (q', \bar{x}')$. From Proposition 3, we need to prove that if $q''(\bar{x}'')$ is a CQ in $\mathsf{GHW}(k)$ such that $(q'', \bar{x}'') \to (q, \bar{x})$, then also $(q'', \bar{x}'') \to (q', \bar{x}')$. This follows directly from Proposition 5.

Assume now that $(q', \bar{x}') \to_k (q, \bar{x})$ and $(q, \bar{x}) \to_k (q', \bar{x}')$. Since $q' \in \mathsf{GHW}(k)$ and $(q', \bar{x}') \to_k (q, \bar{x})$, we have that $(q', \bar{x}') \to (q, \bar{x})$ by Equation (2), and hence $q \subseteq q'$ by Equation (1). In addition, since $(q, \bar{x}) \to_k (q', \bar{x}')$ it follows from

Proposition 3 and Equation (1) that if $q \subseteq q''$ and $q'' \in \mathsf{GHW}(k)$, then $q' \subseteq q''$. That is, there is no q'' in $\mathsf{GHW}(k)$ such that $q \subseteq q'' \subset q'$. Hence q' is a $\mathsf{GHW}(k)$ -overapproximation.

Example 2 (Example 1 cont.) It is now easy to see that the CQ q' in Figure 2 is a GHW(1)-overapproximation of q. In fact, since $q' \to q$, we only need to show that $q \to_1 q'$. The latter is simple and left to the reader.

Next we show that this characterization allows us to show that the identification and evaluation problems for $\mathsf{GHW}(k)$ -overapproximations can be solved in polynomial time.

4.3 Identification and evaluation of GHW(k)-overapproximations

A direct corollary of Proposition 2 and Theorem 2 is that the *identification* problem for $\mathsf{GHW}(k)$ -overapproximations is in polynomial time.

Corollary 2 Fix $k \ge 1$. Given $CQs \ q, q'$ such that $q' \in GHW(k)$, checking if q' is the GHW(k)-overapproximation of q can be solved in polynomial time.

This corresponds to a *promise version* of the problem, as it is given to us that q' is in fact in $\mathsf{GHW}(k)$. Checking the latter is NP-complete for every fixed k > 1 [26,21].

Let us assume now that we are given the *promise* that q has a $\mathsf{GHW}(k)$ -overapproximation q' (but q' itself is not given). How hard is it to evaluate q' over a database \mathcal{D} ? We could try to compute q', but so far we have no techniques to do that. Notably, we can use existential cover games to show that $\mathsf{GHW}(k)$ -overapproximations can be evaluated efficiently, without even computing them. This is based on the next result, which states that evaluating q' over \mathcal{D} boils down to checking $(q, \bar{x}) \to_k (\mathcal{D}, \bar{a})$ for the tuples \bar{a} over \mathcal{D} .

Theorem 3 Consider a fixed $k \geq 1$. Let $q(\bar{x})$ be a CQ with a GHW(k)overapproximation $q'(\bar{x}')$. Then for every \mathcal{D} and \bar{a} it is the case that

$$\bar{a} \in q'(\mathcal{D}) \iff (q', \bar{x}') \to (\mathcal{D}, \bar{a}) \iff (q, \bar{x}) \to_k (\mathcal{D}, \bar{a}).$$

Proof Assume first that $(q, \bar{x}) \to_k (\mathcal{D}, \bar{a})$. Based on the fact that q' is a $\mathsf{GHW}(k)$ -overapproximation of q, we have that $(q', \bar{x}') \to (q, \bar{x})$. Since winning strategies for Duplicator compose and $(q', \bar{x}') \to (q, \bar{x})$ implies $(q', \bar{x}') \to_k (q, \bar{x})$, it is the case that $(q', \bar{x}') \to_k (\mathcal{D}, \bar{a})$. But $q' \in \mathsf{GHW}(k)$, and thus $(q', \bar{x}') \to (\mathcal{D}, \bar{a})$ from Equation (2).

Assume, on the other hand, that $(q', \bar{x}') \to (\mathcal{D}, \bar{a})$. From Theorem 2, we have that $(q, \bar{x}) \to_k (q', \bar{x}')$. By composition and the fact that $(q', \bar{x}') \to (\mathcal{D}, \bar{a})$ implies $(q', \bar{x}') \to_k (\mathcal{D}, \bar{a})$, it follows that $(q, \bar{x}) \to_k (\mathcal{D}, \bar{a})$ holds. \square

As a corollary to Theorem 3 and Proposition 2 we obtain the following.

Corollary 3 Fix $k \geq 1$. Checking if $\bar{a} \in q'(\mathcal{D})$, given a CQ q that has a GHW(k)-overapproximation q', a database \mathcal{D} , and a tuple \bar{a} in \mathcal{D} , can be solved in polynomial time by checking if $(q, \bar{x}) \rightarrow_k (\mathcal{D}, \bar{a})$. Moreover, this can be done without even computing q'.

4.4 More liberal GHW(k)-overapproximations

CQs may not have $\mathsf{GHW}(k)$ -overapproximations, for some $k \geq 1$. We observe in this section that this anomaly can be taken care of by extending the language of queries over which overapproximations are to be found.

An infinite CQ is as a finite one, save that now the number of atoms is countably infinite. We assume that there are finitely many free variables in an infinite CQ. The evaluation of an infinite CQ $q(\bar{x})$ over a database \mathcal{D} is defined analogously to the evaluation of a finite one. Similarly, the generalized hypertreewidth of an infinite CQ is defined as in the finite case, but now tree decompositions can be infinite. We write $\mathsf{GHW}(k)^\infty$ for the class of all CQs, finite and infinite ones, of generalized hypertreewidth at most k. The next result states a crucial relationship between the existential k-cover game and the class $\mathsf{GHW}(k)^\infty$.

Lemma 2 Fix $k \ge 1$. For every CQ q there is a $q'(\bar{x}')$ in $GHW(k)^{\infty}$ such that for every database \mathcal{D} and tuple \bar{a} of constants in \mathcal{D} it is the case that

$$\bar{a} \in q'(\mathcal{D}) \iff (q', \bar{x}') \to (\mathcal{D}, \bar{a}) \iff (q, \bar{x}) \to_k (\mathcal{D}, \bar{a}).$$

This holds even for countably infinite databases \mathcal{D} .

Proof The lemma follows directly from the proof of Lemma 1 by starting with $(T_{\infty}, \lambda_{\infty})$ instead of (T_c, λ_c) , where $(T_{\infty}, \lambda_{\infty})$ is the infinite rooted tree labeled with k-unions of \mathcal{D}_q representing all possible moves of Spoiler on \mathcal{D}_q in the existential k-cover game. All arguments are still valid when \mathcal{D} is infinite. \square

Since we now deal with infinite CQs and databases, we cannot apply Proposition 3 directly in our analysis of $\mathsf{GHW}(k)^\infty$ -overapproximations. Instead, we use the following suitable reformulation of it, which we obtain by inspection of its proof.

Proposition 6 Fix $k \geq 1$. Consider countably infinite databases \mathcal{D} and \mathcal{D}' . Then $(\mathcal{D}, \bar{a}) \rightarrow_k (\mathcal{D}', \bar{b})$ iff for each CQ $q(\bar{x})$ in $\mathsf{GHW}(k)^{\infty}$ we have that if $(q, \bar{x}) \rightarrow (\mathcal{D}, \bar{a})$ then $(q, \bar{x}) \rightarrow (\mathcal{D}', \bar{b})$.

 $\mathsf{GHW}(k)^\infty$ -overapproximations. We expand the notion of overapproximation by allowing infinite CQs. Let $q' \in \mathsf{GHW}(k)^\infty$. Then q' is a $\mathsf{GHW}(k)^\infty$ -overapproximation of CQ q, if $q \subseteq q'$ and there is no $q'' \in \mathsf{GHW}(k)^\infty$ such that $q \subseteq q'' \subset q'$. (Here, \subseteq is still defined with respect to finite databases only). In $\mathsf{GHW}(k)^\infty$, we can provide each CQ q an overapproximation.

Theorem 4 Fix $k \ge 1$. For every CQ q there is a CQ in $GHW(k)^{\infty}$ that is a $GHW(k)^{\infty}$ -overapproximation of q.

Proof We show that the CQ $q'(\bar{x}')$ – as given in Lemma 2 – is a GHW $(k)^{\infty}$ overapproximation of $q(\bar{x})$. Notice first that $(q', \bar{x}') \to (q, \bar{x})$ (by choosing (\mathcal{D}, \bar{a}) as (q, \bar{x}) in Lemma 2), and hence $q \subseteq q'$. This is because the condition

that $(q', \bar{x}') \to (q, \bar{x})$ implies $q \subseteq q'$, expressed in Equation (1), holds even for countably infinite CQs. We also have that $(q, \bar{x}) \to_k (q', \bar{x}')$ (by choosing (\mathcal{D}, \bar{a}) as (q', \bar{x}') in Lemma 2). Proposition 6 then tells us that for each $q''(\bar{x}'')$ in $\mathsf{GHW}(k)^\infty$, if $(q'', \bar{x}'') \to (q, \bar{x})$ then $(q'', \bar{x}'') \to (q', \bar{x}')$. But then $q \subseteq q''$ implies $q' \subseteq q''$. This is because the condition that $q \subseteq q''$ implies $(q'', \bar{x}'') \to (q, \bar{x})$, expressed in Equation (1), continues to hold as long as q (but not necessarily q') is finite. Thus, q' is a $\mathsf{GHW}(k)^\infty$ -overapproximation of q.

Despite the non-computable nature of $\mathsf{GHW}(k)^\infty$ -over approximations, we get from Proposition 2 and the proof of Theorem 4 that they can be evaluated efficiently.

Corollary 4 Fix $k \geq 1$. Checking whether $\bar{a} \in q'(\mathcal{D})$, given a CQ q with $\mathsf{GHW}(k)^{\infty}$ -overapproximation q', a database \mathcal{D} , and a tuple \bar{a} in \mathcal{D} , boils down to checking if $(q, \bar{x}) \to_k (\mathcal{D}, \bar{a})$, and thus it can be solved in polynomial time.

Recall from Section 2 that the "approximation" of the notion of homomorphism provided by the existential k-cover game suffices for evaluating those $\operatorname{CQs} q(\bar{x})$ whose core is in $\operatorname{GHW}(k)$. That is, for every database $\mathcal D$ and tuple \bar{a} of elements in $\mathcal D$, we have that $\bar{a} \in q(\mathcal D) \Leftrightarrow (q,\bar{x}) \to (\mathcal D,\bar{a}) \Leftrightarrow (q,\bar{x}) \to_k (\mathcal D,\bar{a})$. For other CQs the existential k-cover game always provides an "overestimation" of the exact result. Interestingly, Corollary 4 establishes that such an overestimation is not completely arbitrary, as it is the one defined by the CQ in $\operatorname{GHW}(k)^\infty$ that better approximates q over the set of all databases.

5 Existence of Overapproximations

CQs always have $\mathsf{GHW}(k)^\infty$ -overapproximations, but not necessarily finite ones. Here we study when a CQ q has a finite overapproximation. We start with the case k=1, which we show to be decidable in 2EXPTIME (we do not know if this is optimal). For k>1 we leave the decidability open, but provide some explanation about where the difficulty lies.

5.1 The acyclic case

We start with the case of GHW(1)-overapproximations. Recall that GHW(1) is an important class, as it consists precisely of the well-known *acyclic* CQs. Our main result is the following:

Theorem 5 The following statements hold:

- (a) There is a 2Exptime algorithm that checks if a CQ q has a GHW(1)overapproximation and, if one exists, it computes one in triple-exponential
 time.
- (b) If the maximum arity of the schema is fixed, there is an EXPTIME algorithm that does this and computes a GHW(1)-overapproximation of q in double-exponential time.

The general idea behind the proof is as follows. From a CQ q we build a two-way alternating tree automaton [14], or 2ATA, \mathcal{A}_q , such that the language $L(\mathcal{A}_q)$ of trees accepted by \mathcal{A}_q is nonempty if and only if q has a GHW(1)-overapproximation. Intuitively, \mathcal{A}_q accepts those trees that encode a GHW(1)-overapproximation q' of q. The emptiness problems for languages defined by 2ATAs can be solved in exponential time in the number of states [14]. As our automaton \mathcal{A}_q will have exponentially many states, its emptiness can be tested in double-exponential time. In addition, if the maximum arity of the schema is fixed, the number of states in \mathcal{A}_q is polynomial, and hence emptiness can be tested in exponential time.

Before describing the details of the construction, let us shortly recapitulate two-way alternating tree automata. We closely follow the presentation from [14]. The input of an 2ATA is a ranked tree over an alphabet Σ . In each computation step, a 2ATA is in one of finitely many states and visits a node v of the input tree. Depending on the state and the label of the node, it can recursively start a bunch of processes; each of them starting from some state and either one of v's neighbors or from v itself. Whether the computation step is successful depends on a boolean combination of the outcomes of these processes.

Formally, a 2ATA is a tuple $(\Sigma, S, S_0, \delta, F)$ where Σ is the tree alphabet, S is a finite set of states, $S_0 \subseteq S$ is a set of initial states, $F \subseteq S$ is a set of accepting states, and δ is a transition function defined on $S \times \Sigma$ such that if $\sigma \in$ Σ has arity ℓ then $\delta(s,\sigma)$ is a propositional formula with variables from $S \times [\ell]$. Here, $[\ell]$ denotes the set $\{-1,0,1,\ldots,\ell\}$ of directions the automaton can take, where -1 denotes moving to the parent node, 0 denotes staying in the same node, and j > 0 denotes moving to the jth child. A proposition $(s', i) \in \delta(s, \sigma)$ represents that the automaton transitions into state s' and moves to the node represented by i. As an example, if $\delta(s,\sigma) = (r,2) \wedge ((p,1) \vee (q,-1))$ then, when being in state s and reading σ , the automaton starts two processes. One of them starting in state r in the second child; the other starting in state p in the first child or in state q in the parent node. Thus, in particular, using the propositional formula of a transition, a 2ATA can universally or existentially choose a next state. A run of the 2ATA starts in the root of an input tree and in a state from S_0 . Starting from there, a computation tree is spanned by applying the transitions. The input tree is accepted if the automaton has a computation tree whose leaves are accepting and the propositional formulas of transitions taken by the automaton are satisfied. We refer to [14] for the details of the semantics.

We now show how the problem of existence of GHW(1)-overapproximations can be reduced to the emptiness problem for 2ATA.

Proposition 7 There exists an algorithm that takes as input a CQ q and returns a 2ATA A_q such that q has a GHW(1)-overapproximation iff $L(A_q) \neq \emptyset$. Furthermore, the algorithm needs double-exponential time and A_q has exponentially many states. Furthermore:

- From every tree T in $L(\mathcal{A}_q)$ one can construct in polynomial time a $\mathsf{GHW}(1)$ overapproximation of q.
- If the maximum arity of the schema is fixed, then the algorithm needs exponential time and A_q has polynomially many states.

Proof For simplicity we assume that q is Boolean; towards the end of the proof we explain how the construction can be adapted for non-Boolean queries.

Before describing the construction of \mathcal{A}_q , we explain how input trees for \mathcal{A}_q encode CQs in GHW(1). Suppose that the maximum arity of q is r. Then an *encoding* of a GHW(1)-tree decomposition is a tree whose nodes are labeled with (a) variables from the set $\{u_1, \ldots, u_{2r}\}$ and (b) atoms over these variables whose arity is at most r; the only condition being that all variables of a node are covered by one of the atoms.

A CQ q' from GHW(1) with tree decomposition $(T_{q'}, \chi)$ of width one can be encoded as follows. Even though q' can have unbounded many variables, in each node of $T_{q'}$ at most r variables appear, where r is the maximum arity of an atom in q'. Thus, by reusing variables, $(T_{q'}, \chi)$ can be encoded by using 2r variables: the encoding $\operatorname{Enc}(T_{q'}, \chi)$ of $(T_{q'}, \chi)$ is obtained by traversing the nodes of $(T_{q'}, \chi)$ in a top-down fashion. Fresh variables in a node v, i.e. variables not used by its parent node, are encoded by fresh variables from $\{u_1, \ldots, u_{2r}\}$. On the other hand, an encoding of a $\operatorname{GHW}(1)$ -tree decomposition can be decoded in a top-down manner into a $\operatorname{GHW}(1)$ -tree decomposition by assigning a fresh variable name to each new occurrence of a variable u_i , that is, an occurrence of u_i that does not appear in the parent node. Observe that decoding $\operatorname{Enc}(T_{q'}, \chi)$ yields the decomposition of a query identical to q' up to renaming of variables.

The 2ATA \mathcal{A}_q needs to verify that the CQ q' encoded by $T' = \mathsf{Enc}(T_{q'}, \chi)$ is a GHW(1)-overapproximation of q. By Theorem 2, we need to check: (1) $q' \to_1 q$, and (2) $q \to_1 q'$. The 2ATA \mathcal{A}_q will be defined as the intersection of 2ATAs \mathcal{A}_1 and \mathcal{A}_2 , that check conditions (1) and (2), respectively.

Condition (1) is equivalent to $q' \to q$ (since $q' \in \mathsf{GHW}(1)$). The 2ATA \mathcal{A}_1 can guess and verify a homomorphism from q' to q. More precisely, it assumes that T' is annotated by an intended homomorphism $h: q' \to q$, that is, each variable x' in a node of T' is annotated by a variable x in q. The automaton then checks that this annotation encodes a homomorphism, i.e. that (a) all connected occurrences of x' are annotated by the same variable of q, and (b) for each atom $R(\bar{x}')$ labeling a node of T', the image of $R(\bar{x}')$ defined by the annotation is in q. For (a), when processing a node t' of T', the automaton stores the partial homomorphism for the variables of t' and tests that it is consistent with the partial homomorphism of each neighboring node of t'. In particular, \mathcal{A}_1 requires no alternation and has at most exponentially many states. If the maximum arity of the schema is fixed then only polynomially many states are needed, as then the stored partial homomorphisms are over constantly many elements and thus can be stored in $O(\log |q|)$ bits.

We now describe how the automaton A_2 works. First, as mentioned in Section 2.3, $q \to_1 q'$ can be characterized as Duplicator having a compact

winning strategy, which in turn can be characterized as follows [13]. Duplicator has a compact winning strategy on q and q', i.e. $q \to_1 q'$, iff there is a non-empty family \mathcal{F} of partial homomorphisms from q to q' such that: (a) the domain of each $f \in \mathcal{F}$ is a 1-union of q, and (b) if U and U' are 1-unions of q, then each $f \in \mathcal{F}$ with domain U can be extended to U', i.e., there is $f' \in \mathcal{F}$ with domain U' such that f(x) = f'(x) for every $x \in U \cap U'$.

The 2ATA A_2 assumes an annotation of $T' = \mathsf{Enc}(T_{q'}, \chi)$ that encodes the intended strategy \mathcal{F} . This annotation labels each node t' of T' by the set of partial mappings from q to q' whose domain is a 1-union of q, and whose range is contained in the variables from $\{u_1, \ldots, u_{2r}\}$ labeling t'. It can be easily checked from the labelings of T' if each mapping in this annotation is a partial homomorphism.

To check condition (2), the 2ATA A_2 makes a universal transition for each pair (U, U') of 1-unions and each partial mapping g with domain U annotating a node t of T'. Then it checks the existence of a node t' in T' that is annotated with a mapping g' that extends g to U'. The latter means that, for each $x \in U \cap U'$, both g(x) and g'(x) are the same variable of q', that is, g(x) and g(x') are connected occurrences of the same variable in $\{u_1, \ldots, u_{2r}\}$. Thus to check the consistency of g and g', the automaton can store the variables in $\{g(x) \mid x \in U \cap U'\}$, and check that these are present in the label of each node guessed before reaching t'. As this is a polynomial amount of information, A_2 can be implemented using exponentially many states. Again, if the maximum arity of the schema is fixed then only polynomially many states suffice.

The construction above can be easily extended from Boolean to non-Boolean queries (q, \bar{x}) and (q', \bar{x}') . In this case, the encoding $T' = \mathsf{Enc}(T_{q'}, \chi)$ of q' includes atoms that may contain free variables. The automaton \mathcal{A}_1 additionally checks that whenever a node in T' is annotated by an atom $R(\bar{y}')$ then there is an atom $R(\bar{y})$ in q such that $h'(\bar{y}') = \bar{y}$ where h' is the extension of the intended homomorphism h that also maps \bar{x}' to \bar{x} . The automaton \mathcal{A}_2 does an analogous check for the partial homomorphisms.

It is easy to see how Theorem 5 follows from Proposition 7. Checking if a CQ q has a GHW(1)-overapproximation amounts to checking if $L(\mathcal{A}_q) \neq \emptyset$. The latter can be done in exponential time in the number of states of \mathcal{A}_q [14], and thus in double-exponential time in the size of q. If $L(\mathcal{A}_q) \neq \emptyset$, one can construct a tree $T \in L(\mathcal{A}_q)$ in double-exponential time in the size of \mathcal{A}_q , and thus in triple-exponential time in the size of q. From T one then gets in polynomial time (i.e., in 3EXPTIME in the size of q) a GHW(1)-overapproximation of q.

If the maximum arity is fixed, the 2ATA \mathcal{A}_q has polynomially many states and, therefore, $L(\mathcal{A}_q) \neq \emptyset$ can be checked in single-exponential time. If $L(\mathcal{A}_q) \neq \emptyset$, one can then construct a tree $T \in L(\mathcal{A}_q)$ in double-exponential time in the size of q. From T one then gets in polynomial time (i.e., in 2EXPTIME in the size of q) a GHW(1)-overapproximation of q.

5.2 The case of Boolean CQs over binary schemas

For Boolean CQs over schemas of maximum arity two the existence and computation of GHW(1)-overapproximations can be solved in polynomial time. This is of practical importance since data models such as $graph\ databases\ [4]$ and description logic $ABoxes\ [2]$ can be represented using schemas of this kind. It is worth noticing that in this context GHW(1) coincides with the class of CQs of treewidth one [15].

Theorem 6 There is a PTIME algorithm that checks if a Boolean CQ q over a schema of maximum arity two has a GHW(1)-overapproximation q', and computes such a q' if it exists.

We devote the rest of this section to prove Theorem 6. Let q be a Boolean CQ. We define the Gaifman graph $\mathcal{G}(q)$ of q to be the undirected graph whose nodes are the variables of q and the edges are the pairs $\{z,z'\}$ such that $z \neq z'$ and z and z' appear together in some atom of q. A connected component of q is a Boolean CQ associated with a connected component C = (V(C), E(C)) of $\mathcal{G}(q)$, i.e., one whose set of variables is V(C) and contains precisely all the atoms in q induced by variables in V(C). The Boolean CQ q is connected if it has only one connected component, that is, if $\mathcal{G}(q)$ is connected.

When the maximum arity is two, we have that a Boolean CQ q is in GHW(1) iff $\mathcal{G}(q)$ is an *acyclic* (undirected) graph. In particular, if q is connected then $q \in \mathsf{GHW}(1)$ iff $\mathcal{G}(q)$ is a tree.

To prove the theorem, we first show how the problem can be solved in polynomial time for connected Boolean CQs, and then explain how to reduce in polynomial time the problem for general Boolean CQs to connected ones.

5.2.1 A polynomial time algorithm for connected Boolean CQs

We start with the following observation:

Lemma 3 Let us assume that q is a connected Boolean CQ that has an GHW(1)-overapproximation. Then it is the case that q has a connected GHW(1)-overapproximation.

Proof Let q' be an GHW(1)-overapproximation of q. Without loss of generality, we can assume that q' is a core. By contradiction, suppose that q' is not connected. By Theorem 2, we have that $q \to_1 q'$ and $q' \to_1 q$. Since $q' \in \text{GHW}(1)$, the latter is equivalent to $q' \to q$. We claim that there is a connected component q'_0 of q' such that $q \to_1 q'_0$. Recall from Section 2.3 that $q \to_1 q'$ can be witnessed by a compact winning strategy \mathcal{H} of the Duplicator. We make the Spoiler play in an arbitrary non-empty 1-union S_0 of q. Note that in this case, a 1-union is either a singleton or an edge of $\mathcal{G}(q)$. The Duplicator can respond, following \mathcal{H} , with a partial homomorphism h_0 from q to q' with domain S_0 . The elements in $h_0(S_0)$ must belong to some connected component of q', say q'_0 . Starting from this configuration of the game, we assume that Spoiler plays

in a connected manner, i.e., if S and S' are two consecutive moves for Spoiler then $S \cap S' \neq \emptyset$. Then the Duplicator can play indefinitely by following \mathcal{H} . By the way Spoiler plays, all responses of Duplicator must fall in q'_0 . Also, since q is connected, each 1-union of q is eventually played by the Spoiler. This implies that $q \to_1 q'_0$.

We have on the other hand that $q'_0 \to q$, and hence $q' \equiv q'_0$. Since q' is not connected, q'_0 has fewer atoms than q'. This contradicts the fact that q' is a core. We conclude that q' must be connected.

The high-level idea of the construction is to show that whenever a connected Boolean CQ q has an GHW(1)-overapproximation q', then we can assume that q' is a "subquery" of q, or of a slight modification of q. This will allow us to design a polynomial time algorithm that greedily looks for an GHW(1)-overapproximation of q. Note that Lemma 3 tells us that $q' \in \text{GHW}(1)$ can be assumed to be a connected Boolean CQ. In order to show that q' is a "subquery" of q, we first show a key lemma (see Lemma 5 below) about the structure of the endomorphisms of a connected core in GHW(1). In particular, we prove that besides the identity mapping, there can be only one extra endomorphism of a very particular form. Recall that an endomorphism is a homomorphism from the Boolean CQ to itself. For a core, any endomorphism h is actually an isomorphism, i.e., a bijection such that h^{-1} is a homomorphism.

We first need to establish the following auxiliary lemma.

Lemma 4 Let q be a connected Boolean CQ in GHW(1) that is a core. Let u and v be variables of q. Then there is at most one endomorphism of q mapping u to v.

Proof By contradiction, assume that there are two distinct endomorphisms h_1 and h_2 of q with $h_1(u) = h_2(u) = v$. Recall that, since q is a core, h_1 and h_2 are isomorphisms. By hypothesis, $\mathcal{G}(q)$ is a tree and we root it at u. Since $h_1 \neq h_2$, there must be a variable $w \neq u$ of q with $h_1(w) \neq h_2(w)$, which we choose to have minimal distance to u in the tree $\mathcal{G}(q)$. As h_2 is a bijection, there is a unique w' such that $h_2(w') = h_1(w)$. Note that $w' \neq w$. We claim that w and w' have the same parent in $\mathcal{G}(q)$. Let z be the parent of w. Since h_1 is an isomorphism, $h_1(w)$ and $h_1(z)$ are adjacent in $\mathcal{G}(q)$. As $h_1(w) = h_2(w')$ and, by minimality of w, $h_1(z) = h_2(z)$, we have that $h_2(w')$ and $h_2(z)$ are also adjacent. Since h_2 is an isomorphism, it follows that w' and z are adjacent in $\mathcal{G}(q)$. However, w' cannot be the parent of z since $h_1(w') \neq h_2(w')$. Therefore z is the parent of w'.

We define a mapping h from q to itself such that $h(t) = h_2(t)$ if t is a variable that belongs to the subtree of $\mathcal{G}(q)$ rooted at w'; otherwise, $h(t) = h_1(t)$. Note that h is an endomorphism of q. Indeed, the only atoms that in principle are not satisfied by h are those mentioning w' and z. However, since h_2 is an endomorphism and $h_1(z) = h_2(z)$, these atoms are actually satisfied. Finally, observe that

$$h(w) = h_1(w) = h_2(w') = h(w'),$$

and then h is not injective. This is a contradiction to q being a core.

Let $q \in \mathsf{GHW}(1)$ be a Boolean connected core. We say that an endomorphism h of q is a swapping endomorphism for u and v if h(u) = v and $\{u, v\}$ is an edge in $\mathcal{G}(q)$, i.e., u and v are adjacent variables in $\mathcal{G}(q)$. Using the fact that $\mathcal{G}(q)$ is a tree and h is an isomorphism, it can be seen that h(v) = u must also hold in this case (and hence the name "swapping"). Moreover, if such h exists for u and v, by Lemma 4, it must be unique and then we can speak about the swapping endomorphism for u and v. We then have the following:

Lemma 5 Let q be a Boolean connected CQ in GHW(1) that is a core. Then q has at most one endomorphism besides the identity mapping. If such endomorphism exists, it is the swapping endomorphism for some u and v.

Proof Fix a simple path $P = w_0, w_1, \ldots, w_m$ in $\mathcal{G}(q)$ of maximal length (recall that in a simple path all vertices are distinct). Suppose that there exists an endomorphism h of q different from the identity. As q is a core, h is an isomorphism. Then the path $P' = h(w_0), h(w_1), \ldots, h(w_m)$ is a simple path of the same length. Furthermore, P and P' share a vertex. Indeed, if this is not the case, since $\mathcal{G}(q)$ is connected, one can pick w in P and w' in P' such that w and w' are connected by a simple path P'' which is vertex-disjoint from P and P' (except for w and w'), and construct a longer path than P.

We claim that m is odd (recall that m is the size of P). By contradiction, suppose m is even and let $u = w_{m/2}$ be the middle vertex of P. It must be the case that u is also the middle vertex of P', as otherwise $\mathcal{G}(q)$ would contain a path longer than P. In particular, h(u) = u and then h must be the identity mapping by Lemma 4; a contradiction.

Let $u=w_{\lfloor m/2\rfloor}$ and $v=w_{\lceil m/2\rceil}$ be the middle vertices of P. Again, by maximality of P, we have that u and v are also the middle vertices of P', i.e., $\{u,v\}=\{h(u),h(v)\}$. Since h is not the identity and by Lemma 4, it follows that h(u)=v (and hence, h(v)=u). Since u and v are adjacent in $\mathcal{G}(q)$, we conclude that h must be the swapping endomorphism for u and v.

Let $q \in \mathsf{GHW}(1)$ be a connected Boolean CQ. Let u and v be variables adjacent in $\mathcal{G}(q)$. Since $\mathcal{G}(q)$ is a tree, if we remove from q all the atoms that mention u and v simultaneously, we obtain two connected Boolean CQs, one containing u and the other containing v. We denote these CQs by t_u^q and t_v^q , respectively.

We need to introduce some notation. Suppose that q and q' are Boolean CQs and X and X' are subsets of the variables of q and q', respectively. We denote by $(q, X) \to_1 (q', X')$ the fact that the Duplicator has a winning strategy in the existential 1-cover game on q and q' with the property that whenever the Spoiler places a pebble on an element of X in q, then the Duplicator responds with some element of X' in q'. It can be seen that checking whether $(q, X) \to_1 (q', X')$ can still be done in polynomial time.

The following lemma formalizes the idea that an GHW(1)-overapproximation can be assumed to be essentially a subquery of the original query.

Lemma 6 Suppose q is a Boolean CQ and suppose q' is a connected core that is a GHW(1)-overapproximation of q. Then we have the following:

1. If the only endomorphism of q' is the identity mapping, then any homomorphism from q' to q is injective.

- 2. If q' has the swapping endomorphism for some u' and v', then for any homomorphism h from q' to q, we have that
 - (a) $(q, \{h(u'), h(v')\}) \rightarrow_1 (q', \{u', v'\}), \text{ and }$
 - (b) $h(z') \neq h(z'')$ for all pairs of variables z', z'', except maybe for $z' \neq u'$ in $t_{n'}^{q'}$ and $z'' \neq v'$ in $t_{n'}^{q'}$.

Proof Suppose the only endomorphism of q' is the identity and let h be a homomorphism from q' to q. Towards a contradiction, suppose h(z') = h(z'') = b for distinct variables z' and z'' in q'. Let \mathcal{H} be a winning strategy of Duplicator witnessing $q \to_1 q'$. We choose any variable b' in q' such that b' is a possible response of Duplicator according to \mathcal{H} , when Spoiler starts playing on b in q. Suppose that $b' \neq z'$ (the case $b' \neq z''$ is analogous). Then by composing h with \mathcal{H} , we obtain a winning strategy for Duplicator in the game on q' and q' such that b' is a possible response of Duplicator when Spoiler starts playing on z'. Since $q' \in \mathsf{GHW}(1)$, we can define an endomorphism g of g' that maps g' to g'. Then g' is an endomorphism different from the identity, which is a contradiction.

Suppose now that q' has the swapping endomorphism for some u' and v', and let h be a homomorphism from q' to q. First, assume by contradiction that Duplicator's strategy witnessing $q \to_1 q'$ is such that for h(u') (the case for h(v') is analogous), Duplicator responds with $z' \notin \{u', v'\}$. By composing h with this strategy, and using the fact that $q' \in \mathsf{GHW}(1)$, it follows that there is an endomorphism g of q' that maps u' to z'. This endomorphism is different from the identity and from the swapping endomorphism for u' and v', which contradicts Lemma 5. Finally, suppose towards a contradiction that item (2.b) does not hold for some pair $z' \neq z''$ with h(z') = h(z''). We have two cases.

$$\begin{array}{ll} \text{(i)} \ \ z',z'' \in t_{u'}^{q'} \ \text{or} \ z',z'' \in t_{v'}^{q'}; \ \text{or} \\ \text{(ii)} \ \ z' \in t_{u'}^{q'} \ \text{and} \ z'' \in t_{v'}^{q'}, \ \text{and either} \ z' = u' \ \text{or} \ z'' = v'. \end{array}$$

In any case, we can again compose h with the strategy witnessing $q \to_1 q'$ and use the fact that $q' \in \mathsf{GHW}(1)$, to derive an endomorphism of q' that is neither the identity nor the swapping endomorphism for u' and v', which is a contradiction.

Let q be a Boolean CQ and u and v be adjacent variables in $\mathcal{G}(q)$. We define a CQ $q_u \# q_v$ as follows. Denote by $q \setminus v$ the CQ obtained from q by removing all atoms that contain v. Let q_u be the query constructed from $q \setminus v$ by replacing each variable z in $q \setminus v$ by a fresh variable z_u . Similarly, let q_v be the CQ where each variable z in $q \setminus u$ is replaced by a fresh variable z_v . The CQ $q_u \# q_v$ contains all the atoms of q_u and q_v , and additionally, all atoms $R(u_u, v_v)$ or $R(v_v, u_u)$ whenever R(u, v) or R(v, u) is an atom in q, respectively. Note that by mapping variables z_u and z_v to z, we have that $q_u \# q_v \to q$.

Now we are ready to present our algorithm. Observe that Lemma 6 implies that whenever q' is an GHW(1)-overapproximation of q, we can assume that

either q' is a subquery of q (item (1)), or a subquery of $q_u \# q_v$ for some u and v (item (2)). The algorithm then greedily searches through the subqueries of q and $q_u \# q_v$, for all u and v, to find an $\mathsf{GHW}(1)$ -overapproximation of q.

The algorithm. Let q be a connected Boolean CQ. The algorithm first checks whether a subquery of q is an GHW(1)-overapproximation. This is Step 1. In Step 2, the algorithm checks whether a subquery of $q_u \# q_v$ is an GHW(1)-overapproximation, for some u and v in q. If neither step succeed then the algorithm rejects. For a Boolean CQ p and an atom e of p, we denote by $p \setminus e$ the Boolean CQ obtained from p by removing e. Step 1 is as follows:

- 1. Set q_0 to be q.
- 2. While $q_i \notin \mathsf{GHW}(1)$, search for an atom e such that $q_i \to_1 q_i \setminus e$. If there is no such atom then continue with Step 2. Otherwise, set q_{i+1} to be $q_i \setminus e$.
- 3. If $q_i \in GHW(1)$, for some i, then accept and output q_i .

For Step 2, let \mathcal{P} be an enumeration of the pairs (u, v) such that u, v are adjacent in $\mathcal{G}(q)$ and $q \to_1 q_u \# q_v$. Step 2 is as follows:

- 1. Let (u, v) be the first pair in \mathcal{P} .
- 2. Set q_0 to be $q_u \# q_v$.
- 3. While $q_i \notin \mathsf{GHW}(1)$, search for an atom e that does not mention u_u and v_v simultaneously such that $(q_i, \{u_u, v_v\}) \to_1 (q_i \setminus e, \{u_u, v_v\})$. If there is no such atom, let (u, v) be the next pair in \mathcal{P} and repeat from item 2. Otherwise, set q_{i+1} to be $q_i \setminus e$.
- 4. If $q_i \in GHW(1)$, for some i, then accept and output q_i .

Notice that the described algorithm can be implemented in polynomial time. Below we argue that it is correct.

Suppose first that the algorithm, on input q, accepts with output q^* . By construction $q^* \in \mathsf{GHW}(1)$. Assume first that the algorithm accepts in the m-th iteration of Step 1, and thus $q^* = q_m$. By construction, for each $0 \le i < m$, we have that $q_i \to_1 q_{i+1}$ and $q_{i+1} \to_1 q_i$. In particular, $q \to_1 q^*$ and $q^* \to_1 q$, and thus q^* is a $\mathsf{GHW}(1)$ -overapproximation of q. Suppose now that the algorithm accepts in Step 2 for a pair $(u,v) \in \mathcal{P}$, in the m-th iteration. Again we have that $q_i \to_1 q_{i+1}$ and $q_{i+1} \to_1 q_i$, for each $0 \le i < m$, and thus

$$q_u \# q_v \to_1 q^*$$
 and $q^* \to_1 q_u \# q_v$.

Since $(u,v) \in \mathcal{P}$, it follows that $q \to_1 q_u \# q_v$, and then $q \to_1 q^*$. Using the fact that $q_u \# q_v \to q$, we have that $q^* \to_1 q$. Hence, q^* is a GHW(1)-overapproximation of q.

It remains to show that if q has an $\mathsf{GHW}(1)$ -overapproximation q' then the algorithm accepts. Since q is connected, by Lemma 3, we can assume that q' also is. Moreover, we can assume without loss of generality that q' is a core. By Lemma 5, we have two cases:

(1) the only endomorphism of q' is the identity, or

(2) q' has two endomorphisms, namely, the identity and the swapping endomorphism for some variables u' and v'.

First suppose case (1) applies. We show that the algorithm accepts in Step 1. By definition, $q_i \to_1 q_{i+1}$ and $q_{i+1} \to_1 q_i$ (actually $q_{i+1} \to q_i$), for each $0 \le i \le m$, where m is the number of iteration in Step 1. It follows that

$$q_0 = q \to_1 q_m$$
 and $q_m \to_1 q$.

Since the relation \to_1 composes, q' is a GHW(1)-overapproximation of q_m and by using Lemma 6, q' is a subquery of q_m . Now for the sake of contradiction assume that the algorithm does not accept in Step 1. Then $q_m \notin \text{GHW}(1)$ and there is no edge e in q_m such that $q_m \to_1 q_m \setminus e$. Since q' is GHW(1)-overapproximation of q_m , we have that $q_m \to_1 q'$ and, since $q' \in \text{GHW}(1)$, q' is a proper subquery of q_m . It follows that there is an edge e in q_m such that $q_m \to_1 q_m \setminus e$, which is a contradiction.

Suppose case (2) holds. In this case the algorithm accepts in Step 2. Let h be a homomorphism from q' to q, and let u = h(u') and v = h(v'). By Lemma 6, $u \neq v$ and then u and v are adjacent. Also, by Lemma 6, q' is a subquery of $q_u \# q_v$. Since $q \to_1 q'$, it follows that $q \to_1 q_u \# q_v$, and then $(u, v) \in \mathcal{P}$. We claim that the algorithm accepts when (u, v) is chosen from \mathcal{P} . First, note that q' is a GHW(1)-overapproximation of q_m . Indeed, by definition,

$$q_m \to q_u \# q_v$$
, $q_u \# q_v \to q$, and $q \to_1 q'$.

It follows that $q_m \to_1 q'$. On the other hand, we have that

$$(q', (u', v')) \to (q_u \# q_v, (u_u, v_v)),$$

as q' is a subquery of $q_u \# q_v$, and

$$(q_u \# q_v, \{u_u, v_v\}) \to_1 (q_m, \{u_u, v_v\}).$$

It follows that $(q', \{u', v'\}) \to_1 (q_m, \{u_u, v_v\})$, which implies that $(q', (u', v')) \to (q_m, (u_u, v_v))$ via a homomorphism g. Then q' is a GHW(1)-overapproximation of q_m . By applying Lemma 6 to q_m, q' and g, we obtain that

$$(q_m, \{u_u, v_v\}) \to_1 (q', \{u', v'\}),$$

and g satisfies item (2.b). Observe that $g(z') \neq g(z'')$ for all $z' \neq u'$ in $t_{u'}^{q'}$ and $z'' \neq v'$ in $t_{v'}^{q'}$, since $\{u_u, v_v\}$ is a bridge of $\mathcal{G}(q_m)$, i.e., its removal disconnects $\mathcal{G}(q_m)$. We conclude that g is injective and then q' is a subquery of q_m .

Towards a contradiction, assume that the algorithm do not accept when (u, v) is chosen from \mathcal{P} . Then $q_m \notin \mathsf{GHW}(1)$ and there is no edge e that does not mention both u_u, v_v such that $(q_m, \{u_u, v_v\}) \to_1 (q_m \setminus e, \{u_u, v_v\})$. Since

$$(q_m, \{u_u, v_v\}) \to_1 (q', \{u', v'\})$$
 and $(q', (u', v')) \to (q_m, (u_u, v_v))$

via the injective homomorphism g and $q' \in \mathsf{GHW}(1)$, it follows that there is an edge e that does not mention both u_u, v_v such that $(q_m, \{u_u, v_v\}) \to_1 (q_m \setminus e, \{u_u, v_v\})$. This is a contradiction.

5.2.2 Reduction to the connected case

Now we consider the non-connected case. Given a Boolean CQ q with connected components q_1, \ldots, q_m , the algorithm proceeds as follows:

- 1. Start by simplifying q: Compute a minimal subset of Boolean CQs \mathcal{Q} in $\{q_1,\ldots,q_m\}$ such that for each $1\leq i\leq m$, there is a $p\in\mathcal{Q}$ with $q_i\to_1 p$.
- 2. Check whether each $p \in \mathcal{Q}$ has an GHW(1)-overapproximation p' using the algorithm described for connected Boolean CQs in Section 5.2.1. If this is the case then accept and output the disjoint conjunction $\bigwedge_{p \in \mathcal{Q}} p'$.

Clearly, the algorithm can be implemented in polynomial time. For the correctness, suppose first that the algorithm accepts and outputs $q' = \bigwedge_{p \in \mathcal{Q}} p'$. Then

$$q' \to \bigwedge_{p \in \mathcal{Q}} p \to q.$$

We also have that $q \to_1 \bigwedge_{p \in \mathcal{Q}} p$ (by definition of \mathcal{Q}), and $\bigwedge_{p \in \mathcal{Q}} p \to_1 q'$. This implies that q' is a $\mathsf{GHW}(1)$ -overapproximation of q.

Suppose now that q has an GHW(1)-overapproximation q'. Since both $q \to_1 \bigwedge_{p \in \mathcal{Q}} p$ and $\bigwedge_{p \in \mathcal{Q}} p \to_1 q$ hold, it is the case that q' is also an GHW(1)-overapproximation of $\bigwedge_{p \in \mathcal{Q}} p$. By the minimality of \mathcal{Q} , we have that $p \not\to_1 \hat{p}$, for each pair of distinct CQs $p, \hat{p} \in \mathcal{Q}$. Let p be a CQ in \mathcal{Q} . Since $p \to_1 q'$ and p is connected, it follows that $p \to_1 p^*$, where p^* is a connected component of q' (note that this follows by using the same argument we use in the proof of Lemma 3 to show that $q \to_1 q'_0$, for some component q'_0). Also, since $q' \to \bigwedge_{p \in \mathcal{Q}} p$, there is $p_0 \in \mathcal{Q}$ such that $p^* \to p_0$. In particular, $p \to_1 p_0$. It follows that $p_0 = p$, and then p^* is an GHW(1)-overapproximation of p. We conclude that each $p \in \mathcal{Q}$ has an GHW(1)-overpproximation, and thus the algorithm accepts. This finishes the proof of Theorem 6.

Remark The restriction to Boolean CQs in Theorem 6 is used in an essential way in our proof, and it is not clear whether this extends to non-Boolean CQs. The issue is that we do not have an analog of Lemma 3 for the non-Boolean case, and hence, it is not clear how to translate Lemma 5 and Lemma 6 (or some modifications thereof) into a polynomial time algorithm.

5.3 Size of overapproximations

Over binary schemas GHW(1)-overapproximations are of polynomial size. This is optimal as over schemas of arity three there is an exponential lower bound for the size of GHW(1)-overapproximations:

Proposition 8 There is a schema σ with a single ternary relation symbol and a family $(q_n)_{n\geq 1}$ of Boolean CQs over σ , such that (1) q_n is of size O(n), and (2) the size of every GHW(1)-overapproximation of q_n is $\Omega(2^n)$.




Fig. 4 Illustration of the CQs q_3 and q'_3 from Proposition 8. Each triple of variables represents two atoms in the query; e.g., $\{y_0, y_1^1, y_1^2\}$ represents atoms $R(y_0, y_1^1, y_1^2)$ and $R(y_0, y_1^2, y_1^1)$ in q_3' .

Proof The CQ q_n contains the atoms:

 $-R(x_0,x_1^1,x_1^2),\,R(x_0,x_1^2,x_1^1),\,\text{as well as}\\ -R(x_i^j,x_{i+1}^1,x_{i+1}^2)\,\,\text{and}\,\,R(x_i^j,x_{i+1}^2,x_{i+1}^1),\,\text{for each}\,\,1\leq i\leq n-1\,\,\text{and}\,\,j\in\{0,1,2,\ldots,n\}$

Consider now the CQ q'_n with the atoms:

 $\begin{array}{l} - \ R(y_0,y_1^1,y_1^2), \ R(y_0,y_1^2,y_1^1), \ \text{and} \\ - \ R(y_{|w|}^w,y_{|w|+1}^{w1},y_{|w|+1}^{w2}) \ \text{and} \ R(y_{|w|}^w,y_{|w|+1}^{w2},y_{|w|+1}^{w1}), \text{for each word } w \ \text{over} \ \{1,2\} \\ \text{of length} \ 1 \leq |w| \leq n-1. \end{array}$

Figure 4 depicts the CQs q_3 and q'_3 .

Clearly, the mapping $h: q'_n \to q_n$ defined as $h(y_0) = x_0$ and

$$h(y_{|w|+1}^{wj}) = x_{|w|+1}^j,$$

for each word w over $\{1,2\}$ of length $0 \le |w| \le n-1$ and $j \in \{1,2\}$, is a homomorphism. We now show that $q_n \to_1 q'_n$ by building a compact winning strategy for Duplicator (see the proof of Proposition 7) which basically "inverts" the homomorphism h. It contains: (a) Partial homomorphisms $(x_0, x_1^1, x_1^2) \to (y_0, y_1^1, y_1^2)$ and $(x_0, x_1^1, x_1^2) \to (y_0, y_1^2, y_1^1)$, and (b) for each word w over $\{1,2\}$ of length $1 \leq |w| \leq n-1$ and $j \in \{1,2\}$, the partial homomor-

$$\begin{array}{l} -\ (x_{|w|}^j,x_{|w|+1}^1,x_{|w|+1}^2) \to (y_{|w|}^w,y_{|w|+1}^{w1},y_{|w|+1}^{w2}), \text{ and} \\ -\ (x_{|w|}^j,x_{|w|+1}^1,x_{|w|+1}^2) \to (y_{|w|}^w,y_{|w|+1}^{w2},y_{|w|+1}^{w1}). \end{array}$$

It can be seen that this is a winning strategy for Duplicator.

Observe that the size of q'_n is $\Omega(2^n)$. A straightforward case-by-case analysis shows that q'_n is a core, i.e., there is no homomorphism from q'_n to a proper subset of its atoms. We claim that q_n' is the smallest $\mathsf{GHW}(1)$ overapproximation of q_n , from which the proposition follows. Assume, towards a contradiction, that q' is a $\mathsf{GHW}(1)$ -over approximation of q_n with fewer atoms than q'_n . Then, by Corollary 1, we have that $q'_n \equiv q'$. Composing the homomorphisms $h_1: q_n \to q'$ and $h_2: q' \to q'_n$ yields a homomorphism from q'_n to a proper subset of the atoms of q'_n . This is a contradiction since q'_n is a core. \square

5.4 Beyond acyclicity

Theorem 2 characterizes when a CQ has a $\mathsf{GHW}(k)$ -overapproximation. We provide an alternative characterization in terms of a boundedness condition for the existential cover game. This helps understanding where lies the difficulty of determining the decidability status of the problem of existence of $\mathsf{GHW}(k)$ -overapproximations, for k > 1.

Recall that we write $(\mathcal{D}, \bar{a}) \to_k^c (\mathcal{D}', \bar{b})$, for $k \geq 1$ and $c \geq 1$, if Duplicator has a winning strategy in the first c rounds of the (compact) existential k-cover game on (\mathcal{D}, \bar{a}) and (\mathcal{D}', \bar{b}) . The next result establishes that a CQ q has a GHW(k)-overapproximation iff the existential k-cover game played from q is "bounded", i.e., if there is a constant $c \geq 1$ that bounds the number of rounds this game needs to be played in order to determine if Duplicator wins.

Theorem 7 Fix $k \geq 1$. The CQ $q(\bar{x})$ has a $\mathsf{GHW}(k)$ -overapproximation iff there is an integer $c \geq 1$ such that

$$(q, \bar{x}) \to_k (\mathcal{D}, \bar{a}) \iff (q, \bar{x}) \to_k^c (\mathcal{D}, \bar{a}),$$

for each database \mathcal{D} and tuple \bar{a} of elements in \mathcal{D} .

Proof Assume first that there is an integer $c \geq 1$ such that $(q, \bar{x}) \to_k (\mathcal{D}, \bar{a})$ iff $(q, \bar{x}) \to_k^c (\mathcal{D}, \bar{a})$, for each database \mathcal{D} and tuple $\bar{a} \in \mathcal{D}$. Therefore, $(q, \bar{x}) \to_k (\mathcal{D}, \bar{a})$ iff $(q_c, \bar{x}_c) \to (\mathcal{D}, \bar{a})$, for each database \mathcal{D} and $\bar{a} \in \mathcal{D}$, where $q_c(\bar{x}_c)$ is the CQ in $\mathsf{GHW}(k)$ which is defined in Lemma 1. But then, since $q_c \in \mathsf{GHW}(k)$, we obtain from Equation (2) that for each database \mathcal{D} and tuple $\bar{a} \in \mathcal{D}$:

$$(q, \bar{x}) \to_k (\mathcal{D}, \bar{a}) \iff (q_c, \bar{x}_c) \to_k (\mathcal{D}, \bar{a}).$$
 (3)

We show next that q_c is the $\mathsf{GHW}(k)$ -overapproximation of q. From Theorem 2, we need to show that $(q_c, \bar{x}_c) \to_k (q, \bar{x})$ and $(q, \bar{x}) \to_k (q_c, \bar{x}_c)$. Both conditions follow from Equation (3): the former by choosing (\mathcal{D}, \bar{a}) to be (q, \bar{x}) , and the latter by choosing (\mathcal{D}, \bar{a}) to be (q_c, \bar{x}_c) .

Assume on the other hand that $q(\bar{x})$ has a $\mathsf{GHW}(k)$ -overapproximation $q^*(\bar{x}^*)$. From Theorem 3 we obtain that for every (finite) database \mathcal{D} and tuple \bar{a} of elements in \mathcal{D} :

$$(q^*, \bar{x}^*) \to (\mathcal{D}, \bar{a}) \iff (q, \bar{x}) \to_k (\mathcal{D}, \bar{a}).$$

As a first step we prove that this continues to hold for countably infinite databases. We do so by refining the proof of Theorem 3.

Let \mathcal{D} be a countably infinite database. Assume first that $(q, \bar{x}) \to_k (\mathcal{D}, \bar{a})$. Due to the fact that q^* is a $\mathsf{GHW}(k)$ -overapproximation of q, we have that $(q^*, \bar{x}^*) \to (q, \bar{x})$. Proposition 6 then implies that $(q^*, \bar{x}^*) \to (\mathcal{D}, \bar{a})$ since q^* is in $\mathsf{GHW}(k)$. Assume, on the other hand, that $(q^*, \bar{x}^*) \to (\mathcal{D}, \bar{a})$. Suppose, for the sake of contradiction, that $(q, \bar{x}) \not\to_k (\mathcal{D}, \bar{a})$. Proposition 6 then establishes that there is a $\mathsf{CQ}\ q^\#(\bar{x}^\#)$ in $\mathsf{GHW}(k)^\infty$ such that

$$(q^{\#}, \bar{x}^{\#}) \to (q, \bar{x}), \text{ but } (q^{\#}, \bar{x}^{\#}) \not\to (\mathcal{D}, \bar{a}).$$
 (4)

The first fact in Equation (4) implies that $q \subseteq q^{\#}$, as this direction of Equation 1 continues being true as long as the CQ q in the left-hand side of the containment relation is finite. Therefore, $(q^{\#}, \bar{x}^{\#}) \to (q^{*}, \bar{x}^{*})$ since Corollary 1 continues being true for CQs in $\mathsf{GHW}(k)^{\infty}$. We conclude that $(q^{\#}, \bar{x}^{\#}) \to (\mathcal{D}, \bar{a})$ since $(q^{*}, \bar{x}^{*}) \to (\mathcal{D}, \bar{a})$. This contradicts the second fact in Equation (4).

Therefore, if $q'(\bar{x}')$ is the CQ given by Lemma 2 for CQ $q(\bar{x})$, then for every countable database \mathcal{D} and tuple \bar{a} in \mathcal{D} it is the case that

$$(q', \bar{x}') \to (\mathcal{D}, \bar{a}) \iff (q, \bar{x}) \to_k (\mathcal{D}, \bar{a}) \iff (q^*, \bar{x}^*) \to (\mathcal{D}, \bar{a}).$$

In particular, $(q', \bar{x}') \to (q^*, \bar{x}^*)$ and $(q^*, \bar{x}^*) \to (q', \bar{x}')$ by choosing (\mathcal{D}, \bar{a}) as (q^*, \bar{x}^*) and (q', \bar{x}') , respectively, in the previous equation. By composing $(q', \bar{x}') \to (q^*, \bar{x}^*)$ with $(q^*, \bar{x}^*) \to (q', \bar{x}')$, we obtain that $(q', \bar{x}') \to (q'_{\text{fin}}, \bar{x}')$, where q'_{fin} is a CQ with finitely many atoms, all of which belong to q'.

Recall from the proof of Lemma 2 that the atoms of the CQ q' are obtained as the union of the Q_c s, for $c \geq 0$, where Q_c is the set of atoms in the CQ q_c , i.e., the one that describes the first c rounds of the existential k-cover game played from q. Therefore, there must be an integer $c \geq 0$ such that $(q', \bar{x}') \to (q_c, \bar{x}_c)$. For the same reason, it holds that $(q'_c, \bar{x}_c) \to (q', \bar{x}')$. We conclude that for every database \mathcal{D} and tuple \bar{a} of elements in \mathcal{D} :

$$(q', \bar{x}') \to (\mathcal{D}, \bar{a}) \iff (q_c, \bar{x}_c) \to (\mathcal{D}, \bar{a}).$$

In other words, over every database \mathcal{D} and tuple \bar{a} of elements in \mathcal{D} , it is the case that

$$(q, \bar{x}) \to_k (\mathcal{D}, \bar{a}) \iff (q, \bar{x}) \to_k^c (\mathcal{D}, \bar{a}).$$

This finishes the proof of the theorem.

Boundedness conditions are a difficult area of study, with a delicate decidability boundary; e.g., boundedness is decidable for Datalog programs if all intensional predicates are monadic [14], but undecidable if binary intensional predicates are allowed [22]. For least fixed point logic (LFP), undecidability results for boundedness abound with the exception of a few restricted fragments [38,9]. Although the existence of winning Duplicator strategies in existential pebble games is expressible in LFP [35], no result obtained in such context seems to be directly applicable to determining the decidability status of the boundedness condition in Theorem 7.

6 Overapproximations under Constraints

It has been observed that semantic information about the data – in particular, in the form of constraints – enriches the quality of approximations [5]. This is exemplified next.

Example 3 As mentioned before, the Boolean CQ

$$q = \exists x \exists y \exists z (E(x,y) \land E(y,z) \land E(z,x)),$$

shown in Figure 3, has no GHW(1)-overapproximation. On the other hand, if we know that the data satisfies the constraint

$$\forall x \forall y \forall z (E(x,y) \land E(y,z) \rightarrow E(z,x)),$$

then q becomes equivalent to the CQ $q' = \exists x \exists y \exists z (E(x,y) \land E(y,z))$, which is in GHW(1).

In this section we study the notion of GHW(k)-overapproximation under constraints. We consider the two most important classes of database constraints; namely:

1. Tuple-generating dependencies (tgds), i.e., expressions of the form

$$\forall \bar{x} \forall \bar{y} \left(\phi(\bar{x}, \bar{y}) \to \exists \bar{z} \psi(\bar{x}, \bar{z}) \right), \tag{5}$$

where ϕ and ψ are conjunctions of atoms. Notice that the constraint in Example 3 is a tgd.

Tgds subsume the central class of inclusion dependencies (IDs) [17]. For example, assuming that R and P are binary relations, the ID $R[1] \subseteq P[2]$, which states that the set of values occurring in the first attribute of R is a subset of the set of values in the second attribute of P, is expressed via the tgd $\forall x \forall y (R(x,y) \rightarrow \exists z P(z,x))$.

There is a particular subclass of tgds that is expressive enough to subsume IDs and has received considerable attention in the literature; namely, the class of guarded tgds [10]. A tgd is guarded if its body $\phi(\bar{x}, \bar{y})$ contains an atom, called the guard, that mentions all the variables in $(\bar{x} \cup \bar{y})$. Notice that tgds that represent IDs are trivially guarded, as their body consists of a single atom.

2. Equality-generating dependencies (egds), i.e., expressions of the form

$$\forall \bar{x} \left(\phi(\bar{x}) \to y = z \right), \tag{6}$$

where ϕ is a CQ and y, z are variables in \bar{x} . Egds subsume the important classes of keys and functional dependencies (FDs).

For example, assuming that R is a ternary relation, the FD $R: \{1\} \to \{3\}$, i.e., the first attribute of R functionally determines the third attribute of R, is expressed via the egd

$$\forall x \forall y \forall z \forall y' \forall z' (R(x, y, z) \land R(x, y', z') \rightarrow z = z').$$

Notice that FDs that have more than one attribute in the right-hand side, can be expressed via a *set* of egds.

A (potentially infinite) database \mathcal{D} satisfies a tgd of the form given in Equation (5) if the following holds: for each tuple (\bar{a}, \bar{b}) of elements such that $(\phi, \bar{x}, \bar{y}) \to (\mathcal{D}, \bar{a}, \bar{b})$, there is a tuple \bar{c} of elements such that $(\psi, \bar{x}, \bar{z}) \to (\mathcal{D}, \bar{a}, \bar{c})$. Analogously, \mathcal{D} satisfies an egd as in Equation (6) if, for each tuple \bar{a} of elements such that $(\phi, \bar{x}) \to (\mathcal{D}, \bar{a})$ via a homomorphism h, it is the case that h(y) = h(z). Finally, \mathcal{D} satisfies a set \mathcal{L} of constraints if it satisfies every tgd and egd in \mathcal{L} .

CQ containment under constraints. The right notion of containment, under constraints, is measured over those databases that satisfy the constraints only (as we know that our datasets satisfy such constraints). Formally, let q, q' be CQs and Σ a set of constraints. Then q is contained in q' under Σ , denoted $q \subseteq_{\Sigma} q'$, if and only if $q(\mathcal{D}) \subseteq q'(\mathcal{D})$ for each database \mathcal{D} that satisfies Σ . It is worth remarking that, as before, containment is defined over finite databases only. The notion of equivalence is defined analogously, and we write $q \equiv_{\Sigma} q'$.

The chase procedure is a canonical tool for reasoning about CQ containment under constraints [37]. We start by defining a single chase step for tgds. Let q be a CQ and τ a tgd of the form $\forall \bar{x} \forall \bar{y} (\phi(\bar{x}, \bar{y}) \to \exists \bar{z} \psi(\bar{x}, \bar{z}))$. We say that τ is applicable with respect to q, if there exists a tuple (\bar{a}, \bar{b}) of elements in q such that $(\phi, \bar{x}, \bar{y}) \to (q, \bar{a}, \bar{b})$. In this case, the result of applying τ over q with (\bar{a}, \bar{b}) is the CQ q' that extends q with every atom in $\psi(\bar{a}, \bar{c})$, where \bar{c} is the tuple obtained by simultaneously replacing each $z \in \bar{z}$ with a fresh element not occurring in q. For such a single chase step we write $q \xrightarrow{\tau,(\bar{a},\bar{b})} q'$.

Let us now assume that q is a CQ and \varSigma a set of tgds. A chase sequence for q under \varSigma is a sequence

$$q_0 \xrightarrow{\tau_0,(\bar{a}_0,\bar{b}_0)} q_1 \xrightarrow{\tau_1,(\bar{a}_1,\bar{b}_1)} q_2 \dots$$

of chase steps such that:

- 1. $q_0 = q$.
- 2. For each $i \geq 0$ we have that τ_i is a tgd in Σ .
- 3. $q' \models \Sigma$, where q' is the CQ formed by the union of the atoms in the q_i s, for $i \geq 0$.

The (potentially infinite) CQ q' is the *result* of this chase sequence, which always exists.

Although the result of a chase sequence is not unique (up to isomorphism), each such result is equally useful for our purposes since it can be homomorphically embedded into every other result. This is a consequence of the fact that the result q' of a chase sequence for q under Σ is universal, i.e., for every (potentially infinite) CQ q'' such that $q \subseteq q''$ and $q'' \models \Sigma$, there is a homomorphism from q' to q'' [18,16]. Henceforth, we write $chase_{\Sigma}(q)$ for the result of an arbitrary chase sequence for q under Σ .

As for tgds, the chase is a useful tool when reasoning with egds. Let us first define a single chase step for egds. Consider a CQ q and an egd ϵ of the form $\forall \bar{x}(\phi(\bar{x}) \to x_i = x_j)$. We say that ϵ is applicable with respect to q,

if there exists a homomorphism h that witnesses $\phi \to q$ for which it holds that $h(x_i) \neq h(x_j)$. In this case, the result of applying ϵ over q with h is the CQ q' that obtained from q by identifying $h(x_i)$ and $h(x_j)$ everywhere. We can define the notion of the chase sequence for a CQ q under a set Σ of egds analogously as we did for tgds. Notice that such a sequence is finite and unique; thus, we refer to the chase for q under Σ , denoted $chase_{\Sigma}(q)$. Observe that the chase sequence that leads to $chase_{\Sigma}(q)$ gives rise to a homomorphism $h_{q,\Sigma}: q \to chase_{\Sigma}(q)$ such that $h_{q,\Sigma}(q) = chase_{\Sigma}(q)$.

It is well-known that an extended notion of containment under constraints – which is defined over both finite and infinite databases – can be characterized in terms of the notion of homomorphism and the chase procedure. Formally, let q, q' be CQs and Σ a set of constraints. We write $q \subseteq_{\Sigma}^{\infty} q'$ iff $q(\mathcal{D}) \subseteq q'(\mathcal{D})$ for each countable database \mathcal{D} that satisfies Σ . T

Lemma 7 Let $q(\bar{x}), q'(\bar{x}')$ be CQs.

1. If Σ is a set of tyds, then

$$q \subseteq_{\Sigma}^{\infty} q' \iff (q', \bar{x}') \to (chase_{\Sigma}(q), \bar{x}).$$

2. If Σ is a set of egds, then

$$q \subseteq_{\Sigma}^{\infty} q' \iff (q', \bar{x}') \to (chase_{\Sigma}(q), h_{q, \Sigma}(\bar{x})).$$

The sets Σ of constraints for which the notions \subseteq_{Σ} and $\subseteq_{\Sigma}^{\infty}$ coincide are called *finitely controllable*. It is easy to see that any set Σ of egds is finitely controllable, as $chase_{\Sigma}(q)$ is always finite in such a case. On the other hand, arbitrary sets of tgds are not necessarily finitely controllable. An important exception corresponds to the case when Σ is a set of guarded tgds. In fact, a deep result shows that sets of guarded tgds are finitely controllable, i.e., for any CQs q, q' and set Σ of guarded tgds, it holds that $q \subseteq_{\Sigma} q' \Leftrightarrow q \subseteq_{\Sigma}^{\infty} q'$ [3].

6.1 GHW(k)-overapproximations under constraints

In the absence of constraints, $\mathsf{GHW}(k)$ -overapproximations may not exist. However, as discussed in Section 4, by considering the class $\mathsf{GHW}(k)^\infty$ of CQs with countable many atoms and generalized hypertreewidth bounded by k, we can provide each CQ with a $\mathsf{GHW}(k)^\infty$ -overapproximation. We show that this good behavior generalizes to the case when constraints expressed as egds or guarded tgds are present.

Formally, given CQs q and q' such that $q' \in \mathsf{GHW}(k)^{\infty}$, we say that q' is a $\mathsf{GHW}(k)^{\infty}$ -overapproximation of q under Σ , if (i) $q \subseteq_{\Sigma} q'$, and (ii) there is no $q'' \in \mathsf{GHW}(k)^{\infty}$ such that $q \subseteq_{\Sigma} q'' \subsetneq_{\Sigma} q'$.

Theorem 8 Fix $k \geq 1$. For every $CQ\ q(\bar{x})$ and set Σ consisting exclusively of egds or guarded tgds, there is a $CQ\ q'(\bar{x}') \in \mathsf{GHW}(k)^{\infty}$ that is a $\mathsf{GHW}(k)^{\infty}$ -overapproximation of q under Σ .

Proof We only consider the case when Σ is a set of guarded tgds, as for egds the proof is even simpler. As in Lemma 2, we can show that there is a CQ $q'(\bar{x}')$ such that for every countable database \mathcal{D} and tuple \bar{a} in \mathcal{D} :

$$\bar{a} \in q(\mathcal{D}) \iff (q', \bar{x}') \to (\mathcal{D}, \bar{a}) \iff (chase_{\Sigma}(q), \bar{x}) \to_k (\mathcal{D}, \bar{a}).$$
 (7)

Notice that q' is not guaranteed to exist a priori, since Lemma 2 is stated only for a finite CQ q, while $chase_{\Sigma}(q)$ may be infinite. However, the same proof of Lemma 2 applies to the infinite case. We show next that q' is the $\mathsf{GHW}(k)^{\infty}$ -overapproximation of q under Σ .

First, $(q', \bar{x}') \to (chase_{\Sigma}(q), \bar{x})$ by choosing (\mathcal{D}, \bar{a}) as $(chase_{\Sigma}(q), \bar{x})$ in Equation (7), and hence $q \subseteq_{\Sigma}^{\infty} q'$ as this direction of Lemma 7 holds even for CQs with countable many atoms. Therefore, also $q \subseteq_{\Sigma} q'$. On the other hand, $(chase_{\Sigma}(q), \bar{x}) \to_k (q', \bar{x}')$ by choosing (\mathcal{D}, \bar{a}) as (q', \bar{x}') in Equation (7). Proposition 6 then tells us that for each $q''(\bar{x}'')$ in $\mathsf{GHW}(k)^{\infty}$, if $(q'', \bar{x}'') \to (chase_{\Sigma}(q), \bar{x})$ then $(q'', \bar{x}'') \to (q', \bar{x}')$. But then $q \subseteq_{\Sigma} q''$ implies $q' \subseteq q''$. This is because $q \subseteq_{\Sigma} q''$ implies $(q'', \bar{x}'') \to (chase_{\Sigma}(q), \bar{x})$ by finite controllability of Σ . Indeed, in this case $q \subseteq_{\Sigma} q''$ implies $q \subseteq_{\Sigma} q''$, and thus $q(\mathcal{D}) \subseteq q''(\mathcal{D})$ over every countable database \mathcal{D} that satisfies Σ . Since $chase_{\Sigma}(q)$ is one such a database and $\bar{x} \in q(chase_{\Sigma}(q))$, we have that $\bar{x} \in q''(chase_{\Sigma}(q))$. By definition then, $(q'', \bar{x}'') \to (chase_{\Sigma}(q), \bar{x})$.

In summary, $q \subseteq_{\Sigma} q'$ and for each CQ q'' in $\mathsf{GHW}(k)^{\infty}$ we have that

$$q \subseteq_{\Sigma} q'' \implies q' \subseteq q''.$$

This implies that q' is the $GHW(k)^{\infty}$ -overapproximation of q under Σ . \square

The only property of the sets of guarded tgds that we used in the previous proof is finite controllability. Since finite controllability does not hold for general tgds, we cannot extend Theorem 8 to arbitrary sets of constraints. On the other hand, if we change the notion of $\mathsf{GHW}(k)^{\infty}$ -overapproximation under Σ to be defined in terms of $\subseteq_{\Sigma}^{\infty}$, we can mimic the proof of Theorem 8 and provide each CQ with a $\mathsf{GHW}(k)^{\infty}$ -overapproximation under Σ , where Σ is an arbitrary set of tgds. While considering countable databases is not standard in databases, it is a common choice for the semantics of containment and related problems in the area of ontology-mediated query answering; cf., [10,3].

As a corollary to the proof of Theorem 8 we obtain that the $\mathsf{GHW}(k)^{\infty}$ overapproximation of a CQ under Σ , where Σ is a set of egds or guarded tgds,
can be evaluated by applying the existential k-cover game on $chase_{\Sigma}(q)$.

Corollary 5 Fix $k \geq 1$. Consider a $CQ(q(\bar{x}))$ and a set Σ consisting exclusively of egds or guarded tgds, such that the $\mathsf{GHW}(k)^{\infty}$ -overapproximation of q under Σ is $q'(\bar{x}')$. Then for every database \mathcal{D} and tuple \bar{a} in \mathcal{D} it is the case that

$$\bar{a} \in q'(\mathcal{D}) \iff (q', \bar{x}') \to (\mathcal{D}, \bar{a}) \iff (chase_{\Sigma}(q), \bar{x}) \to_k (\mathcal{D}, \bar{a}).$$

Evaluating overapproximations under constraints. While in the absence of constraints the $\mathsf{GHW}(k)^\infty$ -overapproximation of q can be evaluted in polynomial time by applying the existential k-cover game on q, the situation is more complex in the presence of constraints. In fact, as stated in Corollary 5 to evaluate the $\mathsf{GHW}(k)^\infty$ -overapproximation of q under Σ by using the existential k-cover game, we first need to compute the result of the chase on q. For arbitrary egds this might take exponential time, as checking whether $chase_{\Sigma}(q) = q$, when Σ consists of a single egd ϵ , is an NP-complete problem (since we need to detect whether the CQ that defines the body of ϵ is applicable on q). For sets of guarded tgds the result of the chase might be infinite, and thus it is not even computable.

We show that, in spite of the previous observation, if Σ is a set of guarded tgds then the $\mathsf{GHW}(k)^{\infty}$ -overapproximation of q under Σ can be evaluated in polynomial time. This is because when a database \mathcal{D} satisfies Σ , applying the existential k-cover game on q or $chase_{\Sigma}(q)$ is the same.

Theorem 9 Fix $k \geq 1$. Given a CQ q, a set of Σ of guarded tyds, a database \mathcal{D} satisfying Σ , and a tuple \bar{a} , checking whether $\bar{a} \in q'(\mathcal{D})$, where q' is the $\mathsf{GHW}(k)^{\infty}$ -overapproximation of q under Σ , can be solved in polynomial time by simply verifying if $(q, \bar{x}) \rightarrow_k (\mathcal{D}, \bar{a})$.

Proof As stated in [5], if Σ is a set of guarded tgds and \mathcal{D} satisfies Σ , then $(chase_{\Sigma}(q), \bar{x}) \to_1 (\mathcal{D}, \bar{a}) \iff (q, \bar{x}) \to_1 (\mathcal{D}, \bar{a})$ for every CQ q. A slight modification of this proof shows that this property extends to any k > 1. That is, for every $k \geq 1$ it is the case that

$$(chase_{\Sigma}(q), \bar{x}) \to_k (\mathcal{D}, \bar{a}) \iff (q, \bar{x}) \to_k (\mathcal{D}, \bar{a}).$$

The result then follows from this equivalence and Corollary 5.

For egds, on the other hand, we obtain that the problem can be solved in time $|D|^{O(1)} \cdot f(|q|)$, for a computable function $f: \mathbb{N} \to \mathbb{N}$. In parameterized complexity terms, this means that the problem is *fixed-parameter tractable* (FPT), with the parameter being the size of the CQ. This is a positive result, as no FPT for general CQ evaluation is believed to exist [39].

Theorem 10 Fix $k \geq 1$. Given a CQ q, a set of Σ of egds, a database \mathcal{D} satisfying Σ and a tuple \bar{a} , checking whether $\bar{a} \in q'(\mathcal{D})$, where q' is the $\mathsf{GHW}(k)^{\infty}$ -overapproximation of q under Σ , can be solved by an FPT algorithm.

Proof The algorithm first computes $chase_{\Sigma}(q)$ in exponential time in the size of q, and then checks whether $(chase_{\Sigma}(q), \bar{x}) \to_k (\mathcal{D}, \bar{a})$ in polynomial time in the combined size of $chase_{\Sigma}(q)$ and \mathcal{D} . But the size of $chase_{\Sigma}(q)$ is bounded by that of q, and thus the whole procedure can be carried out in time

$$2^{p(|q|)} + (|D| + |q|)^{O(1)},$$

for $p: \mathbb{N} \to \mathbb{N}$ a polynomial. Hence, the algorithm is FPT.

Notice than in case that Σ consists exclusively of FDs, then the problem can be solved in polynomial time. This is because in such a case $chase_{\Sigma}(q)$ can be computed in polynomial time.

Corollary 6 Fix $k \geq 1$. Given a CQ q, a set of Σ of FDs, a database \mathcal{D} satisfying Σ , and a tuple \bar{a} , checking whether $\bar{a} \in q'(\mathcal{D})$, where q' is the $\mathsf{GHW}(k)^{\infty}$ -overapproximation of q under Σ , can be solved in polynomial time.

7 Beyond Under- and Overapproximations: Δ -Approximations

We now turn to Δ -approximations. Recall that a $\mathsf{GHW}(k)$ - Δ -approximation of q is a maximal element in $\mathsf{GHW}(k)$ with respect to the partial order \sqsubseteq_q , where $q' \sqsubseteq_q q''$, for $\mathsf{CQs}\ q', q'' \in \mathsf{GHW}(k)$, iff $\Delta(q(\mathcal{D}), q''(\mathcal{D})) \subseteq \Delta(q(\mathcal{D}), q'(\mathcal{D}))$ for all databases \mathcal{D} . It is worth noticing that $\mathsf{GHW}(k)$ - Δ -approximations generalize over- and underapproximations.

Proposition 9 Fix $k \ge 1$. Let q, q' be CQs such that $q' \in \mathsf{GHW}(k)$. If $q \subseteq q'$ (resp., $q' \subseteq q$), then q' is a $\mathsf{GHW}(k)$ - Δ -approximation of q if and only if q' is an $\mathsf{GHW}(k)$ -overapproximation (resp., $\mathsf{GHW}(k)$ -underapproximation) of q.

Proof We only prove it for the case when $q \subseteq q'$. The proof for the case when $q' \subseteq q$ is analogous.

Suppose first that q' is a $\mathsf{GHW}(k)$ - Δ -approximation of q. Assume, towards a contradiction, that there is a query q'' such that $q \subseteq q'' \subset q'$. Then (i) $q(\mathcal{D}) \subseteq q''(\mathcal{D}) \subseteq q'(\mathcal{D})$ for each database \mathcal{D} , and (2) there is a database \mathcal{D}^* such that $q'(\mathcal{D}^*) \not\subseteq q''(\mathcal{D}^*)$. In particular, $\Delta(q(\mathcal{D}), q''(\mathcal{D})) \subseteq \Delta(q(\mathcal{D}), q'(\mathcal{D}))$ for each database \mathcal{D} , while $\Delta(q(\mathcal{D}^*), q'(\mathcal{D}^*)) \not\subseteq \Delta(q(\mathcal{D}^*), q''(\mathcal{D}^*))$. This is a contradiction to q' being a $\mathsf{GHW}(k)$ - Δ -approximation of q.

Now suppose that q' is a $\mathsf{GHW}(k)$ -overapproximation of q. Assume, towards a contradiction, that there is a query q'' such that (i) $\Delta(q(\mathcal{D}), q''(\mathcal{D})) \subseteq \Delta(q(\mathcal{D}), q'(\mathcal{D}))$ for each database \mathcal{D} , while (ii) for some database \mathcal{D}^* it is the case that $\Delta(q(\mathcal{D}^*), q'(\mathcal{D}^*)) \not\subseteq \Delta(q(\mathcal{D}^*), q''(\mathcal{D}^*))$. Then $q \subseteq q'' \subseteq q'$ as $\Delta(q(\mathcal{D}), q'(\mathcal{D}))$ may only contain tuples in $q'(\mathcal{D}) \setminus q(\mathcal{D})$. Also, $q'(\mathcal{D}^*) \not\subseteq q''(\mathcal{D}^*)$. This is a contradiction to q' being a $\mathsf{GHW}(k)$ -overapproximation of q.

For this reason, we concentrate on the study of $\mathsf{GHW}(k)$ - Δ -approximations that are neither $\mathsf{GHW}(k)$ -under- nor $\mathsf{GHW}(k)$ -overapproximations. Evaluating such Δ -approximations can give us useful information that complements the one provided by under- and overapproximations. But, do these $\mathsf{GHW}(k)$ - Δ -approximations exist at all, i.e., are there $\mathsf{GHW}(k)$ - Δ -approximations that are neither $\mathsf{GHW}(k)$ -under- nor $\mathsf{GHW}(k)$ -overapproximations? In the rest of this section, we settle this question and study complexity questions associated with such $\mathsf{GHW}(k)$ - Δ -approximations.

7.1 Incomparable GHW(k)- Δ -approximations

Let q be a CQ. In view of Proposition 9, the $\mathsf{GHW}(k)$ - Δ -approximations q' of q that are neither $\mathsf{GHW}(k)$ -under nor $\mathsf{GHW}(k)$ -overapproximations must be incomparable with q in terms of containment; i.e., both $q \not\subseteq q'$ and $q' \not\subseteq q$ must hold. Incomparable $\mathsf{GHW}(k)$ - Δ -approximations do not necessarily exist, even when approximating in the set of infinite CQs $\mathsf{GHW}(k)^{\infty}$. A trivial example is any CQ q in $\mathsf{GHW}(k)$, as its only $\mathsf{GHW}(k)$ - Δ -approximation (up to equivalence) is q itself. On the other hand, the following characterization will help us to find CQs that do have incomparable $\mathsf{GHW}(k)$ - Δ -approximations.

Theorem 11 Fix $k \geq 1$. Let $q(\bar{x}), q'(\bar{x}')$ be CQs such that $q' \in \mathsf{GHW}(k)$. Then q' is an incomparable $\mathsf{GHW}(k)$ - Δ -approximation of q iff $(q, \bar{x}) \to_k (q', \bar{x}')$, and both $q \not\subseteq q'$ and $q' \not\subseteq q$ hold.

Proof Suppose that q' is an incomparable $\mathsf{GHW}(k)$ - Δ -approximation of q and assume, by contradiction, that $(q, \bar{x}) \not\to_k (q', \bar{x}')$. By Proposition 3, there is a $q''(\bar{x}'') \in \mathsf{GHW}(k)$ such that $q \subseteq q''$ and $q' \not\subseteq q''$. Recall that $(q'' \land q')(\bar{z})$ denotes the disjoint conjunction of q'' and q' (see Section 4.1 for the precise definition). We show that $q' \sqsubseteq_q (q'' \land q')$, which is a contradiction as $(q'' \land q') \in \mathsf{GHW}(k)$. Assume that $\bar{a} \in \Delta(q(\mathcal{D}), (q'' \land q')(\mathcal{D}))$, for some \mathcal{D} and $\bar{a} \in \mathcal{D}$. If $\bar{a} \not\in q(\mathcal{D})$, then $\bar{a} \in (q'' \land q')(\mathcal{D}) \subseteq q'(\mathcal{D})$, and thus, $\bar{a} \in \Delta(q(\mathcal{D}), q'(\mathcal{D}))$. Otherwise, $\bar{a} \in q(\mathcal{D})$ and $\bar{a} \not\in (q'' \land q')(\mathcal{D})$. Since $q \subseteq q''$, we have $\bar{a} \not\in q'(\mathcal{D})$, and then $\bar{a} \in \Delta(q(\mathcal{D}), q'(\mathcal{D}))$. Hence $q' \sqsubseteq_q (q'' \land q')$. Now, since $q' \not\subseteq q''$, there is a database \mathcal{D}^* such that $q'(\mathcal{D}^*) \not\subseteq q''(\mathcal{D}^*)$, i.e., $\bar{a} \in q'(\mathcal{D}^*)$ but $\bar{a} \not\in q''(\mathcal{D}^*)$, for some tuple \bar{a} in \mathcal{D}^* . In particular $\bar{a} \in \Delta(q(\mathcal{D}^*), q'(\mathcal{D}^*))$ and $\bar{a} \not\in \Delta(q(\mathcal{D}^*), (q'' \land q')(\mathcal{D}^*))$, and thus $(q'' \land q') \not\sqsubseteq_q q'$.

For the converse, we need the following lemma whose proof can be found in the appendix.

Lemma 8 Fix $k \geq 1$. Let $q(\bar{x}), q'(\bar{x}'), q''(\bar{x}'')$ be CQs such that $q'' \in \mathsf{GHW}(k)$. Let $(q' \land q)(\bar{z})$ be the disjoint conjunction of q' and q. Suppose that $(q, \bar{x}) \rightarrow_k (q', \bar{x}')$. Then $(q'', \bar{x}'') \rightarrow (q' \land q, \bar{z})$ implies $(q'', \bar{x}'') \rightarrow (q', \bar{x}')$.

Assume now that $q \nsubseteq q', q' \nsubseteq q$, and $(q, \bar{x}) \to_k (q', \bar{x}')$. By contradiction, suppose that there is a CQ $q'' \in \mathsf{GHW}(k)$ such that $q' \sqsubseteq_q q''$. We show that $q' \equiv q''$, which is a contradiction. Recall that $\mathcal{D}_{(q' \land q)}$ denotes the canonical database of $(q' \land q)(\bar{z})$. Clearly, $\bar{z} \in q(\mathcal{D}_{(q' \land q)})$ and $\bar{z} \in q'(\mathcal{D}_{(q' \land q)})$. It follows that $\bar{z} \not\in \Delta(q(\mathcal{D}_{(q' \land q)}), q'(\mathcal{D}_{(q' \land q)}))$, and by hypothesis, $\bar{z} \not\in \Delta(q(\mathcal{D}_{(q' \land q)}), q''(\mathcal{D}_{(q' \land q)}))$. Hence, $\bar{z} \in q''(\mathcal{D}_{(q' \land q)})$. By Lemma 8, we have $(q'', \bar{x}'') \to (q', \bar{x}')$, that is, $q' \subseteq q''$. For $q'' \subseteq q'$, note that $\bar{x}'' \not\in q(\mathcal{D}_{q''})$; otherwise, $q'' \subseteq q$ would hold, implying that $q' \subseteq q$, which is a contradiction. Since $\bar{x}'' \in q''(\mathcal{D}_{q''})$, we have $\bar{x}'' \in \Delta(q(\mathcal{D}_{q''}), q''(\mathcal{D}_{q''}))$. This implies that $\bar{x}'' \in \Delta(q(\mathcal{D}_{q''}), q'(\mathcal{D}_{q''}))$, and then $\bar{x}'' \in q'(\mathcal{D}_{q''})$, i.e., $q'' \subseteq q'$. Hence, $q' \equiv q''$.

Example~4 Consider again the CQ

$$q \ = \ \exists x \exists y \exists z \big(E(x,y) \land E(y,z) \land E(z,x) \big)$$

from Figure 3. It is easy to prove that $q' = \exists x E(x,x)$ is the unique $\mathsf{GHW}(1)$ -underapproximation of q. Also, as mentioned in Section 4.1, q has no $\mathsf{GHW}(1)$ -overapproximations. Is it the case, on the other hand, that q has incomparable $\mathsf{GHW}(1)$ - Δ -approximations? By applying Theorem 11, we can give a positive answer to this question. In fact, the CQ

$$q'' = \exists x \exists y (E(x,y) \land E(y,x))$$

is an incomparable GHW(1)- Δ -approximation of q.

Therefore, as Example 4 shows, incomparable $\mathsf{GHW}(k)$ - Δ -approximations may exist for some CQs. However, in contrast with overapproximations, they are not unique in general (see the appendix for details).

Proposition 10 There is a CQ with infinitely many (non-equivalent) incomparable GHW(1)- Δ -approximations. In fact, this even holds for the CQ q shown in Figure 2.

Identification, existence and evaluation. A direct consequence of Theorem 11 is that the identification problem, i.e., checking if $q' \in \mathsf{GHW}(k)$ is an incomparable $\mathsf{GHW}(k)$ - Δ -approximation of a CQ q, is in coNP. It suffices to check that $q \not\subseteq q'$ and $q' \not\subseteq q$ — which are in coNP — and $(q, \bar{x}) \to_k (q', \bar{x}')$ — which is in PTIME from Proposition 2. We show next that this bound is optimal (the proof is in the appendix).

Proposition 11 Fix $k \geq 1$. Checking if a given CQ $q' \in GHW(k)$ is an incomparable GHW(k)- Δ -approximation of a given CQ q, is CONP-complete.

As in the case of $\mathsf{GHW}(k)$ -overapproximations, we do not know how to check existence of incomparable $\mathsf{GHW}(k)$ - Δ -approximations, for k>1. Nevertheless, for k=1 we can exploit the automata techniques developed in Section 5.1 and obtain an analogous decidability result.

Proposition 12 There is a 2EXPTIME algorithm that checks if a CQ q has an incomparable GHW(1)- Δ -approximation and, if one exists, it computes one in triple exponential time. The bounds become EXPTIME and 2EXPTIME, respectively, if the maximum arity of the schema is fixed.

Now we study evaluation. Recall that, unlike $\mathsf{GHW}(k)$ -overapproximations, incomparable $\mathsf{GHW}(k)$ - Δ -approximations of a $\mathsf{CQ}\ q$ are not unique. In fact, there can be infinitely many (see Proposition 10). Thus, it is reasonable to start by trying to evaluate at least one of them. It would be desirable, in addition, if the one we evaluate depends only on q (i.e., it is independent of the underlying database \mathcal{D}). Proposition 12 allows us to do so as follows. Given a $\mathsf{CQ}\ q$ with at least one incomparable $\mathsf{GHW}(1)$ - Δ -approximation, we can compute in $3\mathsf{EXPTIME}$ one such an incomparable $\mathsf{GHW}(1)$ - Δ -approximation q'. We can then evaluate q' over a database \mathcal{D} in time $O(|\mathcal{D}| \cdot |q'|)$ [40], which is $O(|\mathcal{D}| \cdot f(|q|))$, for f a triple-exponential function. This means that the




Fig. 5 The CQ $q \in \mathsf{GHW}(2)$ from Example 5. The CQ $(q^* \land q')$ is an incomparable $\mathsf{GHW}(1)^{\infty}$ - Δ -approximation of q. On the other hand, q has no incomparable $\mathsf{GHW}(1)$ - Δ -approximations.

evaluation of such a q' over \mathcal{D} is fixed-parameter tractable, i.e., it can be solved by an algorithm that depends polynomially on the size of the large database \mathcal{D} , but more loosely on the size of the small CQ q. (Recall that this is a desirable property for evaluation, which does not hold in general for the class of all CQs [39]).

Theorem 12 There is a fixed-parameter tractable algorithm that, given a CQ q that has incomparable $\mathsf{GHW}(1)$ - Δ -approximations, a database \mathcal{D} , and a tuple \bar{a} , checks whether $\bar{a} \in q'(\mathcal{D})$, for some incomparable $\mathsf{GHW}(1)$ - Δ -approximation q' of q that depends only on q.

It is worth noticing that the automata techniques are essential for proving this result, and thus for evaluating incomparable $\mathsf{GHW}(1)$ - Δ -approximations. This is in stark contrast with $\mathsf{GHW}(k)$ -overapproximations, which can be evaluated in polynomial time by simply checking if $(q,\bar{x}) \to_k (\mathcal{D},\bar{a})$. It is not at all clear whether such techniques can be extended to allow for the efficient evaluation of incomparable $\mathsf{GHW}(k)$ - Δ -approximations.

The infinite case. All the previous results continue to apply for the class of infinite CQs in $\mathsf{GHW}(k)^\infty$. The following example shows that, as in the case of $\mathsf{GHW}(k)$ -overapproximations, considering $\mathsf{GHW}(k)^\infty$ helps us to obtain better incomparable $\mathsf{GHW}(k)$ - Δ -approximations.

Example 5 Consider the CQ q that asks for the existence of the two oriented paths P_1 and P_2 , as shown in Figure 5. Theorem 11 can be used to show that q has no incomparable $\mathsf{GHW}(1)$ - Δ -approximation. However, q has an incomparable $\mathsf{GHW}(1)^{\infty}$ - Δ -approximation. In fact, let q^* be the CQ which is depicted in Figure 5 (a P_1 -labeled edge represents a copy of the oriented path P_1 , similarly for P_2). It is easy to see that q^* is the $\mathsf{GHW}(1)^{\infty}$ -overapproximation of q. Also, let q' be an arbitrary CQ in $\mathsf{GHW}(1)$ which is incomparable with q (one such q' is shown in Figure 5). Applying the extension of Theorem 11 to the class $\mathsf{GHW}(k)^{\infty}$, we can prove that $(q^* \wedge q')$ is an incomparable $\mathsf{GHW}(1)^{\infty}$ - Δ -approximation of q.

Example 5 also illustrates the following fact: If there is a CQ $q' \in \mathsf{GHW}(k)$ which is incomparable with q, then $(q^* \wedge q')$ is an incomparable $\mathsf{GHW}(k)^{\infty}$ - Δ -approximation of q, where q^* is the $\mathsf{GHW}(k)^{\infty}$ -overapproximation of q. Given a database \mathcal{D} and a tuple \bar{a} in \mathcal{D} , we can check whether \bar{a} belongs to the

evaluation of such a Δ -approximation $(q^* \wedge q')$ over \mathcal{D} as follows. First we compute q', and then we check both $(q, \bar{x}) \to_k (\mathcal{D}, \bar{a})$ and $\bar{a} \in q'(\mathcal{D})$. In other words, we evaluate $(q^* \wedge q')$ via the existential k-cover game – as for the $\mathsf{GHW}(k)^\infty$ -overapproximation –, and then use the incomparable $\mathsf{CQ}\ q'$ to filter out some tuples in the answer. Interestingly, we can easily exploit automata techniques and compute such an incomparable q' (in case one exists). Thus we have the following:

Theorem 13 Fix $k \geq 1$. There is a fixed-parameter tractable algorithm that given a CQ q that has an incomparable q' in GHW(k), a database \mathcal{D} , and a tuple \bar{a} in \mathcal{D} , decides whether $\bar{a} \in \hat{q}(\mathcal{D})$, for some incomparable $GHW(k)^{\infty}$ - Δ -approximation \hat{q} of q that depends only on q.

8 Final Remarks

Several problems remain open: is the existence of $\mathsf{GHW}(k)$ -overapproximations decidable for k>1? What is the precise complexity of checking for the existence of $\mathsf{GHW}(1)$ -overapproximations? In particular, can we improve the $2\mathsf{EX-PTIME}$ upper bound from Theorem 5? What is an optimal upper bound on the size of $\mathsf{GHW}(1)$ -overapproximations? Can we extend to non-Boolean CQs the result that states the tractability of checking for the existence of $\mathsf{GHW}(1)$ -overapproximations over binary schemas?

In the future we plan to study how our notions of approximation can be combined with other techniques to obtain quantitative guarantees. One possibility is to exploit semantic information about the data – e.g., in the form of integrity constraints – in order to ensure that certain bounds on the size of the result of the approximation hold. Another possibility is to try to obtain probabilistic guarantees for approximations based on reasonable assumptions about the distribution of the data.

References

- 1. Abiteboul, S., Hull, R., Vianu, V.: Foundations of Databases. Addison-Wesley (1995)
- Baader, F., Calvanese, D., McGuinness, D.L., Nardi, D., Patel-Schneider, P.F. (eds.): The Description Logic Handbook: Theory, Implementation, and Applications. Cambridge University Press (2003)
- Bárány, V., Gottlob, G., Otto, M.: Querying the guarded fragment. Logical Methods in Computer Science 10(2) (2014)
- 4. Barceló, P.: Querying graph databases. In: PODS, pp. 175–188 (2013)
- 5. Barceló, P., Gottlob, G., Pieris, A.: Semantic acyclicity under constraints. In: PODS, pp. 343–354 (2016)
- Barceló, P., Libkin, L., Romero, M.: Efficient approximations of conjunctive queries. In: PODS, pp. 249–260 (2012)
- Barceló, P., Libkin, L., Romero, M.: Efficient approximations of conjunctive queries. SIAM J. Comput. 43(3), 1085–1130 (2014)
- Barceló, P., Romero, M., Vardi, M.Y.: Semantic acyclicity on graph databases. SIAM J. Comput. 45(4), 1339–1376 (2016)

- Blumensath, A., Otto, M., Weyer, M.: Decidability results for the boundedness problem. Logical Methods in Computer Science 10(3) (2014)
- Calì, A., Gottlob, G., Kifer, M.: Taming the infinite chase: Query answering under expressive relational constraints. In: KR, pp. 70–80 (2008)
- 11. Chandra, A.K., Merlin, P.M.: Optimal implementation of conjunctive queries in relational data bases. In: STOC, pp. 77–90 (1977)
- Chekuri, C., Rajaraman, A.: Conjunctive query containment revisited. Theor. Comput. Sci. 239(2), 211–229 (2000)
- Chen, H., Dalmau, V.: Beyond hypertree width: Decomposition methods without decompositions. In: CP, pp. 167–181 (2005)
- Cosmadakis, S.S., Gaifman, H., Kanellakis, P.C., Vardi, M.Y.: Decidable optimization problems for database logic programs (preliminary report). In: STOC, pp. 477–490 (1988)
- Dalmau, V., Kolaitis, P.G., Vardi, M.Y.: Constraint satisfaction, bounded treewidth, and finite-variable logics. In: CP, pp. 310–326 (2002)
- Deutsch, A., Nash, A., Remmel, J.B.: The chase revisisted. In: PODS, pp. 149–158 (2008)
- Fagin, R.: A normal form for relational databases that is based on domains and keys.
 ACM Trans. Database Syst. 6(3), 387–415 (1981)
- Fagin, R., Kolaitis, P.G., Miller, R.J., Popa, L.: Data exchange: Semantics and query answering. Theor. Comput. Sci. 336(1), 89–124 (2005)
- Fan, W., Li, J., Ma, S., Tang, N., Wu, Y., Wu, Y.: Graph pattern matching: From intractable to polynomial time. PVLDB 3(1), 264–275 (2010)
- 20. Fink, R., Olteanu, D.: On the optimal approximation of queries using tractable propositional languages. In: ICDT, pp. 174–185 (2011)
- 21. Fischl, W., Gottlob, G., Pichler, R.: General and fractional hypertree decompositions: Hard and easy cases. In: PODS, pp. 17–32 (2018)
- Gaifman, H., Mairson, H.G., Sagiv, Y., Vardi, M.Y.: Undecidable optimization problems for database logic programs. J. ACM 40(3), 683–713 (1993)
- 23. Garofalakis, M., Gibbon, P.: Approximate query processing: taming the terabytes. In: VLDB, p. 725 (2001)
- 24. Gottlob, G., Greco, G., Leone, N., Scarcello, F.: Hypertree decompositions: Questions and answers. In: PODS, pp. 57–74 (2016)
- Gottlob, G., Leone, N., Scarcello, F.: Hypertree decompositions and tractable queries.
 J. Comput. Syst. Sci. 64(3), 579–627 (2002)
- Gottlob, G., Miklós, Z., Schwentick, T.: Generalized hypertree decompositions: NP-hardness and tractable variants. J. ACM 56(6) (2009)
- 27. Greco, G., Scarcello, F.: The power of local consistency in conjunctive queries and constraint satisfaction problems. SIAM J. Comput. 46(3), 1111–1145 (2017)
- Grohe, M., Marx, D.: Constraint solving via fractional edge covers. In: SODA, pp. 289–298 (2006)
- Hell, P., Nesetril, J.: The core of a graph. Discrete Mathematics 109(1-3), 117–126 (1992)
- Hell, P., Nesetril, J., Zhu, X.: Complexity of tree homomorphisms. Discrete Applied Mathematics 70(1), 23–36 (1996)
- 31. Hell, P., Nešetril, J.: Graphs and homomorphisms. Oxford University Press (2004)
- 32. Ioannidis, Y.: Approximations in database systems. In: ICDT, pp. 16-30 (2003)
- Kolaitis, P.G., Panttaja, J.: On the complexity of existential pebble games. In: CSL, pp. 314–329 (2003)
- Kolaitis, P.G., Vardi, M.Y.: On the expressive power of datalog: Tools and a case study.
 J. Comput. Syst. Sci. 51(1), 110–134 (1995)
- Kolaitis, P.G., Vardi, M.Y.: Conjunctive-query containment and constraint satisfaction.
 J. Comput. Syst. Sci. 61(2), 302–332 (2000)
- 36. Liu, Q.: Approximate query processing. In: Encyclopedia of Database Systems, pp. 113–119 (2009)
- Maier, D., Mendelzon, A.O., Sagiv, Y.: Testing implications of data dependencies. ACM Trans. Database Syst. 4(4), 455–469 (1979)
- 38. Otto, M.: The boundedness problem for monadic universal first-order logic. In: LICS, pp. 37–48 (2006)




Fig. 6 Directed graphs that satisfy condition (†) for k = 2, 3, 4, respectively.

- Papadimitriou, C.H., Yannakakis, M.: On the complexity of database queries. J. Comput. Syst. Sci. 58(3), 407–427 (1999)
- 40. Yannakakis, M.: Algorithms for acyclic database schemes. In: VLDB, pp. 82-94 (1981)

Appendix

Proof (Theorem 1) Fix k > 1. The CQ q is defined over graphs, i.e., over a schema with a single binary relation symbol E, and consists of k+1 variables v_1, \ldots, v_{k+1} . For every $1 \le i < j \le k+1$ we add either the atom (i.e., edge) $E(v_i, v_j)$ or $E(v_j, v_i)$ to q in such a way that the subgraph of G induced by $\{v_1, v_2, v_3\}$ is a directed cycle and a certain condition (\dagger), defined below, holds. We start introducing some terminology.

Let G be a directed graph on nodes v_1,\ldots,v_{k+1} that contains, for each $1\leq i< j\leq k+1$, either the edge $E(v_i,v_j)$ or $E(v_j,v_i)$. For a $B\subseteq \{v_1,\ldots,v_{k+1}\}$ of size $1\leq \ell\leq k-1$ and a node $v\in \{v_1,\ldots,v_{k+1}\}\setminus B$, we define $\mathsf{conn}(v,B)$ as the tuple $(e_1,\ldots,e_{k+1})\in \{-1,1,\#\}^{k+1}$ such that for each $1\leq p\leq k+1$:

$$e_p \ = \begin{cases} \#, & \text{if } v_p \notin B, \\ 1, & \text{if } v_p \in B \text{ and the edge } E(v, v_p) \text{ is in } G, \\ -1, & \text{otherwise, i.e., } v_p \in B \text{ and } E(v_p, v) \text{ is in } G. \end{cases}$$

In simple terms, conn(v, B) specifies how v connects with the nodes in B. Our condition (†) then establishes the following:

(†) For each $B\subseteq\{v_1,\ldots,v_{k+1}\}$ of size $2\leq\ell\leq k-1$ and each node v in $\{v_1,\ldots,v_{k+1}\}\setminus B$, there is a node $v'\in\{v_1,\ldots,v_{k+1}\}\setminus B$ such that

$$conn(v, B) \neq conn(v', B).$$

That is, for each such B and v we will always be able to find another v' outside B that connects to the nodes in B in a different way than v.

Example 6 The graphs in Figure 6 satisfy this condition for k = 2, 3, 4, respectively. Notice that the directed cycle on nodes $\{v_1, v_2, v_3\}$, shown in the left-hand side, satisfies condition (†) trivially.

The next lemma establishes that for each k>1 there is always a graph that satisfies this condition.

Lemma 9 For each k > 1, there is a directed graph G on nodes v_1, \ldots, v_{k+1} such that the following hold:

- 1. For each $1 \le i < j \le k+1$, either the edge $E(v_i, v_j)$ or $E(v_j, v_i)$ is in G;
- 2. the subgraph of G induced by $\{v_1, v_2, v_3\}$ is a directed cycle; and
- 3. G satisfies condition (\dagger) .

Proof (Lemma 9) For k=2 this is given by the graph in Example 6. For $k \geq 3$ we prove by induction a stronger claim: There is a directed graph G on nodes v_1, \ldots, v_{k+1} such that:

- 1. G contains either the edge $E(v_i, v_j)$ or $E(v_j, v_i)$ for each $1 \le i < j \le k+1$.
- 2. The subgraph of G induced by $\{v_1, v_2, v_3\}$ is a directed cycle.
- 3. G contains the edges $E(v_1, v_2)$ and $E(v_4, v_3)$.
- 4. G satisfies condition (†).

The basis case k=3 is given again by the graph in Example 6. For the inductive case, assume by induction hypothesis that there is a directed graph G on nodes v_1, \ldots, v_{k+1} that satisfies the claim above. A new graph G' is then created from G by adding a new node v_{k+2} and connecting it to the nodes in $\{v_1, \ldots, v_{k+1}\}$ as follows: For each $1 \le i \le k$, if $E(v_i, v_{i+1})$ is in G then we add the edge $E(v_{k+2}, v_i)$ to G', otherwise we add the edge $E(v_i, v_{k+2})$. Moreover, if $E(v_{k+1}, v_1)$ is in G then we add the edge $E(v_{k+2}, v_{k+1})$ to G', otherwise we add the edge $E(v_{k+1}, v_{k+2})$. Notice that G coincides with the subgraph of G' that is induced by nodes v_1, \ldots, v_{k+1} . Moreover, by construction G' satisfies the first three conditions of the claim. We prove next that it also satisfies condition (\dagger) .

Take an arbitrary $B \subseteq \{v_1, \ldots, v_{k+2}\}$ of size $2 \le \ell \le k$ and a node v outside B. We prove that the condition holds by cases:

- $-v_{k+2} \notin B$, $v \in \{v_1, \dots, v_{k+1}\}$, and $2 \le \ell \le k-1$: By inductive hypothesis there is a node $v' \in \{v_1, \dots, v_{k+1}\} \setminus B$ such that $\mathsf{conn}(v, B) \ne \mathsf{conn}(v', B)$.
- $-v_{k+2} \notin B$, $v \in \{v_1, \ldots, v_{k+1}\}$, and $\ell = k$: We set $v' := v_{k+2}$ and claim that the predecessor u of v in $\{v_1, \ldots, v_{k+1}\}$ distinguishes v and v'. Here, the "predecessor" of v_i is v_{i-1} if $1 \le i \le k+1$, and the predecessor of i is i is i in i
- $-v_{k+2} \notin B$ and $v = v_{k+2}$: There must exist some node v' in $\{v_1, \ldots, v_{k+1}\}$ that does not belong to B but its predecessor u in $\{v_1, \ldots, v_{k+1}\}$ does. Then by construction of G', we have that $E(u, v') \in G'$ if and only if $E(v, u) \in G'$. We conclude that $\mathsf{conn}(v, B) \neq \mathsf{conn}(v', B)$.
- $-v_{k+2} \in B$ and $\ell \geq 3$: Then $B' = B \setminus \{v_{k+2}\}$ is of size $2 \leq \ell 1 \leq k 1$. By induction hypothesis, for every node v outside B' there is another node $v' \in \{v_1, \ldots, v_{k+1}\} \setminus B'$ such that $\mathsf{conn}(v, B') \neq \mathsf{conn}(v', B')$. This implies that $\mathsf{conn}(v, B) \neq \mathsf{conn}(v', B)$.
- $-v_{k+2} \in B$ and $\ell = 2$: Then $B = \{v_{k+2}, u\}$ for some $u \in \{v_1, \dots, v_{k+1}\}$. Suppose first that $u \in \{v_1, v_2, v_3\}$. Since the subgraph induced by $\{v_1, v_2, v_3\}$

in G defines a directed cycle, it is the case that E(u,z) holds if and only if E(w,u) holds, where $\{u,w,z\} = \{v_1,v_2,v_3\}$. Therefore, for each $v \in \{v_1,\ldots,v_{k+1}\} \setminus B$ there is a node $v' \in \{z,w\}$ such that $\mathsf{conn}(v,\{u\}) \neq \mathsf{conn}(v',\{u\})$. It follows that $\mathsf{conn}(v,B) \neq \mathsf{conn}(v',B)$. Suppose now that $u \notin \{v_1,v_2,v_3\}$. It suffices to exhibit two nodes v' and v'' outside B such that $E(v',v_{k+2})$ and $E(v_{k+2},v'')$. By induction hypothesis the edges $E(v_1,v_2)$ and $E(v_4,v_3)$ are in G'. Therefore, v_{k+2} is connected via edges $E(v_3,v_{k+2})$ and $E(v_{k+2},v_1)$ in G'.

This concludes the proof of the lemma.

Fix $k \geq 1$. We then take as q any Boolean CQ whose canonical database is a graph G on nodes v_1, \ldots, v_{2k+1} that satisfies the conditions stated in Lemma 9. That is, (1) for each $1 \leq i < j \leq 2k+1$, either the edge $E(v_i, v_j)$ or $E(v_j, v_i)$ is in G, (2) the subgraph of G induced by $\{v_1, v_2, v_3\}$ is a directed cycle, and (3) G satisfies condition (\dagger). It is easy to see that q is in $\mathsf{GHW}(k+1) \setminus \mathsf{GHW}(k)$ as its underlying undirected graph is a clique on 2k+1 elements. In fact, these elements can be covered with (k+1) edges, but not with k.

We claim that q has no $\mathsf{GHW}(\ell)$ -overapproximation for any $1 \le \ell \le k$. The proofs for the cases when $\ell = 1$ and $\ell > 1$ are slightly different. We start with the latter, i.e., when $1 < \ell \le k$. The proof for every such an ℓ is analogous, and thus we concentrate on proving the claim for $\ell = k > 1$. According to Theorem 7, we need to prove that there is no constant $c \ge 0$ such that for every database $\mathcal D$ it holds that

$$q \to_k \mathcal{D} \iff q \to_k^c \mathcal{D}.$$

It is sufficient to show then that for each integer $c \geq 0$ there is a database \mathcal{D} such that

$$q \to_k^c \mathcal{D}$$
 but $q \not\to_k^{c+1} \mathcal{D}$.

Or, equivalently, that for each integer $c \geq 0$ there is a database \mathcal{D} such that

$$q_c \to \mathcal{D}$$
 but $q_{c+1} \not\to \mathcal{D}$,

where q_c , for $c \geq 0$, is the CQ which is defined in Lemma 1, i.e., for every \mathcal{D} it is the case that $q \to_k^c \mathcal{D}$ iff $q_c \to \mathcal{D}$. In view of Equation (1), this boils down to proving that

$$q_{c+1} \not\to q_c$$
, for each $c \ge 0$. (8)

We prove Equation (8) by induction. The claim clearly holds for c=0, as by definition q_0 is empty while q_1 is not. Let us assume now that the claim holds for $c \geq 0$. That is, $q_{c+1} \not\to q_c$. This means, in particular, that the core of q_{c+1} is not contained in q_c . That is, this core contains at least one node w in q_{c+1} that does not belong to q_c .

By the way q is defined, any k-union of q must be of the form $S \subseteq \{v_1, \ldots, v_{2k+1}\}$ with |S| = 2k. Let us consider now (T_{c+1}, β_{c+1}) as defined in the proof of Lemma 1. Since $w \notin q_c$, it must be the case that there is a unique node t of T_{c+1} such that $w \in \beta_{c+1}(t)$. Moreover, this t must be a leaf

of T_{c+1} . Suppose that $\phi_t(w) = v$, for $v \in \{v_1, \ldots, v_{2k+1}\}$, where ϕ_t is as defined in the proof of Lemma 1, i.e., ϕ_t is a bijection between $\beta_{c+1}(t)$ and the k-union $S \subseteq \{v_1, \ldots, v_{2k+1}\}$ of q such that $\lambda_{c+1}(t) = S$. Notice, by definition, that if the parent of t in T_{c+1} is t', then either $\lambda_{c+1}(t') = \emptyset$ – which holds precisely when t' is the root of T_{c+1} –, or $\lambda_{c+1}(t') = S'$, where S' is the subset of $\{v_1, \ldots, v_{2k+1}\}$ which contains all elements save for v. That is, in the latter case we have that S' is obtained from S by replacing some element v' in $\{v_1, \ldots, v_{2k+1}\}$, with $v' \neq v$, by v itself.

From Proposition 1, we can assume that the homomorphism that maps q_{c+1} to its core is a retraction, i.e., it is the identity on the nodes of this core, in particular, on w. On the other hand, w is linked in q_{c+1} exclusively with the remaining nodes that appear in $\beta_{c+1}(t)$. Moreover, the graph induced by the nodes in $\lambda_{c+1}(t)$ is a clique on 2k elements, and thus all the elements in $\beta_{c+1}(t)$ must belong to the core of q_{c+1} .

Recall that $\phi_t(w) = v$. Take an arbitrary node $v'' \in S$ that is not v. Notice that neither v'' = v' as $v'' \in S$, while $v' \notin S$. By definition, T_{c+2} contains a leaf t'' whose parent is t such that $\lambda_{c+2}(t'') = S''$, where S'' is the subset of $\{v_1, \ldots, v_{2k+1}\}$ which is obtained from S by replacing v'' with the unique node in $\{v_1, \ldots, v_{2k+1}\} \setminus S$, namely v'. Let us assume that $\phi_{t''}(v') = w''$. Notice that w'' appears in no other node in (T_{c+2}, β_{c+2}) .

Assume now, for the sake of contradiction, that $q_{c+2} \to q_{c+1}$. Then the core of q_{c+2} is the same than the core of q_{c+1} . Let C be this core. Henceforth, from Proposition 1 there is a retraction h from q_{c+2} to C. Since all elements in $\beta_{c+2}(t) = \beta_{c+1}(t)$ are in C, the homomorphism h must be the identity on them. But then h maps w' to the unique element in q_{c+1} that is linked to exactly the same nodes than w' in q_{c+2} ; namely, $\phi_t(v'') = w''$.

Suppose that v' and v'' represent the nodes v_i and v_j in $\{v_1, \ldots, v_{2k+1}\}$, respectively. By assumption, $i \neq j$. But this implies then that in the canonical database G of q we have that

$$conn(v_i, B) = conn(v_i, B),$$

where $B = \{v_1, \ldots, v_{2k+1}\} \setminus \{v_i, v_j\}$. This is a contradiction since B is of size 2k-1 > 1 and G satisfies condition (†). This concludes our proof that q has no $\mathsf{GHW}(k)$ -overapproximation (and, analogously, that it has no $\mathsf{GHW}(\ell)$ -overapproximation for any $1 < \ell \le k$).

We prove next that q neither has a $\mathsf{GHW}(1)$ -overapproximation. Let us assume, for the sake of contradiction, that q has a $\mathsf{GHW}(1)$ -overapproximation q'. It is an easy observation that the directed graphs in $\mathsf{GHW}(1)$ are precisely those whose underlying undirected graph is acyclic. Notice also that q' has no directed cycles of length two (i.e., atoms of the form E(u,v) and E(v,u)); otherwise, since $q' \to q$, we would have that q also has such a cycle (which we know it does not). Using the fact that $q' \in \mathsf{GHW}(1)$ and has no directed cycles of length two, it is not difficult to show (see e.g. [31]) that there is a sufficiently large integer $n \geq 1$ such that, if \mathbf{P}_n is the directed path on n vertices, then

$$q' \to \mathbf{P}_n$$
 but $\mathbf{P}_n \not\to q'$.

This implies that if q'' is the Boolean CQ which is naturally defined by \mathbf{P}_n , then $q'' \subseteq q'$. Moreover, $\mathbf{P}_n \to G$. This is due to the fact that G contains a directed cycle on $\{v_1, v_2, v_3\}$. We conclude that

$$q \subseteq q'' \subsetneq q'$$
,

and, therefore, that q' is not a GHW(1)-overapproximation of q. This is a contradiction. We then conclude the proof of Theorem 1.

Proof (Lemma 8) Before proving the lemma, we need some terminology and claims. Let \mathcal{D} be a database and (A_1, \ldots, A_n) be a tuple of pairwise-disjoint subsets of elements of \mathcal{D} , where $n \geq 0$. In addition, let \mathcal{D}' be a database and (a_1, \ldots, a_n) a tuple of elements in \mathcal{D}' . Then we write $(\mathcal{D}, (A_1, \ldots, A_n)) \rightarrow (\mathcal{D}', (a_1, \ldots, a_n))$ iff there is a homomorphism h from \mathcal{D} to \mathcal{D}' such that, for each $i \in \{1, \ldots, n\}$ and $a \in A_i$, it is the case that $h(a) = a_i$.

For such a pair $(\mathcal{D}, (A_1, \ldots, A_n))$, with $n \geq 0$, we define its generalized hypertreewidth in the natural way. The intuition is that we see $(\mathcal{D}, (A_1, \ldots, A_n))$ as a "query", where $A_1 \cup \cdots \cup A_n$ are the "free variables" and the rest of the elements are the "existential variables". Formally, a tree decomposition of $(\mathcal{D}, (A_1, \ldots, A_n))$ is a pair (T, χ) , where T is a tree and χ is a mapping that assigns a subset of the elements in $\mathcal{D} \setminus (A_1 \cup \cdots \cup A_n)$ to each node $t \in T$, such that the following statements hold:

- 1. For each atom $R(\bar{a})$ in \mathcal{D} , it is the case that $\bar{a} \cap (\mathcal{D} \setminus (A_1 \cup \cdots \cup A_n))$ is contained in $\chi(t)$, for some $t \in T$.
- 2. For each element a in $\mathcal{D} \setminus (A_1 \cup \cdots \cup A_n)$, the set of nodes $t \in T$ for which a occurs in $\chi(t)$ is connected.

The width of node t in (T, χ) is the minimal number ℓ for which there are ℓ atoms in \mathcal{D} covering $\chi(t)$, i.e., atoms $R(\bar{a}_1), \ldots, R(\bar{a}_\ell)$ in \mathcal{D} such that $\chi(t) \subseteq \bigcup_{1 \leq i \leq \ell} \bar{a}_i$ The width of (T, χ) is the maximal width of the nodes of T. The generalized hypertreewidth of $(\mathcal{D}, (A_1, \ldots, A_n))$ is the minimum width of its tree decompositions.

By mimicking the proof of the forward implication of Proposition 3, we can show the following:

Lemma 10 Fix $k \geq 1$. Let $q(\bar{x}), q'(\bar{x}')$ be CQs, where $\bar{x} = (x_1, \ldots, x_n)$ and $\bar{x}' = (x'_1, \ldots, x'_n)$, for $n \geq 0$. Suppose that $(q, \bar{x}) \rightarrow_k (q', \bar{x}')$. Then, for each database \mathcal{D} and tuple (A_1, \ldots, A_n) of subsets of \mathcal{D} such that $(\mathcal{D}, (A_1, \ldots, A_n))$ has generalized hypertreewidth at most k, it is the case that

$$(\mathcal{D}, (A_1, \dots, A_n)) \to (q, (x_1, \dots, x_n)) \Longrightarrow (\mathcal{D}, (A_1, \dots, A_n)) \to (q', (x'_1, \dots, x'_n)).$$

Proof Let \mathcal{H} be a winning strategy for Duplicator witnessing the fact that $(q, \bar{x}) \to_k (q', \bar{x}')$. Let us assume that $(\mathcal{D}, (A_1, \ldots, A_n))$ has generalized hypertreewidth at most k, and that $(\mathcal{D}, (A_1, \ldots, A_n)) \to (q, (x_1, \ldots, x_n))$ is witnessed via a homomorphism h. Then we can compose h with the strategy \mathcal{H}

to define a homomorphism g witnessing $(\mathcal{D}, (A_1, \ldots, A_n)) \to (q', (x'_1, \ldots, x'_n))$. The mapping g is defined in a top-down fashion over the tree decomposition (T,χ) of width at most k of $(\mathcal{D}, (A_1, \ldots, A_n))$. One starts at the root r of T, and forces Spoiler to play his pebbles over the set $h(\chi(r))$. If Duplicator responds according to \mathcal{H} with a partial homomorphism f_r , we then let $g(a) = f_r(h(a))$, for each $a \in \chi(r)$. We then move to each child of r and so on, until all leaves are reached and g is defined over all elements in $\mathcal{D}\setminus (A_1\cup\cdots\cup A_n)$. Since Duplicator responds to Spoiler's moves with consistent partial homomorphisms, we have that g is actually a well-defined homomorphism from $(\mathcal{D}, (A_1, \ldots, A_n))$ to $(q', (x'_1, \ldots, x'_n))$.

Now we are ready to show our lemma. Suppose that $(q, \bar{x}) \to_k (q', \bar{x}')$, where $\bar{x} = (x_1, \ldots, x_n)$ and $\bar{x}' = (x_1', \ldots, x_n')$, for some $n \geq 0$. Assume that $(q'', \bar{x}'') \to (q' \land q, \bar{z})$ via a homomorphism h, for $q''(\bar{x}'') \in \mathsf{GHW}(k)$, and suppose that $\bar{x}'' = (x_1'', \ldots, x_n'')$ and $\bar{z} = (z_1, \ldots, z_n)$. For each $i \in \{1, \ldots, n\}$, we define V_i to be the set of variables x in q'' such that $h(x) = z_i$. In particular, $x_i'' \in V_i$, for each $i \in \{1, \ldots, n\}$. We define V to be the set of variables x in q'' such that h(x) = y, where y is an existentially quantified variable of q. Similarly, we define V' with respect to the existentially quantified variables of q'. Note that the sets V, V', V_1, \ldots, V_n form a partition of the variables of q''.

Recall that $\mathcal{D}_{q''}$ be the canonical database of q''. Since $q'' \in \mathsf{GHW}(k)$, we know that

$$(\mathcal{D}_{q''}, (\{x_1''\}, \dots, \{x_n''\}))$$

has generalized hypertreewidth at most k, as defined above. Let \mathcal{D}_V be the database induced in $\mathcal{D}_{q''}$ by the set of variables $V \cup V_1 \cup \cdots \cup V_n$, i.e., the set of atoms $R(\bar{t}) \in \mathcal{D}_{q''}$ such that each element in \bar{t} is in $V \cup V_1 \cup \cdots \cup V_n$. We now show that

$$(\mathcal{D}_V,(V_1,\ldots,V_n))$$

has also generalized hypertreewidth at most k. Indeed, let (T,χ) be the tree decomposition of $(\mathcal{D}_{q''}, (\{x_1''\}, \ldots, \{x_n''\}))$ of width at most k. Define χ' such that for each $t \in T$, we have that $\chi'(t) = \chi(t) \cap V$. We claim that (T,χ') is a tree decomposition of $(\mathcal{D}_V, (V_1, \ldots, V_n))$ of width at most k.

In fact, since (T,χ) is a tree decomposition, we have that, for each $a \in V$, it is the case that the set $\{t \in T \mid a \in \chi'(t)\}$ is connected; and for each atom $R(\bar{a}) \in \mathcal{D}_V$, there is a node $t \in T$ such that $\bar{a} \cap V \subseteq \chi'(t)$. To see that the width of (T,χ') is bounded by k, let t be a node in T. Since the width of (T,χ) is at most k, there are ℓ atoms $R(\bar{a}_1), \ldots, R(\bar{a}_\ell)$ in $\mathcal{D}_{q''}$, with $\ell \leq k$, such that $\chi(t) \subseteq \bigcup_{1 \leq i \leq \ell} \bar{a}_i$. Let $R(\bar{a}_{i_1}), \ldots, R(\bar{a}_{i_p})$, where $1 \leq i_1 < \cdots < i_p \leq \ell$ and $p \leq \ell$, be the atoms in $\{R(\bar{a}_1), \ldots, R(\bar{a}_\ell)\}$ that contain an element in $\chi'(t)$. Since $\chi'(t) \subseteq \chi(t)$, it is the case that $\chi'(t) \subseteq \bigcup_{1 \leq j \leq p} \bar{a}_{i_j}$. It suffices to show that each $R(\bar{a}_{i_j})$ is actually an atom in \mathcal{D}_V , for $1 \leq j \leq p$. Towards a contradiction, suppose that this is not the case. Then, there is an atom in $\mathcal{D}_{q''}$ that contains simultaneously one variable in $\chi'(t) \subseteq V$ and one variable in V'. By the definitions of V' and V, and the fact that h is a homomorphism, it follows that there is an atom in $(q' \land q)(\bar{z})$ that mentions simultaneously one

existentially quantified variable from q' and one from q; this contradicts the definition of $(q' \wedge q)(\bar{z})$. We conclude that the generalized hypertreewidth of $(\mathcal{D}_V, (V_1, \ldots, V_n))$ is at most k.

Recall that h is our initial homomorphism from (q'', \bar{x}'') to $(q' \land q, \bar{z})$. Let h_V be the restriction of h to the set $V \cup V_1 \cup \cdots \cup V_n$. By construction,

$$(\mathcal{D}_V, (V_1, \dots, V_n)) \rightarrow (q, (x_1, \dots, x_n))$$

via homomorphism h_V . We can then apply Lemma 10 and obtain that

$$(\mathcal{D}_V, (V_1, \ldots, V_n)) \rightarrow (q', (x'_1, \ldots, x'_n))$$

via a homomorphism h'. We define our required homomorphism g from (q'', \bar{x}'') to (q', \bar{x}') as follows: if $a \in V \cup V_1 \cup \cdots \cup V_n$, then g(a) = h'(a); otherwise, if $a \in V'$, then g(a) = h(a). To see that g is a homomorphism, it suffices to consider an atom $R(\bar{a}) \in \mathcal{D}_{q''}$ such that \bar{a} contains an element in V' and one element not in V', and show that $R(g(\bar{a})) \in \mathcal{D}_{q'}$. Let A be the set of elements in \bar{a} that are not in V'. As mentioned above, there are no atoms in $\mathcal{D}_{q''}$ mentioning elements in V' and V simultaneously, thus $A \subseteq V_1 \cup \cdots \cup V_n$. In particular, h(a) = h'(a), for each $a \in A$. It follows that $R(g(\bar{a})) = R(h(\bar{a}))$, from which we conclude that $R(g(\bar{a})) \in \mathcal{D}_{q'}$.

Proof (Proposition 10) Consider the Boolean CQ q from Figure 2, defined as

$$q = \exists x \exists y \exists z (P_a(x,y) \land P_a(y,x) \land P_a(y,z) \land P_a(z,y) \land P_b(z,x) \land P_b(x,z)),$$

and the CQ q' from the same figure defined by

$$q' = \exists x \exists y_1 \exists y_2 \exists z \big(P_a(x, y_1) \land P_a(y_1, x) \land P_a(y_2, z)$$
$$\land P_a(z, y_2) \land P_b(z, x) \land P_b(x, z) \big).$$

For each $n \geq 1$, we define the CQ

$$q_n = \exists x_1 \cdots \exists x_{n+1} \big(P_a(x_1, x_2) \wedge \cdots \wedge P_a(x_n, x_{n+1}) \wedge P_b(x_1, x_1) \wedge P_b(x_{n+1}, x_{n+1}) \big).$$

Observe that $q' \wedge q_n \in \mathsf{GHW}(1)$, for each $n \geq 1$. We now show that, for each $n \geq 1$, $q' \wedge q_n$ is an incomparable $\mathsf{GHW}(1)$ - Δ -approximation of q. As mentioned in Example 2, we have that $q \to_1 q'$. In particular $q \to_1 (q' \wedge q_n)$. Clearly, $q \not\to (q' \wedge q_n)$. Also, $q_n \not\to q$ since variables x_1 and x_{n+1} of q_n cannot be mapped to any variable in q via a homomorphism. Therefore, $(q' \wedge q_n) \not\to q$. By Theorem 11, it follows that $q' \wedge q_n$ is an incomparable $\mathsf{GHW}(1)$ - Δ -approximation of q.

Now we show that the CQs $\{q' \land q_n\}_{n \geq 1}$ form a family of non-equivalent CQs. First note that $q_n \not\to q'$, for each $n \geq 1$. Also, observe that $q_i \to q_j$ iff i = j, for $i, j \geq 1$. It follows that for each $i, j \geq 1$, such that $i \neq j$, it is the case that $(q' \land q_i) \not\to (q' \land q_j)$ and $(q' \land q_j) \not\to (q' \land q_i)$. In particular, $\{q' \land q_n\}_{n \geq 1}$ is a family of non-equivalent CQs.

Proof (Proposition 11) As already mentioned, the CONP upper bound follows directly from Theorem 11. For the lower bound, we consider the Non-Hom(H) problem, for a fixed directed graph H, which asks, given a directed graph G, whether $G \not\to H$. Let us assume that, for each $k \ge 1$, there is a directed graph H_k such that:

- 1. $H_k \in \mathsf{GHW}(k)$, or more formally, the Boolean CQ q_{H_k} whose canonical database is H_k belongs to $\mathsf{GHW}(k)$.
- 2. Non-Hom (H_k) is coNP-complete even when the input directed graph G satisfies that $H_k \not\to G$.

We later explain how to obtain these graphs H_k 's. Now we reduce from the restricted version of Non-Hom (H_k) given by item (2) above. Let G be a directed graph such that $H_k \not\to G$. We first check in polynomial time whether $G \to_k H_k$. If $G \not\to_k H_k$, we output a fixed pair q_0, q'_0 such that $q'_0 \in \mathsf{GHW}(k)$ and q'_0 is an incomparable $\mathsf{GHW}(k)$ - Δ -approximation of q_0 . In case that $G \to_k H_k$, we output the pair q_G, q_{H_k} , where q_G and q_{H_k} are Boolean CQs whose canonical databases are precisely G and H_k , respectively. Since $q_{H_k} \in \mathsf{GHW}(k)$ by item (1) above, the reduction is well-defined.

Suppose first that $G \not\to H_k$. If $G \not\to_k H_k$, then we are done, since q_0' is an incomparable $\mathsf{GHW}(k)$ - Δ -approximation of q_0 . Otherwise, if $G \to_k H_k$, since $G \not\to H_k$ and $H_k \not\to G$ (item (2) above), Theorem 11 implies that q_{H_k} is an incomparable $\mathsf{GHW}(k)$ - Δ -approximation of q_G . On the other hand, assume that $G \to H_k$. In particular, we have that $G \to_k H_k$, and then, in this case, the reduction outputs the pair q_G, q_{H_k} . Since $G \to H_k$, we conclude that q_{H_k} is not an incomparable $\mathsf{GHW}(k)$ - Δ -approximation of q_G .

It remains to define the directed graph H_k . If $k \geq 2$, it suffices to consider the clique on 2k vertices, that is, the directed graph \mathbf{K}_{2k} whose vertex set is $\{1,\ldots,2k\}$ and whose edges are $\{(i,j)\mid i\neq j, \text{ for } i,j\in\{1,\ldots,2k\}\}$. We have that $\mathbf{K}_{2k}\in\mathsf{GHW}(k)$, and thus item (1) above is satisfied. Also, we can reduce from the non-2k-colorability problem by replacing each undirected edge $\{u,v\}$ of a given undirected graph G, by a directed edge in an arbitrary direction, e.g., from u to v. Clearly, this is a reduction from non-2k-colorability to Non-Hom(\mathbf{K}_{2k}). Also note that the output f(G) of the reduction satisfies that $\mathbf{K}_{2k} \not\to f(G)$, as f(G) has no directed loops nor directed cycles of length 2. Therefore, item (2) above is satisfied. For k=1, it is known from [30] that there is an oriented tree T (i.e., a directed graph whose underlying undirected graph is a tree and has no directed cycles of length 1 (loops) and 2) such that Non-Hom(T) is conP-complete. Since T is an oriented tree then it belongs to $\mathsf{GHW}(1)$, and then item (1) is satisfied. Also, by inspecting the reduction in [30], we have that item (2) also holds.