# Discovery of a Meteor of Interstellar Origin

Amir Siraj<sup>1</sup> and Abraham Loeb<sup>1</sup>

<sup>1</sup>Department of Astronomy, Harvard University, 60 Garden Street, Cambridge, MA 02138, USA

#### ABSTRACT

The first interstellar object, 'Oumuamua, was discovered in the Solar System by Pan-STARRS in 2017, allowing for a calibration of the abundance of interstellar objects of its size  $\sim 100\,$  m. One would expect a much higher abundance of smaller interstellar objects, with some of them colliding with Earth frequently enough to be noticeable. Based on the CNEOS catalog of bolide events, we identify the  $\sim 0.45 \mathrm{m}$  meteor detected at 2014-01-08 17:05:34 UTC as originating from an unbound hyperbolic orbit with 99.999% confidence. We infer that the meteor had an asymptotic speed of  $v_{\infty} \sim 42.1 \pm 5.5 \mathrm{~km~s^{-1}}$  outside of the solar system. Its origin is approximately towards R.A.  $49.4 \pm 4.1^{\circ}$  and declination  $11.2 \pm 1.8^{\circ}$ , implying that its initial velocity vector was  $58 \pm 6 \mathrm{~km~s^{-1}}$  away from the velocity of the Local Standard of Rest (LSR). Its high LSR speed implies a possible origin from the deep interior of a planetary system or a star in the thick disk of the Milky Way galaxy. The local number density of its population is  $10^{6+0.75}_{-1.5} \mathrm{~AU^{-3}}$  or  $9 \times 10^{21}_{-1.5}^{+0.75} \mathrm{~pc^{-3}}$  (necessitating 0.2 - 20 Earth masses of material to be ejected per local star). This discovery enables a new method for studying the composition of interstellar objects, based on spectroscopy of their gaseous debris as they burn up in the Earth's atmosphere.

Keywords: Minor planets, asteroids: general – comets: general – meteorites, meteors, meteoroids

## 1. INTRODUCTION

'Oumuamua was the first interstellar object detected in the Solar System by Pan-STAARS (Meech et al. 2017; Micheli et al. 2018). Several follow-up studies of 'Oumuamua were conducted to better understand its origin and composition (Bannister et al. 2017; Gaidos et al. 2017; Jewitt et al. 2017; Mamajek 2017; Ye et al. 2017; Bolin et al. 2017; Fitzsimmons et al. 2018; Trilling et al. 2018; Bialy & Loeb 2018; Hoang et al. 2018; Siraj & Loeb 2019a,b; Seligman et al. 2019). Its size was estimated to be 20m - 200m, based on Spitzer Space Telescope constraints on its infrared emission given its temperature (Trilling et al. 2018). Forbes & Loeb (2019) predicted that spectroscopy of 'Oumuamua-like objects grazing the Sun could reveal their chemical compositions. Since there should be a higher abundance of interstellar objects smaller than 'Oumuamua, we could observe small interstellar objects impacting the Earth's atmosphere. Spectroscopy of the gaseous debris from such objects as they burn up in the Earth's atmosphere could reveal their composition. This raises the question:

is there definitive evidence of meter-scale interstellar meteors? Previous studies have reported the detections of interstellar dust grains, but as of yet, no study has produced definitive evidence of meter-scale interstellar meteors (Opik 1950; Hajdukova 1994; Taylor et al. 1996; Baggaley 2000; Meisel et al. 2002; Weryk & Brown 2005; Afanasiev et al. 2007; Musci et al. 2012; Engelhardt et al. 2017; Hajdukova et al. 2018).

The CNEOS catalog includes the geocentric velocity components and geographic coordinates for bolides detected by U.S. government sensors.<sup>1</sup> In this *Letter*, we identify a meteor from the CNEOS catalog that is likely of interstellar origin.

### 2. METHODS

We analyzed the bolide events in the CNEOS catalog, and found that the meteor detected at 2014-01-08 17:05:34 UTC had an unusually high heliocentric velocity at impact.<sup>2</sup> Accounting for the motion of the Earth relative to the Sun and the motion of the meteor relative

https://cneos.jpl.nasa.gov/fireballs/

<sup>&</sup>lt;sup>2</sup> The fastest meteor in the CNEOS catalog obtains its high speed from a head-on orbit relative to the Earth and its extrapolated orbit is found to be bound to the Sun. The meteor we focus

to the Earth, we found that the meteor had a heliocentric velocity of  $\sim 60~\rm km\;s^{-1}$  at impact, which implies that the object was unbound. To uncover the kinematic history of this meteor, we integrated its motion from impact backward in time.

The Python code created for this work used the opensource N-body integator software REBOUND<sup>3</sup> to trace the motion of the meteor under the gravitational influence of the Solar System (Rein & Liu 2012).


We initialize the simulation with the Sun, the eight planets, and the meteor, with geocentric velocity vector  $(vx_{obs}, vy_{obs}, vz_{obs}) = (-3.4, -43.5, -10.3) \text{ km s}^{-1}$ , located at 1.3° S 147.6° E, at an altitude of 18.7 km, at the time of impact,  $t_i = 2014\text{-}01\text{-}08 \ 17:05:34 \ \text{UTC}$ , as reported in the CNEOS catalog. We then use the IAS15 adaptive time-step integrator to trace the meteor's motion back in time (Rein & Spiegel 2014).

### 3. RESULTS

## 3.1. Trajectory

There are no substantial gravitational interactions between the meteor and any planet other than Earth for any trajectory within the reported errors. Based on the impact speed reported by CNEOS,  $v_{obs}=44.8~{\rm km~s^{-1}}$ , we find that the meteor was unbound with an asymptotic speed of  $v_{\infty}\sim42.1~{\rm km~s^{-1}}$  outside of the solar system. In order for the object to be bound, the observed speed of  $v_{obs}=44.8~{\rm km~s^{-1}}$  would have to be off by more than 45%, or 20 km s<sup>-1</sup>.

The typical velocity uncertainty for meter-scale impactors in the CNEOS catalog was estimated by Brown et al. (2016) and Granvik & Brown (2018) to be less than 1 km s<sup>-1</sup>, but 2 of 10 events analyzed by Devillepoix et al. (2019) have discrepancies with alternate measurements of up to 28% in speed. These papers underlined the importance of uncertainties and motivated us to retrieve the measurement uncertainties for the 2014-01-08 meteor specifically. The one sigma uncertainties on each of the velocity components are better than  $\pm 10\%$  (M. Heavner (LANL), private communication, May 2019). Given the geocentric velocity vector  $(vx_{obs}, vy_{obs}, vz_{obs}) = (-3.4 \pm 0.34, -43.5 \pm 4.35, -10.3 \pm 0.34, -43.5 \pm 0.34, -10.3 \pm 0.34, -10.34, -10.34, -10.34, -10.34, -10.34, -10.34, -10.34, -10.34, -10.34, -10.34, -10.34, -10.3$ 1.03) km s<sup>-1</sup> and assuming Gaussian statistics, we construct a trivariate normal distribution for the geocentric velocity of the meteor, centered at  $(vx_{obs}, vy_{obs}, vz_{obs}) =$ (-3.4, -43.5, -10.3) km s<sup>-1</sup> and with root-mean-square uncertainties  $(\sigma_x, \sigma_y, \sigma_z) = (0.34, 4.35, 1.03) \text{ km s}^{-1}$ . We also construct a corresponding distribution of helio-



**Figure 1.** Trajectory of the January 8, 2014 meteor (red), shown intersecting with that of Earth (blue) at the time of impact,  $t_i = 2014$ -01-08 17:05:34.

centric impact speeds, and integrate over the probability space where the heliocentric impact speeds would imply a bound origin,  $v \leq 43.6~{\rm km~s^{-1}}$ . The resulting probability that an object with the measured velocities of the 2014-01-08 meteor was actually bound is  $5.5 \times 10^{-8}$ . As a result, we would expect  $10^{-5}$  of the  $\sim 200$  meteors with reported speeds in the CNEOS catalog to be bound objects with incorrectly measured velocities that imply interstellar origins as definitively as the 2014-01-08 meteor. Assuming Poisson statistics for a singular detection, we conclude with 99.999% confidence that the 2014-01-08 meteor was interstellar.

We find that the heliocentric orbital elements of the meteor at time of impact are as follows: semi-major axis,  $a=-0.47\pm0.15$  AU, eccentricity,  $e=2.4\pm0.3$ , inclination  $i=10\pm2^\circ$ , longitude of the ascending node,  $\Omega=108\pm1^\circ$ , argument of periapsis,  $\omega=58\pm2^\circ$ , and true anomaly,  $f=-58\pm2^\circ$ . The trajectory is shown in Fig. 1. The origin is towards R.A.  $49.4\pm4.1^\circ$  and declination  $11.2\pm1.8^\circ$ . The heliocentric incoming velocity at infinity of the meteor in right-handed Galactic coordinates is  $v_\infty(\mathrm{U},\mathrm{V},\mathrm{W})=(32.7\pm5.8,-4.5\pm1.5,26.1\pm2.0)$  km s<sup>-1</sup>, which is  $58\pm6$  km s<sup>-1</sup> away from the velocity of the Local Standard of Rest (LSR), (U, V, W)<sub>LSR</sub> = (-11.1, -12.2, -7.3) km s<sup>-1</sup> (Schonrich et al. 2010).

### 3.2. Size distribution

Given the impact speed of the meteor,  $\sim 44.8~\rm km~s^{-1}$ , and the total impact energy,  $4.6\times 10^{18}$  ergs, the meteor mass was approximately  $4.6\times 10^5$  g. Assuming bulk density values of 1.7 g/cm³ and 0.9 g/cm³ for Type II and Type IIIa objects respectively, we obtain a radius, R, of 0.4m - 0.5m for a spherical geometry (Ceplecha 1988; Palotai et al. 2018).

The CNEOS catalog includes bolide events at a relatively high frequency for the past decade, so we approximate the yearly detection rate of interstellar meteors

on is the second fastest. The orbit of the third fastest meteor in the catalog is possibly bound within uncertainties.

<sup>&</sup>lt;sup>3</sup> https://rebound.readthedocs.io/en/latest/

to be at least  $\sim 0.1~\rm yr^{-1}$ . We estimate the number density of similarly sized interstellar objects by dividing the yearly detection rate by the product of the impact speed of the meteor and the cross sectional area of the Earth, finding the approximate number density of interstellar objects with a size of order  $R \sim 0.45 \rm m$  and a speed  $v \sim 60~\rm km~s^{-1}~km~s^{-1}$  relative to the LSR, to be,

$$n \sim \frac{0.1 \,\mathrm{yr}^{-1}}{(13 \,\mathrm{AU/yr})(5.7 \times 10^{-9} \,\mathrm{AU}^2)} \sim 10^6 \,\mathrm{AU}^{-3}.$$
 (1)

Given 95% Poisson uncertainties, the inferred local number density for interstellar objects of this size is  $n = 10^{6^{+0.75}_{-1.5}} \text{ AU}^{-3}$ . This figure necessitates  $6 \times 10^{22^{+0.75}_{-1.5}}$ similarly size objects, or 0.2 - 20 Earth masses of material, to be ejected per local star. This is at tension with the fact that a minimum-mass solar nebula is expected to have about an Earth mass of total planetesimal material interior to the radius where the orbital speed is  $\sim 60 \text{ km s}^{-1}$  (Desch 2007), with similar values for other planetary systems (Kuchner 2004). Our inferred abundance for interstellar meteors should be viewed as a lower limit since the CNEOS data might have a bias against detection of faster meteors (Brown et al. 2016). Do et al. (2018) estimated the number density of 'Oumuamua-size ( $R \sim 100 \text{m}$ ) objects to be  $0.2 \text{ AU}^{-3}$ . Using this number density, along with our estimated density for  $R \sim 0.45$ m objects, we construct a range of estimates for the slope of the power-law of the size distribution for interstellar objects as shown in Fig. 2. The range of possible power-law slopes, -1.9 to -3.8, is consistent with that inferred for small bodies in the Kuiper belt, -2.5 to -3 (Kenyon & Bromley 2004). The range is also consistent with the lower limits on the flux of  $R \sim 10^{-4}$  m interstellar dust particles calculated by Weryk & Brown (2005), as well as the upper limits on the flux of  $R \sim 10^{-3}$  m interstellar dust particles calculated by Musci et al. (2012), assuming a smooth powerlaw distribution. However, the power-law extrapolation may not hold at all bolide radii down to dust particles.

## 4. DISCUSSION

We presented and analyzed impact data from the meteor detected at 2014-01-08 17:05:34 UTC, showing that it originated from an unbound hyperbolic orbit with 99.999% confidence. The meteor had an




Figure 2. Size distribution of interstellar objects based on the detection of 'Oumuamua and of the meteor detected at 2014-01-08 17:05:34 UTC. Red lines indicate the envelope for possible power-law fits (slopes of -1.9 to -3.8), given 95% Poisson distribution confidence intervals on each number density (based on a single detection for each). The range of possible power-law slopes is consistent with that inferred for small bodies in the Kuiper belt, -2.5 to -3 (Kenyon & Bromley 2004).

asymptotic speed of  $v_{\infty} \sim 42.1 \pm 5.5 \text{ km s}^{-1}$  outside of the Solar System. Its size, trajectory, and excess speed exclude the possibility that it was gravitationally scattered within the Solar System prior to impact (Wiegert 2014). Its  $\sim 58 \pm 6 \text{ km s}^{-1} \text{ deviation}$ from the LSR suggests that it perhaps originated in the thick disk, which has velocity dispersion components of  $(\sigma_{\rm U}, \sigma_{\rm V}, \sigma_{\rm W}) = (50, 50, 50) \, \rm km \, s^{-1} \, relative to the LSR$ (Bland-Hawthorn & Gerhard 2016). However, the ratio of local thick disk stars to thin disk stars is 0.04, making this a minority population. Alternatively, for a parent planetary system with a more typical velocity relative to the LSR, the object could have originated in the deep interior, where the orbital speeds of objects are of the necessary magnitude. Either way, the meteor had an unusual origin. We obtained a range of estimates for the slope of the power-law of the size distribution for interstellar objects implied by the detection of this interstellar meteor and that of 'Oumuamua, which is consistent with that inferred for small bodies in the Kuiper belt. The mass density of interstellar objects of radius  $R \sim 0.45$ m implied by the discovery of this meteor is similar to that of  $R \sim 100$ m objects implied by the discovery of 'Oumuamua, the two mass densities being  $3 \times 10^{25^{+0.75}_{-1.5}}$  kg pc<sup>-3</sup> and  $6 \times 10^{25^{+0.75}_{-1.5}}$  kg pc<sup>-3</sup>, respectively.

The discovery of additional interstellar meteors will serve as an important calibration for population-wide

 $<sup>^4</sup>$  Gravitational focusing by the Earth is negligible since the meteor speed exceeds considerably the escape speed from the Earth. The density enhancement due to gravitational focusing by the Sun is well below the uncertainty in the estimated value of n, so that our inferred range of local values also corresponds to the density outside of the Solar System.

parameters of interstellar objects, including their abundance and origin.

We estimate the impact rate of similarly sized objects with the Earth, given 95% Poisson distribution confidence intervals, to be at least  $0.1^{+0.457}_{-0.097}$  per year. Future meteor surveys could flag incoming objects with excess heliocentric velocities for follow-up pre-impact observations. Spectroscopy of gaseous debris from these objects as they burn up in the Earth's atmosphere would reveal their composition. Given that some isotope ratios are expected to be markedly different for objects of interstellar origin compared to the Solar System, could validate an interstellar origin (Lingam & Loeb 2018a; Forbes & Loeb 2019). Precision tracking with the upcoming Large Synoptic Survey Telescope (LSST<sup>5</sup>) could determine the trajectory of meteors of interstellar origin to their parent systems in the Gaia catalog.<sup>6</sup> Our discovery also implies that at least  $4.5 \times 10^{8^{+0.75}}$  similarly sized interstellar bolide events have occurred over

Earth's lifetime. Potentially, interstellar meteors could deliver life from another planetary system and mediate panspermia (Ginsburg et al. 2018). Interestingly, the high speed for the meteor discussed here implies a likely origin in the habitable zone of the abundant population of dwarf stars, indicating that similar objects could carry life from their parent planetary systems.

### ACKNOWLEDGEMENTS

We thank Alan Hurd and Matthew Heavner for their exceptional help in declassifying and retrieving the velocity measurement uncertainties of the 2014-01-08 meteor. Without their help, the interstellar origin of this meteor would have been enshrouded in uncertainty. We also thank Nathan Golovich, Jeff Kuhn, Matthew Payne, Steinn Sigurdsson, and Peter Veres for helpful comments on the manuscript. This work was supported in part by a grant from the Breakthrough Prize Foundation.

## REFERENCES

Afanasiev V. L., Kalenichenko V. V., Karachentsev I. D., 2007, Astrophysical Bulletin, 62(4), 319

Baggaley W. J., 2000, Journal of Geophysical Research, 105(A5), 10353

Bannister M. T., et al., 2017, The Astrophysical Journal, 851, L38

Bialy S., Loeb A., 2018, The Astrophysical Journal, 868, L1
Bland-Hawthorn J., Gerhard O., 2016, Annual Review of Astronomy and Astrophysics, 54, 529

Bolin B. T., et al., 2017, The Astrophysical Journal Letters, Volume 852, Issue 1, article id. L2, 10 pp. (2018)., 852

Brown P., et al., 2016, Icarus, 266, 96

Ceplecha Z., 1988, Bulletin of the Astronomical Institutes of Czechoslovakia, 39, 221

Do A., Tucker M. A., Tonry J., 2018, The Astrophysical Journal, 855, L10

Desch S.J., 2007, The Astrophysical Journal, 671(1), 878
Devillepoix H.A.R., et al., 2019, Monthly Notices of the Royal Astronomical Society, 483(4), 5166

Engelhardt T., et al., 2017, The Astronomical Journal, 153, 133

Fitzsimmons A., et al., 2018, Nature Astronomy, 2, 133 Forbes J., Loeb A., 2018, Turning up the heat on 'Oumuamua. (arXiv:1901.00508) Gaidos E., Williams J., Kraus A., 2017, Research Notes of the AAS, 1, 13

Ginsburg I., Lingam M., Loeb A., 2018, The Astrophysical Journal, 868(1), L12

Granvik M., Brown P., 2018, Icarus, 311(1), 271

Hajdukova M., Jr., 1994, Astronomy and Astrophysics 288(1), 330

Hajdukova M., Sterken, V., Wiegert, P., 2018, European Planetary Science Congress 12

Hoang T., Loeb A., Lazarian A., Cho J., 2018, The Astrophysical Journal, 860(1), 42

Jewitt D., Luu J., Rajagopal J., Kotulla R., Ridgway S., Liu W., Augusteijn T., 2017, The Astrophysical Journal, 850, L36

Kenyon S.J., Bromley B.C., 2018, The Astronomical Journal, 128(4), 1916

Kuchner M.J., 2004, The Astrophysical Journal, 612, 1147

Lingam M., Loeb A., 2018a, The Astronomical Journal, 156 Mamajek E., 2017, Research Notes of the AAS, 1, 21

Marcos C.F., Marcos R.F., Aarseth S.J., 2015, The

Meech K. J., et al., 2017, Nature, 552, 378

Astrophysical Journal, 812(1), 26

Meisel D. D., Janches D., Mathews J. D., 2002, The Astrophysical Journal, 567, 323

Micheli M., et al., 2018, Nature, 559, 223

Musci R., et al., 2012, The Astrophysical Journal, 745, 161

Opik, E. J., 1950, Irish Astronomical Journal 1(3), 80

<sup>&</sup>lt;sup>5</sup> https://www.lsst.org/

<sup>6</sup> https://gea.esac.esa.int/archive/

- Palotai, C., et al., 2018, Analysis of June 2, 2016 bolide event. (arXiv:1801.05072)
- Rein H., Liu S.-F., 2012, Astronomy & Astrophysics, 537, A128
- Rein H., Spiegel D.S., 2014, Monthly Notices of the Royal Astronomical Society, 446(2), 1424
- Schonrich, R., Binney, J., & Dehnen, W., 2010, Monthly Notices of the Royal Astronomical Society, 403(4), 1829
- Seligman D., Laughlin G., Batygin K., 2019, The Astrophysical Journal Letters,

- Siraj A. & Loeb A., 2019, The Astrophysical Journal Letters, 872(1), L10
- Siraj A. & Loeb A., 2019, Research Notes of the American Astronomical Society, 3(1), 15
- Taylor A. D., Baggaley W. J., Steel D. I., 2018, Nature, 380, 323
- Trilling, D., et al., 2018, The Astronomical Journal, 156, 261
- Weryk R. J., Brown P., 2005, Icarus, 95, 221.
- Wiegert P., 2014, Icarus, 242, 112.
- Ye Q.-Z., Zhang Q., Kelley M. S. P., Brown P. G., 2017, The Astrophysical Journal, 851, L5