

New Kloosterman sum identities from the Helleseth-Zinoviev result on Z_4 -linear Goethals codes

Minglong Qi · Shengwu Xiong

Received: date / Accepted: date

Abstract In the paper of Tor Helleseht and Victor Zinoviev (Designs, Codes and Cryptography, **17**, 269-288(1999)), the number of solutions of the system of equations from Z_4 -linear Goethals codes G_4 was determined and stated in Theorem 4. We found that Theorem 4 is wrong for m even. In this note, we complete Theorem 4, and present a series of new Kloosterman sum identities deduced from Theorem 4. Moreover, we show that several previously established formulas on the Kloosterman sum identities can be rediscovered from Theorem 4 with much simpler proofs.

Keywords Kloosterman sum identities · Z_4 -linear Goethals codes · nonlinear system of equations · exponential sums

Mathematics Subject Classification (2010) 11L05 · 11T23

1 Introduction

Let m be a positive integer, $q = 2^m$, $F := \mathbb{F}_q$ be the finite field of q elements, $F^* := F \setminus \{0\}$, and $F^{**} := F \setminus \{0, 1\}$. The well known Kloosterman sums [1, 2] are defined by

$$K(a) = \sum_{x \in F^*} (-1)^{\text{Tr}(ax+1/x)} \quad (1)$$

where $a \in F^*$, and $\text{Tr}(\cdot)$ is the trace function of F over \mathbb{F}_2 .

Let $b, c \in F$. The problem of finding the coset weight distribution of Z_4 -linear Goethals codes G_4 , is transformed into solving the following nonlinear system of

Communicated by Pascale Charpin, Alexander Pott, Dieter Jungnickel

Minglong Qi · Shengwu Xiong
School of Computer Science and Technology, Wuhan University of Technology
Mafangshan West Campus, 430070 Wuhan City, China
E-mail: mlqiecully@163.com (Minglong Qi)
E-mail: xiongsw@whut.edu.cn (Shengwu Xiong)

equations over F [3]:

$$\begin{cases} x + y + z + u & = 1 \\ u^2 + xy + xz + xu + yz + yu + zu & = b^2 \\ x^3 + y^3 + z^3 + u^3 & = c \end{cases} \quad (2)$$

where x, y, z and u are pairwise distinct elements of F . The number of solutions of (2) (see p.284 of [3]), denoted by $\mu_2(b, c)$, is given by Theorem 4 of [3].

We found that Theorem 4 of [3], which gives the explicit evaluation of $\mu_2(b, c)$, is wrong for m even. It is obvious that the authors of [3] forgot to take account of the fact, that $\text{Tr}(1) = 1$ for m odd and 0 for m even, in the last step of the proof of Theorem 4. The following is the correct version of Theorem 4 of [3]:

Theorem 1 *Let $\mu_2(b, c)$ be the number of different 4-tuples (x, y, z, u) , where x, y, z, u are pairwise distinct elements of F , which are solutions to the system (2) over F , where b, c are arbitrary elements of F .*

(1) *If m is odd and $\text{Tr}(c) = 1$ or m is even and $\text{Tr}(c) = 0$, then*

$$\mu_2(b, c) = \frac{1}{6} (q - 8 + (-1)^{\text{Tr}(b)} (K(k_1 k_2) - 3)). \quad (3)$$

(2) *If m is odd and $\text{Tr}(c) = 0$ or m is even and $\text{Tr}(c) = 1$, then*

$$\mu_2(b, c) = \frac{1}{6} (q - 2 - (-1)^{\text{Tr}(b)} (K(k_1 k_2) + 3)). \quad (4)$$

Where $k_1 = b^2 + c + 1$ and $k_2 = b^2 + b + c + \sqrt{c}$.

In this note, we will show that Theorem 4 of [3] implies not only a series of new Kloosterman sum identities, but almost all previously discovered formulas on the identities for Kloosterman sums, thanks to the following theorem which is a corollary of that theorem, too:

Theorem 2 *Let $b, c \in F, k_1 = b^2 + c + 1$ and $k_2 = b^2 + b + c + \sqrt{c}$. If $k_1 k_2 \neq 0$, then*

$$K(k_1 k_2) = K(k_1 k_2 + k_2). \quad (5)$$

The idea is to affect a special value to k_1 or k_2 and then substitute them into (5) to arrive at a special Kloosterman sum identity. Let $b, c \in F$, and n, k be rational numbers such that b^n, b^k, c^n, c^k are elements of F . In the sequel, k_1 and k_2 are affected to the following values:

$$\begin{cases} k_1 \in \{cb^n, b^2c^n, cb^n + b^k, b^2c^n + c^k\}, \\ k_2 \in \{cb^n + \sqrt{c}, \sqrt{c}b^n + c, b^2c^n + b, bc^n + b^2\}. \end{cases} \quad (6)$$

The previously discovered formulas on the identities for Kloosterman sums are stated in the following theorem:

Theorem 3 *Let $a \in F^{**}$. Then*

- (1) $K(a^3(a+1)) = K(a(a+1)^3)$ [8, 9, 5, 4] (Helleseeth-Zinoviev Formula I).
 (2) $K(a^5(a+1)) = K(a(a+1)^5)$ [4] (Helleseeth-Zinoviev Formula II).
 (3) $K(a^8(a^4+a)) = K((a+1)^8(a^4+a))$ [6] (Hollmann-Xiang Formula).
 (4) $K(a/(a+1)^4) = K(a^3/(a+1)^4)$ [8] (Shin-Kumar-Helleseeth Formula).

Note that Shin-Kumar-Helleseeth Formula can be deduced from Helleseeth-Zinoviev Formula I by the variable change $a = b/(b+1)$. In this note, we will show that except for Helleseeth-Zinoviev Formula II, every formula of Theorem 3 can be obtained from Theorem 4 of [3] with help of Theorem 2.

In Section 2, we at first prove Theorem 2 with help of Theorem 1 (Theorem 4 of [3]) and two lemmas of [5], treat in details some cases of (6) for k_1 and k_2 which bring us new Kloosterman sum identities, prove Helleseeth-Zinoviev Formula I and Hollmann-Xiang Formula, and finally generalize Shin-Kumar-Helleseeth Formula.

2 New identities for Kloosterman sums

2.1 Proof of Theorem 2

Before proving Theorem 2, we need several preliminary lemmas:

Lemma 1 ([1]) *Let $a \in F^*$. Then, $K(a) = K(a^2)$.*

Lemma 2 ([7]) *Let $a, b \in F$. Then,*

- (1) $\text{Tr}(a) = \text{Tr}(a^2)$, $\text{Tr}(a+b) = \text{Tr}(a) + \text{Tr}(b)$.
 (2) $\text{Tr}(1) = 1$ if m is odd, and $\text{Tr}(1) = 0$ if m is even.

The symmetry of the solutions of the nonlinear system (2) is characterized by the following two lemmas:

Lemma 3 [5, Lemma 10] *Let b, c be any elements of F , where F has order 2^m . Let m be even. Let $\mu_2(b, c)$ denote the number of solutions to system (2). Then the following symmetry conditions are valid:*

$$\mu_2(b, c) = \mu_2(b+1, c) \quad (7)$$

and

$$\mu_2(b, c) = \mu_2(b, c+1). \quad (8)$$

Lemma 4 [5, Lemma 11] *Let b, c be any elements of F , where F has order 2^m . Let m be odd. Let $\mu_2(b, c)$ denote the number of solutions to system (2). Then the following symmetry conditions are valid:*

$$\mu_2(b, c) + \mu_2(b+1, c) = \frac{1}{3} \times \begin{cases} (2^m - 2) & \text{if } \text{Tr}(c) = 0, \\ (2^m - 8) & \text{if } \text{Tr}(c) = 1, \end{cases} \quad (9)$$

and

$$\mu_2(b, c) + \mu_2(b, c+1) = \frac{1}{3} \times \begin{cases} (2^m - 2) & \text{if } \text{Tr}(c) = 1, \\ (2^m - 8) & \text{if } \text{Tr}(c) = 0. \end{cases} \quad (10)$$

Now, we are ready to prove Theorem 2. The idea of proof is to substitute the formulas for $\mu_2(b, c)$ from Theorem 1 into the formulas of Lemma 3 for m even, and into the formulas of Lemma 4 for m odd.

Proof (of Theorem 2)

(I) Case for m even.

(a) Subcase that $\text{Tr}(c) = 0$.

Note that $\text{Tr}(1) = 0$ for m even by Lemma 2, and $\text{Tr}(b+1) = \text{Tr}(b) + \text{Tr}(1) = \text{Tr}(b)$. Let $k'_1 := (b+1)^2 + c + 1, k'_2 := (b+1)^2 + (b+1) + c + \sqrt{c}$. It is clear that $k'_1 = k_1 + 1, k'_2 = k_2$ and $k'_1 k'_2 = k_1 k_2 + k_2$, where $k_1 = b^2 + c + 1$ and $k_2 = b^2 + b + c + \sqrt{c}$. From (3) of Theorem 1, we have

$$\begin{aligned}\mu_2(b+1, c) &= \frac{1}{6}(q-8 + (-1)^{\text{Tr}(b+1)}(K(k'_1 k'_2) - 3)) \\ &= \frac{1}{6}(q-8 + (-1)^{\text{Tr}(b)}(K(k_1 k_2 + k_2) - 3)) \text{ and,} \\ \mu_2(b, c) &= \frac{1}{6}(q-8 + (-1)^{\text{Tr}(b)}(K(k_1 k_2) - 3)).\end{aligned}$$

Theorem 2 follows from (7) of Lemma 3 for this subcase.

Let $k''_1 := b^2 + (c+1) + 1, k''_2 := b^2 + b + (c+1) + \sqrt{c+1}$. It is clear that $k''_1 = k_1 + 1, k''_2 = k_2$ and $k''_1 k''_2 = k_1 k_2 + k_2$. Since $\text{Tr}(c+1) = \text{Tr}(c) = 0$, from (3) of Theorem 1, we obtain

$$\begin{aligned}\mu_2(b, c+1) &= \frac{1}{6}(q-8 + (-1)^{\text{Tr}(b)}(K(k''_1 k''_2) - 3)) \\ &= \frac{1}{6}(q-8 + (-1)^{\text{Tr}(b)}(K(k_1 k_2 + k_2) - 3)) \text{ and,} \\ \mu_2(b, c) &= \frac{1}{6}(q-8 + (-1)^{\text{Tr}(b)}(K(k_1 k_2) - 3)).\end{aligned}$$

Again, Theorem 2 follows from (8) of Lemma 3 for this subcase.

(b) Subcase that $\text{Tr}(c) = 1$.

For this subcase, we use (4) of Theorem 1 and (7) or (8) of Lemma 3 to prove Theorem 2, and omit the details due to limited space.

(II) Case for m odd.

For this case, we use (9) or (10) of Lemma 4 and Theorem 1 to prove Theorem 2, which is similar to the case for m even and omitted.

□

2.2 New Kloosterman sum identities and the proof of Theorem 3

Theorem 4 Let $c \in F$, and n be a rational number such that c^n is an element of F . If $c^{4n} + 1 \neq 0$, then

$$K\left(\frac{(c^{n+1} + c^n)^3(c^{n+1} + 1)}{c^{4n} + 1}\right) = K\left(\frac{(c^{n+1} + c^n)(c^{n+1} + 1)^3}{c^{4n} + 1}\right). \quad (11)$$

Proof Let $k_1 = b^2 + c + 1 = b^2 c^n$. Then, $b^2 = (c + 1)/(c^n + 1)$, and $k_1 = (c^{n+1} + c^n)/(c^n + 1)$. Further, we can obtain

$$\begin{aligned} k_2 &= b^2 + b + c + \sqrt{c} = \frac{c+1}{c^n+1} + \sqrt{\frac{c+1}{c^n+1}} + c + \sqrt{c}; \\ k_2^2 &= \frac{c^2+1}{c^{2n}+1} + \frac{c+1}{c^n+1} + c^2 + c = \frac{(c^{n+1}+c^n)(c^{n+1}+1)}{c^{2n}+1}; \\ (k_1 k_2)^2 &= \frac{(c^{n+1}+c^n)^3(c^{n+1}+1)}{c^{4n}+1}; (k_1 k_2)^2 + k_2^2 = \frac{(c^{n+1}+c^n)(c^{n+1}+1)^3}{c^{4n}+1}. \end{aligned}$$

The actual theorem follows from Lemma 1 and Theorem 2. \square

Let $L(n) := (c^{n+1} + c^n)^3(c^{n+1} + 1)/(c^{4n} + 1)$, $R(n) := (c^{n+1} + c^n)(c^{n+1} + 1)^3/(c^{4n} + 1)$, then, $K(L(n)) = K(R(n))$. We now prove Helleseeth-Zinoviev Formula I by Theorem 4.

Proof (of Helleseeth-Zinoviev Formula I)

$$\begin{aligned} L(1) &= \frac{(c^2+c)^3(c^2+1)}{c^4+1} = \frac{c^3(c+1)^5}{(c+1)^4} = c^3(c+1), \\ R(1) &= \frac{(c^2+c)(c^2+1)^3}{c^4+1} = \frac{c(c+1)^7}{(c+1)^4} = c(c+1)^3. \end{aligned}$$

Helleseeth-Zinoviev Formula I follows from the case $K(L(1)) = K(R(1))$ of Theorem 4. \square

Corollary 1 Let $c \in F$ such that the rational functions in c occurring in the following formulas are valid. Then,

- (1) $K(c^6(c^2+c+1)/(c+1)^4) = K(c^2(c^2+c+1)^3/(c+1)^4)$,
- (2) $K(c^9(c+1)^3/(c^8+c^4+1)) = K(c^3(c+1)^9/(c^8+c^4+1))$,
- (3) $K((c+1)^8(c^4+c)) = K(c^3(c^3+1)^3)$,
- (4) $K((c^{11}+c^3)(c^5+1)) = K(c(c^5+1)^3)$,
- (5) $K((c+1)^{20}(c^8+c)) = K((c+1)^4(c^8+c)^3)$.

Proof The first formula of Corollary 1 corresponds to the case $K(L(2)) = K(R(2))$ of Theorem 4, and the second formula to $K(L(3)) = K(R(3))$, which the proofs are analogous to that of Helleseeth-Zinoviev Formula I and omitted. We now prove the third formula:

$$\begin{aligned} L(-1/4) &= \frac{(c^{3/4}+c^{-1/4})^3(c^{3/4}+1)}{c^{-1}+1}, R(-1/4) = \frac{(c^{3/4}+c^{-1/4})(c^{3/4}+1)^3}{c^{-1}+1} \\ L(-1/4)^4 &= \frac{(c^3+c^{-1})^3(c^3+1)}{c^{-4}+1} = \frac{(c^4+1)^3(c^4+c)}{c^4+1} = (c+1)^8(c^4+c), \\ R(-1/4)^4 &= \frac{(c^3+c^{-1})(c^3+1)^3}{c^{-4}+1} = \frac{(c^4+1)(c^4+c)^3}{c^4+1} = c^3(c^3+1)^3. \end{aligned}$$

The third formula follows from $K(L(-1/4)) = K(R(-1/4))$ of Theorem 4 and Lemma 1. The fourth formula arises from the case $K(L(1/4)) = K(R(1/4))$ and the fifth formula from $K(L(-1/8)) = K(R(-1/8))$, which the proofs are similar to that of the third formula and omitted. \square

Proof (of Hollmann-Xiang Formula)

From Helleseeth-Zinoviev Formula I, we obtain $K(c^3(c^3+1)^3) = K(c^9(c^3+1)) = K(c^8(c^4+c))$. From the third formula of Corollary 1, we have $K((c+1)^8(c^4+c)) = K(c^3(c^3+1)^3) = K(c^8(c^4+c))$. \square

Remark that Shin-Kumar-Helleseeth Formula is the specific case $K(L(-2)) = K(R(-2))$ of Theorem 4, which we omit the proof.

2.3 New Kloosterman sum identities and the generalization of Shin-Kumar-Helleseeth Formula

In this subsection, we establish several identities for Kloosterman sums which generalize Shin-Kumar-Helleseeth Formula.

Theorem 5 *Let $b \in F$, and n, k be rational numbers such that b^n, b^k are elements of F . If $b^{4n} + 1 \neq 0$, then,*

$$K\left(\frac{(b^{n+2} + b^k + 1)(b^{n+2} + b^n + b^k)^3}{b^{4n} + 1}\right) = K\left(\frac{(b^{n+2} + b^k + 1)^3(b^{n+2} + b^n + b^k)}{b^{4n} + 1}\right).$$

Proof Let $k_1 = b^2 + c + 1 = cb^n + b^k$, then $c = (b^k + b^2 + 1)/(b^n + 1)$. After an elementary algebraic computation over \mathbb{F}_2 , we get

$$\begin{aligned} k_1^2 &= (b^{n+2} + b^n + b^k)^2 / (b^{2n} + 1), \\ k_2^2 &= (b^{n+2} + b^k + 1)(b^{n+2} + b^n + b^k) / (b^{2n} + 1), \\ (k_1 k_2)^2 &= (b^{n+2} + b^k + 1)(b^{n+2} + b^n + b^k)^3 / (b^{4n} + 1), \\ (k_1 k_2)^2 + k_2^2 &= (b^{n+2} + b^k + 1)^3 (b^{n+2} + b^n + b^k) / (b^{4n} + 1). \end{aligned}$$

The actual theorem follows from Lemma 1 and Theorem 2. \square

We obtain the following corollaries by affecting particular values to n, k in Theorem 5, which the proofs are similar to Corollary 1 and omitted.

Corollary 2 *Set $n = 1, k = 0$ in Theorem 5. Then*

$$K\left(\frac{b^3(b^3 + b + 1)^3}{(1 + b)^4}\right) = K\left(\frac{b^9(b^3 + b + 1)}{(1 + b)^4}\right).$$

Corollary 3 *Set $n = 1, k = 2$ in Theorem 5. Then*

$$K\left(\frac{(b^3 + b^2 + 1)(b^3 + b^2 + b)^3}{(1 + b)^4}\right) = K\left(\frac{(b^3 + b^2 + 1)^3(b^3 + b^2 + b)}{(1 + b)^4}\right).$$

Corollary 4 Set $n = -1$ in Theorem 5. Then

$$K\left(\frac{(b^{k+1} + b^2 + b)(b^{k+1} + b^2 + 1)^3}{(1 + b)^4}\right) = K\left(\frac{(b^{k+1} + b^2 + b)^3(b^{k+1} + b^2 + 1)}{(1 + b)^4}\right).$$

We can obtain analogous results as Theorem 4 and Theorem 5 if we affect one of remainder values from (5) to k_1 or k_2 . For instance, set $k_2 := b^2 + b + c + \sqrt{c} = b^n\sqrt{c} + c$, we obtain

Theorem 6 Let $b \in F$, and n be rational number such that b^n is an element of F . If $b^{4n} + 1 \neq 0$, then,

$$K\left(\frac{(b^{n+1} + b^2)(b^{n+1} + b^n + b^2 + 1)^3}{b^{4n} + 1}\right) = K\left(\frac{(b^{n+1} + b^2)^3(b^{n+1} + b^n + b^2 + 1)}{b^{4n} + 1}\right).$$

Proof The proof for the actual theorem is similar to that of Theorem 4 and Theorem 5 and omitted. □

Set $n = 3$ in the formula of Theorem 6, we obtain an interesting corollary which the proof is similar to Corollary 1 and omitted:

Corollary 5 Let $b \in F$ such that $b^8 + b^4 + 1 \neq 0$. Then,

$$K\left(\frac{(b^3 + b^2)(b^3 + b + 1)^3}{b^8 + b^4 + 1}\right) = K\left(\frac{(b^3 + b^2)^3(b^3 + b + 1)}{b^8 + b^4 + 1}\right).$$

3 Conclusion

In this note, we obtained a series of new Kloosterman sum identities and rediscovered several previously found formulas with much simpler proof from Theorem 4 of [3]. The exception is that the formula, $K(a^5(1 + a)) = K(a(1 + a)^5)$, we believe, could not be deduced from Theorem 4 of [3] and Theorem 2.

References

1. Carlitz, L.: Kloosterman sums and finite field extensions, Acta Arithmetica, **15**(2), 179-193 (1969).
2. Charpin P., Helleseht T., Zinoviev V.: The divisibility modulo 24 of Kloosterman sums on $GF(2^m)$, m odd, Journal of Combinatorial Theory, Series A, **144**, 322-338 (2007).
3. Helleseht T., Zinoviev V.: On Z_4 -Linear Goethals Codes and Kloosterman Sums, Designs, Codes and Cryptography, **17**, 269-288 (1999).
4. Helleseht T., Zinoviev V.: New Kloosterman sums identities over F_{2^m} for all m , Finite Fields Their Appl., **9**, 187-193 (2003).
5. Helleseht T., Zinoviev V.: On a new identity for Kloosterman sums and nonlinear system of equations over finite fields of characteristic 2, Discrete Math., **274**, 109-124 (2004).
6. Hollmann Henk D.L., Xiang Q.: Kloosterman sums identities over F_{2^m} , Discrete Math., **279**, 277-286 (2004).
7. Lidl R., Niederreiter H.: Finite Fields, Encyclopedia of Mathematics and Its Applications, 2nd ed., Cambridge University Press, Vol. 20 (1997).
8. Shin D.J., Kumar P.V., Helleseht T.: 3-Designs from the Z_4 -Goethals codes via a new Kloosterman sum identity, Des. Codes Cryptography, **28**, 247-263 (2003).
9. Shin D.J., Sung W.J.: A new Kloosterman sum identity over F_{2^m} for odd m , Discrete Math., **268**, 337-341 (2003).