Efficient Black-Box Identity Testing for Free Group Algebra

V. Arvind[∗] Abhranil Chatterjee† Rajit Datta‡ Partha Mukhopadhyay§

April 30, 2019

Abstract

Hrubeš and Wigderson [\[HW14\]](#page-16-0) initiated the study of noncommutative arithmetic circuits with division computing a noncommutative rational function in the *free skew field*, and raised the question of rational identity testing. It is now known that the problem can be solved in deterministic polynomial time in the white-box model for noncommutative formulas with inverses, and in randomized polynomial time in the black-box model [\[GGOW16,](#page-16-1) [IQS18,](#page-16-2)[DM18\]](#page-16-3), where the running time is polynomial in the size of the formula.

The complexity of identity testing of noncommutative rational functions remains open in general (when the formula size is not polynomially bounded). We solve the problem for a natural special case. We consider polynomial expressions in the free group algebra $\mathbb{F}\langle X, X^{-1}\rangle$ $\mathbb{F}\langle X, X^{-1}\rangle$ $\mathbb{F}\langle X, X^{-1}\rangle$ ¹ where $X = \{x_1, x_2, \ldots, x_n\}$, a subclass of rational expressions of inversion height one. Our main results are the following.

- 1. Given a degree d expression f in $\mathbb{F}\langle X, X^{-1}\rangle$ as a black-box, we obtain a randomized $\text{poly}(n, d)$ algorithm to check whether f is an identically zero expression or not. We obtain this by generalizing the Amitsur-Levitzki theorem [\[AL50\]](#page-15-0) to $\mathbb{F}\langle X, X^{-1}\rangle$. This also yields a deterministic identity testing algorithm (and even an expression reconstruction algorithm) that is polynomial time in the sparsity of the input expression.
- 2. Given an expression f in $\mathbb{F}\langle X, X^{-1}\rangle$ of degree at most D, and sparsity s, as black-box, we can check whether f is identically zero or not in randomized $\text{poly}(n, \log s, \log D)$ time.

1 Introduction

Noncommutative computation is an important sub-area of arithmetic circuit complexity. In the usual arithmetic circuit model for noncommutative computation, the arithmetic operations are addition and multiplication. However, the multiplication gates respect the input order since the variables are noncommuting. Analogous to commutative arithmetic computation, the central questions are to show lower bounds for explicit polynomials and derandomization of polynomial identity testing (PIT) for noncommutative polynomial rings. Exploiting the limited cancellations, strong lower bounds and PIT results are known for noncommutative computations (in contrast to the commutative setting). Nisan [\[Nis91\]](#page-16-4) has shown that any algebraic branching program (ABP) computing the $n \times n$ noncommutative Determinant or Permanent polynomial requires exponential (in n) size. On the PIT front, Raz and Shpilka [\[RS05\]](#page-16-5) have shown a deterministic polynomial-time PIT for noncommutative ABPs in the white-box model. A quasi-polynomial time derandomization is also known for the black-box model [\[FS12\]](#page-16-6). However, for general circuits there are no better results (either lower bound or PIT) than known in the commutative setting.

The randomized polynomial-time PIT algorithm for noncommutative circuits computing a polynomial of polynomially bounded degree [\[BW05\]](#page-15-1) follows from Amitsur-Levitzki theorem [\[AL50\]](#page-15-0). The Amitsur-Levitzki theorem states that a nonzero noncommutative polynomial $p \in \mathbb{F}\langle X \rangle$ of degree $\langle 2k \rangle$ cannot be an identity for the matrix ring $M_k(\mathbb{F})$. Additionally, it is shown that a nonzero noncommutative polynomial does not vanish on matrices of dimension logarithmic in the sparsity of the polynomial, yielding a randomized polynomial time algorithm for noncommutative circuits computing a nonzero polynomial of exponential degree and exponential sparsity [\[AJMR17\]](#page-15-2).

Hrubeš and Wigderson $[HW14]$ initiated the study of noncommutative computation with inverses. In the commutative world, it suffices to consider additions and multiplications. By Strassen's result [\[Str73\]](#page-17-0) (extended to finite fields [\[HY11\]](#page-16-7)), divisions can be efficiently replaced by polynomially many additions and multiplications. However, divisions in noncommutative computation are more complex [\[HW14\]](#page-16-0). In the same paper [\[HW14\]](#page-16-0) the authors introduce rational identity testing: Given a noncommutative formula involving addition, multiplication and division gates, efficiently check if the resulting rational expression is identically zero in the free skew-field of noncommutative rational functions. They show that the rational identity testing problem reduces to the following SINGULAR problem:

Given a matrix $A_{n\times n}$ where the entries are linear forms over noncommuting variables $\{x_1, x_2, \ldots, x_n\}$, is A invertible in the free skew-field?

In the white-box model the problem is in deterministic polynomial time, and in randomized polynomial time in the black-box model [\[GGOW16,](#page-16-1) [IQS18,](#page-16-2) [DM18\]](#page-16-3). Specifically, for rational formulas of size s, random matrix substitutions of dimension linear in s suffices to test if the rational expression is identically zero [\[DM18\]](#page-16-3).

The complexity of identity testing for general rational expressions remains open. For example, given a noncommutative circuit involving addition, multiplication and division gates, no efficient algorithm is known to check if the resulting rational expression is identically zero in the free skew-field of noncommutative rational functions. In order to precisely formulate the problem, we define classes of rational expressions based on Bergman's definition [\[Ber76\]](#page-15-3) of inversion height which we now recall and elaborate upon with some notation.

Definition 1. [\[Ber76\]](#page-15-3) Let X be a set of free noncommuting variables. Polynomials in the free ring $\mathbb{F}\langle X\rangle$ are defined to be rational expressions of height 0. A rational expression of height $i + 1$ is inductively defined to be a polynomial in rational expressions of height at most i, and inverses of such expressions.

Let $\mathcal{E}_{d,0}$ denote all polynomials of degree at most d in the free ring $\mathbb{F}\langle X\rangle$. We inductively define rational expressions in $\mathcal{E}_{d,i+1}$ as follows: Let f_1, f_2, \ldots, f_r and g_1, g_2, \ldots, g_s be rational expressions in $\mathcal{E}_{d,i}$ in the variables x_1, x_2, \ldots, x_n . Let $f(y_1, y_2, \ldots, y_s, z_1, z_2, \ldots, z_r)$ be a degree-d polynomial in $\mathbb{F}\langle X \rangle$. Then $f(g_1, g_2, \ldots, g_s, f_1^{-1}, f_2^{-1}, \ldots, f_r^{-1})$ is a rational expression (of inversion height $i+1$) in $\mathcal{E}_{d,i+1}$.

Black-box identity testing for rational expressions is not well understood in general. Bergman has shown [\[Ber76,](#page-15-3) Proposition 5.1] that there are rational expressions that are nonzero over a dense subset of 2×2 matrices but evaluate to zero on dense subsets of 3×3 matrices. This makes it difficult to formulate an Amitsur-Levitzki type of theorem [\[AL50\]](#page-15-0) for rational expressions.

Remark 1. In this connection, we note that Hrubes and Wigderson $|HW14|$ have observed that testing if a 'correct' rational expression Φ is not identically zero is equivalent to testing if the rational expression Φ^{-1} is 'correct'. I.e. testing if a correct rational expression of inversion height i is identically zero or not can be reduced to testing if a rational expression of inversion height $i+1$ is correct or not. Furthermore, testing if a rational expression of inversion height one is correct can be done by applying (to each inversion operation in this expression) a theorem of Amitsur (see [\[Row80,](#page-16-8)[LZ09\]](#page-16-9)) which implies that a nonzero degree $2d-1$ noncommutative polynomial evaluated on $d \times d$ matrices will be invertible with high probability. However, this does not yield an efficient randomized identity testing algorithm for rational expressions of inversion height one. Because that seems to require testing correctness of expressions of inversion height two which is a question left open in their paper [\[HW14,](#page-16-0) Section 9].

The Free Group Algebra

This motivates the study of black-box identity testing for rational expressions in the *free group algebra* $\mathbb{F}\langle X, X^{-1}\rangle$.

We consider expressions in the free group algebra $\mathbb{F}\langle X, X^{-1}\rangle$, where $(X, X^{-1})^*$ denotes the free group generated by the n generators $X =$ ${x_1, x_2, \ldots, x_n}$ and their inverses

$$
X^{-1} = \{x_1^{-1}, x_2^{-1}, \dots, x_n^{-1}\}.
$$

Elements of the free group $(X, X^{-1})^*$ are words in X, X^{-1} . The only relations satisfied by the generators is $x_i x_i^{-1} = x_i^{-1} x_i = 1$ for all *i*. Thus, the elements in

the free group $(X, X^{-1})^*$ are the *reduced words* which are words to which the above relations are not applicable.

The elements of the *free group algebra* $\mathbb{F}\langle X, X^{-1} \rangle$ are $\mathbb{F}\text{-linear combinations}$ of the form

$$
f = \sum_{w} \alpha_w w, \ \alpha_w \in \mathbb{F},
$$

where each $w \in (X, X^{-1})^*$ is a reduced word. The *degree* of the expression f is defined as the maximum length of a word w such that $\alpha_w \neq 0$. The expression f is said to have *sparsity* s if there are s many reduced words w such that $\alpha_w \neq 0$ in f. We also use the notation $[w]$ to denote the coefficient α_w of the reduced word w in the expression f .

The free noncommutative ring $\mathbb{F}\langle X\rangle$ is a subalgebra of $\mathbb{F}\langle X, X^{-1}\rangle$. Clearly, the elements of $\mathbb{F}\langle X, X^{-1}\rangle$ are a special case of rational expressions of *inversion* height one. I.e., we note that:

Proposition 1. $\mathbb{F}\langle X, X^{-1}\rangle \subset \cup_{d>0} \mathcal{E}_{d,1}.$

Note that the rational expressions in $\mathbb{F}\langle X, X^{-1}\rangle$ allows inverses only of the variables x_i , whereas the *free skew field* $\mathbb{F}\langle X \rangle$ contains all possible rational expressions (with inverses at any nested level).

Our results

The main goal of the current paper is to obtain black-box identity tests for rational expressions in the free group algebra $\mathbb{F}\langle X, X^{-1}\rangle$.

Our first result is a generalization of the Amitsur-Levitzki theorem [\[AL50\]](#page-15-0) to $\mathbb{F}\langle X, X^{-1}\rangle$. Let A be an associative algebra with identity over F. An expression $f \in \mathbb{F}\langle X, X^{-1}\rangle$ is an *identity* for A if

$$
f(a_1,\ldots,a_n)=0
$$

for all $a_i \in A$ such that a_i^{-1} is defined for each $i \in [n]$.

Theorem 1. Let \mathbb{F} be any field of characteristic zero and $f \in \mathbb{F}\langle X, X^{-1} \rangle$ be a nonzero expression of degree d. Then f is not an identity for the matrix algebra $\mathbb{M}_{2d}(\mathbb{F})$.

The following corollary is immediate.

Corollary 1 (Black-box identity testing for circuits in free group algebra). There is a black-box randomized $poly(n, d)$ identity test for degree d expressions in $\mathbb{F}\langle X, X^{-1}\rangle$.

If the black-box contains a sparse expression, we show efficient deterministic algorithms for identity testing and interpolation algorithm.

Theorem 2 (Black-box identity testing and reconstruction for sparse expressions in free group algebra). Let $\mathbb F$ be any field of characteristic zero and f is an expression in $\mathbb{F}\langle X, X^{-1}\rangle$ of degree d and sparsity s given as black-box. Then we can reconstruct f in deterministic $poly(n, d, s)$ time with matrix-valued queries to the black-box.

Our next result is another generalization of the Amitsur-Levitzki theorem [\[AL50\]](#page-15-0) extending a result of [\[AJMR17\]](#page-15-2) to free group algebras. We show that a nonzero expression $f \in \mathbb{F}\langle X, X^{-1} \rangle$ of degree D and sparsity s does not vanish on $O(\log s)$ dimensional matrices. It yields a randomized polynomial-time identity test if the black-box contains an expression f of exponential degree and exponential sparsity.

Theorem 3. Let \mathbb{F} be any field of characteristic zero. Then, a degree-D expression $f \in \mathbb{F}\langle X, X^{-1} \rangle$ of sparsity s is not an identity for the matrix algebra $\mathbb{M}_k(\mathbb{F})$ for $k = O(\log s)$.

Corollary 2 (Black-box identity testing for expoential sparse expressions with exponential degree in free group algebra). Given a degree-D expression $f \in$ $\mathbb{F}\langle X, X^{-1}\rangle$ of sparsity s as black-box, we can check whether f is identically zero or not in randomized $\text{poly}(n, \log D, \log s)$ time.

Remark 2. We state our results for fields of characteristic zero only for simplicity. However, by suitable modifications, we can extend our results for fields of positive characteristic.

Organization

The paper is organized as follows. In Section [2,](#page-4-0) we prove Theorem [1,](#page-3-0) Corollary [1,](#page-3-1) and Theorem [2.](#page-3-2) In Section [3,](#page-9-0) we prove Theorem [3](#page-4-1) and Corollary [2.](#page-4-2) Finally, in Section [4,](#page-14-0) we discuss suitable modifications to extend our results over finite fields.

2 A Generalization of Amitsur-Levitzki Theorem for Free Group Algebra

The main idea in our proof is to efficiently encode expressions in $\mathbb{F}\langle X, X^{-1}\rangle$ as polynomials in a suitable commutative ring preserving the identity. Let $\mathbb{F}[Y, Z]$ denote the commutative ring $\mathbb{F}[y_{ij}, z_{ij}]_{i \in [n], j \in [d]}$ for $n, d \in \mathbb{N}$, where $Y = \{y_{ij} \mid i \in [n], j \in [d]\}\$ and $Z = \{z_{ij} \mid i \in [n], j \in [d]\}.$

Definition 2. Define a map $\varphi : \mathbb{F}\langle X, X^{-1} \rangle \to \mathbb{F}[Y, Z]$ to be a map such that φ is identity on \mathbb{F} , and for each reduced word $w = x_{i_1}^{b_1}$ $_{i_{1}}^{b_{1}}x_{i_{2}}^{b_{2}}$ $\frac{b_2}{i_2}\cdots x_{i_d}^{b_d}$ $\frac{v_d}{i_d}$,

$$
\varphi(x_{i_1}^{b_1} x_{i_2}^{b_2} \cdots x_{i_d}^{b_d}) = \prod_{j=1}^d (\mathbb{1}_{[b_j=1]} \cdot y_{i_j j} + \mathbb{1}_{[b_j=-1]} \cdot z_{i_j j}),
$$

where $\mathbb{1}_{[b_j=b]}=1$ if $b_j=b$ and $\mathbb{1}_{[b_j=b]}=0$ otherwise.

By linearity the map φ is defined on all expressions in $\mathbb{F}\langle X, X^{-1}\rangle$. We observe the following properties of φ .

1. The map φ is injective on the reduced words $(X, X^{-1})^*$. I.e., it maps each reduced word $w \in (X, X^{-1})^*$ to a unique monomial over the commuting variables $Y \cup Z$.

- 2. Consequently, φ is identity preserving. I.e., an expression f in $\mathbb{F}\langle X, X^{-1}\rangle$ is identically zero if and only if its image $\varphi(f)$ is the zero polynomial in $\mathbb{F}[Y,Z].$
- 3. φ preserves the sparsity of the expression. I.e., f in $\mathbb{F}\langle X, X^{-1}\rangle$ is s-sparse iff $\varphi(f)$ in $\mathbb{F}[Y,Z]$ is s-sparse.
- 4. Given the image $\varphi(f) \in \mathbb{F}[Y, Z]$ in its sparse description (i.e., as a linear combination of monomials), we can efficiently recover the sparse description of $f \in \mathbb{F}\langle X, X^{-1} \rangle$.

Given polynomials $f, f' \in \mathbb{F}[Y, Z]$, we say f and f' are weakly equivalent, if for each monomial m, $[m]f = 0$ if and only if $[m]f' = 0$, where $[m]f$ denotes the coefficient of monomial m in f .

Given a black-box expression f in $\mathbb{F}\langle X, X^{-1}\rangle$, we show how to evaluate it on suitable matrices and obtain a polynomial in $\mathbb{F}[Y, Z]$ that is *weakly equivalent* to $\varphi(f)$ as a specific entry of the resulting matrix. The matrix substitutions are based on automata constructions. Similar ideas have been used earlier to design PIT algorithms for noncommutative polynomials [\[AMS10\]](#page-15-4). However, since we are dealing with rational expressions, some difficulties arise. The matrix substitutions for the variables x_1, \ldots, x_n are obtained as the corresponding transition matrices M_i of the automaton. The matrix substitution for x_i^{-1} will be M_i^{-1} . Therefore, we need to ensure that the transition matrices M_i are invertible and sufficiently structured to be useful for the identity testing.

We first illustrate our construction for an example degree-2 expression $f =$ $x_1x_2^{-1} + x_2x_1^{-1}$, where $X = \{x_1, x_2\}$.

The basic "building block" for the transition matrix M_i is the 2×2 block matrix

$$
\begin{bmatrix} 0 & y_{ij} \\ \frac{1}{z_{ij}} & 0 \end{bmatrix},
$$

$$
\begin{bmatrix} 0 & z_{ij} \\ \frac{1}{y_{ij}} & 0 \end{bmatrix}.
$$

whose inverse is

When the 2×2 block is the jth diagonal block in M_i , the corresponding automaton will go from state $2j-1$ to state $2j$ replacing x_i by y_{ij} (or if x_i^{-1} occurs, it will replace it by z_{ij}).

We will keep the transition matrix M_i for x_i a block diagonal matrix with such 2×2 invertible blocks as the principal minors along the diagonal. In order to ensure this we introduce two new variables $W = \{w_1, w_2\}$ and substitute x_i by the word $w_i x_i w_i$ in the expression. This will ensure that we do not have two consecutive x_i in the resulting reduced words. In fact, between two X variables (or their inverses) we will have inserted exactly two W variables (or their inverses). Now, we define M_i for the above example as

$$
M_i = \begin{bmatrix} 0 & y_{i1} & 0 & 0 \\ \frac{1}{z_{i1}} & 0 & 0 & 0 \\ 0 & 0 & 0 & y_{i2} \\ 0 & 0 & \frac{1}{z_{i2}} & 0 \end{bmatrix}, \qquad M_i^{-1} = \begin{bmatrix} 0 & z_{i1} & 0 & 0 \\ \frac{1}{y_{i1}} & 0 & 0 & 0 \\ 0 & 0 & 0 & z_{i2} \\ 0 & 0 & \frac{1}{y_{i2}} & 0 \end{bmatrix}.
$$

The corresponding transitions of the automaton is shown in Figure [1.](#page-6-0)

Figure 1: The transition diagram of the automaton for x variables

We now describe the transition matrices N_i for w_i . The matrix N_i is also a 4×4 block diagonal matrix. There are three blocks along the diagonal. The first and third are 1×1 blocks of the identity. The second one is a 2×2 block for w_i -transitions from state q_2 to state q_3 . It ensures that for any subword $w_1^{b_1} w_2^{b_2}, b_i \in \{1, -1\}$, in the resulting product matrix $N_1^{b_1} N_2^{b_2}$ the $(1, 2)^{th}$ entry of the 2×2 block is nonzero. The corresponding transitions of the automaton is depicted in Figure [2.](#page-6-1)

Figure 2: The transition diagram of the automaton for w variables

$$
N_i=\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & i & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix},\quad N_i^{-1}=\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & -i & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix},\quad N_i^{b_1}N_j^{b_2}=\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & b_1i+b_2j & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.
$$

Hence, evaluating $f(N_1M_1N_1, N_2M_2N_2)$ we obtain (a polynomial weakly equivalent to) $\varphi(f)$ at the $(1,4)^{th}$ entry. The complete automaton is depicted in figure [3.](#page-7-0)

Figure 3: The transition diagram of the automaton

We now explain the general construction. For $f \in \mathbb{F}\langle X, X^{-1}\rangle$ let $H_{\ell}(f)$ denote the degree- ℓ homogeneous part of f. We will denote by $\varphi(\widehat{H_{\ell}(f)})$ an arbitrary polynomial in $\mathbb{F}[Y, Z]$ weakly equivalent to $\varphi(H_\ell(f)).$

Lemma 1. Let $f \in \mathbb{F}\langle X, X^{-1} \rangle$ be a nonzero expression of degree d. There is an n-tuple of $2d \times 2d$ matrices (M_1, M_2, \ldots, M_n) whose entries are either scalars, or variables $u \in Y \cup Z$, or their inverses $1/u$, such that

$$
(f(M_1,\ldots,M_n))_{1,2d}=\widehat{\varphi(H_d(f))}.
$$

Furthermore, for each degree-d reduced word of $m = x_{i_1}^{b_1}$ $_{i_{1}}^{b_{1}}x_{i_{2}}^{b_{2}}$ $\frac{b_2}{i_2}\cdots x_{i_d}^{b_d}$ $_{i_d}^{b_d}$ in $\mathbb{F}\langle X, X^{-1}\rangle$,

$$
[\varphi(m)]\varphi(\widehat{H_d(f)}) = [m]f \cdot \prod_{j=1}^{d-1} (b_j \cdot i_j + b_{j+1} \cdot i_{j+1}).
$$
\n(1)

Proof. Let e_{ij} , for $i, j \in [k]$, be the $(i, j)^{th}$ elementary matrix in $\mathbb{M}_k(\mathbb{F})$: its (i, j) th entry is 1 and other entries are 0.

We now define the transition matrices of the NFA for variables $\{w_i: 1 \leq i \leq n\}$ n} and $\{x_i : 1 \le i \le n\}$. For each $i \in [n]$, define 2×2 matrix $N'_i = e_{11} + e_{22} + i \cdot e_{12}$. Now N_i is a $2d \times 2d$ matrix defined as the block diagonal matrix,

$$
N_i' = \begin{bmatrix} 1 & i \\ 0 & 1 \end{bmatrix}, \qquad N_i = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & N_i' & 0 & \dots & 0 & 0 \\ 0 & 0 & N_i' & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & N_i' & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 \end{bmatrix}.
$$

$$
N_i'^{-1} = \begin{bmatrix} 1 & -i \\ 0 & 1 \end{bmatrix}, \qquad N_i^{-1} = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & N_i'^{-1} & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & N_i'^{-1} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & N_i'^{-1} & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 \end{bmatrix}.
$$

Figure 4: The transition diagram of the automaton

Each $M_i, 1 \leq i \leq n$ is the $2d \times 2d$ block diagonal matrix where each $2 \times 2d$ block M'_{ij} , $1 \leq j \leq d$ is a 2×2 matrix defined as $M'_{i,j} = y_{ij} \cdot e_{12} + \frac{1}{z_{ij}} \cdot e_{21}$. Their inverses have a similar structure.

$$
M'_{i,p} = \begin{bmatrix} 0 & y_{ip} \\ \frac{1}{z_{ip}} & 0 \end{bmatrix}, \qquad M_i = \begin{bmatrix} M'_{i,1} & 0 & 0 & \dots & 0 \\ 0 & M'_{i,2} & 0 & \dots & 0 \\ 0 & 0 & M'_{i,3} & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & M'_{i,d} \end{bmatrix}.
$$

$$
M'_{i,p}^{-1} = \begin{bmatrix} 0 & z_{ip} \\ \frac{1}{y_{ip}} & 0 \end{bmatrix}, \qquad M_i^{-1} = \begin{bmatrix} M'_{i,1}^{-1} & 0 & 0 & \dots & 0 \\ 0 & M'_{i,2}^{-1} & 0 & \dots & 0 \\ 0 & 0 & M'_{i,3}^{-1} & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & M'_{i,d}^{-1} \end{bmatrix}.
$$

The corresponding NFA is depicted in Figure [4.](#page-8-0) We substitute each x_{i_j} by the $2d \times 2d$ matrix $N_{i_j} N_{i_j} N_{i_j}$. Each $x_{i_j}^{-1}$ is substituted by its inverse matrix $N_{i_j}^{-1} M_{i_j}^{-1} N_{i_j}^{-1}.$

Correctness.

Consider a degree-d reduced word $m = x_{i_1}^{b_1}$ $\frac{b_1}{i_1} x_{i_2}^{b_2}$ $\frac{b_2}{i_2}\cdots x_{i_d}^{b_d}$ $\frac{v_d}{i_d}$.

Following the automaton construction of Figure [4,](#page-8-0) $x_i^{b_i}$ occurring at position j is substituted by $([\mathbb{1}_{[b_i=1]}y_{ij}+\mathbb{1}_{[b_i=-1]}z_{ij})$. Moreover, for each position $j \in [d-1]$, the adjacent pair $x_{i,j}^{b_j}$ $_{i_{j}}^{b_{j}}x_{i_{j+1}}^{b_{j+1}}$ $i_{j+1}^{o_{j+1}}$ produces a scalar factor $(b_j \cdot i_j + b_{j+1} \cdot i_{j+1})$ due to the product $N_{i,j}^{b_j}$ $\frac{b_j}{i_j} N^{b_{j+1}}_{i_{j+1}}$ $i_{j+1}^{o_{j+1}}$. Consequently, it follows that

$$
(m(M_1,\ldots,M_n))_{1,2d}=\prod_{j=1}^{d-1}(b_j\cdot i_j+b_{j+1}\cdot i_{j+1})\prod_{j=1}^d([b_j=1]y_{i_jj}+[b_j=-1]z_{i_jj}).
$$

As φ is a linear map, the lemma follows.

 \Box

2.1 Black-box identity testing for circuits in free group algebra

Theorem [1](#page-7-1) follows easily from Lemma [1.](#page-7-1) Lemma 1 says that if $f \in$ $\mathbb{F}\langle X, X^{-1}\rangle$ is nonzero of degree d then the $(1, 2d)$ entry of the matrix $p(N_1M_1N_1,\ldots,N_nM_nN_n)$ is a nonzero polynomial in $\mathbb{F}[Y,Z]$. Hence f can not be an identity for $M_{2d}(\mathbb{F})$.

It also immediately gives an identity testing algorithm. We can randomly substitute for the variables and apply the Schwartz-Zippel-Demillo-Lipton Theorem [\[Sch80,](#page-17-1) [Zip79,](#page-17-2) [DL78\]](#page-16-10). This completes the proof of the Corollary [1.](#page-3-1)

2.2 Reconstruction of sparse expressions in free group algebra

If the black-box contains an s-sparse expression in $\mathbb{F}\langle X, X^{-1}\rangle$, we give a $poly(s, n, d)$ deterministic interpolation algorithm (which also gives a deterministic identity testing for such expressions). We use a result of Klivans-Spielman [\[KS01,](#page-16-11) Theorem11] that constructs a test set in deterministic polynomial time for sparse commutative polynomials, which is used for the interpolation algorithm.

Proof of Theorem [2](#page-3-2)

Let the black-box expression f be s-sparse of degree d . By Lemma [1,](#page-7-1) a polynomial $\widehat{\varphi(H_d(p))}$ in $\mathbb{F}[Y, Z]$ is obtained at the $(1, 2d)^{th}$ entry of the matrix $f(M_1, \ldots, M_n)$, where $M_i \in M_{2d}(\mathbb{F}[Y, Z])$ is as defined in Lemma [1.](#page-7-1) By Defi-nition [2,](#page-4-3) $\varphi(f) \in \mathbb{F}[Y, Z]$ is s-sparse and has 2nd variables. Let $\mathcal{H}_{2nd,d,s}$ be the corresponding test set from [\[KS01\]](#page-16-11) to interpolate a polynomial of degree d and ssparse over 2nd variables. Querying the black-box on $M_1(\vec{h}), M_2(\vec{h}), \ldots, M_n(\vec{h})$ for each $\vec{h} \in \mathcal{H}_{2nd,d,s}$ we can interpolate the commutative polynomial $\varphi(\overline{H}_d(\overline{f}))$ and obtain an expression for $\widehat{\varphi(H_d(f))} = \sum_{t=1}^s c_{m_t} m_t$ as a sum of monomials.

We now need to adjust the extra scalar factors in $\varphi(H_d(f))$ to obtain $\varphi(H_d(f))$. We can perform this adjustment for each monomial as Lemma [1](#page-7-1) shows that the extra scalar factor for the word $m = x_{i_1}^{b_1}$ $_{i_{1}}^{b_{1}}x_{i_{2}}^{b_{2}}$ $\frac{b_2}{i_2}\cdots x_{i_\ell}^{b_\ell}$ $\frac{b_{\ell}}{i_{\ell}}$ is just $\alpha_m = \prod_{j=1}^{\ell-1} (b_j \cdot i_j + b_{j+1} \cdot i_{j+1}).$ So the algorithm constructs the expression $\varphi(\widehat{H_d(f)}) = \sum_{t=1}^s \frac{c_{m_t}}{\alpha_{m_t}}$ $\frac{c_{m_t}}{c_{m_t}} m_t$. We can remove the factors α_{m_t} for each monomial m_t and *invert* the map φ (using the 4th property of Definition [2\)](#page-4-3) on every monomial m_t to obtain $H_d(f)$ as a sum of degree d reduced words. This yields the expression for highest degree homogeneous component of f . We can repeat the above procedure on $f - H_d(f)$ and reconstruct the remaining homogeneous components of f . \Box

3 Black-box Identity Testing for Expressions of Exponential Degree and Exponential Sparsity

In this section, we prove a different generalization of Amitsur-Levitzki theorem [\[AL50\]](#page-15-0) for free group algebras, based on ideas from [\[AJMR17\]](#page-15-2). We show

that the dimension of the matrix algebra for which a nonzero input expression f does not vanish is logarithmic in the sparsity of f . It yields a randomized $poly(log D, log s, n)$ time identity testing algorithm when the black-box contains an expression of degree D and sparsity s.

We first recall the notion of *isolating index set* from [\[AJMR17\]](#page-15-2).

Definition 3. Let $\mathcal{M} \subseteq \{X, X^{-1}\}^D$ be a subset of reduced words of degree D. An index set $I \subseteq [D]$ is an isolating index set for M if there is a word $m \in \mathcal{M}$ such that for each $m' \in \mathcal{M} \setminus \{m\}$ there is an index $i \in I$ for which $m[i] \neq m'[i]$. I.e. no other word in $\mathcal M$ agrees with m on all positions in the index set I. We say m is an isolated word.

In the following lemma we show that M has an isolating index set of size $log|\mathcal{M}|$. The proof is identical to [\[AJMR17\]](#page-15-2). Nevertheless, we give the simple details for completeness because we deal with both variables and their inverses.

Lemma 2. [\[AJMR17\]](#page-15-2) Let $M \subseteq \{X, X^{-1}\}^D$ be reduced degree-D words. Then M has an isolating index set of size k which is bounded by $log|M|$.

Proof. The words $m \in \mathcal{M}$ are indexed, where $m[i]$ denotes the variable (or the inverse of a variable) in the i^{th} position of m. Let $i_1 \leq D$ be the first index such that not all words agree on the i_1^{th} position. Let

$$
S_j^+ = \{m : m[i_1] = x_j\}
$$

$$
S_j^- = \{m : m[i_1] = x_j^{-1}\}.
$$

For some j, $|S_j^+|$ or $|S_j^-|$ is of size at most $|\mathcal{M}|/2$. Let $S_{i_1}^b$ denote that subset, $b \in \{+, -\}.$ We replace M by $S_{i_1}^b$ and repeat the same argument for at most $log |\mathcal{M}|$ steps. Clearly, by this process, we identify a set of indices $I = \{i_1, \ldots, i'_k\}, k' \leq \log |\mathcal{M}|$ such that the set shrinks to a singleton set $\{m\}.$ Clearly, I is an isolating index set as witnessed by the *isolating word* m . \Box

Proof of Theorem [3](#page-4-1)

Let $k = 4(k'+1)$ where k' is the size of the isolating set I. As in Section [2,](#page-4-0) we substitute each x_i by $w_i x_i w_i$, where $w_i, i \in [n]$ are n new variables. The transition matrices for w_i and x_i are denoted by N_i and M_i respectively.

For $1 \leq i \leq n$, we define $k \times k$ matrix N_i as a block diagonal matrix of k many 4×4 matrices N'_i where $N'_i = I_4 + i(e_{12} + e_{34} + e_{32} + e_{14}).$

$$
N'_i = \begin{bmatrix} 1 & i & 0 & i \\ 0 & 1 & 0 & 0 \\ 0 & i & 1 & i \\ 0 & 0 & 0 & 1 \end{bmatrix}, \qquad N_i = \begin{bmatrix} N'_i & 0 & 0 & \dots & 0 \\ 0 & N'_i & 0 & \dots & 0 \\ 0 & 0 & N'_i & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & N'_i \end{bmatrix},
$$

$$
N_i'^{-1} = \begin{bmatrix} 1 & -i & 0 & -i \\ 0 & 1 & 0 & 0 \\ 0 & -i & 1 & -i \\ 0 & 0 & 0 & 1 \end{bmatrix}, \qquad N_i^{-1} = \begin{bmatrix} N_i'^{-1} & 0 & 0 & \dots & 0 \\ 0 & N_i'^{-1} & 0 & \dots & 0 \\ 0 & 0 & N_i'^{-1} & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & N_i'^{-1} \end{bmatrix}.
$$

Notice that

$$
N_i'^{b_1} N_j'^{b_2} = \begin{bmatrix} 1 & (b_1 i + b_2 j) & 0 & (b_1 i + b_2 j) \\ 0 & 1 & 0 & 0 \\ 0 & (b_1 i + b_2 j) & 1 & (b_1 i + b_2 j) \\ 0 & 0 & 0 & 1 \end{bmatrix}.
$$

We now define the $k \times k$ transition matrix M_i as a block diagonal matrix,

$$
M'_{i,j} = \begin{bmatrix} 0 & y_{ij} \\ \frac{1}{z_{ij}} & 0 \end{bmatrix}, \t M'_{\xi_i} = \begin{bmatrix} 0 & \xi_i \\ \frac{1}{\xi_i} & 0 \end{bmatrix},
$$

$$
M_i = \begin{bmatrix} 1 & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & M_{\xi_1} & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & M'_{i,1} & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & M_{\xi_2} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & M_{\xi_{k'+1}} & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 & 1 \end{bmatrix}
$$

.

These matrices can be seen as the transitions of a suitable NFA. We sketch the construction of this NFA.

Let $I = \{i_1, \ldots, i_{k'}\}$ be an isolating set such that $i_1 < \ldots < i_{k'}$. Intuitively, the NFA does one of two operations on each symbol (a variable or its inverse) of the input expression: a Skip or an Encode. In a Skip stage, the NFA deals with positions that are not part of the (guessed) isolating index set. In this stage, the NFA substitutes the w_i variables by suitable scalars (coming from the N'_i matrices) and x_i variables by block variables $\{\xi_1, \ldots, \xi_{k'+1}\}.$ The NFA nondeterministically decides whether the Skip stage is over and it enters the Encode stage for a guessed index of the isolating set. It substitutes x_i and x_i^{-1} variables by y_{ij} and z_{ij} respectively. Fig. [5](#page-11-0) summarizes the action of the NFA.

Figure 5: The transition diagram of the automaton

Define \hat{f} in $\mathbb{F}(Y, Z, \overline{\xi})$ to be rational function we obtain at the $(1, k)^{th2}$ $(1, k)^{th2}$ $(1, k)^{th2}$ entry by evaluating the expression $f(N_1M_1N_1, \ldots, N_nM_nN_n)$. Notice that, the isolating word m of degree D will be of following form $m =$ $W_1x_{i_1}^{b_{i_1}}W_2x_{i_2}^{b_{i_2}}$ ${}_{i_2}^{b_{i_2}} \cdots W_k' x_{i'_k}^{b_{i'_k}} W_{k'+1}$ where each subword $W_j = x_{j_1}^{b_1}$ $_{j_1}^{b_1}x_{j_2}^{b_2}$ $\frac{b_2}{j_2}\cdots\frac{b_{\ell_j}}{j_{\ell_j}}$ $j_{\ell_j}^{i_j}$ is of length $\ell_j \geq 0$, where some of the W_j could be the empty word as well.

We refer to an NFA transition $q_i \rightarrow q_j$ as a *forward edge* if $i < j$ and a backward edge if $i > j$. We classify the backward edges in three categories based on the substitution on the edge-label. We say, a backward edge is of type A if a variable is substituted by a scalar value; a backward edge is of type B if a variable is substituted by $\frac{1}{\xi_j}$ for some j; a backward edge is of type C if a variable is substituted by $\frac{1}{y_{ij}}$ or $\frac{1}{z_{ij}}$ for some *i*, *j*.

Consider a walk of the NFA on an input word m that reaches state k using only type A backward edges. In that case, m is substituted by $\alpha \cdot \hat{m}$ where \hat{m} is a monomial over ${Y, Z, \xi}$ of same degree,

$$
\hat{m} = \prod_{j=1}^{k'+1} \xi_j^{\ell_j} \cdot \prod_{j=1}^{k'} ([b_{i_j} = 1]y_{i_j j} + [b_{i_j} = -1]z_{i_j j}).
$$

and α is some nonzero constant obtained as a product of $[m]$ with the scalars obtained as substitutions from the edges involving the w_i variables in the *Skip* stages. Indeed, as we can see from the entries of product matrices $N_i^{b_1} \cdot N_j^{b_2}$, where $b_1, b_2 \in \{-1, 1\}$, the scalar α is a product of $[m]$ with terms of the form $b_1i + b_2j$, for $i \neq j$, each of which is nonzero for any reduced word.

Figure 6: The transition diagram of the automaton at *Encode* stage

Claim 1.

$$
[\hat{m}]\hat{f} \neq 0 \text{ iff } [m]f \neq 0.
$$

Proof. It suffices to show that for any word $m' \neq m$, where m' has degree $\leq D$, no walks of the NFA accepting m' generate \hat{m} after substitution. We now argue that no other walks in the NFA can generate \hat{m} . For a computation path J, the monomial m_J in f has two parts, let us call it skip_J and encode_J where $skip_j$ is a monomial over $\{\xi_1, \ldots, \xi_{k'+1}\}\$ and $encode_j$ is a monomial over ${y_{i,j, z_{i,j}}}_{i\in[n],j\in[k']}$. If the computation path J (which is different from the computation path described above for \hat{m}) uses only type A backward edges, then necessarily $m_J \neq \hat{m}$ from the definition of *isolating index set*. This argument is analogous to the argument given in [\[AJMR17\]](#page-15-2).

²Recall that $k = 4(k'+1)$ where k' is the size of an isolating set.

Figure 7: The transition diagram of the automaton at *Skip* stage

Now consider a walk J which involves backward edges of other types. Let us first consider those walks that take backward edges only of type A and type B. Such a walk still produces a monomial over $\{y_{i,j}, z_{i,j}\}_{i \in [n], j \in [k']}$ and $\{\xi_i\}_{1 \leq i \leq k'+1}$ because division only by ξ_i variables occur in the resulting expression. Since \hat{m} is of highest degree, the total degree of these monomials is strictly lesser than degree of \hat{m} . For those walks that take at least one backward edge of type C, a rational expression in $\{y_{i,j}, z_{i,j}\}_{i \in [n], j \in [k']}$ and $\{\xi_i\}_{1 \le i \le k'+1}$ is produced (as there is division by y_{ij} or z_{ij} variables). As the sum of the degree of the numerator and degree of the numerator is bounded by the total degree, the degree of the numerator is smaller than degree of \hat{m} .

Thus the $(1,k)^{th}$ entry of the output matrix is of the form $\sum_{i=1}^{N_1} c_i m_i +$ $\sum_{j=1}^{N_2} r_j$ where $\{m_1, \ldots, m_{N_1}\}\$ are monomials arising from different walks (w.l.o.g. assume that $m_1 = \hat{m}$) and $\{r_1, \ldots, r_{N_2}\}\$ are the rational expressions from the other walks (due to the backward edges of type C). Note that, denominator in each r_j is a monomial over Y, Z of degree at most D. Let $L = \prod_{i=1}^n \prod_{j=1}^{k'} y_{i,j}^D \cdot z_{i,j}^D$. Now, we have,

$$
\sum_{i=1}^{N_1} c_i m_i + \sum_{j=1}^{N_2} r_j = \frac{1}{L} \cdot \left(\sum_{i=1}^{N_1} c_i m_i L + \sum_{j=1}^{N_2} p_j \right).
$$

Since $\hat{m} L \neq m_i L$ for any $i \in \{2, ..., N_1\}$ and degree of each $p_j <$ degree of mL for any $j \in \{1, ..., N_2\}$, the numerator of the final expression is a nonzero polynomial in $\mathbb{F}[Y, Z, \overline{\xi}].$ \Box

The above proof shows that the matrix $f(N_1M_1N_1,\ldots,N_nM_nN_n)$ is nonzero with rational entries in $\mathbb{F}[Y, Z, \overline{\xi}]$. Each entry is a linear combination of terms of the form m_1/m_2 , where m_1 and m_2 are monomials in $Y \cup Z \cup \{\xi_1, \ldots, \xi_{k'+1}\}\$ of degree bounded by D. This completes the proof.

To get an identity testing algorithm, we can do random substitutions.The matrix dimension is $\log s$ and the overall running time of the algorithm is $poly(n, \log s, \log D)$. This also proves Corollary [2.](#page-4-2) \Box

Remark 3. For algorithmic purposes, we note that Theorem [1](#page-3-0) is sometimes preferable to Theorem [3.](#page-4-1) For instance, the encoding used in Theorem [3](#page-4-1) does not preserve the sparsity of the polynomial as required in the sparse reconstruction result (Theorem [2\)](#page-3-2).

4 Adaptation for Fields of Positive Characteristic

Let $\mathbb F$ be any finite field of characteristic p. We need to ensure that for each word m in the free group algebra, the scalar α_m (see Equation [1\)](#page-7-2) produced by the automaton described in Section [2](#page-4-0) is not zero in \mathbb{F} . Recall that, reading $w_i^{b_i} w_j^{b_j}$ j for two consecutive positions, the automaton produces a scalar $(b_i \cdot i + b_j \cdot j)$ where $b_i, b_j \in \{-1, +1\}$. Moreover, this is the only way the automaton produces a scalar and for each m , α_m is a product of such terms. Hence, all we need to ensure is that for each pair $i, j \in [n]$, $(b_i \cdot i + b_j \cdot j) \neq 0$. Similarly, it ensures that the scalar produced by the automaton described in Section [3](#page-9-0) is non-zero.

We note that, if p is more than $2n$ then each term $(b_i \cdot i + b_j \cdot j) \neq 0 \pmod{p}$ where $b_i, b_j \in \{-1, +1\}$ and $i, j \in [n]$. This results in a dependence on the characteristic of the base field for the analogous statements of Theorems [1,](#page-3-0) [3](#page-4-1) over finite field. Additionally, for Theorem [1,](#page-3-0) the $(1, 2d)^{th}$ entry of the output matrix is a polynomial of degree d , and for Theorem [3,](#page-4-1) the degrees of the numerator polynomials in the rational expression of the output matrix is bounded by some scalar multiple of $nD \log s$. This lower bounds the size of the fields in the application. We summarize the above discussion in the following.

Observation 1. We can obtain results analogous to Theorem [1](#page-3-0) and Theorem [3](#page-4-1) over finite fields of characteristic more than $2n$ and sizes at least $d+1$ or $\Omega(nD \log s)$ respectively.

However, the algorithms presented in Theorem [2](#page-3-2) and Corollaries [1,](#page-3-1) [2](#page-4-2) can be modified to work for finite fields of any characteristic. To this end, we first notice the following simple fact.

Proposition 2. Let \mathbb{F} be a finite field of characteristic $p \leq 2n$. In We can find elements $\alpha_1, \alpha_2, \ldots, \alpha_n$ from a suitable (deterministically constructed) small extension field \mathbb{F}' of \mathbb{F} in deterministic poly(n) time, such that for any $b_i \in \{-1, 1\}, 1 \leq i \leq n$ we have

For each
$$
1 \leq i < j \leq n
$$
, $b_i \alpha_i + b_j \alpha_j \neq 0$.

Let $\alpha_1, \alpha_2, \ldots, \alpha_n \in \mathbb{F}'$ as given by the above proposition. We modify the matrix N' _i in the proof of Theorem [2](#page-3-2) and Corollary [1](#page-3-1) as

$$
N_i' = \begin{bmatrix} 1 & \alpha_i \\ 0 & 1 \end{bmatrix},
$$

and in Corollary [2](#page-4-2) we modify N' _i as

$$
N'_{i} = \begin{bmatrix} 1 & \alpha_{i} & 0 & \alpha_{i} \\ 0 & 1 & 0 & 0 \\ 0 & \alpha_{i} & 1 & \alpha_{i} \\ 0 & 0 & 0 & 1 \end{bmatrix}.
$$

For each pair $i, j \in [n]$, $(b_i \cdot \alpha_i + b_j \cdot \alpha_j) \neq 0$ by Proposition [2.](#page-14-1) Thus, for each word m, the scalar α_m produced by the automata are nonzero in the extension field \mathbb{F}' as well. Furthermore, the test set of [\[KS01\]](#page-16-11) works for all fields. Hence Theorem [2](#page-3-2) holds for all finite fields too. To obtain Corollaries [1](#page-3-1) and [2,](#page-4-2) we need to do the random substitution from suitable small degree extension fields and use Schwartz-Zippel-Demillo-Lipton Theorem [\[Sch80,](#page-17-1) [Zip79,](#page-17-2) [DL78\]](#page-16-10). In summary, our algorithms in the paper can be adapted to work over all fields. *Proof of Proposition [2.](#page-14-1)* Define polynomial $g \in \mathbb{F}[x_1, x_2, \ldots, x_n]$ as

$$
g(x_1, x_2, \dots, x_n) = \prod_{1 \leq i < j \leq n} (x_i + x_j) \cdot (x_i - x_j).
$$

We substitute y^i for $x_i, 1 \leq i \leq n$. Then $g(y, y^2, \ldots, y^n) = G(y) \in \mathbb{F}[y]$ is a univariate polynomial of degree at most $2n^3$. Using standard techniques, in deterministic polynomial time we can construct an extension field \mathbb{F}' of \mathbb{F} such that $|\mathbb{F}'|$ is of $poly(n) \geq 2n^3 + 1$ size. We can find an element $\alpha \in \mathbb{F}'$ such that $G(\alpha) \neq 0$ and set $\alpha_i = \alpha^i, 1 \leq i \leq n$. \Box

References

- [AJMR17] Vikraman Arvind, Pushkar S. Joglekar, Partha Mukhopadhyay, and S. Raja. Randomized polynomial time identity testing for noncommutative circuits. In Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2017, Montreal, QC, Canada, June 19-23, 2017, pages 831–841, 2017.
- [AL50] A. S. Amitsur and J. Levitzki. Minimal identities for algebras. Proceedings of the American Mathematical Society, 1(4):449–463, 1950.
- [AMS10] Vikraman Arvind, Partha Mukhopadhyay, and Srikanth Srinivasan. New results on noncommutative and commutative polynomial identity testing. Computational Complexity, 19(4):521–558, 2010.
- [Ber76] George M Bergman. Rational relations and rational identities in division rings. Journal of Algebra, $43(1):252 - 266$, 1976.
- [BW05] Andrej Bogdanov and Hoeteck Wee. More on noncommutative polynomial identity testing. In 20th Annual IEEE Conference on Computational Complexity (CCC 2005), 11-15 June 2005, San Jose, CA, USA, pages 92–99, 2005.
- [DL78] Richard A. Demillo and Richard J. Lipton. A probabilistic remark on algebraic program testing. Information Processing Letters, $7(4):193 - 195, 1978.$
- [DM18] Harm Derksen and Visu Makam. Algorithms for orbit closure separation for invariants and semi-invariants of matrices. CoRR, abs/1801.02043, 2018.
- [FS12] Michael Forbes and Amir Shpilka. Quasipolynomial-time identity testing of non-commutative and read-once oblivious algebraic branching programs. Foundations of Computer Science, 1975., 16th Annual Symposium on, 09 2012.
- [GGOW16] Ankit Garg, Leonid Gurvits, Rafael Mendes de Oliveira, and Avi Wigderson. A deterministic polynomial time algorithm for noncommutative rational identity testing. 2016 IEEE 57th Annual Symposium on Foundations of Computer Science (FOCS), pages 109–117, 2016.
- [HW14] Pavel Hrubes and Avi Wigderson. Non-commutative arithmetic circuits with division. pages 49–66, 01 2014.
- [HY11] Pavel Hrubes and Amir Yehudayoff. Arithmetic complexity in ring extensions. Theory of Computing, 7:119–129, 2011.
- [IQS18] Gábor Ivanyos, Youming Qiao, and K. V. Subrahmanyam. Constructive non-commutative rank computation is in deterministic polynomial time. computational complexity, 27(4):561–593, Dec 2018.
- [KS01] Adam R. Klivans and Daniel Spielman. Randomness efficient identity testing of multivariate polynomials. In Proceedings of the Thirty-third Annual ACM Symposium on Theory of Computing, STOC '01, pages 216–223, New York, NY, USA, 2001. ACM.
- [LZ09] Tsiu-Kwen Lee and Yiqiang Zhou. Right ideals generated by an idempotent of finite rank. *Linear Algebra and its Applications*, 431:2118–2126, 11 2009.
- [Nis91] Noam Nisan. Lower bounds for non-commutative computation (extended abstract). In Proceedings of the 23rd Annual ACM Symposium on Theory of Computing, May 5-8, 1991, New Orleans, Louisiana, USA, pages 410–418, 1991.
- [Row80] Louis Halle Rowen. Polynomial identities in ring theory. Pure and Applied Mathematics. Academic Press, 1980.
- [RS05] Ran Raz and Amir Shpilka. Deterministic polynomial identity testing in non-commutative models. Computational Complexity, 14(1):1–19, 2005.
- [Sch80] Jacob T. Schwartz. Fast probabilistic algorithm for verification of polynomial identities. J. ACM., 27(4):701–717, 1980.
- [Str73] Volker Strassen. Vermeidung von divisionen. Journal fr die reine und angewandte Mathematik, 264:184–202, 1973.
- [Zip79] R. Zippel. Probabilistic algorithms for sparse polynomials. In Proc. of the Int. Sym. on Symbolic and Algebraic Computation, pages 216–226, 1979.