Fusion products of twisted modules in permutation orbifolds

Chongying Dong

Department of Mathematics, University of California, Santa Cruz, CA 95064 USA

Haisheng Li

Department of Mathematical Sciences, Rutgers University, Camden, NJ 08102 USA

Feng Xu

Department of Mathematics, University of California, Riverside, CA 92521 USA

Nina Yu

School of Mathematical Sciences, Xiamen University, Fujian 361005 CHINA

August 6, 2021

Abstract

Abstract: Let V be a vertex operator algebra, k a positive integer and σ a permutation automorphism of the vertex operator algebra $V^{\otimes k}$. In this paper, we determine the fusion product of any $V^{\otimes k}$ -module with any σ -twisted $V^{\otimes k}$ -module.

1 Introduction

This paper is about permutation orbifold vertex operator algebras. The theory of permutation orbifolds studies the representations of the tensor product vertex operator algebra $V^{\otimes k}$ with the natural action of the symmetric group S_k as an automorphism group, where V is a vertex operator algebra and k is a positive integer. The study of permutation orbifolds was initiated in [\[BHS\]](#page-35-0), where the twisted sectors (modules), genus one characters and their modular transformations, as well as the fusion rules for cyclic permutations for affine vertex operator algebras and the Virasoro vertex operator algebras were studied. The genus one characters and modular transformation properties of permutation orbifolds for a general rational conformal field theory were given in [\[Ba\]](#page-35-1).

The study on permutation orbifolds for general vertex operator algebras was started by Barron, Dong and Mason in [\[BDM\]](#page-35-2), where for a general vertex operator algebra V with any positive integer k , an intrinsic connection between twisted modules for tensor product vertex operator algebra $V^{\otimes k}$ with respect to permutation automorphisms and Vmodules was found. More specifically, let σ be a k-cycle which is viewed as an automorphism of $V^{\otimes k}$. Then, for any V-module $(W, Y_W(\cdot, z))$, a canonical σ -twisted $V^{\otimes k}$ -module structure on W was obtained, which was denoted by $T_{\sigma}(W)$. Furthermore, it was proved therein that this gives rise to an isomorphism of the categories of weak, admissible and ordinary V-modules and the categories of weak, admissible and ordinary σ -twisted $V^{\otimes k}$ -modules, respectively. The C_2 -cofiniteness of permutation orbifolds and general cyclic orbifolds was established later in [\[A1,](#page-35-3) [A2,](#page-35-4) [M1,](#page-37-0) [M2\]](#page-37-1). On the other hand, an equivalence of two constructions (see [\[FLM\]](#page-36-0), [\[Le\]](#page-36-1), [\[BDM\]](#page-35-2)) of twisted modules for permutation orbifolds of lattice vertex operator algebras was given in [\[BHL\]](#page-35-5). The permutation orbifolds of the lattice vertex operator algebras with $k = 2$ and $k = 3$ were extensively studied in [\[DXY1,](#page-36-2) [DXY2,](#page-36-3) [DXY3\]](#page-36-4). In particular, the fusion rules for the permutation orbifolds were determined. On the other hand, a study on permutation orbifolds in the context of conformal nets was given in [\[KLX\]](#page-36-5), where irreducible representations of the cyclic orbifolds were determined and fusion rules were given for $k = 2$.

Note that intertwining operator and fusion rule for modules were introduced in [\[FHL\]](#page-36-6), while a theory of tensor products for modules for vertex operator algebras was developed and has been extensively studied by Huang and Lepowsky (see [\[HL1\]](#page-36-7), [\[HL2\]](#page-36-8), [\[H1\]](#page-36-9), [\[H2\]](#page-36-10)). Intertwining operators among three twisted modules were studied by Xu (see [\[X\]](#page-37-2)) and a notion of tensor product for twisted modules was introduced in [\[DLM1\]](#page-35-6).

In this paper, we study fusion products of $V^{\otimes k}$ -modules with σ -twisted $V^{\otimes k}$ -modules for any $\sigma \in S_k$. More specifically, by using the explicit construction of the σ -twisted $V^{\otimes k}$ -modules due to [\[BDM\]](#page-35-2), we completely determine the fusion products. To achieve this goal, we study fusion products of twisted modules and obtain the associativity of the fusion product of twisted modules by using a theorem of [\[KO\]](#page-36-11) and [\[CKM\]](#page-35-7). We also generalize and use a result of [\[Li2\]](#page-36-12) and [\[Li3\]](#page-36-13) on an analogue of Hom-functor. More specifically, given any σ -twisted modules W_2 and W_3 for a vertex operator algebra V with a finite order automorphism σ , we introduce the space $\mathcal{H}(W_2, W_3)$ of generalized intertwining operators from W_2 to W_3 and prove that $\mathcal{H}(W_2, W_3)$ has a natural weak V-module structure. Furthermore, we prove that for any V-module W_1 , the space $\text{Hom}_V(W_1, \mathcal{H}(W_2, W_3))$ is naturally isomorphic to the space of intertwining operators of type $\binom{W_3}{W_1\ W_2}$. As one of the main results, we show that for any V-modules M, N, W, there is a canonical linear isomorphism from the space $I_V \binom{W}{M N}$ of intertwining operators of the indicated type to the space $I_{V^{\otimes k}}\binom{T_{\sigma}(W)}{M^1\,T_{\sigma}(N)}$ of intertwining operators, where $M^1 = M \otimes V^{\otimes (k-1)}$ is viewed as a $V^{\otimes k}$ -module.

Now, we describe the contents of this paper with more details. Let V be any vertex operator algebra and let k be a positive integer. Recall from [\[BDM\]](#page-35-2) that for $\sigma = (12 \cdots k) \in$ Aut $(V^{\otimes k})$, every σ -twisted $V^{\otimes k}$ -module is isomorphic to $T_{\sigma}(W)$ for some V-module W and $T_{\sigma}(W)$ is irreducible if and only if W is irreducible. On the other hand, from [\[FHL\]](#page-36-6), irreducible $V^{\otimes k}$ -modules are classified as $M_1 \otimes \cdots \otimes M_k$, where M_1, \ldots, M_k are irreducible V-modules. Our first goal is to determine the fusion product

$$
(M_1\otimes\cdots\otimes M_k)\boxtimes T_{\sigma}(W).
$$

We first consider the special case where $M_j = V$ for $2 \le j \le k$. While it is natural to start with a special case, actually the main motivation comes from the following observation

$$
(M_1 \otimes \cdots \otimes M_k) \simeq \boxtimes_{i=1}^k (V^{\otimes (i-1)} \otimes M_i \otimes V^{\otimes (k-i)}).
$$
\n(1.1)

The following is our first main result:

Theorem A. Let V be any vertex operator algebra and set $\sigma = (12 \cdots k) \in \text{Aut}(V^{\otimes k})$ with k a positive integer. Let M and N be any V-modules such that a tensor product $M \boxtimes N$ exists. Then

$$
\left(M\otimes V^{\otimes (k-1)}\right)\boxtimes_{V^{\otimes k}}T_{\sigma}(N)\simeq T_{\sigma}\left(M\boxtimes N\right).
$$

To prove this theorem, we generalize and use a result of [\[Li2\]](#page-36-12) and [\[Li3\]](#page-36-13) on an analogue of the classical Homfunctor. Let V be any vertex operator algebra with a finite order automorphism τ . Let W_2 and W_3 be τ -twisted V-modules. We first introduce the space $\mathcal{H}(W_2, W_3)$ of what are called generalized intertwining operators from W_2 to W_3 , where the definition of a generalized intertwining operator $\phi(x)$ from W_2 to W_3 captures the essential properties of $I(w, x)$ for an intertwining operator $I(\cdot, x)$ of type $\begin{pmatrix} W_3 \\ W W_2 \end{pmatrix}$ with W a V-module and with $w \in W$. More specifically, in addition to the lower truncation condition, $\phi(x)$ satisfies the conditions that $[L(-1), \phi(x)] = \frac{d}{dx}\phi(x)$ and that for every $v \in V$, there exists a nonnegative integer n such that

$$
(x_1 - x)^n Y_{W_3}(v, x_1) \phi(x) = (x_1 - x)^n \phi(x) Y_{W_2}(v, x_1).
$$

We prove that $\mathcal{H}(W_2, W_3)$ has a natural weak V-module structure and that for any V-module W_1 , the linear space $\text{Hom}_V(W_1, \mathcal{H}(W_2, W_3))$ is naturally isomorphic to the space of intertwining operators of type ${W_3 \choose W_1\ W_2}$. Coming back to vertex operator algebra $V^{\otimes k}$, let M,N,W be V -modules. By exploiting certain techniques from [\[BDM\]](#page-35-2) we obtain

a linear isomorphism between $I_{V^{\otimes k}}\binom{T_{\sigma}(W)}{M^T T_{\sigma}(N)}$ and $I_V\binom{W}{M N}$, where $M^1 = M \otimes V^{\otimes (k-1)}$. Furthermore, we use this natural isomorphism to obtain the fusion product isomorphism relation of Theorem A.

As it was indicated before, the main idea is to prove Theorem A first and then use the relation [\(1.1\)](#page-1-0) and the associativity of the tensor product to deal with the general case. Using Theorem A and a general result (Lemma [6.4\)](#page-32-0), we show

$$
\left(V^{\otimes (r-1)} \otimes M \otimes V^{\otimes (k-r)}\right) \boxtimes_{V^{\otimes k}} T_{\sigma}(N) \simeq \left(M \otimes V^{\otimes (k-1)}\right) \boxtimes_{V^{\otimes k}} T_{\sigma}(N) \simeq T_{\sigma}(M \boxtimes N)
$$

for $2 \leq r \leq k$.

On the other hand, to carry out this plan we shall need the associativity of the tensor product for twisted modules. Let V be a regular and self-dual vertex operator algebra of CFT type and let G be a finite abelian automorphism group of V. For any g-twisted V-module M and h-twisted V-module N with $q, h \in G$, we establish that a tensor product of M and N exists and the tensor product functor is associative.

This result is achieved by using a theorem of [\[KO\]](#page-36-11) and [\[CKM\]](#page-35-7) and a theorem of [\[H2\]](#page-36-10), where our approach is essentially the same as that in [\[CKM\]](#page-35-7). First, it was proved in [\[CM,](#page-35-8) [M2\]](#page-37-1) that V^G is also a regular and self-dual vertex operator algebra of CFT type. Then by [\[H2\]](#page-36-10), the V^G -module category C_{V^G} is a modular tensor category. On the other hand, a fusion category $\text{Rep}(V)$ was introduced in [\[KO\]](#page-36-11) and [\[CKM\]](#page-35-7) as a subcategory of \mathcal{C}_{V^G} . This in particular implies that the tensor product functor for $\text{Rep}(V)$ is associative. It turns out that for any finite order automorphism g of V, a g-twisted V-module is an object in $\text{Rep}(V)$. Then it is shown that the tensor product of two twisted modules in the sense of [\[X\]](#page-37-2) and [\[DLM1\]](#page-35-6) exists and is equivalent to the tensor product in the category $\text{Rep}(V)$. Consequently, we obtain the existence and the associativity of the tensor product in the sense of [\[X\]](#page-37-2) and [\[DLM1\]](#page-35-6). One can easily check Theorem A holds for intertwining operators with logarithm functions with the same proof.

The following is the second main theorem of this paper:

Theorem B. Let V be a regular and self-dual vertex operator algebra of CFT type, and let $M_1, ..., M_k, N$ be Vmodules. Let σ be a k-cycle permutation. Then

$$
(M_1\otimes\cdots\otimes M_k)\boxtimes_{V^{\otimes k}}T_{\sigma}(N)\simeq T_{\sigma}(M_1\boxtimes_V\cdots\boxtimes_V M_k\boxtimes_V N).
$$

Now, let $\sigma \in S_k$ be an arbitrary permutation. A classical fact is that the conjugacy class of σ is uniquely determined by a partition κ of k, i.e., $\kappa = (k_1, \ldots, k_s)$ is a sequence of positive integers such that

$$
k_1 \geq k_2 \geq \cdots \geq k_s, \quad k_1 + k_2 + \cdots + k_s = k.
$$

Define $\sigma_{\kappa} = \sigma_1 \cdots \sigma_s$, where $\sigma_1 = (12 \cdots k_1)$, $\sigma_2 = ((k_1 + 1) \cdots (k_1 + k_2))$, and so on. Then $\sigma_{\kappa} = \mu \sigma \mu^{-1}$ for some permutation μ . It is known that the categories of σ -twisted modules and σ_k -twisted modules are isomorphic canonically. Using [\[BDM\]](#page-35-2) and [\[FHL\]](#page-36-6), we show that irreducible σ_{κ} -twisted $V^{\otimes k}$ -modules are classified as

$$
T_{\sigma_{\kappa}}(N_1,\ldots,N_s) := T_{\sigma_1}(N_1) \otimes \cdots \otimes T_{\sigma_s}(N_s),\tag{1.2}
$$

where N_1, \ldots, N_s are irreducible V-modules. Furthermore, irreducible σ -twisted $V^{\otimes k}$ -modules are classified as

$$
T_{\sigma}(N_1,\ldots,N_s) := (T_{\sigma_{\kappa}}(N_1,\ldots,N_s))^{\mu},\tag{1.3}
$$

where for any σ_{κ} -twisted $V^{\otimes k}$ -module (W, Y_W) , (W^{μ}, Y_W^{τ}) is a σ -twisted $V^{\otimes k}$ -module with $W^{\mu} = W$ as a vector space and

$$
Y_W^{\mu}(a,x) = Y_W(\mu(a),x) \quad \text{for } a \in V^{\otimes k}.
$$

The following is the third main result of this paper:

Theorem C. Assume that V is a regular and self-dual vertex operator algebra of CFT type. Let k be a positive integer and $\sigma \in S_k \subset \text{Aut}(V^{\otimes k})$. Suppose $\mu \sigma \mu^{-1} = \sigma_{\kappa}$, where $\kappa = (k_1, \ldots, k_s)$ is a partition of k and $\mu \in S_k$. Let M_1, \ldots, M_k and N_1, \ldots, N_s be irreducible V-modules. Then

$$
(M_1 \otimes \cdots \otimes M_k) \boxtimes T_{\sigma_k}(N_1, N_2, \ldots, N_s) \simeq T_{\sigma_k}(M^{[k_1]} \boxtimes N_1, \ldots, M^{[k_s]} \boxtimes N_s)
$$
(1.4)

as a σ_{κ} -twisted $V^{\otimes k}$ -module, and

$$
(M_1 \otimes \cdots \otimes M_k)^{\mu} \boxtimes T_{\sigma}(N_1, N_2, \ldots, N_s) \simeq T_{\sigma}(M^{[k_1]} \boxtimes N_1, \ldots, M^{[k_s]} \boxtimes N_s)
$$
(1.5)

as a σ -twisted $V^{\otimes k}$ -module, where $M^{[k_1]} = \boxtimes_{i=1}^{k_1} M_i$, $M^{[k_2]} = \boxtimes_{i=k_1+1}^{k_1+k_2} M_i$, and so on, are V-modules.

We can replace the regularity assumption in Theorems B and C by the associativity assumption of tensor product in $(V^{\otimes k})^{\langle \sigma \rangle}$ -module category, which is what we really need.

This paper is organized as follows. In Section 2, we review some basics on vertex operator algebras, we also present the analytic function version of the twisted Jacobi Identity. In Section 3, we prove that for a regular, self-dual vertex operator algebra V of CFT type and a finite abelian automorphism group G of V, the tensor product $M\boxtimes_V N$ for any g-twisted V-module M and h-twisted V-module N exists and is associative for $g, h \in G$. In Section 4, we present the weak V-module $\mathcal{H}(M_2, M_3)$ of generalized intertwining operators from a σ -twisted V-module M_2 to another σ twisted V-module M_3 . In Section 5, for any V-modules M, N, W , we construct a linear isomorphism between the spaces $I_V \binom{W}{M \ N}$ and $I_{V^{\otimes k}} \binom{T_{\sigma}(W)}{M^1 \ T_{\sigma}(N)}$, and prove Theorem A: $\left(M \otimes V^{\otimes (k-1)}\right) \boxtimes T_{\sigma}(N) = T_{\sigma} (M \boxtimes_V N)$. Finally, in Section 6, by applying associativity of the tensor product ⊠ in Section 3, we obtain Theorems B and C.

Notations: For this paper, we use \mathbb{Z}_+ to denote the set of nonnegative integers.

2 Basics

Let $(V, Y, 1, \omega)$ be a vertex operator algebra in the sense of [\[FLM\]](#page-36-0) and [\[FHL\]](#page-36-6) (cf. [\[Bo\]](#page-35-9)). First, we recall the definitions of an automorphism of V and a g-twisted V-module for a finite order automorphism g of V (see [\[FLM\]](#page-36-0), [\[DLM1\]](#page-35-6)).

Definition 2.1. An *automorphism* of a vertex operator algebra V is a linear isomorphism g of V such that $g(\omega) = \omega$ and $gY(v, z)g^{-1} = Y(gv, z)$ for any $v \in V$. Denote by Aut (V) the group of all automorphisms of V.

A straightforward consequence is that every automorphism g of V preserves the vacuum vector 1, i.e., $g(1) = 1$. On the other hand, a standard fact is that for any subgroup $G \leq Aut(V)$, the set of G-fixed points

$$
V^G := \{ v \in V \mid g(v) = v \text{ for } g \in G \}
$$

is a vertex operator subalgebra.

Let g be a finite order automorphism of V with a period T in the sense that T is a positive integer such that $g^T = 1$. Then

$$
V = \bigoplus_{r=0}^{T-1} V^r,\tag{2.1}
$$

where $V^r = \{ v \in V \mid gv = e^{2\pi i r/T} v \}$ for $r \in \mathbb{Z}$. Note that for $r, s \in \mathbb{Z}$, $V^r = V^s$ if $r \equiv s \pmod{T}$.

Definition 2.2. A *weak* g*-twisted* V *-module* is a vector space M with a linear map

$$
Y_M(\cdot, z) : V \to (\text{End}M) \left[[z^{1/T}, z^{-1/T}] \right]
$$

$$
v \mapsto Y_M(v, z) = \sum_{n \in \frac{1}{T} \mathbb{Z}} v_n z^{-n-1} \ (v_n \in \text{End}M),
$$

which satisfies the following conditions: For all $u \in V^r$, $v \in V$, $w \in M$ with $0 \le r \le T - 1$,

$$
Y_M(u, z) = \sum_{n \in \frac{r}{T} + \mathbb{Z}} u_n z^{-n-1},
$$

 $u_n w = 0$ for *n* sufficiently large,

$$
Y_M(\mathbf{1},z)=Id_M,
$$

$$
z_0^{-1}\delta\left(\frac{z_1 - z_2}{z_0}\right)Y_M(u, z_1)Y_M(v, z_2) - z_0^{-1}\delta\left(\frac{z_2 - z_1}{-z_0}\right)Y_M(v, z_2)Y_M(u, z_1)
$$

$$
= z_2^{-1}\left(\frac{z_1 - z_0}{z_2}\right)^{-\frac{r}{T}}\delta\left(\frac{z_1 - z_0}{z_2}\right)Y_M\left(Y\left(u, z_0\right)v, z_2\right),\tag{2.2}
$$

where $\delta(z) = \sum_{n \in \mathbb{Z}} z^n$.

Notice that for any weak g-twisted V-module (M, Y_M) , from definition we have

$$
z^{\frac{r}{T}} Y_M(u, z) w \in M((z)) \quad \text{for } u \in V^r, \ r \in \mathbb{Z}, \ w \in M. \tag{2.3}
$$

On the other hand, we have (see [\[DL\]](#page-35-10))

$$
z_2^{-1}\left(\frac{z_1-z_0}{z_2}\right)^{-\frac{r}{T}}\delta\left(\frac{z_1-z_0}{z_2}\right)=z_1^{-1}\left(\frac{z_2+z_0}{z_1}\right)^{\frac{r}{T}}\delta\left(\frac{z_1-z_0}{z_2}\right).
$$

Remark 2.3. It is well known that the twisted Jacobi identity [\(2.2\)](#page-4-0) is equivalent to the following associativity relation

$$
(z_0 + z_2)^{l + \frac{r}{T}} Y_M (u, z_0 + z_2) Y_M (v, z_2) w = (z_2 + z_0)^{l + \frac{r}{T}} Y_M (Y (u, z_0) v, z_2) w
$$
\n(2.4)

for $w \in M$, where l is a nonnegative integer such that $z^{l+\frac{r}{T}} Y_M(u, z) w$ involves only nonnegative integral powers of z, and commutator relation

$$
\begin{aligned} \left[Y_M\left(u,z_1\right), Y_M\left(v,z_2\right)\right] \\ &= \text{Res}_{z_0} z_2^{-1} \left(\frac{z_1 - z_0}{z_2}\right)^{-\frac{r}{T}} \delta\left(\frac{z_1 - z_0}{z_2}\right) Y_M\left(Y\left(u,z_0\right)v,z_2\right). \end{aligned} \tag{2.5}
$$

From [\[DLMi\]](#page-35-11) (cf. Definition 2.7 and Lemma 2.9 of [\[LTW\]](#page-37-3)), [\(2.4\)](#page-4-1) can be equivalently replaced with the property that for $u, v \in V$, there exists a nonnegative integer m such that

$$
(z_1 - z_2)^m Y_M(u, z_1) Y_M(v, z_2) \in \text{Hom}\left(M, M((z_1^{\frac{1}{T}}, z_2^{\frac{1}{T}}))\right)
$$

and

$$
z_0^m Y_M(Y(u, z_0)v, z_2) = (z_1 - z_2)^m Y_M(u, z_1) Y_M(v, z_2)|_{z_1^{\frac{1}{T}} = (z_2 + z_0)^{\frac{1}{T}}}.
$$

Definition 2.4. An *admissible g-twisted V-module* is a weak g-twisted module with a $\frac{1}{T}\mathbb{Z}_+$ -grading $M = \bigoplus_{n\in\frac{1}{T}\mathbb{Z}_+} M(n)$ such that $u_m M(n) \subset M$ (wt $u - m - 1 + n$) for homogeneous $u \in V$ and $m, n \in \frac{1}{T} \mathbb{Z}$.

Note that if $M=\bigoplus_{n\in\frac{1}{T}\mathbb{Z}_+}M(n)$ is an irreducible admissible g-twisted V-module, then there is a complex number λ_M such that $L(0)|_{M(n)} = \lambda_M + n$ for all n. As a convention, we assume $M(0) \neq 0$, and λ_M is called the weight or conformal weight of M.

Definition 2.5. A g-twisted V-module is a weak g-twisted V-module M which carries a C-grading induced by the spectrum of $L(0)$ where $L(0)$ is the component operator of $Y(\omega, z) = \sum_{n \in \mathbb{Z}} L(n)z^{-n-2}$. That is, we have $M = \bigoplus_{\lambda \in \mathbb{C}} M_{\lambda}$, where $M_{\lambda} = \{w \in M \mid L(0)w = \lambda w\}$. Moreover, it is required that $\dim M_{\lambda} < \infty$ for al fixed λ_0 , $M_{\frac{n}{T} + \lambda_0} = 0$ for all small enough integers n.

In case $g = 1$, we recover the notions of weak, ordinary and admissible V-modules (see [\[DLM2\]](#page-35-12)).

Definition 2.6. A vertex operator algebra V is said to be g*-rational* if the admissible g-twisted module category is semisimple. In particular, V is said to be *rational* if V is 1-rational.

It was proved in [\[DLM2\]](#page-35-12) that if V is a q -rational vertex operator algebra, then there are only finitely many irreducible admissible g-twisted V-modules up to isomorphism and any irreducible admissible g-twisted V-module is ordinary.

Definition 2.7. A vertex operator algebra V is said to be *regular* if every weak V-module M is a direct sum of irreducible ordinary V -modules.

Definition 2.8. A vertex operator algebra V is said to be C_2 -cofinite if $V/C_2(V)$ is finite dimensional, where $C_2(V)$ = span ${u_{-2}v \mid u, v \in V}$.

A vertex operator algebra $V = \bigoplus_{n \in \mathbb{Z}} V_n$ is said to be of *CFT type* if $V_n = 0$ for all negative integers n and $V_0 = \mathbb{C}1.$

Remark 2.9. We here recall some known results we shall use extensively.

(1) It was proved in [\[DLM2,](#page-35-12) [Li4\]](#page-36-14) that if V is a C_2 -cofinite vertex operator algebra, then V has only finitely many irreducible admissible modules up to isomorphism.

(2) Assume that V is rational and C_2 -cofinite. It was proved in [\[ADJR\]](#page-35-13) that V is g-rational for any finite automor-phism g, and on the other hand, it was proved in [\[DLM3\]](#page-35-14) that λ_M is a rational number for every irreducible g-twisted V -module M.

(3) It is shown in [\[KL\]](#page-36-15) and [\[ABD\]](#page-35-15) that for CFT type vertex operator algebras, regularity is equivalent to rationality and C_2 -cofiniteness.

Definition 2.10. Let $M = \bigoplus_{n \in \frac{1}{T}\mathbb{Z}_+} M(n)$ be an admissible g-twisted V-module. Set

$$
M' = \bigoplus_{n \in \frac{1}{T}\mathbb{Z}_+} M(n)^*,
$$

the *restricted dual*, where $M(n)^* = \text{Hom}_{\mathbb{C}}(M(n), \mathbb{C})$. For $v \in V$, define a vertex operator $Y_{M'}(v, z)$ on M' via

$$
\langle Y_{M'}(v,z)f,u\rangle = \langle f, Y_M(e^{zL(1)}(-z^{-2})^{L(0)}v,z^{-1})u\rangle,
$$

where $\langle f, w \rangle = f(w)$ is the natural paring $M' \times M \to \mathbb{C}$. On the other hand, if $M = \bigoplus_{\lambda \in \mathbb{C}} M_{\lambda}$ is a g-twisted V-module, we define $M' = \bigoplus_{\lambda \in \mathbb{C}} M_{\lambda}^*$ and define $Y_{M'}(v, z)$ for $v \in V$ in the same way.

The same argument of [\[FHL\]](#page-36-6) gives:

Proposition 2.11. If (M, Y_M) is an admissible g-twisted V-module, then $(M', Y_{M'})$ carries the structure of an admissible g^{-1} -twisted V-module. On the other hand, if (M, Y_M) is a g-twisted V-module, then $(M', Y_{M'})$ carries the structure of a g^{-1} -twisted V-module. Moreover, M is irreducible if and only if M' is irreducible.

In particular, if (M, Y_M) is a V-module, then $(M', Y_{M'})$ is also a V-module. A V-module M is said to be selfdual if M and M' are isomorphic. A vertex operator algebra V is said to be *self-dual* if V and V' are isomorphic V -modules.

We shall need the following result from [\[CM,](#page-35-8) [M2\]](#page-37-1):

Theorem 2.12. Assume that V is a regular and self-dual vertex operator algebra of CFT type. Then for any solvable subgroup G of $\mathrm{Aut}(V)$, V^G is a regular and self-dual vertex operator algebra of CFT type.

Let g_1, g_2, g_3 be mutually commuting automorphisms of V of periods T_1, T_2, T_3 , respectively. In this case, V decomposes into the direct sum of common eigenspaces for g_1 and g_2 :

$$
V = \bigoplus_{0 \le j_1 < T_1, \ 0 \le j_2 < T_2} V^{(j_1, j_2)}
$$

where for $j_1, j_2 \in \mathbb{Z}$,

$$
V^{(j_1,j_2)} = \left\{ v \in V \mid g_s v = e^{2\pi i j_s/T_s}, s = 1,2 \right\}.
$$
 (2.6)

,

For any complex number α , we define

$$
(-1)^{\alpha} = e^{\alpha \pi i}.
$$

Now, we define intertwining operators among weak g_s -twisted modules (M_s, Y_{M_s}) for $s = 1, 2, 3$.

Definition 2.13. An *intertwining operator of type* $\binom{M_3}{M_1 M_2}$ associated with the given data is a linear map

$$
\mathcal{Y}(\cdot, z): M_1 \to (\text{Hom}(M_2, M_3)) \{z\}
$$

$$
w \mapsto \mathcal{Y}(w, z) = \sum_{n \in \mathbb{C}} w_n z^{-n-1}
$$

such that for any $w^1 \in M_1$, $w^2 \in M_2$ and for any fixed $c \in \mathbb{C}$,

$$
w_{c+n}^1 w^2 = 0 \quad \text{for } n \in \mathbb{Q} \text{ sufficiently large},
$$

$$
z_0^{-1} \left(\frac{z_1 - z_2}{z_0}\right)^{j_1/T_1} \delta\left(\frac{z_1 - z_2}{z_0}\right) Y_{M_3}(u, z_1) \mathcal{Y}(w, z_2)
$$

$$
- z_0^{-1} \left(\frac{z_2 - z_1}{z_0}\right)^{j_1/T_1} \delta\left(\frac{z_2 - z_1}{z_0}\right) \mathcal{Y}(w, z_2) Y_{M_2}(u, z_1)
$$

$$
= z_2^{-1} \left(\frac{z_1 - z_0}{z_2}\right)^{-j_2/T_2} \delta\left(\frac{z_1 - z_0}{z_2}\right) \mathcal{Y}\left(Y_{M_1}(u, z_0)w, z_2\right) \tag{2.7}
$$

on M_2 for $u \in V^{(j_1,j_2)}$ with $j_1, j_2 \in \mathbb{Z}$ and $w \in M_1$, and

$$
\frac{d}{dz}\mathcal{Y}(w,z) = \mathcal{Y}(L(-1)w,z).
$$

All intertwining operators of type $\binom{M_3}{M_1 M_2}$ form a vector space, which we denote by $I_V \binom{M_3}{M_1 M_2}$. Set $N_{M_1 M_2}^{M_3}$ = $\dim I_V \binom{M_3}{M_1 \ M_2}$, which is called the *fusion rule*.

Remark 2.14. As it was proved in [\[X\]](#page-37-2), if there are weak g_s -twisted modules (M_s, Y_{M_s}) for $s = 1, 2, 3$ such that $N_{M_1 M_2}^{M_3} > 0$, then $g_3 = g_1 g_2$. In view of this, we shall always assume that $g_3 = g_1 g_2$.

By the same arguments in Chapter 7 of [\[DL\]](#page-35-10) for formulas (7.24) and (7.11) we have:

Proposition 2.15. The twisted Jacobi identity [\(2.7\)](#page-6-0) is equivalent to the *generalized commutativity*: For $u \in V^{(j_1,j_2)}$ and $w \in M_1$, there is a positive integer n such that

$$
(z_1 - z_2)^{n + \frac{j_1}{T_1}} Y_{M_3}(u, z_1) \mathcal{Y}(w, z_2) = (-z_2 + z_1)^{n + \frac{j_1}{T_1}} \mathcal{Y}(w, z_2) Y_{M_2}(u, z_1)
$$

for $u \in V^{(j_1,j_2)}$, $w \in M_1$ and *generalized associativity*: For $u \in V^{(j_1,j_2)}$, $w \in M_1$ and $w_2 \in M_2$, there is a positive integer n depending on u and w_2 only such that

$$
(z_0 + z_2)^{n + \frac{j_2}{T_2}} Y_{M_3}(u, z_1) \mathcal{Y}(w, z_2) w_2 = (z_2 + z_0)^{n + \frac{j_2}{T_2}} \mathcal{Y}(Y_{M_1}(u, z_0)w, z_2) w_2.
$$

Let V_1 and V_2 be vertex operator algebras and let σ_1, σ_2 be finite order automorphisms of V_1, V_2 , respectively. Let W_1 be a V_1 -module, W_2 , W_3 σ_1 -twisted V_1 -modules, and let N_1 be a V_2 -module, N_2 , N_3 σ_2 -twisted V_2 -modules. Note that $W_1 \otimes N_1$ is a $V_1 \otimes V_2$ -module, while $W_2 \otimes N_2$ and $W_3 \otimes N_3$ are σ -twisted $V_1 \otimes V_2$ -modules with $\sigma = \sigma_1 \otimes \sigma_2$. For any intertwining operator $\mathcal{Y}_1(\cdot, z)$ of type $\binom{W_3}{W_1\ W_2}$ and for any intertwining operator $\mathcal{Y}_2(\cdot, z)$ of type $\binom{N_3}{N_1\ N_2}$, define a linear map

$$
(\mathcal{Y}_1 \otimes \mathcal{Y}_2)(\cdot, z): W_1 \otimes N_1 \to (\text{Hom}(W_2 \otimes N_2, W_3 \otimes N_3))\{z\}
$$

by

$$
(\mathcal{Y}_1 \otimes \mathcal{Y}_2)(w_1 \otimes n_1, z)(w_2 \otimes n_2) = \mathcal{Y}_1(w_1, z)w_2 \otimes \mathcal{Y}_2(n_1, z)n_2
$$
\n
$$
(2.8)
$$

for $w_1 \in W_1$, $w_2 \in W_2$, $n_1 \in N_1$, $n_2 \in N_2$. As in [\[ADL\]](#page-35-16), by using the generalized commutativity and associativity, it is straightforward to show that $(\mathcal{Y}_1 \otimes \mathcal{Y}_2)(\cdot, z)$ is an intertwining operator of type ${W_3 \otimes N_3 \choose W_1 \otimes N_1 W_2 \otimes N_2}$. Then we have a linear map

$$
\psi: I_{V_1}\begin{pmatrix} W_3 \\ W_1 \end{pmatrix} \otimes I_{V_2}\begin{pmatrix} N_3 \\ N_1 \end{pmatrix} \to I_{V_1 \otimes V_2}\begin{pmatrix} W_3 \otimes N_3 \\ W_1 \otimes N_1 \end{pmatrix}
$$

$$
(\mathcal{Y}_1, \mathcal{Y}_2) \mapsto \mathcal{Y}_1 \otimes \mathcal{Y}_2.
$$
 (2.9)

The same arguments of [\[ADL\]](#page-35-16) in proving Theorem 2.10 (with minor necessary modifications) give:

Theorem 2.16. Let $V_1, V_2, \sigma_1, \sigma_2, W_1, W_2, W_3, N_1, N_2, N_3$ be given as above. Then the linear map ψ is one-to-one. Furthermore, if either

$$
\dim I_{V^1}\binom{W_3}{W_1\,W_2} < \infty, \text{ or } \dim I_{V^2}\binom{N_3}{N_1\,N_2} < \infty,
$$

then ψ is a linear isomorphism.

Set
$$
\mathbb{D} = \mathbb{C}^2 \setminus \{(z, z), (z, 0), (0, z) | z \in \mathbb{C}\}
$$
. Then we have:

Proposition 2.17. Let V be rational and C_2 -cofinite, and let $G \subset Aut(V)$ be a finite abelian group. Then in the presence of other conditions in Definition [2.13,](#page-6-1) the Jacobi identity [\(2.7\)](#page-6-0) is equivalent to the following properties for $g_1, g_2 \in G$ and $v \in V^{(j_1, j_2)}, w \in M_1, w_2 \in M_2, w'_3 \in M'_3$:

(a) The formal series

$$
\langle w_3', Y_{M_3}(v, z_1)\mathcal{Y}(w, z_2)w_2\rangle(z_1-z_2)^{j_1/T_1}z_1^{j_2/T_2}
$$

converges to a unique multi-valued analytic function $f(z_1, z_2) = \frac{\sum_i h_i(z_1, z_2) z_2^{r_i}}{z_1^r z_2^s (z_1 - z_2)^t}$ for some $h_i(z_1, z_2) \in \mathbb{C}[z_1, z_2]$, $r_i \in \mathbb{Q}$ and $r, s, t \in \mathbb{Z}$ in the domain $|z_1| > |z_2| > 0$.

(b) The formal series

$$
\langle w_3', \mathcal{Y}(w, z_2) Y_{M_2}(v, z_1) w_2 \rangle (-z_2 + z_1)^{j_1/T_1} z_1^{j_2/T_2}
$$

converges to $f(z_1, z_2)$ in the domain $|z_2| > |z_1| > 0$.

(c) The formal series

$$
\langle w_3', \mathcal{Y} (Y_{M_1}(v, z_1 - z_2) w, z_2) w_2 \rangle (z_1 - z_2)^{j_1/T_1} z_1^{j_2/T_2}
$$

converges to $f(z_1, z_2)$ in the domain $|z_2| > |z_1 - z_2| > 0$.

Proof. Since V^G is rational by Theorem [2.12,](#page-5-0) from [\[DLM3\]](#page-35-14) each M_i is a direct sum of finitely many irreducible V^G -modules whose weights are rational. Without loss we can assume that w, w_2, w'_3 are vectors in some irreducible V^G -modules. Then there exists a rational number r such that the powers of z_2 in the formal power series $\langle w_3', Y_{M_3}(v, z_1)\mathcal{Y}(w, z_2)w_2\rangle$ lie in $r + \mathbb{Z}$. The same is true for the corresponding terms in (b) and (c). Now results follow from Proposition [2.15](#page-6-2) and the proofs of Propositions 7.6, 7.8 and 7.9 of [\[DL\]](#page-35-10). \Box

Now we recall a notion of tensor product (see [\[HL1\]](#page-36-7), [\[Li3\]](#page-36-13), [\[DLM1\]](#page-35-6)).

Definition 2.18. Let g_1, g_2 be commuting finite order automorphisms of a vertex operator algebra V and let M_i be a g_i -twisted V-module for $i=1,2.$ A *tensor product* for the ordered pair (M_1,M_2) is a pair $(M,F(\cdot,z))$ where M is a weak g_1g_2 -twisted V-module and $F(\cdot, z)$ is an intertwining operator of type $\binom{M}{M_1 M_2}$ such that the following universal property holds: For any weak g_1g_2 -twisted V-module W and for any intertwining operator $I(\cdot, z)$ of type ${W \choose M_1 \ M_2}$, there exists a unique weak twisted V-module homomorphism ψ from M to W such that $I(\cdot, z) = \psi \circ F(\cdot, z)$.

We shall use $M_1 \boxtimes^V M_2$ to denote a generic tensor product module of M_1 and M_2 , provided that the existence is verified.

Remark 2.19. We here recall Huang-Lepowsky's tautological construction of a tensor product (see [\[HL1\]](#page-36-7)). Suppose that V is a vertex operator algebra with a finite order automorphism τ such that V is τ -rational, namely, every τ -twisted V-module is completely reducible. Let W be a V-module and T a τ -twisted V-module such that $\dim I_V \binom{P}{W_T} < \infty$ for every irreducible τ -twisted V-module P. Let $\{P_\alpha \mid \alpha \in A\}$ be a complete set of equivalence class representatives of irreducible τ -twisted V-modules. Set

$$
M = \bigoplus_{\alpha \in \mathcal{A}} I \binom{P_{\alpha}}{W \, T}^* \otimes P_{\alpha} = \bigoplus_{\alpha \in \mathcal{A}} \text{Hom}\left(I \binom{P_{\alpha}}{W \, T}, P_{\alpha} \right),\tag{2.10}
$$

a τ -twisted V-module. For $\alpha \in A$, define $\mathcal{Y}^{\alpha}(w, z)t \in \text{Hom}\left(I\left(\begin{smallmatrix} P_{\alpha} \\ W^T \end{smallmatrix}\right), P_{\alpha}\right) \{z\}$ by

$$
(\mathcal{Y}^\alpha(w,z)t)(I)=I(w,z)t\quad\text{for }I\in I\binom{P_\alpha}{W\,T}.
$$

Set $\mathcal{Y}(\cdot, z) = \sum_{\alpha \in \mathcal{A}} \mathcal{Y}^{\alpha}(\cdot, z)$. Then (M, \mathcal{Y}) is a tensor product of W and T.

The following is a simple property of fusion rules (numbers) (see [\[FHL\]](#page-36-6), [\[HL1\]](#page-36-7), [\[DLM1\]](#page-35-6):

Lemma 2.20. Let g_1, g_2 be commuting finite order automorphisms of a vertex operator algebra V and let M_i be a g_i-twisted V-module for $i = 1, 2, 3$ with $g_3 = g_1 g_2$. Then $N_{M_1 M_2}^{\overline{M}_3} = N_{M_2 M_1}^{\overline{M}_3}$.

3 Tensor products of twisted modules

Throughout this section, we assume that V is a regular and self-dual vertex operator algebra of CFT type and G is a finite abelian automorphism group of V. The main goal of this section is to prove that for any $q, h \in G$ and for any q-twisted V-module M and h-twisted V-module N, a tensor product of M and N exists and associativity holds. These results will be used later to determine the tensor product of an untwisted module with a twisted module for the study on permutation orbifolds.

Denote by C_V the category of ordinary V-modules and by C_{V^G} the category of ordinary V^G -modules. From [\[H2\]](#page-36-10), both C_V and C_V are modular tensor categories. Furthermore, from [\[KO\]](#page-36-11) and [\[CKM\]](#page-35-7), V is a commutative associative algebra in C_{V^G} as $V = \bigoplus_{\chi \in \text{irr}(G)} V^{\chi}$, where $\text{irr}(G)$ is the set of irreducible characters of G and V^{χ} are irreducible V^G -modules (cf. [\[DJX\]](#page-35-17)), i.e., simple objects in \mathcal{C}_{V^G} .

Recall the following definition from [\[KO\]](#page-36-11) and [\[CKM\]](#page-35-7) (see Proposition 3.46 of [\[CKM\]](#page-35-7)):

Definition 3.1. Denote by $\text{Rep}(V)$ the subcategory of C_{V^G} consisting of every V^G -module W together with a V^G intertwining operator $Y_W(\cdot, z)$ of type $\binom{W}{V_W}$ such that the following conditions are satisfied:

1. (Associativity) For any $u, v \in V$, $w \in W$ and $w' \in W'$, the formal series

$$
\langle w', Y_W(u,z_1)Y_W(v,z_2)w \rangle
$$

and

$$
\langle w', Y_W(Y(u,z_1-z_2)v,z_2))w\rangle
$$

converge on the domains $|z_1| > |z_2| > 0$ and $|z_2| > |z_1 - z_2| > 0$, respectively, to multivalued analytic functions which coincide on their common domain.

2. (Unit)
$$
Y_W(1, z) = Id_W
$$
.

Recall from [\[KO\]](#page-36-11) (see also [\[CKM\]](#page-35-7)) that there is a categorical tensor product functor \mathbb{Z}_V in the category of Rep(V), which is associative. We denote by $M \boxtimes_V N$ the tensor product of M and N in $\text{Rep}(V)$. Then for any two V^G -modules M, N in $\operatorname{Rep}(V), M \boxtimes_V N$ is a quotient of $M \boxtimes^{V^G} N$. Moreover, $\operatorname{Rep}(V)$ is a fusion category. In particular, $\operatorname{Rep}(V)$ is a semisimple category with finitely many inequivalent simple objects.

From the generalized associativity for a twisted module (see Proposition [2.15\)](#page-6-2), we immediately have:

Lemma 3.2. If W is a g-twisted V-module with $g \in G$, then W is an object of $Rep(V)$. Furthermore, if W_i is a g_i-twisted V-module with $g_i \in G$ for $i = 1, 2$, then W_1 and W_2 are equivalent objects in Rep(V) if and only if $g_1 = g_2$ and $W_1 \simeq W_2$ as twisted V-modules.

On the other hand, we have the following lemma (see also [\[K1,](#page-36-16) [K2\]](#page-36-17)).

Lemma 3.3. If W is a simple object in $\text{Rep}(V)$, then W is an irreducible g-twisted V-module for some $g \in G$.

Proof. We need to use the Frobenius-Perron dimension in a fusion category (see [\[BK\]](#page-35-18), [\[ENO\]](#page-36-18), [\[DMNO\]](#page-35-19)). Let C be a fusion category and let $K(\mathcal{C})$ denote the Grothendieck ring of C. According to [\[ENO\]](#page-36-18), there exists a unique ring homomorphism FPdim $c : K(\mathcal{C}) \to \mathbb{R}$ such that FPdim $c(X) > 0$ for all nonzero $X \in \mathcal{C}$, where FPdim $c(X)$ is called the Frobenius-Perron dimension of $X \in \mathcal{C}$. Furthermore, one has the Frobenius-Perron dimension for the category C:

$$
\mathrm{FPdim}(\mathcal{C}) = \sum_{X \in \mathcal{O}(\mathcal{C})} \mathrm{FPdim} \, \mathcal{C}(X)^2,
$$

where $\mathcal{O}(\mathcal{C})$ denotes the equivalence classes of the simple objects in C. Note that both \mathcal{C}_{V^G} and $\text{Rep}(V)$ are fusion categories. It follows from [\[DMNO\]](#page-35-19) and [\[ENO\]](#page-36-18) that

$$
(\text{FPdim}_{\mathcal{C}_{V}G}V)(\text{FPdim}(\text{Rep}(V))) = \text{FPdim}(\mathcal{C}_{V^G}).\tag{3.1}
$$

.

We also need quantum dimensions and global dimensions from [\[DJX\]](#page-35-17) and [\[DRX\]](#page-36-19). Let M be a g-twisted V-module with $g \in G$ and set $ch_q M = \text{tr}|_M q^{L(0)-c/24}$. Then $ch_q M$ converges to a holomorphic function $\chi_M(\tau)$ on the upper half plane with $q = e^{2\pi i \tau}$ (see [\[Z\]](#page-37-4), [\[DLM3\]](#page-35-14)). The quantum dimension of M over V is defined as

$$
\operatorname{qdim}_{V} M = \lim_{y \to 0} \frac{\chi_M(iy)}{\chi_V(iy)}
$$

From [\[ADJR\]](#page-35-13) we have

$$
\operatorname{qdim}_{V} M = \operatorname{FPdim}_{\operatorname{Rep}(V)} M, \text{ and } \operatorname{qdim}_{V^G} W = \operatorname{FPdim}_{\mathcal{C}_{V^G}} W
$$

for any V^G -module W. Denote the equivalence classes of irreducible g-twisted V-modules by $\mathcal{M}(g)$. The global dimension of V is defined as

$$
\text{glob}(V) = \sum_{M \in \mathcal{M}(1)} (\text{qdim}_V M)^2.
$$

It follows that $\text{glob}(V^G) = \text{FPdim}(\mathcal{C}_{V^G})$. Recall from [\[DJX\]](#page-35-17) and [\[DRX\]](#page-36-19) that

$$
qdim_{V^G} V = o(G),
$$

glob $(V^G) = o(G)^2$ glob (V) ,
glob $(V) = \sum_{M \in \mathcal{M}(g)} (qdim_V M)^2$

for any $g \in G$. Then we have

$$
\text{FPdim}_{\mathcal{C}_{V}G} V = \text{qdim}_{V^G} V = o(G), \quad \text{FPdim}(\mathcal{C}_{V^G}) = \text{glob}(V^G) = o(G)^2 \text{glob}(V).
$$

Combining these relations with equation [\(3.1\)](#page-9-0) we obtain

$$
FPdim (Rep(V)) = o(G)glob(V).
$$

Since for any $g \in G$, every irreducible g-twisted module is a simple object in $Rep(V)$ by Lemma [3.2,](#page-9-1) we get

$$
\mathrm{FPdim} \operatorname{Rep}(V) \ge \sum_{g \in G} \sum_{M \in \mathcal{M}(g)} (\mathrm{qdim}_V M)^2 = o(G) \mathrm{glob}(V) = \mathrm{FPdim} \operatorname{Rep}(V).
$$

Thus

$$
\mathrm{FPdim} \operatorname{Rep}(V) = \sum_{g \in G} \sum_{M \in \mathcal{M}(g)} (\mathrm{qdim}_V M)^2,
$$

which implies that every simple object in $\text{Rep}(V)$ is an irreducible g-twisted V-module for some $g \in G$.

Remark 3.4. Note that Lemma [3.3](#page-9-2) holds for any (not necessarily abelian) finite automorphism group G as long as V^G is rational and C_2 -cofinite.

Let g, $h \in G$. For any g-twisted module M and h-twisted module N, as they are objects in Rep(V) by Lemma [3.2,](#page-9-1) $M \boxtimes_V N$ exists in Rep(V). We will show next that the tensor product in the sense of Definition [2.18](#page-7-0) exists and is isomorphic to $M \boxtimes_V N$. We first prove that a categorical intertwining operator is the same as an intertwining operator in the sense of Definition [2.13.](#page-6-1)

The following is a generalization of Theorem 3.53 in [\[CKM\]](#page-35-7) where $g_i = 1$ for $i = 1, 2, 3$ with a similar proof:

Theorem 3.5. Let g_1, g_2, g_3 be commuting finite order automorphisms of V and let M_i be a g_i -twisted V-module for $i = 1, 2, 3$. Assume

$$
\mathcal{Y}(\cdot, z) \cdot : M_1 \otimes M_2 \longrightarrow M_3\{z\}
$$

$$
w_1 \otimes w_2 \longmapsto \mathcal{Y}(w_1, z)w_2 = \sum_{n \in \mathbb{C}} (w_1)_n w_2 z^{-n-1}
$$

is a bilinear map satisfying the following conditions:

1. *Lower truncation:* For any $w_1 \in M_1$, $w_2 \in M_2$ and $\alpha \in \mathbb{C}$, $(w_1)_{\alpha+m} w_2 = 0$ for sufficiently large $m \in \mathbb{N}$.

2. The
$$
L(-1)
$$
-derivative formula: $[L(-1), \mathcal{Y}(w_1, z)] = \frac{d}{dx} \mathcal{Y}(w_1, z) = \mathcal{Y}(L(-1)w_1, z)$ for $w_1 \in M_1$.

- 3. The $L(0)$ -bracket formula: $[L(0), \mathcal{Y}(w_1, z)] = z \frac{d}{dz} \mathcal{Y}(w_1, z) + \mathcal{Y}(L(0)w_1, z)$ for $w_1 \in M_1$.
- 4. For any $w'_3 \in M'_3$, $w_2 \in M_2$, $v \in V$, $w_1 \in M_1$, the following series define multivalued analytic functions on the indicated domains, which coincide on the respective intersections of their domains:

$$
\langle w_3', Y_{M_3}(a, z_1) \mathcal{Y}(w_1, z_2) w_2 \rangle \qquad \text{on } |z_1| > |z_2| > 0,
$$
\n(3.2)

 \Box

$$
\langle w_3', \mathcal{Y}(Y_{M_1}(a, z_1 - z_2)w_1, z_2)w_2 \rangle \qquad \text{on } |z_2| > |z_1 - z_2| > 0,
$$
\n(3.3)

$$
\langle w_3', \mathcal{Y}(w_1, z_2) Y_{M_2}(a, z_1) w_2 \rangle \qquad \text{on } |z_2| > |z_1| > 0. \tag{3.4}
$$

Specifically, the principal branches (that is, $z_2 = e^{\log z_2}$) of the first and second multivalued functions yield single-valued functions which coincide on the simply connected domain

$$
S_1 := \{(z_1, z_2) \in \mathbb{C}^2 \mid \text{Re } z_1 > \text{Re } z_2 > \text{Re}(z_1 - z_2) > 0, \text{ Im } z_1 > \text{Im } z_2 > \text{Im}(z_1 - z_2) > 0\}
$$

and the principal branches of the second and third multivalued functions yield single-valued functions which coincide on the simply connected domain

$$
S_2 := \{ (z_1, z_2) \in \mathbb{C}^2 \mid \text{Re } z_2 > \text{Re } z_1 > \text{Re}(z_2 - z_1) > 0, \text{ Im } z_2 > \text{Im } z_1 > \text{Im}(z_2 - z_1) > 0 \}.
$$

5. There exists a multivalued analytic function g defined on $\{(z_1, z_2) \in \mathbb{C}^2 \mid z_1, z_2, z_1 - z_2 \neq 0\}$ that restricts to the three multivalued functions above on their respective domains.

Then Y is an intertwining operator of type $\binom{M_3}{M_1 M_2}$ in the sense of Definition [2.13.](#page-6-1)

Proof. The proof here is a slight modification of that of Theorem 3.53 in [\[CKM\]](#page-35-7). First, fix $z_2 \in \mathbb{C}^\times$ with Re z_2 , Im $z_2 >$ 0. Then $f_{z_2}(z_1) := g(z_1, e^{\log z_2}) z_1^{j_1/T_1} (z_1 - z_2)^{j_2/T_2}$ yields a possibly multivalued analytic function of z_1 defined on $\{z_1 \in \mathbb{C}^\times \mid z_1 \neq z_2\}$. However, note that f_{z_2} has single-valued restrictions to each of $\{z_1 \in \mathbb{C}^\times \mid |z_1| > |z_2|\}$, ${z_1 \in \mathbb{C}^\times |z_2| > |z_1 - z_2| > 0}$ and ${z_1 \in \mathbb{C}^\times |z_2| > |z_1|}$, and these restrictions coincide on certain simplyconnected subsets of the intersections of these domains. Thus these restrictions define a single-valued analytic function on the union of these domains. This implies that f_{z_2} has a single-valued restriction f_{z_2} to $\mathbb{C}^\times \setminus \{z_2\}$ with possible poles at $0, z_2, \infty$. Thus f_{z_2} is a rational function. It follows from Proposition [2.17](#page-7-1) that Y satisfies the Jacobi identity and hence it is an intertwining operator of type $\binom{M_3}{M_1 M_2}$ in the sense of Definition [2.13.](#page-6-1)

Theorem 3.6. Let $g_1, g_2, g_3 \in G$ and let M_i be g_i -twisted V-modules for $i = 1, 2, 3$. Then the space of intertwining operators of type $\binom{M_3}{M_1\ M_2}$ in the sense of Definition [2.13](#page-6-1) is linearly isomorphic to the space of the categorical Rep(V) intertwining operators of type $\binom{M_3}{M_1\ M_2}$. Furthermore, a tensor product module $M_1\boxtimes^V M_2$ exists and $M_1\boxtimes^V M_2$ and $M_1 \boxtimes_V M_2$ are isomorphic $g_1 g_2$ -twisted V-modules.

Proof. By Theorem 3.44 of [\[CKM\]](#page-35-7), the space of the categorical $Rep(V)$ intertwining operators of type $\binom{M_3}{M_1 \ M_2}$ is isomorphic to the space of bilinear maps

$$
\mathcal{Y}(\cdot, z) \cdot : M_1 \otimes M_2 \longrightarrow M_3 \{z\}
$$

$$
w_1 \otimes w_2 \longmapsto \mathcal{Y}(w_1, z)w_2 = \sum_{n \in \mathbb{C}} (w_1)_n w_2 z^{-n-1}
$$

satisfying the conditions 1-5 in Theorem [3.5.](#page-10-0) By Proposition [2.17](#page-7-1) and Theorem [3.5,](#page-10-0) the latter space is the same as the space of intertwining operators of type $\binom{M_3}{M_1 M_2}$ in the sense of Definition [2.13.](#page-6-1) It follows from Remark [2.14](#page-6-3) that the space of the categorical $\text{Rep}(V)$ intertwining operators of type $\binom{M_3}{M_1 M_2}$ is zero if $g_1g_2 \neq g_3$. So we assume $g_1g_2 = g_3$ for the rest of the proof.

Let $\{M_3^1, \ldots, M_3^r\}$ be a complete set of equivalence class representatives of irreducible g_3 -twisted V-modules. From Lemma [3.2,](#page-9-1) M_3^1, \ldots, M_3^r are inequivalent simple objects in $\text{Rep}(V)$. From the first part, the fusion rules $N_{M_1,M_2}^{M_3^i}$ of the indicated types in the sense of Definition [2.13](#page-6-1) and in $\text{Rep}(V)$ are the same. As $M_1 \boxtimes_V M_2$ is an object in Rep(V) (consisting of some V^G-modules), the fusion numbers $N_{M_1,M_2}^{M_3^i}$ are finite. For $1 \le i \le r$, set $N_i = N_{M_1,M_2}^{M_3^i}$ for short. Set

$$
W = \bigoplus_{i=1}^r (\mathbb{C}^{N_i} \otimes M_3^i),
$$

which is a g_3 -twisted V-module and hence an object in $Rep(V)$. It follows that there is an isomorphism in the category $\mathrm{Rep}(V)$ from $M_1 \boxtimes_V M_2$ to W .

For $1 \le i \le r$, let $\mathcal{Y}_1^i, ..., \mathcal{Y}_{N_i}^i$ be a basis of $I_V \binom{M_3^i}{M_1 M_2}$. Then we have intertwining operators $e_s \otimes \mathcal{Y}_j^i$ for $1 \le i \le r, \ 1 \le s, j \le N_i$ of type ${W \choose M_1 M_2}$, where e_1, \ldots, e_{N_i} are the standard basis vectors of \mathbb{C}^{N_i} . Set

$$
F(\cdot, z) = \sum_{i=1}^{r} \sum_{s,j=1}^{N_i} e_s \otimes \mathcal{Y}_j^i(\cdot, z).
$$
 (3.5)

We now prove that (W, F) is a tensor product of M_1 and M_2 in the sense of Definition [2.18.](#page-7-0) Let M be any g_3 twisted V-module and $I \in I_V \binom{M}{M_1 \ M_2}$. We need to show that there is a unique V-homomorphism $\psi: W \to M$ such that $I = \psi \circ \mathcal{Y}$. Since V is g_3 -rational, M is isomorphic to a direct sum of M_3^i for $1 \le i \le r$. Thus $M =$ $\oplus_{i=1}^r \text{Hom}_V(M_3^i,M) \otimes M_3^i.$ It can be readily seen that

$$
I_V\binom{M}{M_1\ M_2}=\oplus_{i=1}^r\mathrm{Hom}_V(M_3^i,M)\otimes I_V\binom{M_3^i}{M_1\ M_2}.
$$

Then $I = \sum_{i=1}^r \sum_{j=1}^{N_i} \psi_{i,j} \otimes \mathcal{Y}_j^i$ where $\psi_{i,j} \in \text{Hom}_V(M_3^i, M)$. For $1 \le i \le r, 1 \le s \le N_i$, let $p_{i,s}$ be the projection of M onto $e_s \otimes M_3^i$, a V-homomorphism, and let $\pi_{i,s}$ be the identification map $e_s \otimes M_3^i \simeq M_3^i$. Set $\psi=\sum_{i=1}^r\sum_{j,s=1}^{N_i}\psi_{i,j}\pi_{i,s}p_{i,s}$, a V -homomorphism from W to $M.$ It is clear that $I=\psi\circ F.$ Thus (W,F) is a tensor product of M_1 and M_2 in the sense of Definition [2.18.](#page-7-0) Consequently, we have $M_1 \boxtimes^V M_2 \simeq M_1 \boxtimes_V M_2$. \Box Since the tensor product functor \boxtimes_V is associative, we have:

Corollary 3.7. Let M_i be a g_i -twisted V-module with $g_i \in G$ for $i = 1, 2, 3$. Then

$$
(M_1 \boxtimes_V M_2) \boxtimes_V M_3 \simeq M_1 \boxtimes_V (M_2 \boxtimes_V M_3).
$$

4 Weak V-module $\mathcal{H}(M_2, M_3)$ of generalized intertwining operators

Let V be a vertex operator algebra and let $\tau \in$ Aut (V) of period T. In this section, we first prove that for any τ -twisted V-modules M_2 and M_3 , there is a weak module structure on the space $\mathcal{H}(M_2, M_3)$ of generalized intertwining operators. Furthermore, we prove that for any weak V-module M_1 , the space Hom_V $(M_1, \mathcal{H}(M_2, M_3))$ of Vhomomorphisms from M_1 to $\mathcal{H}(M_2, M_3)$ is canonically isomorphic to $I_V\binom{M_3}{M_1M_2}$. Hence giving a V-homomorphism from M_1 to $\mathcal{H}(M_2, M_3)$ is equivalent to giving an intertwining operator of type $\binom{M_3}{M_1 M_2}$.

As before, for $r \in \mathbb{Z}$ set

$$
V^r = \{ v \in V \mid \tau v = \eta_T^r v \},
$$

where

$$
\eta_T = e^{2\pi i/T}
$$

is the principal primitive T -th root of unity.

Definition 4.1. Let (M_2, Y_{M_2}) and (M_3, Y_{M_3}) be weak τ -twisted *V*-modules. A *generalized intertwining operator* from M_2 to M_3 is an element $\phi(x) = \sum_{\alpha \in \mathbb{C}} \phi_{\alpha} x^{-\alpha-1} \in (Hom(M_2, M_3))\{x\}$ satisfying the following conditions:

- (G1) For any $\alpha \in \mathbb{C}$, $w \in M_2$, $\phi_{\alpha+n}w = 0$ for $n \in \mathbb{Z}$ sufficiently large;
- (G2) $[L(-1), \phi(x)] = \frac{d}{dx}\phi(x);$
- (G3) For any $v \in V$, there exists a nonnegative integer k such that

$$
(x_1 - x_2)^k Y_{M_3}(v, x_1)\phi(x_2) = (x_1 - x_2)^k \phi(x_2) Y_{M_2}(v, x_1).
$$
\n(4.1)

Denote by $\mathcal{H}(M_2, M_3)$ the space of all *generalized intertwining operators* from M_2 to M_3 .

Let $\phi(x) \in \mathcal{H}(M_2, M_3)$ and let $u \in V^r$ with $0 \le r < T$. For each $n \in \mathbb{Z}$, we define

$$
u_n^{\mathcal{H}}\phi(x)
$$

= Res_{x₁} $\sum_{j\geq 0}$ $\left(-\frac{r}{j}\right)x^{-\frac{r}{T}-j}x_1^{\frac{r}{T}}\left((x_1-x)^{n+j}Y_{M_3}(u,x_1)\phi(x) - (-x+x_1)^{n+j}\phi(x)Y_{M_2}(u,x_1)\right),$ (4.2)

an element of $(Hom(M_2, M_3))\{x\}.$

Remark 4.2. Here are some explanations about this definition. First, recall that for any weak τ -twisted V-module (W, Y_W) , we have

$$
z^{\frac{r}{T}} Y_W(u, z) w \in W((z)) \quad \text{for } u \in V^r, \ r \in \mathbb{Z}, \ w \in W.
$$

In view of this, the sum in [\(4.2\)](#page-12-0) involves only integer powers of x_1 , so that applying Res_{x_1} to the sum makes sense. Second, due to the weak commutativity [\(4.1\)](#page-12-1), we see that for any $n \in \mathbb{Z}$, the sum contains only finitely many nonzero terms, and $u_n^{\mathcal{H}}\phi(x) = 0$ for *n* sufficiently large. Third, as $\phi(x)$ satisfies the condition (G1), it can be readily seen that $u_n^{\mathcal{H}}\phi(x)$ also satisfies the condition (G1).

Now, set

$$
Y_{\mathcal{H}}(u,z)\phi(x) = \sum_{n \in \mathbb{Z}} u_n^{\mathcal{H}} \phi(x) z^{-n-1}.
$$
\n(4.3)

In terms of this generating function, we have

$$
Y_{\mathcal{H}}(u, z)\phi(x) = \text{Res}_{x_1}\left(\frac{x+z}{x_1}\right)^{-\frac{r}{T}} \left(z^{-1}\delta\left(\frac{x_1-x}{z}\right)Y_{M_3}(u, x_1)\phi(x) - z^{-1}\delta\left(\frac{x-x_1}{-z}\right)\phi(x)Y_{M_2}(u, x_1)\right).
$$
 (4.4)

Using linearity, we obtain a vertex operator map

$$
Y_{\mathcal{H}}(\cdot, z) : V \to (\text{End}(\text{Hom}(M_2, M_3))\{x\}) [[z, z^{-1}]]
$$

$$
u \mapsto Y_{\mathcal{H}}(u, z). \tag{4.5}
$$

First, we present some technical results.

Lemma 4.3. Let $v \in V^r$ with $r \in \mathbb{Z}$ and let $\phi(x) \in \mathcal{H}(M_2, M_3)$. Then

$$
z^{k}(x+z)^{\frac{r}{T}}Y_{\mathcal{H}}(u,z)\phi(x) = \left((x_{1}-x)^{k}x_{1}^{\frac{r}{T}}Y_{M_{3}}(u,x_{1})\phi(x) \right)|_{x_{1}=x+z}, \tag{4.6}
$$

where k is any nonnegative integer such that (4.1) holds. Furthermore, we have

$$
x_1^{-1}\delta\left(\frac{x+z}{x_1}\right)\left(\frac{x+z}{x_1}\right)^{\frac{x}{T}}Y_{\mathcal{H}}(u,z)\phi(x)
$$

= $z^{-1}\delta\left(\frac{x_1-x}{z}\right)Y_{M_3}(u,x_1)\phi(x) - z^{-1}\delta\left(\frac{x-x_1}{-z}\right)\phi(x)Y_{M_2}(u,x_1).$ (4.7)

Proof. Write $r = qT + r_0$ for some $q, r_0 \in \mathbb{Z}$ with $0 \le r_0 < T$. As $Y_H(u, z)\phi(x)$ involves only finitely many negative powers of z, we can multiply both sides of [\(4.4\)](#page-13-0) by $(x + z)^{\frac{r_0}{T}}$ and by doing so we get

$$
z^{k}(x+z)^{\frac{r_{0}}{T}}Y_{\mathcal{H}}(u,z)\phi(x)
$$
\n
$$
= Res_{x_{1}}z^{k}x_{1}^{\frac{r_{0}}{T}}\left(z^{-1}\delta\left(\frac{x_{1}-x}{z}\right)Y_{M_{3}}(u,x_{1})\phi(x)-z^{-1}\delta\left(\frac{x-x_{1}}{-z}\right)\phi(x)Y_{M_{2}}(u,x_{1})\right)
$$
\n
$$
= Res_{x_{1}}z^{-1}\delta\left(\frac{x_{1}-x}{z}\right)\left((x_{1}-x)^{k}x_{1}^{\frac{r_{0}}{T}}Y_{M_{3}}(u,x_{1})\phi(x)\right)
$$
\n
$$
- Res_{x_{1}}z^{-1}\delta\left(\frac{x-x_{1}}{-z}\right)\left((x_{1}-x)^{k}x_{1}^{\frac{r_{0}}{T}}\phi(x)Y_{M_{2}}(u,x_{1})\right)
$$
\n
$$
= Res_{x_{1}}x_{1}^{-1}\delta\left(\frac{x+z}{x_{1}}\right)\left((x_{1}-x)^{k}x_{1}^{\frac{r_{0}}{T}}Y_{M_{3}}(u,x_{1})\phi(x)\right).
$$

Then

$$
z^{k}(x+z)^{\frac{r}{T}}Y_{\mathcal{H}}(u,z)\phi(x)
$$

= $z^{k}(x+z)^{q}(x+z)^{\frac{r_{0}}{T}}Y_{\mathcal{H}}(u,z)\phi(x)$
= Res_{x₁}x₁⁻¹ $\delta\left(\frac{x+z}{x_{1}}\right)(x+z)^{q}\left((x_{1}-x)^{k}x_{1}^{\frac{r_{0}}{T}}Y_{M_{3}}(u,x_{1})\phi(x)\right)$
= Res_{x₁}x₁⁻¹ $\delta\left(\frac{x+z}{x_{1}}\right)x_{1}^{q}\left((x_{1}-x)^{k}x_{1}^{\frac{r_{0}}{T}}Y_{M_{3}}(u,x_{1})\phi(x)\right)$
= Res_{x₁}x₁⁻¹ $\delta\left(\frac{x+z}{x_{1}}\right)\left((x_{1}-x)^{k}x_{1}^{\frac{r}{T}}Y_{M_{3}}(u,x_{1})\phi(x)\right)$
= $\left((x_{1}-x)^{k}x_{1}^{\frac{r}{T}}Y_{M_{3}}(u,x_{1})\phi(x)\right)|_{x_{1}=x+z}$.

This proves the first assertion. Furthermore, we have

$$
z^{k}x_{1}^{\frac{r}{T}}\left(z^{-1}\delta\left(\frac{x_{1}-x}{z}\right)Y_{M_{3}}(u,x_{1})\phi(x)-z^{-1}\delta\left(\frac{x-x_{1}}{-z}\right)\phi(x)Y_{M_{2}}(u,x_{1})\right)
$$

\n
$$
=z^{-1}\delta\left(\frac{x_{1}-x}{z}\right)\left(x_{1}^{\frac{r}{T}}(x_{1}-x)^{k}Y_{M_{3}}(u,x_{1})\phi(x)\right)-z^{-1}\delta\left(\frac{x-x_{1}}{-z}\right)\left(x_{1}^{\frac{r}{T}}(x_{1}-x)^{k}\phi(x)Y_{M_{2}}(u,x_{1})\right)
$$

\n
$$
=x_{1}^{-1}\delta\left(\frac{x+z}{x_{1}}\right)\left((x_{1}-x)^{k}x_{1}^{\frac{r}{T}}Y_{M_{3}}(u,x_{1})\phi(x)\right)
$$

\n
$$
=x_{1}^{-1}\delta\left(\frac{x+z}{x_{1}}\right)\left((x_{1}-x)^{k}x_{1}^{\frac{r}{T}}Y_{M_{3}}(u,x_{1})\phi(x)\right)|_{x_{1}=x+z}
$$

\n
$$
=x_{1}^{-1}\delta\left(\frac{x+z}{x_{1}}\right)z^{k}(x+z)^{\frac{r}{T}}Y_{\mathcal{H}}(u,z)\phi(x),
$$

which is equivalent to [\(4.7\)](#page-13-1).

Note that for any $u \in V$, we have $u = u^0 + u^1 + \cdots + u^{T-1}$, where for $0 \le r \le T-1$,

$$
u^{r} = \frac{1}{T} \sum_{j=0}^{T-1} \eta_{T}^{-jr} \tau^{j}(u) \in V^{r}.
$$

Using this and Lemma [4.3,](#page-13-2) we immediately get the the following generalization:

Proposition 4.4. Let $u \in V$ and let $\phi(x) \in \mathcal{H}(M_2, M_3)$. Then

$$
z^{k}Y_{\mathcal{H}}(u,z)\phi(x) = ((x_1 - x)^{k}Y_{M_3}(u,x_1)\phi(x))\big|_{x_1^{1/T} = (x+z)^{1/T}},
$$
\n(4.8)

where k is any nonnegative integer such that (4.1) holds. Furthermore, we have

$$
\sum_{j=0}^{T-1} x_1^{-1} \delta \left(\eta_T^{-j} \left(\frac{x+z}{x_1} \right)^{\frac{1}{T}} \right) Y_{\mathcal{H}}(\tau^j u, z) \phi(x)
$$

= $z^{-1} \delta \left(\frac{x_1 - x}{z} \right) Y_{M_3}(u, x_1) \phi(x) - z^{-1} \delta \left(\frac{x - x_1}{-z} \right) \phi(x) Y_{M_2}(u, x_1).$ (4.9)

We also have the following generalization of the first part of Lemma [4.3](#page-13-2) (cf. [\[Li5\]](#page-37-5), Lemma 2.15):

Lemma 4.5. Let $v \in V$ and let $\phi_1(x), \dots, \phi_p(x) \in \mathcal{H}(M_2, M_3)$. If

$$
\sum_{j=1}^p f_j(x_1 - x) Y_{M_3}(v, x_1) \phi_j(x) = \sum_{j=1}^p f_j(x_1 - x) \phi_j(x) Y_{M_2}(v, x_1),
$$

where $f_1(x), \ldots, f_p(x) \in \mathbb{C}[x]$, then

$$
\left(\sum_{j=1}^p f_j(x_1-x)Y_{M_3}(v,x_1)\phi_j(x)\right)|_{x_1^{1/T}=(x+z)^{1/T}} = \sum_{j=1}^p f_j(z)Y_{\mathcal{H}}(v,z)\phi_j(x). \tag{4.10}
$$

On the other hand, let $v^1, \dots, v^p \in V$ and let $\phi(x) \in \mathcal{H}(M_2, M_3)$. If

$$
\sum_{j=1}^p f_j(x_1-x)Y_{M_3}(v^j,x_1)\phi(x) = \sum_{j=1}^p f_j(x_1-x)\phi(x)Y_{M_2}(v^j,x_1),
$$

where $f_1(x), \ldots, f_p(x) \in \mathbb{C}[x]$, then

$$
\left(\sum_{j=1}^{p} f_j(x_1 - x) Y_{M_3}(v^j, x_1) \phi(x)\right) \Big|_{x_1^{1/T} = (x+z)^{1/T}} = \sum_{j=1}^{p} f_j(z) Y_{\mathcal{H}}(v^j, z) \phi(x). \tag{4.11}
$$

 \Box

Proof. As $\phi_j(x) \in \mathcal{H}(M_2, M_3)$, there is a nonnegative integer k such that

$$
(x_1 - x)^k Y_{M_3}(v, x_1)\phi_j(x) = (x_1 - x)^k \phi_j(x) Y_{M_2}(v, x_1)
$$

for all $1 \leq j \leq p$. Then

$$
(x_1-x)^k f_j(x_1-x)Y_{M_3}(v,x_1)\phi_j(x)=(x_1-x)^k f_j(x_1-x)\phi_j(x)Y_{M_2}(v,x_1).
$$

In view of Lemma [4.3,](#page-13-2) for $1 \le j \le p$, we have

$$
((x_1 - x)^k Y_{M_3}(v, x_1)\phi_j(x))\big|_{x_1^{1/T} = (x+z)^{1/T}} = z^k Y_{\mathcal{H}}(v, z)\phi_j(x),
$$

so that

$$
((x_1 - x)^k f_j(x_1 - x) Y_{M_3}(v, x_1) \phi_j(x))\big|_{x_1^{1/T} = (x+z)^{1/T}}
$$

= $f_j(x_1 - x)|_{x_1 = x+z} \cdot ((x_1 - x)^k Y_{M_3}(v, x_1) \phi_j(x))\big|_{x_1^{1/T} = (x+z)^{1/T}}$
= $z^k f_j(z) Y_{\mathcal{H}}(v, z) \phi_j(x).$

Then

$$
z^{k}\left(\sum_{j=1}^{p}f_{j}(x_{1}-x)Y_{M_{3}}(v,x_{1})\phi_{j}(x)\right)|_{x_{1}^{1/T}=(x+z)^{1/T}}
$$
\n
$$
=\left((x_{1}-x)^{k}\sum_{j=1}^{p}f_{j}(x_{1}-x)Y_{M_{3}}(v,x_{1})\phi_{j}(x)\right)|_{x_{1}^{1/T}=(x+z)^{1/T}}
$$
\n
$$
=\sum_{j=1}^{p}\left((x_{1}-x)^{k}f_{j}(x_{1}-x)Y_{M_{3}}(v,x_{1})\phi_{j}(x)\right)|_{x_{1}^{1/T}=(x+z)^{1/T}}
$$
\n
$$
=z^{k}\sum_{j=1}^{p}f_{j}(z)Y_{\mathcal{H}}(v,z)\phi_{j}(x),
$$

which amounts to [\(4.10\)](#page-14-0). The second assertion can be proved similarly.

$$
\Box
$$

As the first main result of this section, we have:

Theorem 4.6. Let $\tau \in \text{Aut}(V)$ with $o(\tau) = T$ and let M_2 and M_3 be weak τ -twisted V-modules. Then $(\mathcal{H}(M_2, M_3), Y_{\mathcal{H}})$ carries the structure of a weak V -module.

Proof. First, we prove that $\mathcal{H}(M_2, M_3)$ is closed under the action of V given by [\(4.4\)](#page-13-0), i.e.,

$$
v_n^{\mathcal{H}}\phi(x) \in \mathcal{H}(M_2, M_3) \text{ for } v \in V, n \in \mathbb{Z}, \ \phi(x) \in \mathcal{H}(M_2, M_3).
$$

Assume $v \in V^r$ with $r \in \mathbb{Z}$. As it was pointed out before, $v_n^{\mathcal{H}}\phi(x)$ satisfies condition (G1) for every $n \in \mathbb{Z}$. We now establish condition (G2). Using (G2) for $\phi(x)$ and the corresponding property for $Y_{M_3}(v, x_1)$, we have

$$
[L(-1), Y_{M_3}(v, x_1)\phi(x)] = \left(\frac{\partial}{\partial x_1} + \frac{\partial}{\partial x}\right) Y_{M_3}(v, x_1)\phi(x),
$$

$$
[L(-1), \phi(x)Y_{M_2}(v, x_1)] = \left(\frac{\partial}{\partial x_1} + \frac{\partial}{\partial x}\right)\phi(x)Y_{M_2}(v, x_1).
$$

On the other hand, notice that

$$
\left(\frac{\partial}{\partial x_1} + \frac{\partial}{\partial x}\right) z^{-1} \delta\left(\frac{x_1 - x}{z}\right) = 0, \quad \left(\frac{\partial}{\partial x_1} + \frac{\partial}{\partial x}\right) z^{-1} \delta\left(\frac{x - x_1}{-z}\right) = 0,
$$

$$
\left(\frac{\partial}{\partial x_1} + \frac{\partial}{\partial x}\right) \left(x_1^{-1} \delta\left(\frac{x + z}{x_1}\right) \left(\frac{x + z}{x_1}\right)^{\frac{T}{T}}\right) = 0.
$$

Using these properties and [\(4.7\)](#page-13-1), we get

$$
x_1^{-1}\delta\left(\frac{x+z}{x_1}\right)\left(\frac{x+z}{x_1}\right)^{\frac{x}{T}}[L(-1), Y_{\mathcal{H}}(v, z)\phi(x)]
$$

\n
$$
= z^{-1}\delta\left(\frac{x_1-x}{z}\right)[L(-1), Y_{M_3}(v, x_1)\phi(x)] - z^{-1}\delta\left(\frac{x-x_1}{-z}\right)[L(-1), \phi(x)Y_{M_2}(v, x_1)]
$$

\n
$$
= \left(\frac{\partial}{\partial x_1} + \frac{\partial}{\partial x}\right)\left(z^{-1}\delta\left(\frac{x_1-x}{z}\right)Y_{M_3}(v, x_1)\phi(x) - z^{-1}\delta\left(\frac{x-x_1}{-z}\right)\phi(x)Y_{M_2}(v, x_1)\right)
$$

\n
$$
= \left(\frac{\partial}{\partial x_1} + \frac{\partial}{\partial x}\right)\left(x_1^{-1}\delta\left(\frac{x+z}{x_1}\right)\left(\frac{x+z}{x_1}\right)^{\frac{x}{T}}Y_{\mathcal{H}}(v, z)\phi(x)\right)
$$

\n
$$
= x_1^{-1}\delta\left(\frac{x+z}{x_1}\right)\left(\frac{x+z}{x_1}\right)^{\frac{x}{T}}\left(\frac{\partial}{\partial x_1} + \frac{\partial}{\partial x}\right)(Y_{\mathcal{H}}(v, z)\phi(x))
$$

\n
$$
= x_1^{-1}\delta\left(\frac{x+z}{x_1}\right)\left(\frac{x+z}{x_1}\right)^{\frac{x}{T}}\frac{\partial}{\partial x}(Y_{\mathcal{H}}(v, z)\phi(x)).
$$
 (4.12)

As $Y_H(v, z) \phi(x)$ involves only finitely many negative powers of z, by cancellation we obtain

$$
[L(-1), Y_{\mathcal{H}}(v, z)\phi(x)] = \frac{\partial}{\partial x}Y_{\mathcal{H}}(v, z)\phi(x).
$$

This implies that $v_n^{\mathcal{H}} \phi(x)$ satisfy condition (G2) for all $n \in \mathbb{Z}$.

Next, we consider condition (G3). Let $u \in V$. Then there exists a positive integer k such that

$$
(x_1 - x_2)^k Y_{M_i}(u, x_1) Y_{M_i}(v, x_2) = (x_1 - x_2)^k Y_{M_i}(v, x_2) Y_{M_i}(u, x_1),
$$

$$
(x_1 - x_2)^k Y_{M_3}(u, x_1) \phi(x_2) = (x_1 - x_2)^k \phi(x_2) Y_{M_2}(u, x_1)
$$

for $i = 2, 3$. Using these and [\(4.7\)](#page-13-1) we get

$$
x_1^{-1}\delta\left(\frac{x+z}{x_1}\right)\left(\frac{x+z}{x_1}\right)^{\frac{r}{T}}(y-x)^k(y-x-z)^kY_{M_3}(u,y)(Y_{\mathcal{H}}(v,z)\phi(x))
$$

\n
$$
= x_1^{-1}\delta\left(\frac{x+z}{x_1}\right)\left(\frac{x+z}{x_1}\right)^{\frac{r}{T}}(y-x)^k(y-x_1)^kY_{M_3}(u,y)(Y_{\mathcal{H}}(v,z)\phi(x))
$$

\n
$$
= x_1^{-1}\delta\left(\frac{x+z}{x_1}\right)\left(\frac{x+z}{x_1}\right)^{\frac{r}{T}}(y-x)^k(y-x_1)^k(Y_{\mathcal{H}}(v,z)\phi(x))Y_{M_2}(u,y)
$$

\n
$$
= x_1^{-1}\delta\left(\frac{x+z}{x_1}\right)\left(\frac{x+z}{x_1}\right)^{\frac{r}{T}}(y-x)^k(y-x-z)^k(Y_{\mathcal{H}}(v,z)\phi(x))Y_{M_2}(u,y).
$$

Again, as $(Y_H(v, z)\phi(x))$ involves only finitely many negative powers of z, by cancellation we obtain

$$
(y-x)^{k}(y-x-z)^{k}Y_{M_{3}}(u,y)(Y_{\mathcal{H}}(v,z)\phi(x)) = (y-x)^{k}(y-x-z)^{k}(Y_{\mathcal{H}}(v,z)\phi(x))Y_{M_{2}}(u,y).
$$
 (4.13)

Let $n_0 \in \mathbb{Z}$ such that $v_n^{\mathcal{H}} \phi(x) = 0$ for all $n \ge n_0$. For $m \in \mathbb{N}$, applying $\text{Res}_z z^{n_0-m}$ to both sides we get

$$
\sum_{j=0}^{k} {k \choose j} (-1)^{j} (y-x)^{2k-j} Y_{M_3}(u, y) (v_{n_0-m+j}^{\mathcal{H}} \phi(x))
$$

=
$$
\sum_{j=0}^{k} {k \choose j} (-1)^{j} (y-x)^{2k-j} (v_{n_0-m+j}^{\mathcal{H}} \phi(x)) Y_{M_3}(u, y).
$$
 (4.14)

It then follows from this identity and induction on m that for every $m \in \mathbb{N}$, there exists a positive integer p such that

$$
(y-x)^p Y_{M_3}(u, y) (v_{n_0-m}^{\mathcal{H}} \phi(x)) = (y-x)^p (v_{n_0-m}^{\mathcal{H}} \phi(x)) Y_{M_3}(u, y).
$$

(Note that if $k = 0$, it is clear, and if $k \ge 1$, [\(4.14\)](#page-17-0) gives a recursion formula.) Thus we have proved $v_n^{\mathcal{H}}\phi(x) \in$ $\mathcal{H}(M_2, M_3)$ for $n \in \mathbb{Z}$. Now, we have proved that $Y_{\mathcal{H}}(\cdot, z)$ is a linear map from V to $(\text{End}(\mathcal{H}(M_2, M_3)))[[z, z^{-1}]]$ such that $Y_{\mathcal{H}}(u, z)\phi(x) \in \mathcal{H}(M_2, M_3)((z))$ for $u \in V$, $\phi(x) \in \mathcal{H}(M_2, M_3)$. It can be readily seen from [\(4.4\)](#page-13-0) that $Y_{\mathcal{H}}(1, z) = 1.$

To prove that $\mathcal{H}(M_2, M_3)$ is a weak V-module, it suffices to prove the weak associativity. Let $u \in V^r$, $v \in V^s$ with $r, s \in \mathbb{Z}$ and let $\phi(x) \in \mathcal{H}(M_2, M_3)$. Then there exists a positive integer n such that

$$
(x_1 - x)^n Y_{M_3}(u, x_1) \phi(x) = (x_1 - x)^n \phi(x) Y_{M_2}(u, x_1),
$$

\n
$$
(x_1 - x)^n Y_{M_3}(v, x_1) \phi(x) = (x_1 - x)^n \phi(x) Y_{M_2}(v, x_1),
$$

\n
$$
(x_1 - x)^n Y_{M_3}(u, x_1) Y_{M_3}(v, x) = (x_1 - x)^n Y_{M_3}(v, x) Y_{M_3}(u, x_1).
$$

From the twisted Jacobi identity for (M_3, Y_{M_3}) we have

$$
(y_2 + z_0)^{\frac{r}{T}} Y_{M_3}(Y(u, z_0)v, y_2)
$$

= Res_{y₁} $y_1^{\frac{r}{T}} \left(x_0^{-1} \delta \left(\frac{y_1 - y_2}{z_0} \right) Y_{M_3}(u, y_1) Y_{M_3}(v, y_2) - x_0^{-1} \delta \left(\frac{y_2 - y_1}{z_0} \right) Y_{M_3}(v, y_2) Y_{M_3}(u, y_1) \right).$

Using these properties and the standard delta-function substitutions, we get

$$
(y_2 + z_0)^{\frac{r}{T}} (y_2 - x)^n (y_2 - x + z_0)^n Y_{M_3}(Y(u, z_0)v, y_2) \phi(x)
$$

= $(y_2 + z_0)^{\frac{r}{T}} (y_2 - x)^n (y_2 - x + z_0)^n \phi(x) Y_{M_2}(Y(u, z_0)v, y_2).$

Multiplying both sides by $(y_2 + z_0)^{-\frac{r}{T}}$ (noticing that we are allowed to do so since $Y(u, z_0)v \in V((z_0))$), we obtain

$$
(y_2 - x)^n (y_2 - x + z_0)^n Y_{M_3}(Y(u, z_0)v, y_2) \phi(x)
$$

=
$$
(y_2 - x)^n (y_2 - x + z_0)^n \phi(x) Y_{M_2}(Y(u, z_0)v, y_2).
$$
 (4.15)

Noticing that $u_m v \in V^{r+s}$ for $m \in \mathbb{Z}$, using Lemma [4.5](#page-14-1) we obtain

$$
z_2^n(z_2 + z_0)^n(x + z_2)^{\frac{r+s}{T}} Y_{\mathcal{H}}(Y(u, z_0)v, z_2) \phi(x)
$$

=
$$
\left((x_2 - x)^n (x_2 - x + z_0)^n x_2^{\frac{r+s}{T}} Y_{M_3}(Y(u, z_0)v, x_2) \phi(x) \right) |_{x_2 = x + z_2}.
$$
 (4.16)

On the other hand, we have

$$
(x_2 + z_0)^{\frac{r}{T}} z_0^{n} Y_{M_3}(Y(u, z_0)v, x_2) = ((x_1 - x_2)^n x_1^{\frac{r}{T}} Y_{M_3}(u, x_1) Y_{M_3}(v, x_2)) |_{x_1 = x_2 + z_0},
$$

which gives

$$
z_0^n Y_{M_3}(Y(u,z_0)v,x_2)=(x_2+z_0)^{-\frac{r}{T}}\left((x_1-x_2)^n x_1^{\frac{r}{T}} Y_{M_3}(u,x_1)Y_{M_3}(v,x_2)\right)|_{x_1=x_2+z_0}.
$$

Then

$$
z_{2}^{n}z_{0}^{n}(z_{2}+z_{0})^{n}(x+z_{2})^{\frac{r+s}{T}}Y_{\mathcal{H}}(Y(u,z_{0})v,z_{2})\phi(x)
$$

=
$$
\left((x_{2}-x)^{n}(x_{2}-x+z_{0})^{n}x_{2}^{\frac{r+s}{T}}z_{0}^{n}Y_{M_{3}}(Y(u,z_{0})v,x_{2})\phi(x)\right)|_{x_{2}=x+z_{2}}
$$

=
$$
\left((x_{2}-x)^{n}(x_{2}-x+z_{0})^{n}x_{2}^{\frac{r+s}{T}}(x_{2}+z_{0})^{-\frac{r}{T}}\left((x_{1}-x_{2})^{n}x_{1}^{\frac{r}{T}}Y_{M_{3}}(u,x_{1})Y_{M_{3}}(v,x_{2})\phi(x)\right)|_{x_{1}=x_{2}+z_{0}}\right)|_{x_{2}=x+z_{2}}
$$

=
$$
\left((1+z_{0}/x_{2})^{-\frac{r}{T}}\left((x_{2}-x)^{n}(x_{1}-x)^{n}(x_{1}-x_{2})^{n}x_{1}^{\frac{r}{T}}x_{2}^{\frac{s}{T}}Y_{M_{3}}(u,x_{1})Y_{M_{3}}(v,x_{2})\phi(x)\right)|_{x_{1}=x_{2}+z_{0}}\right)|_{x_{2}=x+z_{2}}
$$

=
$$
\left(1+\frac{z_{0}}{x+z_{2}}\right)^{-\frac{r}{T}}\left(\left((x_{2}-x)^{n}(x_{1}-x)^{n}(x_{1}-x_{2})^{n}x_{1}^{\frac{r}{T}}x_{2}^{\frac{s}{T}}Y_{M_{3}}(u,x_{1})Y_{M_{3}}(v,x_{2})\phi(x)\right)|_{x_{1}=x_{2}+z_{0}}\right)|_{x_{2}=x+z_{2}},
$$

which gives

$$
z_2^n z_0^n (z_2 + z_0)^n (x + z_2)^{\frac{s}{T}} (x + z_2 + z_0)^{\frac{r}{T}} Y_{\mathcal{H}}(Y(u, z_0)v, z_2) \phi(x)
$$

=
$$
\left(\left((x_2 - x)^n (x_1 - x)^n (x_1 - x_2)^n x_1^{\frac{r}{T}} x_2^{\frac{s}{T}} Y_{M_3}(u, x_1) Y_{M_3}(v, x_2) \phi(x) \right) |_{x_1 = x_2 + z_0} \right) |_{x_2 = x + z_2}.
$$
 (4.17)

On the other hand, we consider $Y_{\mathcal{H}}(u, z_1)Y_{\mathcal{H}}(v, z_2)\phi(x)$. With [\(4.13\)](#page-16-0), by Lemma [4.5](#page-14-1) we have

$$
z_1^k (z_1 - z_2)^k (x + z_1)^{\frac{r}{T}} Y_{\mathcal{H}}(u, z_1) Y_{\mathcal{H}}(v, z_2) \phi(x)
$$

=
$$
\left((x_1 - x)^k (x_1 - x - z_2)^k x_1^{\frac{r}{T}} Y_{M_3}(u, x_1) (Y_{\mathcal{H}}(v, z_2) \phi(x)) \right) |_{x_1 = x + z_1}.
$$

We also have

$$
z_2^k(x+z_2)^{\frac{s}{T}} Y_{\mathcal{H}}(v,z_2)\phi(x) = \left((x_2-x)^k x_2^{\frac{s}{T}} Y_{M_3}(v,x_2)\phi(x) \right)|_{x_2=x+z_2}.
$$

Then we get

$$
z_2^k(x + z_2)^{\frac{s}{T}} z_1^k(z_1 - z_2)^k (x + z_1)^{\frac{r}{T}} Y_{\mathcal{H}}(u, z_1) Y_{\mathcal{H}}(v, z_2) \phi(x)
$$

=
$$
\left((x_1 - x)^k (x_1 - x - z_2)^k x_1^{\frac{r}{T}} Y_{M_3}(u, x_1) \left((x_2 - x)^k x_2^{\frac{s}{T}} Y_{M_3}(v, x_2) \phi(x) \right) |_{x_2 = x + z_2} \right) |_{x_1 = x + z_1}
$$

=
$$
\left(\left((x_1 - x)^k (x_1 - x_2)^k (x_2 - x)^k x_1^{\frac{r}{T}} x_2^{\frac{s}{T}} Y_{M_3}(u, x_1) Y_{M_3}(v, x_2) \phi(x) \right) |_{x_2 = x + z_2} \right) |_{x_1 = x + z_1},
$$

which gives

$$
z_2^k(x+z_2)^{\frac{s}{T}}(z_2+z_0)^k z_0^k(x+z_2+z_0)^{\frac{r}{T}} Y_{\mathcal{H}}(u,z_2+z_0) Y_{\mathcal{H}}(v,z_2) \phi(x)
$$

=
$$
\left(\left((x_1-x)^k (x_1-x_2)^k (x_2-x)^k x_1^{\frac{r}{T}} x_2^{\frac{s}{T}} Y_{M_3}(u,x_1) Y_{M_3}(v,x_2) \phi(x) \right) |_{x_2=x+z_2} \right) |_{x_1=x+z_2+z_0}.
$$

Combining this with [\(4.17\)](#page-18-0) (replacing *n* with a larger one so that $n \ge k$) we obtain

$$
z_2^n z_0^n (z_2 + z_0)^n (x + z_2)^{\frac{x}{T}} (x + z_2 + z_0)^{\frac{r}{T}} Y_\mathcal{H}(Y(u, z_0)v, z_2) \phi(x)
$$

=
$$
z_2^n (x + z_2)^{\frac{x}{T}} (z_2 + z_0)^n z_0^n (x + z_2 + z_0)^{\frac{r}{T}} Y_\mathcal{H}(u, z_2 + z_0) Y_\mathcal{H}(v, z_2) \phi(x),
$$

noticing that under the formal series expansion convention, for any $m \in \mathbb{Z}$,

$$
(x_2 + z_0)^m |_{x_2 = x + z_2} = ((x + z_2) + z_0)^m = (x + (z_2 + z_0))^m = x_1^m |_{x_1 = x + z_2 + z_0}.
$$

Multiplying both sides by $z_2^{-n} z_0^{-n} (x + z_2)^{-\frac{s}{T}} (x + z_2 + z_0)^{-\frac{r}{T}}$ we get

$$
(z_2 + z_0)^n Y_{\mathcal{H}}(Y(u, z_0)v, z_2)\phi(x) = (z_2 + z_0)^n Y_{\mathcal{H}}(u, z_2 + z_0)Y_{\mathcal{H}}(v, z_2)\phi(x).
$$

This establishes the weak associativity. Therefore, $(\mathcal{H}(M_2, M_3), Y_\mathcal{H})$ carries the structure of a weak V-module. \Box We end this section with the following technical result:

Lemma 4.7. Let τ , T, M_2 , M_3 be given as before and let $\phi(x) \in (Hom(M_2, M_3))\{z\}$. Suppose that $\phi(x)$ satisfies the conditions (G1) and (G2) and suppose that the condition (G3) holds for any $u \in U$, where U is a subset of V such that $\bigcup_{j=0}^{k-1} \tau^j(U)$ generates V as a vertex algebra. Then $\phi(x) \in \mathcal{H}(M_2, M_3)$.

Proof. Let K consist of all vectors $a \in V$ such that

$$
(x_1 - x)^n Y_{M_3}(a, x_1)\phi(x) = (x_1 - x)^n \phi(x) Y_{M_2}(a, x_1)
$$

for some nonnegative integer n. We must prove $K = V$. As $\cup_{j=0}^{T-1} \tau^j(U)$ generates V as a vertex algebra, it suffices to prove that K is a vertex subalgebra that contains $\bigcup_{j=0}^{T-1} \tau^j(U)$. It is clear that K is a subspace with $1 \in K$, and from assumption we have $U \subset K$. Note that for any $v \in V$, we have

$$
Y_{M_i}(\tau(v), z) = \lim_{z^{1/T} \to \eta_T^{-1} z^{1/T}} Y_{M_i}(v, z)
$$

with $i = 2, 3$. It then follows that $\tau(K) \subset K$. Thus $\cup_{j=0}^{T-1} \tau^{j}(U) \subset K$. Now, it remains to prove that $u_m v \in K$ for any $u, v \in K$, $m \in \mathbb{Z}$. As $\tau(K) \subset K$, it suffices to consider $u \in K \cap V^r$ with $r \in \mathbb{Z}$. Note that

$$
(z_2 + z_0)^{\frac{r}{T}} Y_{M_i}(Y(u, z_0)v, z_2)
$$

= Res_{z₁} $z_1^{\frac{r}{T}} \left(z_0^{-1} \delta \left(\frac{z_1 - z_2}{z_0} \right) Y_{M_i}(u, z_1) Y_{M_i}(v, z_2) - z_0^{-1} \delta \left(\frac{z_2 - z_1}{-z_0} \right) Y_{M_i}(v, z_2) Y_{M_i}(u, z_1) \right)$ (4.18)

holds on M_i for $i = 2, 3$. Let k be a positive integer such that

$$
(z_1 - z)^k Y_{M_3}(a, z_1)\phi(z) = (z_1 - z)^k \phi(z) Y_{M_2}(a, z_1)
$$

for $a \in \{u, v\}$. Using these relations and [\(4.18\)](#page-19-0), we get

$$
(z_2-z)^k(z_2+z_0-z)^k Y_{M_3}(Y(u,z_0)v,z_2)\phi(z)=(z_2-z)^k(z_2+z_0-z)^k\phi(z)Y_{M_2}(Y(u,z_0)v,z_2).
$$

Just as in the first part of the proof of Theorem [4.6,](#page-15-0) it follows from this relation and an induction that $u_m v \in K$ for all $m \in \mathbb{Z}$. Therefore, we conclude $K = V$ and hence $\phi(x) \in \mathcal{H}(M_2, M_3)$. □

Proposition 4.8. Let M_2 , M_3 be weak τ -twisted V-modules and let M be any weak V-module. Then for any intertwining operator $I(\cdot, x)$ of type $\binom{M_3}{M M_2}$, we have $I(w, x) \in \mathcal{H}(M_2, M_3)$ for $w \in M$, and the map f_I defined by $f_I(w) = I(w, x)$ for $w \in M$ is a homomorphism of weak V-modules. Furthermore, the linear map $\theta : I_V \begin{pmatrix} M_3 \\ M_M \end{pmatrix} \to$ $\text{Hom}_V(M, \mathcal{H}(M_2, M_3))$ defined by $\theta(I) = f_I$ is a linear isomorphism.

Proof. Let $I(\cdot, x)$ be any intertwining operator of type $\binom{M_3}{M M_2}$. From definition, we see that for every $w \in M$, $I(w, x)$ satisfies conditions (G1) and (G2). As the Jacobi identity for $I(\cdot, x)$ implies commutator formula which furthermore implies locality, $I(w, x)$ also satisfies (G3). Thus $I(w, x) \in \mathcal{H}(M_2, M_3)$. Then we have a linear map f_I from M to $\mathcal{H}(M_2, M_3)$ defined by $f_I(w) = I(w, x)$ for $w \in M$. For any $u \in V^r$, $w \in M$ with $r \in \mathbb{Z}$, we have

$$
f_I(Y(u, x_0)w)
$$

= $I(Y(u, x_0)w, x)$
= Res_{x₁} $\left(\frac{x + x_0}{x_1}\right)^{-\frac{r}{T}} \left(x_0^{-1}\delta\left(\frac{x_1 - x}{x_0}\right)Y_{M_3}(u, x_1)I(w, x) - x_0^{-1}\delta\left(\frac{x - x_1}{-x_0}\right)I(w, x)Y_{M_2}(u, x_1)\right)$
= $Y_{\mathcal{H}}(u, x_0)I(w, x)$
= $Y_{\mathcal{H}}(u, x_0)f_I(w).$

This proves that f_I is a homomorphism of weak V-modules.

For the second assertion, it is clear that θ is a one-to-one linear map. To prove that it is onto, let f be a Vhomomorphism from M to $\mathcal{H}(M_2, M_3)$. For every $w \in M$, noticing that as an element of $\mathcal{H}(M_2, M_3)$, $f(w)$ is a formal series in a formal variable, we write $f(w)$ as $f(w)(x)$, and then set $I_f(w, x) = f(w)(x) \in \mathcal{H}(M_2, M_3)$. This gives a linear map $I_f(\cdot, x)$ from M to $(Hom_{\mathbb{C}}(M_2, M_3))\{x\}$. We have

$$
[L(-1), I_f(w, z)] = \frac{d}{dz}I_f(w, z).
$$

For any $v \in V$, $w \in M$, as $f(w)(x) \in H(M_2, M_3)$, by condition (G3) there exists a nonnegative integer k such that

$$
(x_1-x)^k Y_{M_3}(v,x_1) f(w)(x) = (x_1-x)^k f(w)(x) Y_{M_2}(v,x_1),
$$

which amounts to

$$
(x_1 - x)^k Y_{M_3}(v, x_1) I_f(w, x) = (x_1 - x)^k I_f(w, x) Y_{M_2}(v, x_1).
$$
\n(4.19)

On the other hand, for $v \in V^r$ with $r \in \mathbb{Z}$ and $w \in M$, with f a homomorphism of weak V-modules, we have

$$
(z_2 + z_0)^{\frac{r}{T}} I_f(Y_M(v, z_0)w, z_2)
$$

= $(z_2 + z_0)^{\frac{r}{T}} f(Y_M(v, z_0)w)(z_2)$
= $(z_2 + z_0)^{\frac{r}{T}} Y_H(v, z_0) f(w)(z_2)$
= Res_{z₁} $z_1^{\frac{r}{T}} \left(z_0^{-1} \delta \left(\frac{z_1 - z_2}{z_0} \right) Y_{M_3}(v, z_1) f(w)(z_2) - z_0^{-1} \delta \left(\frac{z_2 - z_1}{z_0} \right) f(w)(z_2) Y_{M_2}(v, z_1) \right)$
= Res_{z₁} $z_1^{\frac{r}{T}} \left(z_0^{-1} \delta \left(\frac{z_1 - z_2}{z_0} \right) Y_{M_3}(v, z_1) I_f(w, z_2) - z_0^{-1} \delta \left(\frac{z_2 - z_1}{z_0} \right) I_f(w, z_2) Y_{M_2}(v, z_1) \right).$

This together with [\(4.19\)](#page-20-0) gives the Jacobi identity. Therefore, $I_f(\cdot, x)$ is an intertwining operator of type $\binom{M_3}{M M_2}$. It is clear that $\theta(I_f) = f$. This proves that θ is onto, concluding the proof.

5 Permutation orbifolds

In this section, we shall apply the results of Section [4](#page-12-2) to the permutation orbifold model.

5.1 Linear isomorphism between $I_V \begin{pmatrix} W \\ M N \end{pmatrix}$ and $I_{V^{\otimes k}} \begin{pmatrix} T_{\sigma}(W) \\ M^1 T_{\sigma}(N) \end{pmatrix}$ $T_{\sigma}(W)$
 $M^1 T_{\sigma}(N)$

Let V be a vertex operator algebra and let k be a positive integer with $k \geq 2$, which are all fixed throughout this section. Set $\sigma = (1\ 2\ \cdots k)$, which is an automorphism of $V^{\otimes k}$. Let M, N, and W be irreducible V-modules. Set $M^1 = M \otimes V^{\otimes (k-1)}$, which is a $V^{\otimes k}$ -module. Let $T_{\sigma}(N)$ and $T_{\sigma}(W)$ be the σ -twisted $V^{\otimes k}$ -modules constructed in [\[BDM\]](#page-35-2). In this section, we prove that there is a canonical linear isomorphism between the spaces $I_V \binom{W}{M N}$ and $I_{V^{\otimes k}}\binom{T_{\sigma}(W)}{M^1 T_{\sigma}(N)}$. That is, given an intertwining operator of type $\binom{W}{M N}$, we can construct an intertwining operator of type $\binom{T_{\sigma}(W)}{M^1 T_{\sigma}(N)}$. Conversely, given an intertwining operator of type $\binom{T_{\sigma}(W)}{M^1 T_{\sigma}(N)}$, we can construct an intertwining operator of type $\binom{W}{M N}$. As one of the main results, we obtain the following tensor product relation:

$$
(M \otimes V^{\otimes (k-1)}) \boxtimes_{V^{\otimes k}} T_{\sigma}(N) \simeq T_{\sigma}(M \boxtimes_V N).
$$

5.1.1 From $I_V {W \choose M N}$ to $I_{V^{\otimes k}} {T_{\sigma}(W) \choose M^1 \ T_{\sigma}(N)}$

For any $u \in V$, $1 \le j \le k$, we denote by u^j the vector in $V^{\otimes k}$ whose j^{th} tensor factor is u and the other tensor factors are 1:

$$
u^{j} = \mathbf{1}^{\otimes (j-1)} \otimes u \otimes \mathbf{1}^{\otimes (k-j)}.
$$
\n
$$
(5.1)
$$

We have

$$
Y(u^{j}, z) = 1^{\otimes (j-1)} \otimes Y(u, z) \otimes 1^{\otimes (k-j)}.
$$
\n(5.2)

Note that $\sigma u^j = u^{j+1}$ for $1 \le j \le k$, where $u^{k+1} = u^1$ by convention. For any weak $V^{\otimes k}$ -module (W, Y_W) , we have

$$
\left[Y_W(u^i, z_1), Y_W(v^j, z_2)\right] = \text{Res}_{z_0} z_2^{-1} \delta\left(\frac{z_1 - z_0}{z_2}\right) Y_W(Y(u^i, z_0)v^j, z_2)
$$

for $u, v \in V$, $1 \le i, j \le k$. For $n \ge 0$, as $u_n \mathbf{1} = 0$ (in V) we have $(u^i)_n(v^j) = 0$ (in $V^{\otimes k}$) if $i \ne j$. Then we immediately have:

Lemma 5.1. Let (W, Y_W) be any weak $V^{\otimes k}$ -module. Then $[Y_W(u^i, z_1), Y_W(v^j, z_2)] = 0$ for any $u, v \in V, i, j \in$ $\{1, \ldots, k\}$ with $i \neq j$.

Let k be a positive integer. Recall from [\[BDM\]](#page-35-2)

$$
\Delta_k(z) = \exp\left(\sum_{n\geq 1} a_n z^{-\frac{n}{k}} L(n)\right) k^{-L(0)} z^{(1/k-1)L(0)},\tag{5.3}
$$

where the coefficients a_n for $n \geq 1$ are uniquely determined by

$$
\exp\left(\sum_{n\geq 1} -a_n x^{n+1} \frac{d}{dx}\right) x = \frac{1}{k} (1+x)^k - \frac{1}{k}.\tag{5.4}
$$

As in [\[BDM\]](#page-35-2), we shall also use the expression $\Delta_k(z^k)^{-1}$. For convenience, we set

$$
\Phi_k(z) = \Delta_k(z^k)^{-1} = (kz^{k-1})^{L(0)} \exp\left(-\sum_{n\geq 1} a_n z^{-n} L(n)\right).
$$
\n(5.5)

The following results were obtained in [\[BDM\]](#page-35-2) (Proposition 2.2 and Corollary 2.5):

Proposition 5.2. Let (W, Y_W) be any V-module. Then

$$
\Delta_k(x)Y_W(v,z)w = Y_W\left(\Delta_k(x+z)v, (x+z)^{1/k} - x^{1/k}\right)\Delta_k(x)w,
$$
\n(5.6)

$$
\Phi_k(x)Y_W(v, z)w = Y_W \left(\Phi_k(x+z)v, (x+z)^k - x^k\right) \Phi_k(x)w,
$$
\n
$$
\Delta_k(z)L(-1)w - \frac{1}{k}z^{1/k-1}L(-1)\Delta_k(z)w = \frac{d}{dz}\Delta_k(z)w,
$$
\n(5.8)

$$
\Delta_k(z)L(-1)w - \frac{1}{k}z^{1/k-1}L(-1)\Delta_k(z)w = \frac{1}{dz}\Delta_k(z)w,
$$
\n
$$
\Delta_k(z)\omega = \frac{1}{k^2}z^{2(\frac{1}{k}-1)}\omega + z^{-2}\frac{c}{24}(1-k^{-2})\mathbf{1}
$$
\n(5.9)

for $v \in V$, $w \in W$, where ω is the conformal vector of V and c is the central charge.

The following is one of the main results of [\[BDM\]](#page-35-2):

Theorem 5.3. Let V be any vertex operator algebra and let $\sigma = (12 \cdots k) \in \text{Aut}(V^{\otimes k})$. Then for any (weak, admissible) V-module (W, Y_W) , we have a (weak, admissible) σ -twisted $V^{\otimes k}$ -module $(T_{\sigma}(W), Y_{T_{\sigma}(W)})$, where $T_{\sigma}(W) = W$ as a vector space and the vertex operator map $Y_{T_{\sigma}(W)}(\cdot, z)$ is uniquely determined by

$$
Y_{T_{\sigma}(W)}(u^1, z) = Y_W(\Delta_k(z)u, z^{1/k}) \quad \text{for } u \in V,
$$
\n(5.10)

where $u^1 = u \otimes 1^{\otimes (k-1)} \in V^{\otimes k}$. Furthermore, every (weak, admissible) σ -twisted $V^{\otimes k}$ -module is isomorphic to one in this form.

Set $\eta_k = e^{\frac{2\pi i}{k}}$, the principal primitive k-th root of unity.

We shall need the $L(-1)$ -operator for the vertex operator algebra $V^{\otimes k}$ on the σ -twisted $V^{\otimes k}$ -module $T_{\sigma}(W)$. Note that the conformal vector of $V^{\otimes k}$ is

$$
\omega^{(k)} := \omega^1 + \cdots + \omega^k,
$$

where ω denotes the conformal vector of V. Write

$$
Y_W(\omega, z) = \sum_{n \in \mathbb{Z}} L(n) z^{-n-2}, \quad Y_{T_{\sigma}(W)}(\omega^{(k)}, z) = \sum_{n \in \mathbb{Z}} L_{T_{\sigma}}(n) z^{-n-2}.
$$
 (5.11)

Using the covariance property of σ -twisted modules and the expression of $\Delta_k(z)\omega$ from Proposition [5.2,](#page-21-0) we have

$$
Y_{T_{\sigma}(W)}(\omega^{(k)}, z) = \sum_{j=0}^{k-1} Y_{T_{\sigma}(W)}(\sigma^j \omega^1, z) = \sum_{j=0}^{k-1} \lim_{z^{1/k} \to \eta_k^{-j} z^{1/k}} Y_{T_{\sigma}(W)}(\omega^1, z)
$$

\n
$$
= \sum_{j=0}^{k-1} \lim_{z^{1/k} \to \eta_k^{-j} z^{1/k}} Y_W(\Delta_k(z)\omega, z^{1/k})
$$

\n
$$
= \sum_{j=0}^{k-1} \frac{1}{k^2} \eta_k^{-2j} z^{2/k-2} Y_W(\omega, \eta_k^{-j} z^{1/k}) + z^{-2} \frac{c}{24} k(1 - k^{-2})
$$

\n
$$
= \frac{1}{k} \sum_{m \in \mathbb{Z}} L(mk) z^{-m-2} + z^{-2} \frac{c}{24} (k - k^{-1}).
$$

Then

$$
L_{T_{\sigma}}(n) = \begin{cases} \frac{1}{k}L(nk) & \text{for } n \neq 0\\ \frac{1}{k}L(0) + \frac{c}{24}(k - k^{-1}) & \text{for } n = 0. \end{cases}
$$
(5.12)

In particular, we have

$$
L_{T_{\sigma}}(-1) = \frac{1}{k}L(-k). \tag{5.13}
$$

We have the following result which will be used later:

Lemma 5.4. Let (W, Y_W) be any V-module. Then

$$
\frac{d}{dz}\Delta_k(z) = \frac{1}{k} \sum_{i \ge 1} \binom{-k+1}{i} z^{-1 - (i-1)/k} L(i-1) \Delta_k(z) \tag{5.14}
$$

holds on W.

Proof. It is similar to that of Corollary 2.5 of [\[BDM\]](#page-35-2). As in [\[BDM\]](#page-35-2) we set

$$
\Delta_k^x(z) = \exp\left(-\sum_{j\geq 1} a_j z^{-j/k} x^{j+1} \frac{\partial}{\partial x}\right) k^x \frac{\partial}{\partial x} z^{(-1/k+1)x \frac{\partial}{\partial x}},
$$

an element of $(\text{End }\mathbb{C}[x,x^{-1}])[[z^{1/k},z^{-1/k}]]$. As in the proof of Corollary 2.5 of [\[BDM\]](#page-35-2), it suffices to show that

$$
\frac{1}{k} \sum_{i \ge 1} \binom{-k+1}{i} \left(-x^i \frac{\partial}{\partial x}\right) z^{-1-(i-1)/k} \Delta_k^x(z) = \frac{\partial}{\partial z} \Delta_k^x(z) \tag{5.15}
$$

in (End $\mathbb{C}[x,x^{-1}])[[z^{1/k},z^{-1/k}]].$

Note that

$$
\Delta_k^x(z)x = \exp\left(-\sum_{j\geq 1} a_j z^{-j/k} x^{j+1} \frac{\partial}{\partial x}\right) k^x \frac{\partial}{\partial x} z^{(-1/k+1)x \frac{\partial}{\partial x}} x
$$

\n
$$
= \exp\left(-\sum_{j\geq 1} a_j z^{-j/k} x^{j+1} \frac{\partial}{\partial x}\right) k z^{-1/k+1} x
$$

\n
$$
= kz \exp\left(-\sum_{j\geq 1} a_j z^{-j/k} x^{j+1} \frac{\partial}{\partial x}\right) z^{-1/k} x
$$

\n
$$
= kz \exp\left(-\sum_{j\geq 1} a_j (z^{-1/k} x)^{j+1} \frac{\partial}{\partial z^{-1/k} x}\right) z^{-1/k} x
$$

\n
$$
= kz \left(\frac{1}{k} (1 + z^{-1/k} x)^k - \frac{1}{k}\right)
$$

\n
$$
= z \left((1 + z^{-1/k} x)^k - 1\right),
$$

where for the second last equality we are using equation [\(5.4\)](#page-21-1). Then

$$
\frac{1}{k} \sum_{i\geq 1} {\binom{-k+1}{i}} \left(-x^i \frac{\partial}{\partial x} \right) z^{-1-(i-1)/k} \Delta_k^x(z) x
$$
\n
$$
= \frac{1}{k} \sum_{i\geq 1} {\binom{-k+1}{i}} \left(-x^i \frac{\partial}{\partial x} \right) z^{-1-(i-1)/k} z \left((1+z^{-1/k}x)^k - 1 \right)
$$
\n
$$
= -\sum_{i\geq 1} {\binom{-k+1}{i}} x^i z^{-i/k} (1+z^{-1/k}x)^{k-1}
$$
\n
$$
= -\sum_{i\geq 0} {\binom{-k+1}{i}} x^i z^{-i/k} \left((1+z^{-1/k}x)^{k-1} \right) + (1+z^{-1/k}x)^{k-1}
$$
\n
$$
= -1 + (1+z^{-1/k}x)^{k-1}
$$
\n
$$
= \frac{\partial}{\partial z} z \left((1+z^{-1/k}x)^k - 1 \right)
$$
\n
$$
= \frac{\partial}{\partial z} \Delta_k^x(z) x.
$$

From [\[BDM\]](#page-35-2) we have that $\Delta_k^x(z)x^n = (\Delta_k^x(z)x)^n$ for $n \in \mathbb{Z}$. Thus

$$
\frac{1}{k} \sum_{i\geq 1} {\binom{-k+1}{i}} \left(-x^i \frac{\partial}{\partial x} \right) z^{-1-(i-1)/k} \Delta_k^x(z) x^n
$$
\n
$$
= \frac{1}{k} \sum_{i\geq 1} {\binom{-k+1}{i}} \left(-x^i \frac{\partial}{\partial x} \right) z^{-1-(i-1)/k} z^n \left((1+z^{-1/k}x)^k - 1 \right)^n
$$
\n
$$
= -n \sum_{i\geq 1} {\binom{-k+1}{i}} x^i z^{-i/k} (1+z^{-1/k}x)^{k-1} z^{n-1} \left((1+z^{-1/k}x)^k - 1 \right)^{n-1}
$$
\n
$$
= -n \sum_{i\geq 0} {\binom{-k+1}{i}} x^i z^{-i/k} (1+z^{-1/k}x)^{k-1} z^{n-1} \left((1+z^{-1/k}x)^k - 1 \right)^{n-1}
$$
\n
$$
+ n(1+z^{-1/k}x)^{k-1} z^{n-1} \left((1+z^{-1/k}x)^k - 1 \right)^{n-1}
$$
\n
$$
= -nz^{n-1} \left((1+z^{-1/k}x)^k - 1 \right)^{n-1} + n(1+z^{-1/k}x)^{k-1} z^{n-1} \left((1+z^{-1/k}x)^k - 1 \right)^{n-1}
$$
\n
$$
= nz^{n-1} \left((1+z^{-1/k}x)^k - 1 \right)^{n-1} \left(-1 + (1+z^{-1/k}x)^{k-1} \right)
$$
\n
$$
= n(\Delta_k^x(z)x)^{n-1} \frac{\partial}{\partial z} \Delta_k^x(z)
$$
\n
$$
= \frac{\partial}{\partial z} (\Delta_k^x(z)x)^n
$$
\n
$$
= \frac{\partial}{\partial z} \Delta_k^x(z)x^n.
$$

Therefore, equation [\(5.15\)](#page-22-0) holds.

As the first main result of this section we have:

Theorem 5.5. Let M, N and W be weak V-modules and let $\sigma = (1 \ 2 \ \cdots k) \in Aut(V^{\otimes k})$. Let $\mathcal{Y}(\cdot, z)$ be an intertwining operator of type ${W \choose M N}$. For $w \in M$, set

$$
\overline{\mathcal{Y}}(w,z) = \mathcal{Y}(\Delta_k(z)w, z^{1/k}) \in (\text{Hom}(N,W))\{z\}.
$$
\n(5.16)

Then

$$
\overline{\mathcal{Y}}(w, z) \in \mathcal{H}(T_{\sigma}(N), T_{\sigma}(W)) \quad \text{for } w \in M. \tag{5.17}
$$

Furthermore, the linear map $f : M \otimes V^{\otimes (k-1)} \to \mathcal{H}(T_{\sigma}(N), T_{\sigma}(W))$, defined by

$$
f(w\otimes a)=a_{-1}^{\mathcal{H}}\overline{\mathcal{Y}}(w,z)\quad\text{for }w\in M,\;a\in V^{\otimes (k-1)},
$$

is a weak $V^{\otimes k}$ -module homomorphism from $M \otimes V^{\otimes (k-1)}$ to $\mathcal{H}(T_{\sigma}(N), T_{\sigma}(W)).$

Proof. First, we show that $\overline{\mathcal{Y}}(w, z) \in \mathcal{H}(T_{\sigma}(N), T_{\sigma}(W))$ for $w \in M$. It is clear that $\overline{\mathcal{Y}}(w, z)$ satisfies the condition (G1). For condition (G2), note that the following relations hold for any $w \in M$, $m \in \mathbb{Z}$:

$$
\mathcal{Y}(L(-1)w, z) = \frac{d}{dz} \mathcal{Y}(w, z),
$$

\n
$$
[L(m), \mathcal{Y}(w, z)] = \sum_{i \ge 0} {m+1 \choose i} z^{m+1-i} \mathcal{Y}(L(i-1)w, z).
$$

 \Box

Using these relations and Lemma [5.4](#page-22-1) we have

$$
[L_{T_{\sigma}}(-1), \mathcal{Y}(w, z)]
$$
\n
$$
= \frac{1}{k} \Big[L(-k), \mathcal{Y}(\Delta_{k}(z)w, z^{1/k}) \Big]
$$
\n
$$
= \frac{1}{k} \sum_{i \geq 0} {k+1 \choose i} (z^{1/k})^{1-k-i} \mathcal{Y}(L(i-1)\Delta_{k}(z)w, z^{1/k})
$$
\n
$$
= \frac{1}{k} z^{\frac{1}{k}-1} \mathcal{Y}(L(-1)\Delta_{k}(z)w, z^{1/k}) + \frac{1}{k} \sum_{i \geq 1} {k+1 \choose i} z^{-1-(i-1)/k} \mathcal{Y}(L(i-1)\Delta_{k}(z)w, z^{1/k})
$$
\n
$$
= \frac{1}{k} z^{\frac{1}{k}-1} \Big(\frac{\partial}{\partial y^{1/k}} \mathcal{Y}(\Delta_{k}(z)w, y^{1/k}) \Big) |_{y=z}
$$
\n
$$
+ \frac{1}{k} \sum_{i \geq 1} {k+1 \choose i} z^{-1-(i-1)/k} \mathcal{Y}(L(i-1)\Delta_{k}(z)w, z^{1/k})
$$
\n
$$
= \frac{1}{k} \frac{\partial}{\partial y} \mathcal{Y}(\Delta_{k}(z)w, y^{1/k}) \Big) |_{y=z} + \mathcal{Y}(\frac{\partial}{\partial z} \Delta_{k}(z)w, z^{1/k})
$$
\n
$$
= \frac{1}{dz} \mathcal{Y}(\Delta_{k}(z)w, z^{1/k})
$$
\n
$$
= \frac{1}{dz} \overline{\mathcal{Y}}(w, z).
$$

For condition (G3), let $u \in V$, $w \in M$. By Lemma 3.3 in [\[BDM\]](#page-35-2), we have

$$
Y_{T_{\sigma}(W)}(u^{1}, z_{1})\overline{\mathcal{Y}}(w, z_{2}) - \overline{\mathcal{Y}}(w, z_{2})Y_{T_{\sigma}(N)}(u^{1}, z_{1})
$$

\n
$$
= \text{Res}_{z_{0}} \frac{1}{k} z_{2}^{-1} \delta\left(\frac{(z_{1} - z_{0})^{1/k}}{z_{2}^{1/k}}\right) \overline{\mathcal{Y}}(Y_{M}(u, z_{0})w, z_{2})
$$

\n
$$
= \text{Res}_{z_{0}} \frac{1}{k} \sum_{r=0}^{k-1} z_{2}^{-1} \delta\left(\frac{z_{1} - z_{0}}{z_{2}}\right) \left(\frac{z_{1} - z_{0}}{z_{2}}\right)^{-r/k} \overline{\mathcal{Y}}(Y_{M}(u, z_{0})w, z_{2})
$$

\n
$$
= \text{Res}_{z_{0}} \frac{1}{k} \sum_{r=0}^{k-1} z_{1}^{-1} \delta\left(\frac{z_{2} + z_{0}}{z_{1}}\right) \left(\frac{z_{2} + z_{0}}{z_{1}}\right)^{r/k} \overline{\mathcal{Y}}(Y_{M}(u, z_{0})w, z_{2}).
$$
 (5.18)

It follows that there exists a nonnegative integer n such that

$$
(z_1 - z_2)^n Y_{T_{\sigma}(W)}(u^1, z_1) \overline{\mathcal{Y}}(w, z_2) = (z_1 - z_2)^n \overline{\mathcal{Y}}(w, z_2) Y_{T_{\sigma}(N)}(u^1, z_1).
$$
\n(5.19)

Since $\{\sigma^j(u^1) \mid u \in V, 1 \leq j \leq k\}$ generates $V^{\otimes k}$ as a vertex algebra, by Lemma [4.7](#page-19-1) we have $\overline{\mathcal{Y}}(w, z) \in$ $\mathcal{H}(T_{\sigma}(N), T_{\sigma}(W))$. Note that $V^{\otimes k}$ as a vertex algebra is generated by the subset $\{u^j \mid u \in V, 1 \le j \le k\}$.

Second, we show that f is a homomorphism of weak V-modules where $\mathcal{H}(T_{\sigma}(N), T_{\sigma}(W))$ is viewed as a Vmodule through the embedding π_1 of V into $V^{\otimes k}$. Note that for $1 \leq i \leq k$,

$$
Y_{T_{\sigma}(W)}(u^{i}, z_{1}) = Y_{T_{\sigma}(W)}(\sigma^{i-1}u^{1}, z_{1}) = \lim_{z_{1}^{1/k} \to \eta_{k}^{1-i} z_{1}^{1/k}} Y_{T_{\sigma}(W)}(u^{1}, z_{1}).
$$

Then using [\(5.18\)](#page-25-0) (by replacing $z_1^{1/k}$ with $\eta_k^{1-i} z_1^{1/k}$), we get

$$
Y_{T_{\sigma}(W)}(u^{i}, z_{1})\overline{\mathcal{Y}}(w, z_{2}) - \overline{\mathcal{Y}}(w, z_{2})Y_{T_{\sigma}(N)}(u^{i}, z_{1})
$$

= Res_{z₀} $\frac{1}{k} \sum_{r=0}^{k-1} z_{1}^{-1} \delta\left(\frac{z_{2}+z_{0}}{z_{1}}\right) \left(\frac{z_{2}+z_{0}}{z_{1}}\right)^{r/k} \eta_{k}^{r(i-1)} \overline{\mathcal{Y}}\left(Y_{M}(u, z_{0})w, z_{2}\right).$ (5.20)

For $u \in V$, write

$$
u^{i} = u^{i,0} + u^{i,1} + \dots + u^{i,k-1}, \tag{5.21}
$$

where $u^{i,r} \in (V^{\otimes k})^r$ for $0 \le r \le k - 1$. As

$$
Y_{T_{\sigma}(W)}(u^{i,r}, z_1) \in z_1^{-\frac{r}{T}}(\text{End}T_{\sigma}(W))[[z_1, z_1^{-1}]], \quad Y_{T_{\sigma}(N)}(u^{i,r}, z_1) \in z_1^{-\frac{r}{T}}(\text{End}T_{\sigma}(N))[[z_1, z_1^{-1}]],
$$

from (5.20) we get

$$
Y_{T_{\sigma}(W)}(u^{i,r}, z_1) \overline{\mathcal{Y}}(w, z_2) - \overline{\mathcal{Y}}(w, z_2) Y_{T_{\sigma}(N)}(u^{i,r}, z_1)
$$

= Res_{z_0} $\frac{1}{k} z_1^{-1} \eta_k^{(i-1)r} \left(\frac{z_2 + z_0}{z_1}\right)^{r/k} \delta\left(\frac{z_2 + z_0}{z_1}\right) \overline{\mathcal{Y}}(Y_M(u, z_0)w, z_2).$ (5.22)

As $\mathcal{Y}(\cdot, x)$ is an intertwining operator of type ${W \choose M N}$, we have

$$
z_0^{-1}\delta\left(\frac{x_1^{1/k} - x^{1/k}}{z_0}\right) Y_W\left(\Delta_k(x_1)v, x_1^{1/k}\right) \mathcal{Y}\left(\Delta_k(x)w, x^{1/k}\right)
$$

$$
- z_0^{-1}\delta\left(\frac{-x^{1/k} + x_1^{1/k}}{z_0}\right) \mathcal{Y}\left(\Delta_k(x)w, x^{1/k}\right) Y_N\left(\Delta_k(x_1)v, x_1^{1/k}\right)
$$

$$
= x^{-1/k}\delta\left(\frac{x_1^{1/k} - z_0}{x^{1/k}}\right) \mathcal{Y}\left(Y_M(\Delta_k(x_1)v, z_0)\Delta_k(x)w, x^{1/k}\right).
$$
(5.23)

Let p be a nonnegative integer such that $z_0^p Y_M(\Delta_k(x_1)v, z_0) \Delta_k(x)w$ involves only nonnegative (integer) powers of z_0 . Then we have

$$
(x_1-x)^p Y_W\left(\Delta_k(x_1)v, x_1^{1/k}\right) \mathcal{Y}\left(\Delta_k(x)w, x^{1/k}\right) = (x_1-x)^p \mathcal{Y}\left(\Delta_k(x)w, x^{1/k}\right) Y_N\left(\Delta_k(x_1)v, x_1^{1/k}\right),
$$

which is

$$
(x_1 - x)^p Y_{T_\sigma(W)}(v^1, x_1) \overline{\mathcal{Y}}(w, x) = (x_1 - x)^p \overline{\mathcal{Y}}(w, x) Y_{T_\sigma(N)}(v^1, x).
$$

Then

$$
z^{p}Y_{\mathcal{H}}(v^{1}, z)\overline{\mathcal{Y}}(w, x) = ((x_{1} - x)^{p}Y_{T_{\sigma}(W)}(v^{1}, x_{1})\overline{\mathcal{Y}}(w, x))|_{x_{1}^{1/k} = (x+z)^{1/k}}.
$$
\n(5.24)

On the other hand, from [\(5.23\)](#page-26-0) we get

$$
x_1^{-1/k} \delta\left(\frac{x^{1/k} + z_0}{x_1^{1/k}}\right) \left((x_1 - x)^p Y_{T_\sigma(W)}(v^1, x_1) \overline{\mathcal{Y}}(w^1, x)\right)
$$

= $x_1^{-1/k} \delta\left(\frac{x^{1/k} + z_0}{x_1^{1/k}}\right) (x_1 - x)^p \mathcal{Y}\left(Y_M(\Delta_k(x_1)v, z_0) \Delta_k(x) w, x^{1/k}\right).$

Taking the residue with respect to $x_1^{1/k}$, we obtain

$$
\begin{split} & \left((x_1 - x)^p Y_{T_{\sigma}(W)}(v^1, x_1) \overline{\mathcal{Y}}(w, x) \right) \big|_{x_1^{1/k} = x^{1/k} + z_0} \\ & = \left((x_1 - x)^p \mathcal{Y} \left(Y_M(\Delta_k(x_1)v, z_0) \Delta_k(x) w, x^{1/k} \right) \right) \big|_{x_1^{1/k} = x^{1/k} + z_0}. \end{split} \tag{5.25}
$$

Notice that for any

$$
f(x, x_1, z_0) \in x^{\alpha} U((x^{1/T}))((x_1^{1/k}, z_0))
$$

with α a complex number, U a vector space and T a positive integer, we have

$$
\left(f(x,x_1,z_0)|_{x_1^{1/k}=x^{1/k}+z_0}\right)|_{z_0=(x+z)^{1/k}-x^{1/k}}=f(x,x_1,(x+z)^{1/k}-x^{1/k})|_{x_1^{1/k}=(x+z)^{1/k}}.
$$

In view of this, applying substitution $z_0 = (x + z)^{1/k} - x^{1/k}$ to [\(5.25\)](#page-26-1), we get

$$
\begin{split} & \left((x_1 - x)^p Y_{T_\sigma(W)}(v^1, x_1) \overline{\mathcal{Y}}(w, x) \right) \big|_{x_1^{1/k} = (x+z)^{1/k}} \\ & = \left((x_1 - x)^p \mathcal{Y} \left(Y_M(\Delta_k(x_1)v, (x+z)^{1/k} - x^{1/k}) \Delta_k(x)w, x^{1/k} \right) \right) \big|_{x_1^{1/k} = (x+z)^{1/k}} \\ & = z^p \mathcal{Y} \left(Y_M(\Delta_k(x+z)v, (x+z)^{1/k} - x^{1/k}) \Delta_k(x)w, x^{1/k} \right). \end{split} \tag{5.26}
$$

Combining this with (5.24) , and then using (5.6) we get

$$
Y_{\mathcal{H}}(v^1, z)\overline{\mathcal{Y}}(w, x) = \mathcal{Y}\left(Y_M(\Delta_k(x+z)v, (x+z)^{1/k} - x^{1/k})\Delta_k(x)w, x^{1/k}\right)
$$

= $\mathcal{Y}\left(\Delta_k(x)Y_M(v, z)w, x^{1/k}\right)$
= $\overline{\mathcal{Y}}(Y_M(v, z)w, x).$ (5.27)

Furthermore, for any $v \in V$, $a \in \mathbb{C}1 \otimes V^{\otimes (k-1)}$, we have

$$
f(Y(v,z)w\otimes a)=a_{-1}^{\mathcal{H}}\overline{\mathcal{Y}}(Y_M(v,z)w,x)=a_{-1}^{\mathcal{H}}Y_{\mathcal{H}}(v^1,z)\overline{\mathcal{Y}}(w,x)=Y_{\mathcal{H}}(v^1,z)a_{-1}^{\mathcal{H}}\overline{\mathcal{Y}}(w,x)=Y_{\mathcal{H}}(v^1,z)f(w\otimes a),
$$

noticing that $[a_m^{\mathcal{H}},(v^1)_n^{\mathcal{H}}]=0$ for $m,n \in \mathbb{Z}$ as $(v^1)_i a=0$ for all $i \geq 0$. This proves that f is a homomorphism of weak V-modules.

Third, we show that f is a homomorphism of weak $\mathbb{C}1 \otimes V^{\otimes (k-1)}$ -modules. To this end, we first show that if $2 \leq i \leq k$, then

$$
(u^i)_n^{\mathcal{H}} \overline{\mathcal{Y}}(w, x) = 0 \quad \text{for all } u \in V, \ w \in M, \ n \ge 0. \tag{5.28}
$$

Recall

$$
u^{i} = u^{i,0} + u^{i,1} + \cdots + u^{i,k-1},
$$

where $u^{i,r} \in (V^{\otimes k})^r$ for $0 \le r \le k - 1$. By definition, we have

$$
(u^{i,r})_n^{\mathcal{H}} \overline{\mathcal{Y}}(w,x)
$$

=
$$
\sum_{j\geq 0} \left(\frac{-\frac{r}{k}}{j}\right) \text{Res}_{x_1} x^{-\frac{r}{k}-j} x_1^{\frac{r}{T}} \left\{(x_1-x)^{n+j} Y_{T_{\sigma}(W)} (u^{i,r}, x_1) \overline{\mathcal{Y}}(w,x) - (-x+x_1)^{n+j} \overline{\mathcal{Y}}(w,x) Y_{T_{\sigma}(N)} (u^{i,r}, x_1) \right\}
$$

for $n \in \mathbb{Z}$. Assume $n \geq 0$, using [\(5.22\)](#page-26-3) we get

$$
(u^i)_n^{\mathcal{H}} \overline{\mathcal{Y}}(w, x)
$$
\n
$$
= \sum_{r=0}^{k-1} (u^{i,r})_n^{\mathcal{H}} \overline{\mathcal{Y}}(w, x)
$$
\n
$$
= \sum_{r=0}^{k-1} \text{Res}_{x_1} \sum_{j\geq 0} \left(\frac{-\frac{r}{k}}{j} \right) x^{-\frac{r}{k} - j} x_1^{\frac{r}{k}} (x_1 - x)^{n+j} \left(Y_{T_{\sigma}(W)}(u^{i,r}, x_1) \overline{\mathcal{Y}}(w, x) - \overline{\mathcal{Y}}(w, x) Y_{T_{\sigma}(N)}(u^{i,r}, x_1) \right)
$$
\n
$$
= \text{Res}_{x_0} \text{Res}_{x_1} \sum_{r=0}^{k-1} \sum_{j\geq 0} \left(\frac{-\frac{r}{k}}{j} \right) x^{-\frac{r}{k} - j} x_1^{\frac{r}{k}} (x_1 - x)^{n+j}
$$
\n
$$
\cdot \frac{1}{k} \eta_k^{(1-i)r} x_1^{-1} \delta \left(\frac{x_2 + x_0}{x_1} \right) \left(\frac{x_2 + x_0}{x_1} \right)^{r/k} \overline{\mathcal{Y}}(Y_M(u, x_0) w, x)
$$
\n
$$
= \text{Res}_{x_0} \text{Res}_{x_1} \sum_{r=0}^{k-1} \sum_{j\geq 0} \left(\frac{-\frac{r}{k}}{j} \right) x^{-\frac{r}{k} - j} x_1^{\frac{r}{k}} x_0^{n+j}
$$
\n
$$
\cdot \frac{1}{k} \eta_k^{(i-1)r} x_1^{-1} \delta \left(\frac{x + x_0}{x_1} \right) \left(\frac{x + x_0}{x_1} \right)^{r/k} \overline{\mathcal{Y}}(Y_M(u, x_0) w, x)
$$
\n
$$
= \text{Res}_{x_0} \text{Res}_{x_1} \sum_{r=0}^{k-1} \left(\frac{x + x_0}{x_1} \right)^{-r/k} x_0^{n} \cdot \frac{1}{k} \eta_k^{(i-1)r} x_1^{-1} \delta \left(\frac{x + x_0}{x_1} \right) \left(
$$

This proves [\(5.28\)](#page-27-0). Since V^i for $2 \le i \le k$ generate $\mathbb{C}1\otimes V^{\otimes(k-1)}$ as a vertex algebra, from [\[LL\]](#page-37-6), $\overline{\mathcal{Y}}(w, x)$ for $w\in M$ are vacuum-like vectors in $\mathcal{H}(T_\sigma(N),T_\sigma(W))$ viewed as a weak module for $\mathbb{C}1\otimes V^{\otimes(k-1)}$ in the sense of [\[Li1\]](#page-36-20). Then by ([\[LL\]](#page-37-6); Proposition 4.7.7, or Corollary 4.7.8), the linear map f from $M \otimes V^{\otimes (k-1)}$ to $\mathcal{H}(T_{\sigma}(N), T_{\sigma}(N))$, defined by $f(w \otimes a) = a \cdot \frac{\mu}{\nu} \overline{\mathcal{Y}}(w, x)$ for $w \in M$, $a \in V^{\otimes (k-1)}$, is a weak module homomorphism for $\mathbb{C}1 \otimes V^{\otimes (k-1)}$. As V^1 and $\mathbb{C}1 \otimes V^{\otimes (k-1)}$ generate $V^{\otimes k}$ as a vertex algebra, it follows immediately that f is a weak $V^{\otimes k}$ -module homomorphism. This completes the proof. \Box

Given V-modules M, N, and W, we have the space $I_V \begin{pmatrix} W \\ M \end{pmatrix}$ of intertwining operators. Furthermore, we have σ-twisted $V^{\otimes k}$ -modules $T_{\sigma}(N)$ and $T_{\sigma}(W)$, and we have the space $I_{V^{\otimes k}}\binom{T_{\sigma}(W)}{M^1\;T_{\sigma}(N)}$ of intertwining operators of type $\binom{T_{\sigma}(W)}{M^1\;T_{\sigma}(N)}$. On the other hand, in view of Theorem [4.6,](#page-15-0) we have a weak $V^{\otimes k}$ -module $(\mathcal{H}(T_{\sigma}(N), T_{\sigma}(W)), Y_{\mathcal{H}}(\cdot, z))$. Combining Theorem [5.5](#page-24-0) and Proposition [4.8,](#page-19-2) we immediately have:

Corollary 5.6. For every intertwining operator $\mathcal{Y}(\cdot, z)$ of type $\begin{pmatrix} W \\ M \end{pmatrix}$, there exists an intertwining operator $\overline{\mathcal{Y}}(\cdot, z)$ of type $\binom{T_{\sigma}(W)}{M^1\;T_{\sigma}(N)}$, which is uniquely determined by

$$
\overline{\mathcal{Y}}(w^1, z) = \mathcal{Y}(\Delta_k(z)w, z^{1/k}) \quad \text{for } w \in M.
$$
\n(5.29)

In view of Corollary [5.6,](#page-28-0) we have a linear map

$$
\pi: I_V \binom{W}{M \ N} \to I_{V^{\otimes k}} \binom{T_{\sigma}(W)}{M^1 \ T_{\sigma}(N)}; \ \mathcal{Y}(\cdot, z) \mapsto \overline{\mathcal{Y}}(\cdot, z). \tag{5.30}
$$

5.1.2 From $I_{V^{\otimes k}}\binom{T_{\sigma}(W)}{M^1\;T_{\sigma}(N)}$ to $I_{V}\binom{W}{M\;N}$

In this section, we shall prove that the linear map π defined above from $I_V \binom{W}{M \ N}$ to $I_{V \otimes k} \binom{T_{\sigma}(W)}{M^1 \ T_{\sigma}(N)}$ is an isomorphism by constructing its inverse map. The arguments here are similar to those in [\[BDM\]](#page-35-2) (Section 4).

Let V, k, and σ be given as before. On the other hand, let M, N, and W be V-modules. Set $M^1 = M \otimes V^{\otimes (k-1)}$, which is a $V^{\otimes k}$ -module. Recall the σ -twisted $V^{\otimes k}$ -modules $T_{\sigma}(N)$ and $T_{\sigma}(W)$.

Recall $\Delta_k(z)$ from [\(5.3\)](#page-21-3). Here, we shall need its inverse

$$
\Delta_k(z)^{-1} = z^{(1-1/k)L(0)} k^{L(0)} \exp\left(-\sum_{j\geq 1} a_j z^{-j/k} L(j)\right). \tag{5.31}
$$

As before, by convention we define $k^{\alpha} = e^{\alpha \ln k}$ and $(z^k)^{\alpha} = z^{k\alpha}$ for any $\alpha \in \mathbb{C}$. Furthermore, for any formal series

$$
A(x) = \sum_{\alpha \in \mathbb{C}} u_{\alpha} x^{\alpha} \in U\{x\}
$$

with U a vector space, we define

$$
A(z^k) = \sum_{\alpha \in \mathbb{C}} u_{\alpha} z^{k\alpha} \in U\{z\}.
$$

In particular, if w is an $L(0)$ -eigenvector in a V-module with eigenvalue $\alpha \in \mathbb{C}$, we have

$$
z^{(1-1/k)L(0)}k^{L(0)}w = z^{(1-1/k)\alpha}k^{\alpha}w.
$$

Notice that $\Delta_k(z^k)^{-1} \mathbf{1} = \mathbf{1}$.

The following lemma follows from the proof of Lemma 4.1 in [\[BDM\]](#page-35-2):

Lemma 5.7. Let $\overline{\mathcal{Y}}(\cdot, z)$ be an intertwining operator of type $\begin{pmatrix} T_{\sigma}(W) \\ M^T T_{\sigma}(N) \end{pmatrix}$. For $w \in M$, set

$$
\mathcal{Y}(w,z) = \overline{\mathcal{Y}}\left(\left(\Delta_k (z^k)^{-1} w \right)^1, z^k \right),
$$

an element of $(\text{Hom}(N, W))\{z\}$, where $(\Delta_k(z^k)^{-1}w)^1 = \Delta_k(z^k)^{-1}w \otimes \mathbf{1}^{\otimes (k-1)} \in M^1\{z\}$. Then

$$
\mathcal{Y}(L(-1)w, z) = \frac{d}{dz}\mathcal{Y}(w, z)
$$

on N for any $w \in M$.

Furthermore, we have the following commutator formula:

Lemma 5.8. Let $\overline{\mathcal{Y}}(\cdot, z)$ be an intertwining operator of type $\begin{pmatrix} T_{\sigma}(W) \\ M^T T_{\sigma}(N) \end{pmatrix}$. Then

$$
Y_W(u, z_1) \mathcal{Y}(w, z_2) - \mathcal{Y}(w, z_2) Y_N(u, z_1) = \text{Res}_{z_0} z_2^{-1} \delta\left(\frac{z_1 - z_0}{z_2}\right) \mathcal{Y}(Y_M(u, z_0)w, z_2)
$$

for $u \in V$, $w \in M$, where $\mathcal{Y}(w, z)$ for $w \in M$ is defined as in Lemma [5.7.](#page-29-0)

Proof. Let $u \in V$, $w \in M$. Noticing that

$$
(\sigma^j u^1)_n w^1 = (u^{j+1})_n w^1 = 0 \quad \text{for all } 1 \le j \le k - 1, \ n \ge 0,
$$

from the twisted Jacobi identity [\(2.7\)](#page-6-0), we get

$$
Y_{T_{\sigma}(W)}(u^1, z_1) \overline{\mathcal{Y}}(w^1, z_2) - \overline{\mathcal{Y}}(w^1, z_2) Y_{T_{\sigma}(N)}(u^1, z_1)
$$

= Res_{z₀} $\frac{1}{k} z_2^{-1} \delta \left(\frac{(z_1 - z_0)^{1/k}}{z_2^{1/k}} \right) \overline{\mathcal{Y}}(Y(u^1, z_0) w^1, z_2).$

Then we have (see [\[BDM\]](#page-35-2), Section 4)

$$
Y_{W}(u, z_{1})\mathcal{Y}(w, z_{2}) - \mathcal{Y}(w, z_{2})Y_{N}(u, z_{1})
$$
\n
$$
= Y_{T_{\sigma}(W)} \left(\left(\Delta_{k} (z_{1}^{k})^{-1} u \right)^{1}, z_{1}^{k} \right) \overline{\mathcal{Y}} \left(\left(\Delta_{k} (z_{2}^{k})^{-1} w \right)^{1}, z_{2}^{k} \right)
$$
\n
$$
- \overline{\mathcal{Y}} \left(\left(\Delta_{k} (z_{2}^{k})^{-1} w \right)^{1}, z_{2}^{k} \right) Y_{T_{\sigma}(N)} \left(\left(\Delta_{k} (z_{1}^{k})^{-1} u \right)^{1}, z_{1}^{k} \right)
$$
\n
$$
= \text{Res}_{x} \frac{1}{k} z_{2}^{-k} \delta \left(\frac{\left(z_{1}^{k} - x \right)^{1/k}}{z_{2}} \right) \overline{\mathcal{Y}} \left(Y \left(\left(\Delta_{k} (z_{1}^{k})^{-1} u \right)^{1}, x \right) \left(\Delta_{k} (z_{2}^{k})^{-1} w \right)^{1}, z_{2}^{k} \right).
$$

We are going to use the change-of-variable $x = z_1^k - (z_1 - z_0)^k$. Note that for any $n \in \mathbb{Z}$,

$$
\left(z_1^k - x\right)^{n/k}\Big|_{x=z_1^k - (z_1 - z_0)^k} = (z_1 - z_0)^n,\tag{5.32}
$$

where by convention $(z_1^k - x)^{n/k} = \sum_{j \geq 0} {n/k \choose j} (-1)^j z_1^{n-kj} x^j$. Using (3.12) in [\[BDM\]](#page-35-2), we have

$$
Y_{W}(u, z_{1})\mathcal{Y}(w, z_{2}) - \mathcal{Y}(w, z_{2})Y_{N}(u, z_{1})
$$
\n
$$
= Res_{z_{0}} z_{2}^{-k} (z_{1} - z_{0})^{k-1} \delta\left(\frac{z_{1} - z_{0}}{z_{2}}\right) \overline{\mathcal{Y}}\left(Y\left((\Delta_{k}(z_{1}^{k})^{-1}u)^{1}, z_{1}^{k} - (z_{1} - z_{0})^{k}\right)\left(\Delta_{k}(z_{2}^{k})^{-1}w\right)^{1}, z_{2}^{k}\right)
$$
\n
$$
= Res_{z_{0}} z_{2}^{-1} \delta\left(\frac{z_{1} - z_{0}}{z_{2}}\right) \overline{\mathcal{Y}}\left(Y\left((\Delta_{k}(z_{1}^{k})^{-1}u)^{1}, z_{1}^{k} - (z_{1} - z_{0})^{k}\right)\left(\Delta_{k}(z_{2}^{k})^{-1}w\right)^{1}, z_{2}^{k}\right)
$$
\n
$$
= Res_{z_{0}} z_{2}^{-1} \delta\left(\frac{z_{1} - z_{0}}{z_{2}}\right) \overline{\mathcal{Y}}\left(\left(Y\left(\Delta_{k}\left((z_{2} + z_{0})^{k}\right)^{-1}u,(z_{2} + z_{0})^{k} - z_{2}^{k}\right)\Delta_{k}(z_{2}^{k})^{-1}w\right)^{1}, z_{2}^{k}\right).
$$

Then it suffices to prove

$$
Y\left(\Delta_k\left((z_2+z_0)^k\right)^{-1}u,(z_2+z_0)^k-z_2^k\right)\Delta_k(z_2^k)^{-1}=\Delta_k(z_2^k)^{-1}Y(u,z_0),\tag{5.33}
$$

or equivalently

$$
\Delta_k(z_2^k) Y \left(\Delta_k ((z_2 + z_0)^k)^{-1} u, (z_2 + z_0)^k - z_2^k\right) \Delta_k(z_2^k)^{-1} = Y(u, z_0).
$$

This last relation follows from (4.24) ([\[BDM\]](#page-35-2); Proposition 2.2) by replacing u, z and z_0 with $\Delta_k((z_2 + z_0)^k)^{-1}u, z_2^k$ and $(z_2 + z_0)^k - z_2$, respectively. Now, the proof is complete.

As the second main result of this section we have:

Theorem 5.9. Let M, N and W be V-modules and let $\sigma = (1 \ 2 \ \cdots k) \in$ Aut $(V^{\otimes k})$. Set $M^1 = M \otimes V^{\otimes (k-1)}$. Let $\overline{\mathcal{Y}}(\cdot, z)$ be any intertwining operator of type $\binom{T_{\sigma}(W)}{M^T T_{\sigma}(N)}$. For $w \in M$, set

$$
\mathcal{Y}(w,z) = \overline{\mathcal{Y}}\left(\left(\Delta_k (z^k)^{-1} w \right)^1, z^k \right),
$$

an element of $(\text{Hom}(N, W))\{z\}$. Then $\mathcal{Y}(\cdot, z)$ is an intertwining operator of type ${W \choose M N}$.

Proof. As the L(−1)-property has been given in Lemma [5.7,](#page-29-0) it remains to prove the Jacobi identity which is equivalent to the commutator formula which was obtained in Lemma [5.8](#page-29-1) and the weak associativity which states that for any $u \in V$, $w \in M$, $a \in N$, there exists a nonnegative integer n such that

$$
(z_0 + z_2)^n Y_W(u, z_0 + z_2) \mathcal{Y}(w, z_2) a = (z_2 + z_0)^n \mathcal{Y}(Y_M(u, z_0)w, z_2) a.
$$

Let $u \in V$, $w \in M$. Recall

$$
u^1=u\otimes \mathbf{1}^{\otimes (k-1)}\in V^{\otimes k},\quad w^1=w\otimes \mathbf{1}^{\otimes (k-1)}\in M^1.
$$

We have $u^1 = \sum_{r=0}^{k-1} u^{1,r}$, where $u^{i,r} \in (V^{\otimes k})^r$, i.e., $\sigma u^{1,r} = \eta_k^r u^{1,r}$. By the twisted Jacobi identity, there exists a positive integer l such that for $n \geq l$,

$$
(x_0 + x_2)^{r/k+n} Y_{T_{\sigma}(W)}(u^{1,r}, x_0 + x_2) \overline{\mathcal{Y}}(w^1, x_2) a = (x_2 + x_0)^{r/k+n} \overline{\mathcal{Y}}(Y_{M^1}(u^{1,r}, x_0) w^1, x_2) a
$$

for $r = 0, 1, ..., k - 1$. Using change-of-variables $x_2 = z_2^k$ and $x_0 = (z_0 + z_2)^k - z_2^k$, we obtain

$$
(z_0 + z_2)^{r + kn} Y_{T_{\sigma}(W)} \left(u^{1,r}, (z_0 + z_2)^k \right) \overline{\mathcal{Y}} \left(w^1, z_2^k \right) a = (z_2 + z_0)^{r + kn} \overline{\mathcal{Y}} \left(Y_{M^1} \left(u^{1,r}, (z_0 + z_2)^k - z_2^k \right) w^1, z_2^k \right) a.
$$

Thus there exists a positive integer p such that

$$
(z_0 + z_2)^n Y_{T_{\sigma}(W)} (u^{1,r}, (z_0 + z_2)^k) \overline{\mathcal{Y}} (w^1, z_2^k) a = (z_2 + z_0)^n \overline{\mathcal{Y}} (Y_{M^1} (u^{1,r}, (z_0 + z_2)^k - z_2^k) w^1, z_2^k) a
$$

for $n \geq p$, $r = 0, 1, \ldots, k - 1$. Consequently, we have

$$
(z_0 + z_2)^n Y_{T_{\sigma}(W)} (u^1, (z_0 + z_2)^k) \overline{\mathcal{Y}} (w^1, z_2^k) a = (z_2 + z_0)^n \overline{\mathcal{Y}} (Y_{M^1} (u^1, (z_0 + z_2)^k - z_2^k) w^1, z_2^k) a
$$

for $n \geq p$.

Notice that $\Delta_k(z^k)^{-1}u \in V[z, z^{-1}]$ and $\Delta_k(z^k)^{-1}w \in M[z, z^{-1}]$. (Both of them are finite sums.) Then, there exists a positive integer q such that

$$
(z_0 + z_2)^n Y_{T_{\sigma}(W)} ((\Delta_k (z_0 + z_2)^k)^{-1} u)^1, (z_0 + z_2)^k) \overline{\mathcal{Y}} ((\Delta_k (z_2^k)^{-1} w)^1, z_2^k) a
$$

= $(z_2 + z_0)^n \overline{\mathcal{Y}} (Y_{M^1} ((\Delta_k (z_0 + z_2)^k)^{-1} u)^1, (z_0 + z_2)^k - z_2^k) (\Delta_k (z_2^k)^{-1} w)^1, z_2^k) a$

for $n \geq q$. Therefore, we obtain

$$
(z_0 + z_2)^n Y_W(u, z_0 + z_2) \mathcal{Y}(w, z_2) a
$$

= $(z_0 + z_2)^n Y_{T_{\sigma}(W)} \left(\left(\Delta_k \left((z_0 + z_2)^k \right)^{-1} u \right)^1, (z_0 + z_2)^k \right) \overline{\mathcal{Y}} \left(\left(\Delta_k \left(z_2^{k} \right)^{-1} w \right)^1, z_2^{k} \right) a$
= $(z_0 + z_2)^n \overline{\mathcal{Y}} \left(Y_{M^1} \left(\left(\Delta_k \left((z_2 + z_0)^k \right)^{-1} u \right)^1, (z_2 + z_0)^k - z_2^{k} \right) \left(\Delta \left(z_2^{k} \right)^{-1} w \right)^1, z_2^{k} \right) a$
= $(z_0 + z_2)^n \overline{\mathcal{Y}} \left(\left(\Delta_k \left(z_2^{k} \right)^{-1} Y(u, z_0) w \right)^1, z_2^{k} \right) a$
= $(z_0 + z_2)^n \mathcal{Y} \left(Y_M(u, z_0) w, z_2 \right) a$

 \Box

for $n \geq p$, where we are using [\(5.33\)](#page-30-0) for the second last equality. This completes the proof.

To summarize, we have:

Corollary 5.10. The linear map π defined in [\(5.30\)](#page-28-1) is a linear isomorphism from $I_V \begin{pmatrix} W \\ M \end{pmatrix}$ to $I_{V\otimes k} \begin{pmatrix} T_{\sigma}(W) \\ M^1 \end{pmatrix}$.

6 Tensor products of $V^{\otimes k}$ -modules with permutation automorphism twisted $V^{\otimes k}$ -modules

In this section, we study tensor product using the results of Section 5 from the k-cycle σ to an arbitrary permutation in S_k and present the second main theorem of this paper. Some of the results in this section are valid for arbitrary vertex operator algebras.

Let V be a general vertex operator algebra for now. Recall Definition [2.18](#page-7-0) for a tensor product of a q_1 -twisted V-module M with a g_2 -twisted V-module N. As a convention, we shall use $M \boxtimes N$ to denote a generic tensor product module, provided that its existence is justified.

Remark 6.1. Let V be any vertex operator algebra. Note that for any V-module (M, Y_M) , the pair (M, Y_M) is a tensor product of V and M. Furthermore, (M, Y_M^o) is a tensor product of M and V, where $Y_M^o(\cdot, z)$ is defined by

$$
Y_M^o(w, z)v = e^{zL(-1)}Y_M(v, -z)w \quad \text{for } v \in V, w \in M
$$

(see [\[FHL\]](#page-36-6)). Thus $M \boxtimes V \simeq M \simeq V \boxtimes M$.

Remark 6.2. Let U and V be vertex operator algebras and let (W, Y_W) be a U-module and (M, Y_M) a V-module. Then it is straightforward to show that $W \otimes M$ is a tensor product module for $U \otimes V$ -modules $W \otimes V$ and $U \otimes M$, where

$$
\mathcal{Y}(w \otimes v, z)(u \otimes m) = e^{zL(-1)} Y_W(v, -z)w \otimes Y_M(u, z)m \tag{6.1}
$$

for $w \in W$, $v \in V$, $u \in U$, $m \in M$. That is, $W \otimes M \simeq (W \otimes V) \boxtimes (U \otimes M)$.

Let k be a positive integer. Recall from [\[BDM\]](#page-35-2) that for any V-module W, we have a σ -twisted $V^{\otimes k}$ -module $T_{\sigma}(W)$ which equals W as a vector space and a V-module homomorphism from W_1 to W_2 is exactly the same as a σ -twisted $V^{\otimes k}$ -module homomorphism from $T_{\sigma}(W_1)$ to $T_{\sigma}(W_2)$.

Let M and N be V-modules. Assume that $(M \boxtimes N, Y)$ is a tensor product of M and N. That is, $M \boxtimes N$ is a V-module and $\mathcal Y$ is an intertwining operator of type $\binom{M\boxtimes N}{M N}$, satisfying the universal property. From Theorem [5.5,](#page-24-0) we have an intertwining operator \overline{Y} of type $\binom{T_{\sigma}(M\boxtimes N)}{M^1\ T_{\sigma}(N)}$. It follows from Corollary [5.10](#page-31-0) that $(T_{\sigma}(M\boxtimes N), \overline{Y})$ is a tensor product of $M \otimes V^{\otimes (k-1)}$ with $T_{\sigma}(N)$. Thus we have proved:

Theorem 6.3. Let M and N be V-modules. Suppose that there exists a tensor product $(M \boxtimes N, Y)$ of M and N. Then a tensor product of $M \otimes V^{\otimes (k-1)}$ with $T_{\sigma}(N)$ exists and

$$
(M \otimes V^{\otimes (k-1)}) \boxtimes T_{\sigma}(N) \simeq T_{\sigma}(M \boxtimes N)
$$

In particular, we have $(M \otimes V^{\otimes (k-1)}) \boxtimes T_{\sigma}(V) \simeq T_{\sigma}(M)$ and Theorem A (in Introduction) is true.

Next, we shall generalize this result. First, we formulate the following lemma which is straightforward to prove:

Lemma 6.4. Let U be a vertex operator algebra and let σ , τ be automorphisms of U with σ of finite order. Assume that W_1, W_2 are σ -twisted U-modules and W is a V-module on which τ acts such that

$$
\tau Y_W(u, z)w = Y_W(\tau(u), z)\tau w \quad \text{for } u \in U, \ w \in W. \tag{6.2}
$$

If $I(\cdot, z)$ is an intertwining operator of type $\binom{W_2}{W_1},$ then $I^{\tau^{-1}}(\cdot, z)$ is an intertwining operator of type $\binom{W_2}{W^\tau W_1},$ where $W^{\tau} = W$ as a vector space, $Y_{W^{\tau}}(u, z) = Y_W(\tau(u), z)$ for $u \in U$, and

$$
I^{\tau^{-1}}(w, z) = I(\tau^{-1}(w), z) \text{ for } w \in W.
$$

Furthermore, if (P, Y) is a tensor product of W and W_1 , then $(P, Y^{\tau^{-1}})$ is a tensor product of W^{τ} and W_1 . In particular, we have $W \boxtimes W_1 \simeq W^\tau \boxtimes W_1$, provided that one of the two tensor products exists.

We now in a position to prove Theorem B (in Introduction).

Proof. First of all, combining Theorem [6.3](#page-32-1) with Lemma [6.4,](#page-32-0) we have

$$
(V^{\otimes (i-1)} \otimes M \otimes V^{\otimes (k-i)}) \boxtimes T_{\sigma}(N) \simeq T_{\sigma}(M \boxtimes N)
$$

for any V-module M and for $1 \le i \le k$. This shows that this theorem is true if for some $1 \le r \le k$, $M_i = V$ for all $i \neq r$, as $M_1 \boxtimes \cdots \boxtimes M_k \simeq M_r$. For the general case, notice that as $M_i \boxtimes V \simeq M_i \simeq V \boxtimes M_i$ for $1 \leq i \leq k$, we have

$$
M_1 \otimes \cdots \otimes M_k \simeq \boxtimes_{i=1}^k (V^{\otimes (i-1)} \otimes M_i \otimes V^{\otimes (k-i)}),
$$

where we are also using Remark [6.2.](#page-32-2) Then the general case follows from the special case and Theorem [3.7](#page-12-3) (an associativity). □

Now, we consider an arbitrary permutation $\sigma \in S_k$. Recall that a partition of k is a sequence of positive integers k_1, k_2, \ldots, k_s such that

$$
k_1 \geq k_2 \geq \cdots \geq k_s \geq 1 \text{ and } k_1 + \cdots + k_s = k.
$$

Associated to each partition $\kappa = (k_1, k_2, \dots, k_s)$, we have a permutation

$$
\sigma_{\kappa} := \sigma_1 \circ \sigma_2 \circ \cdots \circ \sigma_s,\tag{6.3}
$$

where $\sigma_1 = (12 \cdots k_1)$, $\sigma_2 = ((k_1 + 1) \cdots (k_1 + k_2))$, and so on. A classical fact is that every permutation in S_k is conjugate to σ_{κ} for a uniquely determined partition κ of k.

Remark 6.5. Let U be a vertex operator algebra and let σ , τ be finite order automorphisms of U such that $\tau = \mu \sigma \mu^{-1}$ for some automorphism μ of U. It is straightforward to show that a linear map $Y_W(\cdot, z) : U \to (\text{End}W)\{z\}$ is a τ-twisted U-module structure on W if and only if $Y_W^{\mu}(\cdot, z)$ is a σ-twisted U-module structure on W, where

$$
Y_W^{\mu}(u, z) = Y_W(\mu(u), z) \quad \text{for } u \in U.
$$

If W is a τ -twisted U-module, we denote by W^{τ} the corresponding σ -twisted U-module. On the other hand, if W_1, W_2 are τ -twisted U-modules, a τ -twisted U-module homomorphism from W_1 to W_2 is exactly the same as a σ-twisted U-module homomorphism from W_1^{μ} to W_2^{μ} . Therefore, the category of τ -twisted U-modules is isomorphic to the category of σ -twisted U-modules canonically.

Furthermore, by a straightforward argument we have:

Lemma 6.6. Let V be any vertex operator algebra and let σ and μ be automorphisms of V with σ of finite order. Set $\tau = \mu \sigma \mu^{-1}$. Assume that W is a V-module and T_1, T_2 are τ -twisted V-modules. Then an intertwining operator of type $\binom{T_2}{W^T T_1}$ is exactly the same as an intertwining operator of type $\binom{T_2^{\mu}}{W^{\mu} T_1^{\mu}}$.

Proposition 6.7. Let V be any vertex operator algebra and let σ and μ be automorphisms of V with σ of finite order. Set $\tau = \mu \sigma \mu^{-1}$. Let W be a V-module and T a τ -twisted V-module. Assume that there exists a tensor product $W^{\mu^{-1}} \boxtimes T$ (a τ -twisted V-module). Then there exists a tensor product $W \boxtimes T^{\mu}$ (a σ -twisted V-module) and

$$
W \boxtimes T^{\mu} \simeq (W^{\mu^{-1}} \boxtimes T)^{\mu}.
$$
\n(6.4)

Proof. Recall that T^{μ} is a σ -twisted V-module and $W^{\mu^{-1}}$ is a V-module with W as the underlying space, where the vertex operator map $Y_W^{\mu^{-1}}(\cdot, x)$ is given by

$$
Y_W^{\mu^{-1}}(v, x) = Y_W(\mu^{-1}v, x) \quad \text{ for } v \in V.
$$

From definition, we have an intertwining operator $\mathcal Y$ of type $\binom{W^{\mu^{-1}}\boxtimes T}{W^{\mu^{-1}}T}$, satisfying the universal property. Let P be any σ -twisted V-module and $I(\cdot, z)$ be any intertwining operator of type of $\binom{P}{W T \mu}$. By Lemma [6.6,](#page-33-0) $I(\cdot, z)$ is also an intertwining operator of type $\binom{p^{\mu^{-1}}}{W^{\mu^{-1}}}$ $W^{\mu^{-1}}_{W^{\mu^{-1}}T}$). Then there exists a τ -twisted V-module homomorphism ψ from $W^{\mu^{-1}} \boxtimes T$ to $P^{\mu^{-1}}$ such that $I(w, z)t = \psi(\mathcal{Y}(w, z)t)$ for $w \in W$, $t \in T$. Note that from Remark [6.5,](#page-33-1) ψ is also a σ -twisted V-module homomorphism from $(W^{\mu^{-1}} \boxtimes T)^{\mu}$ to P. This proves that $((W^{\mu^{-1}} \boxtimes T)^{\mu}, Y)$ is a tensor product of W with T^{μ} . Therefore, we have $(W^{\mu^{-1}} \boxtimes T)^{\mu} \simeq W \boxtimes T^{\mu}$, as desired. □

Remark 6.8. Let V_1 and V_2 be vertex operator algebras and let τ_1, τ_2 be finite order automorphisms of V_1, V_2 , respectively. Set $\tau = \tau_1 \otimes \tau_2$, an automorphism of $V_1 \otimes V_2$. Then the same arguments of [\[FHL\]](#page-36-6) show that if W_i is an irreducible τ_i -twisted V_i -module for $i = 1, 2$, then $W_1 \otimes W_2$ is naturally an irreducible τ -twisted $V_1 \otimes V_2$ -module and on the other hand, every irreducible τ -twisted $V_1\otimes V_2$ -module is isomorphic to one in this form. Furthermore, if V_i is τ_i -rational for $i = 1, 2$, then $V_1 \otimes V_2$ is τ -rational (see [\[BDM\]](#page-35-2), Lemma 6.1 and Proposition 6.2).

As before, we view the symmetric group S_k as an automorphism group of $V^{\otimes k}$. From Remarks [6.5](#page-33-1) and [6.8](#page-33-2) and Theorem [5.3](#page-21-4) (due to [\[BDM\]](#page-35-2)) we immediately have:

Proposition 6.9. Let V be any vertex operator algebra and $\sigma \in S_k$ be any permutation. Suppose that $\mu \sigma \mu^{-1} = \sigma_{\kappa}$, where $\mu \in S_k$ and κ is the partition of k which is uniquely determined by σ . Then for any s irreducible V-modules $N_1, ..., N_s$, we have an irreducible σ -twisted $V^{\otimes k}$ -module

$$
T_{\sigma}(N_1,\ldots,N_s) := (T_{\sigma_1}(N_1) \otimes \cdots \otimes T_{\sigma_s}(N_s))^{\mu},\tag{6.5}
$$

where $\sigma_1, \ldots, \sigma_s$ are disjoint cycles as in [\(6.3\)](#page-33-3), and on the other hand, every irreducible σ -twisted $V^{\otimes k}$ -module is isomorphic to one in this form. Furthermore, for irreducible V-modules $N_1, ..., N_s$ and $P_1, ..., P_s, T_{\sigma}(N_1, ..., N_s) \simeq$ $T_{\sigma}(P_1,\ldots,P_s)$ if and only if $N_i \simeq P_i$ for $1 \leq i \leq s$.

To present our general result, we also need the following result:

Proposition 6.10. Let V_1, \ldots, V_s be vertex operator algebras, let τ_1, \ldots, τ_s be finite order automorphisms of V_1, \ldots, V_s , respectively. Assume that V_i is τ_i -rational for $1 \leq i \leq s$. Set

$$
\tau = \tau_1 \otimes \cdots \otimes \tau_s \in \text{Aut}(V_1 \otimes \cdots \otimes V_s).
$$

Assume that W_i is a V_i -module and T_i is a τ_i -twisted V_i -module for $i = 1, \ldots, s$ such that $\dim I_{V_i} {P_i \choose W_i | T_i} < \infty$ for every irreducible τ_i -twisted V_i -module P_i for $1 \leq i \leq s$. Then there exist tensor products $W_1 \boxtimes T_1, \ldots, W_s \boxtimes T_s$, $(W_1 \otimes \cdots \otimes W_s) \boxtimes (T_1 \otimes \cdots \otimes T_s)$, and

$$
(W_1 \boxtimes T_1) \otimes \cdots \otimes (W_s \boxtimes T_s) \simeq (W_1 \otimes \cdots \otimes W_s) \boxtimes (T_1 \otimes \cdots \otimes T_s).
$$
\n
$$
(6.6)
$$

Proof. By induction, it suffices to prove that it is true for $s = 2$. From Remark [2.19,](#page-8-0) there exist tensor products $W_1 \boxtimes T_1$ and $W_2 \boxtimes T_2$. By definition, we have an intertwining operator \mathcal{Y}_1 of type $W_1 \boxtimes T_1$ and an intertwining operator \mathcal{Y}_2 of type $\binom{W_2 \boxtimes T_2}{W_2 T_2}$, satisfying the universal property. Define a linear map

$$
\mathcal{Y}(\cdot,z)\cdot:(W_1\otimes W_2)\otimes(T_1\otimes T_2)\rightarrow((W_1\boxtimes T_1)\otimes(W_2\boxtimes T_2))\{z\}
$$

by

$$
\mathcal{Y}(w_1 \otimes w_2, z)(t_1 \otimes t_2) = \mathcal{Y}_1(w_1, z)t_1 \otimes \mathcal{Y}_2(w_2, z)t_2 \tag{6.7}
$$

for $w_i \in W_i$, $t_i \in T_i$ with $i = 1, 2$. Then \mathcal{Y} is an intertwining operator of type $\binom{(W_1 \boxtimes T_1) \otimes (W_2 \boxtimes T_2)}{W_1 \otimes W_2}$ (see the comments right before Theorem [2.16\)](#page-7-2). We claim that $((W_1 \boxtimes T_1) \otimes (W_2 \boxtimes T_2), \mathcal{Y})$ is a tensor product of $W_1 \otimes W_2$ and $T_1 \otimes T_2$. Let T be any τ -twisted $V_1 \otimes V_2$ -module and let I be any intertwining operator of type $\begin{pmatrix} T \\ W_1 \otimes W_2 \end{pmatrix}$. We must show that there exists a τ -twisted $V_1 \otimes V_2$ -module homomorphism ψ from $(W_1 \boxtimes T_1) \otimes (W_2 \boxtimes T_2)$ to \tilde{T} such that $I = \psi \circ \mathcal{Y}$. As $V_1 \otimes V_2$ is τ -rational, it suffices to consider the case with T an irreducible τ -twisted $V_1 \otimes V_2$ -module. From Remark [6.8,](#page-33-2) $T \simeq P_1 \otimes P_2$, where P_i is an irreducible τ_i -twisted V_i -module for $i = 1, 2$. As $\dim I_{V_i} {P_i \choose W_i | T_i} < \infty$ by assumption, from Theorem [2.16](#page-7-2) (due to [\[ADL\]](#page-35-16)), $I = \sum_{j=1}^{r} I_{1,j} \otimes I_{2,j}$, where $I_{1,j}$ are intertwining operators of type $\binom{P_1}{W_1\ T_1}$ and $I_{2,j}$ are intertwining operators of type $\binom{P_2}{W_2|T_2}$. By the universal property of $(W_i \boxtimes T_i, Y_i)$ for $i = 1, 2$, there exist τ_i twisted V_i -module homomorphisms $\psi_{i,j} : W_i \boxtimes T_i \to P_i$ such that $I_{i,j}(\cdot, z) = \psi_{i,j} \circ \mathcal{Y}_i(\cdot, z)$ for $i = 1, 2, 1 \le j \le r$. Set $\psi = \sum_{j=1}^r \psi_{1,j} \otimes \psi_{2,j}$, which is a τ -twisted $V_1 \otimes V_2$ -module homomorphism from $(W_1 \boxtimes T_1) \otimes (W_2 \boxtimes T_2)$ to T such that $I = \psi \circ \mathcal{Y}$. This proves that $((W_1 \boxtimes T_1) \otimes (W_2 \boxtimes T_2), \mathcal{Y})$ is a tensor product of $(W_1 \otimes W_2)$ with $T_1 \otimes T_2$. Thus tensor product $(W_1 \otimes W_2) \boxtimes (T_1 \otimes T_2)$ exists and $(W_1 \boxtimes T_1) \otimes (W_2 \boxtimes T_2) \simeq (W_1 \otimes W_2) \boxtimes (T_1 \otimes T_2)$. \Box

We now prove Theorem C.

Proof. Using Proposition [6.10](#page-34-0) and Theorem B, we obtain

$$
(M_1 \otimes \cdots \otimes M_k) \boxtimes T_{\sigma_{\kappa}}(N_1, N_2, \ldots, N_s)
$$

= $(M_1 \otimes \cdots \otimes M_k) \boxtimes (T_{\sigma_1}(N_1) \otimes \cdots \otimes T_{\sigma_s}(N_s))$

$$
\simeq (M^{[k_1]} \boxtimes T_{\sigma_1}(N_1)) \otimes \cdots \otimes (M^{[k_s]} \boxtimes T_{\sigma_s}(N_1))
$$

$$
\simeq T_{\sigma_1} (M^{[k_1]} \boxtimes N_1) \otimes \cdots \otimes T_{\sigma_s} (M^{[k_s]} \boxtimes N_s)
$$

= $T_{\sigma_{\kappa}} ((M^{[k_1]} \boxtimes N_1), \ldots, (M^{[k_s]} \boxtimes N_s)),$

proving the first assertion. Then using Proposition [6.7](#page-33-4) we get the second assertion.

 \Box

References

- [A1] T. Abe, C_2 -cofiniteness of the 2-cycle permutation orbifold models of minimal Virasoro vertex operator algebras, *Commun. Math. Phys.* 303 (2011), 825–844.
- [A2] T. Abe, C2-cofiniteness of 2-cyclic permutation orbifold models, *Commun. Math. Phys.* 317 (2013), 425–445.
- [ABD] T. Abe, G. Buhl and C. Dong, Rationality, Regularity, and C_2 -cofiniteness, *Trans. Amer. Math. Soc.* 356 (2004), 3391–3402.
- [ADL] T. Abe, C. Dong and H. Li, Fusion rules for the vertex operator algebras $M(1)^+$ and V_L^+ , *Commun. Math. Phys.* 253 (2005), 171–219.
- [ADJR] C. Ai, C. Dong, X. Jiao and L. Ren, The irreducible modules and fusion rules for the parafermion vertex operator algebras, *Trans. Amer. Math. Soc.* 370 (2018), 5963–5981.
- [Ba] P. Bantay, Characters and modular properties of permutation orbifolds, *Phys. Lett* B419 (1998), 175–178.
- [BK] B. Bakalov and A. Kirillov, Lectures on tensor categories and modular functors. *University Lecture Series* Vol 21, 2000.
- [BDM] K. Barron, C. Dong and G. Mason, Twisted sectors for tensor product vertex operator algebras associated to permutation groups, *Commun. Math. Phys.* 227 (2002), no. 2, 349–384.
- [Bo] R. E. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster. *Proc. Natl. Acad. Sci. USA* 83 (1986), 3068–3071.
- [BHL] K. Barron, Y.-Z.Huang and J. Lepowsky, An equivalence of two constructions of permutation-twisted modules for lattice vertex operator algebras, *J. Pure Appl. Algebra* 210 (2007), no. 3, 797–826.
- [BHS] L. Borisov, M.B. Halpern and C. Schweigert, Systematic approach to cyclic orbifolds, *Int. J. Mod. Phys.* A13 (1998), 125–168.
- [CM] S. Carnahan and M. Miyamoto, Regularity of fixed-point vertex operator subalgebras, [arXiv:1603.05645.](http://arxiv.org/abs/1603.05645)
- [CKM] T. Creutzig, S. Kanade and R. McRae, Tensor categories for vertex operator superalgebra extensions, [arXiv:1705.05017.](http://arxiv.org/abs/1705.05017)
- [DJX] C. Dong, X. Jiao and F. Xu, Quantum dimensions and quantum Galois theory, *Trans. Amer. Math. Soc.* 365 (2013), 6441–6469.
- [DL] C. Dong and J. Lepowsky, Generalized Vertex Algebras and Relative Vertex Operators, *Progress in Math.* Vol. 112, Birkhäuser, Boston, 1993.
- [DLM1] C. Dong, H. Li and G. Mason, Simple currents and extensions of vertex operator algebras, *Commun. Math. Phys.* 180 (1996) 671–707.
- [DLM2] C. Dong, H. Li and G. Mason, Twisted representations of vertex operator algebras, *Math. Ann.* 310 (1998), 571–600.
- [DLM3] C. Dong, H. Li and G. Mason, Modular-invariance of trace functions in orbifold theory and generalized moonshine, *Commun. Math. Phys.* 214 (2000), 1–56.
- [DLMi] B. Doyon, J. Lepowsky and A. Milas, Twisted modules for vertex operator algebras and Bernoulli polynomials, *Int. Math. Res. Not.* 44 (2003), 2391–2408.
- [DMNO] A. Davydov, M. Müger, D. Nikshych and V. Ostrik, The Witt group of non-degenerate braided fusion categories, *J. Reine Angew. Math.* 677 (2013), 135–177.
- [DRX] C. Dong, L. Ren and F. Xu, On orbifold theory, *Adv. Math.* 321 (2017), 1–30.
- [DXY1] C. Dong, F. Xu and N. Yu, 2-cyclic permutations of lattice vertex operator algebras, *Proc. Amer. Math. Soc.* 144 (2016), 3207–3220.
- [DXY2] C. Dong, F. Xu and N. Yu, 2-permutations of lattice vertex operator algebras: higher rank, *J. Algebra* 476 (2017), 1–25.
- [DXY3] C. Dong, F. Xu and N. Yu, The 3-permutation orbifold of a lattice vertex operator algebra, *J. Pure Appl. Algebra* 222 (2018), no. 6, 1316–1336.
- [ENO] P. Etingof, D. Nikshych and V. Ostrik, On fusion categories, *Ann. of Math.* 162 (2005), 581–642.
- [FHL] I. Frenkel, Y.-Z.Huang and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, *Mem. Amer. Math. Soc.* 104 (1993).
- [FLM] I. B. Frenkel, J. Lepowsky and A. Meurman, Vertex operator algebras and the Monster, *Pure and Applied Math.*134, Academic Press, Massachusetts, 1988.
- [H1] Y.-Z. Huang, A theory of tensor products for modules categories for a vertex operator algebra, IV, *J. Pure Appl. Algebra* 100 (1995), 173–216.
- [H2] Y.-Z. Huang, Rigidity and modularity of vertex tensor categories, *Commun. Contemp. Math.*10 (2008), suppl. 1, 871–911.
- [HL1] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for modules categories for a vertex operator algebra, I, II, *Selecta Mathmatica*, New Series 1 (1995), 699–756; 757–786.
- [HL2] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for modules categories for a vertex operator algebra, III, *J. Pure Appl. Algebra* 100 (1995), 141–171.
- [K1] A. Kirillov Jr., Modular categories and orbifold models II, 2001, [arXiv:math/0110221.](http://arxiv.org/abs/math/0110221)
- [K2] A. Kirillov Jr., Modular categories and orbifold models, *Commun. Math. Phys.* 229 (2002), 309–335.
- [KL] M. Karel and H. Li, Certain generating subspaces for vertex operator algebras, *J. Algebra* 217 (1999), 393– 421.
- [KLX] V. Kac, R. Longo and F. Xu, Solitons in affine and permutation orbifolds, *Commun. Math. Phys.* 253 (2005), 723–764.
- [KO] A. Kirillov Jr. and V. Ostrik, On a q-analogue of the McKay correspondence and the ADE classification of sl² conformal field theories, *Adv. Math.* 171 (2002), 183–227.
- [Le] J. Lepowsky, Calculus of twisted vertex operators, *Proc. Natl. Acad. Sci. USA* 82 (1985), 8295–8299.
- [Li1] H. Li, Symmetric invariant bilinear forms on vertex operator algebras, *J. Pure Appl. Algebra* 96 (1994), 279–297.
- [Li2] H. Li, Local systems of twisted vertex operators, vertex operator superalgebras and twisted modules. Moonshine, the Monster, and related topics (South Hadley, MA, 1994), 203–236, *Contemp. Math.* 193, Amer. Math. Soc., Providence, RI, 1996.
- [Li3] H. Li, An analogue of the Hom functor and a generalized nuclear democracy theorem, *Duke Math. J.* 93 (1998), no. 1, 73–114.
- [Li4] H. Li, Some finiteness properties of regular vertex operator algebras, *J. Algebra* 212 (1999), 495–514.
- [Li5] H.-S. Li, Nonlocal vertex algebras generated by formal vertex operators, *Selecta Math. (N.S.)* 11 (2005), 349–397.
- [LL] J. Lepowsky and H. Li, Introduction to vertex operator algebras and their representations, *Progress in Math.* Vol. 227, Birkhäuser, Boston, 2004.
- [LTW] H. Li, S. Tan and Q. Wang, Twisted modules for quantum vertex algebras, *J. Pure Appl. Algebra* 214 (2010), no. 3, 201–220.
- [M1] M. Miyamoto, A \mathbb{Z}_3 -orbifold theory of lattice vertex operator algebra and \mathbb{Z}_3 -orbifold constructions. Symmetries, integrable systems and representations, 319–344, *Springer Proc. Math. Stat.*, 40, Springer, Heidelberg, 2013.
- [M2] M. Miyamoto, C_2 -cofiniteness of cyclic-orbifold models, *Commun. Math. Phys.* 335 (2015), 1279–1286.
- [X] X. Xu, Intertwining operators for twisted modules of a colored vertex operator superalgebra, *J. Algebra* 175 (1995), no. 1, 241–273.
- [Z] Y. Zhu, Modular invariance of characters of vertex operator algebras, *J. Amer. Math. Soc.* 9 (1996), 237–302.

C. Dong: Department of Mathematics, University of California Santa Cruz, CA 95064 USA; dong@ucsc.edu H. Li: Department of Mathematical Sciences, Rutgers University, Camden, NJ 08102 USA; hli@camden.rutgers.edu F. Xu: Department of Mathematics, University of California, Riverside, CA 92521 USA; xufeng@math.ucr.edu N. Yu: School of Mathematical Sciences, Xiamen University, Fujian, 361005, CHINA; ninayu@xmu.edu.cn