

## Radiative leptonic decays on the lattice

---

Christopher Kane<sup>a</sup>, Christoph Lehner<sup>b,c</sup>, Stefan Meinel<sup>\*a,d</sup>, Amarjit Soni<sup>c</sup>

<sup>a</sup>*Department of Physics, University of Arizona, Tucson, AZ 85721, USA*

<sup>b</sup>*Department of Physics, University of Regensburg, 93040 Regensburg, Germany*

<sup>c</sup>*Physics Department, Brookhaven National Laboratory, Upton, NY 11973, USA*

<sup>d</sup>*RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, NY 11973, USA*


*E-mail:* [smeinel@email.arizona.edu](mailto:smeinel@email.arizona.edu)

Adding a hard photon to the final state of a leptonic pseudoscalar-meson decay lifts the helicity suppression and can provide sensitivity to a larger set of operators in the weak effective Hamiltonian. Furthermore, radiative leptonic  $B$  decays at high photon energy are well suited to constrain the first inverse moment of the  $B$ -meson light-cone distribution amplitude, an important parameter in the theory of nonleptonic  $B$  decays. We demonstrate that the calculation of radiative leptonic decays is possible using Euclidean lattice QCD, and present preliminary numerical results for  $D_s^+ \rightarrow \ell^+ \nu \gamma$  and  $K^- \rightarrow \ell^- \bar{\nu} \gamma$ .

*37th International Symposium on Lattice Field Theory - Lattice2019  
16-22 June 2019  
Wuhan, China*

---

\*Speaker.



**Figure 1:** Left: A diagram contributing to  $B^- \rightarrow \ell^- \bar{\nu} \gamma$ , where the green square corresponds to  $W$ -boson exchange in the Standard Model and comes with a factor of  $V_{ub}$ . Right: two diagrams contributing to  $B_s \rightarrow \ell^+ \ell^- \gamma$ , via the operators  $O_{7,9,10}$  (defined, for example, in Ref. [1]).

## 1. Introduction

Radiative leptonic decays of pseudoscalar mesons probe both the weak interaction and the hadronic structure in useful ways. Adding a sufficiently energetic photon to the final state can actually increase the branching fraction [2], as it removes the helicity suppression. Perhaps the most interesting example is  $B^- \rightarrow \ell^- \bar{\nu} \gamma$ , shown in Fig. 1 (left). For large  $E_\gamma^{(0)}$ , this process is the cleanest probe of the first inverse moment of the  $B$ -meson light-cone distribution amplitude,  $1/\lambda_B = \int_0^\infty \frac{\Phi_{B^+}(\omega)}{\omega} d\omega$ , an important input in QCD-factorization predictions for nonleptonic  $B$  decays that is presently poorly determined [3, 4, 5, 6, 7, 8, 9]. A recent search for this decay by Belle gave an upper limit  $\mathcal{B}(B^- \rightarrow \ell^- \bar{\nu} \gamma, E_\gamma^{(0)} > 1 \text{ GeV}) < 3.0 \times 10^{-6}$ , close to the Standard-Model expectation [10]. Lattice QCD results for the  $B^- \rightarrow \ell^- \bar{\nu} \gamma$  form factors could be used to constrain  $\lambda_B$ . Also very interesting are the flavor-changing neutral-current decays  $B^0 \rightarrow \ell^+ \ell^- \gamma$  and  $B_s \rightarrow \ell^+ \ell^- \gamma$  (shown in Fig. 1, right). While the purely leptonic decays are sensitive to  $C_{10,S,P} - C'_{10,S,P}$  only, the radiative leptonic decays probe all Wilson coefficients in the weak effective Hamiltonian, including  $C_9$ , in which global fits of experimental results for other  $b \rightarrow s \ell^+ \ell^-$  decays indicate a deviation from the Standard Model that violates lepton flavor universality (LFU) [1]. Since the radiative leptonic decays are not helicity-suppressed, they are well-suited for testing LFU with light leptons [11, 12]. For the charmed-meson radiative leptonic decays  $D^+ \rightarrow e^+ \nu \gamma$  and  $D_s^+ \rightarrow e^+ \nu \gamma$ , the BESIII collaboration has reported upper limits on the branching fractions with  $E_\gamma^{(0)} > 10 \text{ MeV}$  of  $3.0 \times 10^{-5}$  and  $1.3 \times 10^{-4}$ , respectively [13, 14]. Finally, in contrast to the heavy-meson decays, there are already precise measurements of the differential branching fractions of  $K^- \rightarrow e^- \bar{\nu} \gamma$ ,  $K^- \rightarrow \mu^- \bar{\nu} \gamma$ ,  $\pi^- \rightarrow e^- \bar{\nu} \gamma$ , and  $\pi^- \rightarrow \mu^- \bar{\nu} \gamma$ , as reviewed in Ref. [15]. These decay modes can therefore be used to test the lattice QCD methods.

In the following, we show how radiative leptonic decays can be calculated on a Euclidean lattice, and we present early numerical results. One of us previously reported on this project at the Lattice 2018 conference [16]. At Lattice 2019, radiative leptonic decays were also discussed by G. Martinelli [17].

## 2. Hadronic tensor and form factors

To define the form factors for charged-current radiative leptonic decays of pseudoscalar mesons, we use the notation for  $B^- \rightarrow \ell^- \bar{\nu} \gamma$ . The quark electromagnetic and weak currents are given by  $J_\mu = \sum_q e_q \bar{q} \gamma_\mu q$  and  $J_\mu^{\text{weak}} = \bar{u} \gamma_\mu (1 - \gamma_5) b$ . The decay amplitude depends on the hadronic tensor,

which is defined as

$$T_{\mu\nu} = -i \int d^4x e^{ip_\gamma x} \langle 0 | T (J_\mu(x) J_\nu^{\text{weak}}(0)) | B^-(\mathbf{p}_B) \rangle \quad (2.1)$$

in Minkowski space. Throughout this work, we assume that the photon is real, i.e.,  $p_\gamma^2 = 0$ . The hadronic tensor can be decomposed as [7]

$$T_{\mu\nu} = \varepsilon_{\mu\nu\tau\rho} p_\gamma^\tau v^\rho F_V + i[-g_{\mu\nu}(p_\gamma \cdot v) + v_\mu(p_\gamma)_\nu] F_A - i \frac{v_\mu v_\nu}{p_\gamma \cdot v} m_B f_B + (p_\gamma)_\mu \text{-terms}, \quad (2.2)$$

where  $p_B = m_B v$  and the  $(p_\gamma)_\mu$ -terms will disappear when contracting with the photon polarization vector. The form factors  $F_V$  and  $F_A$  are functions of the photon energy in the  $B$ -meson rest frame,  $E_\gamma^{(0)} = p_\gamma \cdot v = (m_B^2 - q^2)/(2m_B)$ . Also appearing in Eq. (2.2) is the  $B$ -meson decay constant  $f_B$ .

To prepare for the discussion in the next section, it is useful to write down the spectral representation of  $T_{\mu\nu}$  in Minkowski space for the two different time orderings of the currents. By inserting complete sets of energy/momentum eigenstates and performing the time integrals, we find

$$\begin{aligned} T_{\mu\nu}^< &= -i \int_{-\infty(1-i\varepsilon)}^0 dt e^{iE_\gamma t} \int d^3x e^{-i\mathbf{p}_\gamma \cdot \mathbf{x}} \langle 0 | J_\nu^{\text{weak}}(0) J_\mu(t, \mathbf{x}) | B^-(\mathbf{p}_B) \rangle \\ &= -\sum_n \frac{1}{2E_{n,(\mathbf{p}_B-\mathbf{p}_\gamma)}} \frac{\langle 0 | J_\nu^{\text{weak}}(0) | n(\mathbf{p}_B - \mathbf{p}_\gamma) \rangle \langle n(\mathbf{p}_B - \mathbf{p}_\gamma) | J_\mu(0) | B(\mathbf{p}_B) \rangle}{E_\gamma + E_{n,(\mathbf{p}_B-\mathbf{p}_\gamma)} - E_B - i\varepsilon}, \end{aligned} \quad (2.3)$$

$$\begin{aligned} T_{\mu\nu}^> &= -i \int_0^{\infty(1-i\varepsilon)} dt e^{iE_\gamma t} \int d^3x e^{-i\mathbf{p}_\gamma \cdot \mathbf{x}} \langle 0 | J_\mu(t, \mathbf{x}) J_\nu^{\text{weak}}(0) | B^-(\mathbf{p}_B) \rangle \\ &= \sum_m \frac{1}{2E_{m,\mathbf{p}_\gamma}} \frac{\langle 0 | J_\mu(0) | m(\mathbf{p}_\gamma) \rangle \langle m(\mathbf{p}_\gamma) | J_\nu^{\text{weak}}(0) | B(\mathbf{p}_B) \rangle}{E_\gamma - E_{m,\mathbf{p}_\gamma} - i\varepsilon} \end{aligned} \quad (2.4)$$

(in infinite volume, the sums over  $n$  and  $m$  include integrals over the continuous spectrum of multi-particle states).

### 3. Extracting the hadronic tensor from a Euclidean three-point function

In this section, we show that  $T_{\mu\nu}$  can be extracted from the Euclidean three-point function

$$C_{\mu\nu}(t, t_B) = \int d^3x \int d^3y e^{-i\mathbf{p}_\gamma \cdot \mathbf{x}} e^{i\mathbf{p}_B \cdot \mathbf{y}} \left\langle J_\mu(t, \mathbf{x}) J_\nu^{\text{weak}}(0, \mathbf{0}) \phi_B^\dagger(t_B, \mathbf{y}) \right\rangle, \quad (3.1)$$

where  $\phi_B \sim \bar{u}\gamma_5 b$  is an interpolating field for the  $B$  meson, and  $t, t_B$  now denote the Euclidean time. We define the integrals

$$I_{\mu\nu}^<(t_B, T) = \int_{-T}^0 dt e^{E_\gamma t} C_{\mu\nu}(t, t_B), \quad I_{\mu\nu}^>(t_B, T) = \int_0^T dt e^{E_\gamma t} C_{\mu\nu}(t, t_B), \quad (3.2)$$

with a finite integration range  $T$ . Here we take  $t_B$  to be large and negative (with  $t_B < -T$ ), such that ground-state saturation is achieved for the  $B$  meson. Inserting again complete sets of energy/momentum eigenstates, we find, for the first time ordering,

$$\begin{aligned} I_{\mu\nu}^<(t_B, T) &= \langle B(\mathbf{p}_B) | \phi_B^\dagger(0) | 0 \rangle \frac{1}{2E_B} e^{E_B t_B} \\ &\times \sum_n \frac{1}{2E_{n,(\mathbf{p}_B-\mathbf{p}_\gamma)}} \frac{\langle 0 | J_\nu^{\text{weak}}(0) | n(\mathbf{p}_B - \mathbf{p}_\gamma) \rangle \langle n(\mathbf{p}_B - \mathbf{p}_\gamma) | J_\mu(0) | B(\mathbf{p}_B) \rangle}{E_\gamma + E_{n,(\mathbf{p}_B-\mathbf{p}_\gamma)} - E_B} \\ &\times \left( 1 - e^{-(E_\gamma + E_{n,(\mathbf{p}_B-\mathbf{p}_\gamma)} - E_B)T} \right). \end{aligned} \quad (3.3)$$

The sum over states in Eq. (3.3) differs from the sum in Eq. (2.3) by the factor in the last line. However, the exponential  $e^{-(E_\gamma + E_{n,(\mathbf{p}_B - \mathbf{p}_\gamma)} - E_B)T}$  will vanish for large  $T$  if  $E_\gamma + E_{n,(\mathbf{p}_B - \mathbf{p}_\gamma)} > E_B$ . Because the states  $|n(\mathbf{p}_B - \mathbf{p}_\gamma)\rangle$  have the same quark-flavor quantum numbers as the  $B$  meson, we have  $E_{n,(\mathbf{p}_B - \mathbf{p}_\gamma)} \geq E_{B,(\mathbf{p}_B - \mathbf{p}_\gamma)} = \sqrt{m_B^2 + (\mathbf{p}_B - \mathbf{p}_\gamma)^2}$ . Thus, we need  $\sqrt{\mathbf{p}_\gamma^2} + \sqrt{m_B^2 + (\mathbf{p}_B - \mathbf{p}_\gamma)^2} > \sqrt{m_B^2 + \mathbf{p}_B^2}$ . This is in fact always true if  $\mathbf{p}_\gamma \neq 0$ .

For the other time ordering, we find

$$I_{\mu\nu}^>(t_B, T) = -\langle B(\mathbf{p}_B) | \phi_B^\dagger(0) | 0 \rangle \frac{1}{2E_B} e^{E_B t_B} \times \sum_m \frac{1}{2E_{m,\mathbf{p}_\gamma}} \frac{\langle 0 | J_\mu(0) | m(\mathbf{p}_\gamma) \rangle \langle m(\mathbf{p}_\gamma) | J_\nu^{\text{weak}}(0) | B(\mathbf{p}_B) \rangle}{E_\gamma - E_{m,\mathbf{p}_\gamma}} \left( 1 - e^{(E_\gamma - E_{m,\mathbf{p}_\gamma})T} \right). \quad (3.4)$$

The unwanted exponential  $e^{(E_\gamma - E_{m,\mathbf{p}_\gamma})T}$  in the last line goes to zero for large  $T$  if  $E_{m,\mathbf{p}_\gamma} > E_\gamma$ . Because the states  $|m(\mathbf{p}_\gamma)\rangle$  are hadronic and have nonzero masses, their energies are larger than the energy of a photon with the same spatial momentum, showing that this condition is also always satisfied. In summary, for  $\mathbf{p}_\gamma \neq 0$ ,


$$T_{\mu\nu} = -\lim_{T \rightarrow \infty} \lim_{t_B \rightarrow -\infty} \frac{2E_B e^{-E_B t_B}}{\langle B(\mathbf{p}_B) | \phi_B^\dagger(0) | 0 \rangle} I_{\mu\nu}(t_B, T), \quad (3.5)$$

where  $I_{\mu\nu}$  is the integral from  $-T$  to  $T$ . The energy  $E_B$  and the overlap factor  $\langle B(\mathbf{p}_B) | \phi_B^\dagger(0) | 0 \rangle$  can be obtained from the two-point function  $\int d^3x e^{-i\mathbf{p}_B \cdot \mathbf{x}} \langle \phi_B(t, \mathbf{x}) \phi_B^\dagger(0) \rangle$ .


Note that similar nonlocal matrix elements appear in processes with two photons, whose lattice calculation has been discussed, for example, in Refs. [18, 19, 20].

#### 4. Preliminary numerical results


In this section, we present some early numerical results for the  $D_s^+ \rightarrow \ell^+ \nu \gamma$  and  $K^- \rightarrow \ell^- \bar{\nu} \gamma$  form factors. These results are from only 25 configurations of the ‘‘24I’’ RBC/UKQCD ensemble



**Figure 2:** The un-integrated, scaled three-point functions  $-\frac{2E_{D_s} e^{-E_{D_s} t_{D_s}}}{\langle D_s(\mathbf{p}_{D_s}) | \phi_{D_s}^\dagger(0) | 0 \rangle} C_{\mu\nu}(t, t_{D_s})$  as a function of the electromagnetic-current insertion time  $t$ , for  $t_{D_s}/a = -12$  and  $\mathbf{p}_\gamma = (0, 0, 1) \frac{2\pi}{L}$ . The left plot shows a combination of indices sensitive to  $F_V$ , while the right plot shows a combination sensitive to  $F_A$ . The contributions from the  $s$  and  $c$  quark in the electromagnetic current are shown separately, without charge factors.




**Figure 3:** The  $D_s^+ \rightarrow \ell^+ \nu \gamma$  and  $K^- \rightarrow \ell^- \bar{\nu} \gamma$  form factors at  $\mathbf{p}_\gamma = (0, 0, 1) \frac{2\pi}{L}$  as a function of the summation range  $T$ , for two different meson-field insertion times.



**Figure 4:** The  $D_s$  and  $K$  decay constants extracted from  $T_{\mu\nu}$  at  $\mathbf{p}_\gamma = (0, 0, 1) \frac{2\pi}{L}$ , as a function of the summation range  $T$ , for two different meson-field insertion times. For the  $D_s$ , the horizontal line shows the physical value from Ref. [21]. For the  $K$ , the horizontal line shows the value computed on the same ensemble with the standard method in Ref. [22].

[22] with  $2 + 1$  flavors of domain-wall fermions and the Iwasaki gauge action, with  $a^{-1} = 1.785(5)$  GeV and  $m_\pi = 340(1)$  MeV. For the light and strange valence quarks, we use the same domain-wall action as in Ref. [22]. The valence charm quark is implemented with a Möbius domain-wall action with stout-smear gauge links ( $N = 3$ ,  $\rho = 0.1$ ),  $L_5/a = 12$ ,  $aM_5 = 1.0$ ,  $am_f = 0.6$  [23], which approximately corresponds to the physical charm-quark mass. We use local currents with “mostly nonperturbative” renormalization. Gaussian smearing is performed for the lighter quark in the meson interpolating field. We start with a  $\mathbb{Z}_2$  random-wall source at the time slice of the weak current (denoted as time “0” here) and perform sequential inversions through the meson in-



**Figure 5:** The  $D_s^+ \rightarrow \ell^+ \nu \gamma$  and  $K^- \rightarrow \ell^- \bar{\nu} \gamma$  form factors as a function of the photon energy. The results shown here were obtained with  $T/a = 8$  and  $t_{K/D_s}/a = -12$ . Only the statistical uncertainties are given.

terpolating field; disconnected diagrams are presently neglected. All-mode averaging [24] with 16 sloppy and 1 exact samples per configuration is employed; the 16 sloppy samples correspond to 16 different starting time slices. Our initial calculations used  $\mathbf{p}_{K/D_s} = 0$  and  $\mathbf{p}_\gamma^2 \in \{1, 2, 3, 4, 5\} \left(\frac{2\pi}{L}\right)^2$ .

Figure 2 shows examples of the  $D_s^+ \rightarrow \ell^+ \nu \gamma$  three-point functions. Multiplying by  $e^{E_\gamma t}$  and summing over  $t$  gives  $T_{\mu\nu}$  for sufficiently large summation range  $T$ . The form factors  $F_V$  and  $F_A$  extracted from  $T_{\mu\nu}$  (at the lowest photon momentum) are shown as a function of  $T$  in Fig. 3. The results plateau at approximately  $T/a = 8$ . We also extracted the meson decay constants from the  $v^\mu v^\nu$  term in  $T_{\mu\nu}$ . As can be seen in Fig. 4, the results agree with the known values, which is a valuable test of our calculation. Finally, Fig. 5 shows the form factors  $F_V$  and  $F_A$  as a function of the photon energy. Note that, with our current choice of momenta, all of the photon energies are above the physical region for  $K^- \rightarrow \ell^- \bar{\nu} \gamma$ . The results for  $F_A$  are dominated by the point-like contribution equal to  $-e_\ell f_{K/D_s}/E_\gamma^{(0)}$ .

## 5. Conclusions and Outlook

We have shown that the form factors describing radiative leptonic decays can be calculated on the lattice; even though they involve a nonlocal matrix element, the use of imaginary time poses no difficulty in this case. The early results shown here for  $D_s^+ \rightarrow \ell^+ \nu \gamma$  and  $K^- \rightarrow \ell^- \bar{\nu} \gamma$  cover photon energies from approximately 0.5 to 1 GeV. For  $K^- \rightarrow \ell^- \bar{\nu} \gamma$  we need to reach lower photon energies to compare with experiment; this can be achieved by using moving frames (i.e., nonzero  $\mathbf{p}_K$ ) and/or a larger volume. To study the  $B_{(s)}$  radiative leptonic decays with the domain-wall action for the heavy quark, we will need to extrapolate in the mass. We are also considering calculations directly at the physical  $b$ -quark mass using the “relativistic heavy-quark action” [25], but, because this action is only on-shell improved, additional steps are likely needed to remove unphysical behavior occurring when the electromagnetic and weak currents get close to each other.

**Acknowledgments:** We thank the RBC and UKQCD Collaborations for providing the gauge-field configurations. C.K. and S.M. are supported by the US DOE, Office of Science, Office of HEP under award number DE-SC0009913. S.M. is also supported by the RIKEN BNL Research Center. A.S. and C.L. are supported in part by US DOE Contract No. DESC0012704(BNL). During a part of this work, C.L. was also supported by a DOE Office of Science Early Career Award. This work used resources at TACC that are part of XSEDE, supported by NSF grant number ACI-1548562.

- [1] J. Aebischer, W. Altmannshofer, D. Guadagnoli, M. Reboud, P. Stangl, and D. M. Straub, “ $B$ -decay discrepancies after Moriond 2019,” [arXiv:1903.10434](#).
- [2] D. Atwood, G. Eilam, and A. Soni, “Pure leptonic radiative decays  $B^\pm, D_s \rightarrow \ell \nu \gamma$  and the annihilation graph,” *Mod. Phys. Lett.* **A11** (1996) 1061, [arXiv:hep-ph/9411367](#).
- [3] G. P. Korchemsky, D. Pirjol, and T.-M. Yan, “Radiative leptonic decays of  $B$  mesons in QCD,” *Phys. Rev.* **D61** (2000) 114510, [arXiv:hep-ph/9911427](#).
- [4] S. Descotes-Genon and C. T. Sachrajda, “Factorization, the light cone distribution amplitude of the  $B$  meson and the radiative decay  $B \rightarrow \gamma \ell \nu_\ell$ ,” *Nucl. Phys.* **B650** (2003) 356, [arXiv:hep-ph/0209216](#).
- [5] E. Lunghi, D. Pirjol, and D. Wyler, “Factorization in leptonic radiative  $B \rightarrow \gamma e \nu$  decays,” *Nucl. Phys.* **B649** (2003) 349, [arXiv:hep-ph/0210091](#).
- [6] V. M. Braun and A. Khodjamirian, “Soft contribution to  $B \rightarrow \gamma \ell \nu_\ell$  and the  $B$ -meson distribution amplitude,” *Phys. Lett.* **B718** (2013) 1014, [arXiv:1210.4453](#).
- [7] M. Beneke, V. M. Braun, Y. Ji, and Y.-B. Wei, “Radiative leptonic decay  $B \rightarrow \gamma \ell \nu_\ell$  with subleading power corrections,” *JHEP* **07** (2018) 154, [arXiv:1804.04962](#).
- [8] Y.-M. Wang and Y.-L. Shen, “Subleading-power corrections to the radiative leptonic  $B \rightarrow \gamma \ell \nu$  decay in QCD,” *JHEP* **05** (2018) 184, [arXiv:1803.06667](#).
- [9] M. Beneke, G. Buchalla, M. Neubert, and C. T. Sachrajda, “QCD factorization for  $B \rightarrow \pi\pi$  decays: Strong phases and CP violation in the heavy quark limit,” *Phys. Rev. Lett.* **83** (1999) 1914, [arXiv:hep-ph/9905312](#).
- [10] Belle Collaboration, M. Gelb *et al.*, “Search for the rare decay of  $B^+ \rightarrow \ell^+ \nu_\ell \gamma$  with improved hadronic tagging,” *Phys. Rev.* **D98** (2018) 112016, [arXiv:1810.12976](#).
- [11] D. Guadagnoli, M. Reboud, and R. Zwicky, “ $B_s^0 \rightarrow \ell^+ \ell^- \gamma$  as a test of lepton flavor universality,” *JHEP* **11** (2017) 184, [arXiv:1708.02649](#).
- [12] F. Dettori, D. Guadagnoli, and M. Reboud, “ $B_s^0 \rightarrow \mu^+ \mu^- \gamma$  from  $B_s^0 \rightarrow \mu^+ \mu^-$ ,” *Phys. Lett.* **B768** (2017) 163, [arXiv:1610.00629](#).
- [13] BESIII Collaboration, M. Ablikim *et al.*, “Search for the radiative leptonic decay  $D^+ \rightarrow \gamma e^+ \nu_e$ ,” *Phys. Rev.* **D95** (2017) 071102, [arXiv:1702.05837](#).
- [14] BESIII Collaboration, M. Ablikim *et al.*, “Search for the decay  $D_s^+ \rightarrow \gamma e^+ \nu_e$ ,” *Phys. Rev.* **D99** (2019) 072002, [arXiv:1902.03351](#).
- [15] Particle Data Group Collaboration, M. Tanabashi *et al.*, “Review of Particle Physics,” *Phys. Rev.* **D98** (2018) 030001.
- [16] A. Soni, “Flavor anomalies & the lattice,” 2018. Presentation at Lattice 2018, East Lansing, MI, USA.
- [17] RM123 Collaboration, G. Martinelli, “Electromagnetic Corrections to Decay Amplitudes: Real Emissions in Leptonic Decays,” 2019. Presentation at Lattice 2019, Wuhan, China.
- [18] X.-d. Ji and C.-w. Jung, “Studying hadronic structure of the photon in lattice QCD,” *Phys. Rev. Lett.* **86** (2001) 208, [arXiv:hep-lat/0101014](#).
- [19] J. J. Dudek and R. G. Edwards, “Two Photon Decays of Charmonia from Lattice QCD,” *Phys. Rev. Lett.* **97** (2006) 172001, [arXiv:hep-ph/0607140](#).
- [20] X. Feng, S. Aoki, H. Fukaya, S. Hashimoto, T. Kaneko, J.-i. Noaki, and E. Shintani, “Two-photon decay of the neutral pion in lattice QCD,” *Phys. Rev. Lett.* **109** (2012) 182001, [arXiv:1206.1375](#).
- [21] Flavour Lattice Averaging Group Collaboration, S. Aoki *et al.*, “FLAG Review 2019,” [arXiv:1902.08191](#).
- [22] RBC/UKQCD Collaboration, Y. Aoki *et al.*, “Continuum Limit Physics from 2+1 Flavor Domain Wall QCD,” *Phys. Rev.* **D83** (2011) 074508, [arXiv:1011.0892](#).
- [23] RBC/UKQCD Collaboration, P. A. Boyle, L. Del Debbio, N. Garron, A. J ajjtner, A. Soni, J. T. Tsang, and O. Witzel, “SU(3)-breaking ratios for  $D_{(s)}$  and  $B_{(s)}$  mesons,” [arXiv:1812.08791](#).
- [24] E. Shintani, R. Arthur, T. Blum, T. Izubuchi, C. Jung, and C. Lehner, “Covariant approximation averaging,” *Phys. Rev.* **D91** (2015) 114511, [arXiv:1402.0244](#).
- [25] N. H. Christ, M. Li, and H.-W. Lin, “Relativistic Heavy Quark Effective Action,” *Phys. Rev.* **D76** (2007) 074505, [arXiv:hep-lat/0608006](#).