# **P**<sub>CN</sub> calculations for Z=111 to Z=118

W. Loveland<sup>1</sup> and Liangyu Yao<sup>1</sup>

<sup>1</sup> Oregon State University, Corvallis, OR 97331 USA

Abstract. In previous publications [1,2] we presented evidence for the importance of spin in determining capture and evaporation residue cross sections in the synthesis of heavy nuclei. We extend the previous calculations which dealt with nuclei where  $Z_{CN} \le 110$  to the region of  $Z_{CN} = 111-118$ . We deduce a new systematics of the fusion probability  $P_{CN}$  for these reactions.

**Keywords:** P<sub>CN</sub>, superheavy element synthesis, spin mediated capture and evaporation residue cross sections.

## 1 Introduction

The cross section for producing a heavy evaporation residue in a complete fusion reaction can be written as a non-separable product of three factors, which express the capture cross section, the fusion probability and the survival probability.

$$\sigma_{EVR} = \frac{\pi h^2}{2\mu E} \sum_{\ell=0} (2\ell+1) T(E,\ell) P_{CN} W_{sur}(E,\ell)$$
(1)

Each of these factors is dependent on the spin, but the survival probability,  $W_{sur}$ , is zero or very small for higher spin values, effectively limiting the capture and fusion terms. Many partial waves contribute to the capture cross sections, but the higher partial waves result in non-surviving events. In this work, we examine the impact of restrictions on spin placed by the survival probabilities for compound nuclear reactions resulting in the synthesis of superheavy nuclei with  $Z_{CN}$ =111-118. In doing so, we extend the previous work [1,2] to treat the synthesis of the heaviest nuclei with  $Z_{CN}$ =111-118.

## 2 Methodology

As explained in [1], the formalism for calculating the survival, against fission, of a highly excited nucleus is relatively well-understood [3]. One starts with a single particle model [4] of the level density in which one allows the level density parameter to be a function of the excitation energy. Masses and shell corrections are taken from [5]. The deformation dependent collective enhancement of the level density is taken from [6]. The decay widths for decay by neutron, charged particle and  $\gamma$ -emission are calculated with standard formulas. Corrections for Kramers effects [7] are made to

the fission widths. The fission barrier heights are calculated using liquid drop barriers and excitation energy dependent shell corrections.

We begin with the compilation of Duellmann of evaluated evaporation residue cross sections for reactions that produce nuclei with  $Z_{CN}$  =111-118 [8]. For each reaction (projectile, target and beam energy), we calculated the spin dependent evaporation residue cross section assuming  $P_{CN}$ =1using the "Empirical Model" of [3]. In [1], we presented evidence that this procedure results in a reasonable agreement between the calculated and measured spin dependence of the evaporation residue formation cross sections for the test case of <sup>176</sup>Yb(<sup>48</sup>Ca,4n) <sup>220</sup>Th reaction and for the <sup>48</sup>Ca + <sup>208</sup>Pb reaction. Loveland [9] has made a detailed examination of the strengths and weaknesses of models such as [3] and placed limits on how well these models work.

### 3. Results

There are 28 cases we have examined. A summary of the measured and calculated evaporation residue cross sections is given in Table 1. The fusion probability,  $P_{CN}$ , is taken as the ratios of the calculated to the measured evaporation residue cross sections since we have assumed  $P_{CN}$  =1 in our calculations. As expected, the  $P_{CN}$  values for the "cold fusion" reactions (1 n out) are orders of magnitude smaller than those for the hot fusion (2n-4n out) reactions. The deduced values of  $P_{CN}$ generally get smaller as the product of the atomic numbers of the colliding nuclei,  $Z_1Z_2$ , increase.

Beam	Target	Channel	$\sigma_{meas}(pb)$	$\sigma_{calc}(pb)$	P <sub>CN</sub>	Ref
<sup>64</sup> Ni	<sup>209</sup> Bi	ln	$3.5^{+1.9}_{-1.3}$	6910	0.000507	10
<sup>65</sup> Cu	<sup>208</sup> Pb	ln	$1.7^{+3.9}_{-1.4}$	20500	8.3e-05	11
<sup>48</sup> Ca	<sup>238</sup> U	3n	$2.5^{+1.8}_{-1.1}$	60	0.0417	12
<sup>48</sup> Ca	<sup>238</sup> U	4n	$0.7^{+0.6}_{-0.3}$	425	0.00169	13
<sup>48</sup> Ca	<sup>238</sup> U	4n	$0.6^{+1.6}_{-0.5}$	7	0.0857	12
<sup>70</sup> Zn	<sup>208</sup> Pb	ln	$0.5^{+1.1}_{-0.4}$	5e+06	1e-07	14
<sup>48</sup> Ca	<sup>237</sup> Np	3n	0.9	5	0.18	15
<sup>70</sup> Zn	<sup>209</sup> Bi	ln	$0.022^{+0.020}$	940000	2.34e-08	16
			0.013			
<sup>48</sup> Ca	<sup>239</sup> Pu	3n	0.23	16	0.0144	17
<sup>48</sup> Ca	<sup>240</sup> Pu	3n	$2.5^{+2.9}_{-1.4}$	62	0.0403	17
<sup>48</sup> Ca	<sup>242</sup> Pu	2n	0.5	244	0.00205	
<sup>48</sup> Ca	<sup>242</sup> Pu	3n	$3.6^{+3.4}_{-1.7}$	78	0.0463	12
<sup>48</sup> Ca	<sup>242</sup> Pu	4n	$4.5^{+3.6}_{-1.9}$	129	0.0349	12
<sup>48</sup> Ca	<sup>242</sup> Pu	5n	$0.6^{+0.9}$ -0.5	11.6	0.0517	18

Table 1. Measured and calculated evaporation residue cross sections for  $Z_{CN} = 111-118$ 

<sup>48</sup> Ca	<sup>244</sup> Pu	3n	8 <sup>+7.4</sup> -4.5	180	0.0444	19
<sup>48</sup> Ca	<sup>244</sup> Pu	4n	$9.8^{+3.9}_{-3.1}$	220	0.0445	19
<sup>48</sup> Ca	<sup>244</sup> Pu	5n	$1.1^{+2.6}_{-0.9}$	9.2	0.120	20
<sup>48</sup> Ca	<sup>243</sup> Am	2n	$2.5^{+2.7}_{-1.5}$	15.4	0.162	21
<sup>48</sup> Ca	<sup>243</sup> Am	3n	$8.5^{+6.4}_{-3.7}$	660	0.0129	22
<sup>48</sup> Ca	<sup>243</sup> Am	4n	$0.9^{+3.2}_{-0.8}$	169	0.00533	23
<sup>48</sup> Ca	<sup>245</sup> Cm	2n	0.9	6.89	0.131	20
<sup>48</sup> Ca	<sup>245</sup> Cm	3n	$3.7^{+3.6}_{-1.8}$	229	0.0162	24
<sup>48</sup> Ca	<sup>245</sup> Cm	4n	0.8	95	0.00842	24
<sup>48</sup> Ca	<sup>248</sup> Cm	3n	1.2	166	0.00723	12
<sup>48</sup> Ca	<sup>248</sup> Cm	4n	3.4	652	0.00522	25
<sup>48</sup> Ca	<sup>249</sup> Bk	3n	$1.1^{+1.2}_{-0.6}$	1660	0.000663	26
<sup>48</sup> Ca	<sup>249</sup> Bk	4n	$2.4^{+3.3}_{-1.4}$	333	0.00721	26
<sup>48</sup> Ca	<sup>249</sup> Cf	2n	0.9	50.9	0.0177	24

In Figure 1, we show the  $P_{CN}$  values, sorted by exit channel for the hot fusion reactions, as a function of the simple scaling variable,  $Z_1Z_2$ , the product of the atomic numbers of the reacting nuclei.



Figure 1. The calculated values of  $P_{CN}$  for various exit channels as a function of the scaling variable  $Z_1Z_2.$ 

The use of other scaling variables, such as  $x_{CN}$ ,  $x_{eff}$  and  $x_m$  does not significantly improve the description of the data.  $x_{CN}$  is defined as

$$x_{CN} = \frac{Z_{CN}^2 / A_{CN}}{50.883(1 - 1.7826(\frac{A_{CN} - 2Z_{CN}}{A_{CN}})^2)} (2)$$

 $x_{\rm eff}$  is defined as

$$x_{eff} = \frac{\frac{4Z_P Z_T}{A_p^{\frac{1}{3}} A_T^{\frac{1}{3}} \left(A_P^{\frac{1}{3}} + A_T^{\frac{1}{3}}\right)}}{50.883(1 - 1.7826(\left(\frac{A_{CN} - 2Z_{CN}}{A_{CN}}\right)^2$$

 $x_m$  is defined as

 $x_m = 0.25 x_{CN} + 0.75 x_{eff}$ 

All of these scaling variables seek to relate  $P_{CN}$  to the balance of attractive and repulsive forces in the reaction entrance channel. Clearly there is a certain amount of "spatter" in the plots of  $P_{CN}$  vs.  $Z_1Z_2$ . In part, this "spatter" is due to the uncertainties in the measured evaporation residue cross sections which are typically uncertain to the measured value. (Loveland [9] has shown that these uncertainties in  $P_{CN}$  can lead to order of magnitude uncertainties in estimations of the production cross sections for elements 119 and 120, challenging experimentalists dealing with fb production cross sections.)

If we use the simple  $Z_1Z_2$  scaling factor for the 3n and 4n reactions, then we can write a simple formula for the 3n channel as  $P_{CN}(3n) = -0.019Z_1Z_2 + 35.0$  and for the 4n channel  $P_{CN}(4n) = -0.013Z_1Z_2 + 23.2$ .

We can ask how well these new values of  $P_{CN}$  agree with previous measurements and theoretical predictions. Kozulin et al. [27] have reported measurements of  $P_{CN}$ based upon mass-energy distributions of fission-like fragments from a variety of reactions. In Figure 2, we compare our values of  $P_{CN}$  with the Kozulin et al. measurements. Given the intrinsic large uncertainties in our deduced  $P_{CN}$  values, the agreement between the measurements seems satisfactory.

How do our measured values of  $P_{CN}$  compare with various theoretical predictions of  $P_{CN}$ ? Given our methodology, there is no surprise that our deduced values of  $P_{CN}$ agree well with the predictions of Zagrebaev [29]. How about other predictions? In Figure 3, we compare our deduced values of  $P_{CN}$  with predictions of Nasirov et al.[28]. For the hot fusion reactions ( $Z_1Z_2 = 1800-2000$ ), the agreement seems reasonable but there is a stark disagreement for the cold fusion cases.



Figure 2. Comparison of the measurements of  $P_{CN}$  in this work with that of [27].



Figure 3. Comparison of our measured values of  $P_{CN}$  with the predictions of [28]

# 4. Conclusions

What have we learned from this study? We have extended the systematics of  $P_{CN}$  to cases involving the synthesis of elements 111-118. We have parameterized the

new values of P<sub>CN</sub> with a simple linear fit that might be useful in predictions of cross sections for the synthesis of elements 119 and 120. We have compared our measurements with previous measurements and theoretical predictions.

#### 5. Acknowledgments

This work was supported in part by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under Grant No. DE-SC0014380.

### References

- 1. L. Yao and W. Loveland, Survival-mediated capture and fusion cross sections for heavy element synthesis, Phys. Rev. C 97, 014608 (2018).
- 2. W. Loveland and L. Yao, Survival mediated heavy element capture cross sections, EPJA Web of Conferences, 163, 00033 (2017)
- 3. Nuclear reactions video project, statistical model of the decay of excited nuclei, nrv.jinr.ru
- 4. A.V. Ignatyuk, IAEA Report No. INDC(CCP)-233/L, 1985, Unpublished.
- 5. P. Moller, , A.J. Sierk, T. Ichikawa, and H. Sagawa, Nuclear ground state masses and deformations-FRDN-2012, At. Data and Nucl. Data Tables 109-110, 1 (2016).
- 6. V.I. Zagrebaev, Y. Aritomo, M.G. Itkis, Y.T. Oganessian and M. Ohta, Synthesis of superheavy nuclei: How accurately can we describe it and calculate the cross sections?, Phys. Rev. C 65, 014607 (2007).
- 7. H.A. Kramers, Brownian motion in a field of force and the diffusion model of chemical reactions, Physica (Amsterdam) 7, 284 (1940).
- 8. Ch.E. Duellmann, (private communication).
- 9. W. Loveland, An experimentalist's view of the uncertainties in understanding heavy element synthesis, Eur. J. Phys. A 51, 120 (2015).
- 10. S. Hofmann, et al., New results on elements 111 and 112, Eur. Phys. J. A. 14, 147 (2002)
- 11. C.M. Folden et al., Development of an Odd-Z-Projectile Reaction for Heavy Element Synthesis: <sup>208</sup>Pb(<sup>64</sup>Ni,n)<sup>271</sup>Ds and <sup>208</sup>Pb(<sup>65</sup>Cu,n) <sup>272</sup>111.Phys. Rev. Lett. **93**, 212702 (2004).
- 12. Y.T. Oganessian, et al., Measurement of cross sections and decay properties of the isotopes of elements 112, 114, and 116 produced in the fusion reactions <sup>233,238</sup>U, <sup>242</sup>Pu and  $^{248}$ Cm +  $^{48}$ Ca, Phys. Rev. C **70**, 064609 (2004). 13. S. Hofmann, et al., The reaction  $^{48}$ Ca +  $^{238}$ U $\rightarrow$   $^{286}$ 112<sup>\*</sup> studied at the GSI-SHIP, Eur. Phys.
- J. A. 32, 251 (2007).
- 14. S. Hofmann, et al, New results on element 111 and 112, Eur. Phys. J. A. 14, 147 (2002).
- 15. Y.T. Oganessian et al, Synthesis of the isotope <sup>282</sup>113 in the <sup>237</sup>Np + <sup>48</sup>Ca fusion reaction, Phys. Rev. C 76, 011601 (2007).
- 16. K. Morita et al., New Result in the Production and Decay of an isotope, <sup>278</sup>113, of the 113<sup>th</sup> Element, J. Phys. Soc. Jpn 81, 103201 (2012).
- 17. V.K. Utyonkov, et al., Experiments on the synthesis of superheavy nuclei <sup>284</sup>Fl and <sup>285</sup>Fl in the <sup>239,240</sup>Pu +<sup>48</sup>Ca reactions, Phys. Rev. C **92**, 034609 (2015).
- 18. P.A. Ellison et al., New Superheavy Element Isotopes: <sup>242</sup>Pu(<sup>48</sup>Ca, 5n) <sup>285</sup>114, Phys. Rev. Lett 105, 182701 (2010)
- 19. Ch. E. Duellmann, et al. Production and Decay of Element 114: High Cross Sections and the New Nucleus <sup>277</sup>Hs, Phys. Rev. Lett. **104**, 252701 (2010).

- Y.T. Oganessian et al., Measurements of cross sections for the fusion-evaporation reactions: <sup>244</sup>Pu(<sup>48</sup>Ca,xn) <sup>292-x</sup>114 and <sup>245</sup>Cm(<sup>48</sup>Ca,xn)<sup>293-x</sup>116, Phys. Rev. C 69, 054607 (2004).
   Y.T. Oganessian et al, Investigation of the <sup>243</sup>Am + <sup>48</sup>Ca reaction products previously ob-
- served in the experiments on elements 113, 115, and 117, Phys. Rev. C 87, 014302 (2013).
  22. Y.T. Oganessian et al., New Insights into the <sup>243</sup>Am + <sup>48</sup>Ca Reaction Products Previously
- Observed in the Experiments on Elements 113,115, and 117, Phys. Rev. Lett, 108, 022502 (2012).
- 23. Y.T. Oganessian et al., Experiments on the synthesis of element 115 in the reaction <sup>243</sup>Am(<sup>48</sup>Ca,xn)<sup>291-x</sup>115, Phys. Rev. C **69**, 021601 (2004).
- 24. Y.T. Oganessian et al., Synthesis of the isotopes of elements 118 and 116 in the <sup>249</sup>Cf and <sup>245</sup>Cm + <sup>48</sup>Ca fusion reactions, Phys. Rev. C74, 044602 (2006).
  25. S. Hofmann et al., The reaction <sup>48</sup>Ca + <sup>248</sup>Cm→<sup>296</sup>116\* studied at the GSI-SHIP, Eur. Phys.
- J. A 48, 62 (2012).
- 26. Y.T. Oganessian et al., Experimental studies of the <sup>249</sup>Bk + <sup>48</sup>Ca reaction including decay properties and excitation function for isotopes of element 117 and discovery of the new isotope <sup>277</sup>Mt., Phys. Rev. C 87, 054621 (2013).
- 27. E.M. Kozulin et al., Fission and quasifission of composite systems with Z=108-120: Transition from heavy-ion reactions involving S and Ca to Ti and Ni ions, Phys. Rev. C 94, 054613 (2016).
- 28. A.K. Nasirov, G. Mandaglio, G. Giardina, A. Sobiczewski, and A.I. Muminov, Effects of the entrance channel and fission barrier in the synthesis of superheavy element Z=120, Phys. Rev. C 84, 044612 (2011).
- 29. V.I. Zagrebaev, Synthesis of superheavy nuclei: Nucleon collectivization as a mechanism for compound nucleus formation, Phys. Rev. C 64, 034606 (2001).