

INJECTIVE ENVELOPES OF TRANSITION SYSTEMS AND FERRERS LANGUAGES †

MUSTAPHA KABIL AND MAURICE POUZET

† Dedicated to the memory of Maurice Nivat

ABSTRACT. We consider reflexive and involutive transition systems over an ordered alphabet A equipped with an involution. We give a description of the injective envelope of any two-element set in terms of Galois lattice, from which we derive a test of its finiteness. Our description leads to the notion of Ferrers language.

1. INTRODUCTION AND PRESENTATION OF THE MAIN RESULTS

This paper is about involutive and reflexive transition systems from a metric point of view. This point of view, inspired from the work of Quilliot (1983), and applied first to posets and graphs, was initiated by the second author [29] and developed through the theses of Jawhari (1983), Misane (1984) and several papers [17] (1986), [30] (1994), [33] (1992), [23] (1998), [24] (2018). It consists to view arbitrary transition systems as metric spaces. The distance between two states is a language instead of a non-negative real. To a transition system $M := (Q, T)$, with set of states Q and set of transitions T over an alphabet A we associate a map d_M from $Q \times Q$ into the set $\mathcal{P}(A^*)$ of languages over A . The value $d_M(x, y)$ is the language accepted by the automaton $\mathcal{A} := (M, \{x\}, \{y\})$ having x as an initial state and y as a final state. The set $\mathcal{P}(A^*)$ is an ordered monoid, the monoid operation being the concatenation of languages (with neutral element $\{\square\}$, the language reduced to the empty word \square) and the order the reverse of inclusion. The map d_M has similar properties of an ordinary distance (e.g. it satisfies the triangular inequality). Hence, we may use concepts and techniques of the theory of metric spaces in the study of transition systems as well as classes of transition systems. Concepts of *balls*, *hyperconvex metric space* and *non-expansive* maps between metric spaces extend to transition systems and more generally to metric spaces over $\mathcal{P}(A^*)$. Due to the fact that joins exist in the set of values, the category of metric spaces with the non-expansive maps as morphisms has products. Then, one may also define *retractions* and *coretractions*, and by considering *isometries* as approximations of coretractions, *injective* metric spaces and *absolute retracts*. In ordinary metric spaces, the distance is symmetric. To be closer to this situation, it is convenient to suppose that the value of $d_M(x, y)$ determines the value of $d_M(y, x)$; for that, we suppose that the alphabet is equipped with an involution – and our transition systems M are *involutive*, in the sense that $(x, \alpha, y) \in T$ if and only if $(y, \alpha, x) \in T$. Once the involution is extended to A^* and then to $\mathcal{P}(A^*)$, we have $d_M(x, y) = d_M(y, x)$. Then, one can extend the definition of metric spaces to transition systems in a natural way, see [30].

Date: July 5, 2019.

1991 Mathematics Subject Classification. Primary 06A15, 06D20, 46B85, 68Q70; Secondary 68R15 .

Key words and phrases. Metric spaces, Injective envelopes, Transition systems, Ferrers languages, Ordered sets, Interval orders, Well-quasi-order.

This work is a continuation of the work published in [22, 23, 24]. We require that transition systems M are *reflexive*, that is every letter occurs to every vertex: $(x, \alpha, x) \in T$ for every $x \in Q$ and $\alpha \in A$. In this case, distances values are final segments of A^* equipped with the subword ordering, that is subsets F of A^* such that $u \in F$ and $u \leq v$ for the subword ordering imply $v \in F$. It turns out that several properties of involutive and reflexive systems and more generally metric spaces over the set $\mathbf{F}(A^*)$ of final segments of A^* rely almost uniquely on the structure of $\mathbf{F}(A^*)$. According to the terminology of Kaarli and Radeleczki [19], this structure is the dual of an *integral involutive quantale* (I^2Q for short); here we stick to the name of *Heyting algebra* that we used in a series of papers. This is a complete lattice \mathcal{H} with a monoid operation (not necessarily commutative) and an involution $-$ which is isotone and reverses the operation. In order to be closer to the operation of concatenation of languages we denote by \cdot the monoid operation. We suppose that the neutral element of the monoid, that we denote 1, is the least element of the ordering and we suppose that the distributivity law below holds

$$(1) \quad \bigwedge \{p_\alpha \cdot q : \alpha \in I\} = \bigwedge \{p_\alpha : \alpha \in I\} \cdot q.$$

As shown in [17], the notions of injective, absolute retract and hyperconvex spaces over a Heyting algebra coincide, this being essentially due to the fact that the set of the values of the distance, being an Heyting algebra, can be equipped with a distance and that every metric space can be embedded into a power of that metric space. Furthermore, every space has an injective envelope.

In particular, every metric space over the Heyting algebra $\mathbf{F}(A^*)$ has an injective envelope. The study of such injective envelope was initiated in [23]. It is based on the properties of the injective envelope of two-element metric spaces. A large account of its properties was given in [22] and [24]. In this paper, we look at the many facets of this object which have not been published yet.

If F is a final segment of A^* , the injective envelope \mathcal{S}_F of the two-element space $\{x, y\}$ such that $d(x, y) = F$ is associated to an involutive and reflexive transition system. Let M_F be this transition system and \mathcal{A}_F be the automaton $(M_F, \{x\}, \{y\})$.

We characterize first this injective envelope in terms of reflexive and involutive transition systems.

Theorem 1. *Let $\mathcal{A} := (M, \{x\}, \{y\})$ be a reflexive and involutive automaton accepting a final segment F of A^* . Then \mathcal{A} is isomorphic to \mathcal{A}_F iff for every reflexive and involutive automaton $\mathcal{A}' := (M', \{x'\}, \{y'\})$ which accepts F , the following properties hold:*

- (i) *Every automata morphism $f : \mathcal{A} \rightarrow \mathcal{A}'$, if any, is an isometric embedding;*
- (ii) *The map $g : \{x', y'\} \rightarrow M$ such that $g(x') = x$ and $g(y') = y$ extends to a morphism of automata from \mathcal{A}' to \mathcal{A} .*

We will obtain this result as a consequence of a characterization of the injective envelope among generalized metric spaces (Theorem 4) given in Section 2.

Then, we develop an approach in terms of Galois correspondence.

To a final segment F of A^* we associate the incidence structure $R := (A^*, \rho_F, A^*)$ where ρ is the binary relation on A^* defined by $u\rho_F v$ if the concatenation uv of u and v belongs to F . For each $v \in A^*$, let $R^{-1}(v) := \{u \in A^* : u\rho_F v\}$ and for $V \subseteq A^*$, let $R_\wedge^{-1}(V) := \bigcap_{v \in V} \rho_F^{-1}(v)$.

The collection of all $R_\wedge^{-1}(V)$ for $V \subseteq A^*$, once ordered by inclusion, forms a complete lattice $\text{Gal}(R)$, called the *Galois lattice* associated to the incidence structure R . As a subset of $\mathcal{H} := \mathbf{F}(A^*)$, it inherits the metric structure $d_{\mathcal{H}}$ of \mathcal{H} . Containing $x := A^*$ and $y := F$, two elements which verify $d_{\mathcal{H}}(x, y) = F$, this metric space is the injective envelope of $\{x, y\}$.

For concrete examples, suppose that F is a finite union of final segments $F_0, \dots, F_i, \dots, F_{k-1}$ and that each F_i is generated by X_i , a set of words u_i of the same length n_i , all of the form $u_i := a_0^i a_1^i \dots a_{n_i-1}^i$ with $a_j^i \in X_{i,j} \subseteq A$. Let $\mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1}$ be the direct product of k chains $\mathbf{n}_0, \dots, \mathbf{n}_{k-1}$ where $\mathbf{n}_i := \{0, \dots, n_i - 1\}$ is equipped with the natural ordering, and let $\mathbf{F}(\mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1})$ be the collection of final segments of $\mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1}$ ordered by inclusion.

We prove (see Section 5):

Theorem 2. *As a lattice, the injective envelope \mathcal{S}_F can be identified with an intersection closed subset of the set $\mathbf{F}(\mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1})$. Moreover if $\downarrow u_i \cap \downarrow u_j = \{\square\}$ whenever $u_i \in X_i$, $u_j \in X_j$, and $i \neq j$, then \mathcal{S}_F identifies to the full set $\mathbf{F}(\mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1})$.*

We recall that a poset P is *well-quasi-ordered*, in brief w.q.o., if well-founded (every non empty subset contains a minimal element) and contains no infinite antichain. A fundamental result of G.Higman [14] asserts that the free ordered monoid A^* is w.q.o. whenever the alphabet A is w.q.o.. The set of final segments of a w.q.o. set, once ordered by reverse of the inclusion, is well-founded [14]. Hence, if our alphabet A is w.q.o., every final segment F of A^* is generated by finitely many words u_0, \dots, u_{k-1} , hence has the form mentioned above. Consequently, the corresponding injective envelope is finite. Concerning its size, let us mention that if $k = 2$, then $\mathbf{F}(\mathbf{n}_0 \otimes \mathbf{n}_1)$ has size $\binom{n_0 \cdot n_1}{n_0}$ [1] (3.16, p. 80). If $n_0 = \dots = n_{k-1} = 2$, then $(\mathbf{F}(\mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1}), \supseteq)$ is isomorphic to $FD(k)$, the free distributive lattice with k generators. It is a famous problem, raised by Dedekind, to give an explicit and workable formula for $FD(k)$. The largest exact value known is $FD(8)$ [38]. An asymptotic formula was given by Korshunov in 1981[25].

For an example, on the two-letter alphabet $A = \{a, b\}$, the words $u_0 := aa$ and $u_1 := bb$ give the lattice (ordered by reverse of inclusion) and graph represented on Figure 1.




Figure 1

Structural properties of transition systems rely upon algebraic properties of languages and conversely. In fact, transition systems can be viewed as geometric objects interpreting these algebraic properties. An illustration of this claim is given by the following result (see Corollary 4.9 [23]).

Theorem 3. *Let F be a nonempty final segment of A^* . If F is the concatenation of final segments F_1, \dots, F_n then the automaton $\mathcal{A}_F := (M_F, \{x\}, \{y\})$ associated to the injective*

envelope \mathcal{S}_F is the concatenation $\mathcal{A}_{F_1} \cdot \dots \cdot \mathcal{A}_{F_n}$ of automata $\mathcal{A}_{F_i} := (M_{F_i}, \{x_i\}, \{y_i\})$ associated to the injective envelope S_{F_i} , this concatenation being obtained by identifying each y_i with x_{i+1} .

The fact that an automaton decomposes into such a concatenation can be viewed directly by looking at states which disconnect the underlying graph. From this follows the uniqueness of such a decomposition. This uniqueness amounts to the fact that the monoid $\mathbf{F}(A^*) \setminus \{\emptyset\}$ is free. A purely algebraic proof of this result is given in [24].

We discuss then the relationship between the minimal deterministic automaton accepting a final segment F , say Min_F , and the automaton \mathcal{A}_F associated to the injective envelope \mathcal{S}_F . This minimal automaton is part of \mathcal{A}_F , but not in an isometric way (Min_F being deterministic cannot be reflexive, in general it is not involutive). Among involutive and reflexive transition systems accepting a given final segment F , we consider those with a minimum number of states and among those, the ones with a maximal number of transitions, that we call Minmax automata. Exemples given by Mike Main and communicated by Maurice Nivat [27] show that contrarily to the case of deterministic automata, these automata are not unique.

We introduce Ferrers languages. A language L over A^* is *Ferrers* if

$$(2) \quad xx' \in L \text{ and } yy' \in L \text{ imply } xy' \in L \text{ or } yx' \in L \text{ for all } x, x', y, y' \in A^*.$$

The class of Ferrers languages is closed under complement but not under concatenation. Still, if F_1, \dots, F_n are Ferrers and each F_i is a final segment of A^* then the concatenation $F_1 \dots F_n$ is Ferrers (Corollary 19).

We prove that a final segment F of A^* is Ferrers if and only if the injective envelope \mathcal{S}_F is totally orderable, that is there is a linear order \leq on \mathcal{S}_F such that $d(x, z) \leq d(x, y)$ and $d(z, y) \leq d(x, y)$ for all $x \leq z \leq y$. (Theorem 23).

Over a finite alphabet A^* , Boolean combinations of final segments of A^* are called *piecewise testable languages*. They have been characterized by Simon [35] by the fact that their syntactical monoid is \mathcal{J} -trivial. The Boolean algebra of piecewise testable languages is included into the Boolean algebra generated by rational Ferrers languages. Indeed, over a finite alphabet, every final segment is a finite union of rational Ferrers languages. But, on an alphabet with at least two letters, there are rational Ferrers languages which are not piecewise testable (e.g., $L := A^*b$ on $A := \{a, b\}$). We do not know if they are dot-depth one.

This paper is organized as follows. Properties of metric spaces over a Heyting algebra and their injective envelopes are summarized in section 2. In section 3, we introduce the Heyting algebra $\mathbf{F}(A^*)$. In section 4 we consider transition systems as metric spaces. In section 5 we describe the injective envelope of a two-element metric spaces over $\mathbf{F}(A^*)$; we prove Theorem 1 and 2 and conclude the section by a counterexample about Minmax automata due to M.Main. Ferrers languages are introduced in section 6.

The results developed here have been presented at the International Conference on Discrete Mathematics and Computer Science (DIMACOS'11) organized by A. Boussaïri, M. Kabil, and A. Taïk in Mohammedia (Morocco) May, 5-8, 2011. They were never published; a part of it was included into the Thèse d'État defended by the first author [22].

2. METRIC SPACES OVER A HEYTING ALGEBRA

2.1. Basic facts. The following is extracted from [24] (for more details, see [23]). Let \mathcal{H} be a Heyting algebra and let E be a set. A \mathcal{H} -distance on E is a map $d : E^2 \rightarrow \mathcal{H}$ satisfying the following properties for all $x, y, z \in E$:

$$(1) \quad d(x, y) = 1 \iff x = y,$$

- (2) $d(x, y) \leq \overline{d(x, z)} \cdot d(z, y)$,
 (3) $d(x, y) = \overline{d(y, x)}$.

The pair (E, d) is called a \mathcal{H} -metric space. If there is no danger of confusion we will denote it E . A \mathcal{H} -distance can be defined on \mathcal{H} . This fact relies on the classical notion of *residuation*. Let $v \in \mathcal{H}$. Given $\beta \in \mathcal{H}$, each of the sets $\{r \in \mathcal{H} : v \leq r \cdot \beta\}$ and $\{r \in \mathcal{H} : v \leq \beta \cdot r\}$ has a least element, that we denote respectively $[v \cdot \beta^{-1}]$ and $[\beta^{-1} \cdot v]$ (note that $[\beta^{-1} \cdot v] = [\bar{v} \cdot (\bar{\beta})^{-1}]$). It follows that for all $p, q \in \mathcal{H}$, the set

$$D(p, q) := \{r \in \mathcal{H} : p \leq q \cdot \bar{r} \text{ and } q \leq p \cdot r\}$$

has a least element, namely $[\bar{p} \cdot (\bar{q})^{-1}] \vee [p^{-1} \cdot q]$, that we denote $d_{\mathcal{H}}(p, q)$. As shown in [17], the map $(p, q) \rightarrow d_{\mathcal{H}}(p, q)$ is a \mathcal{H} -distance.

Let (E, d) and (E', d') be two \mathcal{H} -metric spaces. Recall that a map $f : E \rightarrow E'$ is a *contraction* (or a non-expansive map) from (E, d) to (E', d') provided that $d'(f(x), f(y)) \leq d(x, y)$ holds for all $x, y \in E$. The map f is an *isometry* if $d'(f(x), f(y)) = d(x, y)$ for all $x, y \in E$. We say that E and E' are *isomorphic*, a fact that we denote $E \cong E'$, if there is a surjective isometry from E onto E' .

Let $((E_i, d_i))_{i \in I}$ be a family of \mathcal{H} -metric spaces. The *direct product* $\prod_{i \in I} (E_i, d_i)$, is the metric space (E, d) where E is the cartesian product $\prod_{i \in I} E_i$ and d is the "sup" (or ℓ^∞) distance defined by $d((x_i)_{i \in I}, (y_i)_{i \in I}) = \bigvee_{i \in I} d_i(x_i, y_i)$.

For a \mathcal{H} -metric space E , $x \in E$ and $r \in \mathcal{H}$, we define the *ball* $B_E(x, r)$ as the set $\{y \in E : d(x, y) \leq r\}$. We say that E is *convex* if the intersection of two balls $B_E(x_1, r_1)$ and $B_E(x_2, r_2)$ is non-empty provided that $d(x_1, x_2) \leq r_1 \cdot \bar{r}_2$. We say that E is *hyperconvex* if the intersection of every family of balls $(B_E(x_i, r_i))_{i \in I}$ is non-empty whenever $d(x_i, x_j) \leq r_i \cdot \bar{r}_j$ for all $i, j \in I$. For an example, $(\mathcal{H}, d_{\mathcal{H}})$ is a *hyperconvex \mathcal{H} -metric space and every \mathcal{H} -metric space embeds isometrically into a power of $(\mathcal{H}, d_{\mathcal{H}})$* [17]. This is due to the fact that for every \mathcal{H} -metric space (E, d) and for all $x, y \in E$ the following equality holds:

$$d(x, y) = \bigvee_{z \in E} d_{\mathcal{H}}(d(z, x), d(z, y)).$$

The space E is a *retract* of E' , in symbols $E \triangleleft E'$, if there are two contractions $f : E \rightarrow E'$ and $g : E' \rightarrow E$ such that $g \circ f = id_E$ (where id_E is the identity map on E). In this case, f is a *coretraction* and g a *retraction*. If E is a subspace of E' , then clearly E is a retract of E' if there is a contraction from E' to E such $g(x) = x$ for all $x \in E$. We can easily see that every coretraction is an isometry. A metric space is an *absolute retract* if it is a retract of every isometric extension. The space E is said to be *injective* if for all \mathcal{H} -metric space E' and E'' , each contraction $f : E' \rightarrow E$ and every isometry $g : E' \rightarrow E''$ there is a contraction $h : E'' \rightarrow E$ such that $h \circ g = f$. We recall that *for a metric space over a Heyting algebra \mathcal{H} , the notions of absolute retract, injective, hyperconvex and retract of a power of $(\mathcal{H}, d_{\mathcal{H}})$ coincide* [17].

2.2. Injective envelope. A contraction $f : E \rightarrow E'$ is *essential* if for every contraction $g : E' \rightarrow E''$, the map $g \circ f$ is an isometry if and only if g is isometry (note that, in particular, f is an isometry). An essential contraction f from E into an injective \mathcal{H} -metric space E' is called an *injective envelope* of E . We will rather say that E' is an *injective envelope* of E . We can view an injective envelope of a metric space E as a minimal injective \mathcal{H} -metric space containing (isometrically) E . Two injective envelopes of E are isomorphic via an isomorphism which is the identity over E . This allows to talk about "the" injective envelope of E ; we will denote it by $\mathcal{N}(E)$. A particular injective envelope of E will be

called a *representation* of $\mathcal{N}(E)$. The construction of injective envelope is based upon the notion of *minimal metric form*. A *weak metric form* is every map $f : E \rightarrow \mathcal{H}$ satisfying $d(x, y) \leq f(x) \cdot \overline{f(y)}$ for all $x, y \in E$. This is a *metric form* if in addition $f(x) \leq d(x, y) \cdot f(y)$ for all $x, y \in E$. Equivalently, f is a metric form if and only if $d_{\mathcal{H}}(d(x, y), f(x)) \leq f(y)$ for all $x, y \in E$. A (weak) metric form is *minimal* if there is no other (weak) metric form g satisfying $g \leq f$ (that is $g(x) \leq f(x)$ for all $x \in E$). Since every weak metric form majorizes a metric form, the two notions of minimality coincide. As shown in [17] *every \mathcal{H} -metric space has an injective envelope; the space of minimal metric forms is a representation of it, (cf. Theorem 2.2 of [23])*.

We give below a new characterization of the injective envelope.

Theorem 4. *Let (E, d) be a metric space over \mathcal{H} and $X \subseteq E$. Then (E, d) is isomorphic to the injective envelope of $(X, d_{\uparrow X})$ iff for every metric space (E', d') , $X' \subseteq E'$, every isometry $f : (X, d_{\uparrow X})$ onto $(X', d'_{\uparrow X'})$:*

- (i) *Every non-expansive map $\bar{f} : (E, d) \rightarrow (E', d')$, if any, which extends f is an isometric embedding of (E, d) into (E', d') ;*
- (ii) *The isometry f^{-1} extends to a non-expansive map from (E', d') into (E, d) .*

Proof. Suppose that (E, d) is isomorphic to the injective envelope of $(X, d_{\uparrow X})$. Since (E, d) is injective, the isometry f^{-1} extends to a non-expansive map from (E', d') into (E, d) . This proves that (ii) holds. The proof that (i) holds relies on the properties of metric forms. To prove that f is an isometry embedding amounts to prove the equality:

$$(3) \quad d'(f(x), f(y)) = d(x, y)$$

for all $x, y \in E$. Set $X'' := f(X)$. For every $x \in E$, let $h_x : X \rightarrow \mathcal{H}$ and $g_x : X \rightarrow \mathcal{H}$ be the maps defined by setting $h_x(z) = d(z, x)$ and $g_x(z) = d'(f(z), f(x))$ for $z \in X$. Since E is the injective envelope of X , h_x is a minimal metric form; since f is non-expansive and induces an isometry from X onto X'' , the map g_x is a metric form below h_x , hence $h_x = g_x$. It follows that

$$(4) \quad d(z, x) = d(f(z), f(x))$$

for every $z \in X$.

Let $x, y \in E$. By construction of the injective envelope, its elements identify to minimal metric forms over X , hence x and y identify respectively to h_x and h_y and

$$(5) \quad d(x, y) = d_{\mathcal{H}}(h_x, h_y) := \text{Sup}\{d_{\mathcal{H}}(d(z, x), d(z, y)) : z \in X\}.$$

Set $\alpha := d'(f(x), f(y))$. Let $z \in E$. By definition, $d_{\mathcal{H}}(d'(f(z), f(x)), d'(f(z), f(y))) = \text{Min} D(d'(f(z), f(x)), d'(f(z), f(y)))$ where $D(d'(f(z), f(x)), d'(f(z), f(y))) = \{r \in \mathcal{H} : d'(f(z), f(x)) \leq d'(f(z), f(y)) \cdot \bar{r} \text{ and } d'(f(z), f(y)) \leq d'(f(z), f(x)) \cdot r\}$. From the triangular inequality, we have $d'(f(z), f(y)) \leq d'(f(z), f(x)) \cdot \alpha$ and $d'(f(z), f(x)) \leq d'(f(z), f(y)) \cdot \bar{\alpha}$. Hence, $\alpha \in D(d'(f(z), f(x)), d'(f(z), f(y)))$. The inequality

$$d_{\mathcal{H}}(d'(f(z), f(x)), d'(f(z), f(y))) \leq \alpha$$

follows. Consequently,

$$\text{Sup}\{d_{\mathcal{H}}(d'(f(z), f(x)), d'(f(z), f(y))) : z \in E\} \leq \alpha.$$

With Equalities (4) and (5) we get: $d(x, y) = \text{Sup}\{d_{\mathcal{H}}(d(z, x), d(z, y)) : z \in E\} \leq \alpha$. Since f is non-expansive, we have $\alpha \leq d(x, y)$. Thus $d(x, y) = \alpha$ that is Equality (3) holds.

Conversely, suppose that (i) and (ii) hold. Let (E', d') be the injective envelope of $(X, d_{\uparrow X})$ and f be the identity map from $(X, d_{\uparrow X})$ onto itself. Applying (ii), the map f^{-1} extends to a non-expansive map g from (E', d') into (E, d) . Since (E', d') is injective, the

map f extends to a non-expansive map \bar{f} from (E, d) into (E', d') . The map $f \circ g$ is non-expansive and is the identity on X . Since (E', d') is the injective envelope of $(X, d_{\uparrow X})$, $f \circ g$ is the identity on E' (note that elements of E' identify to minimal metric forms over X), hence g is injective and f is surjective. Now by (i), f is an isometry on its image. Hence f is an isometry of (E, d) onto (E', d') . Thus (E, d) is the injective envelope of $(X, d_{\uparrow X})$ as claimed. \square

Up to Theorem 6, we include the few facts we need about injective envelopes of two-element metric spaces (see [23] for proofs).

Let \mathcal{H} be a Heyting algebra and $v \in \mathcal{H}$. Let $E := \{x, y\}$ be a two-element \mathcal{H} -metric space such that $d(x, y) = v$. We denote by \mathcal{N}_v the injective envelope of E . We give three representations of it. For the first one, we consider the set of minimal metric forms over E . That is, in this case, the set of minimal pairs $h := (h_x, h_y) \in \mathcal{H}^2$ such that $h_x \cdot \bar{h}_y \geq v$, the set \mathcal{H}^2 being equipped with the product ordering. Each element $z \in \mathcal{N}_v$ identifies to the pair $(d_{\mathcal{N}_v}(x, z), d_{\mathcal{N}_v}(y, z))$; in particular, x and y identify to $(1, \bar{v})$ and to $(v, 1)$ respectively. We equip \mathcal{H}^2 with the supremum distance:

$$d_{\mathcal{H}^2}((u_1, u_2), (u'_1, u'_2)) := d_{\mathcal{H}}(u_1, u'_1) \vee d_{\mathcal{H}}(u_2, u'_2).$$

With the induced distance, \mathcal{N}_v becomes a metric space. If (h_x, h_y) and (h'_x, h'_y) are two elements of \mathcal{N}_v , their distance is $d_{\mathcal{H}}(h_x, h'_x) \vee d_{\mathcal{H}}(h_y, h'_y)$. In fact,

$$(6) \quad d_{\mathcal{H}}(h_x, h'_x) = d_{\mathcal{H}}(h_y, h'_y).$$

The proof is easy: We prove that if $h_x \leq h'_x \cdot \bar{r}$ for some $r \in \mathcal{H}$, then $h'_y \leq h_y \cdot r$. This and the corresponding inequality with y replacing x will lead to (6). Suppose that $h_x \leq h'_x \cdot \bar{r}$. We have $v \leq h_x \cdot \bar{h}_y \leq h'_x \cdot \bar{r} \cdot \bar{h}_y$ and thus $r \leq h'_x \cdot \bar{r} \cdot \bar{h}_y$. Since $v \leq h'_x \cdot \bar{h}'_y$, we have $v \leq h'_x \cdot (\bar{h}'_y \wedge (\bar{r} \cdot \bar{h}_y))$ by distributivity. Since (h'_x, h'_y) is a minimal metric form above v , we have $\bar{h}'_y = \bar{h}'_y \wedge (\bar{r} \cdot \bar{h}_y)$, that is $h'_y \leq h_y \cdot r$.

Due to the fact that in a minimal metric form (h_x, h_y) each component determines the other, we may prefer an other presentation of \mathcal{N}_v as a subset of \mathcal{H} . Set $\mathcal{S}_v := \{[v \cdot \beta^{-1}] : \beta \in \mathcal{H}\}$; equipped with the ordering induced by the ordering over \mathcal{H} this is a complete lattice. A pair (h_x, h_y) belongs to \mathcal{N}_v if and only if $h_x = [v \cdot \bar{h}_y^{-1}]$ and $\bar{h}_y = [h_x^{-1} \cdot v]$. This yields a correspondence between \mathcal{N}_v and \mathcal{S}_v .

Now, in several instances, e.g. in the case of the sum of two metric spaces (see subsection 2.3), it is preferable to consider the set \mathcal{C}_v of all pairs $(u_1, u_2) \in \mathcal{H}^2$ such that $v \leq u_1 \cdot u_2$. Once equipped with the ordering induced by the product ordering on \mathcal{H}^2 , we can consider the set \mathcal{N}'_v of its minimal elements. Each minimal element (u_1, u_2) yields the minimal metric form (u_1, \bar{u}_2) (and conversely). The distance over \mathcal{H}^2 is different from the previous case. We have to equip \mathcal{H}^2 with the product of the distance $d_{\mathcal{H}}$ with the distance $d'_{\mathcal{H}}$ defined on \mathcal{H} by $d'_{\mathcal{H}}(u_2, u'_2) := d_{\mathcal{H}}(\bar{u}_2, \bar{u}'_2)$.

Lemma 1. (Lemma 2.3, Proposition 2.7 of [23]) *The space \mathcal{N}_v equipped with the supremum distance and the set \mathcal{S}_v equipped with the distance induced by the distance over \mathcal{H} are injective envelopes of the two-element metric spaces $\{(1, v), (v, 1)\}$ and $\{1, v\}$ respectively. These spaces are isometric to the injective envelope of $E := \{x, y\}$.*

Theorem 5. (Theorem 2.9 [23]) *Let E be a \mathcal{H} -metric space. If E is the injective envelope of a two-element set then E contains no proper isometric subspace.*

Proposition 1. (Corollary 3.3 [23]) *The following properties are equivalent:*

- (i) *The injective envelope of any finite metric space is finite;*
- (ii) *The injective envelope of any two-element metric space is finite.*

A metric space is *linearly orderable* if there is a linear ordering \leq on E such that $d(x, z) \leq d(x, y)$ and $d(z, y) \leq d(x, y)$ for all $x, y, z \in E$ with $x \leq z \leq y$.

Proposition 2. (Fact 5 [23]) *Let $u \in \mathcal{H}$. The space \mathcal{S}_u is linearly orderable if and only if the ordering is induced by the order on \mathcal{H} or by its reverse.*

We say that a metric space is *finitely indecomposable* if for every finite family $(E_i)_{i \in I}$ of metric spaces, $E \triangleleft \prod_{i \in I} E_i$ implies $E \triangleleft E_i$ for some $i \in I$. This notion was used by E. Corominas for posets [7].

Theorem 6. (Theorem 3.8 [23]) *Let E be a finite absolute retract. The following properties are equivalent:*

- (i) *E is finitely indecomposable;*
- (ii) *E is the injective envelope of a two-element metric space $\{x, y\}$ such that the distance $d(x, y)$ is join-irreducible in \mathcal{H} .*

2.3. Sum of metric spaces. Let (E_1, d_1) and (E_2, d_2) be two disjoint \mathcal{H} -metric spaces and let $x_1 \in E_1, x_2 \in E_2$. If we endow the set $\{x_1, x_2\}$ with a \mathcal{H} -distance d' , then we can define a \mathcal{H} -distance d on $E := E_1 \cup E_2$ as follows:

- If $x, y \in E_i$ with $i \in \{1, 2\}$ then $d(x, y) = d_i(x, y)$;
- If $x \in E_i, y \in E_j$ with $i, j \in \{1, 2\}$ and $i \neq j$, then $d(x, y) = d_i(x, x_i) \cdot d'(x_i, x_j) \cdot d_j(x_j, y)$.

In particular, we can identify x_1 and x_2 which amounts to set $d'(x_1, x_2) = 0$ in the above formula.

If E_1 and E_2 are not disjoint, we replace it by two disjoint copies E'_1, E'_2 (eg $E'_i := E_i \times \{i\}$). Identifying the corresponding elements x'_1, x'_2 , we obtain a \mathcal{H} -metric space that we denote $(E_1, d_1) \cdot (E_2, d_2)$. Alternatively, we may suppose that E_1 and E_2 have only one element in common, say $z_{1,2}$, and we define the distance d on $E_1 \cup E_2$ by setting $d(x, y) := d_i(x, z_{1,2}) \cdot d_j(z_{1,2}, y)$ if $x \in E_i, y \in E_j, i \neq j$, and $d(x, y) := d_i(x, y)$ if $x, y \in E_i$.

We consider now objects consisting of a \mathcal{H} -metric space and two distinguished elements. Given two such objects, say $\underline{E}_1 := ((E_1, d_1), x_1, y_1)$ and $\underline{E}_2 := ((E_2, d_2), x_2, y_2)$, set $\underline{E}_1 \cdot \underline{E}_2 := ((E, d), x, y)$ where (E, d) is the space obtained by taking disjoint copies (E'_1, d'_1) and (E'_2, d'_2) of (E_1, d_1) and (E_2, d_2) , respectively, and by identifying the corresponding elements y'_1 and x'_2 and setting $x := x'_1$ and $y := y'_2$.

Definition 7. *Let \mathcal{H} be Heyting. A pair $(v_1, v_2) \in \mathcal{H}^2$ is summable if \mathcal{S}_{v_1, v_2} is isomorphic to $\mathcal{S}_{v_1} \cdot \mathcal{S}_{v_1}$.*

Definition 8. *Let \mathcal{H}' be an initial segment of \mathcal{H} which is also a submonoid. We say that \mathcal{H}' has the decomposition property if every pair $(v_1, v_2) \in \mathcal{H}' \times \mathcal{H}'$ is summable.*

Several examples of metric spaces over a Heyting algebra are given in [17]. We briefly examine some of these examples w.r.t. to their injective envelopes and the sum operation. Ordinary metric spaces enter in his frame. Add a largest element $+\infty$ to the set \mathbb{R}^+ of non-negative reals, extend the $+$ operation in the natural way, take the identity for the involution. Then $\mathbb{R}^+ \cup \{+\infty\}$ becomes a Heyting algebra and the metric spaces over it are just direct sums of ordinary metric spaces. The injective envelope of such a space is the direct sum of the injective envelope of its factors. The injective envelope of a two-element metric spaces $E := \{x, y\}$ with $r := d(x, y)$ is isometric to the segment $[0, r]$ si $r < +\infty$ and to E if $r = +\infty$. Trivially, \mathbb{R}^+ has the decomposition property. The sum of two convex (resp.

injective) metric spaces with a common vertex is convex (resp. injective). The reader will find in [9] a description of injective envelopes of finite ordinary metric spaces and interesting combinatorial properties as well (see also [10]).

Now, let $\mathcal{H} := \{a, b, 0, 1\}$ ordered by $0 < a, b < 1$ and a incomparable to b . The operation is the join, the involution exchange a and b . Metric spaces over \mathcal{H} correspond to ordered sets. As shown by Banaschewski and Bruns, every poset P has an injective envelope, namely its MacNeille completion [1]. Hence, if P has two elements, its injective envelope is P whenever these two elements are comparable, otherwise this is P augmented of a smallest and a largest element. Convexity property does not hold for \mathcal{H} , hence the sum of two injective with a common element does not need to be injective.

Next, suppose that \mathcal{H} is a complete meet-distributive lattice, the operation is the join and the involution is the identity. For example, if $\mathcal{H} := \mathbb{R} \cup \{+\infty\}$, metric spaces over \mathcal{H} are direct sums of ultrametric spaces. Metric spaces over Boolean algebras have been introduced by Blumenthal [3]. Let B be a Boolean algebra, let the operation be the supremum and the involution be the identity. Although B is not necessary complete, residuation allows to define a distance on B setting $d_B(p, q) := p\Delta q$ where Δ denotes the symmetric difference. From this follows that the interval $[0, u] := \{v \in B : 0 \leq v \leq u\}$ with the distance induced by d_B is the injective envelope of every pair $\{p, q\}$ of vertices of B such that $p \wedge q = 0$ and $p \vee q = u$. Hence if B is finite, the number of elements of the injective envelope of a 2-element metric space is a power of 2 hence the decomposition property does not hold. In Section 3, we give an example for which this decomposition property holds, namely the algebra $\mathbf{F}^\circ(A^*)$ (for more examples of generalisations of metric spaces, see [4], [5]).

3. THE HEYTING ALGEBRA $\mathbf{F}(A^*)$

Let A be a set. Considering A as an *alphabet* whose members are *letters*, we write a word α with a mere juxtaposition of its letters as $\alpha = a_0a_1\dots a_{n-1}$ where a_i are letters from A for $0 \leq i < n$. The integer n is the *length* of the word α and we denote it $|\alpha|$. Hence we identify letters with words of length 1. We denote by \square the empty word, which is the unique word of length zero. The *concatenation* of two word $\alpha := a_0\dots a_{n-1}$ and $\beta := b_0\dots b_{m-1}$ is the word $\alpha\beta := a_0\dots a_{n-1}b_0\dots b_{m-1}$. We denote by A^* the set of all words on the alphabet A . Once equipped with the concatenation of words, A^* is a monoid, whose neutral element is the empty word, in fact A^* is the *free monoid* on A . A *language* is any subset X of A^* . We denote by $\mathcal{P}(A^*)$ the set of languages. We will use capital letters for languages. If $X, Y \in \mathcal{P}(A^*)$ we may set $XY := \{\alpha\beta : \alpha \in X, \beta \in Y\}$ (and use Xy and xY instead of $X\{y\}$ and $\{x\}Y$). With this operation, which extends the concatenation operation on A^* , the set $\mathcal{P}(A^*)$ is a monoid (the set $\{\square\}$ is the neutral element). Ordered by inclusion, this is a (join) lattice ordered monoid. Indeed, concatenation distributes over arbitrary union, namely:

$$\left(\bigcup_{i \in I} X_i\right)Y = \bigcup_{i \in I} X_iY.$$

This monoid is residuated. Let $X, Y, F \in \mathcal{P}(A^*)$. As it is customary, we set $X^{-1}F := \{y \in A^* : Xy \subseteq F\}$ and $FY^{-1} := \{x \in A^* : xY \subseteq F\}$. We recall that

$$(7) \quad X(X^{-1}F) \subseteq F \text{ and } (FY^{-1})Y \subseteq F.$$

Set $\rho_F := \{(x, y) \in A^* \times A^*\}$ and $R := (A^*, \rho_F, A^*)$, then $R(x) = x^{-1}F$ and $R^{-1}(y) = FY^{-1}$. Thus, $\text{Gal}(R)$, the Galois lattice of R , is the set $\{FY^{-1} : Y \subseteq A^*\}$ ordered by inclusion. This is a complete lattice. The meet is the intersection, the largest element is A^* .

In the sequel, we study the metric structure of $\text{Gal}(R)$ when F is a final segment of the monoid A^* , this monoid being equipped with the Higman ordering.

We suppose from now that the alphabet A is ordered and equipped with an involution $\bar{}$ preserving the order. The involution extends to A^* : we set for every $\alpha := a_0 \cdots a_{n-1}$, $\bar{\alpha} := \bar{a}_{n-1} \cdots \bar{a}_0$. Note that $\overline{\alpha\beta} = \bar{\beta}\bar{\alpha}$ for all $\alpha, \beta \in A^*$. We order A^* with the Higman ordering: if α and β are two elements in A^* such $\alpha := a_0 \cdots a_{n-1}$ and $\beta := b_0 \cdots b_{m-1}$ then $\alpha \leq \beta$ if there is an injective and increasing map h from $\{0, \dots, n-1\}$ to $\{0, \dots, m-1\}$ such that for each i , $0 \leq i \leq n-1$, we have $a_i \leq b_{h(i)}$. Then A^* becomes an ordered monoid with respect to the concatenation of words. Let $\mathbf{F}(A^*)$ be the collection of final segments of A^* (that is $X \in \mathbf{F}(A^*)$ if $X \subseteq A^*$ and $\alpha \leq \beta, \alpha \in X$ implies $\beta \in X$). The set $\mathbf{F}(A^*)$ is stable w.r.t. the concatenation of languages: if $X, Y \in \mathbf{F}(A^*)$, then $XY \in \mathbf{F}(A^*)$. Clearly, the neutral element is A^* . The set $\mathbf{F}(A^*)$ ordered by inclusion is a complete lattice (the join is the union, the meet is the intersection). Concatenation distributes over union. Order $\mathbf{F}(A^*)$ by reverse of the inclusion, denote $X \leq Y$ instead of $X \supseteq Y$, extend the involution $\bar{}$ to $\mathbf{F}(A^*)$, set $\bar{X} = \{\bar{\alpha} : \alpha \in X\}$, denote by $X \cdot Y$ the concatenation XY and set $1 := A^*$ then:

Theorem 9. *The set $\underline{\mathbf{F}}(A^*) := (\mathbf{F}(A^*), \leq, \cdot, 1, \bar{})$ is a Heyting algebra.*

We may then define metric spaces over $\underline{\mathbf{F}}(A^*)$ and study injective objects and particularly injective envelopes.

According to Corollary 4.9 p.177 of [23] we have:

Theorem 10. *Let F, F_1, F_2 be final segment of A^* such that $F = F_1 \cdot F_2$. If $F \neq \emptyset$ then*

$$F = F_1 \cdot F_2 \iff \mathcal{S}_F \cong \mathcal{S}_{F_1} \cdot \mathcal{S}_{F_2}$$

According to Definition 8, this means that $\mathbf{F}^\circ(A^*) := \mathbf{F}(A^*) \setminus \{\emptyset\}$ has the decomposition property.

Among metric spaces over $\mathbf{F}(A^*)$ are those coming from reflexive and involutive transition systems. They are introduced in the next section.

4. TRANSITION SYSTEMS AS METRIC SPACES

Let A be a set. A *transition system* on the alphabet A is a pair $M := (Q, T)$ where $T \subseteq Q \times A \times Q$. The elements of Q are called *states* and those of T *transitions*. Let $M := (Q, T)$ and $M' := (Q', T')$ be two transition systems on the alphabet A . A map $f : Q \rightarrow Q'$ is a *morphism* of transition systems if for every transition $(p, a, q) \in T$, we have $(f(p), a, f(q)) \in T'$. When f is bijective and f^{-1} is a morphism from M' to M , we say that f is an *isomorphism*. The collection of transition systems over A , equipped with these morphisms, form a category. This category has products. If $(M_i)_{i \in I}$ is a family of transition systems, $M_i := (Q_i, T_i)$, then their product M is the transition system (Q, T) where Q is the direct product $\prod_{i \in I} Q_i$ and T is defined as follows: if $x := (x_i)_{i \in I}$ and $y := (y_i)_{i \in I}$ are two elements of Q and a is a letter, then $(x, a, y) \in T$ if and only if $(x_i, a, y_i) \in T_i$ for every $i \in I$.

An *automaton* \mathcal{A} on the alphabet A is given by a transition system $M := (Q, T)$ and two subsets I, F of Q called the set of *initial* and *final states*. We denote the automaton as a triple (M, I, F) . A *path* in the automaton $\mathcal{A} := (M, I, F)$ is a sequence $c := (e_i)_{i < n}$ of consecutive transitions, that is of transitions $e_i := (q_i, a_i, q_{i+1})$. The word $\alpha := a_0 \cdots a_{n-1}$ is the *label* of the path, the state q_0 is its *origin* and the state q_n its *end*. One agrees to define for each state q in Q a unique null path of length 0 with origin and end q . Its label is the empty word \square . A path is *successful* if its origin is in I and its end is in F . Finally, a word α on the alphabet A is *accepted* by the automaton \mathcal{A} if it is the label of some successful path. The *language accepted* by the automaton \mathcal{A} , denoted by $L_{\mathcal{A}}$, is the set of all words accepted by

\mathcal{A} . Let $\mathcal{A} := (M, I, F)$ and $\mathcal{A}' := (M', I', F')$ be two automata. A *morphism* from \mathcal{A} to \mathcal{A}' is a map $f : Q \rightarrow Q'$ satisfying the two conditions:

- (1) f is morphism from M to M' ;
- (2) $f(I) \subseteq I'$ and $f(F) \subseteq F'$.

If, moreover, f is bijective, $f(I) = I'$, $f(F) = F'$ and f^{-1} is also a morphism from \mathcal{A}' to \mathcal{A} , we say that f is an *isomorphism* and that the two automata \mathcal{A} and \mathcal{A}' are *isomorphic*.

To a metric space (E, d) over $\underline{F}(A^*)$, we may associate the transition system $M := (E, T)$ having E as set of states and $T := \{(x, a, y) : a \in d(x, y) \cap A\}$ as set of transitions. Notice that such a transition system has the following properties: for all $x, y \in E$ and every $a, b \in A$ with $b \geq a$:

- 1) $(x, a, x) \in T$;
- 2) $(x, a, y) \in T$ implies $(y, \bar{a}, x) \in T$;
- 3) $(x, a, y) \in T$ implies $(x, b, y) \in T$.

We say that a transition system satisfying these properties is *reflexive* and *involutive* (cf. [33, 23]). Clearly if $M := (Q, T)$ is such a transition system, the map $d_M : Q \times Q \rightarrow \mathbf{F}(A^*)$ where $d_M(x, y)$ is the language accepted by the automaton $(M, \{x\}, \{y\})$ is a distance. We have the following:

Lemma 2. *Let (E, d) be a metric space over $\underline{F}(A^*)$. The following properties are equivalent:*

- (i) *The map d is of the form d_M for some reflexive and involutive transition system $M := (E, T)$;*
- (ii) *For all $\alpha, \beta \in A^*$ and $x, y \in E$, if $\alpha \cdot \beta \in d(x, y)$, then there is some $z \in E$ such that $\alpha \in d(x, z)$ and $\beta \in d(z, y)$.*

The category of reflexive and involutive transition systems with the morphisms defined above identify to a subcategory of the category having as objects the metric spaces and morphisms the contractions. Indeed:

Fact 1. *Let $M_i := (Q_i, T_i)$ ($i = 1, 2$) be two reflexive and involutive transition systems. A map $f : Q_1 \rightarrow Q_2$ is a morphism from M_1 to M_2 if and only if f is a contraction from (Q_1, d_{M_1}) to (Q_2, d_{M_2}) .*

From this fact, we can observe that if $(M_i)_{i \in I}$ is a family of transition systems $M_i := (Q_i, T_i)$ then the metric space (Q, d) associated to the transition system $\prod_{i \in I} M_i$, product of the M_i 's, is the product of metric spaces (Q_i, d_i) associated to the transition systems (Q_i, T_i) .

Injective objects satisfy the convexity property stated in (ii) of Lemma 2. Hence, if F is a final segment of A^* , the distance d on the injective envelope \mathcal{S}_F coincide with the distance d_F associated with the transition system $M_F := (S_F, T_F)$ where

$$T_F := \{(p, a, q) : a \in d(p, q) \cap A\}.$$

We denote by \mathcal{A}_F the automaton $(M_F, \{x\}, \{y\})$, where $x := A^*$ and $y := F$.

From the existence of the injective envelope, we get:

Theorem 11. *For every $F \in \mathbf{F}(A^*)$ there is an involutive and reflexive transition system $M := (Q, T)$, an initial state x and a final state y such that the language accepted by the automaton $\mathcal{A} = (M, \{x\}, \{y\})$ is F . Moreover, if A is well-quasi-ordered then we may choose Q to be finite.*

Proof. Take $M := M_F$, $x := A^*$ and $y := F$. If A is well-quasi-ordered then A^* is also well-quasi-ordered (Higman[14]), hence the final segment F has a finite basis, that is, there are finitely many words $\alpha_0, \dots, \alpha_{n-1}$ such that $F = \{\alpha : \alpha_i \leq \alpha \text{ for some } i < n\}$. \square

Since injective objects in the category of metric spaces satisfy the convexity property stated in (ii) of Lemma 2, their distance is the distance associated with a transition system. Thus we may reproduce Theorem 4 almost verbatim. We get:

Theorem 12. *Let $M := (Q, T)$ be a transition system, $X \subseteq Q$. Then (M, d_M) is isomorphic to the injective envelope of $(X, d_{M \upharpoonright X})$ iff for every reflexive and involutive transition system $M' := (Q', T')$, $X' \subseteq Q'$, every isometry $f : (X, d_{M \upharpoonright X})$ onto $(X', d_{M' \upharpoonright X'})$:*

- (i) *Every non-expansive map $\bar{f} : (M, d_M) \rightarrow (M', d_{M'})$, if any, which extends f is an isometric embedding of (M, d_M) into $(M', d_{M'})$;*
- (ii) *The isometry f^{-1} extends to a non-expansive map from $(M', d_{M'})$ into (M, d_M) .*

Taking for X a 2-element subset $\{x, y\}$ of Q , Theorem 4 translates to Theorem 1 stated in the introduction.

4.1. Minimal automaton and Minmax automata. We suppose now that the alphabet A is finite. We refer to [34] Subsection 3.3 p. 111-118 for the construction of the minimal state deterministic automaton. If $F \in \mathbf{F}(A^*)$ and $u \in A^*$, the *left residual* is $u^{-1}F := \{v \in A^* : uv \in F\}$. The minimal automaton Min_F recognizing F has $Q_F := \{u^{-1}F : u \in A^*\}$ as set of states. Its initial state is $x := F$, its final state is $y := A^*$ and the transition function δ_F associate to the pair $(u^{-1}F, a) \in Q_F \times A$ the state $(ua)^{-1}F$.

One can check that:

Lemma 3. *The map $i : Min_F \rightarrow \mathcal{A}_F$ defined by $i(Y) := \bigcap_{y \in Y} Fy^{-1}$ is a morphism of automata.*

Proof. Suppose that $(u^{-1}F, a, (ua)^{-1}F)$ is a transition in the automaton Min_F . We prove that $(i(u^{-1}F), a, i((ua)^{-1}F))$ is a transition in the automaton \mathcal{A}_F . This is equivalent to $a \in d(i(u^{-1}F), i((ua)^{-1}F))$. The last condition amounts to 1) $i(u^{-1}F)a \subseteq i((ua)^{-1}F)$ and 2) $i((ua)^{-1}F\bar{a} \subseteq i(u^{-1}F)$. For 1), let $v \in i(u^{-1}F)$, we claim that $va \in i((ua)^{-1}F)$, that is for each word w , if $uaw \in F$, then $vaw \in F$. Since $v \in i(u^{-1}F)$, from $uaw \in F$, we have $vaw \in F$, as required. The inclusion 2) follows directly from the fact that F is a final segment. The equalities $i(F) = A^*$ and $i(A^*) = F$ are obvious. \square

In general, the transition system associated to Min_F is neither reflexive nor involutive. Among all involutive and reflexive transition systems $M := (Q, T)$ with initial state x and final state y such that the language accepted by the automaton $(M, \{x\}, \{y\})$ is F , we select those with the least number of states and among those transition systems, we select those with a maximum number of transitions; we call *minmax transition systems* these transition systems. The automaton corresponding to a minmax transition system is a *minmax automaton*. In other terms, an automaton $\mathcal{A} := (M, \{x\}, \{y\})$ is minmax if (a) it is reflexive and involutive, (b) for every reflexive and involutive automaton $\mathcal{A}' = (M', \{x'\}, \{y'\})$ such that $d_M(x, y) = d_{M'}(x', y')$, one has $|Q| \leq |Q'|$, moreover (c) $|Q| = |Q'|$ implies $|T| \geq |T'|$.

Proposition 3. *Let $M := (Q, T)$ be transition system and F be a final segment of A^* . If $\mathcal{A} := (M, \{x\}, \{y\})$ is a minmax automaton which accepts F , then \mathcal{A} is isomorphic to an induced automaton \mathcal{A}' of \mathcal{A}_F , that is there is some $Q' \subseteq \mathcal{S}_F$ containing A^* and F such that the automaton $\mathcal{A}' := (M_{F \upharpoonright Q'}, \{A^*\}, \{F\})$ is isomorphic to \mathcal{A} .*

Proof. To the automaton $\mathcal{A} := (M, \{x\}, \{y\})$, we associate the metric space (Q, d_M) induced by M . Identifying x with A^* and y with $F = d_M(x, y)$, the space \mathcal{S}_F identifies with the injective envelope of the 2-element space $\{x, y\}$. Let i be the identity mapping on $\{x, y\}$. This is a partial non-expansive mapping from (Q, d_M) into \mathcal{S}_F . Since \mathcal{S}_F is injective, this partial map extends to a non-expansive mapping f defined on Q . Since f is non-expansive, this map is an automata morphism. Let $Q' := f(Q)$ and consider the transition system $M' := (Q', \mathcal{S}_F \upharpoonright Q')$. We have $F = d(x, y) \leq d_{M'}(x, y) \leq d_M(x, y) = F$. Thus $d_{M'}(x, y) = F$. Since $|Q|$ is minimal, it follows that $|Q| = |Q'|$ proving that f is injective. Now since $|T|$ is maximum, we have $|T_{F \upharpoonright Q'}| = |T|$ proving that f is an automata isomorphism from \mathcal{A} to $\mathcal{A}' := (M', \{x\}, \{y\})$. \square

Let $M_1 := (Q_1, T_1)$ and $M_2 := (Q_2, T_2)$ be two transition systems such that Q_1 and Q_2 have only the element y in common. Let $\mathcal{A}_1 := (M_1, \{x\}, \{y\})$ and $\mathcal{A}_2 := (M_2, \{x\}, \{z\})$ be two automata. We denote $\mathcal{A}_1 \cdot \mathcal{A}_2$ the automaton $(M, \{x\}, \{y\})$ where $M = (Q, T)$ $Q = Q_1 \cup Q_2$ and $T = T_1 \cup T_2$.

Theorem 13. *Let F be a non-empty final segment of A^* . If $F = F_1F_2$, with F_1 and F_2 final segments of A^* , then an automaton \mathcal{A} with initial state x and final state y accepting F is minmax if and only if it decomposes into $\mathcal{A}_1 \cdot \mathcal{A}_2$ where \mathcal{A}_1 and \mathcal{A}_2 are two minmax automata accepting respectively F_1 between x and z and F_2 between z and y .*

Proof. From Theorem 10, we have $\mathcal{S}_{F_1} \cdot \mathcal{S}_{F_2} \cong \mathcal{S}_{F_1F_2}$. According to Proposition 3, a minmax automaton \mathcal{A} accepting F is a subautomaton of $\mathcal{A}_F = \mathcal{A}_{F_1} \cdot \mathcal{A}_{F_2}$. Thus \mathcal{A} decomposes into $\mathcal{A}_1 \cdot \mathcal{A}_2$ where \mathcal{A}_1 is a subautomaton of \mathcal{A}_{F_1} and \mathcal{A}_2 is a subautomaton of \mathcal{A}_{F_2} . Since \mathcal{A} is minmax, both \mathcal{A}_1 and \mathcal{A}_2 are minmax. Conversely, assume that \mathcal{A}_1 and \mathcal{A}_2 are minmax. Let \mathcal{A} be a minmax automaton accepting $F_1 \cdot F_2$. Then \mathcal{A} decomposes into $\mathcal{A}'_1 \cdot \mathcal{A}'_2$ where \mathcal{A}'_1 and \mathcal{A}'_2 are minmax. The automaton \mathcal{A}_1 and \mathcal{A}'_1 (resp. \mathcal{A}_2 and \mathcal{A}'_2) have the same number of states and transitions. That is $\mathcal{A}_1 \cdot \mathcal{A}_2$ is minmax. \square

Example 14. *(Mike Main, 1989, communicated by Maurice Nivat [27]). We give an example of two non-isomorphic minmax automata accepting the same language $L \in F(A^*)$.*

Let $A = \{a, b, c, a', b', c'\}$ with $\bar{a} = a'$, $\bar{b} = b'$ and $\bar{c} = c'$. Consider the automata represented on Figure 2. To each of these automata, we associate the involutive and reflexive automata obtained by replacing each transition (p, α, q) by (p, α, q) and $(q, \bar{\alpha}, p)$ and adding a loop at every vertex. The language accepted by each of these automata between x and y is $L = \uparrow \{ab, ac, ba, bc, ca, cb\}$. As it is easy to check, these automata are minmax but not isomorphic.




Figure 2

5. A DESCRIPTION OF THE INJECTIVE ENVELOPE

We describe the injective envelope \mathcal{S}_F in Galois lattice terms. We derive a test of its finiteness which leads to the notion of Ferrers language.

5.1. Incidence structures. We recall first some basic facts about incidence structures, Galois lattices and Ferrers relations. We follow the exposition given in [31].

An *incidence structure* R is a triple (V, ρ, W) where ρ is a subset of the product $V \times W$. We set $\rho^{-1} := \{(x, y) : (y, x) \in \rho\}$ and $R^{-1} := (W, \rho^{-1}, V)$, that we call the *dual* of R . We denote by $-\rho$ the relation $V \times W \setminus \rho$ and $-R$ the resulting incidence structure.

A subset of $V \times W$ of the form $X \times Y$ is a *rectangle*. Let $(X, Y) \in \mathcal{P}(V) \times \mathcal{P}(W)$. We set $R_\wedge(X) := \{y \in W : x\rho y \text{ for all } x \in X\}$, $R_\wedge^{-1}(Y) := \{x \in V : x\rho y \text{ for all } y \in Y\}$. We recall that the set $X \times Y$ is a maximal rectangle included into ρ if and only if $X = R_\wedge^{-1}(Y)$ and $Y = R_\wedge(X)$. The *Galois lattice* $\text{Gal}(R)$ of R is the collection, ordered by inclusion, of subsets of V of the form $R_\wedge^{-1}(Y)$ for $Y \in \mathcal{P}(W)$. This is a complete lattice; the largest element is $V (= R_\wedge^{-1}(\emptyset))$. Then, $\text{Gal}(R^{-1})$ is the collection, ordered by inclusion, of subsets of W of the form $R_\wedge(X)$ for $X \in \mathcal{P}(V)$. We recall the important fact that $\text{Gal}(R)$ and $\text{Gal}(R^{-1})$ are dually isomorphic. Since $\text{Gal}(R)$ consists of intersections of sets of the form $R^{-1}(y)$ for $y \in W$, $\text{Gal}(R)$ is finite if and only if the set of $R^{-1}(y)$ for $y \in W$ is finite; since $\text{Gal}(R)$ is dually isomorphic to $\text{Gal}(R^{-1})$, it is finite if and only if the set of $R(x)$ for $x \in V$ is finite.

We give two examples from the theory of ordered sets.

Fact 2. *If $R := (P, \leq, P)$ is an ordered set, then $\text{Gal}(R)$, the MacNeille completion of P , is a complete lattice in which every member is a join and a meet of elements of P . And $\text{Gal}(-R)$ is the set of final segments of P ordered by inclusion.*

Now, we mention the facts we need.

Let $R := (V, \rho, W)$ and $R' := (V', \rho', W')$ be two incidence structures. According to Bouchet [2], a *coding* from R into R' is a pair of maps $f : V \rightarrow V', g : W \rightarrow W'$ such that

$$x\rho y \iff f(x)\rho'g(y).$$

Bouchet's Coding Theorem [2] below is a striking illustration of the links between coding and embedding.

Theorem 15. *Let (T, \leq) be a complete lattice and R be an incidence structure. Then R has a coding into (T, \leq, T) if and only if $\text{Gal}(R)$ is embeddable in T .*

The basic facts about coding we need are the following:

Fact 3. *Let (f, g) be a coding from R into R' .*

- (a) *(f, g) is a coding from $\neg R$ to $\neg R'$;*
- (b) *$\text{Gal}(R)$ is embeddable into $\text{Gal}(R')$;*
- (c) *If f is surjective, then $\text{Gal}(R)$ identifies with an intersection closed subset of $\text{Gal}(R')$.*

Proof. (a) Immediate consequence of the definition.

(b) Follows from Bouchet's Theorem.

(c) The map $X \rightarrow X' = R'_{\wedge}{}^{-1}(R'_{\wedge}(X))$ is an embedding from $\text{Gal}(R)$ into $\text{Gal}(R')$ which preserves non-empty intersections. If f is surjective, then the least element of the Galois lattice is preserved, hence $\text{Gal}(R)$ identifies with an intersection closed subset of $\text{Gal}(R')$. \square

Fact 4. *Let $(f_i, g_i)_i$ be a family of codings from (V, ρ_i, W) into (V_i, θ_i, W_i) . Then, the pair $(\Pi_i f_i, \Pi_i g_i)$ is a coding from $(V, \cup_i \rho_i, W)$ into $(\Pi_i V_i, \neg \Pi_i \theta_i, \Pi_i W_i)$.*

Proof. Clearly $(\Pi_i f_i, \Pi_i g_i)$ is a coding from $(V, \cap_i \rho_i, W)$ into $(\Pi_i V_i, \Pi_i \theta_i, \Pi_i W_i)$. Indeed, if $x := (x_i)_i$ and $y := (y_i)_i$, then $x \Pi_i \theta_i y$ means $x_i \theta_i y_i$ for all i . From Fact 3 (a) we get that $(\Pi_i f_i, \Pi_i g_i)$ is a coding from $(V, \cup_i \rho_i, W)$ into $(\Pi_i V_i, \neg \Pi_i \theta_i, \Pi_i W_i)$. \square

Corollary 16. *If for every i , $V_i = W_i$ and θ_i is of the form $\neg \leq_i$ for some ordering \leq_i on V_i , then there is a coding from $(V, \cup_i \rho_i, W)$ into $(\Pi_i V_i, \neg \leq, \Pi_i V_i)$ where \leq is the product ordering on $\Pi_i V_i$.*

Fact 5. *Let $R := (V, \rho, W)$ be an incidence structure. Then $\text{Gal}(R)$ is finite if and only if $\text{Gal}(\neg R)$ is finite.*

Proof. For each $y \in V$, $(\neg R)^{-1}(y) = V \setminus R^{-1}(y)$. Since $\text{Gal}(R)$ is made of intersections of sets of the form $R^{-1}(y)$, if $\text{Gal}(R)$ is finite, the collection of such sets is finite, hence the collection of sets of the form $(\neg R)^{-1}(y)$ is finite too. Since $\text{Gal}(\neg R)$ is made of intersections of these sets, it is finite. \square

Fact 6. *Let $R_i := (V_i, \rho_i, W_i)$ be a family of incidence structures. Then $\text{Gal}(\Pi_i R_i)$ embeds into $\Pi_i \text{Gal}(R_i)$.*

Proof. Let $y := (y_i)_i \in \Pi_i W_i$. We have $(\Pi_i R_i)^{-1}(y) = \Pi_i R_i^{-1}(y_i)$. Hence, if $p_j : \Pi_i V_i \rightarrow V_j$ denotes the j^{th} projection, then for each $X \in \text{Gal}(\Pi_i R_i)$, we have $(p_i(X))_i \in \Pi_i \text{Gal}(R_i)$. This defines an embedding, proving our claim. If each member X of $\text{Gal}(\Pi_i R_i)$ is non-empty, then this embedding is an isomorphism. \square

Fact 7. *Suppose $V_i = V, W_i = W$ for each $i \in I$ and I finite. If $\text{Gal}(R_i)$ is finite for each $i \in I$, then $\text{Gal}((V, \cup_i \rho_i, W))$ is finite.*

Proof. According to Fact 5, this amounts to prove that $\text{Gal}((V, \cap_i \neg \rho_i, W))$ is finite. Fact 4 yields that $(V, \cup_i \rho_i, W)$ has a coding into $(V^V, \neg \Pi_i \neg \rho_i, W^W)$. According to (a) of Fact 3, $(V, \cap_i \neg \rho_i, W)$ has a coding into $(V^V, \Pi_i \neg \rho_i, W^W) \equiv \Pi_i (V, \neg \rho_i, W)$. According to (b) of Fact 3, $\text{Gal}((V, \cap_i \neg \rho_i, W))$ embeds into $\text{Gal}(\Pi_i (V, \neg \rho_i, W))$, which in turns embeds into $\Pi_i \text{Gal}(V, \neg \rho_i, W)$ from Fact 6. From Fact 5, each $\text{Gal}(V, \neg \rho_i, W)$ is finite. The result follows. \square

5.2. Languages and their Galois lattices. Galois lattices arose from group theory and geometry. We show here how they interact with language theory.

We represent subsets of the free monoid by incidence structures as follows.

Let A be an alphabet and L be a subset of A^* . Denoting by xy the concatenation of the words $x, y \in A^*$, define a binary relation ρ_L on A^* by:

$$x\rho_L y \iff xy \in L$$

and set $R_L := (A^*, \rho_L, A^*)$.

We mention without proof some properties of this association.

It preserves Boolean operations, that is:

Fact 8. (1) $\rho_{A^* \setminus L} = A^* \times A^* \setminus \rho_L$. Hence, $R_{A^* \setminus L} = \neg R_L$.
 (2) $\rho_{\cup_i L_i} = \cup_i \rho_{L_i}$ and $\rho_{\cap_i L_i} = \cap_i \rho_{L_i}$

Not every binary relation ρ on A^* can be of the form ρ_L . For an example, if ρ contains a pair (u, v) with $u, v \in A^*$, then ρ contains $\rho_{\{w\}}$ where $w := uv$. In fact, if ρ is a binary relation on A^* , set $\pi_\rho := \{uv : (u, v) \in \rho\}$; then ρ_{π_ρ} is the least binary relation of the form ρ_L containing ρ . In particular, $\rho_{\{w\}}$ is the least relation of the form ρ_L containing a pair (u, v) such that $w = uv$.

In the Boolean lattice made of relations of the form ρ_L , the atoms are of the form $\rho_{\{w\}}$ where w is any word in A^* . They form a partition of $A^* \times A^*$. Every binary relation of the form ρ_L is an union of some blocks of this partition.

We will say more about this association in Section 6.

We conclude this subsection by the following property.

If B is a subset of A^* , we set $R_L \upharpoonright B := (B, \rho_L \cap B \times B, B)$.

Fact 9. *The Galois lattices of R_L and $R_L \upharpoonright A^* \setminus L$ are isomorphic provided that L is a final segment of A^* .*

Proof. Observe that for every subset X of A^* , we have: $R_\wedge^{-1}(R_\wedge(X)) = R_\wedge^{-1}(R_\wedge(X \setminus L))$. \square

5.3. Proof of the first part of Theorem 2. From Corollary 16, and *c*) of Fact 3, in order to prove the first part of Theorem 2, namely that \mathcal{S}_F can be identified to an intersection closed subset of the set $\mathbf{F}(\mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1})$, all that we need is to prove that for each i , there is a coding (f_i, g_i) , with f_i surjective, from (A^*, ρ_{F_i}, A^*) into $(\mathbf{n}_i, \neg \leq, \mathbf{n}_i)$. But, this is false. From Fact 9, a coding from $(A^* \setminus F, \rho_{F_i}, A^* \setminus F)$ into $(\mathbf{n}_i, \neg \leq, \mathbf{n}_i)$ suffices. Let X_0, \dots, X_{n-1} be non-empty subsets of A and let $X := X_0 \cdots X_{n-1}$ (that is the set of words $x_0 \cdots x_{n-1}$ with $x_i \in X_i$) and let $F := \uparrow X$. Let $\mathbf{n} := \{0, \dots, n-1\}$ be equipped with the natural ordering. Let $f : A^* \setminus F \rightarrow \mathbf{n}$ and $g : A^* \setminus F \rightarrow \mathbf{n}$ defined as follows: For $v \in A^* \setminus F$, if there is $m \in \mathbb{N}$ such that $v \in \uparrow (X_0 \cdots X_{m-1})$ then $f(v)$ is the largest m having this property, otherwise $f(v) = 0$. For $w \in A^* \setminus F$, if there is $p \in \mathbb{N}$ such that $v \in \uparrow \{X_{p-1} \cdots X_{n-1}\}$ then $g(v)$ is the least p having this property, otherwise $g(v) = n-1$. By a straightforward verification, we have the following:

Fact 10. *The map f is surjective and the pair (f, g) form a coding from $(A^* \setminus F, \rho_F, A^* \setminus F)$ into $(\mathbf{n}, \neg \leq, \mathbf{n})$.*

This proves the first part of Theorem 2.

5.4. Proof of the second part of Theorem 2. Let $F := F_0 \cup \dots \cup F_{k-1}$, let $X_i \subseteq F$ such that $F_i = \uparrow X_i$ and let $X_{i_0}, \dots, X_{i_{n_i-1}} \subseteq A$ such that $X_{i_0} \cdots X_{i_{n_i-1}} = X_i$. From Fact 10 above, for each i , we have a coding (f_i, g_i) from $(A^* \setminus F_i, \rho_{F_i}, A^* \setminus F_i)$ into $(\mathbf{n}_i, \neg \leq, \mathbf{n}_i)$. The restrictions of f_i and g_i to $A^* \setminus F$ give a coding that we still denote (f_i, g_i) from $(A^* \setminus F_i, \rho_{F_i}, A^* \setminus F_i)$ into $(\mathbf{n}_i, \neg \leq, \mathbf{n}_i)$ and f_i is still surjective. From Fact 4, $(\Pi f_i, \Pi g_i)$ is a coding from $R_F := (A^* \setminus F, \cup \rho_{F_i}, A^* \setminus F)$ into $R' := (\mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1}, \neg \leq, \mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1})$ where \leq is the natural ordering on the direct product $\mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1}$. Since for each i , f_i is surjective, Πf_i is surjective and from (c) of Fact 3, $\text{Gal}(R_F)$ is identified with an intersection closed subsets of $\text{Gal}(R')$ which is $\mathbf{F}(\mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1})$ by Fact 2. \square

5.5. An explicit isomorphism. For reader's convenience, let us describe explicitly the isomorphism between \mathcal{S}_F and an intersection closed subset of $\mathbf{F}(\mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1})$. For sake of simplicity, let us suppose that F is finitely generated (which is the case if A is w.q.o.). Let $u_0, \dots, u_{k-1} \in A^*$, $n_0 := |u_0|, \dots, n_{k-1} := |u_{k-1}|$, and $F := \uparrow \{u_0, \dots, u_{k-1}\}$. Let $v \in A^*$; for each $i \in \{0, \dots, k-1\}$, let v'_i be the largest suffix of u_i such that $v''_i \leq v$ and let v'_i be the unique prefix of u_i such that $u_i = v'_i v''_i$. Set $s(v) := \{x := (x_0, \dots, x_{k-1}) \in \mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1} \text{ such that } x_i < |v_i| \text{ for all } i\}$ and $\tau(v) := \mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1} \setminus s(v)$. Define $\varphi : \mathcal{S}_F \rightarrow \mathbf{F}(\mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1})$ by setting

$$\varphi(X) := \bigcap_{v \in A^*, X \subseteq \rho^{-1}(v)} \tau(v) \quad .$$

For $w := w_0 \dots w_m \in A^*$ and $\ell < |w|$, let $w|_\ell = w_0 \dots w_{\ell-1}$ be the restriction of w to the first ℓ letters. Let $x := (x_0, \dots, x_{k-1}) \in \mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1}$, let $\mu(x) = \uparrow \{u_i|_{x_{i-1}} : 0 \leq i \leq k-1\}$ be the final segment of A^* generated by words of the form $u_i|_{x_{i-1}}$.

Define $\Psi : \mathbf{F}(\mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1}) \rightarrow \mathcal{S}_F$ by setting

$$\Psi(Y) := \bigcap_{x \in \mathbf{n}_0 \otimes \dots \otimes \mathbf{n}_{k-1} \setminus Y} \mu(x)$$

The maps φ and Ψ preserve intersections and $\Psi \circ \varphi = id_{\mathcal{S}_F}$.

5.6. Finiteness of the injective envelope.

Proposition 4. *Let F be a final segment of A^* . The following conditions are equivalent:*

- (i) *The injective envelope \mathcal{S}_F is finite;*
- (ii) *F is finite union $F = F_0 \cup \dots \cup F_i \cup \dots \cup F_{k-1}$ of final segments, each F_i generated by a set X_i of words u_i of the same length n_i all of the form $u_i = u_{i_0} \dots u_{i_j} \dots u_{i_{n_i-1}}$ with $u_{i_j} \in X_{i_j} \subseteq A$.*

Proof. (i) \Rightarrow (ii). Let us suppose that \mathcal{S}_F is finite. Let \mathcal{A}_F be the corresponding reflexive and involutive automaton with an initial state $x := A^*$ and a final state $y := F$. Let Q_F^* be the set of finite sequences $s := (s_0, \dots, s_n)$ such that all s_i are distinct states, $s_0 := x, s_n := y, (s_i, a_i, s_{i+1}) \in T_F$ for some letter a_i . Since \mathcal{S}_F is finite this set is finite too. For each $s = (s_0, \dots, s_n)$ let $X_{s_j} := \{a \in A : (s_j, a, s_{j+1}) \in T_F\}$ and let $X_s := \{a_0 \dots a_{n-1} : a_j \in X_{s_j}\}$ and let $F_s := \uparrow X_s$. It is easy to check that $F = \bigcup_{s \in Q_F^*} F_s$, hence has the form mentioned above.

(ii) \Rightarrow (i). Apply Theorem 2. \square

Theorem 17. *Let $\mathcal{H} := \mathbf{F}(A^*)$ be the Heyting algebra made of final segments of A^* . The following conditions are equivalent:*

- (i) *A is well-quasi-ordered;*

(ii) *The injective envelope $\mathcal{N}(E)$ of every finite metric space E is finite.*

Proof. $\neg(i) \implies \neg(ii)$. Let F be a final segment of A^* . According to Proposition 4, the injective envelope \mathcal{S}_F is infinite whenever for each decomposition $F = \cup F_i$, some F_i cannot be generated by a set of words having a bounded length. This is the case if F is generated by an infinite antichain X made of words of unbounded length. If A is not w.q.o., then there is an infinite bad sequence of letters, say a_0, \dots, a_n, \dots . The set $X = \{a_0, a_1a_2, a_3a_4a_5, a_6a_7a_8a_9, \dots\}$ is such an example.

$(i) \implies (ii)$. According to Proposition 1, it suffices to show that the injective envelope of a two-element metric space is finite. Let F be a final segment of A^* . If A is w.q.o., F satisfies condition (ii) of Proposition 4, hence \mathcal{S}_F is finite. \square

Comments. Proposition 4 and Theorem 17 are special instances of two basic facts of language theory concerning rational languages, namely Myhill-Nerode Theorem (see [34] Theorem 2.3, p.247) and a result of Ehrenfeucht-Haussler-Rozenberg (Theorem 3.3 of [12], see [34] Theorem 5.3, p.296). For brevity, let us put these results together in the context of monoids. Let us recall that if L is a subset of a monoid M , the *residuals* of L are sets of the form $Lv^{-1} := \{u \in M : uv \in L\}$ for $v \in M$ and of the form $u^{-1}L := \{v \in M : uv \in L\}$ for $u \in M$. The *syntactic congruence* of L is the largest congruence \equiv_L on M for which L is an union of classes. The set L is *recognisable* if it is the union of classes of a congruence on M which has finitely many classes (alias finite index). The aforementioned results read as follows:

Theorem 18. *For a subset L of a monoid M , the following properties are equivalent:*

- (i) *L is recognisable;*
- (ii) *The set of residuals $\{Lv^{-1} : v \in M\}$ is finite;*
- (iii) *The set of residuals $\{u^{-1}L : u \in M\}$ is finite;*
- (iv) *The syntactic congruence \equiv_L has finite index;*
- (v) *L is a final segment of a well-quasi-ordered set on M which is compatible with the monoid operation.*

The equivalences from (i) to (iv) is Myhill-Nerode Theorem. The equivalence with (v) is Ehrenfeucht-Haussler-Rozenberg Theorem.

Let $R := (M, \rho_L, M)$ be the incidence structure where $\rho_L := \{(u, v) \in M \times M : uv \in L\}$. Then $u\rho_L v \iff uv \in L$ and $R(u) = u^{-1}L$, $R^{-1}(v) = Lv^{-1}$ for all $u, v \in M$. Conditions (ii) and (iii) in the theorem above express both that the Galois lattice $\text{Gal}(R)$ is finite.

According to Higman's Theorem, if the alphabet A is w.q.o., A^* equipped with the Higman ordering is w.q.o. Since this ordering is compatible with concatenation, we may apply (v) to every final segment F of A^* , obtaining that the Galois lattice $\text{Gal}(R)$ is finite. Since the domain of this lattice is \mathcal{S}_F , implication (i) \implies (ii) of Theorem 17 follows.

6. INTERVAL ORDERS, FERRERS RELATIONS AND INJECTIVE ENVELOPE

We record the characterization of interval orders and Ferrers relations. These two notions are intertwined. Interval orders are those orderings for which the irreflexive part is a Ferrers relation. Ferrers relations have been introduced by J.Riguet [32]. Interval orders are studied by Fishburn in [13]. Part of the characterization of these relations given below is due to Wiener [39]. Let C be a chain; an *interval* of C is any subset of I such that

$$x \in I, y \in I \text{ and } x \leq z \leq y \text{ imply } z \in I. \quad (1)$$

The collection $\text{Int } C$ of non-empty intervals of C is ordered as follows:

$$X < Y \text{ if } x < y \text{ for all } x \in X, y \in Y. \quad (2)$$

Let P be a poset. The ordering on P , or P itself, is an *interval order* if P is order isomorphic to a collection of non-empty intervals of some chain C , ordered by condition (2). To each incidence structure $R := (V, \rho, W)$ we associate a poset $B(R) := (P, \leq)$ defined as follows:

The domain of P is $V \times \{0\} \cup W \times \{1\}$, for $u = (x, i), v = (y, j) \in P$, the order relation is defined by:

$$u < v \text{ if } i < j \text{ and } x\rho y. \quad (3)$$

Let $R := (V, \rho, W)$ be an incidence structure. We say that R is *Ferrers* or ρ is a *Ferrers relation* if R satisfies one of the following conditions (see [13]):

Proposition 5. *Let $R := (V, \rho, W)$ be an incidence structure. The following conditions are equivalent:*

- (i) *The set $\{R(x) : x \in V\}$ is totally ordered by inclusion;*
- (ii) *The set $\{R^{-1}(y) : y \in W\}$ is totally ordered by inclusion;*
- (iii) *The Galois lattice $\text{Gal}(R)$ is totally ordered by inclusion;*
- (iv) *R has a coding into a chain;*
- (v) *The poset $B(R)$ does not embed the direct sum $2 \oplus 2$ of two copies of the 2-element chain 2;*
- (vi) *The ordering on $B(R)$ is an interval order;*
- (vii) *$x\rho y$ and $x'\rho y'$ imply $x\rho y'$ or $x'\rho y$ for all $x, x' \in V, y, y' \in W$.*

We say that a language L is *Ferrers* if the relation ρ_L is Ferrers.

According to condition (vii) of Proposition 5, this amounts to the following condition:

$$xx' \in L \text{ and } yy' \in L \text{ imply } xy' \in L \text{ or } yx' \in L \text{ for all } x, x', y, y' \in A^*. \quad (4)$$

The study of Ferrers relations leads to the notion of Ferrers dimension of a binary relation: the least number of Ferrers relations whose intersection is this relation. The study of this notion, initiated by Bouchet in his thesis [2], yields numerous interesting results (e.g., see [6, 8]). This suggest to look at the same direction in the theory of languages. But, we may notice that contrarily to the case of relations, not every language is Ferrers, or is an intersection of Ferrers languages. In fact, if w is any word, $\rho_{\{w\}}$ is a Ferrers relation only if $w = \square$ (indeed, let $R := (A^*, \rho_{\{w\}}, A^*)$, then $\text{Gal}(R) = \{\{u\} : u \text{ prefix of } w\} \cup \{\emptyset, A^*\}$, thus if $w \neq \square$, the Galois lattice has at least two incomparable elements, namely $\{w\}$ and $\{\square\}$ plus a top and a bottom, thus it is not a chain). Thus, if $w \neq \square$, $\{w\}$ is not a Ferrers language and since it is a singleton, it is not a union of Ferrers languages, hence its complement $A^* \setminus \{w\}$ is not an intersection of Ferrers languages.

Since the complement of a Ferrers relation is Ferrers, the complement of a Ferrers language is Ferrers (apply (1) of Fact 8). The concatenation of two Ferrers languages is not Ferrers in general. For a simple minded example, let $A := \{a, b\}$, let $U := \{a^n : n \geq 2\}$, $U' := \{b^n : n \geq 2\}$. Each of these languages is Ferrers, but the concatenation UU' is not: let $x := a^2b, y := b$ and $x' := a, y' := ab^2$. Then $xy = x'y' = a^2b^2 \in UU'$ but neither $xy' = a^2bab^2$ nor $x'y = ab$ belong to UU' .

Recall that for two words u and v , u is a *prefix* of v if $v = uw$ for some word w ; similarly, u is a *suffix* of v if $v = wu$ for some word w .

Fact 11. *Let U and U' be two subsets of A^* . If U, U' are Ferrers and U is a final segment for the prefix ordering or U' is a final segment for the suffix ordering then the concatenation UU' is Ferrers.*

Proof. Let $L := UU'$, let $xx' \in L$ and $yy' \in L$. We prove that either xy' or $x'y$ belong to L . There are four cases to consider; we only consider two, the others being similar.

Case 1. $x = x_1x_2, y = y_1y_2$ with $x_1, y_1 \in U$ and $x_2x' \in U'$ and $y_2y' \in U'$. Since U' is Ferrers, either $x_2y' \in U'$ or $y_2x' \in U'$. In the former case $xy' \in UU'$ whereas in the latter $yx' \in UU'$.

Case 2. $x = x_1x_2, y' = y'_1y'_2$ with $x_1 \in U, yy'_1 \in U, x_2x' \in U'$ and $y'_2 \in U'$. If U is a final segment for the prefix ordering, then $x_1x_2y'_1 \in U$ since x_1 belongs to U and is a prefix of $x_1x_2y'_1$. Thus $xy' = x_1x_2y'_1y'_2 \in UU'$. If U' is a final segment for the suffix ordering, then $x_2y'_1y'_2 \in U'$ since y'_2 belongs to U' and is a suffix of y'_2 . Thus $xy' = x_1x_2y'_1y'_2 \in UU'$. \square

Corollary 19. *The concatenation of finitely many Ferrers final segments of A^* is a Ferrers final segment.*

We recall that an *ideal* of an ordered set P is any non-empty initial segment \mathcal{I} which is up-directed (that is any two elements x and y of \mathcal{I} have an upper bound z in \mathcal{I}). Filters are defined dually. Ideals of P are the join-irreducible elements of the lattice $I(P)$ of initial segments of P . Ideals of the poset A^* equipped with the Higman ordering have been described when A is finite by Jullien [18] and by us [20] when A is an ordered alphabet possibly infinite. According to Jullien, an *elementary ideal* of A^* is any set of the form $J \cup \{\square\}$ for some non empty ideal J of A , a *star-ideal* is any set of the form I^* for some initial segment I of A . Products of ideals are ideals. It is proved in [20] that every ideal is a finite product of elementary and star-ideals if and only if the alphabet is well-quasi-ordered.

Fact 12. *Ideals and filters of A^* are Ferrers.*

Proof. Let \mathcal{I} be an ideal of A^* . Suppose $xx', yy' \in \mathcal{I}$. We prove that xy' or $yx' \in \mathcal{I}$. Let $z \in \mathcal{I}$ such that $xx', yy' \leq z$. Let z_1 be the least prefix of z such that $x, y \leq z_1$ and z_2 the corresponding suffix, i.e., $z = z_1z_2$. If $x' \leq z_2$ or $y' \leq z_2$, then since $x, y \leq z_1$ then xy' or $yx' \in \mathcal{I}$. If not, then since $xx' \leq z$ and $yy' \leq z$, we have $x, y \leq z_1^-$, where z_1^- is obtained from z_1 by deleting its last letter. This contradicts the choice of z_1 . Hence, \mathcal{I} is Ferrers. Let \mathcal{F} be a filter of A^* . Let $xx', yy' \in \mathcal{F}$. Let $z \in \mathcal{F}$ such that $z \leq xx'$ and $z \leq yy'$. Write $z = z^x z^{x'}$ with $z^x \leq x, z^{x'} \leq x'$ and $z = z^y z^{y'}$ with $z^y \leq y, z^{y'} \leq y'$. Either $z^{y'} \leq z^{x'}$ or $z^{x'} \leq z^{y'}$. In the first case, since $z^y \leq y$ and $z^{y'} \leq z^{x'} \leq x'$ we have $z = z^y z^{y'} \leq yx'$, hence $yx' \in \mathcal{F}$. In the second case we obtain similarly $xy' \in \mathcal{F}$. Hence \mathcal{F} is Ferrers. \square

For every $u \in A^*$, the initial segment $\downarrow u$ of A^* is an ideal and the final segment $\uparrow u$ of A^* is a filter, hence:

Corollary 20. *For every $u \in A^*$, the initial segment $\downarrow u$ of A^* and the final segment $\uparrow u$ of A^* are Ferrers.*

Since $\{u\} = \downarrow u \cap \uparrow u$, we have:

Corollary 21. *For every $u \in A^*$, $\{u\}$ is an intersection of two Ferrers languages and $A^* \setminus \{u\}$ is an union of two Ferrers languages.*

If A is w.q.o., every final segment is finitely generated and every ideal is a finite union of ideals, hence from the second part of Corollary 20 and from the first part of Fact 12, we obtain.

Corollary 22. *If A is w.q.o., every final segment is a finite union and a finite intersection of Ferrers languages.*

From Corollary 19 and Fact 12 follows:

Proposition 6. *Final segments of A^* which are finite product of complement of ideals of A^* are Ferrers.*

According to Corollary 21, for every $u \in A^*$, $A^* \setminus \{u\}$ is an union of two Ferrers languages, hence:

Proposition 7. *Every language is a union, possibly infinite, of a family of intersections of two Ferrers languages.*

According to Corollary 22, if the alphabet is w.q.o. (and particularly, if it is finite), Boolean combinations of final segments, alias piecewise testable languages, are Boolean combination of rational Ferrers languages.

• *If the alphabet A consists of one letter, say a , these two Boolean algebras coincide.* Indeed, if L is Ferrers then with respect to the natural order on $A^* := \{a^n : n \in \mathbb{N}\}$, it is convex. Otherwise, there are $n < p < m \in \mathbb{N}$ such that $a^n, a^m \in L$, $a^p \notin L$. Choosing n, m with the difference $m - n$ minimum, we have $a^q \notin L$ for $n < p < m$. Let $X := (a^n)^{-1}L$ and $Y := (a^{m-1})^{-1}L$. Then \square and $a^{m-n} \in X$ but $\square \notin X$, whereas $a \in Y$ but $\square \notin Y$. Hence, X and Y are incomparable with respect to inclusion, contradicting the fact that L is Ferrers. Being convex, L is the intersection of an initial segment with a final segment, thus it is piecewise testable.

• *If $A := \{a, b\}$, with $a \neq b$, then $L := A^*b$ is rational and Ferrers and not piecewise testable.* Indeed, let $Q_L := \{u^{-1}L : u \in A^*\}$, then Q_L has two elements, namely L and $L' := \{\square\} \cup L$ (in fact $a^{-1}L = L$, $b^{-1}L = L'$, $a^{-1}L' = L$, $b^{-1}L' = L'$). The fact that L is not piecewise testable follows from Stern's criterium ([36] Theorem 1.2): the sequence $(u_n)_{n \in \mathbb{N}}$ defined by $u_{2n} := (ba)^n b$ and $u_{2n+1} := (ba)^{n+1}$ is increasing for the subword ordering while $a_{2n} \in L$ and $a_{2n+1} \notin L$ for all $n \in \mathbb{N}$.

The language L above has dot-depth one. Is this general? That is:

Question. *Do rational Ferrers languages have dot-depth one?*

We relate Ferrers piecewise testable languages and structural properties of transition systems.

Theorem 23. *Let F be a final segment of A^* . The following conditions are equivalent:*

- (1) *F is Ferrers;*
- (2) *The space \mathcal{S}_F is linearly orderable.*

Proof. Proposition 5 and Proposition 2. □

Corollary 24. *The finitely indecomposable absolute retracts are linearly orderable.*

Proof. Let E be a finitely indecomposable absolute retract. From Theorem 6, E is isomorphic to \mathcal{S}_F where F is join-irreducible in the lattice $\mathbf{F}(A^*)$ ordered by reverse of inclusion. The fact that F is join-irreducible amounts to the fact that $A^* \setminus F$ is an ideal of A^* . According to Fact 12, $A^* \setminus F$ is Ferrers. Hence, its complement F is Ferrers. The result follows from Theorem 23. □

REFERENCES

- [1] B. Banaschewski, G. Bruns, *Categorical characterization of the MacNeille completion*, Arch. Math. (Basel) 18 (1967) 369–377.
- [2] A. Bouchet, *Codages et dimensions de relations binaires*, Annals of discrete Mathematics 23, Orders: Description and roles, (M. Pouzet, D. Richard Eds), 387-396. 1984.

- [3] L.M. Blumenthal, *Boolean geometry*, I. Rend. Circ. Mat. Palermo (2) 1 (1952), 343–360.
- [4] L.M. Blumenthal, *Theory and applications of distance geometry*, Second edition Chelsea Publishing Co., New York 1970 xi+347 pp.
- [5] L.M. Blumenthal, K. Menger, *Studies in geometry*, W. H. Freeman and Co., San Francisco, Calif. 1970 xiv+512 pp.
- [6] O. Cogis, *On the Ferrers dimension of a digraph*, Discrete Math. 38 (1982) 47–52.
- [7] E. Corominas, *Sur les ensembles ordonnés projectifs et la propriété du points fixe*, C.R.Acad.Sc. Paris, Série A311 (1990), 199–204.
- [8] J.P. Doignon, A. Ducamp, J.C. Falmagne, *On realizable biorders and the biorder dimension of a relation*, J. Math. Psych. 28(1984) 73–109.
- [9] A.W.N. Dress, *Trees, tight extensions of metric spaces, and the cohomological dimension of certain groups, a note on combinatorial properties of metric spaces*, Adv. in Math. 53 (3)(1984), 321–402.
- [10] A.W.N. Dress, *Towards a classification of transitive group actions on finite metric spaces*, Adv. in Math. 74 2 (1989),163–189.
- [11] D. Duffus, I. Rival, *A structure theory of ordered sets*, J.of Discrete Math. 35 (1981), 53–118.
- [12] A. Ehrenfeucht, D. Haussler, G. Rozenberg, *On regularity of context-free languages*,Theoret. Comput. Sci. 27 (1983), no. 3, 311–332.
- [13] P.C. Fishburn, *Interval Orders and Interval Graphs*, Wiley, 1985.
- [14] G. Higman *Ordering by divisibility in abstract algebra*, Proc. London Math. Soc (3) 2 (1952), 326–336.
- [15] A. Hudry, *Rétractions, corétractions et envelope injective d’une algèbre de transitions*, Discrete Math. 247 (2002), no. 1-3, 117–134.
- [16] A. Hudry, *Injective envelope and parallel decomposition of a transition system*, Discrete Math. 289 (2004), no. 1-3, 45–61.
- [17] E. Jawhari, D. Misane, M. Pouzet, *Retracts graphs and ordered sets from the metric point of view*, (I.Rival ed) Contemporary Mathematics, Vol 57,175–226. 1986.
- [18] P. Jullien, *Sur un théorème d’extension dans la théorie des mots*, C. R. Acad. Sci. Paris Série A-B 266 (1968) 851–854.
- [19] K. Kaarli, S. Radeleczki: *Representation of integral quantales by tolerances*. Algebra Universalis 79 (5) (2018), <https://doi.org/10.1007/s00012-018-0484-1>
- [20] M. Kabil, M. Pouzet, *Une extension d’un théorème de P. Jullien sur les âges de mots*, RAIRO Inform. Théor. Appl. 26 (1992), no. 5, 449–482.
- [21] M. Kabil, M. Pouzet, *Indécomposabilité et irréductibilité dans la variété des rétractes absolus des graphes réflexifs*, C.R.Acad.Sci. Paris Série A 321 (1995), 499–504.
- [22] M. Kabil, *Une approche métrique de l’indécomposabilité et de l’irréductibilité dans la variété des rétractes absolus des graphes et des systèmes de transitions*, Thèse de doctorat d’État, Université Hassan II Aïn Chock, Casablanca, 19 Décembre 1996.
- [23] M. Kabil, M. Pouzet, *Injective envelope of graphs and transition systems*, Discrete Math.192 (1998), 145–186.
- [24] M. Kabil, M. Pouzet, I.G. Rosenberg, *Free monoids and metric spaces*, To the memory of Michel Deza, Europ. J. of Combinatorics, Online publication complete: April 2, 2018, <https://doi.org/10.1016/j.ejc.2018.02.008>. arXiv: 1705.09750v1, 27 May 2017.
- [25] A.D. Korshunov, *The number of monotonic Boolean functions*, Problemy Kibern, 38 (1981), 5–108 (Russian), 272, MR 83h : 06012.
- [26] M. Lothaire, *Combinatorics on words*, Encyclopedia Math. Appl., 17, Addison-Wesley, 1983.
- [27] M.Nivat, Personnel communication, 1989.
- [28] R. Nowakowski, I. Rival, *The smallest graph variety containing all paths*, J. of Discrete Math. 43 (1983), 185–198.
- [29] M. Pouzet, *Une approche métrique de la rétraction dans les ensembles ordonnés et les graphes*, (French) [A metric approach to retraction in ordered sets and graphs] Proceedings of the conference on infinitistic mathematics (Lyon, 1984), 59–89, Publ. Dép. Math. Nouvelle Sér. B, 85-2, Univ. Claude-Bernard, Lyon, 1985.
- [30] M. Pouzet, I.G. Rosenberg, *General metrics and contracting operations*, Graphs and combinatorics (Lyon, 1987; Montreal, PQ, 1988). Discrete Math. 130 (1994),103–169.
- [31] M. Pouzet, H. Si Kaddour, N. Zaguia, *Finite dimensional scattered posets*, European J. of Combinatorics, 37(2014), 79–99.
- [32] J. Riguet, *Les relations de Ferrers*, C.R.Acad.Sc. Paris Série A 232 (1951), 1729–1730.
- [33] F. Saïdane, *Graphe et languages: une approche métrique*, Thèse de doctorat, Université Claude-Bernard, Lyon, Novembre 1991.

- [34] J. Sakarovitch, *Elements of automata theory*, Translated from the 2003 French original by Reuben Thomas. Cambridge University Press, Cambridge, 2009.
- [35] I. Simon, *Piecewise testable events*, Automata theory and formal languages (Second GI Conf., Kaiserslautern, 1975), pp. 214-222. Lecture Notes in Comput. Sci., Vol. 33, Springer, Berlin, 1975.
- [36] J. Stern, *Characterizations of some classes of regular events*, Theoretical Computer Science 35 (1985) 17-42.
- [37] G. Viennot, *Factorisations des monoïdes*, Thèse de 3^{ème} cycle, Paris, Janvier 1971.
- [38] D. Wiedemann, *A computation of the eight Dedekind number*, Order 8 (1991), 5-6.
- [39] N. Wiener, *A contribution to the theory of relative position*, Proc. Camb. Philos. Soc. 17 (1914), 441-449.

LABORATOIRE MATHÉMATIQUES ET APPLICATIONS, DÉPARTEMENT DE MATHÉMATIQUES, FACULTÉ DES SCIENCES ET TECHNIQUES, UNIVERSITÉ HASSAN II -CASABLANCA, BP 146 MOHAMMEDIA, MOROCCO.
E-mail address: `kabilfstm@gmail.com`

UNIV. LYON, UNIVERSITÉ CLAUDE-BERNARD LYON1, CNRS UMR 5208, INSTITUT CAMILLE JORDAN, 43 BD. 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX, FRANCE AND MATHEMATICS & STATISTICS DEPARTMENT, UNIVERSITY OF CALGARY, CALGARY, ALBERTA, CANADA T2N 1N4
E-mail address: `pouzet@univ-lyon1.fr`, `mpouzet@ucalgary.ca`