

HAYDN

High-precision Asteroseismology of DeNse stellar fields

Andrea Miglio · Léo Girardi · Frank Grundahl · Benoit Mosser · Nate Bastian · Angela Bragaglia · Karsten Brogaard · Gaël Buldgen · William Chantereau · William Chaplin · Cristina Chiappini · Marc-Antoine Dupret · Patrick Eggenberger · Mark Gieles · Robert Izzard · Daisuke Kawata · Christoffer Karoff · Nadège Lagarde · Ted Mackereth · Demetrio Magrin · Georges Meynet · Eric Michel · Josefina Montalbán · Valerio Nascimbeni · Arlette Noels · Giampaolo Piotto · Roberto Ragazzoni · Igor Soszyński · Eline Tolstoy · Silvia Toonen · Amaury Triaud · Fiorenzo Vincenzo

Received: date / Accepted: date

A list of people who have expressed interest for HAYDN is available here:
<https://www.asterochronometry.eu/haydn/people.html>.

Andrea Miglio
School of Physics and Astronomy, University of Birmingham, UK E-mail: andrea.miglio@unibo.it
Dipartimento di Fisica e Astronomia, Università degli Studi di Bologna, Bologna, Italy
INAF-OAS Bologna, Italy

Léo Girardi
INAF Osservatorio Astronomico di Padova, Italy

Frank Grundahl
Stellar Astrophysics Centre (SAC), Aarhus, Denmark

Benoit Mosser
LESIA, Observatoire de Paris, Université PSL, CNRS, Sorbonne Université, Université de Paris, 92195 Meudon, France

Nate Bastian
ARI, Liverpool John Moores University, Liverpool, UK

Angela Bragaglia
INAF-OAS Bologna, Italy

Karsten Brogaard
Stellar Astrophysics Centre (SAC), Aarhus, Denmark
Astronomical Observatory, Institute of Theoretical Physics and Astronomy, Vilnius Univer-

sity, 10257 Vilnius, Lithuania

Gaël Buldgen
Observatoire de Genève, Université de Genève, Switzerland

William Chantereau
ARI, Liverpool John Moores University, Liverpool, UK

William Chaplin
School of Physics and Astronomy, University of Birmingham, UK

Cristina Chiappini
AIP, Potsdam, Germany

Marc-Antoine Dupret
STAR Institute, University of Liège, Belgium

Patrick Eggenberger
Observatoire de Genève, Université de Genève, Switzerland

Mark Gieles
ICREA, Pg. Lluís Companys 23, 08010 Barcelona, Spain
Institut de Ciències del Cosmos (ICCUB), Universitat de Barcelona (IEEC-UB), E08028
Barcelona, Spain

Robert Izzard
Astrophysics Research Group, University of Surrey, Guildford, UK

Daisuke Kawata
MSSL, University College London, Surrey, UK

Christoffer Karoff
Stellar Astrophysics Centre (SAC), Aarhus, Denmark

Nadège Lagarde
Institut UTINAM, UMR6213, OSU THETA, Observatoire de Besançon, France

Ted Mackereth
School of Physics and Astronomy, University of Birmingham, UK
Canadian Institute for Theoretical Astrophysics, University of Toronto, Toronto, Canada
Dunlap Institute for Astronomy and Astrophysics, University of Toronto, Toronto, Canada
David A. Dunlap Department for Astronomy and Astrophysics, University of Toronto,
Toronto, Canada

Demetrio Magrin
INAF Osservatorio Astronomico di Padova, Italy

Georges Meynet
Observatoire de Genève, Université de Genève, Switzerland

Eric Michel
LESIA, Observatoire de Paris, Meudon, France

Josefina Montalbán
School of Physics and Astronomy, University of Birmingham, UK

Valerio Nascimbeni
INAF Osservatorio Astronomico di Padova, Italy

Arlette Noels
STAR Institute, University of Liège, Belgium

Giampaolo Piotto
Dipartimento di Fisica e Astronomia, Università di Padova, Italy

Roberto Ragazzoni

Abstract In the last decade, the *Kepler* and CoRoT space-photometry missions have demonstrated the potential of asteroseismology as a novel, versatile and powerful tool to perform exquisite tests of stellar physics, and to enable precise and accurate characterisations of stellar properties, with impact on both exoplanetary and Galactic astrophysics. Based on our improved understanding of the strengths and limitations of such a tool, we argue for a new small/medium space mission dedicated to gathering high-precision, high-cadence, long photometric series in dense stellar fields. Such a mission will lead to breakthroughs in stellar astrophysics, especially in the metal poor regime, will elucidate the evolution and formation of open and globular clusters, and aid our understanding of the assembly history and chemodynamics of the Milky Way’s bulge and a few nearby dwarf galaxies.

Keywords Stars: low-mass · globular clusters · Galaxy: bulge · Galaxies: dwarf · Asteroseismology

1 Context and motivation

Primarily driven and motivated by the impetus behind the search for transiting exoplanets, high-precision, high-cadence, long photometric surveys from space have also led to major advances in the study and interpretation of global, resonant pulsation modes in stars (asteroseismology).

The discoveries enabled by asteroseismic studies with data from CoRoT, *Kepler*, K2, and now TESS, not only are revolutionising our knowledge of stellar evolution, but have also demonstrated the relevance and potential of this novel technique well beyond the realm of stellar physics. The key role of asteroseismology in providing precise and accurate characterisations of planet-host stars is widely recognised and, e.g., is deeply rooted into the core programme of the future ESA PLATO mission. Moreover, asteroseismic constraints for ensembles of stars (in particular red giants) have given us the ability to measure masses (hence infer ages) of tens of thousands of stars across the Galaxy.

INAF Osservatorio Astronomico di Padova, Italy

Igor Soszyński
Astronomical Observatory, University of Warsaw, Poland

Eline Tolstoy
Kapteyn Astronomical Institute, University of Groningen, NL

Silvia Toonen
School of Physics and Astronomy, University of Birmingham, UK
Anton Pannekoek Institute, University of Amsterdam, Amsterdam, the Netherlands

Amaury Triaud
School of Physics and Astronomy, University of Birmingham, UK

Fiorenzo Vincenzo
School of Physics and Astronomy, University of Birmingham, UK
Department of Astronomy & Center for Cosmology and AstroParticle Physics, The Ohio State University, Columbus, USA

Coupling asteroseismic constraints to precise and accurate astrometric data from *Gaia*, and data from large-scale spectroscopic surveys, enables the study of the chemical and dynamical evolution of our Galaxy with unprecedented temporal resolution.

To some extent, results from the exploitation of asteroseismic data went even beyond the expectations of those who, starting from the early 1980s, have been advocating for a European space mission dedicated to stellar seismology (see e.g. Roxburgh, 2002, for a review on the formidable foresight and decades-long efforts leading eventually to space-based asteroseismology). As discussed in more detail below, results from *Kepler* and CoRoT have primarily demonstrated the potential of asteroseismology as a versatile and powerful tool. With the benefit of hindsight, i.e. having now a better understanding of the strengths and limitations of such a tool, we can propose a mission design that will lead to breakthroughs in three broad areas:

1. high-precision stellar astrophysics, especially in the metal poor regime,
2. evolution and formation of stellar clusters,
3. assembly history and chemical evolution of the Milky Way's bulge and a few nearby dwarf galaxies.

We argue that a mission dedicated to high-precision, high-cadence, long photometric time series to dense stellar fields is needed to address specific, long-standing key questions in those three areas, which we describe in detail in Sec. 2.

For instance, high-quality asteroseismic constraints for large samples of coeval and initially-chemically-homogeneous stars in open and globular clusters, and in different evolutionary stages, will then be coupled with spectroscopic and astrometric data to assemble high-precision stellar physics laboratories to test stellar evolution models to unprecedented precision.

The paper is organised as follows: in Sec. 2 we describe in detail the proposed science questions. We then translate scientific requirements into a list of targets and we provide a possible mission design aided by simulations / calibrations using *Kepler* data (Sec. 3). Before delving into the science objectives, we provide below a succinct summary about the inferences on stellar structure that one can make from the observation of solar-like oscillations (Sec. 1.1), and clarify the limitations of present and planned asteroseismic missions (Sec. 1.2).

1.1 Preamble - what can we infer from high-precision, high-cadence, long photometric time series of solar-like oscillators?

Among the various classes of pulsating stars, we focus here primarily on those showing rich spectra of solar-like oscillations, which are global oscillation modes excited and intrinsically damped by turbulence in the outermost layers of convective envelopes.

These oscillations are ubiquitous in stars with sufficiently deep convective envelopes, hence they can be detected in low- and intermediate-mass stars




Fig. 1 Power spectra based on *Kepler*'s photometric observations of a red-giant stars (KIC12008916). The *upper-left panel* shows examples of the average seismic parameters large frequency separation ($\Delta\nu$) between radial modes and have illustrated the oscillation envelope of power (red thick line), the maximum of which is labelled as ν_{\max} . The *lower-left panel* shows the complex, rich spectrum of individual mode frequencies, which can be however robustly interpreted and used to make detailed inferences about both global and local stellar features (see Sec. 1.1). Symbols represent the radial modes (red squares), the dipole mixed pressure-gravity modes (green triangles with extended bars), quadrupole modes (blue pentagons), and octupole modes (yellow hexagons). The *right panel* shows a cartoon of the internal structure of a red-giant-branch star, together with an example of the perturbation associated to the radial displacement of a mixed pressure-gravity mode, which is able to probe both the envelope and the core of such a star.

belonging to different evolutionary stages (main sequence, subgiant phase, along the evolution on the red-giant branch, in the core-He burning phase and the early asymptotic giant branch (AGB)). This means that we can explore large mass (approximately $0.8\text{-}3 M_{\odot}$) and age intervals, including the low-mass, low-metallicity domain, which is key to studies of the early Universe. At the same time, e.g. within a given cluster, we can study the structures and properties of a wide range of coeval stars, using a homogeneous sample.

Moreover, solar-like oscillations show very rich, yet relatively simple frequency patterns that one can robustly interpret aided by well-established stellar pulsation theory. The information on global and local features of stellar interiors that can be extracted from such spectra is succinctly presented below (see Chaplin and Miglio, 2013; Hekker and Christensen-Dalsgaard, 2017; Garcia and Ballot, 2019, for recent, comprehensive, reviews on the subject).

- Inference on *global stellar properties*. High-precision mean densities, masses, radii, ages, and envelope He mass fraction can be inferred via detailed forward and inverse modelling techniques. This can be done using asteroseismic constraints of different quality: from average seismic parameters to individual-mode frequencies (see Fig. 1). Moreover, seismically inferred radii coupled with effective temperature allow inferring distances, with typical uncertainties of a few percent, which make them competitive with Gaia for the faint targets.
- Constraints on the *internal rotation rate* of stars. These are possible once rotational frequency splittings are measured, which is generally possible both in main-sequence (MS) stars (rotational splitting of pressure (p)

modes, probing the envelope/surface rotation) and in subgiants and giants (core rotation inferred from the splitting of gravity (g) and mixed modes, which we do observe in evolved stars). In evolved stars (e.g., stars showing mixed p-g modes) it is primarily the core rotation we can measure, with perhaps subtleties on more detailed, local inferences on the rotational profile (Beck et al., 2012; Di Mauro et al., 2016; Deheuvels et al., 2012; Mosser et al., 2013; Gehan et al., 2018). In MS stars, it is the envelope rotation which is accessible to observations, including information on latitudinal differential rotation (Lund et al., 2014; Nielsen et al., 2017; Bazot et al., 2019; Benomar et al., 2018).

- The *inclination angles of the star’s spin axis* with respect to the line of sight can potentially be inferred from measuring the relative amplitudes of the different components of rotationally split frequencies (e.g., Gizon and Solanki, 2003; Chaplin et al., 2013; Huber et al., 2013; Corsaro et al., 2017; Kuszlewicz et al., 2019).
- Detailed constraints on the *internal structure* of stars can be inferred by looking at specific signatures in the oscillation frequencies. To a first approximation, from the (mixed) oscillation modes patterns one can infer average quantities characteristics of high-order p and g modes, i.e., respectively, large frequency spacing ($\Delta\nu$) and period spacing (ΔP). $\Delta\nu$ (p modes) is primarily a proxy for the stellar mean density, while ΔP is determined by the thermal and chemical stratification of the radiative, near-core regions. The latter can be used to e.g. infer the mass of the inert helium core on the red giant branch (RGB), and the detailed density/chemical composition profiles of the inner regions in core-He burning stars (e.g. Montalbán et al., 2013; Bossini et al., 2015; Constantino et al., 2015; Cunha et al., 2015). Moreover from the observational signature of the coupling between the pressure- and gravity-modes cavity, one can infer very local properties of the evanescent region between the two coupled propagation cavities (Takata, 2016; Mosser et al., 2017b).
- Deviations from the expected, approximated, frequency patterns of p and g modes can be used to detect and infer detailed properties of *localised gradients* in e.g. the chemical composition / thermal stratification / sound speed profile (referred to as “glitches”). Examples of such localised features include convective boundaries (regions of strong thermal and chemical gradients) and He-ionisation region where the adiabatic index, hence sound speed, has a local minimum, and whose signature can be used to infer the envelope He abundance (e.g. Vorontsov, 1988; Monteiro et al., 1994; Miglio et al., 2010; Mazumdar et al., 2014; Broomhall et al., 2014; Verma et al., 2019; McKeever et al., 2019).
- Determination of model independent properties through *seismic inversions* are possible with high-quality (i.e. *Kepler*-like) data. These approaches can be used to robustly infer global properties (e.g. mean density, see Reese et al., 2012; Buldgen et al., 2019), but also to build specific proxies (Roxburgh and Vorontsov, 2002; Buldgen et al., 2015b,a, 2018) of stellar

structure designed to test key ingredients of stellar physics related to, e.g., microscopic diffusion, convective boundary mixing (see Sec. 2.1).

- The signature of *activity cycles* can be found in frequency shifts / modes' lifetimes changes as a function of time (e.g., García et al., 2010; Kiefer et al., 2017; Santos et al., 2018; Salabert et al., 2018). In some cases a coarse latitudinal dependence of the magnetic field may be inferred from the angular dependence of the visibility of the rotationally split modes (Karoff et al., 2018; Thomas et al., 2019).

The precision and robustness with which one can achieve such detailed constraints on stars is strongly dependent on the length and quality of the photometric time-series available. The analysis requires, for instance, long-duration time series to obtain the requisite frequency resolution for extracting clear signatures of rotation in the oscillation spectrum. Examples of how the different duration of a dataset affects the seismic inferences is presented e.g. in Davies and Miglio (2016) and, extensively, in Mosser et al. (2019).

The mapping from the science questions outlined in Sec. 2 into observational tests and, finally, into a possible mission design (Sec. 3) will in fact be greatly aided by the knowledge acquired on the analysis and interpretation of existing seismic data.

1.2 Limitations of present/planned asteroseismic missions

In spite of being mainly planet-hunting missions, CoRoT (Baglin et al., 2009) and *Kepler* (Borucki et al., 2010) have marked a milestone in stellar astrophysics, allowing the inference of stellar properties (global parameters and internal structure) for a few thousands of stars, and opening the way to the synergy between asteroseismology and stellar population studies. Due to different telescope characteristics and observation strategies, the asteroseismic databases differ in spatial and frequency resolution, signal-to-noise level, and sampled regions of the sky. While *Kepler* observed during four years a 116 deg² field towards Cygnus-Lyra, CoRoT collected 3-6 months light-curves from more than ten 4 deg² fields in the Galactic disc in the Center and Anticenter directions. K2 (Howell et al., 2014) has observed 19 fields close to the Ecliptic, during 75 days each.

The *Kepler* and CoRoT databases comprise currently more than 20000 G-K red giants (and 600 dwarfs and subgiants) for which at least the seismic global parameters $\Delta\nu$ and ν_{\max} (and hence stellar mass and radius) have been measured (e.g. Yu et al., 2018; Mosser et al., 2010; de Assis Peralta et al., 2018; Chaplin et al., 2011; Davies et al., 2016; Michel et al., 2008; Anders et al., 2017). For a third of them, period spacing (ΔP) and coupling factor values have been also estimated (e.g. Vrad et al., 2016; Mosser et al., 2017a). Finally, the internal rotation profile has been derived for a few main-sequence pulsators (i.e. Benomar et al., 2015; Aerts, 2015) and larger samples of evolved stars (Beck et al., 2012; Deheuvels et al., 2012, 2014; Di Mauro et al., 2016; Huber et al., 2013; Mosser et al., 2012). CoRoT data (and K2), although

having lower frequency resolution and signal-to-noise than *Kepler* data, are good enough to derive $\Delta\nu$ and ν_{\max} as well as ΔP for some of the evolved stars (Mosser et al., 2011).

The quality of seismic data already collected sufficed to demonstrate their capability in inferring global and internal properties of stars, and in revealing hidden problems in current stellar modelling. However, to unambiguously answer some of the outstanding questions in stellar physics we need to disentangle the contribution from different quantities and uncertainties. A study of simpler populations, covering different stellar parameters would be necessary.

Kepler observations of NGC 6791 and NGC 6819 – two Galactic open clusters with ages of ~ 8 Gyr and ~ 2.4 Gyr, respectively – have shown an enormous potential clearly revealing, for instance, the presence of non-standard evolution processes (Handberg et al., 2017). However, there are only four stellar clusters in the *Kepler* database, with the two young and sparse clusters NGC 6811 and NGC 6866 providing less stringent constraints than NGC 6791 and NGC 6819. In addition, the clusters M67 and M4 have been targeted for 70 days with K2, but such data do not allow detailed asteroseismic studies of stellar structures (Stello et al., 2016; Miglio et al., 2016). Overall, we have *detailed* asteroseismology for just two clusters, covering a small region of the age–metallicity plane.

Moreover, the distribution of *Kepler* and CoRoT targets peaks at slightly sub-solar metallicity, and their selection functions are complex since they have been shaped by planetary science. Unfortunately, the subsequent mission, NASA’s TESS, although having the advantage of looking at much wider sections of the sky, has a much larger pixel size and shorter observation periods. It does not allow much more than global stellar properties to be obtained for the bulk of their targets (see e.g. Mosser et al., 2019; Davies and Miglio, 2016). The upcoming ESA PLATO mission, instead, will have much improved signal-to-noise and extended observations for its two continuing view zones, allowing measurements of individual frequencies and global parameters for more than 260000 targets (ESA, 2017). However, given the target selection aims mainly at finding nearby planetary systems characterizable through follow-up spectroscopy, PLATO detailed asteroseismic information will be available mainly for bright, nearby field F5-K7 dwarfs and subgiants. Moreover, the large pixel sizes of PLATO (projected size in sky: 15 arcsec, Samadi et al., 2019) will hamper obtaining detailed information for stars in crowded fields, including star clusters, the Milky Way’s bulge, and nearby galaxies.

2 Scientific objectives

Having in mind the inferences on stellar structure possible with constraints from solar-like oscillations, and taking into account the limitations of present/planned asteroseismic missions, we identify a series of open, key science questions that a mission dedicated to high-precision, high-cadence, long photometric time series to dense stellar fields will be able to address.

2.1 Science Objective 1: Unanswered questions in *stellar astrophysics*

Research in stellar astrophysics is now re flourishing, thanks to the availability of the long-sought combination of high-precision spectroscopic and astrometric data, and to the advent of exquisite asteroseismic constraints from space-borne telescopes. However, while *Kepler* data have started exposing the limitations of current models, robust and more stringent tests of stellar structure are currently limited by the composite nature of the stellar population observed, and by the scarce number of low-metallicity stars with asteroseismic constraints.

High-precision tests of stellar physics need to be performed in “controlled environments”, and clusters are the best stellar physics laboratories we have access to. Constraints from star counts, photospheric chemical abundances, and asteroseismology, together with strong priors on population-level parameters (e.g. distance, initial chemical composition, age) will enable percent-level tests of predictions by stellar evolution, as presented below.

Precise asteroseismic data for a set of open and, crucially, globular clusters will aid the theoretical development of a new generation of stellar models, which will benefit - in the next few decades - from an improved description of micro physics and 2D/3D numerical simulations of transport processes. Such developments will have direct impact on most areas of astrophysical research, from the interpretation of the photometry of resolved galaxies (see Tolstoy et al., 2009) and integrated spectra of high-redshift galaxies (see Conroy, 2013, and references therein), to the detailed characterisation of exoplanets and their environment (e.g. see Christensen-Dalsgaard and Aguirre, 2018).

Below we list open questions in stellar evolution which the proposed mission has the potential to address.

2.1.1 *Transport of chemical elements in the stellar interior*

As described in detail e.g. in the recent review by Salaris (2016), the predictive power of current stellar models is limited by the simplified treatment of (and interaction between) various processes that concur at determining how chemical elements are transported throughout the stellar interior.

Current shortcomings in the modelling of convection, atomic diffusion, mixing induced by thermohaline instabilities, gravity waves, and rotation affect the predicted evolutionary timescales, global stellar properties (e.g. surface temperatures, luminosities), internal structures, and surface abundances. These systematic uncertainties necessarily bias any model-based inference we make on e.g. stellar ages, also when seismic constraints are available.

Moreover, the use of stellar photospheric abundances in studies of galactic chemical evolution relies on our understanding of internal mixing, as surface abundances may have changed compared to the initial value due to nuclear processes, and brought to the surface by internal mixing. Our limited understanding of such processes also affects the robustness of spectroscopic proxies of stellar ages like $[C/N]$ (Masseron and Gilmore, 2015; Martig et al., 2016;

Mackereth et al., 2017; Casali et al., 2019), which is based on how the efficiency of transport processes in convective and radiative regions depend on the mass and metallicity (hence age) of red giant stars (Salaris and Cassisi, 2017; Lagarde et al., 2019).

The properties of the convective boundary mixing layer All stars with masses above approximately $1.1 M_{\odot}$ have convective cores during their MS, on the top of which a “convective boundary mixing layer” may develop. In classical 1D models of stellar structure, this layer is commonly known as the “overshooting region”. Importantly, the efficiency of convective mixing (or overshooting extent in 1D models) affects the relation between the initial mass and MS age for all stars more massive than $1.1 M_{\odot}$, or, alternatively, it affects all age determinations of stars younger than about 8 Gyr.

The best hopes to model the extent and structure of the convective mixing layer reside on modern 2D and 3D hydrodynamical models of stellar structure (e.g. Arnett et al., 2015; Baraffe et al., 2017), which require robust observational constraints from asteroseismology. While, e.g. in red giants, there are a few approaches to measure the efficiency of convective mixing with *Kepler* data alone (Montalbán et al., 2013; Bossini et al., 2017), they are complicated by a few factors, including e.g. the wide range of metallicities spanned by *Kepler* field stars, and its complex target selection. Detailed modelling of individual MS stars (e.g. Deheuvels et al., 2016; Michielsen et al., 2019) still has not produced unequivocal results. On the other hand, the distribution of gravity-modes period spacings in the *Kepler* clusters NGC 6819 and NGC 6791 seems to constrain quite well both the extent and the temperature gradient in the convective mixing layer of core-He burning stars (Bossini et al., 2017). The problem in this case is that present measurements regard just two clusters, and just the He-burning phase.

Indeed, *Kepler*-like data for open clusters could provide crucial information for a more definitive assessment of the impact of convective mixing layer on stellar evolution (and hence the “stellar clock”) as a function of stellar mass. Other advantages of extending the cluster sample, probably still not evident from present *Kepler* data, are: 1) Fast rotation is likely to be the main contributor to the increased cores observed in post-MS phases (Costa et al., 2019); fortunately, this effect can be assessed via the observation of rotational velocities in the star clusters themselves, via either accurate photometry or high-resolution spectroscopy of stars on the MS (e.g. Cordoni et al., 2018; Bastian et al., 2018). 2) In star clusters, the consequences of having larger/smaller cores can be directly observed also in the most advanced evolutionary phases – including for instance its possible impact on the luminosity of early-AGB stars (Bossini et al., 2015), and on white dwarf (WD) masses (i.e. the initial-to-final mass relation; Cummings et al., 2019). These can provide precious consistency checks for the overall efficiency of convective mixing and rotation, a step change compared to the loose constraints usually obtained from field stars.

Transport processes in radiative regions There is clear observational evidence for the occurrence of transport processes of chemical elements beyond convective instabilities. Several physical mechanisms have been proposed in the literature to explain such observations, including atomic diffusion and radiative levitation, rotation-induced mixing, mixing by internal gravity waves, thermohaline mixing, and the complex interaction between these processes (e.g. Maeder et al., 2013; Mathis, 2009).

For instance, helioseismology provides strong evidence that including atomic diffusion in the models reduces the tension with the seismically inferred sound speed profile (e.g. see the review by Christensen-Dalsgaard, 2002). Evidence for diffusion from spectroscopy in clusters has also been reported – e.g. in NGC 6397 (Korn et al., 2007; Husser et al., 2016), and from the APOGEE (Souto et al., 2018) and Gaia-ESO (e.g Smiljanic et al., 2016; Lagarde et al., 2019) surveys. Albeit this evidence, a quantitative understanding and calibration of the efficiency of such processes, is still lacking. Promising results on testing such effects on a star-by-star basis have been reported using high-quality asteroseismic data on main sequence stars (Deal et al., 2017; Verma et al., 2019) and by combining seismic constraints on masses and spectroscopic observations in two open clusters (Lagarde et al., 2015; Szigeti et al., 2018).

The way forward is to explore these effects in stars in clusters, in particular in those where we expect to see an effect of diffusion changing with position on the colour-magnitude diagram (CMD) such as, e.g., in Globular Clusters (GCs) and M67. Precise and accurate constraints on masses can be used to break the degeneracy between models with and without diffusion that affects the CMD (Salaris, 2016). Even more stringent constraints on the effect of diffusion on e.g. local chemical composition gradients, can be inferred for stars near and at the turnoff, providing an unprecedented calibration of the efficiency of transport processes. This is of paramount relevance if one wishes to improve the accuracy of age determinations for globular clusters (e.g. see Chaboyer et al. 1992), which are reference “time pieces” for the study of the early Universe (see also Sec. 2.2.1).

As mentioned earlier, one of the challenges that one has to face is to design observational tests that are specific to a given process. We know that for the case of rotation-induced transport processes, for instance, one would have to explain not only chemical composition profiles, but also the redistribution of angular momentum in stars. The latter can be probed by inferences on the rotational profile made using seismic constraints on large-scale velocity fields in the stellar interior, as illustrated below.

2.1.2 Core rotation and transport of angular momentum

The observations of solar-like oscillations in the Sun, in sun-like stars, and evolved stars have brought important constraints on their internal rotation rates (Thompson et al., 1996; Benomar et al., 2015; Beck et al., 2012; Deheuvels et al., 2012; Mosser et al., 2012). Comparisons between seismically inferred rotational profiles and models have demonstrated that an efficient transport

mechanism is needed in addition to the transport of angular momentum (AM) by meridional circulation and shear instability (e.g., see Eggenberger et al., 2012; Ceillier et al., 2013; Marques et al., 2013; Cantiello et al., 2014; Aerts et al., 2018).

Candidates for such an additional transport process include internal gravity waves and magnetic instabilities and their complex interplay with thermal and chemical gradient in the evolving stellar interior (Eggenberger et al., 2005; Cantiello et al., 2016; Pinçon et al., 2017; Eggenberger et al., 2019; Fuller et al., 2019). First-principles, numerical simulations of such processes performed under realistic stellar conditions are, however, particularly difficult to achieve, hence direct observational constraints on the internal rotation of stars are of prime importance to inform and test the modelling of angular momentum transport in stellar interiors. An improved understanding of AM transport processes is also key to understand the core rotation rates of compact objects (Kawaler, 2015; Heger et al., 2005).

To make further progress one needs more stringent and robust constraints on the evolution of AM transport, and to extend the constraints to the low-metallicity regime: this can be achieved by measuring rotational splittings of pulsation frequencies in stars in clusters. For instance, with accurate asteroseismic data e.g. on M67, we will have available direct observational evidence of how angular momentum is being redistributed along the evolution, from solar analogues through the MS turnoff, the subgiant phase and to the giant branches (see Sec. 3). Moreover, the detection of rotational splittings of mixed modes in stars in GCs will enable exploring metallicity effects on the transport of angular momentum, and testing models of AM transport in a regime which is closer to the one relevant for the first stars.

2.1.3 Mass loss on the red-giant branch

While mass loss close to the tip of the RGB has been suspected for long, direct measurements of the mass loss efficiency are characterised by uncomfortably large uncertainties. The best determination so far seems to be based on asteroseismology, more specifically on the measurement of the mass difference between RGB and red-clump (RC) stars in the clusters NGC 6819 and NGC 6791, observed by *Kepler*, and M67 using K2 data (Miglio et al., 2012; Handberg et al., 2017; Stello et al., 2016). These measurements however do not provide the dependencies of mass loss with key stellar parameters such as luminosity and metallicity, and do not even allow us to identify the basic mechanism responsible for the mass loss. Estimates of the mass loss in GC stars are also controversial, as demonstrated by the case of 47 Tuc: while its horizontal-branch (HB) morphology requires stars to have lost about $0.2 M_{\odot}$ on the RGB (e.g. Salaris et al., 2016), measurements both from spectra and infrared photometry do not give clearcut confirmation/disproval of mass loss (Origlia et al., 2010; Boyer et al., 2010), and have even been challenged by dynamical arguments (Parada et al., 2016).

On top of measuring the integrated RGB mass loss, oscillation frequencies can provide direct constraints on the mass of stars in the upper part of the RGB, where it is expected that the mass loss rates are the highest. The subclass of long-period variables known as OSARGs (Soszyński et al., 2004) are potentially very useful in this regard.

OSARGs are compatible with stars in the RGB phase, just below the tip, and also with the AGB phase (Tabur et al. 2010). Since their first detection it was suggested that they could be in fact solar-like oscillators (see also Dziembowski and Soszyński, 2010). Recent works by Takayama et al. (2013) and Mosser et al. (2013) clearly support that OSARGs are the high-luminosity counterpart of solar-like oscillators observed by *Kepler* and CoRoT at lower luminosity. Although with smaller number of detected modes, the oscillation spectra of OSARGs are compatible with the presence of radial and non-radial acoustic modes, just as the low-luminosity RGB stars. The low radial order of the modes appearing in OSARGs and the low number of modes do not allow to define quantities such as $\Delta\nu$ and ν_{\max} , however, as shown in Takayama et al. (2013), combining the information on the sequence at which this dominant mode belong with the luminosity, would allow to characterise the mass of the star. Given their large amplitudes, and periods of few tens of days, these oscillations can be easily detected in all the clusters considered here.

In addition to mass loss, variations in the helium content Y among the multiple stellar populations of GCs, are now believed to play a key role in determining the variety of observed HB morphologies. Therefore, a robust calibration of how much mass is lost on RGB stars at different metallicities, as allowed by asteroseismology, would contribute to open a clearer view into GC formation and evolution (Sect. 2.2.3).

Robust measurements of integrated stellar mass loss, would also be key to determine accurate ages for the huge numbers of field core-He burning stars already observed by *Kepler* and K2, and those that will be observed by PLATO: while their present masses are provided by asteroseismology, the *initial masses* are needed for age determinations (e.g. Miglio et al., 2013; Casagrande et al., 2016; Anders et al., 2017). Better estimates of mass loss on the RGB would also impact the initial conditions assumed for evolutionary models of AGB and post-AGB stars, used to characterise extra-galactic stellar populations.

2.1.4 Occurrence of mergers / products of binary evolution

Thanks to constraints on stellar mass from asteroseismology in the open clusters observed by *Kepler*/K2, there is now clear evidence for over- and under-massive stars (Brogaard et al., 2015, 2018; Stello et al., 2016; Handberg et al., 2017) which are the likely products of mass exchange with a companion. There is also likely evidence of such products among field stars that are enriched in α elements, hence supposedly belonging to an old population (Martig et al., 2015; Chiappini et al., 2015; Jofré et al., 2016; Yong et al., 2016; Izzard et al., 2018; Hekker and Johnson, 2019).

The direct measurement of stellar masses could provide a better understanding of the observational abundance of blue stragglers and other types of rejuvenated stars, i.e. stars that previously accreted material from a companion star through a stable mass transfer event or a merger. An extension of the asteroseismic sample to a wide range of cluster ages and metallicities would provide invaluable constraints to the study of tidal interactions (e.g. Beck et al., 2018), and to models of binary populations, e.g. event rates of binary interaction, initial-mass-ratio/period distributions (e.g., see Moe and Di Stefano, 2017) and constraints on key parameters in the modelling of interacting binaries.

For instance, robust inferences on the number of peculiar-mass stars will enable constraints on the α parameter that determines the efficiency of the transfer of orbital energy of the inspiraling cores to the envelope during the common-envelope phase of evolution (e.g., see Izzard et al., 2018). This parameter is the key to many processes in stellar astrophysics e.g. BHBH/BHNS/NSNS/WDWD mergers (hence sources of gravitational waves), type Ia supernovae, cataclysmic variables, FK Comae stars, X-ray binaries, post-(A)GB stars, and helium white dwarfs.

Moreover, testing the dependency with metallicity is also crucial. There are indications that binary interactions are more common at low metallicities (e.g., see Badenes et al., 2018). Do we see that? Are the structures of the rejuvenated stars different at different metallicities?

On top of characterising evolved stars that are products of mass transfer, blue stragglers near the turnoff of old-open clusters will also be accessible to a precise asteroseismic inference, and will contribute to understanding e.g. how angular momentum is redistributed, so that it prevents the accreting stars to quickly spin up to critical rotation (see also Sec. 2.1.2).

2.2 Science Objective 2: Unanswered questions in *cluster formation and evolution*

2.2.1 *Globular clusters formation from absolute ages*

Most GCs form in the first few Gyrs of the evolution of their host galaxy and they therefore contain valuable information about the earliest phases of galaxy formation. Because the stars in a GC are believed to have the same age, precise stellar (core) masses allow absolute age dating of GCs, with an age resolution that is sufficient to resolve the age differences among GCs. Combined with their metallicities, absolute ages allow us to distinguish accreted from in-situ formed GCs (Forbes and Bridges, 2010), derive the accretion history of the Milky Way (Leaman et al., 2013; Massari et al., 2019) and discover individual ancient ($\gtrsim 10$ Gyr) accretion events (e.g. Myeong et al., 2018, 2019). Accurate ages also serve as a valuable benchmark for other age dating methods, such as isochrone fitting and age dating from the white dwarf cooling sequence.

2.2.2 Thresholds for the onset of multiple populations

Another unexpected trend is that only clusters older than 2 Gyr and massive enough, with a threshold close to $10^5 M_{\odot}$ (e.g. Martocchia et al. 2019), appear to host multiple populations (i.e. with the chemical signature) in the Magellanic Clouds. In the Milky Way only the really old and massive clusters, i.e. those classically called GC, show the presence of multiple populations (see Bragaglia et al. 2017 for a detailed list). Apparently, no open cluster, even among the oldest (e.g. NGC 6791, Berkeley 39), was massive enough at birth and/or formed in an environment favourable to the development of multiple populations. While these trends need to be confirmed by larger samples, the availability of asteroseismic masses/rotation for a few clusters will provide new light on this subject.

2.2.3 Measuring helium content in GCs with asteroseismology

At present, the surface helium content Y can be determined for giants in a GC from photometry using a combination of filters comprising the UV bands (essentially, only with HST, see e.g. Milone et al. 2018), or from the luminosity of the RGB bump (e.g. Bragaglia et al., 2010; Lagioia et al., 2018). For HB stars, it can be derived modelling the extension of the HB, or directly through spectroscopy in a limited temperature range (e.g. Villanova et al., 2012; Marino et al., 2014). In red giants, the only accessible He line is chromospheric, and its measurement is complicated (e.g. Pasquini et al., 2011; Dupree and Avrett, 2013). A further method to determine Y for massive samples of giant stars in GC would then be very important.

As already mentioned in Sect. 2.1.3, a range of initial He abundances is mandatory to reproduce the HB morphology of old and intermediate-age stellar clusters and, in turn, to constrain the efficiency of mass-loss on the RGB (e.g., Salaris et al., 2006, 2016; Chantreau et al., 2019; Tailo et al., 2019). To determine the mass of stars (through asteroseismology) is another indirect way to measure the He content since there is an anti-correlation between the initial He abundance and the initial mass of stars (e.g., Salaris et al., 2006; Chantreau et al., 2015). In addition, it is imperative to know the precise initial He abundance variation (and distribution) to greatly constrain scenarios trying to explain the origin of multiple populations. However, present determinations of the He content are still subject to large uncertainties and vary as a function of the method used. For instance, the determined He spread between Salaris et al. (2016) and Lagioia et al. (2018) in 47 Tuc differ by a factor 3, making any progress in the multiple populations field difficult.

Therefore, *direct* asteroseismic constraints to Y in cluster stars would be invaluable. In open clusters of different metallicity, they would also provide an empirical recipe for the He-enrichment law in the Galaxy.

2.2.4 Globular clusters masses and dynamics

Precise masses of giants and an assessment of (photometric) binarity are valuable input for dynamical models of GCs. GCs are collisional systems and two-body relaxation evolves the system towards equipartition and ‘mass segregation’, such that the more massive stars and stellar remnants are more centrally concentrated. Dynamical mass models of GCs that include an equipartition description (e.g. Da Costa and Freeman, 1976; Gunn and Griffin, 1979; Gieles and Zocchi, 2015) can be used to infer the underlying GC mass profile from the number density profile. For this, precise stellar masses help to ‘anchor’ the models with multiple mass components. In addition, mass estimates at the different evolutionary stages can be combined with dynamical models to estimate at which moments the mass is lost (Heyl et al., 2015; Parada et al., 2016). The properties of binaries are the result of the primordial binary properties and the dynamical evolution. For example, the inverse trend of the binary fraction with GC mass (e.g. Milone et al., 2012) is likely the result of binary destruction in massive GCs. Photometric binary properties, complemented by the wealth of multi-epoch radial velocity surveys with Integral Field Units (IFUs) (e.g. Kamann et al., 2018), can be used to derive a full census of binaries in GCs and can be included in dynamical modelling exercises of GCs. Finally, it allows us to understand the interplay between binaries and the dynamical evolution of GCs and leads indirectly to insight in the formation properties (e.g. density and mass) of GCs.

2.2.5 Black holes

Based on an extrapolation of the ‘ $M - \sigma$ relation’ (Gebhardt et al., 2000; Ferrarese and Merritt, 2000), GCs may host the sought-after intermediate-mass black holes (IMBHs, $\sim 10^{3-5} M_{\odot}$), filling the gap between stellar-mass BHs and supermassive BHs. The existence of IMBHs is highly debated, and from radio and X-ray observations, stringent upper limits of $\sim 10^3 M_{\odot}$ were found for several Milky Way GCs (Tremou et al., 2018). However, GCs are gas free and the absence of an accretion signal may not yet be evidence for the absence of IMBHs. Various claims have been made for an IMBH in ω Cen (e.g. Noyola et al., 2008; Baumgardt, 2017), 47 Tuc (Kızıltan et al., 2017), NGC 6397 (Kamann et al., 2016) and M54 (Ibata et al., 2009). These findings are based on kinematics of stars in the centre of the cluster, and dynamical mass modelling. A complication with this method is that the IMBH signal is (partially) degenerate with other – more plausible – effects, such as radial orbit anisotropy (e.g. Zocchi et al., 2017) and a population of stellar-mass BHs (Zocchi et al., 2019; Lützgendorf et al., 2013). An alternative approach is to look for gravitational-lensing effects (Kains et al., 2018). In the search for IMBHs, various stellar-mass BHs have been reported in several GCs (e.g. Strader et al., 2012), including 47 Tuc and ω Cen. High-precision and high-cadence photometry could be used for micro-lensing events and light-curve modelling of BHs with stellar companions (see Giesers et al., 2018, for an ex-

ample in NGC 3201) to more firmly establish the presence of stellar-mass BHs in GCs, which impacts on our understanding of the role of GCs as gravitational wave factories.

2.2.6 Do stars in clusters rotate faster than field stars?

One of the most surprising results of high-precision HST photometry of Magellanic Cloud clusters regards the presence of a large fraction of main sequence stars rotating close to their critical break-up velocities, Ω_c . For instance, in the ~ 200 Myr LMC cluster NGC 1866, about 2/3 of the MS stars appear to be rotating at $\sim 0.9 \Omega_c$ (Milone et al., 2017). Moreover, extended main sequence turn-offs (MSTOs) are an ordinary feature in intermediate-age star clusters (Milone et al., 2009; Cordoni et al., 2018) and probably largely due to the presence of fast rotators (Bastian and de Mink, 2009; Bastian et al., 2018). Although the distributions of rotational velocities in field stars (Royer et al., 2007) is more difficult to interpret, projected rotational velocities seem to indicate that young clusters have more fast rotators than their neighbouring fields (see Huang and Gies, 2006; Strom et al., 2005; Dufton et al., 2006). Therefore, the question arises whether this is a general trend, applying to all stars born in clusters. Some authors (Kamann et al., 2019; Corsaro et al., 2017, see also Sec. 2.2.7) also suggest that cluster stars largely inherit the initial angular momentum of the molecular cloud from which they form, hence spinning faster and at preferential orientations as compared to the field. These suggestions could be verified with more data on rotational velocities added to the asteroseismic information (rotational velocities and spin orientations), in and around a few star clusters.

Given that stellar evolution models are usually calibrated with observations of star clusters, and later applied to all galaxy components irrespective of their density, this question may be very important in the context of stellar evolution and stellar population synthesis of galaxies. A very basic assumption of these models, namely that stars in clusters form and evolve in the same way as stars in the field, may be wrong.

Faster rotation can also arise from more mergers among young stars in a dense cluster environment, which may lead to more rotating (proto)stars. This might be worth investigating with the additional constraints (including stellar masses and core rotation rates) provided by asteroseismology.

2.2.7 Spin alignment in clusters

Recent results have suggested that the alignment of stellar rotation axes in clusters is not isotropically distributed, but rather has a preferential inclination angle (Corsaro et al., 2017). This result is based on asteroseismology analysis of *Kepler* data for stars in NGC 6791 and NGC 6819. The generally advocated explanation for the alignment of stellar rotation axes in clusters is that the stars were born within a giant molecular cloud that contained significant amounts of angular momentum, which was inherited by the stars as

they formed (e.g. Corsaro et al., 2017; Rey-Raposo and Read, 2018). While some caution needs to be taken in interpreting these results, due to potential systematic effects in inferring the stellar inclination angle from *Kepler* analyses (Kamiaka et al., 2018; Mosser et al., 2018; Kuzlewicz et al., 2019), the results potentially offer an unexpected and intriguing insight into the formation of stellar clusters. Looking at these two clusters, Kamann et al. (2019) found that the cluster NGC 6791 rotated with an axis that was consistent with the spin alignment of the stars, suggesting that angular momentum has transferred from the large scale (cloud/cluster) to stars during the early phases of formation. Expanding this type of study to a statistically meaningful sample would be a major step forward.

2.3 Science Objective 3: Assembly history and chemodynamics of the Milky Way and dwarf galaxies

One of our main goals is to improve the quantity and quality of asteroseismic constraints at low metallicities, with new GC observations. This alone would produce an improvement in stellar models at low metallicity (see e.g. Sec. 2.1.1), possibly impacting the interpretation of data for low-metallicity dwarf galaxies, and the resolved halos of more massive galaxies, presently observed out to the Virgo cluster. Even more certain would be the impact coming from the *direct* observation of a few galaxy components, such as:

2.3.1 ω Cen

Rather than a normal globular cluster, ω Cen is probably the heart of a dwarf galaxy whose periphery has been dispersed by the Milky Way. It contains multiple stellar populations, with a large range of metallicities (i.e. heavy element content) that betray a formation over an extended period of time. Interestingly, ω Cen is close enough for asteroseismic observations to reveal accurate masses of its RGB and red-HB stars (with long time series allowing even the detection of internal rotation; see Fig. 2). Given the available HST multi-band photometry and spectroscopy (e.g. Hilker et al., 2004; Piotto et al., 2005; Villanova et al., 2014), the accurate seismic masses would allow a much more complete picture of the star formation and chemical evolution history in this unique object (Romano et al., 2007).

2.3.2 *The Sagittarius Dwarf Spheroidal and M54*

Among the many dwarf spheroidal galaxies, the Sgr dSph is both one of the closest, and more interesting ones. It contains a nuclear star cluster at its core, M54, which is expected to evolve similarly to ω Cen (Carretta et al., 2010). Its distance (25 kpc away) represents the main difficulty for asteroseismic observations; however, our simulations (Sec. 3.2) indicate that solar-like oscillations

are detectable at the level of the RC and above, provided a mirror with diameter $\gtrsim 90\text{cm}$ and at least 6 months of observations. These observations could represent the first asteroseismic measurements for an external galaxy. They would allow a determination of the Sgr dSph star formation history, which then may be used to infer possible age gradients and to date when Sgr dSph entered within the sphere of influence of our Galaxy.

2.3.3 *Magellanic Clouds*

The LMC and SMC galaxies have always been fundamental testbeds for the theories of stellar evolution and stellar populations. One could conceivably detect solar-like oscillations in the upper RGB, provided a large enough mirror size could be considered (see Sec. 3.2). In the context of the proposed mission, one could run a pilot asteroseismic study of the Magellanic Clouds. For instance, this is easily done in the case of the SMC, which lays in the background of two populous GCs, namely 47 Tuc and NGC 362. This would, at the very least, enable to extend studies based on large-amplitude pulsators such as OSARGs (Soszyński et al. 2007, see also Sec. 2.1.3) to stars with lower luminosities and lower pulsation amplitudes.

2.3.4 *The Milky Way bulge*

The Milky Way bulge still holds many unanswered questions despite the modern data (see Barbuy et al., 2018, for a recent review): not only is it a region where several galactic components co-exist (spheroidal, bar, innermost parts of the thick and thin disc, and residuals from merger accretions), but it is also not easily reachable given the large extinctions, large distances involved as well as crowding. Regarding its spatial structure and kinematics, added to the general idea of it being a box/peanut bulge, there are still many uncertainties regarding, for instance, the fraction of bulge which is actually in a bar, the different spatial distributions of metal-rich and metal-poor components, and the nature of its X-shaped feature (see e.g. Zoccali et al., 2017). Regarding its stellar populations, its age distribution is still a matter of debate, with some observations suggesting that the Milky Way bulge is not simply “old” but contains a significant fraction of intermediate-age populations (e.g. Saha et al., 2019), whose fractions are far from being established. For instance, while the precise HST photometry from Brown et al. (2010) indicates only old stars, the spectroscopic masses of microlensing dwarfs from Bensby et al. (2017) reveal a $\sim 15\%$ fraction of young stars.

While large spectroscopic surveys are providing excellent kinematics and chemical abundances, the weak point of present bulge studies resides in the lack of accurate distances and ages. For distances, even by combining Gaia DR2 and APOGEE DR16, the Bayesian spectrophotometric code StarHorse (Anders et al. 2019, Queiroz et al. 2020) still has distance uncertainties of the order of 10% in that region – therefore not only insufficient to disentangle the co-existing stellar populations in the bulge region, but also to get large enough

samples of stars. The principal difficulties stem from the high extinction values, its unknown distribution along the line-of-sight, and the position-dependent non-standard extinction curves (e.g Nataf et al., 2016; Saha et al., 2019). Such problems will persist even with Gaia parallaxes, which will not be able to peer through the dust to reveal the real stellar distances over most of the bulge. An even more incomplete situation is the one for age estimates in the bulge region. In this specific front, asteroseismology can be of great help, by providing accurate masses and radii of bulge red giants, and hence their ages and distances. Combined with Gaia’s proper motion data, these data enable us to map the age of stars depending on the orbits. Stars in the bar structures, such as the long bar and the shot X-shaped feature, are in the trapped orbit, and the age distribution of stars in the orbit of each feature tells us the formation epoch of each component of the bar structure. Recently, Bovy et al. (2019) made the first attempt to age date the stars in the Galactic bar and claim that the bar is dominated by old stars, and therefore formed at the early epoch. However, this is relying on spectroscopic distances and ages, which are calibrated with the currently limited asteroseismic data, and rely on current stellar model tracks. Direct and precise seismically inferred ages and distances of the bulge stars will unambiguously answer the formation epoch of the bar, even the difference of the formation epochs of the long bar and the X-shaped feature, which is considered to form by “buckling” that occurred more recently.

Of course, asteroseismic observations of the bulge should also meet some basic requirements, such as being sensitive to redder wavelengths and avoiding too-crowded regions. A careful field selection will be needed in this case, and will be aided by combining Gaia data with ongoing/near-future spectroscopic surveys (such as 4MOST, APOGEE South, and MOONS). The observations would be complementary and have a clear synergy with the recently selected Japanese-led SMALL-JASMINE mission (planned launch in mid 2020s, <http://jasmine.nao.ac.jp/index-en.html>), which will provide near-infrared astrometry and time-series photometry data in the Galactic nucleus region (within $\sim 0.7^\circ$).

2.4 Additional/complementary science from high-cadence, high-precision long photometric monitoring

2.4.1 Solar twins and “solar evolution” from M67 observations

With its solar metallicity and ~ 4 Gyr age, the star cluster M67 can provide extraordinary insights on the evolution of the Sun. It contains a handful of “solar twins”, stars whose spectra are very close to the Sun’s (Pasquini et al., 2008) and which indicate a nearly-identical chemical composition (Liu et al., 2016). Asteroseismic analysis of these stars is granted to provide an interesting comparison with helioseismology. In addition, M67 could offer additional, population-level constraints on the evolution of rotation, activity, and diffu-

sion, in stars which were initially similar to the Sun. As discussed in Sec. 3, M67 is close enough to allow detailed asteroseismic analysis of MS stars in addition to its numerous red giants. Given its position close to the Ecliptic, PLATO observations of M67 (which are still to be defined) will be limited to short time series, and to the cluster’s outskirts.

2.4.2 Exoplanets in clusters / bulge

Characterising planetary systems in star clusters is crucially important to constrain our theories on planetary formation and subsequent migration. Clusters provide us with a homogeneous environment where fundamental parameters such as chemical abundances and age can be reliably measured, in a much better way than how it is done on field stars, putting the detected planets in an accurate evolutionary context. While there is a growing number of exoplanet surveys focusing on low mass stars and binary systems, only very few attempts were, and are being made to explore alternate Galactic environments. So far most exoplanetary systems have been detected orbiting single, Sun-like stars, with only a handful of planets detected in open clusters by *Kepler*, K2 and TESS (Fujii and Hori, 2019). Quite surprisingly, the occurrence of planets within open clusters (OCs) has been suggested to be similar or even slightly higher than for field stars, although the small sample size severely limits the statistical significance of the results (Meibom et al., 2013; Brucalassi et al., 2016, the latter found a $5.6^{+5.4}_{-2.6}\%$ frequency of hot-Jupiter planets around solar-metallicity stars). Unfortunately: 1) only four OCs were located within the *Kepler* field, all of them too distant for a planet search to be fully effective on main sequence stars; 2) K2 and TESS were/are not particularly helpful in enlarging the sample, given their short baseline (~ 70 and 27 days, respectively) that limits the discovery space mostly to hot planets; 3) PLATO will be limited by its large pixel scale ($\sim 15''$, only slightly smaller than TESS), which implies that confusion will badly impact its photometric performances on crowded fields, also boosting the occurrence of false-positive detections. Confirming (or disproving) a higher rate of hot Jupiters in OCs would favour (or reject) the *planet-planet scattering* scenario, which is one of the competing theories to explain the origin of hot Jupiters (Dawson and Johnson, 2018). On the other hand, evidence for small (rocky) planets in OCs is still fragmentary and to be explored.

The situation is even less clear when it comes to globular clusters. For instance, HST observations of 47 Tuc revealed no transiting hot Jupiter (Gilliland et al., 2000) and also subsequent efforts on ω Cen and NGC 6397 were unfruitful (Weldrake et al., 2008; Nascimbeni et al., 2012). The interpretation of these results is difficult. It remains unclear whether this absence is caused by the low metallicity of the cluster, by the intense density of stellar systems (which could either reduce disc and therefore planet formation, or could lead to system-wide instabilities post-formation), or both. It is also debated whether the null result on 47 Tuc is actually statistically significant (Masuda and Winn, 2017).

Microlensing surveys sometimes detect exoplanetary systems into the bulge (Zhu et al., 2017), but those planets usually orbit fairly far from their host, preventing any direct statistical comparison to transit and Doppler surveys. Altogether, this calls for additional, more detailed investigations. In other words, we need a new mission to constrain the Initial Radius/Mass Function for exoplanets in different (and largely unexplored) environments.

A new *asteroseismic* mission, conducting photometric observations as precise as *Kepler* in the Milky Way’s bulge, and clusters of various ages and metallicities, would beautifully fulfil this role, by being able to detect transiting planets within these multiple stellar environments. It would allow us to verify whether the statistics of planet detection match one another between each of these environments, and whether they match the detections of *Kepler*, *TESS* and of all the ground-based results. It would allow us to seek similarities and differences caused 1) by environment (high stellar densities might lead to more frequently unstable systems, decreasing how packed planetary systems are found (e.g. Malmberg et al., 2011), 2) by metallicity (confirming how it correlates to the presence of gas-giants and of super-Earths (e.g. Sousa et al., 2019), 3) by metal abundances (asking whether α -enhanced stars can produce a different planet population), 4) by age (since we expect planet populations to change with time (Pu and Wu, 2015), and 5) by history (by tracking systems that have migrated within the Milky Way, moving from one environment to another). The length of each stare will determine up to which orbital period transiting planets can be detected. Microlensing might also be attempted. With enough transits, masses for exoplanets can be determined from transit-timing variations. Foreground white dwarfs, and very late M dwarfs might fall within each field, with the potential to discover Earth-sized planets. Particularly interesting would be the possibility of discovering “Solar System twins” orbiting the M67 solar twins.

2.4.3 Activity / flares

It has long been established that solar p-mode frequencies respond to the Sun’s changing levels of magnetic activity and hence provide a probe of the underlying physical changes driving these variations, i.e. stellar dynamos. Seismic detections of stellar-cycle-related variability are now being extended to other solar-like oscillators thanks to precise, long-timebase photometric data from CoRoT and *Kepler* (see Sec. 1.1). However, these detections are in field main-sequence stars. Here, there is a great scientific opportunity in extending studies to counterparts in markedly different stellar environments. The additional constraints provided by common cluster membership opens the exciting possibility to probe the action of stellar dynamos as a function of stellar mass.

Stellar activity is likely to have a large influence on the habitability of exoplanets. Therefore in order to understand the chances for finding life in the Universe, we need to understand the nature and evolution of activity of Sun-like stars. Over the last decades we have learned that cycles like the 11-year sunspot cycle are common in Sun-like stars (Baliunas et al., 1995),

that young stars tend to have enhanced levels of UV and X-ray emission (Güdel, 2007) and that some stars, including Sun-like stars, host devastating superflares (Maehara et al., 2012).

The proposed mission will allow us to study the photometric effect of stellar activity in a large number of stars with well constrained parameters. This will be especially important for the studies of superflares as it is still an open question if the Sun would be capable of hosting superflares (Karoff et al., 2016; Notsu et al., 2019). In order to answer this question we need to either detect or rule out superflares on truly Sun-like stars with respect to mass, age, rotation, and chemical composition. Here focusing on stars in stellar clusters will be a great advance.

Finally, the high-precision, high-cadence photometric monitoring of stars along the HB in GCs, will also shed light on the nature of the variability of blue horizontal branch (BHB) stars, and its relation to chromospheric activity / chemical peculiarities (see e.g. Paunzen et al., 2019; Montenegro et al., 2019).

2.4.4 Additional science from stellar pulsations

The driving of pulsation modes occurs typically near the surface, and is therefore mainly a function of the effective temperature. The part of the HR diagram going from the classical instability strip to the red-giant branch (main targets of the proposed mission) corresponds to stars with a convective envelope. By one way or another, convective motions and ionization zones always drive pulsations. This is clearly seen on the main sequence: going from blue to red, we encounter δ Sct stars (HeII ionization zone driving), γ Dor stars (driving at the base of the convective envelope, see e.g. Dupret et al., 2005) and solar-like oscillators (stochastic driving by convective motions). Going up to post-main sequence stars we observe on one side RR Lyrae and Cepheids and on the other side solar-like oscillations of red giants. There is no obvious reason for the absence of pulsation in-between. The only obstacle to non-radial pulsations in evolved stars is radiative damping in the core. However, partial reflections in the evanescent zone usually leads to the trapping of some non-radial modes in the envelope, making their detection possible. As long as the core-envelope density contrast is not too high, mixed modes probing deeper layers are also predicted (Dupret et al., 2009) and observed (Bedding et al., 2011) in red giants, increasing considerably their seismic potential: for instance, probing of the core rotation and density. Note also that the instability strips associated with different types of mode driving processes always intersect, producing hybrid stars with even higher seismic potential. We expect mixed modes and hybrid stars to be also present in hotter stars, making their very detailed seismic characterization possible for the first time.

2.4.5 Additional targets in the field

Asteroseismic surveys carried out in the 2020–2030 era will benefit from extensive new data from wide-area surveys such as Gaia, eROSITA, the Rubin

Observatory (previously known as the LSST), SDSS-V, which will provide a wide variety of new targets for more detailed time-series, high-cadence photometry, both within and outside star clusters. It is hard to foresee all the possibilities here, but we might expect:

- better characterization of classical pulsators, eclipsing binaries, and close binary systems uncovered by “low-cadence surveys” such as Gaia and the Rubin Observatory;
- improved statistics of FK Com stars, which are rapidly rotating giants probably resulting from recent stellar mergers. *Kepler* allowed direct measurements of rotation periods for a handful of them (Costa et al., 2015) and confirmed their association with bright X-ray sources with high levels of chromospheric activity and flaring behaviour (Howell et al., 2016), hence narrowly opening the way to link models (of individual objects or of populations) to their observations.
- follow-up lightcurves of unusual transients, such as those found by *Kepler* and which still remain without satisfactory explanations (e.g. KIC 8462852; Boyajian et al., 2016).

3 Translating top-level science objectives into target selection and a possible mission profile.

Given the main scientific objectives outlined in Sec. 2, we discuss here the targets needed to achieve our goals and give indications about the key factors that will concur to define the mission profile.

3.1 Targets

We make a preliminary selection of suitable targets considering proximity and diversity as the main selection criteria. Table 1 gives an example of a target list, which includes three globular clusters (with different chemical compositions, morphologies of the HB, and properties of multiple populations), ω Cen (likely the nucleus of a satellite galaxy, with extended star formation history), an old open cluster (M67), and a field with a significant bulge population. We have also listed the Sgr dwarf spheroidal galaxy among the targets worthy of consideration.

3.2 Seismic performance

As presented in Mosser et al. (2019), the plethora of data and analyses performed on photometric time series from several space-borne telescopes allows us to define and quantify a robust seismic performance indicator. For the targets in Table 1, we adopt the method described in detail by Mosser et al. (2019) to quantify the performance expected when varying the mirror size and

Table 1 Approximate characteristics of the targets for which we assess the seismic performance in Sec. 3.2. This list is indicative only, alternative options are available.

Object name	d [kpc]	$m_{V, RC/ HB}$	[Fe/H]	Age [Gyr]	ang. size
NGC 104 (47 Tuc)	4	14	-0.8	13	31'
NGC 6121 (M4)	2.2	13	-1.5	12.2	26'
NGC 6397	2.4	BHB	-1.8	13.4	32'
NGC 2682 (M67)	0.9	10.5	0.0	4.5	30'
NGC 5139 (ω Cen)	4.9	14	broad	broad	36'
Milky Way's bulge	6–10	$I > 14$	broad	broad	wide
Sgr dSph	25	18–20	broad	broad	450' \times 216'




Fig. 2 Hertzsprung-Russell diagram showing stellar populations representative of ω Cen (*left panel*) and M67 (*right panel*). Each star in the population is coloured according to the seismic information that can be extracted: blue, solar-like oscillations are detectable; yellow, information on the gravity-mode period spacing can also be inferred; and green, rotationally split pulsation modes can be measured, hence information on the internal rotational profile can also be inferred. We assume a mirror diameter of 90cm and a duration of the observations of 2 years (ω Cen) and 1 year (M67).

duration of the observations. The latter are the two key parameters impacting on the seismic yields, provided e.g. that the pixel size of the detectors on the sky is such as to mitigate the effect of contamination from nearby sources¹ and a cadence appropriate for the stars of interest is chosen. The noise model, as described in Mosser et al. (2019), includes a systematic component originating from e.g. the sky background, the jitter, or the readout noise, and it is assumed to be similar to *Kepler*'s. We assume a duty cycle of 95% and consider a broad, white-light photometric filter.

For the targets listed in Table 1 we quantify our ability not only to detect solar-like oscillations, but also to measure the detailed properties of mixed

¹ Depending on the specific choice of the target, a pixel size of the order of 1" or less is needed, compared to 4" of *Kepler*, 20" of TESS and 15" of PLATO




Fig. 3 Same as Fig. 2, this time for one cluster (M4) but changing the diameter of the telescope (*left panel*) or the duration of the observations (*right panel*).

modes, and to resolve rotationally split frequencies, which are needed e.g. to achieve the science objectives presented in Sec. 2.1. To simulate the stellar population in the proposed fields, we use TRILEGAL (Girardi et al., 2005) with literature-based estimates for the age, chemical composition, distance, and extinction for each target. Since the seismic performance indicator is largely determined by global stellar properties and the targets’ distance, the dependence on the details of the stellar models used to simulate the populations is not a source of concern.

Examples of the detection thresholds obtained with a different combination of telescope size and duration of the observations, for some of the key targets in Table 1, are illustrated in Figs. 2 and 3. We present three cases: a telescope with a diameter of 30cm, 90cm, and 150cm and vary the duration of the observations from 1 month to 2 years. Since our aim is not only to infer global stellar properties (e.g. mass) but, at least for some of the targets, to perform high-precision tests of stellar structure models, we argue that a collecting surface corresponding to a ~ 90 cm diameter and a combination of 24-, 12-, and 6-months long campaigns for the candidate targets in Table 1 would enable reaching the key objectives presented in Sec. 2. For instance, if one aims at a precise and accurate characterisation of the rotational profile and to infer detailed, local properties of stellar interiors using the information from individual mixed p-g modes, observations longer than ~ 1 yr are needed (e.g. Scientific Objective 1, Sec. 2.1). On the other hand, 6-months long observations are enough in most cases to enable robust and precise inferences on global stellar properties (e.g. masses, ages) to a level which would enable breakthroughs in studies of stellar populations (Scientific Objective 3, Sec. 2.3).

Larger mirrors would be needed if one wishes to fully characterise giants at and below the luminosity of the RC in satellite galaxies, e.g. in the LMC ($\mu \simeq 18.5$). A small-size mirror (e.g. 30cm) would still enable detection and characterisation of stars in the closest globular clusters, yet limit the high-

precision tests of stellar physics, especially at the turnoff of globular clusters, and on the MS of open clusters.

3.3 Preliminary considerations on telescope design

As evinced from the typical angular size of our prime targets (see Table 1), an appropriate diameter of the field of view is $\lesssim 1$ deg. A suitable field for the bulge will be chosen based on detailed studies enabled by Gaia and in synergy with SMALL-JASMINE (Sec. 2.3.4). Similarly, for the Sgr dSphe, a suitable sub-field can be identified by a detailed study of its stellar content and foreground contamination.

As discussed in Sec. 3.2, a 90cm aperture telescope would allow us to reach our main goals in terms of seismic performance. Assuming pixel sizes of about $5\mu\text{m}$, the telescope focal length will depend on the adopted pixel scale. Pixel scales of the order of 0.25 arcsec/px (to be compared to the 4 arcsec/px of *Kepler*) are certainly well suited to separate stars in open clusters and in the outskirts of globular clusters and the bulge. They would imply a focal ratio of about 5, hence a reasonably-compact telescope design. However, detailed simulations (using available images of our targets) are necessary to assess the optimal pixel scale for all targets, considering all the other constraints on telescope dimensions and weight.

4 Summary

In the last decade, the *Kepler* and CoRoT space-photometry missions have demonstrated the potential of asteroseismology as a novel, versatile, and powerful tool to perform exquisite tests of stellar physics, and to enable precise and accurate characterisations of stellar properties, with impact on both exoplanetary and galactic astrophysics.

Based on our improved understanding of the strengths and limitations of such a tool, we argue for a new small/medium space mission dedicated to gathering high-precision, high-cadence, long photometric series in dense stellar fields. These crowded fields are being avoided in all present and planned exoplanet searches, which instead favour large pixel sizes and nearby bright stars.

The proposed science case / mission will lead to breakthroughs in stellar astrophysics, especially in the metal poor regime, by performing high-precision tests of stellar physics in open and, crucially, globular clusters (see Sec. 2.1). Moreover, precise and accurate asteroseismic constraints on global stellar properties, together with complementary astrometric and spectroscopic data, will address unanswered, long-standing questions about the evolution and formation of open and globular clusters (see Sec. 2.2), and aid our understanding of the assembly history and chemical evolution of the Milky Way's bulge and few nearby dwarf galaxies (see Sec. 2.3).

We use the knowledge acquired from the analysis and interpretation of existing seismic data to propose a possible mission design, including a list of potential targets. The details of what can be achieved depend on the telescope collecting area, on the length and cadence of observations, and the distance of the selected targets. With a few reasonable assumptions, we identify the following priorities, which we show can be reached with a ~ 90 cm-wide telescope with a $\sim 1^\circ$ field-of-view, and a combination of 24-, 12-, and 6-months long observing runs on the targets identified in Table 1:

1. to detect solar-like oscillations in stars with luminosities lower than the red clump, enabling precise and accurate characterisation of global properties (masses, radii, distances, ages) for stars belonging to a few globular clusters of varying metallicity, open clusters of varying age, and building blocks of the Milky Way;
2. in globular clusters, calibration of poorly understood processes of stellar evolution, like diffusive processes, mass loss on the red-giant branch, characterisation of horizontal branches and identification of the physical mechanisms determining their morphologies, evolution, and occurrence of interacting binaries;— in general, calibration of stellar models at low metallicities; in addition, direct measurements of their helium content and ages, tight constraints to dynamical models, and to the origin of their multiple populations;
3. in the open clusters, characterization of convective mixing zones and core growth across the HR diagram, distributions and evolution of rotational velocities, orientation of spin axes, transport of elements and angular momentum; in M67, detailed asteroseismology of solar twins and their parents/progeny; — in general, the ultimate data for calibrating stellar models at near-solar metallicities;
4. to characterize the stellar populations and their age distributions in the core of the nearby dwarf galaxies ω Cen and Sgr dSph, and in the Milky Way’s bulge.

Moreover, the high-precision, high-cadence, long-duration photometric data gathered by such a mission will also enable us to advance our understanding of stellar activity, flares, products of binary interactions, and to constrain the occurrence and properties of exoplanets in different (and largely unexplored) environments, widening the impact of the proposed science case even further.

Acknowledgements AM, JM and FV acknowledge support from the European Research Council Consolidator Grant funding scheme (project ASTEROCHRONOMETRY, G.A. n. 772293, <http://www.asterochronometry.eu>). AM, BM and LG are grateful to the International Space Science Institute (ISSI) for support provided to the asteroSTEP ISSI International Team. AM and WJC acknowledge the support of the UK Science and Technology Facilities Council (STFC). GB is sponsored by the Swiss National Science Foundation (project number 200020 – 172505). WC acknowledges funding from the Swiss National Science Foundation under grant P400P2.183846. We thank G. R. Davies for providing the power spectra used in Fig. 1. N.L. acknowledges financial support from ”Programme National de Physique Stellaire” (PNPS) of CNRS/INSU, France. CC acknowledges partial support from DFG

Grant CH1188/2-1 and from the ChETEC COST Action (CA16117), supported by COST (European Cooperation in Science and Technology).

References

- Aerts C (2015) The age and interior rotation of stars from asteroseismology. *AN* 336(5):477, DOI 10.1002/asna.201512177, 1503.06690
- Aerts C, Mathis S, Rogers T (2018) Angular Momentum Transport in Stellar Interiors. arXiv e-prints arXiv:1809.07779, 1809.07779
- Anders F, Chiappini C, Minchev I, Miglio A, Montalbán J, Mosser B, Rodrigues TS, Santiago BX, Baudin F, Beers TC, da Costa LN, García RA, García-Hernández DA, Holtzman J, Maia MAG, Majewski S, Mathur S, Noels-Grotsch A, Pan K, Schneider DP, Schultheis M, Steinmetz M, Valentini M, Zamora O (2017) Red giants observed by CoRoT and APOGEE: The evolution of the Milky Way's radial metallicity gradient. *A&A*600:A70, DOI 10.1051/0004-6361/201629363, 1608.04951
- Anders F, Khalatyan A, Chiappini C, Queiroz AB, Santiago BX, Jordi C, Girardi L, Brown AGA, Matijevič G, Monari G, Cantat-Gaudin T, Weiler M, Khan S, Miglio A, Carrillo I, Romero-Gómez M, Minchev I, de Jong RS, Antoja T, Ramos P, Steinmetz M, Enke H (2019) Photo-astrometric distances, extinctions, and astrophysical parameters for Gaia DR2 stars brighter than $G = 18$. arXiv e-prints arXiv:1904.11302, 1904.11302
- Arnett WD, Meakin C, Viallet M, Campbell SW, Lattanzio JC, Mocák M (2015) Beyond Mixing-length Theory: A Step Toward 321D. *ApJ*809(1):30, DOI 10.1088/0004-637X/809/1/30, 1503.00342
- Badenes C, Mazzola C, Thompson TA, Covey K, Freeman PE, Walker MG, Moe M, Troup N, Nidever D, Allende Prieto C (2018) Stellar Multiplicity Meets Stellar Evolution and Metallicity: The APOGEE View. *ApJ*854(2):147, DOI 10.3847/1538-4357/aaa765, 1711.00660
- Baglin A, Auvergne M, Barge P, Deleuil M, Michel E, CoRoT Exoplanet Science Team (2009) CoRoT: Description of the Mission and Early Results. In: Pont F, Sasselov D, Holman MJ (eds) *Transiting Planets*, IAU Symposium, vol 253, pp 71–81, DOI 10.1017/S1743921308026252
- Baliunas SL, Donahue RA, Soon WH, Horne JH, Frazer J, Woodard-Eklund L, Bradford M, Rao LM, Wilson OC, Zhang Q, Bennett W, Briggs J, Carroll SM, Duncan DK, Figueroa D, Lanning HH, Misch T, Mueller J, Noyes RW, Poppe D, Porter AC, Robinson CR, Russell J, Shelton JC, Soyumer T, Vaughan AH, Whitney JH (1995) Chromospheric Variations in Main-Sequence Stars. II. *ApJ*438:269, DOI 10.1086/175072
- Baraffe I, Pratt J, Goffrey T, Constantino T, Folini D, Popov MV, Walder R, Viallet M (2017) Lithium Depletion in Solar-like Stars: Effect of Overshooting Based on Realistic Multi-dimensional Simulations. *ApJ*845(1):L6, DOI 10.3847/2041-8213/aa82ff, 1707.09498

- Barbuy B, Chiappini C, Gerhard O (2018) Chemodynamical History of the Galactic Bulge. *ARA&A*56:223–276, DOI 10.1146/annurev-astro-081817-051826, 1805.01142
- Bastian N, de Mink SE (2009) The effect of stellar rotation on colour-magnitude diagrams: on the apparent presence of multiple populations in intermediate age stellar clusters. *MNRAS*398(1):L11–L15, DOI 10.1111/j.1745-3933.2009.00696.x, 0906.1590
- Bastian N, Kamann S, Cabrera-Ziri I, Georgy C, Ekström S, Charbonnel C, de Juan Ovelar M, Usher C (2018) Extended main sequence turnoffs in open clusters as seen by Gaia - I. NGC 2818 and the role of stellar rotation. *MNRAS*480(3):3739–3746, DOI 10.1093/mnras/sty2100, 1807.10779
- Baumgardt H (2017) N -body modelling of globular clusters: masses, mass-to-light ratios and intermediate-mass black holes. *MNRAS*464:2174–2202, DOI 10.1093/mnras/stw2488, 1609.08794
- Bazot M, Benomar O, Christensen-Dalsgaard J, Gizon L, Hanasoge S, Nielsen M, Petit P, Sreenivasan KR (2019) Latitudinal differential rotation in the solar analogues 16 Cygni A and B. *A&A*623:A125, DOI 10.1051/0004-6361/201834594, 1902.01676
- Beck PG, Montalbán J, Kallinger T, De Ridder J, Aerts C, García RA, Hekker S, Dupret MA, Mosser B, Eggenberger P (2012) Fast core rotation in red-giant stars as revealed by gravity-dominated mixed modes. *Nature*481(7379):55–57, DOI 10.1038/nature10612, 1112.2825
- Beck PG, Mathis S, Gallet F, Charbonnel C, Benbakoura M, García RA, do Nascimento JD (2018) Testing tidal theory for evolved stars by using red giant binaries observed by Kepler. *MNRAS*479(1):L123–L128, DOI 10.1093/mnrasl/sly114, 1806.07208
- Bedding TR, Mosser B, Huber D, Montalbán J, Beck P, Christensen-Dalsgaard J, Elsworth YP, García RA, Miglio A, Stello D, White TR, De Ridder J, Hekker S, Aerts C, Barban C, Belkacem K, Broomhall AM, Brown TM, Buzasi DL, Carrier F, Chaplin WJ, di Mauro MP, Dupret MA, Frandsen S, Gilliland RL, Goupil MJ, Jenkins JM, Kallinger T, Kawaler S, Kjeldsen H, Mathur S, Noels A, Silva Aguirre V, Ventura P (2011) Gravity modes as a way to distinguish between hydrogen- and helium-burning red giant stars. *Nature*471(7340):608–611, DOI 10.1038/nature09935, 1103.5805
- Benomar O, Takata M, Shibahashi H, Ceillier T, García RA (2015) Nearly uniform internal rotation of solar-like main-sequence stars revealed by space-based asteroseismology and spectroscopic measurements. *MNRAS*452(3):2654–2674, DOI 10.1093/mnras/stv1493, 1507.01140
- Benomar O, Bazot M, Nielsen MB, Gizon L, Sekii T, Takata M, Hotta H, Hanasoge S, Sreenivasan KR, Christensen-Dalsgaard J (2018) Asteroseismic detection of latitudinal differential rotation in 13 Sun-like stars. *Science* 361(6408):1231–1234, DOI 10.1126/science.aao6571, 1809.07938
- Bensby T, Feltzing S, Gould A, Yee JC, Johnson JA, Asplund M, Meléndez J, Lucatello S, Howes LM, McWilliam A, Udalski A, Szymański MK, Soszyński I, Poleski R, Wyrzykowski L, Ulaczyk K, Kozłowski S, Pietrukowicz P, Skowron J, Mróz P, Pawlak M, Abe F, Asakura Y, Bhattacharya A, Bond

- IA, Bennett DP, Hirao Y, Nagakane M, Koshimoto N, Sumi T, Suzuki D, Tristram PJ (2017) Chemical evolution of the Galactic bulge as traced by microlensed dwarf and subgiant stars. VI. Age and abundance structure of the stellar populations in the central sub-kpc of the Milky Way. *A&A*605:A89, DOI 10.1051/0004-6361/201730560, 1702.02971
- Borucki WJ, Koch D, Basri G, Batalha N, Brown T, Caldwell D, Caldwell J, Christensen-Dalsgaard J, Cochran WD, DeVore E (2010) Kepler Planet-Detection Mission: Introduction and First Results. *Science* 327(5968):977, DOI 10.1126/science.1185402
- Bossini D, Miglio A, Salaris M, Pietrinferni A, Montalbán J, Bressan A, Noels A, Cassisi S, Girardi L, Marigo P (2015) Uncertainties on near-core mixing in red-clump stars: effects on the period spacing and on the luminosity of the AGB bump. *MNRAS*453(3):2290–2301, DOI 10.1093/mnras/stv1738, 1507.07797
- Bossini D, Miglio A, Salaris M, Vrad M, Cassisi S, Mosser B, Montalbán J, Girardi L, Noels A, Bressan A (2017) Kepler red-clump stars in the field and in open clusters: constraints on core mixing. *MNRAS*469(4):4718–4725, DOI 10.1093/mnras/stx1135, 1705.03077
- Bovy J, Leung HW, Hunt JAS, Mackereth JT, Garcia-Hernandez DA, Roman-Lopes A (2019) Life in the fast lane: a direct view of the dynamics, formation, and evolution of the Milky Way’s bar. arXiv e-prints arXiv:1905.11404, 1905.11404
- Boyajian TS, LaCourse DM, Rappaport SA, Fabrycky D, Fischer DA, Gandolfi D, Kennedy GM, Korhonen H, Liu MC, Moor A (2016) Planet Hunters IX. KIC 8462852 - where’s the flux? *MNRAS*457(4):3988–4004, DOI 10.1093/mnras/stw218, 1509.03622
- Boyer ML, van Loon JT, McDonald I, Gordon KD, Babler B, Block M, Bracker S, Engelbracht C, Hora J, Indebetouw R (2010) Is Dust Forming on the Red Giant Branch in 47 Tuc? *ApJ*711(2):L99–L103, DOI 10.1088/2041-8205/711/2/L99, 1002.1348
- Bragaglia A, Carretta E, Gratton R, D’Orazi V, Cassisi S, Lucatello S (2010) Helium in first and second-generation stars in globular clusters from spectroscopy of red giants. *A&A*519:A60, DOI 10.1051/0004-6361/201014702, 1005.2659
- Bragaglia A, Carretta E, D’Orazi V, Sollima A, Donati P, Gratton RG, Lucatello S (2017) NGC 6535: the lowest mass Milky Way globular cluster with a Na-O anti-correlation? Cluster mass and age in the multiple population context. *A&A*607:A44, DOI 10.1051/0004-6361/201731526, 1708.07705
- Brogaard K, Sandquist E, Jessen-Hansen J, Grundahl F, Frandsen S (2015) Exploiting the Open Clusters in the Kepler and CoRoT Fields. In: *Asteroseismology of Stellar Populations in the Milky Way*, vol 39, p 51, DOI 10.1007/978-3-319-10993-0_6, 1409.2271
- Brogaard K, Christiansen SM, Grundahl F, Miglio A, Izzard RG, Tauris TM, Sandquist EL, VandenBerg DA, Jessen-Hansen J, Arentoft T (2018) The blue straggler V106 in NGC 6791: a prototype progenitor of old single giants masquerading as young. *MNRAS*481(4):5062–5072, DOI

- 10.1093/mnras/sty2504, 1809.00705
- Broomhall AM, Miglio A, Montalbán J, Eggenberger P, Chaplin WJ, Elsworth Y, Scufflaire R, Ventura P, Verner GA (2014) Prospects for asteroseismic inference on the envelope helium abundance in red giant stars. *MNRAS*440(2):1828–1843, DOI 10.1093/mnras/stu393, 1403.7045
- Brown TM, Sahu K, Anderson J, Tumlinson J, Valenti JA, Smith E, Jeffery EJ, Renzini A, Zoccali M, Ferguson HC, VandenBerg DA, Bond HE, Casertano S, Valenti E, Minniti D, Livio M, Panagia N (2010) The WFC3 Galactic Bulge Treasury Program: Metallicity Estimates for the Stellar Population and Exoplanet Hosts. *ApJ*725(1):L19–L23, DOI 10.1088/2041-8205/725/1/L19, 1011.0457
- Brucalassi A, Pasquini L, Saglia R, Ruiz MT, Bonifacio P, Leão I, Canto Martins BL, de Medeiros JR, Bedin LR, Biazzo K, Melo C, Lovis C, Randich S (2016) Search for giant planets in M67. III. Excess of hot Jupiters in dense open clusters. *A&A*592:L1, DOI 10.1051/0004-6361/201527561, 1606.05247
- Buldgen G, Reese DR, Dupret MA (2015a) Using seismic inversions to obtain an indicator of internal mixing processes in main-sequence solar-like stars. *A&A*583:A62, DOI 10.1051/0004-6361/201526390, 1507.06466
- Buldgen G, Reese DR, Dupret MA, Samadi R (2015b) Stellar acoustic radii, mean densities, and ages from seismic inversion techniques. *A&A*574:A42, DOI 10.1051/0004-6361/201424613, 1411.2416
- Buldgen G, Reese DR, Dupret MA (2018) Constraining convective regions with asteroseismic linear structural inversions. *A&A*609:A95, DOI 10.1051/0004-6361/201730693, 1711.05031
- Buldgen G, Salmon SJAJ, Noels A, Scufflaire R, Montalban J, Baturin VA, Eggenberger P, Gryaznov VK, Iosilevskiy IL, Meynet G, Chaplin WJ, Miglio A, Oreshina AV, Richard O, Starostin AN (2019) Combining multiple structural inversions to constrain the solar modelling problem. *A&A*621:A33, DOI 10.1051/0004-6361/201833971, 1809.08958
- Cantiello M, Mankovich C, Bildsten L, Christensen-Dalsgaard J, Paxton B (2014) Angular Momentum Transport within Evolved Low-mass Stars. *ApJ*788(1):93, DOI 10.1088/0004-637X/788/1/93, 1405.1419
- Cantiello M, Fuller J, Bildsten L (2016) Asteroseismic Signatures of Evolving Internal Stellar Magnetic Fields. *ApJ*824(1):14, DOI 10.3847/0004-637X/824/1/14, 1602.03056
- Carretta E, Bragaglia A, Gratton RG, Lucatello S, Bellazzini M, Catanzaro G, Leone F, Momany Y, Piotto G, D’Orazi V (2010) M54 + Sagittarius = ω Centauri. *ApJ*714(1):L7–L11, DOI 10.1088/2041-8205/714/1/L7, 1002.1963
- Casagrande L, Silva Aguirre V, Schlesinger KJ, Stello D, Huber D, Serenelli AM, Schönrich R, Cassisi S, Pietrinferni A, Hodgkin S, Milone AP, Feltzing S, Asplund M (2016) Measuring the vertical age structure of the Galactic disc using asteroseismology and SAGA. *MNRAS*455(1):987–1007, DOI 10.1093/mnras/stv2320, 1510.01376
- Casali G, Magrini L, Tognelli E, Jackson R, Jeffries RD, Lagarde N, Taut-

- vaisiene G, Masseron T, Degl'Innocenti S, Prada Moroni PG, Kordopatis G, Pancino E, Randich S, Feltzing S, Sahlholdt C, Spina L, Friel E, Roccatagliata V, Sanna N, Bragaglia A, Drazdauskas A, Mikolaitis S, Minkeviciute R, Stonkute E, Chorniy Y, Bagdonas V, Jimenez-Esteban F, Martell S, Van der Swaelmen M, Gilmore G, Vallenari A, Bensby T, Koposov SE, Korn A, Worley C, Smiljanic R, Bergemann M, Carraro G, Damiani F, Prisinzano L, Bonito R, Franciosini E, Gonneau A, Hourihane A, Jofre P, Lewis J, Morbidelli L, Sacco G, Sousa SG, Zaggia S, Lanzafame AC, Heiter U, Frasca A, Bayo A (2019) The Gaia-ESO survey: Calibrating a relationship between Age and the [C/N] abundance ratio with open clusters. arXiv e-prints arXiv:1907.07350, 1907.07350
- Ceillier T, Eggenberger P, García RA, Mathis S (2013) Understanding angular momentum transport in red giants: the case of KIC 7341231. *A&A*555:A54, DOI 10.1051/0004-6361/201321473, 1305.5630
- Chaboyer B, Sarajedini A, Demarque P (1992) Ages of Globular Clusters and Helium Diffusion. *ApJ*394:515, DOI 10.1086/171602
- Chantereau W, Charbonnel C, Decressin T (2015) Evolution of long-lived globular cluster stars. I. Grid of stellar models with helium enhancement at [Fe/H] = -1.75. *A&A*578:A117, DOI 10.1051/0004-6361/201525929, 1504.01878
- Chantereau W, Salaris M, Bastian N, Martocchia S (2019) Helium enrichment in intermediate-age Magellanic Clouds clusters: towards an ubiquity of multiple stellar populations? *MNRAS*484(4):5236–5244, DOI 10.1093/mnras/stz378, 1902.01806
- Chaplin WJ, Miglio A (2013) Asteroseismology of Solar-Type and Red-Giant Stars. *ARA&A*51(1):353–392, DOI 10.1146/annurev-astro-082812-140938, 1303.1957
- Chaplin WJ, Kjeldsen H, Christensen-Dalsgaard J, Basu S, Miglio A, Appourchaux T, Bedding TR, Elsworth Y, García RA, Gilliland RL (2011) Ensemble Asteroseismology of Solar-Type Stars with the NASA Kepler Mission. *Science* 332(6026):213, DOI 10.1126/science.1201827, 1109.4723
- Chaplin WJ, Sanchis-Ojeda R, Campante TL, Handberg R, Stello D, Winn JN, Basu S, Christensen-Dalsgaard J, Davies GR, Metcalfe TS, Buchhave LA, Fischer DA, Bedding TR, Cochran WD, Elsworth Y, Gilliland RL, Hekker S, Huber D, Isaacson H, Karoff C, Kawaler SD, Kjeldsen H, Latham DW, Lund MN, Lundkvist M, Marcy GW, Miglio A, Barclay T, Lissauer JJ (2013) Asteroseismic Determination of Obliquities of the Exoplanet Systems Kepler-50 and Kepler-65. *ApJ*766(2):101, DOI 10.1088/0004-637X/766/2/101, 1302.3728
- Chiappini C, Anders F, Rodrigues TS, Miglio A, Montalbán J, Mosser B, Girardi L, Valentini M, Noels A, Morel T, Minchev I, Steinmetz M, Santiago BX, Schultheis M, Martig M, da Costa LN, Maia MAG, Allende Prieto C, de Assis Peralta R, Hekker S, Thewissen N, Kallinger T, García RA, Mathur S, Baudin F, Beers TC, Cunha K, Harding P, Holtzman J, Majewski S, Mészáros S, Nidever D, Pan K, Schiavon RP, Shetrone MD, Schneider DP, Stassun K (2015) Young $[\alpha/\text{Fe}]$ -enhanced stars discovered by CoRoT and

- APOGEE: What is their origin? *A&A*576:L12, DOI 10.1051/0004-6361/201525865, 1503.06990
- Christensen-Dalsgaard J (2002) Helioseismology. *Reviews of Modern Physics* 74(4):1073–1129, DOI 10.1103/RevModPhys.74.1073, astro-ph/0207403
- Christensen-Dalsgaard J, Aguirre VS (2018) Ages for Exoplanet Host Stars, p 184. DOI 10.1007/978-3-319-55333-7_184
- Conroy C (2013) Modeling the Panchromatic Spectral Energy Distributions of Galaxies. *ARA&A*51(1):393–455, DOI 10.1146/annurev-astro-082812-141017, 1301.7095
- Constantino T, Campbell SW, Christensen-Dalsgaard J, Lattanzio JC, Stello D (2015) The treatment of mixing in core helium burning models - I. Implications for asteroseismology. *MNRAS*452(1):123–145, DOI 10.1093/mnras/stv1264, 1506.01209
- Cordoni G, Milone AP, Marino AF, Di Criscienzo M, D’Antona F, Dotter A, Lagioia EP, Tailo M (2018) Extended Main-sequence Turnoff as a Common Feature of Milky Way Open Clusters. *ApJ*869(2):139, DOI 10.3847/1538-4357/aaedc1, 1811.01192
- Corsaro E, Lee YN, García RA, Hennebelle P, Mathur S, Beck PG, Mathis S, Stello D, Bouvier J (2017) Spin alignment of stars in old open clusters. *Nature Astronomy* 1:0064, DOI 10.1038/s41550-017-0064, 1703.05588
- Costa AD, Canto Martins BL, Bravo JP, Paz-Chinchón F, das Chagas ML, Leão IC, Pereira de Oliveira G, Rodrigues da Silva R, Roque S, de Oliveira LLA (2015) Kepler Rapidly Rotating Giant Stars. *ApJ*807(2):L21, DOI 10.1088/2041-8205/807/2/L21, 1506.06644
- Costa G, Girardi L, Bressan A, Marigo P, Rodrigues TS, Chen Y, Lanza A, Goudfrooij P (2019) Mixing by overshooting and rotation in intermediate-mass stars. *MNRAS*485(4):4641–4657, DOI 10.1093/mnras/stz728, 1903.04368
- Cummings JD, Kalirai JS, Choi J, Georgy C, Tremblay PE, Ramirez-Ruiz E (2019) A Novel Approach to Constrain Rotational Mixing and Convective-core Overshoot in Stars Using the Initial-Final Mass Relation. *ApJ*871(1):L18, DOI 10.3847/2041-8213/aafc2d, 1901.02904
- Cunha MS, Stello D, Avelino PP, Christensen-Dalsgaard J, Townsend RHD (2015) Structural Glitches near the Cores of Red Giants Revealed by Oscillations in g-mode Period Spacings from Stellar Models. *ApJ*805(2):127, DOI 10.1088/0004-637X/805/2/127, 1503.09085
- Da Costa GS, Freeman KC (1976) The structure and mass function of the globular cluster M3. *ApJ*206:128–137, DOI 10.1086/154363
- Davies GR, Miglio A (2016) Asteroseismology of red giants: From analysing light curves to estimating ages. *AN* 337(8-9):774, DOI 10.1002/asna.201612371, 1601.02802
- Davies GR, Silva Aguirre V, Bedding TR, Handberg R, Lund MN, Chaplin WJ, Huber D, White TR, Benomar O, Hekker S (2016) Oscillation frequencies for 35 Kepler solar-type planet-hosting stars using Bayesian techniques and machine learning. *MNRAS*456(2):2183–2195, DOI 10.1093/mnras/stv2593, 1511.02105

- Dawson RI, Johnson JA (2018) Origins of Hot Jupiters. *ARA&A*56:175–221, DOI 10.1146/annurev-astro-081817-051853, 1801.06117
- de Assis Peralta R, Samadi R, Michel E (2018) A new method for extracting seismic indices and granulation parameters: Results for more than 20,000 CoRoT and Kepler red giants. *AN* 339(134):134–157, DOI 10.1002/asna.201813469, 1805.04296
- Deal M, Escobar ME, Vauclair S, Vauclair G, Hui-Bon-Hoa A, Richard O (2017) Asteroseismology of the exoplanet-host F-type star 94 Ceti: Impact of atomic diffusion on the stellar parameters. *A&A*601:A127, DOI 10.1051/0004-6361/201629825, 1702.03170
- Deheuvels S, García RA, Chaplin WJ, Basu S, Antia HM, Appourchaux T, Benomar O, Davies GR, Elsworth Y, Gizon L (2012) Seismic Evidence for a Rapidly Rotating Core in a Lower-giant-branch Star Observed with Kepler. *ApJ*756(1):19, DOI 10.1088/0004-637X/756/1/19, 1206.3312
- Deheuvels S, Doğan G, Goupil MJ, Appourchaux T, Benomar O, Bruntt H, Campante TL, Casagrande L, Ceillier T, Davies GR (2014) Seismic constraints on the radial dependence of the internal rotation profiles of six Kepler subgiants and young red giants. *A&A*564:A27, DOI 10.1051/0004-6361/201322779, 1401.3096
- Deheuvels S, Brandão I, Silva Aguirre V, Ballot J, Michel E, Cunha MS, Lebreton Y, Appourchaux T (2016) Measuring the extent of convective cores in low-mass stars using Kepler data: toward a calibration of core overshooting. *A&A*589:A93, DOI 10.1051/0004-6361/201527967, 1603.02332
- Di Mauro MP, Ventura R, Cardini D, Stello D, Christensen-Dalsgaard J, Dziembowski WA, Paternò L, Beck PG, Bloemen S, Davies GR (2016) Internal Rotation of the Red-giant Star KIC 4448777 by Means of Asteroseismic Inversion. *ApJ*817(1):65, DOI 10.3847/0004-637X/817/1/65, 1511.06160
- Dufton PL, Smartt SJ, Lee JK, Ryans RSI, Hunter I, Evans CJ, Herrero A, Trundle C, Lennon DJ, Irwin MJ (2006) The VLT-FLAMES survey of massive stars: stellar parameters and rotational velocities in NGC 3293, NGC 4755 and NGC 6611. *A&A*457(1):265–280, DOI 10.1051/0004-6361:20065392, astro-ph/0606409
- Dupree AK, Avrett EH (2013) Direct Evaluation of the Helium Abundances in Omega Centauri. *ApJ*773(2):L28, DOI 10.1088/2041-8205/773/2/L28, 1307.5860
- Dupret MA, Grigahcène A, Garrido R, Gabriel M, Scufflaire R (2005) Convection-pulsation coupling. II. Excitation and stabilization mechanisms in δ Sct and γ Dor stars. *A&A*435(3):927–939, DOI 10.1051/0004-6361:20041817
- Dupret MA, Belkacem K, Samadi R, Montalbán J, Moreira O, Miglio A, Godart M, Ventura P, Ludwig HG, Grigahcène A, Goupil MJ, Noels A, Caffau E (2009) Theoretical amplitudes and lifetimes of non-radial solar-like oscillations in red giants. *A&A*506(1):57–67, DOI 10.1051/0004-6361/200911713, 0906.3951
- Dziembowski WA, Soszyński I (2010) Acoustic oscillations in stars near the tip of the red giant branch. *A&A*524:A88, DOI 10.1051/0004-6361/201015205,

1009.5989

- Eggenberger P, Maeder A, Meynet G (2005) Stellar evolution with rotation and magnetic fields. IV. The solar rotation profile. *A&A*440(1):L9–L12, DOI 10.1051/0004-6361:200500156, [astro-ph/0508455](#)
- Eggenberger P, Montalbán J, Miglio A (2012) Angular momentum transport in stellar interiors constrained by rotational splittings of mixed modes in red giants. *A&A*544:L4, DOI 10.1051/0004-6361/201219729, 1207.1023
- Eggenberger P, Deheuvels S, Miglio A, Ekström S, Georgy C, Meynet G, Lagarde N, Salmon S, Buldgen G, Montalbán J (2019) Asteroseismology of evolved stars to constrain the internal transport of angular momentum. I. Efficiency of transport during the subgiant phase. *A&A*621:A66, DOI 10.1051/0004-6361/201833447, 1812.04995
- ESA (2017) Plato definition study report (red book). <http://sci.esa.int/plato/59252-plato-definition-study-report-red-book/>
- Ferrarese L, Merritt D (2000) A Fundamental Relation between Supermassive Black Holes and Their Host Galaxies. *ApJ*539:L9–L12, DOI 10.1086/312838, [astro-ph/0006053](#)
- Forbes DA, Bridges T (2010) Accreted versus in situ Milky Way globular clusters. *MNRAS*404:1203–1214, DOI 10.1111/j.1365-2966.2010.16373.x, 1001.4289
- Fujii MS, Hori Y (2019) Survival rates of planets in open clusters: the Pleiades, Hyades, and Praesepe clusters. *A&A*624:A110, DOI 10.1051/0004-6361/201834677, 1811.08598
- Fuller J, Piro AL, Jermyn AS (2019) Slowing the spins of stellar cores. *MNRAS*485(3):3661–3680, DOI 10.1093/mnras/stz514, 1902.08227
- García RA, Ballot J (2019) Asteroseismology of solar-type stars. arXiv e-prints arXiv:1906.12262, 1906.12262
- García RA, Mathur S, Salabert D, Ballot J, Régulo C, Metcalfe TS, Baglin A (2010) CoRoT Reveals a Magnetic Activity Cycle in a Sun-Like Star. *Science* 329(5995):1032, DOI 10.1126/science.1191064, 1008.4399
- Gebhardt K, Bender R, Bower G, Dressler A, Faber SM, Filippenko AV, Green R, Grillmair C, Ho LC, Kormendy J, Lauer TR, Magorrian J, Pinkney J, Richstone D, Tremaine S (2000) A Relationship between Nuclear Black Hole Mass and Galaxy Velocity Dispersion. *ApJ*539:L13–L16, DOI 10.1086/312840, [astro-ph/0006289](#)
- Gehan C, Mosser B, Michel E, Samadi R, Kallinger T (2018) Core rotation braking on the red giant branch for various mass ranges. *A&A*616:A24, DOI 10.1051/0004-6361/201832822, 1802.04558
- Gieles M, Zocchi A (2015) A family of lowered isothermal models. *MNRAS*454:576–592, DOI 10.1093/mnras/stv1848, 1508.02120
- Giesers B, Dreizler S, Husser TO, Kamann S, Anglada Escudé G, Brinchmann J, Carollo CM, Roth MM, Weilbacher PM, Wisotzki L (2018) A detached stellar-mass black hole candidate in the globular cluster NGC 3201. *MNRAS*475:L15–L19, DOI 10.1093/mnrasl/slx203
- Gilliland RL, Brown TM, Guhathakurta P, Sarajedini A, Milone EF, Albrow MD, Baliber NR, Bruntt H, Burrows A, Charbonneau D (2000) A

- Lack of Planets in 47 Tucanae from a Hubble Space Telescope Search. *ApJ*545(1):L47–L51, DOI 10.1086/317334
- Girardi L, Groenewegen MAT, Hatziminaoglou E, da Costa L (2005) Star counts in the Galaxy. Simulating from very deep to very shallow photometric surveys with the TRILEGAL code. *A&A*436:895–915, DOI 10.1051/0004-6361:20042352, [astro-ph/0504047](#)
- Gizon L, Solanki SK (2003) Determining the inclination of the rotation axis of a sun-like star. *The Astrophysical Journal* 589(2):1009–1019, DOI 10.1086/374715, URL <https://doi.org/10.1086%2F374715>
- Güdel M (2007) The Sun in Time: Activity and Environment. *Living Reviews in Solar Physics* 4(1):3, DOI 10.12942/lrsp-2007-3, 0712.1763
- Gunn JE, Griffin RF (1979) Dynamical studies of globular clusters based on photoelectric radial velocities of individual stars. I - M3. *AJ*84:752–773, DOI 10.1086/112477
- Handberg R, Brogaard K, Miglio A, Bossini D, Elsworth Y, Slumstrup D, Davies GR, Chaplin WJ (2017) NGC 6819: testing the asteroseismic mass scale, mass loss and evidence for products of non-standard evolution. *MNRAS*472(1):979–997, DOI 10.1093/mnras/stx1929, 1707.08223
- Heger A, Woosley SE, Spruit HC (2005) Presupernova Evolution of Differentially Rotating Massive Stars Including Magnetic Fields. *ApJ*626(1):350–363, DOI 10.1086/429868, [astro-ph/0409422](#)
- Hekker S, Christensen-Dalsgaard J (2017) Giant star seismology. *A&A Rev.*25(1):1, DOI 10.1007/s00159-017-0101-x, 1609.07487
- Hekker S, Johnson JA (2019) Origin of α -rich young stars: clues from C, N, and O. *MNRAS*487(3):4343–4354, DOI 10.1093/mnras/stz1554, 1906.10830
- Heyl J, Kalirai J, Richer HB, Marigo P, Antolini E, Goldsbury R, Parada J (2015) When do stars in 47 Tucanae lose their mass? *ApJ*810(2):127, DOI 10.1088/0004-637X/810/2/127, 1502.07306
- Hilker M, Kayser A, Richtler T, Willemsen P (2004) The extended star formation history of ω Centauri. *A&A*422:L9–L12, DOI 10.1051/0004-6361:20040188, [astro-ph/0406017](#)
- Howell SB, Sobeck C, Haas M, Still M, Barclay T, Mullally F, Troeltzsch J, Aigrain S, Bryson ST, Caldwell D (2014) The K2 Mission: Characterization and Early Results. *PASP*126(938):398, DOI 10.1086/676406, 1402.5163
- Howell SB, Mason E, Boyd P, Smith KL, Gelino DM (2016) Rapidly Rotating, X-Ray Bright Stars in the Kepler Field. *ApJ*831(1):27, DOI 10.3847/0004-637X/831/1/27, 1608.07828
- Huang W, Gies DR (2006) Stellar Rotation in Young Clusters. I. Evolution of Projected Rotational Velocity Distributions. *ApJ*648(1):580–590, DOI 10.1086/505782, [astro-ph/0510450](#)
- Huber D, Carter JA, Barbieri M, Miglio A, Deck KM, Fabrycky DC, Montet BT, Buchhave LA, Chaplin WJ, Hekker S (2013) Stellar Spin-Orbit Misalignment in a Multiplanet System. *Science* 342(6156):331–334, DOI 10.1126/science.1242066, 1310.4503
- Husser TO, Kamann S, Dreizler S, Wendt M, Wulff N, Bacon R, Wisotzki L, Brinchmann J, Weilbacher PM, Roth MM, Monreal-Ibero A (2016)

- MUSE crowded field 3D spectroscopy of over 12 000 stars in the globular cluster NGC 6397. I. The first comprehensive HRD of a globular cluster. *A&A*588:A148, DOI 10.1051/0004-6361/201526949, 1602.01649
- Ibata R, Bellazzini M, Chapman SC, Dalessandro E, Ferraro F, Irwin M, Lanzoni B, Lewis GF, Mackey AD, Miocchi P, Varghese A (2009) Density and Kinematic Cusps in M54 at the Heart of the Sagittarius Dwarf Galaxy: Evidence for a $10^4 M_{\text{sun}}$ Black Hole? *ApJ*699(2):L169–L173, DOI 10.1088/0004-637X/699/2/L169, 0906.4894
- Izzard RG, Preece H, Jofre P, Halabi GM, Masseron T, Tout CA (2018) Binary stars in the Galactic thick disc. *MNRAS*473(3):2984–2999, DOI 10.1093/mnras/stx2355, 1709.05237
- Jofré P, Jorissen A, Van Eck S, Izzard RG, Masseron T, Hawkins K, Gilmore G, Paladini C, Escorza A, Blanco-Cuaresma S, Manick R (2016) Cannibals in the thick disk: the young α -rich stars as evolved blue stragglers. *A&A*595:A60, DOI 10.1051/0004-6361/201629356, 1603.08992
- Kains N, Calamida A, Sahu KC, Anderson J, Casertano S, Bramich DM (2018) A Search for Black Hole Microlensing Signatures in Globular Cluster NGC 6656 (M22). *ApJ*867(1):37, DOI 10.3847/1538-4357/aae311, 1810.01417
- Kamann S, Husser TO, Brinchmann J, Emsellem E, Weilbacher PM, Wisotzki L, Wendt M, Krajinović D, Roth MM, Bacon R, Dreizler S (2016) MUSE crowded field 3D spectroscopy of over 12 000 stars in the globular cluster NGC 6397. II. Probing the internal dynamics and the presence of a central black hole. *A&A*588:A149, DOI 10.1051/0004-6361/201527065, 1602.01643
- Kamann S, Husser TO, Dreizler S, Emsellem E, Weilbacher PM, Martens S, Bacon R, den Brok M, Giesers B, Krajinović D, Roth MM, Wendt M, Wisotzki L (2018) A stellar census in globular clusters with MUSE: The contribution of rotation to cluster dynamics studied with 200 000 stars. *MNRAS*473:5591–5616, DOI 10.1093/mnras/stx2719, 1710.07257
- Kamann S, Bastian NJ, Gieles M, Balbinot E, Hénault-Brunet V (2019) Linking the rotation of a cluster to the spins of its stars: the kinematics of NGC 6791 and NGC 6819 in 3D. *MNRAS*483(2):2197–2206, DOI 10.1093/mnras/sty3144, 1811.08476
- Kamiaka S, Benomar O, Suto Y (2018) Reliability of stellar inclination estimated from asteroseismology: analytical criteria, mock simulations, and Kepler data analysis. *MNRAS*479(1):391–405, DOI 10.1093/mnras/sty1358, 1805.07044
- Karoff C, Knudsen MF, De Cat P, Bonanno A, Fogtman-Schulz A, Fu J, Frasca A, Inceoglu F, Olsen J, Zhang Y, Hou Y, Wang Y, Shi J, Zhang W (2016) Observational evidence for enhanced magnetic activity of superflare stars. *Nature Communications* 7:11058, DOI 10.1038/ncomms11058
- Karoff C, Metcalfe TS, Santos ARG, Montet BT, Isaacson H, Witzke V, Shapiro AI, Mathur S, Davies GR, Lund MN, Garcia RA, Brun AS, Salabert D, Avelino PP, van Saders J, Egeland R, Cunha MS, Campante TL, Chaplin WJ, Krivova N, Solanki SK, Stritzinger M, Knudsen MF (2018) The Influence of Metallicity on Stellar Differential Rotation and Magnetic Activity. *ApJ*852(1):46, DOI 10.3847/1538-4357/aaa026, 1711.07716

- Kawaler SD (2015) Rotation of White Dwarf Stars. In: Dufour P, Bergeron P, Fontaine G (eds) 19th European Workshop on White Dwarfs, Astronomical Society of the Pacific Conference Series, vol 493, p 65, 1410.6934
- Kiefer R, Schad A, Davies G, Roth M (2017) Stellar magnetic activity and variability of oscillation parameters: An investigation of 24 solar-like stars observed by Kepler. *A&A*598:A77, DOI 10.1051/0004-6361/201628469, 1611.02029
- Kızıltan B, Baumgardt H, Loeb A (2017) Astrophysical parameters of Galactic open clusters. *Nature*4542:203–205
- Korn AJ, Grundahl F, Richard O, Mashonkina L, Barklem PS, Collet R, Gustafsson B, Piskunov N (2007) Atomic Diffusion and Mixing in Old Stars. I. Very Large Telescope FLAMES-UVES Observations of Stars in NGC 6397. *ApJ*671(1):402–419, DOI 10.1086/523098, 0709.0639
- Kuzlewicz JS, Chaplin WJ, North TSH, Farr WM, Bell KJ, Davies GR, Campante TL, Hekker S (2019) Bayesian hierarchical inference of asteroseismic inclination angles. *MNRAS*1650, DOI 10.1093/mnras/stz1689, 1907.01565
- Lagarde N, Miglio A, Eggenberger P, Morel T, Montalbán J, Mosser B, Rodrigues TS, Girardi L, Rainer M, Poretti E, Barban C, Hekker S, Kallinger T, Valentini M, Carrier F, Hareter M, Mantegazza L, Elsworth Y, Michel E, Baglin A (2015) Models of red giants in the CoRoT asteroseismology fields combining asteroseismic and spectroscopic constraints. *A&A*580:A141, DOI 10.1051/0004-6361/201525856, 1505.01529
- Lagarde N, Reylé C, Robin AC, Tautvaisiene G, Drazdauskas A, Mikolaitis Š, Minkeviciute R, Stonkute E, Chorniy Y, Bagdonas V (2019) The Gaia-ESO Survey: impact of extra mixing on C and N abundances of giant stars. *A&A*621:A24, DOI 10.1051/0004-6361/201732433, 1806.01868
- Lagioia EP, Milone AP, Marino AF, Cassisi S, Aparicio AJ, Piotto G, Anderson J, Barbuy B, Bedin LR, Bellini A (2018) The Hubble Space Telescope UV Legacy Survey of Galactic Globular Clusters - XII. The RGB bumps of multiple stellar populations. *MNRAS*475(3):4088–4103, DOI 10.1093/mnras/sty083, 1801.03395
- Leaman R, VandenBerg DA, Mendel JT (2013) The bifurcated age-metallicity relation of Milky Way globular clusters and its implications for the accretion history of the galaxy. *MNRAS*436(1):122–135, DOI 10.1093/mnras/stt1540, 1309.0822
- Liu F, Asplund M, Yong D, Meléndez J, Ramírez I, Karakas AI, Carlos M, Marino AF (2016) The chemical compositions of solar twins in the open cluster M67. *MNRAS*463(1):696–704, DOI 10.1093/mnras/stw2045, 1608.03788
- Lund MN, Miesch MS, Christensen-Dalsgaard J (2014) Differential Rotation in Main-sequence Solar-like Stars: Qualitative Inference from Asteroseismic Data. *ApJ*790(2):121, DOI 10.1088/0004-637X/790/2/121, 1406.7873
- Lützgendorf N, Baumgardt H, Kruijssen JMD (2013) N-body simulations of globular clusters in tidal fields: Effects of intermediate-mass black holes. *A&A*558:A117, DOI 10.1051/0004-6361/201321927, 1309.0451

- Mackereth JT, Bovy J, Schiavon RP, Zasowski G, Cunha K, Frinchaboy PM, García Perez AE, Hayden MR, Holtzman J, Majewski SR, Mészáros S, Nidever DL, Pinsonneault M, Shetrone MD (2017) The age-metallicity structure of the Milky Way disc using APOGEE. *MNRAS*471(3):3057–3078, DOI 10.1093/mnras/stx1774, 1706.00018
- Maeder A, Meynet G, Lagarde N, Charbonnel C (2013) The thermohaline, Richardson, Rayleigh-Taylor, Solberg-Høiland, and GSF criteria in rotating stars. *A&A*553:A1, DOI 10.1051/0004-6361/201220936, 1303.3230
- Maehara H, Shibayama T, Notsu S, Notsu Y, Nagao T, Kusaba S, Honda S, Nogami D, Shibata K (2012) Superflares on solar-type stars. *Nature*485(7399):478–481, DOI 10.1038/nature11063
- Malmberg D, Davies MB, Heggie DC (2011) The effects of fly-bys on planetary systems. *MNRAS*411(2):859–877, DOI 10.1111/j.1365-2966.2010.17730.x, 1009.4196
- Marino AF, Milone AP, Przybilla N, Bergemann M, Lind K, Asplund M, Cassisi S, Catelan M, Casagrande L, Valcarce AAR (2014) Helium enhanced stars and multiple populations along the horizontal branch of NGC 2808: direct spectroscopic measurements. *MNRAS*437(2):1609–1627, DOI 10.1093/mnras/stt1993, 1310.4527
- Marques JP, Goupil MJ, Lebreton Y, Talon S, Palacios A, Belkacem K, Ouazzani RM, Mosser B, Moya A, Morel P (2013) Seismic diagnostics for transport of angular momentum in stars. I. Rotational splittings from the pre-main sequence to the red-giant branch. *A&A*549:A74, DOI 10.1051/0004-6361/201220211, 1211.1271
- Martig M, Rix HW, Silva Aguirre V, Hekker S, Mosser B, Elsworth Y, Bovy J, Stello D, Anders F, García RA, Tayar J, Rodrigues TS, Basu S, Carrera R, Ceillier T, Chaplin WJ, Chiappini C, Frinchaboy PM, García-Hernández DA, Hearty FR, Holtzman J, Johnson JA, Majewski SR, Mathur S, Mészáros S, Miglio A, Nidever D, Pan K, Pinsonneault M, Schiavon RP, Schneider DP, Serenelli A, Shetrone M, Zamora O (2015) Young α -enriched giant stars in the solar neighbourhood. *MNRAS*451(2):2230–2243, DOI 10.1093/mnras/stv1071, 1412.3453
- Martig M, Foesneau M, Rix HW, Ness M, Mészáros S, García-Hernández DA, Pinsonneault M, Serenelli A, Silva Aguirre V, Zamora O (2016) Red giant masses and ages derived from carbon and nitrogen abundances. *MNRAS*456(4):3655–3670, DOI 10.1093/mnras/stv2830, 1511.08203
- Martocchia S, Dalessandro E, Lardo C, Cabrera-Ziri I, Bastian N, Kozhurina-Platais V, Salaris M, Chantreau W, Geisler D, Hilker M (2019) The search for multiple populations in Magellanic Clouds clusters - V. Correlation between cluster age and abundance spreads. *MNRASp* 1528, DOI 10.1093/mnras/stz1596, 1906.03273
- Massari D, Koppelman HH, Helmi A (2019) The origin of the Galaxy’s system of globular clusters. arXiv:190608271 arXiv:1906.08271, 1906.08271
- Masseron T, Gilmore G (2015) Carbon, nitrogen and α -element abundances determine the formation sequence of the Galactic thick and thin discs. *MNRAS*453(2):1855–1866, DOI 10.1093/mnras/stv1731, 1503.00537

- Masuda K, Winn JN (2017) Reassessment of the Null Result of the HST Search for Planets in 47 Tucanae. *AJ*153(4):187, DOI 10.3847/1538-3881/aa647c, 1703.06136
- Mathis S (2009) Transport by gravito-inertial waves in differentially rotating stellar radiation zones. I - Theoretical formulation. *A&A*506(2):811–828, DOI 10.1051/0004-6361/200810544
- Mazumdar A, Monteiro MJPF, Ballot J, Antia HM, Basu S, Houdek G, Mathur S, Cunha MS, Silva Aguirre V, García RA, Salabert D, Verner GA, Christensen-Dalsgaard J, Metcalfe TS, Sanderfer DT, Seader SE, Smith JC, Chaplin WJ (2014) Measurement of Acoustic Glitches in Solar-type Stars from Oscillation Frequencies Observed by Kepler. *ApJ*782(1):18, DOI 10.1088/0004-637X/782/1/18, 1312.4907
- McKeever JM, Basu S, Corsaro E (2019) The Helium Abundance of NGC 6791 from Modeling of Stellar Oscillations. *ApJ*874(2):180, DOI 10.3847/1538-4357/ab0c04, 1903.00497
- Meibom S, Torres G, Fressin F, Latham DW, Rowe JF, Ciardi DR, Bryson ST, Rogers LA, Henze CE, Janes K, Barnes SA, Marcy GW, Isaacson H, Fischer DA, Howell SB, Horch EP, Jenkins JM, Schuler SC, Crepp J (2013) The same frequency of planets inside and outside open clusters of stars. *Nature*499(7456):55–58, DOI 10.1038/nature12279, 1307.5842
- Michel E, Baglin A, Auvergne M, Catala C, Samadi R, Baudin F, Appourchaux T, Barban C, Weiss WW, Berthomieu G (2008) CoRoT Measures Solar-Like Oscillations and Granulation in Stars Hotter Than the Sun. *Science* 322(5901):558, DOI 10.1126/science.1163004, 0812.1267
- Michielsen M, Pedersen MG, Augustson KC, Mathis S, Aerts C (2019) Probing the shape of the mixing profile and of the thermal structure at the convective core boundary through asteroseismology. arXiv e-prints arXiv:1906.05304, 1906.05304
- Miglio A, Montalbán J, Carrier F, De Ridder J, Mosser B, Eggenberger P, Scuflaire R, Ventura P, D’Antona F, Noels A, Baglin A (2010) Evidence for a sharp structure variation inside a red-giant star. *A&A*520:L6, DOI 10.1051/0004-6361/201015442, 1009.1024
- Miglio A, Brogaard K, Stello D, Chaplin WJ, D’Antona F, Montalbán J, Basu S, Bressan A, Grundahl F, Pinsonneault M (2012) Asteroseismology of old open clusters with Kepler: direct estimate of the integrated red giant branch mass-loss in NGC 6791 and 6819. *MNRAS*419(3):2077–2088, DOI 10.1111/j.1365-2966.2011.19859.x, 1109.4376
- Miglio A, Chiappini C, Morel T, Barbieri M, Chaplin WJ, Girardi L, Montalbán J, Valentini M, Mosser B, Baudin F (2013) Galactic archaeology: mapping and dating stellar populations with asteroseismology of red-giant stars. *MNRAS*429(1):423–428, DOI 10.1093/mnras/sts345, 1211.0146
- Miglio A, Chaplin WJ, Brogaard K, Lund MN, Mosser B, Davies GR, Handberg R, Milone AP, Marino AF, Bossini D, Elsworth YP, Grundahl F, Arntoft T, Bedin LR, Campante TL, Jessen-Hansen J, Jones CD, Kuszlewicz JS, Malavolta L, Nascimbeni V, Sandquist EL (2016) Detection of solar-like oscillations in relics of the Milky Way: asteroseismology of K giants in

- M4 using data from the NASA K2 mission. *MNRAS*461(1):760–765, DOI 10.1093/mnras/stw1555, 1606.02115
- Milone AP, Bedin LR, Piotto G, Anderson J (2009) Multiple stellar populations in Magellanic Cloud clusters. I. An ordinary feature for intermediate age globulars in the LMC? *A&A*497(3):755–771, DOI 10.1051/0004-6361/200810870, 0810.2558
- Milone AP, Piotto G, Bedin LR, Aparicio A, Anderson J, Sarajedini A, Marino AF, Moretti A, Davies MB, Chaboyer B, Dotter A, Hempel M, Marín-Franch A, Majewski S, Paust NEQ, Reid IN, Rosenberg A, Siegel M (2012) The ACS survey of Galactic globular clusters. XII. Photometric binaries along the main sequence. *A&A*540:A16, DOI 10.1051/0004-6361/201016384, 1111.0552
- Milone AP, Marino AF, D’Antona F, Bedin LR, Piotto G, Jerjen H, Anderson J, Dotter A, di Criscienzo M, Lagioia EP (2017) Multiple stellar populations in Magellanic Cloud clusters - V. The split main sequence of the young cluster NGC 1866. *MNRAS*465(4):4363–4374, DOI 10.1093/mnras/stw2965, 1611.06725
- Milone AP, Marino AF, Renzini A, D’Antona F, Anderson J, Barbuy B, Bedin LR, Bellini A, Brown TM, Cassisi S (2018) The Hubble Space Telescope UV legacy survey of galactic globular clusters - XVI. The helium abundance of multiple populations. *MNRAS*481(4):5098–5122, DOI 10.1093/mnras/sty2573, 1809.05006
- Moe M, Di Stefano R (2017) Mind Your Ps and Qs: The Interrelation between Period (P) and Mass-ratio (Q) Distributions of Binary Stars. *ApJS*230(2):15, DOI 10.3847/1538-4365/aa6fb6, 1606.05347
- Montalbán J, Miglio A, Noels A, Dupret MA, Scuflaire R, Ventura P (2013) Testing Convective-core Overshooting Using Period Spacings of Dipole Modes in Red Giants. *ApJ*766(2):118, DOI 10.1088/0004-637X/766/2/118, 1302.3173
- Monteiro MJPF, Christensen-Dalsgaard J, Thompson MJ (1994) Seismic study of overshoot at the base of the solar convective envelope. *A&A*283(1):247–262
- Montenegro K, Minniti D, Alonso-García J, Hempel M, Saito RK, Beers TC, Brown D (2019) VVV Survey of Blue Horizontal Branch Stars in the Bulge-Halo Transition Region of the Milky Way. *ApJ*872(2):206, DOI 10.3847/1538-4357/aae379, 1810.00316
- Mosser B, Belkacem K, Goupil MJ, Miglio A, Morel T, Barban C, Baudin F, Hekker S, Samadi R, De Ridder J (2010) Red-giant seismic properties analyzed with CoRoT. *A&A*517:A22, DOI 10.1051/0004-6361/201014036, 1004.0449
- Mosser B, Barban C, Montalbán J, Beck PG, Miglio A, Belkacem K, Goupil MJ, Hekker S, De Ridder J, Dupret MA (2011) Mixed modes in red-giant stars observed with CoRoT. *A&A*532:A86, DOI 10.1051/0004-6361/201116825, 1105.6113
- Mosser B, Goupil MJ, Belkacem K, Marques JP, Beck PG, Bloemen S, De Ridder J, Barban C, Deheuvels S, Elsworth Y (2012) Spin down of the core

- rotation in red giants. *A&A*548:A10, DOI 10.1051/0004-6361/201220106, 1209.3336
- Mosser B, Dziembowski WA, Belkacem K, Goupil MJ, Michel E, Samadi R, Soszyński I, Vrad M, Elsworth Y, Hekker S, Mathur S (2013) Period-luminosity relations in evolved red giants explained by solar-like oscillations. *A&A*559:A137, DOI 10.1051/0004-6361/201322243, 1310.0839
- Mosser B, Belkacem K, Pinçon C, Takata M, Vrad M, Barban C, Goupil MJ, Kallinger T, Samadi R (2017a) Dipole modes with depressed amplitudes in red giants are mixed modes. *A&A*598:A62, DOI 10.1051/0004-6361/201629494, 1610.03872
- Mosser B, Pinçon C, Belkacem K, Takata M, Vrad M (2017b) Period spacings in red giants. III. Coupling factors of mixed modes. *A&A*600:A1, DOI 10.1051/0004-6361/201630053, 1612.08453
- Mosser B, Gehan C, Belkacem K, Samadi R, Michel E, Goupil MJ (2018) Period spacings in red giants IV: Toward a complete description of the mixed-mode pattern. arXiv e-prints arXiv:1807.08301, 1807.08301
- Mosser B, Michel E, Samadi R, Miglio A, Davies GR, Girardi L, Goupil MJ (2019) Seismic performance. *A&A*622:A76, DOI 10.1051/0004-6361/201834607, 1812.11906
- Myeong GC, Evans NW, Belokurov V, Sanders JL, Koposov SE (2018) The Sausage Globular Clusters. *ApJ*863(2):L28, DOI 10.3847/2041-8213/aad7f7, 1805.00453
- Myeong GC, Vasiliev E, Iorio G, Evans NW, Belokurov V (2019) Evidence for Two Early Accretion Events That Built the Milky Way Stellar Halo. arXiv:190403185 arXiv:1904.03185, 1904.03185
- Nascimbeni V, Bedin LR, Piotto G, De Marchi F, Rich RM (2012) An HST search for planets in the lower main sequence of the globular cluster NGC 6397. *A&A*541:A144, DOI 10.1051/0004-6361/201118655, 1202.5112
- Nataf DM, Gonzalez OA, Casagrande L, Zasowski G, Wegg C, Wolf C, Kunder A, Alonso-Garcia J, Minniti D, Rejkuba M, Saito RK, Valenti E, Zoccali M, Poleski R, Pietrzyński G, Skowron J, Soszyński I, Szymański MK, Udalski A, Ulaczyk K, Wyrzykowski L (2016) Interstellar extinction curve variations towards the inner Milky Way: a challenge to observational cosmology. *MNRAS*456(3):2692–2706, DOI 10.1093/mnras/stv2843, 1510.01321
- Nielsen MB, Schunker H, Gizon L, Schou J, Ball WH (2017) Limits on radial differential rotation in Sun-like stars from parametric fits to oscillation power spectra. *A&A*603:A6, DOI 10.1051/0004-6361/201730896, 1705.10517
- Notsu Y, Maehara H, Honda S, Hawley SL, Davenport JRA, Namekata K, Notsu S, Ikuta K, Nogami D, Shibata K (2019) Do Kepler Superflare Stars Really Include Slowly Rotating Sun-like Stars?—Results Using APO 3.5 m Telescope Spectroscopic Observations and Gaia-DR2 Data. *ApJ*876(1):58, DOI 10.3847/1538-4357/ab14e6, 1904.00142
- Noyola E, Gebhardt K, Bergmann M (2008) Gemini and Hubble Space Telescope Evidence for an Intermediate-Mass Black Hole in ω Centauri. *ApJ*676:1008-1015, DOI 10.1086/529002, 0801.2782

- Origlia L, Rood RT, Fabbri S, Ferraro FR, Fusi Pecci F, Rich RM, Dalessandro E (2010) Dust is Forming Along the Red Giant Branch of 47 Tuc. *ApJ*718(1):522–526, DOI 10.1088/0004-637X/718/1/522, 1005.4618
- Parada J, Richer H, Heyl J, Kalirai J, Goldsbury R (2016) Dynamical Estimate of Post-main-sequence Stellar Masses in 47 Tucanae. *ApJ*826(1):88, DOI 10.3847/0004-637X/826/1/88, 1605.05740
- Pasquini L, Biazzo K, Bonifacio P, Randich S, Bedin LR (2008) Solar twins in M 67. *A&A*489:677–684, DOI 10.1051/0004-6361/200809714, 0807.0092
- Pasquini L, Mauas P, Käufel HU, Cacciari C (2011) Measuring helium abundance difference in giants of NGC 2808. *A&A*531:A35, DOI 10.1051/0004-6361/201116592, 1105.0346
- Paunzen E, Bernhard K, Hümmerich S, Janík J, Semenko EA, Yakunin IA (2019) Search for stellar spots in field blue horizontal-branch stars. *A&A*622:A77, DOI 10.1051/0004-6361/201833098, 1812.07543
- Pinçon C, Belkacem K, Goupil MJ, Marques JP (2017) Can plume-induced internal gravity waves regulate the core rotation of subgiant stars? *A&A*605:A31, DOI 10.1051/0004-6361/201730998, 1705.10101
- Piotto G, Villanova S, Bedin LR, Gratton R, Cassisi S, Momany Y, Recio-Blanco A, Lucatello S, Anderson J, King IR, Pietrinferni A, Carraro G (2005) Metallicities on the Double Main Sequence of ω Centauri Imply Large Helium Enhancement. *ApJ*621(2):777–784, DOI 10.1086/427796, [astro-ph/0412016](#)
- Pu B, Wu Y (2015) Spacing of Kepler Planets: Sculpting by Dynamical Instability. *ApJ*807(1):44, DOI 10.1088/0004-637X/807/1/44, 1502.05449
- Queiroz ABA, Anders F, Chiappini C, Khalatyan A, Santiago BX, Steinmetz M, Valentini M, Miglio A, Bossini D, Barbey B, Minchev I, Minniti D, García Hernández DA, Schultheis M, Beaton RL, Beers TC, Bizyaev D, Brownstein JR, Cunha K, Fernández-Trincado JG, Frinchaboy PM, Lane RR, Majewski SR, Nataf D, Nitschelm C, Pan K, Roman-Lopes A, Sobeck JS, Stringfellow G, Zamora O (2020) From the bulge to the outer disc: StarHorse stellar parameters, distances, and extinctions for stars in APOGEE DR16 and other spectroscopic surveys. *A&A*638:A76, DOI 10.1051/0004-6361/201937364, 1912.09778
- Reese DR, Marques JP, Goupil MJ, Thompson MJ, Deheuvels S (2012) Estimating stellar mean density through seismic inversions. *A&A*539:A63, DOI 10.1051/0004-6361/201118156, 1201.1844
- Rey-Raposo R, Read JI (2018) The alignment is in their stars: on the spin-alignment of stars in star clusters. *MNRAS*481(1):L16–L20, DOI 10.1093/mnrasl/sly150, 1808.02830
- Romano D, Matteucci F, Tosi M, Pancino E, Bellazzini M, Ferraro FR, Limongi M, Sollima A (2007) The chemical evolution of Omega Centauri's progenitor system. *MNRAS*376(1):405–415, DOI 10.1111/j.1365-2966.2007.11446.x, [astro-ph/0701162](#)
- Roxburgh IW (2002) Background to the Eddington mission. In: Battrick B, Favata F, Roxburgh IW, Galadi D (eds) *Stellar Structure and Habitable Planet Finding*, ESA Special Publication, vol 485, pp 11–15

- Roxburgh IW, Vorontsov SV (2002) Inversion for the structure of a star of $1.45 M_{\text{solar}}$ using the internal phase shift. In: Battrick B, Favata F, Roxburgh IW, Galadi D (eds) *Stellar Structure and Habitable Planet Finding*, ESA Special Publication, vol 485, pp 341–343
- Royer F, Zorec J, Gómez AE (2007) Rotational velocities of A-type stars. III. Velocity distributions. *A&A*463(2):671–682, DOI 10.1051/0004-6361:20065224, [astro-ph/0610785](#)
- Saha A, Vivas AK, Olszewski EW, Smith V, Olsen K, Blum R, Valdes F, Claver J, Calamida A, Walker AR, Matheson T, Narayan G, Soraisam M, Cunha K, Axelrod T, Bloom JS, Cenko SB, Frye B, Juric M, Kaleida C, Kunder A, Miller A, Nidever D, Ridgway S (2019) Mapping the Interstellar Reddening and Extinction toward Baade’s Window Using Minimum Light Colors of ab-type RR Lyrae Stars: Revelations from the De-reddened Color-Magnitude Diagrams. *ApJ*874(1):30, DOI 10.3847/1538-4357/ab07ba, [1902.05637](#)
- Salabert D, Régulo C, Pérez Hernández F, García RA (2018) Frequency dependence of p-mode frequency shifts induced by magnetic activity in Kepler solar-like stars. *A&A*611:A84, DOI 10.1051/0004-6361/201731714, [1712.05691](#)
- Salaris M (2016) Age dating of old field stars: Challenges from the uncertain efficiency of atomic diffusion. *AN* 337(8-9):805, DOI 10.1002/asna.201612375
- Salaris M, Cassisi S (2017) Chemical element transport in stellar evolution models. *Royal Society Open Science* 4(8):170192, DOI 10.1098/rsos.170192, [1707.07454](#)
- Salaris M, Weiss A, Ferguson JW, Fusilier DJ (2006) On the Primordial Scenario for Abundance Variations within Globular Clusters: The Isochrone Test. *ApJ*645(2):1131–1137, DOI 10.1086/504520, [astro-ph/0604137](#)
- Salaris M, Cassisi S, Pietrinferni A (2016) On the red giant branch mass loss in 47 Tucanae: Constraints from the horizontal branch morphology. *A&A*590:A64, DOI 10.1051/0004-6361/201628181, [1604.02874](#)
- Samadi R, Deru A, Reese D, Marchiori V, Grolleau E, Green JJ, Pertenais M, Lebreton Y, Deheuvels S, Mosser B (2019) The PLATO Solar-like Light-curve Simulator. A tool to generate realistic stellar light-curves with instrumental effects representative of the PLATO mission. *A&A*624:A117, DOI 10.1051/0004-6361/201834822, [1903.02747](#)
- Santos ARG, Campante TL, Chaplin WJ, Cunha MS, Lund MN, Kiefer R, Salabert D, García RA, Davies GR, Elsworth Y, Howe R (2018) Signatures of Magnetic Activity in the Seismic Data of Solar-type Stars Observed by Kepler. *ApJS*237(1):17, DOI 10.3847/1538-4365/aac9b6, [1806.00136](#)
- Smiljanic R, Romano D, Bragaglia A, Donati P, Magrini L, Friel E, Jacobson H, Randich S, Ventura P, Lind K (2016) The Gaia-ESO Survey: Sodium and aluminium abundances in giants and dwarfs. Implications for stellar and Galactic chemical evolution. *A&A*589:A115, DOI 10.1051/0004-6361/201528014, [1602.03289](#)
- Soszyński I, Udalski A, Kubiak M, Szymański M, Pietrzyński G, Zeburń K, Szweczyk O, Wyrzykowski L (2004) The Optical Gravitational Lensing Ex-

- periment. Small Amplitude Variable Red Giants in the Magellanic Clouds. *Acta Astron.*54:129–152, [astro-ph/0407057](#)
- Soszyński I, Dziembowski WA, Udalski A, Kubiak M, Szymański MK, Pietrzyński G, Wyrzykowski L, Szewczyk O, Ulaczyk K (2007) The Optical Gravitational Lensing Experiment. Period–Luminosity Relations of Variable Red Giant Stars. *Acta Astron.*57:201–225, [0710.2780](#)
- Sousa SG, Adibekyan V, Santos NC, Mortier A, Barros SCC, Delgado-Mena E, Demangeon O, Israelian G, Faria JP, Figueira P (2019) The metallicity–period–mass diagram of low-mass exoplanets. *MNRAS*485(3):3981–3990, DOI [10.1093/mnras/stz664](#), [1903.04937](#)
- Souto D, Cunha K, Smith VV, Allende Prieto C, García-Hernández DA, Pinsonneault M, Holzer P, Frinchaboy P, Holtzman J, Johnson JA, Jönsson H, Majewski SR, Shetrone M, Sobeck J, Stringfellow G, Teske J, Zamora O, Zasowski G, Carrera R, Stassun K, Fernandez-Trincado JG, Villanova S, Minniti D, Santana F (2018) Chemical Abundances of Main-sequence, Turnoff, Subgiant, and Red Giant Stars from APOGEE Spectra. I. Signatures of Diffusion in the Open Cluster M67. *ApJ*857(1):14, DOI [10.3847/1538-4357/aab612](#), [1803.04461](#)
- Stello D, Cantiello M, Fuller J, Huber D, García RA, Bedding TR, Bildsten L, Silva Aguirre V (2016) A prevalence of dynamo-generated magnetic fields in the cores of intermediate-mass stars. *Nature*529(7586):364–367, DOI [10.1038/nature16171](#), [1601.00004](#)
- Strader J, Chomiuk L, MacCarone TJ, Miller-Jones JCA, Seth AC (2012) Two stellar-mass black holes in the globular cluster M22. *Nature*490:71–73, DOI [10.1038/nature11490](#), [1210.0901](#)
- Strom SE, Wolf SC, Dror DHA (2005) B Star Rotational Velocities in η and χ Persei: A Probe of Initial Conditions during the Star Formation Epoch? *AJ*129(2):809–828, DOI [10.1086/426748](#), [astro-ph/0410337](#)
- Szigeti L, Mészáros S, Smith VV, Cunha K, Lagarde N, Charbonnel C, García-Hernández DA, Shetrone M, Pinsonneault M, Allende Prieto C, Fernández-Trincado JG, Kovács J, Villanova S (2018) $^{12}\text{C}/^{13}\text{C}$ isotopic ratios in red-giant stars of the open cluster NGC 6791. *MNRAS*474(4):4810–4817, DOI [10.1093/mnras/stx3027](#), [1711.08183](#)
- Tabur V, Bedding TR, Kiss LL, Giles T, Derekas A, Moon TT (2010) Period–luminosity relations of pulsating M giants in the solar neighbourhood and the Magellanic Clouds. *MNRAS*409(2):777–788, DOI [10.1111/j.1365-2966.2010.17341.x](#), [1007.2974](#)
- Tailo M, Milone AP, Marino AF, D’Antona F, Lagioia EP, Cordoni G (2019) Mass Loss of Different Stellar Populations in Globular Clusters: The Case of M4. *ApJ*873(2):123, DOI [10.3847/1538-4357/ab05cc](#), [1902.03803](#)
- Takata M (2016) Asymptotic analysis of dipolar mixed modes of oscillations in red giant stars. *PASJ*68(6):109, DOI [10.1093/pasj/psw104](#)
- Takayama M, Saio H, Ita Y (2013) On the pulsation modes of OGLE small amplitude red giant variables in the LMC. *MNRAS*431(4):3189–3195, DOI [10.1093/mnras/stt398](#), [1303.7059](#)

- Thomas AEL, Chaplin WJ, Davies GR, Howe R, Santos ARG, Elsworth Y, Miglio A, Campante T, Cunha MS (2019) Asteroseismic constraints on active latitudes of solar-type stars: HD 173701 has active bands at higher latitudes than the Sun. *MNRAS*485(3):3857–3868, DOI 10.1093/mnras/stz672, 1903.04998
- Thompson MJ, Toomre J, Anderson ER, Antia HM, Berthomieu G, Burdonclay D, Chitre SM, Christensen-Dalsgaard J, Corbard T, DeRosa M, Genovese CR, Gough DO, Haber DA, Harvey JW, Hill F, Howe R, Korzennik SG, Kosovichev AG, Leibacher JW, Pijpers FP, Provost J, Rhodes EJ, Schou J, Sekii T, Stark PB, Wilson PR (1996) Differential rotation and dynamics of the solar interior. *Science* 272(5266):1300–1305, DOI 10.1126/science.272.5266.1300, URL <https://science.sciencemag.org/content/272/5266/1300>, <https://science.sciencemag.org/content/272/5266/1300.full.pdf>
- Tolstoy E, Hill V, Tosi M (2009) Star-Formation Histories, Abundances, and Kinematics of Dwarf Galaxies in the Local Group. *ARA&A*47(1):371–425, DOI 10.1146/annurev-astro-082708-101650, 0904.4505
- Tremou E, Strader J, Chomiuk L, Shishkovsky L, Maccarone TJ, Miller-Jones JCA, Tudor V, Heinke CO, Sivakoff GR, Seth AC, Noyola E (2018) The MAVERIC Survey: Still No Evidence for Accreting Intermediate-mass Black Holes in Globular Clusters. ArXiv:180600259 1806.00259
- Verma K, Raodeo K, Basu S, Silva Aguirre V, Mazumdar A, Mosumgaard JR, Lund MN, Ranadive P (2019) Helium abundance in a sample of cool stars: measurements from asteroseismology. *MNRAS*483(4):4678–4694, DOI 10.1093/mnras/sty3374, 1812.02751
- Villanova S, Geisler D, Piotto G, Gratton RG (2012) The Helium Content of Globular Clusters: NGC 6121 (M4). *ApJ*748(1):62, DOI 10.1088/0004-637X/748/1/62, 1201.3241
- Villanova S, Geisler D, Gratton RG, Cassisi S (2014) The Metallicity Spread and the Age-Metallicity Relation of ω Centauri. *ApJ*791(2):107, DOI 10.1088/0004-637X/791/2/107, 1406.5069
- Vorontsov SV (1988) A Search of the Effects of Magnetic Field in the Solar 5-Minute Oscillations. In: Christensen-Dalsgaard J, Frandsen S (eds) *Advances in Helio- and Asteroseismology*, IAU Symposium, vol 123, p 151
- Vrard M, Mosser B, Samadi R (2016) Period spacings in red giants. II. Automated measurement. *A&A*588:A87, DOI 10.1051/0004-6361/201527259, 1602.04940
- Weldrake DTF, Sackett PD, Bridges TJ (2008) The Frequency of Large-Radius Hot and Very Hot Jupiters in ω Centauri. *ApJ*674(2):1117–1129, DOI 10.1086/524917, 0710.3461
- Yong D, Casagrande L, Venn KA, Chené AN, Keown J, Malo L, Martioli E, Alves-Brito A, Asplund M, Dotter A, Martell SL, Meléndez J, Schlesinger KJ (2016) GRACES observations of young $[\alpha/\text{Fe}]$ -rich stars. *MNRAS*459(1):487–495, DOI 10.1093/mnras/stw676, 1603.07034
- Yu J, Huber D, Bedding TR, Stello D, Hon M, Murphy SJ, Khanna S (2018) Asteroseismology of 16,000 Kepler Red Giants: Global Oscillation Param-

- eters, Masses, and Radii. *ApJS*236(2):42, DOI 10.3847/1538-4365/aaaf74, 1802.04455
- Zhu W, Udalski A, Calchi Novati S, Chung SJ, Jung YK, Ryu YH, Shin IG, Gould A, Lee CU, Albrow MD (2017) Toward a Galactic Distribution of Planets. I. Methodology and Planet Sensitivities of the 2015 High-cadence Spitzer Microlens Sample. *AJ*154(5):210, DOI 10.3847/1538-3881/aa8ef1, 1701.05191
- Zoccali M, Vasquez S, Gonzalez OA, Valenti E, Rojas-Arriagada A, Minniti J, Rejkuba M, Minniti D, McWilliam A, Babusiaux C, Hill V, Renzini A (2017) The GIRAFFE Inner Bulge Survey (GIBS). III. Metallicity distributions and kinematics of 26 Galactic bulge fields. *A&A*599:A12, DOI 10.1051/0004-6361/201629805, 1610.09174
- Zocchi A, Gieles M, Hénault-Brunet V (2017) Radial anisotropy in ω Cen limiting the room for an intermediate-mass black hole. *MNRAS*468:4429–4440, DOI 10.1093/mnras/stx316, 1702.00725
- Zocchi A, Gieles M, Hénault-Brunet V (2019) The effect of stellar-mass black holes on the central kinematics of ω Cen: a cautionary tale for IMBH interpretations. *MNRAS*482:4713–4725, DOI 10.1093/mnras/sty1508, 1806.02157