

Cosmology from the two-dimensional renormalization group acting as the Ricci flow

Daniel Friedan*

*New High Energy Theory Center and Department of Physics and Astronomy,
Rutgers, The State University of New Jersey,
Piscataway, New Jersey 08854-8019 U.S.A. and*

Science Institute, The University of Iceland, Reykjavik, Iceland

(Dated: September 3, 2019)

The two-dimensional renormalization group acting as the Ricci flow $\Lambda \frac{\partial}{\partial \Lambda} g_{\mu\nu} = R_{\mu\nu}$ produces a specific 1+3 dimensional space-time metric which describes an expanding universe that starts with a big bang $a \sim t^{1/\sqrt{3}}$ then decelerates until $z = 0.2$ then accelerates until ending at $t_{\max} = 1.6 t_H$ with a big blowup $a \sim (t_{\max} - t)^{-1/\sqrt{3}}$. The only free parameters are the overall time scale and the value of the present time t_0 . These are fixed by the Hubble constant $H_0 = t_H^{-1}$ and the present deceleration parameter $q(t_0)$. This crude calculation of cosmology omits all but the gravitational field. The only energy-momentum is purely gravitational dark matter and energy. This is a preliminary exploration towards a specific, comprehensive, testable calculation of cosmology from a fundamental theory in which physics is produced by a quantum version of the two-dimensional renormalization group.

The renormalization group of the two-dimensional general nonlinear model acts at leading order as the Ricci flow [1–3]

$$\Lambda \frac{\partial}{\partial \Lambda} g_{\mu\nu} = R_{\mu\nu} \quad (1)$$

The field of the nonlinear model takes its values in a Riemannian manifold M . The Riemannian metric $g_{\mu\nu}$ encodes the couplings of the model. The beta-function at leading order is the Ricci tensor $R_{\mu\nu}$. The renormalization group drives $g_{\mu\nu}$ to a fixed point (modulo at most a finite number of relevant parameters). The fixed points are the solutions of

$$R_{\mu\nu} = \nabla_\mu v_\nu + \nabla_\nu v_\mu \quad (2)$$

where $v^\mu(x)\partial_\mu$ is a vector field on M . A vector field is an infinitesimal reparametrization $\dot{x}^\mu = v^\mu(x)$ of M . This is a field redefinition in the nonlinear model equivalent to the redundant perturbation $\dot{g}_{\mu\nu} = \nabla_\mu v_\nu + \nabla_\nu v_\mu$. Thus the fixed point equation (2) expresses physical 2-d scale invariance¹.

The 2-d quantum field theory is manifestly well defined when $g_{\mu\nu}$ has euclidean signature. Assume that M is a four-dimensional Riemannian manifold of the form $I \times S^3$ where I is a real interval and S^3 is the unit 3-sphere. Assume $\text{SO}(4)$ invariance. Parametrize $I \times S^3$ as a spherical shell in \mathbb{R}^4 . The flat euclidean metric is $\delta_{\mu\nu} dx^\mu dx^\nu = dr^2 + r^2 ds_{S^3}^2$ where $ds_{S^3}^2$ is the round metric on the unit 3-sphere. The general $\text{SO}(4)$ -invariant metric has the form

¹ The general nonlinear model (also called the *nonlinear sigma model*) was constructed in [1–3] in $2+\epsilon$ dimensions with fixed point equation $R_{\mu\nu} - \epsilon g_{\mu\nu} = \nabla_\mu v_\nu + \nabla_\nu v_\mu$. The solutions were called quasi-Einstein metrics. The Ricci flow was introduced in Mathematics in [4]. The solutions of $R_{\mu\nu} - \epsilon g_{\mu\nu} = \nabla_\mu v_\nu + \nabla_\nu v_\mu$ have been called *Ricci solitons* or *Ricci flow solitons* in the Mathematics literature.

$g_{\mu\nu}dx^\mu dx^\nu = F_1(r)^2 dr^2 + F_2(r)^2 ds_{S^3}^2$. The metric can be made conformally flat

$$g_{\mu\nu}dx^\mu dx^\nu = a^2 (d\tau^2 + ds_{S^3}^2) \quad a = e^{f(\tau)} \quad (3)$$

by reparametrizing $r \rightarrow \tau(r)$ with $dr/d\tau = F_2/F_1$. The general SO(4)-invariant vector field is $v^\tau(\tau)\partial_\tau$. Then [5]

$$\begin{aligned} R_{\mu\nu}dx^\mu dx^\nu &= (-3\partial_\tau f_\tau) d\tau^2 + (-\partial_\tau f_\tau + 2 - 2f_\tau^2) ds_{S^3}^2 \\ (\nabla_\mu v_\nu + \nabla_\nu v_\mu)dx^\mu dx^\nu &= (2\partial_\tau v_\tau - 2v_\tau f_\tau) d\tau^2 + (2v_\tau f_\tau) ds_{S^3}^2 \\ f_\tau &= \partial_\tau f \quad v_\tau = g_{\tau\tau}v^\tau = a^2 v^\tau \end{aligned} \quad (4)$$

The fixed point equation (2) becomes the ordinary differential equation

$$\partial_\tau f_\tau = -2f_\tau v_\tau - 2f_\tau^2 + 2 \quad \partial_\tau v_\tau = 4f_\tau v_\tau + 3f_\tau^2 - 3 \quad (5)$$

We shall see below that there is an essentially unique solution of (5) which analytically continues in τ to real time $T = i^{-1}\tau$

$$ds^2 = a^2 (-dT^2 + ds_{S^3}^2) \quad (6)$$

The real-time ode is

$$\begin{aligned} \partial_T f_T &= -2f_T v_T - 2f_T^2 - 2 \quad \partial_T v_T = 4f_T v_T + 3f_T^2 + 3 \\ \partial_T &= i\partial_\tau \quad f_T = if_\tau = \partial_T a/a \quad v_T = iv_\tau \end{aligned} \quad (7)$$

The solution is

$$f_T = \frac{\cos 2T + \sqrt{3}}{\sin 2T} \quad v_T = \frac{-\sqrt{3}}{\sin 2T} \quad a = t'_0 \sin^{1+\nu} T \cos^{-\nu} T \quad T \in (0, \pi/2) \quad (8)$$

with $\nu = \sqrt{3}/2 - 1/2 = 0.3660\dots$. The only free parameter is the overall time scale t'_0 . In co-moving time t

$$\begin{aligned} ds^2 &= -dt^2 + a^2 ds_{S^3}^2 \quad dt = adT \\ \frac{t}{t'_0} &= \int_0^T \sin^{1+\nu} T' \cos^{-\nu} T' dT' = \frac{1}{2} B_{\sin^2 T} \left(\frac{2+\nu}{2}, \frac{1-\nu}{2} \right) \\ t \in (0, t_{\max}) \quad \frac{t_{\max}}{t'_0} &= \frac{1}{2} B \left(\frac{2+\nu}{2}, \frac{1-\nu}{2} \right) = 1.470\dots \end{aligned} \quad (9)$$

$B(p, q)$ is the Euler beta function, $B_x(p, q)$ the incomplete beta function. At the limits

$$\begin{aligned} T \rightarrow 0 \quad \frac{t}{t'_0} &\rightarrow \frac{T^{2+\nu}}{2+\nu} \quad a \rightarrow t'_0 (2+\nu)^{1/\sqrt{3}} \left(\frac{t}{t'_0} \right)^{1/\sqrt{3}} \\ T \rightarrow \frac{\pi}{2} \quad \frac{t_{\max} - t}{t'_0} &\rightarrow \frac{\left(\frac{\pi}{2} - T \right)^{1-\nu}}{1-\nu} \quad a \rightarrow t'_0 (1-\nu)^{-1/\sqrt{3}} \left(\frac{t_{\max} - t}{t'_0} \right)^{-1/\sqrt{3}} \end{aligned} \quad (10)$$

This is an expanding universe which begins with a big bang $a \sim t^{0.577\dots}$ and ends with a big blowup $a \sim (t_{\max} - t)^{-0.577\dots}$. The Hubble parameter $H = \partial_t a/a = f_T/a$ is

$$H = \frac{1}{t'_0} (\cos^2 T + \nu) \sin^{-2-\nu} T \cos^{-1+\nu} T \quad (11)$$

The deceleration parameter $q = -a\partial_t^2 a/(\partial_t a)^2 = -\partial_T f_T/f_T^2$ is

$$q = \frac{2(\sqrt{3}\cos 2T + 1)}{(\cos 2T + \sqrt{3})^2} \quad T_{q=0} = \frac{1}{2} \arccos(-1/\sqrt{3}) = (0.6959\dots)\frac{\pi}{2} \quad (12)$$

The expansion decelerates ($q > 0$) until $T = T_{q=0}$ then accelerates ($q < 0$) until the end.

For a first estimate of the time scale t'_0 and the present time t_0 use the deceleration parameter at the present time $q_0 = q(T_0) \approx -0.6$ in equation (12) to obtain $T_0 = 0.77\pi/2$. Then equate the Hubble constant H_0 to $H(T_0)$ in (11) to obtain $t'_0 = 1.1t_H$ where $t_H = 1/H_0 \approx 1.4 \times 10^{10}y$. Then (9) gives $t_{\max} = 1.6t_H$, $t_0 = 0.73t_H$ and (8) gives $a_0 = a(t_0) = 1.5t_H$.

Einstein's equation

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = 8\pi G T_{\mu\nu} \quad (13)$$

is satisfied with energy-momentum tensor

$$T_{\mu\nu} = \frac{1}{8\pi G} (\nabla_\mu v_\nu + \nabla_\nu v_\mu - \nabla_\sigma v^\sigma g_{\mu\nu}) \quad (14)$$

which is that of a perfect fluid of density $\rho(t)$ and pressure $p(t)$

$$\begin{aligned} T_{\mu\nu} dx^\mu dx^\nu &= \rho(t) dt^2 + p(t) a^2 ds_{S^3}^2 \\ 8\pi G \rho(t) &= a^{-2} (3f_T^2 + 3) \quad 8\pi G p(t) = a^{-2} (4f_T v_T + 3f_T^2 + 3) \end{aligned} \quad (15)$$

The energy-momentum is purely gravitational, tautologically dark. The equation-of-state parameter $w(t) = p/\rho$ and the density parameter $\Omega(t) = 8\pi G \rho/3H^2 = 1 + 1/f_T^2$ are

$$w = \frac{\cos 2T}{3\cos 2T + 2\sqrt{3}} \quad \Omega = 1 + \left(\frac{\sin 2T}{\cos 2T + \sqrt{3}} \right)^2 \quad (16)$$

The estimate $T_0 = 0.77\pi/2$ gives present values $w_0 = -0.6$, $\Omega_0 = 1.5$. The cosmological parameters are graphed as functions of t in Figure 1. Past values for a selection of redshifts $z = a_0/a - 1$ are listed in Table I.

We analyze the real-time ode (7) by adapting some of the methods that were used in [7] to analyze the ode analogous to (5) for the euclidean-signature $SO(3)$ -invariant fixed point equation in three dimensions. The real-time ode (7) has time-reflection symmetry $(T, f_T, v_T) \rightarrow (-T, -f_T, -v_T)$ and constant of motion

$$C = \frac{1}{a^2} \left(\frac{2}{3} v_T^2 + 2f_T v_T + f_T^2 + 1 \right) \quad \partial_T C = 0 \quad (17)$$

Changing variables to $h_\pm = f_T + (1 \pm 1/\sqrt{3})v_T$ the constant of motion and ode become

$$h_+ h_- + 1 = Ca^2 \quad \partial_T h_\pm = -(1 + h_\pm^2) + \lambda_\pm (h_+ h_- + 1) \quad \lambda_\pm = 2 \pm \sqrt{3} \quad (18)$$




FIG. 1. The cosmological parameters as functions of the co-moving time t . The present is t_0 .

TABLE I. Cosmological parameters at selected values of z

	z	H/H_0	q	w	Ω
$z \gg 1$	$\gg 1$	$0.75 z^{1.7}$	0.73	0.16	1.0
	1000	1.2×10^5	0.73	0.16	1.0
	100	2.2×10^3	0.73	0.15	1.0
	10	48	0.74	0.15	1.0
$T = \frac{1}{2}T_0$	1.7	4.1	0.74	0.08	1.2
	1.0	2.4	0.69	0.02	1.3
$q = 0$	0.2	1.1	0	-0.3	1.5
$t = t_0$	0	1	-0.6	-0.6	1.5

The cosmological solution (8) is $C = 0$, $h_- = \cot T$, $h_+ = -\tan T$.

The phase portrait of the ode is shown in Figure 2. Every $C \neq 0$ trajectory asymptotes to a $C = 0$ trajectory. The separatrix S and its time-reflection S' each does so at one end; all the other $C \neq 0$ trajectories do so at both ends. The asymptotic behavior is one of

$$\begin{aligned}
 T \rightarrow 0^\pm \quad h_- &\rightarrow \frac{1}{T} & h_+ &\rightarrow -T - \frac{1}{3}T^3 + cT(T^2)^{1+\nu} & \frac{c}{C} &> 0 \\
 & & h_- &\rightarrow c'T(T^2)^{-\nu} & \frac{c'}{C} &> 0
 \end{aligned} \tag{19}$$

The presence of an irrational power of T , namely $(T^2)^{1+\nu}$ or $(T^2)^{-\nu}$, implies that such a solution cannot be the analytic continuation of a solution in imaginary time $\tau = iT$.




FIG. 2. Phase portrait of the real-time ode (7,18) in terms of (f_T, v_T) on the left, (h_-, h_+) on the right. The trajectories are the solutions. The arrows point in the direction of increasing T . The interval between points in the trajectories is $\Delta T = 0.01$. The universe is expanding when $f_T = \partial_t a$ is positive. The expansion is accelerating when f_T is increasing. The two red trajectories are the $C = 0$ solutions. The one in the lower right quadrant labeled $C=0$ is the cosmological solution (8). Its time reflection is labeled $C=0'$. The separatrix is labeled S . Its time-reflection is labeled S' .

The trajectories that start from $h_- = \infty$, $h_+ = 0$ at $T = 0$ are characterized by the numerical invariant

$$c = (h_+ h_- + 1)(h_-^2)^{1+\nu} e^{\int_0^T \frac{2 dT'}{h_-(T')}} = \lim_{T \rightarrow 0} (h_+ h_- + 1)(h_-^2)^{1+\nu} \quad (20)$$

The separatrix S has $c_S = 0.284\dots$. The $c < c_S$ trajectories — the trajectories below S — are all roughly realistic as cosmologies. They all start with a big bang $a \sim t^{1/\sqrt{3}}$ then decelerate then accelerate to a big blow-up $a \sim t^{-1/\sqrt{3}}$, all driven by purely gravitational dark energy-momentum. The gap between the $C=0$ cosmological solution and the separatrix S means that the cosmological solution (8) is stable against real-time perturbations. The zone of stability on the side $C > 0$ shrinks to zero as $T \rightarrow 0$.

The two-dimensional renormalization group was developed into a comprehensive fundamental theory of physics in which the 2-d renormalization group acts not on a space of classical space-time fields but rather on measures on that space, producing not a solution of the classical field equations but rather a measure — a functional integral — expressing a solution of the quantum field equations [8–10]. The present exploratory calculations of cosmology from the 2-d renormalization group will need to be connected to this fundamental theory. Assumptions such as $SO(4)$ space-time symmetry will need to be justified by properties of the 2-d renormalization group flow — by stability properties of the fixed point and/or as the dynamical result of a distinguished trajectory of the rg flow.

The analyticity assumption needs precise characterization and justification. The assumption here that $f_T(T)$ and $v_T(T)$ should be analytic in T singles out the $C = 0$ cosmological solution. Analyticity serves as a selection principle providing specificity. It should express a physical principle.

The next exploratory steps will add standard model fields to the calculation, then spatial fluctuations and quantum effects, hoping to find analytic solutions of 2-d renormalization group fixed point equations that capture more and more qualitative features of the cosmology of the real world, aiming to find a solution that can be tested in quantitative detail. This would provide support for the fundamental theory [8] in which a quantum version of the two-dimensional renormalization group acts mechanically to produce physics and would provide guidance for deriving real world cosmology from that fundamental theory.

This work was supported by the Rutgers New High Energy Theory Center and by the generosity of B. Weeks. I am grateful to the Mathematics Division of the Science Institute of the University of Iceland for its hospitality.

* dfriedan@gmail.com; www.physics.rutgers.edu/~friedan/

- [1] D. Friedan, Nonlinear Models in $2 + \epsilon$ Dimensions, Phys. Rev. Lett. **45**, 1057 (1980).
- [2] Daniel Harry Friedan, *Nonlinear Models in $2 + \epsilon$ Dimensions*, Tech. Rep. (Lawrence Berkeley Laboratory LBL-11517, 1980) U.C. Berkeley PhD thesis (1980).
- [3] Daniel Harry Friedan, Nonlinear Models in $2 + \epsilon$ Dimensions, Ann. Phys. **163**, 318 (1985), republication of [2].
- [4] Richard S. Hamilton, Three-manifolds with positive Ricci curvature, J. Differential Geom. **17**, 255 (1982).
- [5] Calculations are shown in the accompanying supplemental material which consists of a note *Calculations for "Cosmology ..."* and three SageMath [6] notebooks of numerical calculations. The supplemental material can be found at http://www.physics.rutgers.edu/pages/friedan/papers/2019/Cosmology_I_supplementary_material/ or at https://share.cocalc.com/share/6a7035ba-9879-4b05-b3d0-688b1309a21c/Cosmology_I_supplementary_material/?viewer=share/.
- [6] The Sage Developers, *SageMath, the Sage Mathematics Software System (Version 8.8)* (2019), <https://www.sagemath.org>.
- [7] Robert L. Bryant, Ricci flow solitons in dimension 3 with $SO(3)$ -symmetries, unpublished (2005), <https://math.duke.edu/~bryant/3DRotSymRicciSolitons.pdf>.
- [8] D. Friedan, A tentative theory of large distance physics, JHEP **10**, 063, arXiv:hep-th/0204131.
- [9] D. Friedan, A loop of $SU(2)$ gauge fields stable under the Yang-Mills flow, in *Perspectives in mathematics and physics: Essays dedicated to Isadore Singer's 85th birthday*, Surveys in Differential Geometry, Vol. 15, edited by T. Mrowka and S.-T. Yau (International Press of Boston, 2010) pp. 131–204, arXiv:1008.1189 [hep-th].
- [10] D. Friedan, A pragmatic approach to formal fundamental physics, (2018), arXiv:1810.09508 [hep-th].