

Interstellar comet C/2019 Q4 (Borisov)

Piotr Guzik^{1*}, Michał Drahus^{1*}, Krzysztof Rusek², Waław Waniak¹, Giacomo Cannizzaro^{3,4}, Inés Pastor-Marazuela^{5,6}

¹ Astronomical Observatory, Jagiellonian University, ul. Orła 171, 30-244 Kraków, Poland

² AGH University of Science and Technology, al. Mickiewicza 30, Kraków 30-059, Poland

³ SRON, Netherlands Institute for Space Research, Sorbonnelaan, 2, NL-3584CA Utrecht, the Netherlands

⁴ Department of Astrophysics/IMAPP, Radboud University, P.O. Box 9010, 6500 GL Nijmegen, the Netherlands

⁵ Anton Pannekoek Institute for Astronomy, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands

⁶ ASTRON, Netherlands Institute for Radio Astronomy, Oude Hoogeveensedijk 4, 7991 PD Dwingeloo, The Netherlands

* These authors contributed equally to this work.

Interstellar comets penetrating through the Solar System were anticipated for decades^{1,2}. The discovery of the asteroidal ‘Oumuamua by Pan-STARRS was therefore a huge surprise and puzzle. Furthermore, its physical properties turned out to be impossible to reconcile with Solar System objects³⁻⁵, which radically changed our view on interstellar minor bodies. Here, we report the identification and early characterization of a new interstellar object which has an evidently cometary appearance. The body was identified by our data mining code in publicly available astrometric data. The data clearly show significant systematic deviation from what is expected for a parabolic orbit and are consistent with an enormous orbital eccentricity of 3.16 ± 0.13 . Images taken on 10 and 13 September 2019 with the William Herschel Telescope and Gemini North Telescope show an extended coma and a faint, broad tail – the canonical signatures of cometary activity. The observed g’ and r’-band magnitudes imply the g’-r’ color index of 0.63 ± 0.02 , essentially the same as measured for the native Solar System comets. We have also estimated the ‘photometric’ nucleus diameter to be ~ 500 m. The nucleus is thus quite small and has a ~ 10% chance of experiencing

rotational disruption. Based on these early characteristics, and putting its hyperbolic orbit aside, C/2019 Q4 (Borisov) appears indistinguishable from the native Solar System comets. The discovery of this object indicates that interstellar comets might be common and creates a tremendous opportunity to study the first such object in great detail.

On 8 September 2019 at 04:15 UT, we were alerted by our software “Interstellar Crusher” (see Methods) of a new hyperbolic object. The body was discovered by Gennady Borisov on 30 August 2019 at 01:03 UT and received a provisional label gb00234 which was subsequently changed to C/2019 Q4 (Borisov)⁶. On 11 September 2019, publicly available astrometric data allowed us to confirm the hyperbolic orbit of this object at 15- σ significance. Using 129 positions obtained over a 12.42-day interval we found a preliminary osculating hyperbolic orbit consistent with the solution published by the Minor Planet Center around the same time⁶. Our solution is presented in Table 1 and the observers that contributed the data that we used for this computation are listed in Supplementary Table 1. For the most up-to-date orbital parameters readers are encouraged to visit the JPL Small Body Database Browser. The enormous eccentricity of 3.16 ± 0.13 , together with the perihelion distance of 1.96 ± 0.04 au, imply a hyperbolic excess speed of ~ 30 km s⁻¹. The body entered the Solar System from the direction $\sim 70^\circ$ off the Solar apex with the asymptotic radiant at RA = 02^h 08^m and Dec = 59.7° in the constellation of Cassiopeia. In Figure 1 we present astrometric residuals of C/2019 Q4 (Borisov) calculated for hyperbolic (left panel) and parabolic (right panel) orbital solutions. While a systematic trend is clearly visible for the parabolic solution, the residuals are much smaller and evenly distributed in the hyperbolic case. Since the body arrives from the direction off the ecliptic plane, the hyperbolic orbit cannot be explained by planetary perturbations inside the Solar System, and the residuals calculated for parabolic fit

are few orders of magnitude too large to be accounted to non-gravitational forces observed in Solar System comets.

We observed this object using the 4.2 m William Herschel Telescope (WHT) on La Palma and the 8.2 m Gemini North telescope at Maunakea. WHT data were obtained with the Auxiliary-port CAMera (ACAM) on 10 September 2019 at 05:38 UT and on 12 September 2019 at 05:47 UT (observation mid-points). The first set comprises 10 sidereal-tracked 60-sec exposures, of which 5 were obtained in g' and 5 in r' bands⁷, whereas the second set contains 40 sidereal-tracked 20-sec exposures, half in g' and half in r' . Gemini data were collected with the Gemini Multi-Object Spectrograph (GMOS-N) on 10 September 2019 at 14:57 UT (mid-point) with non-sidereal tracking and comprise four g' -band and four r' -band exposures taken with 60-sec integration time. The datasets were obtained at low elevation (22° to 31°) in morning twilight. At the time of the observations, the helio- and geocentric distances of C/2019 Q4 (Borisov) were equal to 2.8 and 3.4 au, respectively, and the phase angle was $\sim 15^\circ$. The images were corrected for overscan, bias and flatfield in the standard fashion, and then a global background level was subtracted from each frame.

In Figure 2 we show median stacked g' -band and r' -band images from Gemini, which have a slightly better signal-to-noise ratio than the WHT images. The images reveal an extended coma and a broad, short tail emanating in roughly antisolar direction. We see no clear difference in morphology in the two bands. The comet was also measured photometrically in each individual image from the two telescopes and two nights. We consistently used a 2.0-arcsec radius photometric aperture and determined the brightness against background stars available in the Sloan Digital Sky Survey photometric catalogue⁸. As a result, we obtained

dataset-averaged AB magnitudes $g' = 19.32 \pm 0.02$ and $r' = 18.69 \pm 0.02$ for the WHT observations on 10 September 2019 at 05:38 UT, $g' = 19.30 \pm 0.01$ and $r' = 18.69 \pm 0.01$ for the Gemini observations on 10 September 2019 at 14:57 UT, and $g' = 19.21 \pm 0.02$ and $r' = 18.56 \pm 0.02$ for the WHT observations on 13 September 2019 at 05:47 UT. Photometric magnitudes can be used to estimate the nucleus size. Following two independent approaches^{2,9} we have found that the nucleus of C/2019 Q4 (Borisov) is most likely ~ 0.5 km in diameter (see Methods). However, it should be noted that these approaches are inherently very uncertain. Our photometric measurements give a very consistent $g'-r'$ color index, with an averaged value of 0.63 ± 0.02 mag, i.e. slightly redder than the solar $g'-r' = 0.45 \pm 0.02$ mag¹⁰. Within the errors, the same color index is obtained for other photometric apertures as well (we made measurements for the apertures ranging from 1.0 to 10.0 arcsec in radius), implying a very uniform color of the coma. This, and the similarity of morphology in both bands, are both indicative of dust-dominated activity. Monte-Carlo modeling of the object's dust environment with our established code¹¹ has revealed the power-law particle size-distribution exponent of -3.5 ± 0.7 and a 40 m s^{-1} ejection velocity of $\beta = 1$ particles.

The dynamical properties of C/2019 Q4 (Borisov) make it clear that the body is the second-known (after 'Oumuamua³) interstellar minor body identified in the Solar System, whereas its morphology demonstrably shows it is the first certain case of an interstellar comet. Evidently, the extended coma and the broad tail stand in stark contrast with the purely asteroidal appearance of 'Oumuamua. The estimated nucleus size is small, but not uncommon among Solar System's long period comets⁹. Such small and active nuclei are prone to rapid rotation-period changes leading to rotational instability and disruption^{12,13}. Adopting the formalism of rotational disruption probability¹⁴ and assuming typical properties of Solar System comets, we have estimated the probability of rotational disruption of C/2019 Q4 (Borisov) during the

Solar System flyby to be of ~ 10%. We note that the measured color is perfectly consistent with the colors of the Solar System's long period comets¹⁵ and falls right in the center of the observed distribution (Figure 3). The same similarity can be noticed for the dust coma parameters¹¹. These facts are remarkable in and of themselves, and especially remarkable after 'Oumuamua, the multiple peculiarities of which^{3,4,5} prompted us to rethink our entire view of the nature of interstellar interlopers¹⁶. However, comet C/2019 Q4 (Borisov) is completely similar to the native Solar System's comets (putting its hyperbolic orbit aside).

Comet C/2019 Q4 (Borisov) was discovered on its way to perihelion (9 December 2019 at 2.0 au) and before the closest approach to Earth (28 December 2019 at 1.9 au), thus the overall visibility will be gradually improving. The body is destined for an intensive observing campaign lasting many months (Supplementary Figure 1) that will allow us to gain sensational insights into the physical properties of interstellar comets and exosolar planetary systems in general. This discovery is in line with the detection statistic of one interstellar object per year proposed by several authors^{17,18} after the discovery of 'Oumuamua, which is an order of magnitude higher than the most optimistic pre-'Oumuamua estimates¹⁹. It also shows that such bodies might be common and thus the former expectations^{1,2} were probably valid. More discoveries are expected in near future thanks to LSST.

Methods

Interstellar Crusher

“Interstellar Crusher” is a custom Python3 code working on Windows Subsystem for Linux (WSL). It operates on Bill Gray’s Find_Orb. While continuously monitoring astrometric positions reported on the Possible Comet Confirmation Page, it is performing a real-time search for hyperbolic orbits among newly discovered minor bodies. The detection of a possible interstellar object raises an alarm that is sent via an e-mail.

Orbit

To calculate the orbit we used Bill Gray’s Find_Orb, accommodating planetary perturbations. At the time of writing, 154 astrometric positions obtained at 22 different stations were publicly available. Three of those stations reported positions grossly inconsistent with rest of the data. We assumed that the data from those three stations contained some systematic errors and thus rejected them. Therefore, to compute the orbital parameters of C/2019 Q4 (Borisov) we used 129 measurements taken at 19 different stations. We computed both parabolic and hyperbolic, purely gravitational solutions and then plotted the residuals for both orbits.

Nucleus size estimation

The diameter D (in km) of a bare cometary nucleus can be calculated from²⁰:

$$\log D = 3.1236 - 0.5 \log (p_v) - 0.2 H,$$

Provided that its absolute magnitude (H) and albedo (p_v) are known. Interstellar comet C/2019 Q4 (Borisov) is already active, thus any measurement of its nucleus brightness is contaminated by coma signal. Boe et al.⁹ found that for active comets their absolute “nuclear

magnitudes” reported formerly by JPL are roughly 5.5 mag brighter than their true absolute magnitudes. Absolute “nuclear magnitudes” given by JPL were calculated based on “nuclear magnitudes” reported as a by-product of astrometric measurements. They were in fact representing an inner coma brightness (measured in an aperture of the order of the PSF). JPL reported those absolute nuclear magnitudes as Visual-band magnitudes. We assume that they were equivalent to V-band magnitudes.

The typical albedo of a cometary nucleus is ~ 0.04 , thus we adopted this value. To estimate the size of the nucleus of C/2019 Q4 (Borisov) we measured the brightness of the comet in a 1-arcsec radius photometric aperture which allowed us to estimate its absolute magnitude $H = 20.35$ (taking into account the 5.5 mag shift). Substituting this value to the equation above, we obtained the diameter of the order of ~ 550 meters.

Engelhardt et al.², based on the previous works^{21,22,23} gives an estimate of an apparent magnitude of a fresh long period active comet (or interstellar comet) if the entire sunlit hemisphere of the object is active:

$$V = 1.6 H + 6.2 \log (r) + 5 \log (\Delta) - 21.1,$$

which is equivalent to:

$$H = (V - 6.2 \log (r) - 5 \log (\Delta) + 21.1)/1.6$$

Using different photometric apertures, we found from our data $H = 20.86$ mag. This value indicates the nucleus diameter of ~ 450 meters, consistent with the estimation presented above.

One should be aware of some unavoidable shortcomings of these approaches. The former one is burdened with significant uncertainties originating from heterogeneity in the measurements of “nuclear magnitude”, the latter is based on the assumption that the cometary activity is

driven by water ice sublimation, which may or may not be true for C/2019 Q4 (Borisov).

Both are obtained under the assumed albedo of 0.04. Still, however, both gives consistent results, suggesting that the interstellar comet's nucleus is probably sub-kilometer sized.

References

1. Sekanina, Z. A probability of encounter with interstellar comets and the likelihood of their existence. *Icarus* **27**, 123–133 (1976).
2. Engelhardt, T. et al. An Observational Upper Limit on the Interstellar Number Density of Asteroids and Comets. *Astron. J.* **153**, 133 (2017).
3. Meech, K. J. et al. A brief visit from a red and extremely elongated interstellar asteroid. *Nature* **552**, 378–381 (2017).
4. Drahus, M. et al. Tumbling motion of 1I/Oumuamua and its implications for the body's distant past. *Nature Astron.* **2**, 407–412 (2018).
5. Micheli, M. et al. Non-gravitational acceleration in the trajectory of 1I/2017 U1 (Oumuamua). *Nature* **559**, 223–226 (2018).
6. MPEC 2019-R106: COMET C/2019 Q4 (Borisov). Available at: <https://www.minorplanetcenter.net/mpec/K19/K19RA6.html>. (Accessed: 16th September 2019)
7. Fukugita, M. et al. The Sloan Digital Sky Survey photometric system. *Astron. J.* **111**, 1748–1756 (1996).
8. Eisenstein, D. J. et al. SDSS-III: Massive Spectroscopic Surveys of the Distant Universe, the Milky Way, and Extra-Solar Planetary Systems. *Astron. J.* **142**, 72 (2011).
9. Boe B., et al., The Orbit and Size-Frequency Distribution of Long Period Comets Observed by Pan-STARRS1. *Icarus* **333**, 252–272 (2019).
10. Holmberg, J., Flynn, C. & Portinari, L. The colours of the Sun. *Mon. Not. R. Astron. Soc.* **367**, 449–453 (2006).
11. Waniak, W., Zola, S. & Krzesinski, J. Dust Emission for Comets Shoemaker-Levy 1991a1 and McNaught-Russell 1993v. *Icarus*, **136**, 280–297 (1998).
12. Drahus, M. et al., Rotation State of Comet 103P/Hartley 2 from Radio Spectroscopy at 1 mm. *Astrophys. J. Lett.* **734**, L4 (2011).
13. Jewitt, D. & Luu, J. Disintegrating In-Bound Long-Period Comet C/2019 J2. *ArXiv1909.01964 Astro-Ph* (2019).
14. Drahus, M. Rotational Disruption of Comets with Parabolic Orbits. *46th DPS meeting, id.200.04*
15. Jewitt, D. Color systematics of comets and related bodies. *Astron. J.* **150**, 201 (2015).
16. The Oumuamua ISSI Team, The natural history of Oumuamua. *Nature Astron.* **3**, 594–602 (2019).
17. Jewitt, D. et al. Interstellar Interloper 1I/2017 U1: Observations from the NOT and WIYN Telescopes. *Astrophys. J. Lett.* **850**, L36 (2017).
18. Do, A., Tucker, M. A. & Tonry, J. Interstellar Interlopers: Number Density and Origin of Oumuamua-like Objects. *Astrophys. J. Lett.* **855**, L10 (2018).
19. Moro-Martín, A., Turner, E. L. & Loeb, A. Will the Large Synoptic Survey Telescope Detect Extra-Solar Planetesimals Entering the Solar System? *Astrophys. J.* **704**, 733–742 (2009).

20. Harris, A., On the Revision of Radiometric Albedos and Diameters of Asteroids. *Icarus*, **126**, 450-454 (1997).
21. Covan, J., A'Hearn, M., Vaporization of Comet Nuclei: Light Curves and Life Times. *M&P*, **21**, 155 (1979).
22. Jorda, L., Crovisier, J., Green, D. W. E., The Correlation Between Visual Magnitudes and Water Production Rates. *Asteroids, Comets, Meteors*, 2008.
23. Meisel D., Morris, C., Comet head photometry - Past, present, and future. in *Comets* (ed. L. L. Wilkening) 413-432 (1982).

Author information

Correspondence and requests for materials should be addressed to P.G.

(piotr.guzik@doctoral.uj.edu.pl) or M.D. (drahus@oa.uj.edu.pl).

Acknowledgements

Based in part on observations obtained at the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the National Research Council (Canada), CONICYT (Chile), Ministerio de Ciencia, Tecnología e Innovación Productiva (Argentina), and Ministério da Ciência, Tecnologia e Inovação (Brazil). The William Herschel Telescope is operated on the island of La Palma by the Isaac Newton Group of Telescopes in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. We thank the staff of both observatories for assistance and are indebted to the Director of Gemini Observatory for allocating Gemini North time for this program. M.D. and P.G. are grateful for support from the National Science Centre of Poland through SONATA BIS grant 2016/22/E/ST9/00109 and Polish Ministry of Science and Higher Education grant DIR/WK/2018/12. G.C.

acknowledges support from European Research Council Consolidator Grant 647208. I.P.M.

acknowledges funding from the Netherlands Research School for Astronomy (NOVA5-

NW3-10.3.5.14).

Competing Financial Interests

The authors declare that they have no competing financial interests.

Figure Legends

Figure 1 | Astrometric residuals of C/2019 Q4 (Borisov) calculated for parabolic and unrestricted (hyperbolic) solutions. Top and bottom left panels show the astrometric residuals calculated for hyperbolic fit in Right Ascension and Declination respectively. Right panels show the astrometric residuals calculated for best parabolic fit in Right Ascension and Declination accordingly. Mean residuals are 0.59 arcsec and 1.30 arcsec for hyperbolic and parabolic solution respectively.

Figure 2 | Median-stacked images of C/2019 Q4 (Borisov) from Gemini North. Left panel shows the image in the g' band and right panel shows the image in r' band. Both panels subtend 1.0×1.0 arcmin, north is to the top and east is to the left. The images were scaled logarithmically.


Figure 3 | Color of C/2019 Q4 (Borisov) in the context of Solar System's long-period comets. Histogram shows the g'-r' color index distribution for 31 measurements of 25 comets¹⁵, calculated from the B-V and V-R colors using standard transformation formulae. Solid blue line is a fitted normal distribution, which has an average value of 0.628 mag and a standard deviation 0.074 mag, and dashed red line indicates the g'-r' color index of Comet Borisov.



Tables

Table 1 | Hyperbolic orbital elements of C/2019 Q4 (Borisov) calculated for the osculation epoch 2019 Sep. 11.0 TT.

T	2019 Dec. 9.745968 \pm 0.94 TT
e	3.1551235 \pm 0.13
q	1.95954186 \pm 0.035 au
ω	210.22059 \pm 0.90 deg (2000.0)
Ω	307.71365 \pm 0.33 deg (2000.0)
i	44.49503 \pm 0.34 deg (2000.0)

Figures





Supplementary Materials

Supplementary Table 1 | List of observers who contributed observations that were used to find hyperbolic orbit of C/2019 Q4 (Borisov).

Station code	Observer(s)	# of observations
L51	Gennady Borisov	34
595	E. Pettarin	3
H06	H. Sato	4
186	A. Novichonok	3
L01	K. Korlević, A. Radica, F. Valentine	3
J95	P. Birtwhistle	6
C10	J.-F. Soulier	3
A71	M. Jaeger, E. Prosperi, S. Prosperi	2
G40	B. Lutkenhoner, G. Gasparovic, Y. Chen	5
N55	T. Chen	16
568	K. J. Meech, J. Kleyna	5
121	I. Slyusarev	3
L06	G. Ventre	3
C23	A. Diepvens	2
104	P. Bacci, M. Maestripieri, R. Bacci	7
T05	L. Denneau, J. Tonry, A. Heinze, H. Weiland, H. Flewelling	15
L04	L. Hudin	6
585	A. Baransky, N. Kaplina	7
H36	G. Hug	2

Supplementary Figure 1 | Visibility prospects of C/2019 Q4 (Borisov) in the upcoming months. Top panel illustrates basic geometric circumstances and bottom panel shows r'-band magnitude extrapolated from our measurement for different photometric models. We adopted the canonical form of photometric models for active comets $m = H + 5.0 \log (\Delta) + 2.5 n \log (r)$, where m is the observed magnitude, H is the absolute magnitude (at 1 au from Earth and Sun), n is a free parameter (typically $n \sim 4$), Δ (au) is the geocentric, distance and r (au) is the heliocentric distance.

