

SCUBA-2 observations of candidate starbursting protoclusters selected by *Planck* and *Herschel*-SPIRE

T. Cheng,^{1*} D.L. Clements,¹ J. Greenslade,¹ J. Cairns,¹ P. Andreani,² M. Bremer,³
 L. Conversi,⁴ A. Cooray,⁵ H. Dannerbauer,^{6,7} G. De Zotti,⁸ S. Eales,⁹
 J. González-Nuevo,¹⁰ E. Ibar,¹¹ L. Leeuw,¹² J. Ma,⁵ M. J. Michałowski,¹³
 H. Nayyeri,⁵ D. A. Riechers,^{14,15} D. Scott,¹⁶ P. Temi,¹⁷ M. Vaccari,^{18,19}
 I. Valtchanov,²⁰ E. van Kampen² and L. Wang^{21,22}

¹*Astrophysics Group, Imperial College London, Blackett Laboratory, Prince Consort Road, London SW7 2AZ, UK*

²*European Southern Observatory, Karl-Schwarzschild-Strae 2, 85748 Garching, Germany*

³*HH Wills Physics Laboratory, University of Bristol, Tyndall Avenue, Bristol, BS8 1TL, UK*

⁴*European Space Agency / ESAC, Camino Bajo del Castillo, 28692 Villanueva de la Cañada (Madrid), Spain*

⁵*Department of Physics and Astronomy, University of California, Irvine, CA 92697, USA*

⁶*Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain*

⁷*Universidad de La Laguna Dpto. Astrofísica, E-38206 La Laguna, Tenerife, Spain*

⁸*INAF-Osservatorio astronomico di Padova, Vicolo dell'Osservatorio 5, I-35122 Padova, Italy*

⁹*School of Physics and Astronomy, Cardiff University, The Parade, Cardiff CF24 3AA, UK*

¹⁰*Departamento de Física, Universidad de Oviedo, C. Federico García Lorca 18, 33007 Oviedo, Spain*

¹¹*Instituto de Física y Astronomía, Universidad de Valparaíso, Avda. Gran Bretaña 1111, Valparaíso, Chile*

¹²*College of Graduate Studies, University of South Africa, PO Box 392, UNISA, 0003, South Africa*

¹³*Astronomical Observatory Institute, Faculty of Physics, Adam Mickiewicz University, ul. Słoneczna 36, 60-286 Poznań*

¹⁴*Department of Astronomy, Cornell University, Space Sciences Building, Ithaca, NY 14853, USA*

¹⁵*Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany*

¹⁶*Department of Physics and Astronomy, University of British Columbia, Vancouver, BC V6T1Z1, Canada*

¹⁷*Astrophysics Branch, NASA Ames Research Center, Moffett Field, CA 94035, USA*

¹⁸*Department of Physics and Astronomy, University of the Western Cape, Private Bag X17, Bellville 7535, Cape Town, South Africa*

¹⁹*INAF - Istituto di Radioastronomia, via Gobetti 101, 40129 Bologna, Italy*

²⁰*Telespazio Vega UK for ESA, European Space Astronomy Centre, Operations Department, E-28691 Villanueva de la Cañada, Spain*

²¹*SRON Netherlands Institute for Space Research, Landleven 12, 9747 AD, Groningen, The Netherlands*

²²*Kapteyn Astronomical Institute, University of Groningen, Postbus 800, 9700 AV, Groningen, The Netherlands*

Accepted 2019 September 17. Received 2019 August 19; in original form 2019 April 25

ABSTRACT

We present SCUBA-2 850- μm observations of 13 candidate starbursting protoclusters selected using *Planck* and *Herschel* data. The cumulative number counts of the 850- μm sources in 9/13 of these candidate protoclusters show significant overdensities compared to the field, with the probability $<10^{-2}$ assuming the sources are randomly distributed in the sky. Using the 250-, 350-, 500- and 850- μm flux densities, we estimate the photometric redshifts of individual SCUBA-2 sources by fitting spectral energy distribution (SED) templates with an MCMC method. The photometric redshift distribution, peaking at $2 < z < 3$, is consistent with that of known $z > 2$ protoclusters and the peak of the cosmic star-formation rate density (SFRD). We find that the 850- μm sources in our candidate protoclusters have infrared luminosities of $L_{\text{IR}} \gtrsim 10^{12} L_{\odot}$ and star-formation rates of $\text{SFR} = (500\text{--}1,500) M_{\odot} \text{yr}^{-1}$. By comparing with results in the literature considering only *Herschel* photometry, we conclude that our 13 candidate protoclusters can be categorised into four groups: six of them being high-redshift starbursting protoclusters, one being a lower-redshift cluster/protocluster, three being protoclusters that contain lensed DSFG(s) or are rich in 850- μm sources, and three regions without significant *Herschel* or SCUBA-2 source overdensities. The total SFRs of the candidate protoclusters are found to be comparable or higher than those of known protoclusters, suggesting our sample contains some of the most extreme protocluster population. We infer that cross-matching *Planck* and *Herschel* data is a robust method for selecting candidate protoclusters with overdensities of 850- μm sources.

Key words: galaxies: high-redshift – galaxies: starburst – submillimetre: galaxies

1 INTRODUCTION

Protoclusters are defined as structures that are expected to collapse into galaxy clusters before the present epoch (e.g. Overzier 2016). Normally at $z > 2$, their hot gas halos may not yet be virialized. Without the virialized hot gas, protoclusters are difficult to find via traditional cluster-detection methods, such as X-rays or the Sunyaev-Zeldovich Effect (SZE). They also may not have a significant number of red sequence galaxies which would allow them to be identified on the optical colour-magnitude diagram (CMD), where red sequence galaxies at a fixed redshift cluster together due to similar stellar populations (Brodwin et al. 2007; Eisenhardt et al. 2008; Newman et al. 2014; Andreon et al. 2014). Recent observations of high- z (proto)clusters have found some red sequence galaxies (Martinache et al. 2018), but such cases are rare.

Most protoclusters are found through optical/near-infrared surveys, using overdensities of Lyman- α emitters (LAEs), Lyman-break galaxies (LBGs) or H- α emitters (HAEs) identified in the field (e.g. The Hyper Suprime-Cam Subaru Strategic Program, HSC-SSP: Aihara et al. 2018b,a; Douglas et al. 2010) or around rare objects such as QSOs or radio galaxies (Pentericci et al. 2000; Kurk et al. 2004; Verhamme et al. 2008; Tanaka et al. 2011; Hayashi et al. 2012; Husband et al. 2013; Casey et al. 2015). Thus QSOs and radio galaxies are often used as beacons when searching for protoclusters.

Dusty star-forming galaxies (DSFGs) are galaxies heavily obscured by dust and forming stars rapidly (Smail et al. 1997; Hughes et al. 1998). Submillimetre galaxies (SMGs) are a subsample of DSFGs selected in submillimetre surveys. The brightest and most luminous of these DSFGs can have luminosities exceeding $10^{13}L_{\odot}$ (Chapman et al. 2005; Gruppioni et al. 2015; Rowan-Robinson et al. 2018; Hill et al. 2018), which corresponds to star-formation rates (SFRs) of thousands of solar masses per year (Casey et al. 2014), under standard calibrations (Zhang et al. 2018). In galaxy formation models, DSFGs are thought to be the progenitors of elliptical, early-type galaxies residing in the cores of today’s massive galaxy clusters (Farrar et al. 2006; Lapi et al. 2006; Fan et al. 2008; Cook et al. 2010; Lapi et al. 2011; Wang et al. 2013; Cai et al. 2013; Lapi et al. 2014; Toft et al. 2014; Aversa et al. 2015; Wilkinson et al. 2017), so in principle we should see DSFGs in cluster progenitors, such as protoclusters.

Although it is statistically possible for line-of-sight overdensities of DSFGs to occur (Negrello et al. 2017), there are existing observations of protoclusters hosting DSFGs (Chapman et al. 2009; Daddi et al. 2009; Dannerbauer et al. 2014; Umehata et al. 2015; Casey et al. 2015) with SFRs as high as $3,000 M_{\odot}\text{yr}^{-1}$. Some of the DSFGs in protoclusters are also found to be formed of multiple sources by using higher resolution imagers, e.g. ALMA, VLT and HST (Bussmann et al. 2015; Wang et al. 2016; Gómez-Guijarro et al. 2018; Oteo et al. 2018; Kneissl et al. 2018; Gómez-Guijarro et al. 2019). There are also cases where overdensities of DSFGs in the sky are line-of-sight projections of two protocluster structures at different redshifts (Flores-Cacho et al. 2016). The redshifts of these protoclusters range from $z \approx 2$ to $z \approx 5$ (Capak et al. 2011; Walter et al. 2012; Hodge et al. 2013), and their angular sizes vary from <1 arcminute to around 0.5

degree, probing regimes of size from cluster cores to large-scale structures such as filaments (Hayashino et al. 2004; Matsuda et al. 2005; Chiang et al. 2013; Wang et al. 2016; Oteo et al. 2018).

According to some galaxy-formation models, during the formation of a galaxy cluster, member galaxies are expected to undergo a starbursting phase (Granato et al. 2004; Casey 2016), making them so-called “starbursting galaxies”. The timescale of this starbursting phase is short compared to the formation of a galaxy cluster and this phase might start and end at different times for different member galaxies; therefore, the probability that we observe a protocluster hosting a large number of DSFGs at the same time is very low (Chiang et al. 2013; Casey 2016). The fact that a number of protoclusters containing DSFGs have been observed (Steidel et al. 1998; Chapman et al. 2009; Dannerbauer et al. 2014; Casey et al. 2015) suggests there is an inconsistency between model predictions and observations.

Some models suggest that these starbursting galaxies might be mainly driven by mergers (Joseph & Wright 1985; Hopkins et al. 2008; Sparre & Springel 2016), perhaps explaining why there are more DSFG-rich protoclusters than predicted. Alternatively, Casey (2016) proposes that the starbursting phase in cluster galaxies can happen simultaneously based on observations of gas depletion times of known protoclusters, and hence, might explain why there are observations of DSFG-rich protoclusters.

Even though some optical surveys have been dedicated to protocluster searches, the total number of confirmed protoclusters to date is still limited to approximately 40 (Overzier 2016). More protoclusters are needed, especially at high redshifts, in order to understand this population, and to resolve the inconsistency between observations and models. Searches for protoclusters in the submillimetre or far-infrared (FIR) bands are especially important because they directly look for DSFGs, and a galaxy’s flux density can be as bright at $z \sim 8$ as at $z = 0$, due to the negative submm K-correction (see Casey et al. 2014).

Negrello et al. (2005) proposed a method that uses two FIR imaging instruments, one low and another high resolution, to identify candidate protoclusters. They show that the total flux density within the beam of a low-resolution instrument is in fact often the sum of a clump of several structures, which can be distinguished with a high-resolution instrument. Using this method with the *Planck*¹ High Frequency Instrument (HFI, Lamarre et al. 2010) as the low resolution instrument and *Herschel*² SPIRE (Spectral and Photometric Imaging Receiver, Griffin et al. 2010) as the high resolution instrument, a number of candidate protoclusters have been found (Herranz et al. 2013; Clements et al. 2014, 2016; Planck Collaboration et al. 2016c; Greenslade et al.

¹ *Planck* (<http://www.esa.int/Planck>) is a project of the European Space Agency (ESA) with instruments provided by two scientific consortia funded by ESA member states (in particular the lead countries France and Italy), with contributions from NASA (USA) and telescope reflectors provided by a collaboration between ESA and a scientific consortium led and funded by Denmark.

² *Herschel* (Pilbratt et al. 2010) is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.

2018) using maps from H-ATLAS (Eales et al. 2010) and HerMES (Oliver et al. 2012) surveys (as well as targeted observations, Planck Collaboration et al. 2016b). This method is ideal for searching for high-redshift protoclusters, probing their formation periods and covering the epoch where the cosmic star-formation density peaks (Hopkins & Beacom 2006; Clements et al. 2014).

Even though a number of candidate protoclusters have been identified using this method, their total cluster flux densities are found to be greater than in simulations by about a factor of 3 at 350 μm (Granato et al. 2015; Greenslade et al. 2018). Such an inconsistency between observations and simulations in Granato et al. (2015) cannot simply be explained by starbursts with higher SFRs, since this will not give the correct total cluster stellar mass expected at $z=0$ in the simulations. In order to resolve this issue, observations need to be conducted on more protoclusters in order to feed into simulations.

Follow-up observations of candidate protoclusters selected using the above method have already been conducted. Observations using LABOCA (LARGE APEX BOlometer CAmera, Siringo et al. 2009) and SCUBA-2 (Holland et al. 2013) have found overdensities of submillimetres sources in a candidate protocluster which hosts a lensed DSFG at $z=3.26$ (Clements et al. 2016). Photometric redshifts of a few other candidate protoclusters using ancillary data suggest redshifts from 1 to 3 (Clements et al. 2014). In this paper we present follow-up observations of 13 candidate protoclusters selected using the above method. We study their number counts, far-infrared colours and photometric redshifts using observations from SCUBA-2 and *Herschel*-SPIRE.

The content of this paper is as follows. In Section 2 we describe the *Herschel*-SPIRE and SCUBA-2 observations of these 13 candidate protoclusters. In Section 3 we present results on number counts, FIR colours and photometric redshifts. We discuss and conclude in Sections 4 and 5, respectively. The standard concordance cosmology of $H_0 = 70\text{kms}^{-1}\text{Mpc}^{-1}$, $\Omega_M = 0.3$, and $\Omega_\Lambda = 0.7$ is used throughout this paper.

2 OBSERVATIONS AND DATA REDUCTION

2.1 Selection of candidate protoclusters

The selection process for the candidate protoclusters, including those observed with SCUBA-2 described in this paper, is defined in Greenslade et al. (2018) (hereafter Gr18). Here we briefly summarise the selection process.

Using a protocluster survey technique, as introduced in Negrello et al. (2005), sources from *Planck* catalogues of compact sources (beam FWHM 5') were selected. Subsequently these were examined in *Herschel*-SPIRE maps (beam FWHM 18–36'') and catalogues from H-ATLAS and HerMES surveys to exclude local galaxies, Galactic cirrus or lensed objects. SPIRE has bands at 250-, 350-, and 500- μm , and is often used to select dusty and far-infrared-bright sources at $z > 2$ (Riechers et al. 2013; Dowell et al. 2014; Asboth et al. 2016) or lensed DSFGs (Wardlow et al. 2013; Nayyeri et al. 2016).

After the above selection criteria, some of the remaining *Planck* compact sources (Herranz et al. 2013) were targeted with other observations (Clements et al. 2014, 2016).

Gr18 further identified 27 *Planck* compact sources as candidate high-redshift protoclusters where their *Herschel* source overdensities were $>3\sigma$ in at least one SPIRE band.

Starting with the *Planck* catalogues of compact sources, we looked for overdensities of *Herschel*-SPIRE sources in the 250, 350, and/or 500 μm bands (Greenslade et al. (2018)) from H-ATLAS (Eales et al. (2010)) and HerMES (Oliver et al. (2012)) surveys. One of the aims of these surveys are to look at fields in the sky which are also covered by other multi-wavelength and extragalactic surveys. The total area of sky covered by H-ATLAS and HerMES are 570 deg^2 and 70 deg^2 , respectively. The instrument we use is SPIRE, which has bands in 250, 350, and 500 μm , covering the far-infrared range and is often used to select dusty and far-infrared-bright sources at $z > 2$ (Riechers et al. (2013); Dowell et al. (2014); Asboth et al. (2016)). Such dusty sources have rest-frame spectral energy distributions (SEDs) peaking at $\sim 100 \mu\text{m}$ due to re-radiated dust-obscured UV lights, and these dust-SED peaks are redshifted to SPIRE bands.

The SCUBA-2 observations were carried out for 13 fields of candidate protoclusters selected in Gr18 which are accessible by JCMT. Their properties (coordinates and overdensities) are listed in Table 1. The overdensity values are reported from Gr18, in which they counted the number of *Herschel*/SPIRE sources with flux densities above 25.4 mJy at 350 or 500 μm within the *Planck* beam and compared with the expected number counts from Clements et al. (2010) and Valiante et al. (2016).

Note that five candidate protocluster fields were in fact not classified as candidate protoclusters in Gr18 because their *Herschel*-SPIRE source overdensities are below 3σ . However, we still describe them as candidate protoclusters throughout this paper, since Gr18 only applied their 3σ overdensity cut to sources in *Herschel*-SPIRE bands; protocluster galaxies that are brighter at other bands (particularly 850 μm), might be missed. Section 4.1 discusses this additional selection effect.

The specific candidate protoclusters labelled Bootes1, EGS and Lockman were studied in Clements et al. (2014) and have redshift estimates of 2.27 ± 0.24 , 0.76 ± 0.10 , and 2.05 ± 0.09 , respectively. The candidate protocluster G12 is also well studied (Clements et al. 2016) and is believed to be at $z = 3.26$, consistent with the lensed DSFG in the centre of the field, which has a spectroscopic redshift (Fu et al. 2012).

2.2 SCUBA-2 Observations

The JCMT/SCUBA-2 observations for these 13 candidate protoclusters took place between 8 and 12 April, 2013 (Project number M13AU12, PI D. Clements). The data were obtained at both 450 and 850 μm simultaneously, but the 450- μm maps did not reach the sensitivity needed to detect protocluster galaxies due to the weather conditions, so we only study the 850- μm maps here. Each candidate protocluster field was observed for approximately 2 hours, with several pointings using the CV Daisy mode (Holland et al. 2013), giving rms noise levels of approximately 2.25 mJy/beam. Weather conditions were good throughout these observations (precipitable water vapour between 0.83 mm and 2.58 mm) and standard calibrations were conducted. Fig.1 shows an example of the noise distribution in one of our candidate

Table 1. List of 13 candidate protoclusters observed with SCUBA-2 in this paper. Overdensity values are quoted from Gr18, in which *Herschel* sources were selected with flux densities above 25.4 mJy at 350 or 500 μm within the *Planck* beam, and compared with expected field number counts from Clements et al. (2010) and Valiante et al. (2016). Note that G12, NGP2, NGP3, NGP6 and NGP9 have overdensities below 3σ in all SPIRE bands so were not selected as candidate protoclusters in Gr18 but we still refer to them as candidate protoclusters in this paper. The last column shows the category after comparing the overdensities of candidate protoclusters in Gr18 and in this paper, which will be discussed in Section 4.1.

Name	RA (J2000)	Dec (J2000)	Overdensity at 250 μm (σ_{250})	Overdensity at 350 μm (σ_{350})	Overdensity at 500 μm (σ_{500})	Category (Section 4.1)
Bootes1	14:34:18.1	+35:33:20.0	1.0	4.5	5.7	I
EGS	14:24:35.8	+52:56:42.0	4.7	5.8	2.4	I
G12	11:46:33.6	-00:11:15.0	0.8	2.8	2.4	III
Lockman	10:33:26.9	+59:10:09.1	5.4	4.5	4.7	II
NGP1	13:24:25.5	+28:44:47.6	2.3	2.0	3.3	IV
NGP2	13:19:37.2	+26:28:01.6	1.0	2.2	2.9	III
NGP3	13:31:42.9	+23:46:16.9	1.5	1.4	1.6	III
NGP4	13:14:26.0	+26:30:35.6	2.5	3.3	4.0	I
NGP5	13:40:41.4	+32:37:17.1	2.1	3.5	3.3	I
NGP6	13:23:12.2	+33:23:11.9	2.3	0.8	1.1	IV
NGP7	13:37:06.7	+32:07:55.3	0.8	2.5	3.3	I
NGP8	13:29:26.2	+28:13:25.4	1.7	3.8	4.0	I
NGP9	12:59:15.5	+31:35:40.7	0.5	1.7	2.9	IV




Figure 1. The noise map of the candidate protocluster Bootes1. Colorbar units are in mJy. The green circle indicates the central region with 4 arcminute radius where the sensitivity is highest. North is up and east is left.

protoclusters. The green circle shows the central region with 4 arcminute radius, indicating the most sensitive region.

The data were reduced using the SCUBA-2 pipeline SMURF package (Chapin et al. 2013). The raw SCUBA-2 data are composed of 30-second chunks of observations, for each of the four sub-arrays at each wavelength. They are in the STARLINK NDF format (.sdf). The first step is to use the MAKEMAP command to create a map combining all sub-arrays and observing chunks for each scan. Under the MAKEMAP command, the ITERATE method is used, which fits models for the noise and the instrumental behaviour. Next, the PICARD

recipe MOSAIC_JCMT_IMAGES is used, which co-adds separate scans into a single map and removes contaminant signals such as cosmic rays.

Since we expect these 850- μm sources to be point sources and unresolved due to their high redshifts, we used PICARD’s SCUBA2_MATCHED_FILTER recipe. This recipe first subtracts the background by convolving (smoothing) the maps and the PSF with a 30” FWHM Gaussian kernel. Then the signal maps are convolved with the PSF to produce the matched-filtered signal map. The noise maps are also convolved with the PSF to produce the variance map. This process essentially performs the maximum likelihood fit of the PSF on every pixel of the map, and is beneficial at finding sources with angular scales of the telescope beam. This process gives an effective beam FWHM of 14.6 arcseconds (Dempsey et al. 2013) surrounded by a shallow negative ring.

The last step in the reduction is to crop the maps to a diameter of 700 arcsec (11.7 arcmin) where they are most sensitive using PICARD’s CROP_SCUBA2_IMAGES recipe and produce signal-to-noise maps using the MAKESNR command. The standard flux conversion factor (FCF) of 537 Jy/pW at 850 μm was used for the calibration from detected power units to flux density (Dempsey et al. 2013). According to JCMT/SCUBA-2 website³, the calibration error is approximately 5 percent. We include the 5% calibration error and add it in quadrature to the uncertainties of the flux densities of the 850 μm sources.

2.3 Source Detection

In order to build our SCUBA-2 850- μm source catalogue, we find peaks in the S/N maps, which were obtained by dividing the beam-convolved map with the noise map. Nearby, connected pixels having S/N > 3.5 are regarded as part of the

³ <https://www.eaobservatory.org/jcmt/instrumentation/continuum/scuba-2/calibration/>

same source and the highest S/N pixel is used to record the flux density and noise of the source.

A correction to the flux densities and noise is needed, due to flux boosting. We apply the correction from equation 5 of Geach et al. (2017), in which the flux density and noise correction is a power-law function of the observed signal-to-noise ratio (SNR):

$$\frac{S_{obs}}{S_{true}} = 1 + 0.2 \left(\frac{SNR}{5} \right)^{-2.3}, \quad (1)$$

where S_{obs} is the observed flux density and noise; S_{true} is the de-boosted flux density and noise.

We study the reliability of the sources by inverting the flux density maps, which are background subtracted, and using the same extraction method as above. By doing this, negative sources due to negative noise peaks will be found. We assume there is a similar number of spurious sources in the original maps due to positive noise peaks. By calculating the ratio between the detected number of positive sources and the expected number of spurious sources, we find a reliability of 80 percent at 3.5σ , consistent with similar SCUBA-2 studies of candidate protoclusters, such as in MacKenzie et al. (2017).

Since the beam deviates from a Gaussian, the number of positive and negative noise spikes may be different, which may cause incorrect reliability estimates using the above method. Hence we also produce a series of jackknife maps, where the data are separated into two halves alternately. The data of one half is then inverted and co-add with the other half, producing what is essentially the maps with pure noise. By doing source extraction on these jackknife maps, we found the reliability of all 13 candidate protocluster fields being significantly better than 80 percent at 3.5σ .

Using a higher S/N detection threshold than 3.5σ would improve the reliability of the sources while decreasing the total number of sources. According to our reliability studies there are chances of spurious sources in our catalogue. For example, those with photometric redshifts of $z > 6$ (see Table B1; discussed in Section 3.3) are likely to be spurious due to their bad photometry. To test how much change different S/N values will incur, we impose different S/N cuts up to 5 σ and perform the cumulative number count analysis (discussed in Section 3.1). We find no significant change in the conclusions we make in this paper.

The completeness of the sources is estimated by inserting fake sources from 2 to 20 mJy to the flux density maps of each candidate protocluster field, and then calculating the fraction of these sources recovered using the same source detection algorithm. The shape of the fake sources is a circular 2D Gaussian, with the standard deviation being half of the SCUBA-2 beam FWHM at $850 \mu\text{m}$, i.e. 14.6 arcsec. We found that although the completeness levels vary between different fields, all but three candidate protocluster fields show a completeness level above 50 percent at 8 mJy, after flux-deboosting. If we restrict the fake sources to be within the 4 arcmin radius, all candidate protocluster fields have a completeness level above 50 percent at ≥ 8 mJy. In the three fields (EGS, NGP3, NGP8) where completeness falls below 50 percent at 8 mJy, only 2 sources lie outside the 4 arcmin radius region, and only in NFP3 field. We note these 2 sources in the source catalogue in Table B1 since

they have a lower completeness level. Overall we conclude our SCUBA-2 sources have a reasonable completeness level above 8 mJy and calculate the probability of the observed number of sources above this flux density in the studies of number counts (Section 3.1).

Maps showing the detected sources in all 13 candidate protocluster fields are shown in Fig A1. Catalogues showing the de-boosted flux densities of the detected SCUBA-2 $850\text{-}\mu\text{m}$ sources for all 13 candidate protocluster fields are given in Table B1.

3 RESULTS

3.1 Cumulative Number Counts of SCUBA-2 Sources

In order to study the existence of overdensities of $850\text{-}\mu\text{m}$ sources, cumulative number counts are estimated for these 13 fields, based on their de-boosted flux densities.

Fig.2 shows the cumulative number counts. We sort the flux densities of the sources from bright to faint and count the cumulative number of sources at each 2-mJy step, and plot as blue points. The results from Geach et al. (2017) (hereafter Ge17) are shown in red, which is from the SCUBA-2 Cosmology Legacy Survey (S2CLS) and is composed of approximately 5 deg^2 of blank field.

These cumulative counts need to be corrected for the changing sensitivity across the maps. The effective area to detect each source is smaller than the entire map due to the attenuated sensitivity near the edge. We have corrected for this overestimated area by dividing the number of sources by the effective area (rather than the entire map area) corresponding to different sensitivities. Hence the number of sources in Fig 2 and Table 2 is greater than that listed in Table B1.

The cumulative number counts of the 13 candidate protoclusters and of the field in S2CLS from Ge17 are listed in Table 2. We now describe the number count behaviour of each candidate protocluster field and calculate the probability⁴ of the observed number of sources above 8 mJy, assuming that the sources are randomly distributed. All the observed number counts are quoted after the variable sensitivity is corrected for in the effective size of each individual map. In the last row of Table 2 we list the expected number counts from Ge17 scaled to the size of each candidate protocluster field, which is approximately 0.03 deg^2 .

Bootes1: The cumulative number counts show overdensities ($> 5\sigma$) of $850 \mu\text{m}$ sources at 6.0, 8.0 and 10.0 mJy, compared to Ge17. We observe 6 sources with flux densities above 8.0 mJy in the map. Compared to the expected number from Ge17, and assuming that the sources are randomly distributed, the probability of observing this number of sources in a random field ($P(\geq 6)$) is 0.02, following a Poisson distribution.

EGS: An overdensity of $850\text{-}\mu\text{m}$ sources is seen from 8.0 to 12.0 mJy in the cumulative number counts. We observe 14 sources with flux densities > 8.0 mJy in the map. The

⁴ This is the *upper tail* of the probability density function following a Poisson distribution, calculated using R function `ppois(observed-1, lambda=expected, lower=FALSE)`.

Table 2. Cumulative number counts of our 13 candidate starbursting protoclusters and in the field from the S2CLS survey (Geach et al. 2017). Cumulative number counts represent the number of sources with flux densities >4 mJy, >6 mJy, >8 mJy, >10 mJy and >12 mJy. The uncertainties in the candidate protocluster fields represent the completeness error and the uncertainties in the S2CLS survey represent Poissonian errors. The number of sources is corrected for the variable sensitivity and is compared with field surveys. Flux densities of the two lensed sources in G12 and NGP1 are de-magnified based on the magnification factor (see text) and counted. The last row shows the expected number counts scaled to the area of each map, approximately 0.03 deg^2 , in order to compare with the actual number of sources in the maps, as discussed in the text. The P_{random} column shows the probability of the observed number of sources compared with the expected number in Geach et al. (2017) at 8 mJy, assuming the sources are randomly distributed. $N_{\text{overdensity}}$ and $P_{\text{overdensity}}$ columns are the number of overdense random regions and the overdense level (fraction of regions showing an overdensity, see text), after putting 10,000 random regions in the S2CLS/COSMOS field to study the potential cosmic variance effect.

Name	>4 mJy	>6 mJy	>8 mJy	>10 mJy	>12 mJy	P_{random} (at 8 mJy)	$N_{\text{overdensity}}$	$P_{\text{overdensity}}$
Bootes1	49±35	24±0.4	6.0±0.07	2.1±0.02	N/A	0.02	920	0.092
EGS1	14±0.6	14±0.6	14±0.6	1.4±0.02	1.4±0.02	2.44×10^{-8}	0	$< 10^{-4}$
G12	33±0.3	33±0.3	8.7±0.09	2.1±0.02	N/A	9.97×10^{-4}	318	0.0318
Lockman	3.7±0.08	3.7±0.08	1.7±0.02	N/A	N/A	0.86	4422	0.4422
NGP1	16±0.2	16±0.2	3.8±0.04	N/A	N/A	0.32	1825	0.1825
NGP2	24±10	24±10	6.5±0.1	2.8±0.03	1.2±0.01	0.02	616	0.0616
NGP3	5.1±0.2	5.1±0.2	5.1±0.2	2.3±0.03	2.3±0.03	0.05	920	0.092
NGP4	22±7	22±7	17±0.3	3.7±0.04	1.1±0.01	4.46×10^{-11}	0	$< 10^{-4}$
NGP5	19±3	19±3	5.7±0.06	5.7±0.06	1.0±0.01	0.05	920	0.092
NGP6	18±0.3	18±0.3	1.0±0.01	1.0±0.01	N/A	0.86	7233	0.7233
NGP7	46±6	46±6	13±0.2	6.1±0.08	3.4±0.04	1.75×10^{-7}	17	0.0017
NGP8	16±0.9	16±0.9	16±0.9	4.1±0.06	N/A	3.88×10^{-10}	0	$< 10^{-4}$
NGP9	6.1±0.1	6.1±0.1	3.5±0.03	N/A	N/A	0.32	1825	0.1825
S2CLS (expected)	22.6±0.34	6.3 $^{+0.16}_{-0.15}$	1.97 $^{+0.09}_{-0.08}$	0.61±0.05	0.21±0.03			

probability of observing this number of sources in a random field ($P(\geq 14)$) is 2.44×10^{-8} .

Lockman: There is no overdensity in the observed flux density range. There are 1.7 sources with flux densities above 8.0 mJy in the map. The probability of observing this number of sources in a random field ($P(\geq 1.7)$) is 0.86.

G12: Overdensities in cumulative number counts are seen from 6.0 to 10.0 mJy. We observe 8.7 sources with flux densities above 8.0 mJy in the map. The probability of observing this number of sources in a random field ($P(\geq 8.7)$) is 9.97×10^{-4} . In the central region of G12, there is a strongly lensed system, which is at $z=3.26$ (Fu et al. 2012; Herranz et al. 2013), and has an observed SCUBA-2 flux density of 79.8 ± 4.2 mJy. The $850 \mu\text{m}$ flux density of this source is de-magnified according to the magnification factor $\mu(880 \mu\text{m}) = 7.6 \pm 1.5$ from Fu et al. (2012) and considered in the cumulative number counts. This candidate protocluster was already studied in Clements et al. (2016), and is believed to have a similar redshift to the spectroscopically confirmed lensed DSFG at $z=3.26$. Note that the number of observed FIR sources presented here is higher than that in Clements et al. (2016) because the quoted number of sources has been corrected for changing sensitivity.

NGP1: There is a bright SCUBA-2 source in the central region of this candidate protocluster with a flux density of 42.8 ± 2.5 mJy, which is a $z=1.676$ lensed source (Bussmann et al. 2013; Calanog et al. 2014; Timmons et al. 2016). The flux density of this source is de-magnified according to the magnification factor of $\mu(\text{dust}) = 4.9 \pm 1.8$ from Timmons et al. (2016) and considered in the cumulative number counts. Only slight overdensities can be seen at 6.0 and 8.0 mJy in the cumulative number counts. We observe 3.8

sources with flux densities above 8.0 mJy in the map. The probability of observing this number of sources in a random field ($P(\geq 3.8)$) is 0.32.

NGP2: Overdensities are seen from 6.0 to 12.0 mJy in the cumulative number counts. We observe 6.5 sources with flux densities above 8.0 mJy in the map. The probability of observing this number of sources in a random field ($P(\geq 6.5)$) is 0.02.

NGP3: Overdensities are seen at 8.0, 12.0 and 16.0 mJy for cumulative number counts. We observe 5.1 sources with flux densities above 8.0 mJy in the map. The probability of observing this number of sources in a random field ($P(\geq 5.1)$) is 0.05.

NGP4: Overdensities are seen from 6.0 to 12.0 mJy in the cumulative number counts. We observe 17 sources with flux densities above 8.0 mJy in the map. The probability of observing this number of sources in a random field ($P(\geq 17)$) is 4.46×10^{-11} .

NGP5: Cumulative number counts show overdensities from 6.0 to 14.0 mJy. We observe 5.7 sources with flux densities above 8.0 mJy in the map. The probability of observing this number of sources in the field ($P(\geq 5.7)$) is 0.05.

NGP6: No overdensity is seen at 10.0 mJy and slight overdensities can be seen at 6.0 mJy, in the cumulative number counts. We observe one source with a flux density above 8.0 mJy in the map. The probability of observing this number of sources in a random field ($P(\geq 1)$) is 0.86. We do not include the flux density bin of 4.0 mJy since the effective area correction becomes unreliable at fainter flux densities, and the number counts need further investigation.

Among the SCUBA-2 detected sources, NGP6.02 is a Herschel-SPIRE dropout, which has no *Herschel* counter-

part and is believed to be either a $z > 6$ SMG or a cool $z = 4$ DSFG (Greenslade et al., submitted). Whether this source is associated with any protocluster or its line-of-sight overlaps needs further investigation.

NGP7: The cumulative number counts show overdensities from 6.0 to 14.0 mJy. We observe 13 sources with flux densities above 8.0 mJy in the map. The probability of observing this number of sources in a random field ($P(\geq 13)$) is 1.75×10^{-7} .

NGP8: Cumulative number counts show overdensities at 8.0 and 10.0 mJy. We observe 16 sources with flux densities above 8.0 mJy in the map. The probability of observing this number of sources in a random field ($P(\geq 16)$) is 3.88×10^{-10} .

NGP9: A slight overdensity is seen in at 8.0 mJy, in cumulative number counts. We observe 3.5 sources with flux densities above 8.0 mJy in the map. The probability of observing this number of sources in a random field ($P(\geq 3.5)$) is 0.32.




Figure 2. Cumulative number counts of the SCUBA-2 sources in our candidate protocluster fields (blue) and in random fields (red) from Geach et al. (2017) (Ge17). Note that the flux densities of the two known strongly lensed objects in G12 and NGP1 are de-magnified and considered. The errorbars of the SCUBA-2 sources in our candidate protocluster fields (blue) are the completeness error, and the errorbars of random fields (red) from Ge17 are Poissonian. The number of sources are scaled to the size of each candidate protocluster field, which is approximately 0.03 deg^2 .

We have estimated the probability of obtaining the observed number of sources in our candidate protoclusters compared to the field assuming the sources are randomly distributed. However, in reality they may not be randomly distributed and may be clustering due to, for example, cosmic variance. We therefore perform a test where 10,000 regions with the same size as the protocluster candidate maps (i.e. $\sim 0.03 \text{ deg}^2$) are randomly placed in the S2CLS/COSMOS field. The same source extraction method is used and we count the number of detections in each random region. We then count the regions that are overdense, i.e. containing more than the observed number of sources in each candidate protocluster fields (e.g. 6 sources $> 8 \text{ mJy}$ in Bootes1). We list the number of these overdense regions ($N_{\text{overdensity}}$) and the overdensity levels, i.e. the fraction of regions more overdense than the protocluster candidates ($P_{\text{overdensity}} = N_{\text{overdensity}}/10,000$), in Table 2.

We found that for the candidate protocluster fields where there are overdensities of $850 \mu\text{m}$ sources, i.e. Bootes1, EGS, G12, NGP2, NGP3, NGP4, NGP5, NGP7 and NGP8, the overdensity level is < 0.1 . Hence even under the potential clustering effect due to cosmic variance, these fields still have significant overdensities of $850 \mu\text{m}$ sources. The result from Clements et al. (2016), which used submm sources in G12, found the overdensity level of 2.5×10^{-2} within the two arcmin radius, consistent with our result of 3.18×10^{-2} .

The uncertainties in flux boosting correction and completeness correction may also affect the significance of number counts. Nonetheless, it is found that the flux boosting correction is consistent among different methods (Geach et al. 2017), and the uncertainties are generally within 1 mJy , especially at brighter flux density bins ($> 5 \text{ mJy}$). The completeness uncertainties are below 6 percent among all candidate protocluster fields, so should not have significant effects on the result of number counts and the overall conclusions made in this paper.

3.2 SCUBA2-Herschel Colours

We estimate the 250-, 350-, and $500\text{-}\mu\text{m}$ flux densities of our SCUBA-2 sources in all 13 candidate protoclusters using photometry data from *Herschel* in the H-ATLAS (Data Release 1 for G12 and Data Release 2 for NGP fields) and HerMES (Data Release 4⁵, except for Bootes1, where we use Data Release 2, due to missing $500\text{-}\mu\text{m}$ photometry information in DR4) surveys (Smith et al. 2012; Roseboom et al. 2010; Wang et al. 2014).

Firstly we match the positions of our SCUBA-2 sources with those in the H-ATLAS or HerMES catalogues, using an 18 arcsec search radius, which matches the SPIRE $250\text{-}\mu\text{m}$ beam from which the *Herschel*-SPIRE catalogues are derived. We then add a random offset to the original SCUBA-2 source positions and then the same matching algorithm is conducted. We find that the search radius with minimum number of spurious matches is $9''$. There are 10 SCUBA-2 sources which have matches in the *Herschel* catalogues with separation between $9''$ and $18''$. We mark these 10 sources (as *cat**) in the source catalogue in Table B1.

If no match is found for a SCUBA-2 source beyond $18''$, we use the flux densities and noise from the maps at 250, 350 and $500 \mu\text{m}$ at the positions of the SCUBA-2 sources.

Using the above method we are able to constrain the *Herschel*-SPIRE flux densities of each SCUBA-2 source in the candidate protocluster fields and extract the 4-band (250, 350, 500, and $850 \mu\text{m}$) photometry. The 250-, 350-, 500- and $850\text{-}\mu\text{m}$ flux densities of all SCUBA-2 sources in all 13 candidate protocluster fields are listed in Table B1.

Once the $850\text{-}\mu\text{m}$ and SPIRE flux densities are obtained, we can derive the colours, a model-independent approach to identify high-redshift DSFGs. Figure 3 shows the $250\mu\text{m}/350\mu\text{m}$ versus $350\mu\text{m}/850\mu\text{m}$ colour-colour plots of the SCUBA-2 sources from the 13 candidate protocluster fields. We also plot the colours derived from template SEDs of the local ULIRG Arp220 (Donley et al. 2007; Rangwala et al. 2011), average SMGs from the ALMA-LABOCA ECDFS Submm Survey (ALESS) (da Cunha et al. 2015), the high- z source HFLS3 (Riechers et al. 2013) and the Cosmic Eyelash (Swinbank et al. 2010), by redshifting these SEDs from $z=0$ to $z=4.5$. These template SEDs are representative of starbursting galaxies or SMGs at various redshifts, and Table 3 summarises their properties.

Based on the redshift track of the template SEDs, we identify the potential $z \geq 2$ sources to have colours $S_{250}/S_{350} \leq 1.3$ and $S_{350}/S_{850} \leq 6.0$, and shown as the red region in Fig.3. Similarly we identify potential $1 < z < 2$ sources to have colours as shown in the blue region in Fig.3.

Among the 13 candidate protocluster fields, Bootes1, G12 and NGP7 all show a number of sources having colours lying within the $z \geq 2$ region. The EGS, Lockman, NGP2, NGP3, NGP4, NGP5, NGP8, NGP9 fields have fewer sources, but their colours still suggest $z \geq 2$. Sources in NGP1 and NGP6, on the other hand, suggest redshifts between 1 and 2.

⁵ The HerMES source catalogues can be downloaded from HeDaM (<http://hedam.lam.fr>).

Table 3. Properties of the model SEDs used in this paper.

Name of model SED	dust temperature	redshift	nature of source
Arp220	66 K	~ 0.018 ^a	local ULIRG
HFLS3	56^{+9}_{-12} K	6.34	starbursting galaxy
ALESS	~ 40 K	1.33-6.12	SMGs
Cosmic Eyelash	30-60 K	2.3	SMG

^a [de Vaucouleurs et al. \(1991\)](#).




Figure 3. Submillimetre colour-colour plots, specifically $250\mu\text{m}/350\mu\text{m}$ versus $350\mu\text{m}/850\mu\text{m}$ plots of 13 candidate protocluster fields observed with SCUBA-2 in this paper. Red points and error bars represent the $850\text{-}\mu\text{m}$ sources in the candidate protocluster fields. Errors are propagated from the 250- , 350- , and $850\text{-}\mu\text{m}$ flux densities of individual sources. Blue, green, purple and grey curves are the redshift tracks on this colour-colour plot from $z=0.0$ to 4.5 with 0.25 steps, using templates of Arp220, ALESS, HFLS3 and Cosmic Eyelash, respectively. Based on the redshift tracks of the template SEDs, we identify colours where sources are potentially at $z \geq 2$ as the red region, and colours where sources are potentially at $1 < z < 2$ as the blue region.

3.3 SED fitting and photometric redshifts

We use the SPIRE 250-, 350-, 500- and SCUBA-2 850- μm flux densities derived in Section 3.2 to estimate the photometric redshifts of the SCUBA-2 sources in the protocluster fields. We use the same model SEDs as in Table 3 to perform the fitting, as these templates are representative of DSFGs with a variety of dust temperatures. Due to the lack of information on the dust temperature of our sources, we expect to obtain different redshift estimates when using different template SEDs. Using a warmer dust-temperature template SED (e.g. Arp220), the resulting redshift is higher, while using a colder template (e.g. ALESS), the resulting redshift is lower. This is referred to as the temperature-redshift degeneracy (Blain 1999b,a), which prevents accurate redshift estimates if observed flux densities only rely on the (redshifted) thermal dust (modified blackbody) SED. Nonetheless, for redder DSFGs (which describes the majority of our SCUBA-2 sources), this effect is minor and only a slight scatter is seen if different template SEDs are used (Iverson et al. 2016). Iverson et al. (2016) also suggest that even with a limited number of templates, the estimated redshifts of the sources can still be accurate.

We perform χ^2 -minimisation using `emcee` (Foreman-Mackey et al. 2013), which is a Python implementation of the affine-invariant ensemble sampler for a Markov chain Monte Carlo (MCMC) procedure (Goodman & Weare 2010). We fit two parameters, redshift z and the normalization factor a . The normalization factor a is the factor by which template flux density values are multiplied and is associated with the luminosity of a source. We constrain the range of parameters to be $0 < z < 15.0$ and $10^{-2} < a < 10^2$, and place flat priors on them. The MCMC chains are composed of 100 walkers, each having 5,000 steps, and a 1,000-step burn-in time is allowed.

In order to obtain initial parameter values for the MCMC fitting, we first perform a χ^2 gridding search over z and a and the parameters with minimised χ^2 are the initial values in the MCMC fitting. Figure 4 shows an example of this χ^2 gridding search. The upper left panel presents the redshifted and normalized SEDs with the observed flux densities and uncertainties. The upper right panel shows likelihood values on the parameter space, z and a . The lower two panels show the probability density functions (PDFs) with respect to z and a , respectively.

In order to marginalize over the four templates and save computational time, we randomly choose 200 samples from the MCMC chains, giving a total of 800 samples for each source. Figure 5 shows the posterior distribution of the parameters, after marginalization over the templates, of a source in the Bootes1 candidate protocluster (Bootes1.04). Black dashed vertical lines in the posterior distribution show the 16th, 50th and 84th percentiles of the distribution. We use the 50th percentile, or the median, as the best-fit value in this study. The 16th and 84th percentiles of the distribution are given as the range of uncertainty.

Photometric redshift estimates with $z > 6$ are regarded as not being robust, due to poor photometry or the redshift-temperature degeneracy. At $z > 6$, the SPIRE and SCUBA-2 bands are also no longer near the peak of the modified black-body SED. Hence, we do not include such sources in the discussion of redshift and of infrared luminosity.

In addition to z and a , we estimate the infrared luminosity (L_{IR}) and the star-formation rate (SFR) for each of the 800 samples, and thereby determine the posterior distribution, best-fit values and uncertainties for each source. The infrared luminosity is included in the triangle diagram in Fig 5. The infrared luminosity is estimated by integrating the template SEDs from 8 to 1,000 μm in the rest frame, given the z and a values for each sample. SFRs are estimated assuming a linear relation between the infrared luminosity (Kennicutt 1998; Kennicutt & Evans 2012), in which the constant factor of $3.89 \times 10^{-44} M_{\odot} \text{yr}^{-1} \text{erg}^{-1} \text{s}$ and a double-power law stellar initial mass function (IMF, Kroupa 2001) are used.

Fig 5 and Table B1 show that even though the four template SEDs span a variety of different dust temperatures, there are robust estimates in terms of z , L_{IR} and SFR for each source. We can see four distinct peaks in the marginalized posterior distribution of a , but these can be attributed to the different normalizations of each template SED and do not affect the conclusions made in this paper.

We notice that there are some sources whose redshifts are poorly estimated, due to the poor photometric data and/or redshift-temperature degeneracy (e.g. Bootes1.07, see Table B1). Nevertheless, they still have robust L_{IR} and SFR estimates, as suggested in Greenslade (2018) (PhD thesis). Although our sources need spectroscopic verifications of their true redshifts, their robust L_{IR} and SFR estimates indicate that they are likely the most luminous DSFGs ($10^{12} L_{\odot} < L_{\text{IR}} < 10^{13} L_{\odot}$) with the most extreme star-formation activity ($100 M_{\odot} \text{yr}^{-1} < \text{SFR} < 1,500 M_{\odot} \text{yr}^{-1}$) within these protoclusters. More photometric data, such as those from millimetre and radio observations of these far-infrared sources would also help to narrow down the photometric redshifts, and on-going observations are being taken or proposed.

The photometric redshift distribution of all the sources in these 13 candidate protoclusters is shown in Fig 6, in black. The results from MacKenzie et al. (2017) are also shown in magenta; this consists of 46 PHz sources (Planck Collaboration et al. 2016c) with *Herschel* overdensities, likely in protoclusters. It can be seen that a significant fraction of our sources lie in the range $2 < z < 3$, which is consistent with the expected peak of cosmic star-formation rate density either in protoclusters or in the field (Clements et al. 2014; Madau & Dickinson 2014). This redshift distribution is also consistent with that of the radio/mid-infrared counterparts of 850 μm sources in the field (Michałowski et al. 2017).

We also apply our MCMC χ^2 -minimisation method to SCUBA-2 maps of a known DSFG-rich protocluster at $z \sim 2$, PCL1002 (Casey et al. 2015; Hung et al. 2016), and 256 sources distributed in the COSMOS field in the S2CLS survey. The original COSMOS map is cropped to the central $1 \text{ deg} \times 1 \text{ deg}$ region in order to exclude the lower sensitivity edges. The resultant photometric redshift distribution is also shown in Fig 6. We note that the photometric redshift distribution peak of our sources in the 13 candidate protoclusters is similar to that of the known $z \sim 2$ protocluster PCL1002 and the field sources in the S2CLS/COSMOS field, all showing peaks at $2 < z < 3$. Again this is expected from the studies of the peak of the cosmic star-formation rate density. Sources in MacKenzie et al. (2017), on the other




Figure 4. An example showing the χ^2 gridding procedure for one of the sources (Bootes1.04). **Upper Left:** Redshifted and normalized SEDs of the model templates with the minimum χ^2 values. Blue is Arp220, green is ALESS, purple is HFLS3, and grey is Cosmic Eyelash. The observed flux densities from SCUBA-2 and SPIRE and their associated uncertainties are shown in red. **Upper Right:** Likelihood values of this source (Bootes1.04) on the parameter space (z and a). **Bottom:** Relative probability density functions (PDFs) as functions of z and a . The vertical black dashed lines represent the 16-th, median, and 84-th quantiles of the PDFs.

hand, show a peak at $3 < z < 4$, but a significant number of sources lie also within $2 < z < 3$.

We note that PCL1002 has the highest surface density of sources in the range $2 < z < 3$, at approximately $0.074 \text{ arcmin}^{-2}$. Our sources in the 13 candidate starbursting protoclusters correspond to approximately $0.027 \text{ arcmin}^{-2}$, which is slightly higher than the number of sources in the S2CLS/COSMOS field. SCUBA-2-detected sources in MacKenzie et al. (2017) peak at only approximately $0.005 \text{ arcmin}^{-2}$.

4 DISCUSSION

4.1 Selection Effects

We compare the results in Gr18 with our results of cumulative number counts shown in Section 3.1. In Gr18, 27 candidate DSFG-rich protoclusters were selected based on overdensities of *Herschel*-SPIRE sources at 250, 350 and 500 μm , whereas in this paper we study number counts at 850 μm .

In order to investigate if different methods select different populations of candidate protoclusters, we classify the 13 candidate protoclusters in this paper into four categories: (i) those selected as candidate protoclusters in Gr18, and having an overdensity of 850- μm sources in this paper; (ii) those selected as candidate protoclusters in Gr18, but not having an overdensity of 850- μm sources in this paper; (iii) those not selected as candidate protoclusters in Gr18, but having an overdensity of 850- μm sources in this paper; and (iv) those neither selected as candidate protoclusters in Gr18,

nor having an overdensity of 850 μm sources in this paper. The category for each of the 13 candidate protoclusters in this paper is listed in Table 1.

Bootes1, EGS, NGP4, NGP5, NGP7, and NGP8 are category (i). These candidate protoclusters all have *Herschel* source overdensities at 350 μm (EGS, NGP5, see Table 1) or 500 μm (Bootes1, NGP4, NGP7, NGP8), along with their 850- μm overdensities, suggesting they are high-redshift protoclusters. The colour-colour plot in Fig 3 and the 4-band photometric redshift estimates also suggest that they are bona-fide protoclusters.

Lockman is in category (ii), since it is selected as a candidate protocluster in Gr18 but shows no 850- μm overdensity. Optical/near-infrared studies on Lockman, using red sequence galaxies, suggest a photometric redshift of $z = 2.05 \pm 0.09$ (Clements et al. 2014). After cross-matching with *Herschel* sources, the 350 μ /850 μm versus 250 μ /350 μm colour-colour plot suggests that the two SCUBA-2 sources are at $z > 2$ (Fig 3), and their photometric redshift estimates suggest $z = 2.5^{+0.7}_{-0.6}$ and $z = 3.1^{+0.7}_{-0.4}$. However, there are only two 850- μm sources in this field, and for *Herschel* sources alone it is most overdense at 250 μm (5.4σ , see Table 1). We conclude that Lockman is a lower-redshift ($z < 2$) cluster/protocluster, which has an overdensity of 250- μm sources but such sources become too faint to be detected by SCUBA-2 at 850 μm . There is also a possibility that our SCUBA-2 maps are not deep enough to detect all of its 850- μm sources.

G12, NGP2 and NGP3 are in category (iii), showing no significant overdensity in any *Herschel*-SPIRE band (250, 350 or 500 μm) in Gr18, but they are overdense at




Figure 5. Posterior distribution of the parameters z , a and L_{IR} for the example source (Bootes1.04) after marginalizing over the four template SEDs. 100 walkers are used, each having 5,000 steps in the constrained parameter ranges $0 < z < 15$ and $10^{-2} < a < 10^2$. We choose to implement a burn-in period of 1,000, which are not considered in the fitting. The posterior distributions show 200 randomly selected samples for each template, resulting in 800 samples in total for each source. The black vertical dashed lines indicate the 16th, 50th and 84th percentiles. The best-fit values and uncertainties are given above each posterior distribution.

$850\mu\text{m}$ seen in this paper. According to the colour-colour plot (Fig 3) and photo- z estimates (Table B1), we suspect they are high-redshift ($z > 2$) protoclusters, rich in $850\text{-}\mu\text{m}$ sources. G12 has a spectroscopically confirmed lensed DSFG at $z=3.26$ at the centre of the field (Fu et al. 2012; Herranz et al. 2013), so is classified as a lensed source in Gr18, and only has a 2.8σ overdensity at $350\mu\text{m}$. However, Clements et al. (2016) found that G12 is more overdense at submillimetre wavelengths using SCUBA-2 ($850\mu\text{m}$) and LABOCA ($870\mu\text{m}$) observations, consistent with the results shown in this paper. Hence we suspect G12 is a DSFG-rich protocluster where one of the DSFGs happens to be lensed by foreground galaxies and contributes to the bright far-infrared emission seen in *Herschel*-SPIRE and SCUBA-2. NGP2 and NGP3, on the other hand, are probably also high-redshift ($z > 2$) protoclusters that are rich in $850\text{-}\mu\text{m}$ sources.

NGP1, NGP6 and NGP9 are in category (iv), having neither a *Herschel* source overdensity nor $850\text{-}\mu\text{m}$ overdensity. They might, nevertheless, still be protoclusters, with source overdensities that are too faint to be seen with the observed sensitivity of our *Herschel* or SCUBA-2 observations. Only deeper observations and/or multi-wavelength follow-

up observations can determine whether they are protoclusters. Note that even though NGP1 has a 3.2σ overdensity at $500\mu\text{m}$, it is classified as a lensed far-infrared source in Gr18 due to the bright *Herschel* source in the central region of this field, which is also a confirmed lensed object (Bussmann et al. 2013; Calanog et al. 2014; Timmons et al. 2016). Thus NGP1 is not classified as (ii).

4.2 Infrared Luminosities and Star-formation Rates

As discussed in Section 3.3, we estimate the infrared luminosity and SFR for each SCUBA-2 source by integrating the SED from 8 to $1,000\mu\text{m}$ (see Table B1). Fig 7 shows the L_{IR} distribution for all the sources in our 13 candidate protoclusters, compared with a known DSFG-rich protocluster at $z \sim 2$, PCL1002, and sources in candidate protoclusters from MacKenzie et al. (2017). We have scaled the number of sources to per arcminute^2 in order to compare fields with different areas.

While PCL1002 has a peak in the bin of $12.5 <$




Figure 6. Photometric redshift distribution of sources in 13 candidate protocluster fields in this paper (black), a known DSFG-rich protocluster at $z \sim 2$, PCL1002 (cyan), 256 sources in the COSMOS field in the S2CLS survey (red), and sources from another *Planck*-based protocluster candidate survey (magenta, MacKenzie et al. 2017). $z > 6$ sources are not included because their photometry no longer covers the peak of the redshifted SED, and their photometric redshifts are considered not robust.




Figure 7. Distribution of infrared luminosity (L_{IR}) of sources in our 13 candidate protoclusters (black), a known DSFG-rich protocluster at $z \sim 2$, PCL1002 (cyan), and sources in candidate protoclusters in MacKenzie et al. (2017) (magenta). SFRs are also shown in the upper horizontal axis following a linear relation with the L_{IR} (Kennicutt 1998).

$\log_{10}(L_{\text{IR}}/L_{\odot}) < 12.75$ ($3.2 \times 10^{12} < L_{\text{IR}} < 5.6 \times 10^{12} L_{\odot}$), our sources in 13 candidate protoclusters show a higher luminosity peak. Nonetheless, a significant number of sources in our candidate protoclusters have infrared luminosities in the same range as the peak in PCL1002. Sources in MacKenzie et al. (2017) tend to be more luminous. We conclude that the $850 \mu\text{m}$ sources in our candidate protoclusters are as luminous as those of known protoclusters (such as PCL1002), and have representative infrared luminosities just below $10^{13} L_{\odot}$. We do not totally rule out the possibility of selection effects in these different sets of (candidate) protoclusters (Cheng et al. in prep.), but the study of such selection effect is beyond the scope of this paper.

As discussed in Section 3.3, we assume a linear relation between the L_{IR} and the star-formation rate (SFR) (Kennicutt 1998; Kennicutt & Evans 2012), with the constant factor being $3.89 \times 10^{-44} M_{\odot} \text{yr}^{-1} \text{erg}^{-1} \text{s}$. In the upper horizontal axis in Fig 7, we also quote the SFR values. Similarly the SFRs of the $850 \mu\text{m}$ sources in our candidate protoclusters are consistent with the known protocluster PCL1002, and have representative SFRs of $500 M_{\odot} \text{yr}^{-1} < \text{SFR} < 1,500 M_{\odot} \text{yr}^{-1}$. There are also a substantial number of sources that have $\text{SFR} > 1,000 M_{\odot} \text{yr}^{-1}$, which might constitute galaxies with extreme star-formation activity or lensed objects.

In confirmed protoclusters at $z > 2$, including PCL1002, there are cases of spectroscopically confirmed DSFGs or starbursts having L_{IR} and SFR values within the range we find here (Dannerbauer et al. 2014; Casey et al. 2015; Umehata et al. 2015). This suggests that even without accurate redshift estimates, approximate infrared luminosities and SFRs can be used to select starbursting members in candidate protoclusters, and this might be applied in general to protoclusters at $z > 2$ (Greenslade, 2018, PhD Thesis ⁶).

4.3 Total SFR and protocluster size

We also estimate the properties of the candidate protoclusters as a whole, by first selecting candidate protocluster member galaxies that have photometric redshifts $z < 6$ and estimating their weighted-mean redshifts. Table 4 shows the derived properties of the 13 candidate protoclusters studied in this paper. The weighted-mean redshifts are between $1.8 \leq z \leq 3.2$, consistent with them being high-redshift protoclusters.

The total SFRs are estimated by summing up the individual SFR of the candidate member galaxies. Since the SFRs are derived using FIR and submm flux densities, the total SFRs may be underestimated due to (partial) dust obscuration. These SFRs should also be considered lower limits since we may miss faint sources in the background that are undetected by SCUBA2. We also estimate the upper limit of the total SFR of each candidate protocluster field by first extracting their *Planck* flux densities at 857, 545, and 352 GHz bands (i.e. 350, 550, and $850 \mu\text{m}$, respectively) from the *Planck* Early Release Compact Source Catalog (ERCSC Planck Collaboration et al. 2011) and the *Planck* Catalogues of Compact Sources (PCCS, PCCS2, Planck Collaboration et al. 2014, 2016a). We then use this 3-band photometry to estimate their SFRs as is performed for the candidate protocluster fields.

The total obscured SFRs of the candidate protoclusters range from approximately 1,600 to $22,500 M_{\odot} \text{yr}^{-1}$ and upper limits for the total SFRs range from approximately 15,400 to $44,000 M_{\odot} \text{yr}^{-1}$. As a reference, the total SFRs of two known protoclusters at $2 < z < 3$ are between 4,900 and $12,500 M_{\odot} \text{yr}^{-1}$ (Lacaille et al. 2019). SFRs of other known protoclusters are $6,500 M_{\odot} \text{yr}^{-1}$ for DRC ($z \sim 4$, Oteo et al. 2018), $> 1,500 M_{\odot} \text{yr}^{-1}$ for AzTEC-3 ($z \sim 5.3$, Capak et al. 2011), and $\sim 3,400 M_{\odot} \text{yr}^{-1}$ for CLJ1001 ($z \sim 2.5$, Wang et al. 2016). The four candidate protoclusters studied in Clements et al. (2014) have SFRs between 620–11,632 $M_{\odot} \text{yr}^{-1}$. Three

⁶ <https://spiral.imperial.ac.uk/handle/10044/1/65836>

Table 4. Properties of the 13 candidate protoclusters as derived from individual photometric redshifts and SFRs. **N**: number of candidate members selected as $z < 6$ sources (see Table B1). z : The weighted-mean redshift from the photometric redshift estimates of individual candidate members. Errors are the standard errors on the weighted mean. **Total SFRs** are estimated by summing up the individual SFRs and **upper limits** are estimated by fitting the photometry of the *Planck* sources (see text). **Angular size**, in arcmin, is estimated by the maximum separation of the candidate members. **Physical size**, in Mpc, is estimated assuming the candidate protocluster is at the weighted-mean redshift. **Physical volume** is estimated assuming the candidate protoclusters are spherical and the physical size as the diameter. **SFRD** and its upper limits are estimated by $\text{SFR}/(\text{physical volume})$.

Name	N	z	Total SFR (upper limit) [$M_{\odot}\text{yr}^{-1}$]	Angular size [arcmin]	Physical size [Mpc]	Physical volume [Mpc^3]	SFRD (upper limit) [$M_{\odot}\text{yr}^{-1}Mpc^{-3}$]
Bootes1	8	2.5 ± 0.32	7117^{+1821}_{-1394} (43946)	7.0	3.4	14.9	478 (2949)
EGS	4	2.6 ± 0.41	5263^{+1475}_{-1121} (20150)	2.8	1.4	1.0	5279 (20150)
G12	12	2.7 ± 0.25	22471^{+7311}_{-5365} (35145)	8.8	4.2	29.1	772 (1207)
Lockman	2	2.8 ± 0.6	1644^{+651}_{-500} (15415)	8.6	4.1	26.6	61 (579)
NGP1	4	1.8 ± 0.41	8019^{+6114}_{-6701} (24437)	4.0	2.0	3.3	2399 (7405)
NGP2	5	2.8 ± 0.41	6079^{+1949}_{-1166} (18019)	4.9	2.3	4.7	1292 (3833)
NGP3	1	2.6 ± 0.88	1885^{+1236}_{-668} (18911)	N/A	N/A	N/A	N/A (N/A)
NGP4	5	3.2 ± 0.48	6333^{+1672}_{-1237} (19083)	3.7	1.7	1.8	3556 (10601)
NGP5	5	3.0 ± 0.4	7052^{+1759}_{-1314} (30761)	4.0	1.9	2.5	2826 (12304)
NGP6	6	2.0 ± 0.32	4128^{+1236}_{-1100} (24847)	5.5	2.8	8.2	502 (3030)
NGP7	10	2.9 ± 0.3	11527^{+2278}_{-2167} (19153)	6.3	3.0	10.2	1131 (1877)
NGP8	4	2.4 ± 0.45	4319^{+1586}_{-1097} (21935)	6.2	3.0	11.1	389 (1976)
NGP9	3	3.2 ± 0.65	2806^{+923}_{-866} (19567)	7.0	3.2	12.6	222 (1552)

of them are also studied in this paper (Bootes1, EGS, Lockman). The SFRs of Bootes1 and Lockman are lower than those estimated in Clements et al. (2014) whereas the SFR of EGS is higher. We suspect this discrepancy is due to the different methods and sources used to estimate the SFRs. In Clements et al. (2014) the SFRs are estimated by fitting a modified blackbody with the dust emissivity of $\beta = 2$, and the *Herschel* sources with 250, 250 and 500 μm flux densities are used.

Due to limited source counts, we approximate the size of each candidate protocluster field by the largest angular separation between any pair of candidate member galaxies, and then converting that angular separation to the physical separation at the weighted-mean redshift. The angular separations of the candidate protocluster fields range from 2.8 to 8.8 arcmin, which corresponds to physical separations of 1.4 to 4.2 Mpc. Note that NGP3 has only one candidate member galaxy so we cannot estimate its size. Our candidate protoclusters are likely to span a wide range of physical sizes, which is consistent with the observed diversity of protoclusters in the literature, from the smallest cores of approximately 80 kpc (CLJ1001, $z \sim 2.5$, Wang et al. 2016) to large scale filament of approximately 60 Mpc (comoving scale) (SSA22, $z \sim 3.1$, Hayashino et al. 2004).

Assuming the candidate protocluster fields are spherical, we also estimate their physical volumes by applying the physical separation as the diameter. The SFRDs and their upper limits are also estimated from the total obscured SFRs and their upper limits. The physical volumes of the candidate protoclusters range from 1.0 to 29.1 Mpc^3 . The SFRDs of the candidate protoclusters range from 61 to 5,279 $M_{\odot}\text{yr}^{-1}Mpc^{-3}$ with upper limits of 579-20,150 $M_{\odot}\text{yr}^{-1}Mpc^{-3}$.

The SFRDs of our candidate protoclusters are higher than low-redshift or local galaxy clusters, which is consistent with the peak of field SFRD at $2 < z < 3$ (Fig.15 in Clements et al. (2014)). These SFRDs indicate that our sample contains some of the most extreme protocluster pop-

ulation, possibly contributing to a large fraction of cosmic SFR at $z > 2$.

5 CONCLUSIONS

Combining *Planck* and *Herschel* data has revealed a number of candidate protoclusters hosting multiple dusty star-forming galaxies (DSFGs). The abundance of these sources makes these protoclusters some of the most extreme star-forming environments in the Universe. DSFGs are thought to be the progenitors of massive elliptical galaxies residing in the cores of today's massive galaxy clusters. Hence studies of these protocluster galaxies are important in understanding the formation of galaxy clusters and elliptical galaxies. However, only a few high- z protoclusters have been found to date. In this paper we examined 13 candidate protoclusters selected using *Planck* and *Herschel* data and then observed with SCUBA-2.

We calculate the cumulative number counts of SCUBA-2 sources in these candidate protocluster fields. Compared to studies of random fields, nine of our candidate protocluster fields show overdensities of DSFGs.

Combining with the 250-, 350- and 500- μm flux densities of these SCUBA-2 sources, we estimate their 250 μm /350 μm versus 350 μm /850 μm colours. 11/13 of the candidate protoclusters have colours similar to those of the template SEDs of known starbursts (Arp220, ALESS, HFSL3 and Cosmic Eyelash) redshifted to $z > 2$.

We estimate photometric redshifts using the 250-, 350-, 500-, and 850- μm flux densities using the same template SEDs and a χ^2 -minimization method. We estimate the infrared luminosity (L_{IR}) of each source by integrating the template SEDs from 8 to 1,000 μm and determine star-formation rates (SFRs) assuming a linear relation to the infrared luminosity. The redshift distributions of all our sources peak at $2 < z < 3$, which is consistent with the redshift distribution of a known protocluster and the peak of star-formation rate density. We found that the infrared lumi-

nosities and SFRs of the sources in our candidate protoclusters are also consistent with those of known protoclusters, and have representative values of $3 \times 10^{12} < L_{\text{IR}} < 10^{13} L_{\odot}$ and $500 M_{\odot} \text{yr}^{-1} < \text{SFR} < 1,500 M_{\odot} \text{yr}^{-1}$, respectively. A substantial number of sources in our candidate protoclusters have $\text{SFR} > 1,000 M_{\odot} \text{yr}^{-1}$, suggesting they are starbursting galaxies with some of the most extreme star-forming activity.

We compare our 13 candidate protocluster sample with the 27 candidate protoclusters selected in Gr18. Six of our 13 candidate protoclusters are also selected in Gr18, suggesting they are the most likely bona-fide high-redshift starbursting protoclusters. Lockman is selected by Gr18 but does not have an 850- μm source overdensity, suggesting it is a lower-redshift protocluster. Three of our 13 candidate protoclusters, which also show overdensities of 850- μm sources, are not selected in Gr18. One of these has a lensed DSFG in the field (G12) and the other two (NGP2, NGP3) appear to be protoclusters rich in 850- μm sources at similarly high redshift. Three other candidate protoclusters are not selected in Gr18 and do not have significant overdensities of 850- μm sources, so are less likely to be true protoclusters, or perhaps the *Herschel* or SCUBA-2 observations are not deep enough to detect associated overdensities of far-infrared or submillimeter sources.

The total obscured SFRs of the candidate protoclusters are estimated and range from approximately 1,600 to 22,500 $M_{\odot} \text{yr}^{-1}$ with upper limits of approximately 15,000 to 44,000 $M_{\odot} \text{yr}^{-1}$. We also estimate their physical sizes and their SFRDs, concluding our sample contains some of the most extreme protocluster population, possibly contributing a large fraction of cosmic star-formation rate at $z > 2$. Future deeper, higher-resolution, multi-wavelength observations (e.g. ALMA, VLT and HST) will help us to understand these early stages of forming galaxy clusters. Those observations include looking for overdensities of optical/near-infrared and mid-infrared sources in these candidate protoclusters, studying the multiplicity rate of the FIR sources, possible weak lensing effects, and potential line-of-sight overlaps of multiple protoclusters in dense DSFGs in the sky. Spectroscopic verifications are also necessary to confirm their protocluster memberships.

ACKNOWLEDGEMENTS

The author appreciates the comments from anonymous referees, M. Zemcov and other collaborators for their insightful comments.

The James Clerk Maxwell Telescope is operated by the East Asian Observatory on behalf of the National Astronomical Observatory of Japan, Academia Sinica Institute of Astronomy and Astrophysics, the Korea Astronomy and Space Science Institute, and the Operation, Maintenance and Upgrading Fund for Astronomical Telescopes and Facility Instruments, budgeted from the Ministry of Finance (MOF) of China and administrated by the Chinese Academy of Sciences (CAS), as well as the National Key R&D Program of China (No. 2017YFA0402700). Additional funding support is provided by the Science and Technology Facilities Council of the United Kingdom and participating universities in the United Kingdom and Canada.

The *Herschel* spacecraft was designed, built, tested,

and launched under a contract to ESA managed by the *Herschel/Planck* Project team by an industrial consortium under the overall responsibility of the prime contractor Thales Alenia Space (Cannes), and including Astrium (Friedrichshafen) responsible for the payload module and for system testing at spacecraft level, Thales Alenia Space (Turin) responsible for the service module, and Astrium (Toulouse) responsible for the telescope, with in excess of a hundred subcontractors.

SPIRE has been developed by a consortium of institutes led by Cardiff University (UK) and including: Univ. Lethbridge (Canada); NAOC (China); CEA, and LAM (France); IFSI, Univ. Padua (Italy); IAC (Spain); Stockholm Observatory (Sweden); Imperial College London, RAL, UCL-MSSL, UKATC, and Univ. Sussex (UK); and Caltech, JPL, NHSC, and Univ. Colorado (USA). This development has been supported by national funding agencies: CSA (Canada); NAOC (China); CEA, CNES, CNRS (France); ASI (Italy); MCINN (Spain); SNSB (Sweden); STFC, UKSA (UK); and NASA (USA).

This research has made use of data from HerMES project (<http://hermes.sussex.ac.uk/>). HerMES is a *Herschel* Key Programme utilising Guaranteed Time from the SPIRE instrument team, ESAC scientists and a mission scientist. The HerMES data was accessed through the *Herschel* Database in Marseille (HeDaM - <http://hedam.lam.fr>) operated by CeSAM and hosted by the Laboratoire d’Astrophysique de Marseille.

GDZ gratefully acknowledges financial support from ASI/INAF agreement n. 2014-024-R.1 for the *Planck* LFI Activity of Phase E2 and from the ASI/Physics Department of the university of Roma–Tor Vergata agreement n. 2016-24-H.0

JGN acknowledges financial support from the I+D 2015 project AYA2015- 65887-P (MINECO/FEDER) and from the Spanish MINECO for a “Ramon y Cajal” fellowship (RYC-2013-13256).

E.I. acknowledges partial support from FONDECYT through grant N° 1171710.

M.J.M. acknowledges the support of the National Science Centre, Poland through the POLONEZ grant 2015/19/P/ST9/04010 and SONATA BIS grant 2018/30/E/ST9/00208; this project has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 665778.

D.R. acknowledges support from the National Science Foundation under grant number AST-1614213.

H.D. acknowledges financial support from the Spanish Ministry of Science, Innovation and Universities (MICIU) under the 2014 Ramón y Cajal program RYC-2014-15686 and AYA2017-84061-P, the later one co-financed by FEDER (European Regional Development Funds).

REFERENCES

- Aihara H., et al., 2018a, *PASJ*, **70**, S4
- Aihara H., et al., 2018b, *PASJ*, **70**, S8
- Andreon S., Newman A. B., Trinchieri G., Raichoor A., Ellis R. S., Treu T., 2014, *A&A*, **565**, A120
- Asboth V., et al., 2016, *MNRAS*, **462**, 1989

- Aversa R., Lapi A., de Zotti G., Shankar F., Danese L., 2015, *ApJ*, **810**, 74
- Blain A. W., 1999a, in Weymann R., Storrie-Lombardi L., Sawicki M., Brunner R., eds, *Astronomical Society of the Pacific Conference Series Vol. 191, Photometric Redshifts and the Detection of High Redshift Galaxies*. p. 255 ([arXiv:astro-ph/9906141](https://arxiv.org/abs/astro-ph/9906141))
- Blain A. W., 1999b, *MNRAS*, **304**, 669
- Brodwin M., Gonzalez A. H., Moustakas L. A., Eisenhardt P. R., Stanford S. A., Stern D., Brown M. J. I., 2007, *ApJ*, **671**, L93
- Bussmann R. S., et al., 2013, *ApJ*, **779**, 25
- Bussmann R. S., et al., 2015, *ApJ*, **812**, 43
- Cai Z.-Y., et al., 2013, *ApJ*, **768**, 21
- Calanog J. A., et al., 2014, *ApJ*, **797**, 138
- Capak P. L., et al., 2011, *Nature*, **470**, 233
- Casey C. M., 2016, *ApJ*, **824**, 36
- Casey C. M., Narayanan D., Cooray A., 2014, *Phys. Rep.*, **541**, 45
- Casey C. M., et al., 2015, *ApJ*, **808**, L33
- Chapin E. L., Berry D. S., Gibb A. G., Jenness T., Scott D., Tilanus R. P. J., Economou F., Holland W. S., 2013, *MNRAS*, **430**, 2545
- Chapman S. C., Blain A. W., Smail I., Ivison R. J., 2005, *ApJ*, **622**, 772
- Chapman S. C., Blain A., Ibatova R., Ivison R. J., Smail I., Morrison G., 2009, *ApJ*, **691**, 560
- Chiang Y.-K., Overzier R., Gebhardt K., 2013, *ApJ*, **779**, 127
- Clements D. L., et al., 2010, *A&A*, **518**, L8
- Clements D. L., et al., 2014, *MNRAS*, **439**, 1193
- Clements D. L., et al., 2016, *MNRAS*, **461**, 1719
- Cook M., Barausse E., Evoli C., Lapi A., Granato G. L., 2010, *MNRAS*, **402**, 2113
- Daddi E., et al., 2009, *ApJ*, **694**, 1517
- Dannerbauer H., et al., 2014, *A&A*, **570**, A55
- Dempsey J. T., et al., 2013, *MNRAS*, **430**, 2534
- Donley J. L., Rieke G. H., Pérez-González P. G., Rigby J. R., Alonso-Herrero A., 2007, *ApJ*, **660**, 167
- Douglas L. S., Bremer M. N., Lehnert M. D., Stanway E. R., Milvang-Jensen B., 2010, *MNRAS*, **409**, 1155
- Dowell C. D., et al., 2014, *ApJ*, **780**, 75
- Eales S., et al., 2010, *PASP*, **122**, 499
- Eisenhardt P. R. M., et al., 2008, *ApJ*, **684**, 905
- Fan L., Lapi A., De Zotti G., Danese L., 2008, *ApJ*, **689**, L101
- Farrah D., et al., 2006, *ApJ*, **641**, L17
- Flores-Cacho I., et al., 2016, *A&A*, **585**, A54
- Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, *PASP*, **125**, 306
- Fu H., et al., 2012, *ApJ*, **753**, 134
- Geach J. E., et al., 2017, *MNRAS*, **465**, 1789
- Gómez-Guijarro C., et al., 2018, *ApJ*, **856**, 121
- Gómez-Guijarro C., et al., 2019, *ApJ*, **872**, 117
- Goodman J., Weare J., 2010, *Communications in Applied Mathematics and Computational Science*, Vol. 5, No. 1, p. 65-80, 2010, **5**, 65
- Granato G. L., De Zotti G., Silva L., Bressan A., Danese L., 2004, *ApJ*, **600**, 580
- Granato G. L., Ragone-Figueroa C., Domínguez-Tenreiro R., Obreja A., Borgani S., De Lucia G., Murante G., 2015, *MNRAS*, **450**, 1320
- Greenslade J., 2018, The nature and evolution of high redshift dusty star forming galaxies, <https://spiral.imperial.ac.uk/8443/handle/10044/1/65836>
- Greenslade J., et al., 2018, *MNRAS*, **476**, 3336
- Griffin M. J., et al., 2010, *A&A*, **518**, L3
- Grupponi C., et al., 2015, *MNRAS*, **451**, 3419
- Hayashi M., Kodama T., Tadaki K.-i., Koyama Y., Tanaka I., 2012, *ApJ*, **757**, 15
- Hayashino T., et al., 2004, *AJ*, **128**, 2073
- Herranz D., et al., 2013, *A&A*, **549**, A31
- Hill R., et al., 2018, *MNRAS*, **477**, 2042
- Hodge J. A., Carilli C. L., Walter F., Daddi E., Riechers D., 2013, *ApJ*, **776**, 22
- Holland W. S., et al., 2013, *MNRAS*, **430**, 2513
- Hopkins A. M., Beacom J. F., 2006, *ApJ*, **651**, 142
- Hopkins P. F., Hernquist L., Cox T. J., Kereš D., 2008, *ApJS*, **175**, 356
- Hughes D. H., et al., 1998, *Nature*, **394**, 241
- Hung C.-L., et al., 2016, *ApJ*, **826**, 130
- Husband K., Bremer M. N., Stanway E. R., Davies L. J. M., Lehnert M. D., Douglas L. S., 2013, *MNRAS*, **432**, 2869
- Ivison R. J., et al., 2016, *ApJ*, **832**, 78
- Joseph R. D., Wright G. S., 1985, *MNRAS*, **214**, 87
- Kennicutt Jr. R. C., 1998, *ARA&A*, **36**, 189
- Kennicutt R. C., Evans N. J., 2012, *ARA&A*, **50**, 531
- Kneissl R., et al., 2018, arXiv e-prints, p. arXiv:1804.06581
- Kroupa P., 2001, *MNRAS*, **322**, 231
- Kurk J. D., Pentericci L., Overzier R. A., Röttgering H. J. A., Miley G. K., 2004, *A&A*, **428**, 817
- Lacaille K. M., et al., 2019, *MNRAS*, **488**, 1790
- Lamarre J. M., et al., 2010, *A&A*, **520**, A9
- Lapi A., Shankar F., Mao J., Granato G. L., Silva L., De Zotti G., Danese L., 2006, *ApJ*, **650**, 42
- Lapi A., et al., 2011, *ApJ*, **742**, 24
- Lapi A., Raimundo S., Aversa R., Cai Z.-Y., Negrello M., Celotti A., De Zotti G., Danese L., 2014, *ApJ*, **782**, 69
- MacKenzie T. P., et al., 2017, *MNRAS*, **468**, 4006
- Madau P., Dickinson M., 2014, *ARA&A*, **52**, 415
- Martinache C., et al., 2018, preprint, ([arXiv:1810.07330](https://arxiv.org/abs/1810.07330))
- Matsuda Y., et al., 2005, *ApJ*, **634**, L125
- Michałowski M. J., et al., 2017, *MNRAS*, **469**, 492
- Nayyeri H., et al., 2016, *ApJ*, **823**, 17
- Negrello M., González-Nuevo J., Magliocchetti M., Moscardini L., De Zotti G., Toffolatti L., Danese L., 2005, *MNRAS*, **358**, 869
- Negrello M., et al., 2017, *MNRAS*, **470**, 2253
- Newman A. B., Ellis R. S., Andreon S., Treu T., Raichoor A., Trinchieri G., 2014, *ApJ*, **788**, 51
- Oliver S. J., et al., 2012, *MNRAS*, **424**, 1614
- Oteo I., et al., 2018, *ApJ*, **856**, 72
- Overzier R. A., 2016, *A&ARv*, **24**, 14
- Pentericci L., et al., 2000, *A&A*, **361**, L25
- Pilbratt G. L., et al., 2010, *A&A*, **518**, L1
- Planck Collaboration et al., 2011, *A&A*, **536**, A7
- Planck Collaboration et al., 2014, *A&A*, **571**, A28
- Planck Collaboration et al., 2016a, *A&A*, **594**, A26
- Planck Collaboration et al., 2016b, *A&A*, **594**, A27
- Planck Collaboration et al., 2016c, *A&A*, **596**, A100
- Rangwala N., et al., 2011, *ApJ*, **743**, 94
- Riechers D. A., et al., 2013, *Nature*, **496**, 329
- Roseboom I. G., et al., 2010, *MNRAS*, **409**, 48
- Rowan-Robinson M., et al., 2018, *A&A*, **619**, A169
- Siringo G., et al., 2009, *A&A*, **497**, 945
- Smail I., Ivison R. J., Blain A. W., 1997, *ApJ*, **490**, L5
- Smith A. J., et al., 2012, *MNRAS*, **419**, 377
- Sparre M., Springel V., 2016, *MNRAS*, **462**, 2418
- Steidel C. C., Adelberger K. L., Dickinson M., Giavalisco M., Pettini M., Kellogg M., 1998, *ApJ*, **492**, 428
- Swinbank A. M., et al., 2010, *Nature*, **464**, 733
- Tanaka I., et al., 2011, *PASJ*, **63**, 415
- Timmons N., et al., 2016, *ApJ*, **829**, 21
- Toft S., et al., 2014, *ApJ*, **782**, 68
- Umehata H., et al., 2015, *ApJ*, **815**, L8
- Valiante E., et al., 2016, *MNRAS*, **462**, 3146
- Verhamme A., Schaerer D., Atek H., Tapken C., 2008, *A&A*, **491**, 89
- Walter F., et al., 2012, *Nature*, **486**, 233
- Wang L., et al., 2013, *MNRAS*, **431**, 648


- Wang L., et al., 2014, *MNRAS*, 444, 2870
Wang T., et al., 2016, *ApJ*, 828, 56
Wardlow J. L., et al., 2013, *ApJ*, 762, 59
Wilkinson A., et al., 2017, *MNRAS*, 464, 1380
Zhang Z.-Y., Romano D., Ivison R. J., Papadopoulos P. P., Matteucci F., 2018, *Nature*, 558, 260
da Cunha E., et al., 2015, *ApJ*, 806, 110
de Vaucouleurs G., de Vaucouleurs A., Corwin Jr. H. G., Buta R. J., Paturel G., Fouqué P., 1991, Third Reference Catalogue of Bright Galaxies. Volume I: Explanations and references. Volume II: Data for galaxies between 0^h and 12^h . Volume III: Data for galaxies between 12^h and 24^h .

APPENDIX A: SCUBA-2 850 μ M FLUX MAPS


Fig. A1 shows SCUBA-2 850- μ m flux maps of the 13 candidate protocluster fields discussed in this paper.

APPENDIX B: SOURCE CATALOGUES

Table B1 shows the SCUBA-2 source catalogue of the 13 candidate protoclusters discussed in this paper.




(a) Bootes1




(b) EGS

Figure A1. SCUBA-2 850- μm flux maps of the 13 candidate prorcluster fields. Detected sources ($>3.5 \sigma$) are shown as red circles, labelled with their names. Black contours are *Herschel* 350 μm flux densities, with levels of 10, 20, 30, 40 and 50 mJy.




(c) G12




(d) Lockman

Figure A1. Continued.




(e) NGP1




(f) NGP2

Figure A1. Continued.




(g) NGP3




(h) NGP4

Figure A1. Continued.




(i) NGP5




(j) NGP6

Figure A1. Continued.



(k) NGP7



(l) NGP8

Figure A1. Continued.




Figure A1. Continued.

Table B1. Catalogue of SCUBA-2 sources in the 13 candidate protocluster fields. **Column 4:** signal-to-noise ratio at 850 μm . **Column 5:** $S_{850}(\sigma_{850})$ de-boosted flux densities and uncertainties of the detected SCUBA-2 sources at 850 μm , after the confusion noise of 0.7 mJy and the 5 percent calibrator error are added in quadrature. **Column 6–8:** S_{250} , S_{350} , S_{500} (σ_{250} , σ_{350} , σ_{500}), SPIRE flux densities and uncertainties; if determined from the SPIRE maps (class = map), the uncertainties are quoted as the confusion limits at the respective bands. **Column 9:** if class=*cat*, the SPIRE flux densities and uncertainties are quoted from the *Herschel* catalogues, and *cat** means that the *Herschel* counterpart has a separation between 9" and 18", which has a higher rate of being a spurious match as discussed in 3.2; if class=*map*, the SPIRE flux densities are quoted in the maps at the positions of the SCUBA-2 sources. **Column 10–12:** z , $\log(L_{\text{IR}})$ and $\log(\text{SFR})$, marginalized photometric redshift, (log of) infrared luminosity and (log of) star-formation rates, respectively. These are found by fitting the 250, 350, 500 and 850 flux densities, using the MCMC method. The best-fit is given as the median and errors are quoted as 16th and 84th percentiles. (#): sources in NGP3 where the completeness is below 50 percent.

Name	RA (J2000)	Dec (J2000)	S/N (at 850 μm)	S_{850} (σ_{850}) [mJy]	S_{250} (σ_{250}) [mJy]	S_{350} (σ_{350}) [mJy]	S_{500} (σ_{500}) [mJy]	Class	Redshift	$\log(L_{\text{IR}}[L_{\odot}])$	$\log(\text{SFR}[M_{\odot}\text{yr}^{-1}])$
Bootes1											
Bootes1.01	218.56919	35.54917	7.8	9.9 (1.4)	132.6 (6.3)	96.0 (6.5)	65.9 (6.9)	cat	1.43 ^{+0.58} _{-0.8}	12.94 ^{+0.41} _{-0.35}	3.11 ^{+0.41} _{-0.35}
Bootes1.02	218.52002	35.56137	5.1	7.4 (1.5)	23.8 (6.3)	32.7 (6.5)	32.2 (7.1)	cat	2.72 ^{+0.77} _{-0.44}	12.83 ^{+0.28} _{-0.15}	3.0 ^{+0.28} _{-0.15}
Bootes1.03	218.55963	35.56139	4.6	5.3 (1.2)	9.7 (6.3)	18.7 (6.6)	15.7 (7.2)	map	3.41 ^{+0.88} _{-0.76}	12.67 ^{+0.22} _{-0.19}	2.85 ^{+0.22} _{-0.19}
Bootes1.04	218.65115	35.57692	4.5	7.5 (1.7)	36.2 (6.3)	33.2 (6.5)	21.7 (6.9)	cat	2.22 ^{+0.59} _{-0.58}	12.74 ^{+0.29} _{-0.17}	2.91 ^{+0.29} _{-0.17}
Bootes1.05	218.54048	35.60583	4.5	9.0 (2.1)	17.9 (6.3)	18.4 (6.6)	27.9 (7.2)	map	3.54 ^{+0.9} _{-0.58}	12.91 ^{+0.2} _{-0.18}	3.09 ^{+0.2} _{-0.18}
Bootes1.06	218.60744	35.56916	4.4	5.4 (1.3)	10.1 (6.3)	11.9 (6.6)	9.0 (7.2)	map	4.09 ^{+2.31} _{-1.21}	12.68 ^{+0.23} _{-0.22}	2.85 ^{+0.23} _{-0.22}
Bootes1.07	218.59377	35.55583	4.3	4.6 (1.1)	-9.4 (6.3)	16.7 (6.6)	4.1 (7.2)	map	8.09 ^{+4.66} _{-3.34}	12.82 ^{+0.3} _{-0.29}	2.99 ^{+0.3} _{-0.29}
Bootes1.08	218.57329	35.57917	4.1	4.6 (1.1)	25.8 (6.3)	25.1 (6.6)	26.9 (7.3)	cat	2.15 ^{+0.78} _{-0.71}	12.58 ^{+0.33} _{-0.19}	2.75 ^{+0.33} _{-0.19}
Bootes1.09	218.61564	35.57694	4.1	5.4 (1.4)	-3.1 (6.3)	4.1 (6.6)	3.3 (7.2)	map	10.18 ^{+3.29} _{-3.78}	12.97 ^{+0.26} _{-0.28}	3.15 ^{+0.26} _{-0.28}
Bootes1.10	218.58830	35.49250	4.0	7.1 (1.8)	3.0 (6.3)	1.8 (6.6)	-2.6 (7.2)	map	11.14 ^{+2.89} _{-3.65}	13.12 ^{+0.25} _{-0.3}	3.29 ^{+0.25} _{-0.3}
Bootes1.11	218.65387	35.55692	4.0	5.7 (1.5)	2.8 (6.3)	2.1 (6.6)	-0.6 (7.2)	map	10.54 ^{+3.15} _{-4.01}	13.01 ^{+0.26} _{-0.32}	3.18 ^{+0.26} _{-0.32}
Bootes1.12	218.56374	35.49139	4.0	6.2 (1.6)	23.1 (6.3)	32.6 (6.6)	24.5 (7.2)	cat	2.54 ^{+0.86} _{-0.51}	12.74 ^{+0.32} _{-0.16}	2.91 ^{+0.32} _{-0.16}
EGS1											
EGS.01	216.15980	52.94400	6.4	12.3 (2.0)	43.5 (3.5)	40.4 (4.1)	32.2 (4.6)	cat	2.48 ^{+0.62} _{-0.39}	12.96 ^{+0.23} _{-0.18}	3.13 ^{+0.23} _{-0.18}
EGS.02	216.11187	52.93178	4.5	8.3 (1.9)	24.5 (3.5)	30.4 (4.1)	25.9 (4.4)	cat	2.94 ^{+0.73} _{-0.42}	12.9 ^{+0.22} _{-0.2}	3.08 ^{+0.22} _{-0.2}
EGS.03	216.11369	52.97287	4.5	8.8 (2.0)	31.2 (3.5)	39.2 (4.1)	36.7 (4.4)	cat	2.92 ^{+0.68} _{-0.31}	13.03 ^{+0.2} _{-0.19}	3.2 ^{+0.2} _{-0.19}
EGS.04	216.17088	52.96288	4.0	7.3 (1.9)	59.8 (3.5)	57.9 (4.1)	38.9 (4.6)	cat	2.01 ^{+0.64} _{-0.68}	12.88 ^{+0.33} _{-0.15}	3.05 ^{+0.33} _{-0.15}
G12											
G12.01	176.65833	-0.19222	59.6	79.8 (4.2)	316.0 (6.6)	357.9 (7.4)	291.8 (7.7)	cat	2.6 ^{+0.83} _{-0.15}	13.92 ^{+0.25} _{-0.18}	4.1 ^{+0.25} _{-0.18}
G12.02	176.61722	-0.18556	6.0	9.3 (1.6)	49.0 (6.5)	55.1 (7.4)	27.5 (7.8)	cat	2.11 ^{+0.62} _{-0.66}	12.84 ^{+0.23} _{-0.15}	3.02 ^{+0.23} _{-0.15}
G12.03	176.67389	-0.22556	5.8	8.8 (1.6)	48.7 (6.6)	43.7 (7.5)	40.0 (7.7)	cat	2.2 ^{+0.79} _{-0.82}	12.86 ^{+0.34} _{-0.19}	3.03 ^{+0.34} _{-0.19}
G12.04	176.68611	-0.15778	5.3	9.0 (1.7)	15.9 (7.3)	20.0 (8.1)	22.0 (8.6)	map	3.57 ^{+0.93} _{-0.55}	12.88 ^{+0.21} _{-0.16}	3.05 ^{+0.21} _{-0.16}
G12.05	176.64833	-0.18111	5.0	5.8 (1.2)	54.9 (6.5)	55.4 (7.5)	33.9 (7.8)	cat	1.7 ^{+0.49} _{-1.07}	12.71 ^{+0.34} _{-0.43}	2.88 ^{+0.34} _{-0.43}
G12.06	176.60722	-0.23444	5.0	8.8 (1.8)	13.6 (7.3)	19.5 (8.1)	15.8 (8.6)	map	3.73 ^{+1.01} _{-0.64}	12.87 ^{+0.21} _{-0.2}	3.04 ^{+0.21} _{-0.2}
G12.07	176.68167	-0.13667	4.7	8.5 (1.9)	7.5 (7.3)	21.9 (8.1)	22.3 (8.6)	map	3.99 ^{+0.93} _{-0.64}	12.9 ^{+0.18} _{-0.17}	3.07 ^{+0.18} _{-0.17}
G12.08	176.64167	-0.21889	4.4	5.9 (1.4)	14.8 (7.3)	21.6 (8.1)	20.7 (8.6)	map	3.09 ^{+0.89} _{-0.6}	12.72 ^{+0.24} _{-0.18}	2.89 ^{+0.24} _{-0.18}
G12.09	176.65167	-0.21222	4.4	5.1 (1.2)	1.8 (7.3)	4.6 (8.1)	12.9 (8.6)	map	7.03 ^{+4.54} _{-2.53}	12.82 ^{+0.26} _{-0.25}	2.99 ^{+0.26} _{-0.25}
G12.10	176.61611	-0.12667	4.3	7.5 (1.8)	25.0 (6.1)	20.1 (7.4)	4.8 (7.7)	cat*	2.57 ^{+0.88} _{-0.71}	12.67 ^{+0.25} _{-0.2}	2.84 ^{+0.25} _{-0.2}
G12.11	176.67500	-0.14889	4.2	6.6 (1.6)	24.5 (6.0)	24.1 (7.4)	16.6 (7.7)	cat*	2.61 ^{+0.76} _{-0.61}	12.7 ^{+0.25} _{-0.18}	2.88 ^{+0.25} _{-0.18}
G12.12	176.61611	-0.17222	4.1	6.0 (1.5)	17.5 (7.3)	24.4 (8.1)	11.7 (8.6)	map	2.81 ^{+0.85} _{-0.69}	12.67 ^{+0.26} _{-0.17}	2.84 ^{+0.26} _{-0.17}
G12.13	176.58833	-0.16333	4.0	7.5 (1.9)	0.5 (7.3)	1.6 (8.1)	-10.2 (8.6)	map	11.91 ^{+2.27} _{-3.36}	13.18 ^{+0.15} _{-0.29}	3.35 ^{+0.15} _{-0.29}
G12.14	176.66611	-0.23444	4.0	5.5 (1.4)	-7.3 (7.3)	-3.5 (8.1)	1.1 (8.6)	map	11.31 ^{+2.61} _{-3.51}	13.04 ^{+0.23} _{-0.38}	3.21 ^{+0.23} _{-0.38}
G12.15	176.69611	-0.11778	3.8	6.7 (1.8)	31.0 (6.5)	31.7 (7.5)	27.2 (7.8)	cat	2.39 ^{+0.72} _{-0.64}	12.74 ^{+0.2} _{-0.17}	2.92 ^{+0.2} _{-0.17}
Lockman											
Lockman.01	158.26016	59.15165	4.2	7.1 (1.7)	21.1 (3.8)	18.7 (4.8)	9.4 (4.6)	cat	2.52 ^{+0.69} _{-0.57}	12.61 ^{+0.25} _{-0.2}	2.78 ^{+0.25} _{-0.2}
Lockman.02	158.48582	59.23719	4.0	6.7 (1.7)	20.0 (3.8)	23.4 (4.8)	26.3 (4.7)	cat	3.05 ^{+0.73} _{-0.48}	12.84 ^{+0.23} _{-0.17}	3.02 ^{+0.23} _{-0.17}
NGP1											
NGP1.01	201.11372	28.74656	33.8	42.8 (2.5)	342.3 (5.6)	371.0 (5.9)	250.9 (6.9)	cat	2.07 ^{+0.8} _{-1.54}	13.62 ^{+0.42} _{-0.47}	3.79 ^{+0.42} _{-0.46}
NGP1.02	201.06683	28.74321	5.3	8.5 (1.7)	3.7 (5.6)	4.2 (5.9)	3.5 (7.1)	map	10.01 ^{+3.47} _{-3.44}	13.17 ^{+0.26} _{-0.3}	3.34 ^{+0.26} _{-0.3}

Table B1 – *continued*

Name	RA (J2000)	Dec (J2000)	S/N (at 850 μ m)	S_{850} (σ_{850}) [mJy]	S_{250} (σ_{250}) [mJy]	S_{350} (σ_{350}) [mJy]	S_{500} (σ_{500}) [mJy]	Class	Redshift	$\log(L_{\text{IR}}[L_{\odot}])$	$\log(SFR[M_{\odot}\text{yr}^{-1}])$
NGP1.03	201.09470	28.77211	5.1	7.0 (1.4)	10.0 (5.6)	4.3 (5.9)	9.7 (7.1)	map	6.83 ^{+4.06} _{-0.37}	12.95 ^{+0.22} _{-0.15}	3.12 ^{+0.22} _{-0.15}
NGP1.04	201.12385	28.73989	4.9	5.1 (1.1)	40.9 (5.8)	25.9 (6.0)	0.5 (7.1)	cat	1.53 ^{+0.71} _{-0.39}	12.47 ^{+0.38} _{-0.15}	2.64 ^{+0.38} _{-0.15}
NGP1.05	201.08455	28.78989	4.4	6.1 (1.4)	74.3 (5.7)	53.7 (5.8)	26.0 (7.0)	cat	1.4 ^{+0.5} _{-0.79}	12.67 ^{+0.39} _{-0.35}	2.84 ^{+0.39} _{-0.35}
NGP1.06	201.07569	28.76544	4.3	6.0 (1.4)	-6.8 (5.6)	1.0 (5.9)	4.8 (7.1)	map	10.55 ^{+3.04} _{-3.19}	13.05 ^{+0.24} _{-0.25}	3.22 ^{+0.24} _{-0.25}
NGP1.08	201.05287	28.76543	4.1	6.0 (1.5)	21.1 (5.6)	17.8 (5.9)	14.8 (7.1)	cat*	2.75 ^{+0.85} _{-0.63}	12.66 ^{+0.24} _{-0.19}	2.84 ^{+0.24} _{-0.19}
NGP2											
NGP2.01	199.87603	26.49655	5.9	12.5 (2.2)	69.8 (4.3)	72.9 (4.7)	58.7 (5.9)	cat	2.28 ^{+0.87} _{-0.64}	13.09 ^{+0.34} _{-0.15}	3.26 ^{+0.34} _{-0.15}
NGP2.02	199.88225	26.41544	4.6	9.5 (2.1)	25.9 (4.4)	38.8 (4.8)	33.6 (5.9)	cat	3.05 ^{+0.72} _{-0.45}	13.01 ^{+0.21} _{-0.21}	3.18 ^{+0.21} _{-0.21}
NGP2.03	199.90210	26.47433	4.2	6.3 (1.5)	16.0 (4.4)	17.5 (4.8)	12.6 (6.0)	cat*	3.05 ^{+0.82} _{-0.6}	12.69 ^{+0.23} _{-0.2}	2.86 ^{+0.23} _{-0.2}
NGP2.04	199.87232	26.42766	4.0	7.2 (1.8)	53.5 (4.4)	54.9 (4.7)	50.4 (6.2)	cat*	2.12 ^{+0.79} _{-0.81}	12.86 ^{+0.38} _{-0.16}	3.03 ^{+0.38} _{-0.16}
NGP2.05	199.88969	26.44433	4.0	6.7 (1.7)	7.3 (5.6)	19.0 (5.9)	23.1 (7.1)	map	3.83 ^{+0.93} _{-0.75}	12.8 ^{+0.2} _{-0.19}	2.97 ^{+0.2} _{-0.19}
NGP3											
NGP3.01(#)	202.91419	23.69914	4.2	16.3 (4.0)	-3.0 (5.6)	3.3 (5.9)	6.0 (7.1)	map	11.48 ^{+2.38} _{-2.94}	13.51 ^{+0.23} _{-0.27}	3.69 ^{+0.23} _{-0.27}
NGP3.02	202.96153	23.74802	4.1	8.5 (2.1)	-3.2 (5.6)	1.8 (5.9)	9.6 (7.1)	map	9.96 ^{+3.25} _{-3.13}	13.16 ^{+0.25} _{-0.29}	3.33 ^{+0.25} _{-0.29}
NGP3.03(#)	202.86200	23.71912	4.1	11.1 (2.8)	51.2 (5.6)	63.5 (5.8)	57.5 (7.2)	cat	2.56 ^{+0.78} _{-0.38}	13.1 ^{+0.28} _{-0.15}	3.28 ^{+0.28} _{-0.15}
NGP4											
NGP4.01	198.55701	26.49932	5.3	11.7 (2.3)	26.7 (5.8)	38.3 (5.8)	31.8 (7.5)	cat	3.16 ^{+0.76} _{-0.42}	13.04 ^{+0.21} _{-0.17}	3.21 ^{+0.21} _{-0.17}
NGP4.02	198.59425	26.55044	5.2	10.9 (2.1)	47.8 (5.8)	51.1 (5.9)	44.2 (7.1)	cat	2.41 ^{+0.76} _{-0.61}	12.97 ^{+0.31} _{-0.15}	3.14 ^{+0.31} _{-0.15}
NGP4.03	198.56570	26.50822	4.8	8.3 (1.8)	17.0 (5.6)	21.2 (5.9)	28.9 (7.1)	map	5.65 ^{+4.21} _{-1.82}	12.91 ^{+0.38} _{-0.24}	3.08 ^{+0.38} _{-0.24}
NGP4.04	198.57936	26.49155	4.5	7.2 (1.6)	15.4 (5.5)	7.4 (6.0)	4.5 (7.0)	cat	4.63 ^{+3.5} _{-1.39}	12.81 ^{+0.24} _{-0.22}	2.99 ^{+0.24} _{-0.22}
NGP4.05	198.59798	26.50267	4.4	9.7 (2.3)	35.0 (5.8)	31.4 (5.8)	30.1 (6.9)	cat	12.47 ^{+1.39} _{-2.33}	13.32 ^{+0.22} _{-0.28}	3.49 ^{+0.22} _{-0.28}
NGP4.06	198.55701	26.51710	4.2	8.0 (1.9)	-4.5 (5.6)	7.6 (5.9)	-0.4 (7.1)	map	3.36 ^{+0.82} _{-0.51}	12.88 ^{+0.21} _{-0.16}	3.06 ^{+0.21} _{-0.16}
NGP4.07	198.67126	26.54599	4.1	6.7 (1.7)	-10.7 (5.6)	-14.6 (5.9)	-13.6 (7.1)	map	10.9 ^{+2.99} _{-3.28}	13.08 ^{+0.27} _{-0.28}	3.25 ^{+0.27} _{-0.28}
NGP4.08	198.60419	26.52156	3.9	8.1 (2.1)	5.2 (5.6)	25.3 (5.9)	30.8 (7.1)	map	9.88 ^{+3.25} _{-3.05}	13.12 ^{+0.24} _{-0.26}	3.29 ^{+0.24} _{-0.26}
NGP4.09	198.54955	26.52377	3.8	7.2 (1.9)	-1.2 (5.6)	-6.9 (5.9)	-16.9 (7.1)	map	12.76 ^{+1.54} _{-2.9}	13.16 ^{+0.23} _{-0.3}	3.33 ^{+0.23} _{-0.3}
NGP5											
NGP5.01	205.17118	32.61808	8.2	13.8 (1.8)	30.6 (4.8)	53.5 (5.5)	50.0 (6.5)	cat*	3.19 ^{+0.7} _{-0.34}	13.16 ^{+0.22} _{-0.17}	3.33 ^{+0.22} _{-0.17}
NGP5.02	205.18570	32.64919	5.9	10.3 (1.8)	34.0 (4.9)	32.8 (5.2)	30.6 (6.3)	cat	2.69 ^{+0.68} _{-0.41}	12.91 ^{+0.25} _{-0.17}	3.09 ^{+0.25} _{-0.17}
NGP5.03	205.17646	32.59808	5.2	9.1 (1.8)	0.8 (5.6)	13.0 (5.9)	37.7 (7.1)	map	4.79 ^{+1.17} _{-0.91}	13.0 ^{+0.17} _{-0.19}	3.17 ^{+0.17} _{-0.19}
NGP5.04	205.10917	32.63362	4.4	9.3 (2.2)	36.0 (4.9)	38.1 (5.3)	33.8 (6.4)	cat	2.58 ^{+0.68} _{-0.48}	12.92 ^{+0.25} _{-0.17}	3.09 ^{+0.25} _{-0.17}
NGP5.05	205.19624	32.61030	4.2	6.5 (1.6)	-2.9 (5.6)	-0.2 (5.9)	9.9 (7.1)	map	9.86 ^{+3.32} _{-3.38}	13.03 ^{+0.25} _{-0.29}	3.21 ^{+0.25} _{-0.29}
NGP5.06	205.18174	32.66031	4.0	6.6 (1.7)	30.7 (5.1)	39.3 (5.2)	36.0 (6.4)	cat	2.46 ^{+0.89} _{-0.71}	12.81 ^{+0.34} _{-0.15}	2.98 ^{+0.34} _{-0.15}
NGP5.07	205.20811	32.60363	3.8	6.9 (1.8)	12.6 (4.8)	3.3 (5.2)	1.5 (6.4)	cat*	7.49 ^{+4.7} _{-3.31}	12.9 ^{+0.33} _{-0.32}	3.08 ^{+0.33} _{-0.32}
NGP6											
NGP6.01	200.81323	33.38511	5.4	4.5 (0.9)	5.6 (5.6)	-3.4 (5.9)	10.0 (7.1)	map	9.45 ^{+3.74} _{-4.18}	12.87 ^{+0.26} _{-0.3}	3.05 ^{+0.26} _{-0.3}
NGP6.02	200.74136	33.40399	5.0	10.1 (2.1)	-5.5 (5.6)	3.6 (5.9)	7.5 (7.1)	map	10.35 ^{+3.0} _{-2.94}	13.26 ^{+0.25} _{-0.27}	3.43 ^{+0.25} _{-0.27}
NGP6.03	200.80258	33.38511	4.4	5.3 (1.2)	49.0 (5.6)	45.5 (5.8)	18.3 (7.2)	cat	1.61 ^{+0.45} _{-0.97}	12.62 ^{+0.35} _{-0.35}	2.79 ^{+0.35} _{-0.35}
NGP6.04	200.83452	33.39400	4.2	4.7 (1.1)	35.8 (5.5)	23.3 (5.9)	14.8 (7.2)	cat	1.72 ^{+0.69} _{-0.53}	12.40 ^{+0.35} _{-0.17}	2.67 ^{+0.35} _{-0.17}
NGP6.05	200.80791	33.40956	4.2	5.4 (1.3)	30.1 (5.8)	32.9 (6.0)	17.0 (7.4)	cat	2.1 ^{+0.65} _{-0.53}	12.62 ^{+0.3} _{-0.16}	2.79 ^{+0.3} _{-0.16}
NGP6.06	200.76264	33.43288	4.1	6.6 (1.6)	92.5 (5.6)	67.6 (5.8)	40.2 (6.9)	cat	1.4 ^{+0.53} _{-0.76}	12.78 ^{+0.36} _{-0.36}	2.96 ^{+0.36} _{-0.36}
NGP6.09	200.86513	33.40065	3.9	5.7 (1.5)	3.8 (5.6)	15.8 (5.9)	32.1 (7.1)	map	3.89 ^{+0.96} _{-0.71}	12.78 ^{+0.19} _{-0.17}	2.95 ^{+0.19} _{-0.17}
NGP6.12	200.76397	33.42399	3.8	6.9 (1.9)	33.0 (5.4)	20.7 (6.0)	14.8 (7.1)	cat	2.12 ^{+0.64} _{-0.64}	12.62 ^{+0.27} _{-0.23}	2.8 ^{+0.27} _{-0.23}

Table B1 – *continued*

Name	RA (J2000)	Dec (J2000)	S/N (at 850 μ m)	S_{850} (σ_{850}) [mJy]	S_{250} (σ_{250}) [mJy]	S_{350} (σ_{350}) [mJy]	S_{500} (σ_{500}) [mJy]	Class	Redshift	$\log(L_{\text{IR}}[L_{\odot}])$	$\log(SFR[M_{\odot}\text{yr}^{-1}])$
NGP7											
NGP7.01	204.25680	32.14564	6.6	11.6 (1.8)	59.2 (5.6)	74.5 (5.7)	67.1 (7.3)	cat	2.35 ^{+0.98} _{-0.8}	13.09 ^{+0.34} _{-0.19}	3.26 ^{+0.34} _{-0.19}
NGP7.02	204.23449	32.13785	4.8	11.4 (2.4)	30.3 (5.8)	29.9 (5.8)	16.6 (7.4)	cat	2.79 ^{+0.79} _{-0.54}	12.9 ^{+0.21} _{-0.23}	3.07 ^{+0.21} _{-0.22}
NGP7.03	204.29617	32.18119	4.7	9.2 (2.0)	24.1 (5.7)	26.5 (5.8)	20.3 (7.6)	cat	2.97 ^{+0.77} _{-0.54}	12.86 ^{+0.22} _{-0.2}	3.03 ^{+0.22} _{-0.2}
NGP7.04	204.27517	32.14119	4.6	6.6 (1.5)	23.6 (5.7)	27.7 (5.8)	17.7 (7.4)	cat	2.56 ^{+0.85} _{-0.54}	12.71 ^{+0.3} _{-0.19}	2.88 ^{+0.3} _{-0.19}
NGP7.05	204.34604	32.15229	4.6	10.8 (2.4)	19.1 (5.6)	32.0 (5.9)	29.8 (7.1)	map	3.47 ^{+0.77} _{-0.51}	13.01 ^{+0.19} _{-0.19}	3.19 ^{+0.19} _{-0.19}
NGP7.06	204.31060	32.13675	4.6	7.6 (1.7)	4.8 (5.6)	12.2 (5.9)	11.8 (7.1)	map	5.59 ^{+2.88} _{-1.54}	12.9 ^{+0.23} _{-0.22}	3.07 ^{+0.23} _{-0.22}
NGP7.07	204.23058	32.08896	4.3	8.1 (2.0)	6.4 (5.6)	1.6 (5.9)	-13.1 (7.1)	map	11.85 ^{+2.27} _{-3.18}	13.22 ^{+0.21} _{-0.3}	3.4 ^{+0.21} _{-0.3}
NGP7.08	204.29222	32.07564	4.2	13.1 (3.2)	17.8 (5.6)	23.1 (6.0)	10.1 (7.1)	cat*	3.8 ^{+0.99} _{-0.93}	12.96 ^{+0.21} _{-0.27}	3.13 ^{+0.21} _{-0.27}
NGP7.09	204.28173	32.16675	4.1	6.4 (1.6)	2.7 (5.6)	12.7 (5.9)	28.0 (7.1)	map	4.35 ^{+1.35} _{-0.89}	12.82 ^{+0.2} _{-0.19}	2.99 ^{+0.2} _{-0.19}
NGP7.10	204.23973	32.16008	4.0	6.7 (1.7)	36.6 (5.7)	31.4 (5.9)	19.9 (7.1)	cat	2.08 ^{+0.36} _{-0.59}	12.67 ^{+0.3} _{-0.15}	2.85 ^{+0.3} _{-0.15}
NGP7.11	204.29485	32.15342	4.0	5.7 (1.5)	46.6 (5.8)	55.7 (5.9)	55.7 (7.6)	cat	2.01 ^{+0.67} _{-1.41}	12.81 ^{+0.33} _{-0.57}	2.98 ^{+0.33} _{-0.57}
NGP7.12	204.30403	32.11897	3.9	6.0 (1.6)	1.7 (5.6)	0.8 (5.9)	3.3 (7.1)	map	10.3 ^{+3.31} _{-3.74}	13.0 ^{+0.28} _{-0.3}	3.17 ^{+0.28} _{-0.3}
NGP7.13	204.24760	32.18563	3.9	7.7 (2.0)	-11.0 (5.6)	-11.3 (5.9)	-17.0 (7.1)	map	12.74 ^{+1.64} _{-2.75}	13.21 ^{+0.22} _{-0.31}	3.38 ^{+0.22} _{-0.31}
NGP8											
NGP8.01	202.31798	28.21872	5.0	10.2 (2.1)	29.2 (5.7)	14.0 (6.2)	-0.9 (7.4)	cat*	2.53 ^{+0.88} _{-0.71}	12.69 ^{+0.22} _{-0.27}	2.86 ^{+0.22} _{-0.27}
NGP8.02	202.36968	28.23650	4.5	7.7 (1.8)	49.6 (5.7)	64.7 (6.0)	65.6 (6.9)	cat	2.27 ^{+0.97} _{-0.96}	12.97 ^{+0.36} _{-0.25}	3.15 ^{+0.36} _{-0.25}
NGP8.03	202.39237	28.18538	4.1	9.1 (2.3)	36.4 (5.6)	38.3 (6.0)	35.8 (7.1)	cat	2.56 ^{+0.77} _{-0.47}	12.91 ^{+0.27} _{-0.17}	3.09 ^{+0.27} _{-0.17}
NGP8.04	202.33564	28.19761	3.8	7.6 (2.0)	5.4 (5.6)	10.7 (5.9)	-9.4 (7.1)	map	10.49 ^{+3.24} _{-4.02}	13.08 ^{+0.18} _{-0.33}	3.26 ^{+0.18} _{-0.33}
NGP8.05	202.29527	28.24426	3.7	8.2 (2.3)	40.6 (5.7)	41.8 (6.0)	25.4 (6.9)	cat	2.21 ^{+0.7} _{-0.55}	12.81 ^{+0.3} _{-0.16}	2.99 ^{+0.3} _{-0.16}
NGP9											
NGP9.01	194.74984	31.58490	3.9	8.1 (2.1)	5.1 (5.6)	10.9 (5.9)	13.9 (7.1)	map	5.7 ^{+3.18} _{-1.52}	12.93 ^{+0.24} _{-0.24}	3.11 ^{+0.24} _{-0.24}
NGP9.02	194.87636	31.59490	3.9	7.3 (1.9)	27.7 (5.6)	25.4 (6.2)	26.8 (7.7)	cat	2.63 ^{+0.7} _{-0.57}	12.78 ^{+0.24} _{-0.17}	2.95 ^{+0.24} _{-0.17}
NGP9.03	194.86857	31.64379	3.9	6.0 (1.6)	18.9 (5.5)	6.2 (5.7)	9.4 (6.8)	cat	3.3 ^{+1.59} _{-1.07}	12.63 ^{+0.23} _{-0.25}	2.8 ^{+0.23} _{-0.25}
NGP9.04	194.82157	31.53603	3.8	8.2 (2.2)	17.4 (5.8)	-5.5 (6.0)	-7.6 (7.7)	cat*	11.42 ^{+2.59} _{-4.49}	13.19 ^{+0.26} _{-0.39}	3.36 ^{+0.26} _{-0.39}

This paper has been typeset from a \TeX/L\AA\TeX file prepared by the author.