

## The Zwicky Transient Facility Bright Transient Survey I: Spectroscopic Classification and the Redshift Completeness of Local Galaxy Catalogs

U. C. FREMLING,<sup>1</sup> A. A. MILLER,<sup>2,3</sup> Y. SHARMA,<sup>1</sup> A. DUGAS,<sup>1</sup> D. A. PERLEY,<sup>4</sup> K. TAGGART,<sup>4</sup> J. SOLLERMAN,<sup>5</sup> A. GOOBAR,<sup>6</sup> M. L. GRAHAM,<sup>7</sup> J. D. NEILL,<sup>1</sup> J. NORDIN,<sup>8</sup> M. RIGAUT,<sup>9</sup> R. WALTERS,<sup>1,10</sup> I. ANDREONI,<sup>1</sup> A. BAGDASARYAN,<sup>1</sup> J. BELICKI,<sup>10</sup> C. CANNELLA,<sup>11</sup> ERIC C. BELLM,<sup>12</sup> S. B. CENKO,<sup>13</sup> K. DE,<sup>1</sup> R. DEKANY,<sup>10</sup> S. FREDERICK,<sup>14</sup> V. ZACH GOLKHOV,<sup>12,15</sup> M. GRAHAM,<sup>1</sup> G. HELOU,<sup>16</sup> A. Y. Q. HO,<sup>1</sup> M. KASLIWAL,<sup>1</sup> T. KUPFER,<sup>17</sup> RUSS R. LAHER,<sup>16</sup> A. MAHABAL,<sup>1,18</sup> FRANK J. MASCI,<sup>16</sup> R. RIDDLE,<sup>10</sup> BEN RUSHOLME,<sup>16</sup> S. SCHULZE,<sup>19</sup> DAVID L. SHUPE,<sup>16</sup> R. M. SMITH,<sup>10</sup> LIN YAN,<sup>10</sup> Y. YAO,<sup>1</sup> Z. ZHUANG,<sup>1</sup> AND S. R. KULKARNI<sup>1</sup>

<sup>1</sup>*Division of Physics, Mathematics and Astronomy, California Institute of Technology, Pasadena, CA 91125, USA*

<sup>2</sup>*Center for Interdisciplinary Exploration and Research in Astrophysics and Department of Physics and Astronomy, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208, USA*

<sup>3</sup>*The Adler Planetarium, Chicago, IL 60605, USA*

<sup>4</sup>*Astrophysics Research Institute, Liverpool John Moores University, Liverpool Science Park, 146 Brownlow Hill, Liverpool L35RF, UK*

<sup>5</sup>*Department of Astronomy, The Oskar Klein Center, Stockholm University, AlbaNova, 10691 Stockholm, Sweden*

<sup>6</sup>*Department of Physics, The Oskar Klein Center, Stockholm University, AlbaNova, 10691 Stockholm, Sweden*

<sup>7</sup>*Department of Astronomy, University of Washington, Box 351580, U.W., Seattle, WA 98195, USA*

<sup>8</sup>*Institut für Physik, Humboldt-Universität zu Berlin, Newtonstr. 15, 12489, Berlin, Germany*

<sup>9</sup>*Université Clermont Auvergne, CNRS/IN2P3, Laboratoire de Physique de Clermont, F-63000 Clermont-Ferrand, France.*

<sup>10</sup>*Caltech Optical Observatories, California Institute of Technology, Pasadena, CA 91125, USA*

<sup>11</sup>*Department of Electrical and Computer Engineering, Duke University, Durham, North Carolina 27708, United States*

<sup>12</sup>*DIRAC Institute, Department of Astronomy, University of Washington, 3910 15th Avenue NE, Seattle, WA 98195, USA*

<sup>13</sup>*Astrophysics Science Division, NASA Goddard Space Flight Center, 8800 Greenbelt Road, Greenbelt, MD 20771, USA*

<sup>14</sup>*Department of Astronomy, University of Maryland, College Park, MD 20742, USA*

<sup>15</sup>*The eScience Institute, University of Washington, Seattle, WA 98195, USA\**

<sup>16</sup>*IPAC, California Institute of Technology, 1200 E. California Blvd, Pasadena, CA 91125, USA*

<sup>17</sup>*Kavli Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106, USA*

<sup>18</sup>*Center for Data Driven Discovery, California Institute of Technology, Pasadena, CA 91125, USA*

<sup>19</sup>*Ben-Ziyyo Center for Astrophysics, The Weizmann Institute of Science, Rehovot 76100, Israel*

(Received October 30, 2019; Revised; Accepted)

Submitted to ApJ

### ABSTRACT

The Zwicky Transient Facility (ZTF) is performing a three-day cadence survey of the visible Northern sky ( $\sim 3\pi$  steradian). The transient candidates found in this survey are announced via public alerts. As a supplementary product ZTF is also conducting a large spectroscopic campaign: the ZTF Bright Transient Survey (BTS). The goal of the BTS is to spectroscopically classify all extragalactic transients brighter than 18.5 mag in either the  $g_{\text{ZTF}}$  or  $r_{\text{ZTF}}$ -filters at peak brightness and immediately announce those classifications to the public. Extragalactic discoveries from ZTF are predominantly Supernovae (SNe). The BTS is the largest flux-limited SN survey to date. Here we present a catalog of the 761 BTS SNe that were classified during the first nine months of the survey (2018 Apr. 1 to 2018 Dec. 31). The BTS SN catalog contains redshifts based on SN template matching and spectroscopic host galaxy redshifts when available. Based on this data we perform an analysis of the redshift completeness of local galaxy catalogs, dubbed as the Redshift Completeness Fraction (RCF; the number of SN host galaxies with known spectroscopic redshift prior to SN discovery divided by the total number of SN hosts). In total, we identify the host galaxies of 512 Type Ia supernovae, 227 of which have known spectroscopic redshifts, yielding an RCF estimate of  $44\% \pm 1\%$  (90% confidence interval). We find a steady decrease in the RCF with increasing distance in the local universe. For  $z \lesssim 0.05$ , or  $\sim 200$  Mpc, we

find  $\text{RCF} \approx 0.6$ , which has important ramifications when searching for multimessenger astronomical events. Prospects for dramatically increasing the RCF are limited to new multi-fiber spectroscopic instruments that can catalog  $\gtrsim 10$  million galaxies in the local universe, or wide-field narrowband surveys. We find that existing galaxy redshift catalogs are only 50% complete at  $r \approx 16.9$  mag (AB). Pushing this limit several magnitudes deeper will pay huge dividends when searching for electromagnetic counterparts to gravitational wave events or sources of ultra high energy cosmic rays or neutrinos.

*Keywords:* supernovae: general — galaxies: distances and redshifts — catalogs — surveys

## 1. INTRODUCTION

Fritz Zwicky and Walter Baade first hypothesized that supernovae (SNe) were the transition of normal stars into neutron stars (Baade & Zwicky 1934). To test this hypothesis, Zwicky used the 18-inch Schimdt telescope commissioned on Palomar mountain in 1936, to carry out the first systematic SN survey (Zwicky 1938a,b, 1942). This survey was carried out by visually inspecting photographic plates of nebulae,<sup>1</sup> and identifying new point-sources. Twelve SNe were identified by Zwicky between 1936 Sept. 5 to 1940 Jan. 1.

Since the pioneering efforts by Zwicky, a variety of SN types have been identified through spectroscopy (see e.g., Filippenko 1997). Thermonuclear SNe (SNe Ia) in particular have proven to be invaluable tools in order to measure cosmological distances (e.g., Goobar & Leibundgut 2011), and the study of SNe Ia eventually led to the remarkable discovery of the accelerating expansion of the universe (Riess et al. 1998; Perlmutter et al. 1999). Studies of core-collapse (CC) SNe have led to considerable insights in massive star evolution; extragalactic neutrinos were detected in SN 1987A (Hirata et al. 1987), a  $\gamma$ -ray burst was associated with SN 1998bw (Galama et al. 1998), direct evidence for binary-star driven mass loss was seen in SN 1993J (e.g., Schmidt et al. 1993; Fox et al. 2014).

In order to constrain cosmological models and to characterize both SNe in general and the various SN types and their host galaxies, a large number of SN surveys have been carried out since Zwicky’s time. The scope of these surveys largely traces the progress made in both automation and detector technology during the last few decades. The first systematic search for SNe using a charge-coupled device (CCD) was performed on the 1.5-m telescope at La Silla (Norgaard-Nielsen et al. 1989). The Field-of-View (FoV) of this telescope and CCD was  $2.5' \times 4'$ , and the survey was designed to find a thermonuclear supernova at high-redshift. Two SNe, one SN Ia and one probable SN II, were found in two years. More recent examples of SN surveys that have also been able to systematically classify their supernova candidates using

spectroscopy include for example: the Lick Observatory Supernova Search (LOSS; Li et al. 2000), the Nearby Supernova Factory (SNfactory; Aldering et al. 2002), the Sloan Digital Sky Survey-II (SDSS-II) SN Survey (Frieman et al. 2008), and the Supernova Legacy Survey (SNLS; Astier et al. 2006). In the last few years, based on statistics on the Transient Name Server (TNS<sup>2</sup>), several surveys are discovering hundreds of SNe that are also being spectroscopically classified, including the Palomar Transient Factory (PTF; Law et al. 2009), the Asteroid Terrestrial-impact Last Alert System (ATLAS; Tonry et al. 2018), the All-Sky Automated Survey for SuperNovae (ASAS-SN; Shappee et al. 2014), and the Panoramic Survey Telescope and Rapid Response System (Pan-STARRS1, hereafter PS1; Chambers et al. 2016) Medium Deep Survey.

The past few decades have seen a growing complexity in SN search surveys, with the general trend being an increase in volumetric survey speed (e.g., Bellm 2016) and consequently the number of SN discoveries. Given the scarcity of spectroscopic resources for SN follow-up observations, the increase in SN discoveries has resulted in a smaller fraction of the SNe being classified with time. Of the on-going surveys, only ASAS-SN is able to maintain close to complete spectroscopic coverage ( $95 \pm 3\%$  for  $m_{\text{peak}} < 16.5$ ; Holoien et al. 2019), largely since ASAS-SN only detects very bright SNe. Otherwise, the typical strategies are to either: (i) focus entirely on the most nearby galaxies (LOSS employed this strategy and maintained a nearly complete survey for  $\sim 10$  yr), (ii) focus observations on likely SNe Ia to study cosmology (e.g., SDSS-II, SNLS), or (iii) target only a subset of SN candidates (e.g., PTF, ATLAS). Any of these choices result in major systematic ambiguities underlying any attempt to derive SN rates and demographics, or to use SNe from these surveys as population probes of galaxies. Nevertheless, these compromises have been necessary given the resources at hand.

With the Zwicky Transient Facility (ZTF; Bellm et al. 2019a,b; Graham et al. 2019) in combination with the fully automated Spectral Energy Distribution Machine (SEDM; Ben-Ami et al. 2012; Blagorodnova et al. 2018; Rigault et al.

\* Moore-Sloan, WRF Innovation in Data Science, and DIRAC Fellow

<sup>1</sup> At the time, the term nebulae encompassed any diffuse astronomical object, including galaxies.

<sup>2</sup> <https://wis-tns.weizmann.ac.il/stats-maps>

2019), a low-resolution ( $R \sim 100$ ) Integral-Field-Unit (IFU) spectrograph mounted on the robotic Palomar 60-inch telescope (P60; Cenko et al. 2006), we have set out to address the lack of spectroscopic completeness described above. We aim to monitor the entire visible sky at moderate cadence while being complete in terms of spectroscopic classification. The  $47 \text{ deg}^2$  field of view of the ZTF camera, along with upgrades to the Palomar 48-inch (P48) telescope and dome, achieves a survey speed of  $3750 \text{ deg}^2 \text{ hr}^{-1}$ , to a  $5\sigma$  depth of  $\sim 20.5 \text{ mag}$  in  $r_{\text{ZTF}}$  using 30 s exposures. This allows most of the sky visible from Palomar to be imaged at a 3-day cadence (see §2 for details). Furthermore, SEDM is capable of classifying  $> 10$  SNe in the 18.5 – 19 magnitude range every night. A significant amount of time is also allocated to this project on the Palomar 200-inch telescope (P200), Keck I, the Liverpool Telescope (LT), Apache Point Observatory (APO) and the Nordic Optical Telescope (NOT). These resources are used to supplement our SEDM observations when SEDM classification is not possible, and the combination enables the ZTF Bright Transient Survey (BTS): a SN survey of unprecedented scale and spectroscopic completeness in the local universe.

The primary goal of the BTS is to spectroscopically classify and publicly report every extragalactic  $r < 18.5 \text{ mag}$  transient in the Northern sky covered by the public ZTF surveys,<sup>3</sup> producing the first large, fully magnitude-complete sample to  $r_{\text{peak}} < 18.5 \text{ mag}$ .<sup>4</sup> In this paper we will focus on SNe, but the BTS also finds and classifies Tidal Disruption Events (TDEs), which will be analysed separately, and other extragalactic phenomena such as massive active galactic nucleus (AGN) flares, and fast and highly energetic transients such as AT 2018cow<sup>5</sup> (Prentice et al. 2018; ?; Ho et al. 2019), whose nature remains mysterious. Here we present a catalog of the 761 BTS SNe classified during the first 9 months of the survey (2018 Apr. 1 to 2018 Dec. 31; Table 1). Our catalog contains redshifts based on SN template matching (Blondin & Tonry 2007) and spectroscopic host galaxy redshifts when available. WISE (Wright et al. 2010)  $W1$ -band and PS1 (Chambers et al. 2016)  $i$ -band host galaxy magnitudes are also reported.

We expect that this sample, and its ongoing extension through 2019 and 2020, will be useful for a wide variety of topics within supernova astrophysics, some of which will be the focus of follow-up papers. In this paper, we focus on an analysis of the redshift completeness of local galaxy catalogs (§5), dubbed as the Redshift Completeness Fraction (RCF; the number of SN host galaxies with known redshift prior to SN discovery divided by the total number of SNe). The

methodology for this analysis closely follows that of Kulkarni et al. (2018).

## 2. SURVEY DESIGN

Transient candidates for the BTS are provided by the public ZTF surveys: the Northern Sky Survey (NSS) and the Galactic Plane Survey (GPS). These surveys are made possible by an award from the NSF Mid-Scale Innovations Program (MSIP), and we henceforth refer to them as the ZTF MSIP surveys (see Bellm et al. 2019b for details). The NSS covers  $\approx 13,000 \text{ deg}^2$  of the Northern sky at a 3-day cadence in the  $g_{\text{ZTF}}$  and  $r_{\text{ZTF}}$  filters, while using 34% of the P48 telescope time. The GPS covers  $\approx 1,500 \text{ deg}^2$  of the galactic plane at a 1-day cadence, also in the  $g_{\text{ZTF}}$  and  $r_{\text{ZTF}}$  filters, and uses 6% of the P48 telescope time.

The BTS avoids low Galactic latitudes by design; we reject all transient candidates found within  $7^\circ$  of the Galactic plane (see §2.1). The combination of significant Galactic extinction and the BTS Galactic plane cut means that the vast majority of the transients in the BTS originate from the NSS. However, due to the large FoV and the fixed main field-grid used by ZTF (Masci et al. 2019), some of the GPS fields still allow transients to be found and monitored after a  $7^\circ$  galactic plane cut. During 2018 we classified two SNe within GPS fields.

The images from the ZTF MSIP surveys are processed and analysed at IPAC by an automated pipeline (Masci et al. 2019), that uses the Zackay et al. (2016) difference-imaging method. The pipeline produces transient alert packets from the difference images in the Apache Avro<sup>TM</sup> format.<sup>6</sup> The Avro alert packets are distributed through the University of Washington (UW) as a Kafka data stream.<sup>7</sup> The alert stream originating from the ZTF MSIP surveys is the data source for BTS transient candidates. The full ZTF Alert Distribution System (ZADS) is described in detail in Patterson et al. (2019).

Figure 1 shows the  $g_{\text{ZTF}}$  and  $r_{\text{ZTF}}$ -band coverage maps of the ZTF MSIP surveys, between 2018 April 1 and 2018 December 31. The distribution of re-visit times (cadence) for the NSS for each field during the same time period, excluding 2018 September 29 to 2018 October 31 when the P48 was undergoing maintenance, is shown in Figure 2. Approximately 70% of the NSS observations were carried out at the planned 3 d cadence, and  $\sim 90\%$  of re-visits occurred within  $\leq 6 \text{ d}$  during 2018. For the GPS, around 80% of the observations were carried out at a 1-day cadence. However, since only two BTS SNe were found and classified during 2018 in the GPS fields, the GPS cadence is not representative for the BTS.


<sup>3</sup> Excluding the galactic plane ( $\pm 7^\circ$ ).

<sup>4</sup> The saturation limit of ZTF is  $\sim 14 \text{ mag}$ .

<sup>5</sup> The internal ZTF designation for AT 2018cow is ZTF18abcfcoo.

<sup>6</sup> <https://avro.apache.org>

<sup>7</sup> <https://kafka.apache.org>



**Figure 1.** Coverage maps for the ZTF MSIP surveys, in the  $g_{ZTF}$ - (left panel) and  $r_{ZTF}$ -band (right panel) between 2018 Apr. 1 and 2018 Dec. 31. The colored rectangles represent the fixed ZTF main field grid. The color intensity indicates the number of observations during this time period, truncated to a maximum of 65.

In order to identify bright SN candidates within the raw ZTF alert stream produced by the MSIP surveys we utilize the filtering capability within the GROWTH<sub>Marshal</sub> framework (Kasliwal et al. 2019) to apply the candidate filter described below (§2.1). The GROWTH<sub>Marshal</sub> is also used to organize BTS sources and the corresponding spectroscopic follow-up efforts (§§2.2, 2.3).

### 2.1. Supernova Candidate Filter

The BTS filter used between 2018 Apr. 1st and 2018 Dec. 31st was deliberately designed to be simple, in order to minimize the risk of false negatives (i.e., real transients that are not saved as part of the program). Using the GROWTH<sub>Marshal</sub> alert filtering system, we applied the following cuts to the raw ZTF alert stream in order to identify bright SN candidates:

- Alerts at low Galactic latitudes ( $|b| \leq 7^\circ$ ) are rejected.
- Alerts with a random forest based machine learning real-bogus score ( $rb_{score}$ ; Mahabal et al. 2019) of less than 0.2 are rejected. This choice results in a completeness (i.e., 1 - false negative rate) of  $> 99\%$  (figure 16 in Duev et al. 2019).
- Alerts produced at the position of known stars, as identified in the catalog created by Tachibana & Miller (2018), are rejected. A small fraction of galaxies (estimated to be  $< 0.5\%$  in Tachibana & Miller 2018), and thus nuclear SNe will be missed as a result of this cut.
- Alerts produced close to very bright stars have been rejected ( $< 20''$  for  $< 15$  mag stars;  $\sim 1\%$  loss of survey area<sup>8</sup>).

<sup>8</sup> Based on randomly injecting 10,000 SNe within the survey footprint and matching against PS1 stars.

- Alerts that do not include at least two detections separated by a minimum of 30 min are rejected (moving object filter).
- Alerts with negative flux relative to the reference image are rejected.
- The alert must include at least one epoch with  $m > 19$  mag (in  $g_{ZTF}$ - or  $r_{ZTF}$ -band), otherwise it is rejected.

The BTS filter effectively passes all alerts brighter than 19 mag that are not consistent with stellar events (star detected in PS1) or moving objects for human vetting (scanning; §2.2).

### 2.2. Human candidate vetting

On a typical night in 2018 a few hundred alerts passed the BTS filter. These BTS transient candidates were visually inspected by a team of scanners on a daily basis. To organize this effort we use the GROWTH<sub>Marshal</sub> (Kasliwal et al. 2019), where the light curves and image cutouts (science, reference, and subtraction) contained in the Avro packets of the passing alerts are collected on a scanning page for each night. Supplementary information is also displayed, such as: PS1 and SDSS color composite cutouts centered on the position of the transient, the star-galaxy separation score ( $sg_{score}$ ), which gives a probability that the closest PS1 counterpart is an extended galaxy or a point-like star (Tachibana & Miller 2018), multi-band photometry of this PS1 counterpart, a cross-check for known near earth objects (NEOs), and information about if and when there have been previous ZTF alerts at the same position that are not part of the 30-day history contained in the alert packet itself. External catalog cross-matches (e.g., NED, TNS, SIMBAD, VizieR) are also linked through the GROWTH<sub>Marshal</sub> to provide additional contextual information for each potential SN candidate.

The human vetting process essentially consisted of inspecting the information contained in the Avro packets for each alert that passed our filter and also taking into account any relevant supplementary information available in order to rule out variability from a stellar counterpart, and to reject alerts produced by known AGN.

Among the passing alerts, 5–15 SN candidates are typically identified by the human scanners as good SN candidates and assigned for spectroscopic followup, per night.<sup>9</sup> The two main contaminants in our scanning process are cataclysmic variables (CVs) that are too faint to be seen in PS1 in their quiescent phase, and therefore lack an `sgscore`, and AGN. Both of these must be avoided, given our limited spectroscopic resources. We have found that the vast majority of CVs can be avoided by monitoring the lightcurve behavior until  $\sim 1$  week after the initial outburst and comparing the evolution with typical CV lightcurves.<sup>10</sup> This is especially effective since ZTF produces both  $g_{ZTF}$ - and  $r_{ZTF}$ -band photometry for virtually all transients that are detected.


Filtering AGN is more challenging: excluding all AGN from the BTS could inadvertently reject a SN that has exploded near the nucleus of a galaxy that harbors an AGN. Filtering on past variability is a very effective way of excluding AGN, but for the sample presented here the baseline of ZTF observations was only a few weeks or months, which is not always a sufficient amount of time for the AGN to change in flux enough to be recognized as a variable object. We have generally not triggered spectroscopic follow-up for alerts that are positionally coincident with known AGN (e.g., the ALLWISE mid-infrared AGN catalog; [Secrest et al. 2015](#), the Million Quasar catalog; [Flesch 2015](#)), unless the photometric evolution of the transient is very similar to that of a SN. Thus, the BTS is incomplete for SNe near AGN (a focused survey with the specific goal of discovering SNe in galaxies with AGN is needed to address this).

### 2.2.1. Completeness of the BTS filter

To assess the completeness of our BTS GROWTH Marshal filter (§2.1) and human scanning effort (§2.2) we have re-processed and re-filtered all public ZTF alerts between 2018 Apr. 1 to 2018 Dec. 31 using the AMPEL system ([Nordin et al. 2019](#); [Soumagnac & Ofek 2018](#)). Two filters were applied for this exercise: a variant of our BTS GROWTH Marshal filter converted to work within

<sup>9</sup> This number strongly depends on the weather; after a period of bad weather a large number of SNe will be recovered when observations are resumed. Good weather periods produce a more constant number each night.

<sup>10</sup> CVs feature a fast evolution (rise time of  $\sim 1$ –2 d, decline of  $\sim 7$ –10 d) and their  $g_{ZTF} - r_{ZTF}$  colors are persistently blue. We do not follow-up events with these characteristics that also lack an obvious host-galaxy counterpart.



**Figure 2.** Cadence distribution for the ZTF NSS, in the  $g_{ZTF}$ - (blue bars) and  $r_{ZTF}$ -band (red bars), truncated at six days. Cumulative distributions are shown as a blue solid line for the  $g$  band and a red dashed line for the  $r$  band.  $N/N_{\text{tot}}$  is the fraction of observations at a specific cadence compared to the total number of observations between 2018 Apr. 1 and 2018 Dec. 31.

AMPEL<sup>11</sup>, and the AMPEL TNS channel filter described in [Nordin et al. \(2019\)](#).

After the filtering step, we combined the passing candidates from both filters and imposed a cut requiring: at least 5 detections, a peak mag  $< 18.5$ , no more than one negative detection, and a time between the first and last detection more than 5 days and less than 90 days<sup>12</sup>. Finally, we also required that the candidate passes a version of the GROWTH Marshal filter that checks all associated Avro packets for that candidate. All of the remaining objects that were not saved on the GROWTH Marshal were then vetted individually to remove any remaining CVs, AGNs, classical novae, and artifacts.

This singled out 17 likely SNe that our BTS filter scanning efforts had not picked up. Among these 9 were saved by other ZTF science programs or AMPEL, and were spectroscopically classified. The remaining 8 are unclassified. The BTS sample contains 520 SNe peaking at  $< 18.5 \pm 0.05$  mag. This implies a completeness in our scanning of the BTS filter in 2018 of 97% for candidates with peak mag  $\lesssim 18.5$ . The 17 objects identified here are not included in our sam-

<sup>11</sup> We have confirmed that this filter passes all candidates we found and classified in 2018

<sup>12</sup> A subset of the  $>90$  day events were also vetted and no SNe were found.

ple or analysis but will be included and analysed in our next data-release. We also note that [Nordin et al. \(2019\)](#) showed that ZTF has been complete with respect to detecting SNe reported to TNS by other groups when they fall on active ZTF CCD regions.

### 2.3. Spectroscopic followup assignment

The primary classification instrument of BTS candidates is the SEDM ([Blagorodnova et al. 2018](#); [Rigault et al. 2019](#)). For candidates that pass the BTS filter (§2.1) and visual inspection (§2.2), we assign spectroscopic SEDM observations with the following priorities: transients that are, or are likely to become, brighter than 18.5 mag are scheduled with the highest priority (P3). Transients that appear likely to peak between 18.5 and 18.75 mag are assigned a lower priority (P2), and transients expected to peak fainter than 18.75 mag are triggered at the lowest priority (P1). For a typical BTS source ( $m \approx 18.5$  mag) observed under typical observing conditions for 1800 s, SEDM obtains  $\text{SNR} \approx 12$  per resolution element, and an integrated  $\text{SNR} \approx 50$  in the region covered by the  $r_{\text{ZTF}}$  filter.

The SEDM queue is designed so that lower priority targets (e.g., P1, P2) are only observed if no higher priority targets are present in the queue that could be observed within the same observation-time window. The purpose of this priority scheme is two-fold. First, it ensures that we can reach a high level of completeness for  $m_{\text{peak}} < 18.5$  mag sources, by allowing some margin for error on the fainter end of 18.5 mag. Second, when the queue allows (e.g., periods of consistently good weather), significantly fainter targets, including as faint as  $\sim 19$  mag, can also be observed in addition to our high priority (P3) targets.

By default, triggers that enter the SEDM queue remain active for 7 d. If the transient has not been observed at this time, the candidate is re-assigned to SEDM, or to larger telescopes if the candidate has become too faint for SEDM ( $m \gtrsim 19$  mag). Larger telescopes are also used if classification with SEDM is unsuccessful, which typically happens only if the candidate was observed in poor sky conditions, there is strong host galaxy contamination, or if higher resolution is needed for a secure classification. For this purpose we have primarily used P200 and Keck I, but supporting observing programs at LT, NOT and APO have contributed significantly as well. Community efforts (e.g., ePESSTO; [Smartt et al. 2015](#)) have also contributed through TNS (see §3.1 for details).

#### 2.3.1. Spectroscopic completeness

A key goal for the BTS is to obtain high spectroscopic completeness for all events passing our basic selection criteria (essentially,  $m < 18.5$  mag and extragalactic, see §2.1). While we made every effort to spectroscopically classify every transient saved to the BTS program, our efforts were in-

evitably imperfect. Following the conclusion of 2018 (i.e., the period covered in this early-release paper), we conducted two independent tests of our completeness as described below.

As an initial test of the completeness of BTS spectroscopic follow-up, we compiled every object that was saved following visual inspection (§2.2) and applied additional filters designed to remove most variables and AGN using a more sophisticated point-source coincidence check and the long-term light-curve history.<sup>13</sup> We restrict these candidates to those brighter than  $m < 18.5$  in at least one observation. Every such object without a formal spectroscopic classification (74 in total) was then visually examined by our team of scanners. Most of these events are clearly not transients (e.g., subtraction artifacts, AGN, stars) based on their full light curves. However, 31 events had properties consistent with SNe and were unclassified. If each of these events is a genuine SN, this would suggest a completeness of  $\sim 96\%$ .

We separately examined the filtered subset of alerts described above and estimated the characteristic rise and fade times (from, and to, 0.75 mag below peak, respectively) for every event using an automated procedure. For events with sufficient P48 data around peak to accurately constrain the rise and fade times, and with rise times between 4–100 d and fade times between 10–200 d (i.e., probable SNe), our classification completeness rate is 100% to  $m_{\text{peak}} < 16.5$  mag, 98.8% to  $m_{\text{peak}} < 17.5$  mag, 93.6% to  $m_{\text{peak}} < 18.5$  mag, and 88.8% to  $m_{\text{peak}} < 19.0$  mag. This method, which is easily automated (and could be applied to samples larger than what can be manually inspected), is consistent with the results from our visual inspection described above.

## 3. BTS SN CLASSIFICATIONS

### 3.1. Classification Method

As previously noted, spectroscopic observations of BTS SN candidates are primarily obtained with SEDM. In cases where the candidates were too faint to be observed with SEDM, or scheduling conflicts prevented SEDM observations, or the SEDM spectra proved to be ambiguous, spectra were obtained with spectrographs on larger aperture telescopes: the Double Spectrograph (DBSP; [Oke & Gunn 1982](#)) on P200, the Low Resolution Imaging Spectrometer (LRIS; [Oke et al. 1995](#)) on the Keck 1 telescope, the Dual Imaging Spectrograph (DIS) on the APO 3.5 m telescope, the Spectrograph for the Rapid Acquisition of Transients (SPRAT; [Piascik et al. 2014](#)) on the 2.0 m LT, and the Alhambra Faint Object Spectrograph and Camera (ALFOSC) on the 2.56 m NOT.

<sup>13</sup> All but 3 of the early-release SNe passed these additional filters. These 3 were rejected due to their proximity to bright stars, meaning the additional filters still find genuine SNe with high fidelity.

SEDM data were reduced by the pipeline described in Blagorodnova et al. (2018), for data until August 2018, and the automatic PYSED pipeline described in Rigault et al. (2019) for data after August 2018. DIS data were reduced with the pyDIS package (Davenport et al. 2016). DBSP data were reduced using the PyRAF (Science Software Branch at STScI 2012) based pipeline pyraf-dbsp (Bellm & Sesar 2016). ALFOSC data were reduced using standard procedures and tools based on IRAF (Tody 1986). LRIS data were reduced using the LRIS automated reduction pipeline (LPipe; Perley 2019).

For SN candidates where we could not obtain spectroscopic observations from any of the above mentioned telescopes, public TNS spectra were analysed from the following instruments: the Asiago Faint Object Spectrograph and Camera (AFOSC) on the Asiago Ekar 182 cm telescope, FLOYDS on the Faulkes Telescope South (FTS) in Australia operated by Las Cumbres Observatory (LCO), the Wide Field Reimaging CCD (WFCCD) on LCO’s duPont telescope, the ESO Faint Object Spectrograph and Camera v.2 (EFOSC2) on the New Technology Telescope (NTT), the Dolores (Device Optimized for the Low REsolution) on the Telescopio Nazionale Galileo (TNG), the Intermediate-dispersion Spectrograph and Imaging System (ISIS) on the William Herschel Telescope (WHT), and the DeVeney spectrograph on the Discovery Channel Telescope (DCT).

Preliminary classifications are made via SuperNova IDentification (SNID; Blondin & Tonry 2007) template matching *and* visual inspection. SNID is automatically applied to all SEDM spectra, whereas for all other instruments SNID is applied to the spectra by the observer. These preliminary classifications are annotated and recorded within the BTS program on the GROWTH *Marshal*, and subsequently sent to TNS within one to two days.<sup>14</sup>

We revisit each of the preliminary classifications for this study in order to develop a homogeneous classification scheme. For this purpose we developed a custom process to spectroscopically classify the 761 SNe in the BTS sample. For each BTS SN, we identified the top 15 spectral matches ( $r_{\text{lap}_{\text{min}}} \geq 5$ ) from SNID to the most recent spectrum available on the GROWTH *Marshal*. We used the latest spectrum from the *Marshal* under the assumption that BTS targets only received additional spectroscopic observations when the initial classification was inconclusive. The SNID templates used for this process include the developer defaults,<sup>15</sup> as well as SNe Ia and a few non SN templates

from the Berkeley SN Ia program (BSNIP; Silverman et al. 2012a), SN Ib/c templates from Modjaz et al. (2014, 2016); Liu et al. (2016), and Williamson et al. (2019), and SN IIP templates from Gutiérrez et al. (2017).

Following the SNID matching, we produced plots showing a comparison between the observed BTS spectrum and the (redshift-corrected) template spectrum from SNID. These plots were visually inspected to identify the best matching template. In practice, the sample was split into 6 groups, and each group was inspected by a member of our team (CF, AAM, AD, YS, KT, AG). While identifying the best-matching template, we recorded the SN type and redshift, as well as the name and phase of the template SN spectrum. In cases where the same classification was reported for all 15 matches, we recorded the type, redshift, and phase from the top match from the SNID output. We otherwise selected the best match based on common prominent SN spectral features (H, He, Si, Ca, Fe, etc.). If the top 15 matches from SNID proved ambiguous, we used either the ZTF light curve or alternative spectra to remove the ambiguity. For example, in cases with multiple matches to both SNe Ia and SNe Ic, the telltale secondary near-infrared (nIR) peak of SNe Ia can typically be seen in ZTF  $r_{\text{ZTF}}$ -band light curves. The secondary nIR peak is unique to SNe Ia, as explained in Kasen (2006). It occurs following a recombination transition of the iron-group elements in the ejecta, whereby the strength of the Fe III and Co III lines decreases and there is a corresponding strengthening of the Fe II and Co II lines (see also Blondin et al. 2015). While these papers have mostly considered  $\lambda \gtrsim 7500 \text{ \AA}$ , a related “shoulder” in the  $r$ -band lightcurve is also observed at approximately the same time (Papadogiannakis et al. 2019a,b).

If at this stage a classification still proved ambiguous, then the SN was examined by another member of the team. For consistency, a final check of all ambiguous classifications was performed by two members of the team (CF, AAM). Ultimately, we have classified 761 SNe via their spectra and light curves (Table 1). Out of these, 503 were classified using SEDM spectra, 86 using P200-DBSP, 76 using Keck 1-LRIS, 20 using LT-SPRAT, 11 using APO-DIS, and 9 using NOT-ALFOSC. Finally, 56 were classified based on publicly available spectra on TNS.

We note that the positions reported in Table 1 are obtained by taking the weighted average of the position of the SN in every image in which the SN is detected (i.e., for every alert associated with the SN). The updated positions are more accurate than those reported to TNS, which typically only include a single low signal-to-noise ratio (SNR) detection of the SN.

<sup>14</sup> The BTS classifications on TNS should also be considered preliminary. They reflect the initial classifications on the GROWTH *Marshal*, since turnaround speed is important, but they are subsumed by the efforts described in this paper.

<sup>15</sup> <https://people.lam.fr/blondin.stephane/software/snid/#Download>

### 3.2. Classifications

**Table 1.** ZTF BTS SNe

ZTF	$\alpha_{\text{SN}}$	$\delta_{\text{SN}}$	IAU	TNS Internal	Discovered	$E(B-V)^b$	SN	$z_{\text{SN}}^c$	Observed time and mag at maximum brightness <sup>a</sup>					
									$t_g$	$m_g$	$\sigma_{m_g}$	$t_r$	$m_r$	$\sigma_{m_r}$
Name	(J2000.0)	(J2000.0)	Name	Name	by	(mag)	Type		(JD') <sup>d</sup>	(mag)	(mag)	(JD') <sup>d</sup>	(mag)	(mag)
ZTF18aabssth	11:00:45.38	+22:17:15.0	SN 2018aex	ZTF18aabssth	ZTF	0.015	II	0.026	494.05	20.44	0.19	218.71	18.74	0.05
ZTF18aabxslsv	10:29:51.62	+09:00:46.6	SN 2018aks	ASASSN-18ga	ASAS-SN	0.025	Ib	0.055	605.71	20.33	0.19	224.68	18.64	0.04
ZTF18aaemivw	10:33:42.69	+39:29:26.6	SN 2018hus	ZTF18aaemivw	ZTF	0.012	Ia	0.065	423.96	19.00	0.16	534.87	17.95	0.15
ZTF18aagpzjk	07:59:25.01	+16:25:34.5	SN 2018afm	...	POSS	0.031	II	0.013	...	...	...	217.66	17.46	0.02
ZTF18aagrdfs	14:33:19.98	+41:16:02.3	SN 2018alc	ASASSN-18ge	ASAS-SN	0.012	Ib	0.024	547.91	17.37	0.05	217.90	16.67	0.01
ZTF18aagrtps	13:14:25.46	+50:58:39.7	SN 2018amo	ASASSN-18gi	ASAS-SN	0.010	Ia	0.018	214.76	16.52	0.02	214.73	16.46	0.02
ZTF18aagstdc	15:50:03.56	+42:05:18.5	SN 2018apn	ASASSN-18gs	ASAS-SN	0.018	Ia	0.038	210.86	17.24	0.02	539.01	17.27	0.06
ZTF18aagtcxj	16:32:11.55	+42:42:48.3	SN 2018aqm	...	TNTS	0.011	Ia	0.033	539.03	19.01	0.19	219.96	18.09	0.03
ZTF18aahesrp	08:35:45.43	+28:16:12.9	SN 2018aqy	ATLAS18mzs	ATLAS	0.036	Ia	0.051	...	...	...	217.69	18.54	0.07
ZTF18aaahfeiy	10:17:15.57	+43:31:24.2	SN 2018loy	ZTF18aaahfeiy	ZTF	0.010	Ia	0.071	429.99	20.53	0.21	214.71	18.15	0.05

NOTE— This table is available in its entirety in a machine-readable form in the online journal. A portion is shown here for guidance regarding its form and content.

References for SNe that were recovered by ZTF and discovered elsewhere (“Discovered by” in the table) are as follows: ASAS-SN (Shappee et al. 2014), ATLAS (Tonry et al. 2018), the Corona Borealis Observatory Supernova Survey (CSNS; Sun et al. 2018), G. Cortini (Cortini 2018), DLT40 (Tartaglia et al. 2018), *Gaia* (Hodgkin et al. 2013), K. Itagaki (Itagaki 2018a,b), the Italian Supernovae Search Project (ISSP; <http://italiansupernovae.org/>), LOSS (Filippenko et al. 2001), the Mobile Astronomical System of Telescope Robots (MASTER; Gorbvskoy et al. 2013), the Puckett Observatory Supernova Search (POSS; <http://www.possupernova.com/>), PS1 (Chambers et al. 2016), J. Grzegorzec (Grzegorzec 2018), the PMO-Tsinghua Supernova Survey (PTSS, <http://www.cneost.org/ptss/>), the Great Supernova Hunt (SNhunt; <http://nessi.cacr.caltech.edu/SNhunt/>), the Tsinghua University-National Astronomical Observatories, Chinese Academy of Sciences Transient Survey (TNTS; Zhang et al. 2015), the Xinming Observatory Supernova Survey (XOSS; (Zhang et al. 2018)), Y. Tanaka (Tanaka 2018)

<sup>a</sup> Time and magnitude of maximum brightness are determined directly from the observations, as available in the AVRO alert packets. “...” is used when the SN was not detected in either the  $g_{\text{ZTF}}$  or  $r_{\text{ZTF}}$  filter. No correction for extinction has been applied.

<sup>b</sup>  $E(B-V)$  is determined using the Schlafly & Finkbeiner (2011) updates to the Schlegel et al. (1998) maps.

<sup>c</sup> Determined from SNID (see text).

<sup>d</sup>  $\text{JD}' = \text{JD} - 2,458,000$ .

We broadly classify all BTS SNe as belonging to one of 4 different classes: SNe Ia, SNe II, SNe Ib/c, and super luminous SNe (SLSNe). As detailed above, these classifications are primarily made via the SN spectra, however, in some cases the photometric evolution of the SN also informs the classification. This is especially true of the SLSNe, which are defined by their luminosity (typically  $M < -21$  mag, Gal-Yam 2012; although ZTF adopts  $M < -20$  mag).

Of the 761 SNe, we find that 547 are SNe Ia, 155 are SNe II, 40 are SNe Ib/c, and 19 are SLSNe. The fraction of discoveries belonging to each of these classes is in agreement with the results from Li et al. (2011) for a magnitude-limited survey, as shown in Figure 3.<sup>16</sup> Figure 3 also shows the relative rate of SNe found by ASAS-SN (Holoien et al. 2017a,b,c, 2019), which, like the ZTF BTS and unlike LOSS, does not target specific galaxies when searching for transients.<sup>17</sup> Both ZTF and ASAS-SN find a higher fraction

of SNe II than LOSS, although these estimates all agree to within the uncertainties. By targeting massive galaxies, including a significant fraction of passive elliptical galaxies, LOSS may have been slightly biased against finding CC SNe (see Taubenberger 2017 and references therein). The relative rate of SLSNe is somewhat higher for ZTF compared to ASAS-SN, but still consistent within the uncertainties (see §3.3.4).

Table 2 summarizes the relative fraction of SNe in each class for the ZTF BTS, ASAS-SN, and LOSS. For LOSS we directly use the estimates from Li et al. (2011), while for the BTS and ASAS-SN we assume the observations are drawn from a multinomial distribution and estimate 95% confidence intervals on the true rate via the approximate method of Goodman (1965) as implemented in the `MultinomCI` (Signorell & et al. 2019) package in R. The true uncertainties on these fractions require a detailed estimate of the completeness of the BTS, which is beyond the scope of this paper, and will be addressed in future work (Nordin et al., in prep.).

<sup>16</sup> The ZTF BTS utilizes a 3-d cadence (see §2), whereas Table 7 in Li et al. (2011) reports results for surveys with a 1-d and 5-d cadence. The results in Li et al. (2011) are identical for 1-d and 5-d cadences, and thus we assume an intermediate 3-d cadence would also yield identical relative fractions of SNe.

<sup>17</sup> To calculate the relative rates of SNe found by ASAS-SN (and host-galaxy offsets, which are discussed below) we include both ASAS-SN dis-


coveries, and SNe recovered by ASAS-SN, as all discovered and recovered SNe are included in our analysis of the ZTF BTS. Therefore, the numbers shown here differ slightly from what is shown in e.g., Figure 1 of Holoien et al. (2019), which only considers SNe discovered by ASAS-SN.



**Table 2.** Relative SN rates

Survey	$\mathcal{R}(\text{Ia})$	$N_{\text{Ia}}$	$\mathcal{R}(\text{II})$	$N_{\text{II}}$	$\mathcal{R}(\text{Ibc})$	$N_{\text{Ibc}}$	$\mathcal{R}(\text{SLSN})$	$N_{\text{SLSN}}$
LOSS	$0.792 \pm_{0.055}^{0.044}$	...	$0.166 \pm_{0.039}^{0.050}$	...	$0.041 \pm_{0.013}^{0.016}$	...	...	...
ASAS-SN	$0.742 \pm_{0.045}^{0.040}$	607	$0.211 \pm_{0.037}^{0.043}$	173	$0.043 \pm_{0.016}^{0.024}$	35	$0.004 \pm_{0.003}^{0.012}$	3
ZTF BTS	$0.719 \pm_{0.048}^{0.043}$	547	$0.204 \pm_{0.038}^{0.043}$	155	$0.053 \pm_{0.018}^{0.027}$	40	$0.025 \pm_{0.012}^{0.021}$	19

NOTE— Relative rates  $\mathcal{R}$  of SNe Ia, II, Ibc, and SLSN from a flux limited search. Results for LOSS are taken directly from Table 7 in Li et al. (2011) for a 1-d cadence and are based on an assumed luminosity function and Monte Carlo simulations. Thus, the total number of each type of SN is not relevant and therefore not reported. Results for ASAS-SN use all discovered and recovered SNe reported in Holoien et al. (2017a,b,c, 2019). Uncertainties for both ASAS-SN and the ZTF BTS include 90% confidence intervals (see text).



**Figure 3.** Relative fractions of SNe Ia, II, Ib/c, and SLSNe in the flux-limited ZTF BTS survey. The orange, hatched bars show the results for the BTS, while the light grey, open bars show results from LOSS (Li et al. 2011), and the blue double-hatched bars show the results from ASAS-SN (Holoien et al. 2019). The lightly shaded regions show the uncertainty on each estimate (see text for further details). To within the uncertainties, the results agree for all 3 surveys (Li et al. 2011 did not report results for the SLSN class – see text).

As a test of our classification accuracy, we compare our final classifications to those made by ePESSTO (Smartt et al. 2015).<sup>18</sup> There are 26 sources in common between our BTS classifications and those made by ePESSTO, and the classifications are in agreement for all but 2 sources: ZTF18abmrhom (SN 2018ffi) and ZTF18abtswj (SN 2018gfx). We classify ZTF18abmrhom as a SN Ia, whereas the ePESSTO spectrum is classified as having just galaxy light. The ZTF spectrum of ZTF18abmrhom,

<sup>18</sup> We adopt ePESSTO for comparison because (i) it is the survey with the most overlap with the ZTF BTS, and (ii) all ePESSTO classifications are made with the 3.6 m NTT, which, on average, will perform better than SEDM for SN classification.

which was obtained 2 nights prior to the ePESSTO spectrum, shows clear and strong Si II absorption. The same broad feature can be seen in the ePESSTO spectrum, which is otherwise dominated by emission from the host galaxy. Furthermore, the  $r_{\text{ZTF}}$ -band light curve shows a “shoulder” a few weeks after maximum light. Taken together, it is very likely that ZTF18abmrhom is a SN Ia. The classification of ZTF18abtswj is more challenging: the spectrum clearly shows narrow emission lines, and we have classified the event as a SN II, whereas ePESSTO classified the spectrum as an active galactic nucleus (AGN). The photometric evolution of ZTF18abtswj is far more reminiscent of SNe than AGN: the transient exhibits a  $\sim 30$  d rise, followed by a very slow, monotonic decline (the transient has exhibited a monotonic fade by  $\sim 1.5$  mag over the period covering 2018 and 2019) as is characteristic of many SNe II (e.g., Turatto et al. 1993). Furthermore, the WISE colors for the host galaxy are not consistent with AGN (Jarrett et al. 2011). Thus, based on our comparison to ePESSTO, we conclude that our classifications are of high fidelity.

There were 3 BTS sources for which we attempted spectroscopic classification, but the nature of the transients remains unresolved. Each of the 3 candidates are positionally coincident with the nucleus of their host galaxies. ZTF18aaqkdwf (SN 2018fhd) exhibits a broad rise and decline over  $\sim 200$  d and significant P-Cygni-like feature around  $H\alpha$ . However, variability at the location of ZTF18aaqkdwf was detected  $\sim 3.5$  yr prior to the 2018 variability by iPTF, and the host galaxy is a bright point source in the radio (Helfand et al. 2015). Thus the variability may be due to an AGN. The light curve of ZTF18abuqhje (SN 2018gki) has poor coverage with only 4  $r_{\text{ZTF}}$  detections that show a decline of  $\sim 0.7$  mag over  $\sim 15$  d. We obtained 2 spectra of ZTF18abuqhje, both of which show many narrow lines that could be consistent either with an AGN or a SN II. However, we detected variability from this nucleus  $\sim 2.5$  yr prior to 2018 with iPTF, which would be consistent with an AGN. The general evolution of the ZTF18aarchg (AT 2018boa)  $r_{\text{ZTF}}$  light curve is consis-

tent with a SN: the transient rises by  $\sim 1.5$  mag over  $\sim 20$  d, followed by a monotonic decline. Spectra of ZTF18aarcchg exhibit a blue continuum with superposed narrow emission from the Balmer series, [O III], and [S II], all of which are consistent with star formation (the host is classified as a star-forming galaxy; Maraston et al. 2013). Without discernible SNe features in the spectrum, we cannot classify this transient. We exclude these 3 candidates from the subsequent analysis.

### 3.3. Sub-type Classifications

The primary purpose of this study is to improve the measurement of the RCF, and to extend the redshift coverage of this measurement beyond what is presented in Kulkarni et al. (2018). To that end, we only need to separate SNe Ia from all other transients in the ZTF BTS. Nevertheless, we attempt to make sub-type classifications, based on the SNID matches described above, as detailed for each class below. We caution, however, that the vast majority of these classifications are made with very low resolution ( $R \approx 100$ ) SEDM spectra, and as a result there are significant uncertainties on the subclass of any individual SN. Furthermore, any biases in the SNID template set can influence the final BTS classifications. The SNID templates were compiled from heterogeneous sources and do not perfectly reflect the discovery-space of an un-targeted transient survey, meaning the SNID templates themselves are a further source of uncertainty for the final sub-classifications presented in Table 1 (see Blondin & Tonry 2007 for further details).

#### 3.3.1. SNe Ia

The vast majority of SNe Ia discovered in magnitude-limited surveys are considered “normal.” It is argued in Li et al. (2011) that the most common subclass is SN 1991T-like (SN Ia-91T in Table 1), which are slightly over-luminous relative to normal SNe Ia, followed by SN 1991bg-like (SN Ia-91bg in Table 1), which are under-luminous and decline faster than normal SNe Ia. These rates are derived from LOSS, which targeted relatively massive, high star-formation rate galaxies. SNe Ia-91T seem to prefer late-type galaxies and may be associated with young stellar populations (e.g., Howell 2001). Thus, the relative rate of SNe Ia-91T may have been over-estimated (see e.g., Taubenberger 2017; Silverman et al. 2012b).

In the ZTF BTS we identify 504 normal SNe Ia, 31 91T-like SNe, and 6 91bg-like SNe. This represents significantly less 91T-like ( $\sim 6\%$ ) and 91bg-like ( $\sim 1\%$ ) SNe than one would expect based on the LOSS results for a magnitude limited survey ( $\sim 18\%$  91T-like and  $\sim 3\%$  91bg-like SNe; Li et al. 2011), however, it does agree with what is found by ASAS-SN ( $\sim 6\%$  91T-like and  $\sim 1\%$  91bg-like SNe; Holoiien et al. 2017a,b,c, 2019).

We caution that sub-type classification can be difficult with a single low-resolution SEDM spectrum, as is the case for the majority of the SNe in our sample. For example, the most distinguishing feature of 91T-like SNe is weak Si II and Ca II absorption prior to maximum light (e.g., Filippenko 1997; Branch et al. 2006). At  $R \approx 100$  even relatively strong absorption can be smeared out, and as a result SNID frequently identifies both normal and 91T-like SNe Ia as the best matches for the SNe Ia in our sample. Furthermore, it is difficult to separate normal and 91T-like SNe in post-maximum spectra. Thus, we conservatively label SNe Ia as normal, unless there is strong spectroscopic (very weak Si II and Ca II) or photometric (over-luminous *and* a slow decline) evidence to support a 91T-like classification. Similarly for 91bg-like SNe, unless there is strong spectroscopic (weak Fe II, strong Ti II; Filippenko 1997) and photometric (under-luminous *and* a fast decline) evidence, we label the SN as normal.

In addition to the above subclasses, we additionally identify 3 SN 2002cx-like SNe, 1 SN Ia that shows signs of CSM interaction (SN Ia-CSM), and 2 SNe that appear to be super-Chandrasekhar mass explosions.

#### 3.3.2. SNe II

We have identified 162 H-rich SNe in the ZTF BTS. Of these, we make no effort to distinguish between Type IIP and IIL SNe, which are photometrically defined subtypes. We classify SNe II as either “normal” (119 of the 162), I Ib (15), I In (19), SLSNe-II (7, see also §3.3.4), or SN 1987A-like (2; SN II-87A in Table 1). The relative fraction of SNe I Ib and SNe I In is significantly smaller in the BTS than that reported from LOSS (Li et al. 2011). This could be a direct consequence of the different targeting strategies, although there is one important caveat for SNe I In: for SEDM spectra, the SNe I In represent the most difficult subclass to positively identify because the narrow emission that is the hallmark of SNe I In (see Schlegel 1990) can easily be confused with emission lines from the host galaxy. When SEDM spectra indicate the presence of strong narrow H emission, we have generally attempted to obtain higher resolution spectra. However, some SNe I In may not have strong enough narrow lines to be noticed in a SEDM spectrum. These would be classified as SNe II. In conclusion, SNe identified as Type I In in the BTS all have clear evidence for strong H $\alpha$  emission lines that are significantly broader than would be expected from a galaxy or H II region.

SNe for which there were both Type II and I Ib SNID matches have been manually inspected and classified by comparing the absolute and relative strengths of the H and He features (both emission and absorption) to those seen in typical hydrogen rich SN II spectra. The two SN II-87A events exhibit a nearly identical spectroscopic evolution as

SN 1987A itself, as well as the highly unusual light curve with an initial decline followed by a  $\sim 100$  d long rise (see e.g., Arnett et al. 1989; McCray 1993).

### 3.3.3. *SNe Ibc*

There are 40 H-poor core-collapse SNe in the BTS (excluding 12 SLSNe-I; see §3.3.4). We classify these sources as either SNe Ib (11), Ic (18), Ic-BL (5), Ib/c (3), Ibn (2), or Ic-pec (1). The relative ratio of SNe Ib to Ic in the BTS sample is in agreement with that found by LOSS (Li et al. 2011), though we note that for both surveys the total number of stripped-envelope SNe discovered is relatively small and thus the uncertainties on the relative rates are high.

The SNe Ib clearly show He in their spectra, whereas the SNe Ic do not. There are three events that clearly lack H emission, but where we cannot distinguish between either a Ib or Ic classification (designated as SN Ib/c). The five SNe Ic-BL show very broad absorption features, similar to SN 1998bw (Patat et al. 2001), while the two SNe Ibn display the hallmark narrow He emission lines that define the subtype (Pastorello et al. 2007; Foley et al. 2007). Finally, there is a single event, ZTF18aceqrrs (SN 2018ijp), that lacks both H and He, but has a highly unusual spectroscopic and photometric evolution (Tartaglia et al., in prep.). Thus, we refer to ZTF18aceqrrs as a “peculiar” SN Ic (Ic-pec in Table 1).

### 3.3.4. *SLSNe*

A population of so-called “superluminous” supernovae, with peak optical luminosities ( $M_V$  up to  $-23$  mag) greatly in excess of any known supernova at the time, was first identified in the late 2000s (Quimby et al. 2007) and quickly recognized to occur in both hydrogen-rich (SLSN-II) and hydrogen-free (SLSN-I) varieties (Smith et al. 2007; Gal-Yam et al. 2009; Quimby et al. 2011). More recent surveys have shown that the luminosity distributions of both types of SLSNe overlap with “ordinary” Type II<sub>n</sub> and Ic SNe (De Cia et al. 2018; Angus et al. 2019), and the spectral properties may also form a continuum. Thus, the identification of a particular luminous transient as a superluminous SN, versus merely a luminous SN Ic or II<sub>n</sub> is a challenge. Work by Quimby et al. (2018) does indicate that SLSN-I can be classified with spectral features alone without any luminosity cut. Quimby et al. (2018) show several SLSN-I events with peak luminosity below the traditional  $-21$  mag threshold. However it is still true that post-peak spectra for SLSNe-I and SNe Ic can be very similar (Pastorello et al. 2010). For the purposes of this analysis we use spectroscopic matches to previous “unambiguous” SLSNe-I as the primary determinant, but also place any transient with  $M_g < -20$  mag in this category even if it is well-fit by ordinary SNe (as is the case for nearly all SLSNe-II, which are good matches to SNe II<sub>n</sub>).

In total, we identify 19 SLSNe in the BTS sample. Of these, 12 are classified as H-poor (SLSN-I), and the re-

maining 7 have H emission lines (SLSN-II). This represents  $2.5\% \pm_{1.2}^{2.1}$  of the sample—which is, not surprisingly, a far higher fraction than what was found in galaxy-targeted surveys such as LOSS. LOSS found a single SLSN (SN 2006gy; Foley et al. 2006), which was not recognized as part of a separate class in Li et al. (2011). SLSNe are volumetrically rare, and best found via untargted wide-area surveys (Quimby et al. 2011). The relative rate of SLSNe in ASAS-SN is  $0.4\% \pm_{0.3}^{1.2}$ . This is lower but still consistent with our BTS estimate within the uncertainties on both measurements.

The true fraction of SLSNe may be even higher than what we report here: over this early-survey period our selection methods were biased against SLSNe, because these very long-lived and slow-rising transients were often present in the reference image, such that at peak the subtraction of their own pre-max flux made them appear fainter than they really were and thus less likely to pass the BTS filter.

The detailed analysis of the four SLSN-I discovered during the ZTF commission phase has been submitted for publication by (Lunnan et al. 2019), and a thorough investigation of the full ZTF SLSN sample is underway.

## 4. SN DISTANCES AND HOST GALAXIES

As in Kulkarni et al. (2018), our aim is to measure the RCF of local galaxy catalogs, in this case using SNe from the ZTF BTS. In order to make this measurement we need to both identify the host galaxy for every SN, and measure the SN redshift (for cases where the host redshift is unknown). Using this information it is then possible to calculate the RCF.

### 4.1. *SN redshifts*


In addition to providing SN spectral types, SNID estimates the redshift of the SN it is attempting to classify. We adopt the redshift of the best-matching SNID template as the redshift of the SN,  $z_{\text{SN}}$ .<sup>19</sup> The redshift distribution for BTS SNe is shown in Figure 4, where we adopt the redshift of the host galaxy ( $z_{\text{host}}$ ) when known, otherwise we show  $z_{\text{SN}}$ .

We can estimate the accuracy of the SNID redshift measurements using the subset of BTS SNe that have host galaxies with known redshift (for more on the identification of BTS host galaxies, see §4.2). We find that  $z_{\text{SN}}$  is a good estimator of the host galaxy redshift,  $z_{\text{host}}$ , as summarized in Figure 5 for the 345 ZTF BTS host galaxies with known redshifts.


The main panel in Figure 5 shows the difference between  $z_{\text{SN}}$  and  $z_{\text{host}}$  as a function of  $z_{\text{host}}$ . The residuals show that for SNe Ia there is relatively small scatter ( $\sigma \approx 0.0037$ )<sup>20</sup>

<sup>19</sup> No corrections for Heliocentric, Galactocentric, or host-galaxy-rotation velocity are applied to  $z_{\text{SN}}$ .

<sup>20</sup> We use a robust estimate of scatter by taking half of the difference between the 84<sup>th</sup> and 16<sup>th</sup> percentiles. The sample standard deviation



**Figure 4.** Redshift distribution of BTS SNe Ia (top) and CC SNe (bottom) shown via thin dashed lines. SNe associated with host galaxies of known redshift are shown as solid histograms. If the host galaxy redshift is unknown we adopt  $z_{\text{SN}}$  as the host galaxy redshift, as shown by the thick solid lines. SLSNe are not shown.



**Figure 5.** Difference between host-galaxy redshifts ( $z_{\text{host}}$ ) and SNID-derived redshifts ( $z_{\text{SN}}$ ) as a function of host-galaxy redshift. CC SNe are shown as crosses, while SNe Ia are shown as open circles. The scatter in the difference is nearly constant as a function of redshift. *Top:*  $z_{\text{SN}}$  vs.  $z_{\text{host}}$ . *Right:* A Gaussian KDE of the PDF of the residuals for CC SNe (red) and SNe Ia (blue). The typical scatter is  $\sim 0.004$  (see text), while 95% of all SNe Ia have  $z_{\text{SN}}$  within 0.01 of  $z_{\text{host}}$ .

and virtually no bias in the estimates of  $z_{\text{SN}}$ . The scatter is somewhat higher for CC SNe ( $\sigma \approx 0.0047$ ), where SNID appears to systematically overestimate the true redshift, as is seen in the Gaussian Kernel Density Estimate (KDE) shown in the right panel of Figure 5. We find some evidence for an increased scatter at higher redshifts. For SN Ia hosts with  $z \leq 0.04$ ,  $\sigma \approx 0.0032$ , while for hosts with  $0.08 \leq z < 0.12$   $\sigma \approx 0.0045$  (though the 90<sup>th</sup> percentile widths are nearly identical in these two regions). This increased scatter makes sense as the signal-to-noise ratio typically decreases for higher redshift SNe; for SNe outside the redshift range shown in Figure 5 the typical uncertainty on any individual redshift may be larger than  $\sim 0.004$ . We also find that the scatter is not appreciably larger when restricting the sample to only those SNe that have been observed by SEDM. In the analysis that follows, we assume that  $z_{\text{host}} = z_{\text{SN}}$  for normal SNe Ia, and that the typical uncertainty on  $z_{\text{SN}}$  is 0.004. We note that (i) this is very consistent with the uncertainty ( $\sigma = 0.005$ ) reported in figure 19 in Blondin & Tonry (2007), even though a very different redshift range was used ( $z = 0.1 - 0.8$ ), and (ii) our estimated redshift uncertainty on  $z_{\text{SN}}$  is much larger than the uncertainty in the wavelength calibration of SEDM (figure 13 in Rigault et al. 2019). These two findings indicate that the accuracy of our SNID based redshifts ( $z_{\text{SN}}$ ) is not limited by the low resolution of SEDM.

#### 4.2. Host Galaxy Identification

Correctly associating a newly discovered transient with its host galaxy is a challenging problem, especially when the redshifts of the host candidates are unknown. Simply identifying the closest galaxy (in angular offset) is likely to produce a significant number of misidentifications, especially in the case of nearby SNe for which angular offsets relative to the host nuclei may be quite large.

We use a combination of automated procedures and visual inspection to identify hosts for ZTF BTS SNe. As an initial pass, we query the NASA Extragalactic Database (NED)<sup>21</sup> for all galaxies within  $2'$  of the SN position. Within this list, the galaxy with the smallest angular separation from the SN AND a cataloged redshift in NED is automatically assigned as the host. In cases where there are no cataloged galaxy redshifts within  $2'$  of the SN, the galaxy with the smallest angular separation from the SN is assigned as the host. In cases where the  $z_{\text{host}}$  and  $z_{\text{SN}}$  significantly differ ( $|z_{\text{SN}} - z_{\text{host}}| > 0.05$ ), the galaxy with the smallest angular separation from the SN is assigned as the host. From here we calculated the projected separation,  $d_p$ , between the SN and the host galaxy using the redshift of the SN (see above). We visually inspect host candidates with a projected separation  $d_p \geq 19$  kpc. In

is  $\sim 0.005$ . The sample standard deviation for the recession velocity is  $\Delta v/v \approx 0.14$ .

<sup>21</sup> <https://ned.ipac.caltech.edu/>

most of these cases it is clear that the automated procedure identified a background galaxy that is clearly not the host, in which case we update the host with the NED galaxy with the smallest angular separation from the SN. Following this procedure, there were a total of 12 SN host identifications with separations  $d_p \geq 19$  kpc (9 SNe Ia and 3 SNe II). In 11 of these 12, the host redshift is known *and* that redshift matches that of the SN, providing confidence in these associations. We cannot rule out the possibility that these SNe occurred in faint dwarf galaxies that are associated with the bright galaxy that has been identified as the host. For the purposes of the RCF calculation below, we assume each of these identifications to be correct. For the remaining SN, ZTF18arcetn (SN 2018jag), there is a very bright ( $r' = 14.3$  mag) elliptical galaxy, PSO J015.9596+10.5902, in the field of the SN. The SN is  $\sim 3.2$  Petrosian radii from the galaxy which has an SDSS photometric redshift,  $0.052 \pm 0.012$  (Abolfathi et al. 2018), that agrees with the SN redshift determined by SNID, 0.053 (which gives  $d_p \approx 24$  kpc). For the calculations below we assume that this identification is correct.

Following this procedure, we use deep *i*-band stack images from PS1 to visualize the position of each SN relative to its host galaxy. In the vast majority of cases these images confirmed a clear association between the SN and host galaxy. In some cases, the putative host was extremely faint and a significantly brighter galaxy with only slightly larger angular separation was selected at the likely host. Finally, there were a handful of cases where the association was ambiguous or with a low signal-to-noise ratio PS1 detection. We find that for 40 SNe in the sample, the host identification is ambiguous. We exclude these SNe from the host galaxy analysis below. We thus identify host galaxies for 721 of the 761 SNe in the ZTF BTS sample. Properties of the BTS SNe host galaxies are summarized in Table 3, including notes on each of the SNe identified as having ambiguous hosts. For the 40 ambiguous cases, we find that 29 SNe have no discernible host, including 17 SLSNe, for which host galaxies are typically not found in imaging at the depth of PS1 in this redshift range (e.g., Quimby et al. 2011). The remaining 11 are either roughly equidistant between multiple galaxies of the same brightness, or very close to a faint galaxy, with a significantly brighter galaxy at a similar redshift of the SN residing at much larger angular separation.

The host galaxy coordinates available via NED come from a heterogeneous set of catalogs and surveys, resulting in an astrometric offset between NED and ZTF SN positions, which are measured relative to *Gaia* (Gaia Collaboration et al. 2016). We crossmatch the host positions against the PS1 catalog, which is also calibrated against *Gaia*, in order to place the BTS SNe and host galaxies on the same relative astrometric system. 715 of our initial host positions have counterparts within 2 arcsec in the PS1 DR1 MeanObject table,

which is astrometrically calibrated against *Gaia*. Four of the hosts, those associated with ZTF18aapgrxo (SN 2018bym), ZTF18aayjyub (SN 2018cod), ZTF18acaeous (SN 2018hbu), and ZTF18acrknyn (SN 2018jef), are too faint to be included in the PS1 MeanObject table, and instead we use PS1 positions from the StackObjectThin table. The last two hosts, associated with ZTF18acbzoyn (SN 2018hqu) and ZTF18acdwohd (SN 2018ids) are not detected in the PS1 catalog, and we instead use their positions from SDSS and the *Galaxy Evolution Explorer* (GALEX; Martin et al. 2005), respectively, in Table 3. Following this update of the positions, we recalculate the distribution of host-galaxy separations, which is shown in Figure 6. We find a nearly identical distribution in projected separation for CC SNe and SNe Ia (middle panel of Figure 6). We use a two-sample Kolmogorov-Smirnov (KS) test and a  $\chi^2$  test for independence to determine the statistical difference between the two distributions and find no significant difference between the projected offsets of CC SNe and SNe Ia.

Using the newly identified host offsets, we can additionally examine whether or not there is a bias against finding nuclear SNe in ZTF, as has been found for other surveys (see Holoien et al. 2019). Figure 7 compares the cumulative distribution of angular offsets for bright ( $m_{\text{peak}} \leq 17$  mag) BTS SNe and ASAS-SN.<sup>22</sup> We find remarkably similar distributions between ZTF and ASAS-SN. Both a two-sample KS test and a  $\chi^2$  test for independence do not show a statistically significant difference between the two samples. This suggests that ZTF is not significantly biased against finding nuclear SNe. In contrast, other surveys that detect bright SNe are biased away from galaxy nuclei (Holoien et al. 2019).

Following host identification, we need to determine the absolute magnitude of the host galaxies in order to measure the RCF (see Kulkarni et al. 2018 and below for further details). In this study we focus on the mid-infrared (mid-IR) flux of the host galaxies, primarily for 2 reasons: (i) a galaxy’s mid-IR absolute magnitude serves as a good proxy for the total galactic stellar mass (Wen et al. 2013), and (ii) mid-IR photons are mostly transparent to dust in the Milky Way, meaning that significant reddening corrections are not needed to estimate a galaxy’s absolute magnitude in the mid-IR. The *Wide-field Infrared Survey Explorer* (WISE; Wright et al. 2010) satellite conducted an all-sky survey in the mid-IR, and we use WISE images to determine the brightness of ZTF BTS SN host galaxies at  $3.4 \mu\text{m}$ .

The largest WISE source catalogs (e.g., unWISE; Schlafly et al. 2019) utilize an unresolved (i.e. stellar) point-spread-function (PSF) to measure source flux. This PSF is not appropriate for many of the low-*z* galaxies in our sample, which

<sup>22</sup> As shown in Figure 6 the BTS sample needs to be restricted for a fair comparison to ASAS-SN as high-*z* SNe have smaller angular offsets.

**Table 3.** BTS Host Galaxies

ZTF	Host	$\alpha_{\text{host}}$	$\delta_{\text{host}}$	$z_{\text{host}}$	SN offset	$d_p$	$m_g$	$m_r$	$m_i$	$m_z$	$m_y$	$m_{W1}$	$m_{W2}$	$E(B-V)^a$
Name	Name	(J2000.0)	(J2000.0)		(arcsec)	(kpc)	(mag)	(mag)	(mag)	(mag)	(mag)	(mag)	(mag)	(mag)
ZTF18aabssth	PSO J165.1878+22.2877	11:00:45.07	+22:17:15.8	0.02291	4.44	2.06	16.89	16.49	16.38	16.38	16.61	15.79	16.38	0.015
ZTF18aabxslsv	PSO J157.4639+09.0106	10:29:51.32	+09:00:38.1	0.04797	9.53	8.96	16.17	15.48	15.27	15.02	15.02	14.82	15.30	0.025
ZTF18aaemivw	PSO J158.4280+39.4908	10:33:42.72	+39:29:26.8	0.06807	0.33	0.43	16.66	16.03	15.67	15.53	15.35	14.94	15.08	0.012
ZTF18aagpzjk	PSO J119.8484+16.4214	07:59:23.61	+16:25:16.9	0.01631	26.83	8.91	14.96	14.49	14.33	14.49	14.87	13.85	14.43	0.031
ZTF18aagrdds	PSO J218.3331+41.2658	14:33:19.95	+41:15:56.9	0.01814	5.43	2.00	18.53	17.62	17.45	18.11	18.86	17.32	18.00	0.012
ZTF18aagrtxs	PSO J198.6087+50.9792	13:14:26.08	+50:58:45.0	0.02966	7.94	4.72	...	...	...	13.77	13.68	13.70	14.36	0.010
ZTF18aagstdc	...	...	...	...	...	...	...	...	...	...	...	...	...	...
ZTF18aagtcxj	PSO J248.0475+42.7139	16:32:11.40	+42:42:50.0	0.03240	2.40	1.55	16.72	15.70	15.31	15.00	14.82	14.28	14.75	0.011
ZTF18aahesrp	PSO J128.9390+28.2705	08:35:45.35	+28:16:13.7	...	1.36	1.35	19.43	18.99	18.82	18.69	18.68	19.14	19.86	0.036
ZTF18aahfeiy	PSO J154.3116+43.5219	10:17:14.78	+43:31:18.8	0.07126	10.09	13.71	17.13	16.69	16.38	16.43	16.50	16.19	16.61	0.010

NOTE— This table is available in its entirety in a machine-readable form in the online journal. A portion is shown here for guidance regarding its form and content. Optical *grizy* photometry is taken from PS1 Kron magnitude measurements, while mid-IR *W1* and *W2* photometry are from *Tractor* and *WISE* (see text). All magnitudes are reported in the AB system, and no correction for extinction has been applied.

<sup>a</sup>  $E(B-V)$  is determined using the Schlafly & Finkbeiner (2011) updates to the Schlegel et al. (1998) maps.

are clearly resolved in *WISE* imaging. The recent development of *The Tractor* (Lang et al. 2016a), enables “forced” *WISE* flux measurements, where the mid-IR apertures are determined via sources detected in (relatively) high-resolution optical images. *The Tractor* was used to measure the mid-IR flux of  $\sim 400$  million *WISE* sources that were detected by SDSS (Lang et al. 2016b), and now is also being applied to Legacy Survey (Dey et al. 2019) images. There are 371 BTS host galaxies that have *Tractor* forced mid-IR photometry from both Legacy Survey and SDSS images, while an additional 157 hosts have detections in just Legacy Survey images and another 90 hosts have detections in just SDSS<sup>23</sup> (hereafter we refer to this aperture-matched forced photometry as *Tractor* photometry). We include *Tractor* photometry based on both Legacy Survey and SDSS images in our analysis of the RCF below. For the 371 sources detected both in the Legacy Survey and SDSS we compare the *Tractor* photometry derived from each set of images and measure a sample standard deviation of 0.008 in  $\Delta \text{flux}/\text{flux}$ . This small difference suggests that there are no systematic effects introduced by combining photometry from the two different optical catalogs. Ultimately, this results in 618 host galaxies with  $3.4 \mu\text{m}$  flux measurements that we can use in the analysis of the RCF. Including only the brightest mid-IR galaxies in our sample will bias our final measurement of the RCF, as discussed below.

## 5. THE REDSHIFT COMPLETENESS FACTOR

To calculate the Redshift Completeness Factor (RCF), we follow the methodology originally outlined in Kulkarni et al.

<sup>23</sup> We only retain galaxies with a signal-to-noise ratio  $>5$  in the *W1* filter from the forced photometry catalogs.


(2018). The RCF captures the probability that a random galaxy will have a catalogued spectroscopic redshift as a function of its redshift and IR luminosity. To estimate the RCF we use only SNe Ia, as they occur in both star-forming and passive galaxies, whereas CC SNe would only trace star-forming galaxies. When a ZTF BTS SN has a previously catalogued host galaxy redshift, we consider that a “hit” ( $\text{NED}_z$ ), and when the host does not have a known redshift that is considered a “miss” ( $!\text{NED}_z$ ).

By raw number, there are 512 SNe Ia with known hosts in the BTS,<sup>24</sup> and 227 of them are “hits” (have known redshifts). Thus, over the redshift range sampled by the BTS, the RCF =  $44\% \pm 1\%$  (90% confidence interval).<sup>25</sup> This estimate is significantly lower than what was found for a lower-redshift sample (Kulkarni et al. 2018 estimated RCF  $\approx 75\%$ ). The difference in RCF estimates can be entirely understood by the differing redshift distributions of the two samples. If we restrict our analysis to SNe with  $z \leq 0.03$ , we find the RCF =  $69\% \pm 4\%$  (90% confidence interval), which is consistent with the results reported in Kulkarni et al. (2018). At face value these results show that the RCF decreases as redshift increases, an unsurprising result.

We can further constrain the RCF as a function of redshift and galaxy luminosity by estimating the joint distribution for a galaxy to have a cataloged redshift given its red-

<sup>24</sup> We exclude Ia-02cx, Ia-csm, and Ia-SC events from the RCF calculations. These “peculiar” events account for  $\sim 1\%$  of the BTS SN Ia sample, and would not substantially change our analysis.


<sup>25</sup> As in Kulkarni et al. (2018) we find that the RCF as traced by CC SNe is much higher than that traced by SNe Ia. This suggests that redshift catalogs are more complete for star-forming galaxies than passive galaxies.



**Figure 6.** *Top:* Projected physical separation  $d_p$ , in kpc, between ZTF BTS SNe and their respective host galaxies. SNe Ia are shown via a solid histogram, while CC SNe are shown via a thick, crimson line. The general shape of these distributions are similar to what was found during PTF, with roughly an order of magnitude fewer SNe at  $d_p \approx 10$  kpc, as there are with  $d_p < 1$  kpc (Kasliwal et al. 2012). *Middle:* Cumulative distribution of  $d_p$  for BTS SNe and their hosts. The distribution for SNe Ia and CC SNe is nearly identical; the small discrepancies observed between  $\sim 10$ – $15$  kpc are likely due to small number statistics. *Bottom:* Cumulative distribution of the angular offset, in arcsec, between BTS SNe and their hosts. Given that the average SN Ia is at higher redshift than the average CC SN in the BTS (Figure 4), but the physical separations are the same, it makes sense that SNe Ia, on average, have a smaller angular offset than CC SNe.

shift and  $M_{W1}$ ,  $\text{RCF}(z, M_{W1})$ .<sup>26</sup> A detailed summary of the joint probability RCF calculation is included in Appendix A. To estimate  $\text{RCF}(z, M_{W1})$ , we include only those SNe Ia with identified host galaxies that have a measured IR brightness. This reduces the sample to 442 SNe, of which 213 are

<sup>26</sup> We convert observed  $W1$  magnitudes to absolute magnitude by calculating the distance modulus with `astropy` (Astropy Collaboration et al. 2013), assuming a concordance cosmology with  $\Omega_\Lambda = 0.7$ ,  $\Omega_M = 0.3$ , and  $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ . We also correct for Milky Way extinction, a small effect, using  $E(B - V)$  from Schlafly & Finkbeiner (2011),  $R_V = 3.1$ , and the extinction law from Fitzpatrick & Massa (2007).




**Figure 7.** Cumulative distribution of the angular offset between  $m_{\text{peak}} \leq 17$  mag SNe and their host galaxies found by the ZTF BTS (solid line) and ASAS-SN (dashed line). The distributions are generally very similar, with ASAS-SN doing slightly better at small separations ( $< 5''$ ), however, these differences may simply be due to small number statistics.

“hits.” An estimate of this joint distribution is shown in Figure 8, as well as one-dimensional probabilities  $\text{RCF}(z)$  and  $\text{RCF}(M_{W1})$ . All 512 SNe Ia are used to constrain  $\text{RCF}(z)$ , as a host galaxy identification or brightness measurement is not necessary for that calculation.


From Figure 8 it is clear that the analysis in Kulkarni et al. (2018) was significantly limited by the lower redshift sample that was available at that time. For example, there are no “hits” for  $z > 0.12$ , and the  $\text{RCF}(z)$  tends towards zero at high redshifts, whereas Kulkarni et al. (2018) only found mild evidence that the  $\text{RCF}(z)$  decreases with  $z$ . Unsurprisingly, we still find that low- $z$  and massive galaxies are the most likely to be catalogued. At higher redshifts ( $z > 0.1$ ), only very massive galaxies ( $M_{W1} \lesssim -22$  mag AB, comparable to that of the Milky Way) are catalogued.

The decline in  $\text{RCF}(z)$  as a function of  $z$  has important ramifications for the electromagnetic (EM) follow-up of gravitational wave events. The typical localization areas for 2 and 3 detector networks is several hundred to several thousand  $\text{deg}^2$  (Kasliwal & Nissanke 2014). One strategy to mitigate against these large areas that would be impossible to search even with modest field-of-view instruments, is to target known galaxies within the LIGO localization volume (e.g., Gehrels et al. 2016). During O3, the horizon distance for binary neutron star (BNS) mergers is  $\sim 200$  Mpc (see O3 alerts for S190425z, S190510g, S190901ap, S190910h; Ligo Scientific Collaboration & VIRGO Collaboration 2019a,b,c,d), roughly corresponding to  $z \approx 0.05$ . According to the BTS, for  $z \leq 0.05$  the  $\text{RCF} \approx 63\%$ , while integrating our best model inference from  $z = 0$  to 0.05 yields  $\text{RCF} \approx 57\%$ . Thus, targeted efforts to identify EM radiation from BNS mergers are likely to miss 1/3, or more, of all potential host galaxies for the EM transient. These numbers become significantly worse for events discovered at a distance  $z > 0.05$ . Any future efforts to quantify the rate of EM counterparts to gravitational wave



**Figure 8.** Absolute  $W_1$ -band magnitude,  $M_{W_1, \text{host}}$ , vs. redshift,  $z$ , for the host galaxies of SNe Ia in the ZTF BTS. Galaxies with known redshifts (from NED or other databases) prior to SN discovery are shown as magenta pluses, while those lacking redshifts (!NED $_z$ ) are shown as gold circles. The dashed line roughly corresponds to the *WISE* detection limit  $m_{W_1, \text{limit}} \approx 20.629$  mag [Schlafly et al. 2019](#). The shaded background shows the probability of a host galaxy having a cataloged redshift given its redshift and  $M_{W_1}$  ( $\text{RCF}(z, M_{W_1})$ ), based on 442 galaxies with *WISE* detections. The top and right plots show the probability of a host galaxy having a cataloged redshift given *only* its redshift,  $\text{RCF}(z)$ , or  $M_{W_1}$ ,  $\text{RCF}(M_{W_1})$ , respectively. In these two plots the solid lines show the median value of the RCF, while the shaded region corresponds to the 90% credible region on the RCF.





**Figure 9.** Redshift completeness factor as a function of *apparent* host galaxy brightness, as traced by the  $r_{\text{PS1}}$  Kron mag measurement. The solid purple curve shows the median value for all hosts in PS1 (95% of all identified ZTF BTS hosts), while the green, dot-dashed curve shows hosts within the SDSS imaging footprint (72% of all hosts), and the orange, dashed curve shows hosts outside SDSS (23% of all hosts). The RCF is higher within the SDSS footprint, likely due to the many SDSS spectroscopic redshift surveys. For all galaxies within PS1, the RCF = 0.5 at  $r_{\text{PS1}} \approx 16.9$  mag, while within the SDSS footprint, the RCF = 0.5 at  $r_{\text{PS1}} \approx 17.4$  mag, and outside SDSS the RCF = 0.5 at  $r_{\text{PS1}} \approx 15.5$  mag.

events should account for the fraction of “missing” galaxies that we have identified with the BTS.

### 5.1. Catalog Completeness as a Function of Galaxy Brightness

To date SDSS has been the most prolific survey in terms of spectroscopically measuring galaxy redshifts, with  $\sim 2.8$  million catalogued galaxies and counting (Aguado et al. 2019). At this stage, significant improvements to the RCF will require tens of millions of new redshift measurements. Fortunately, within the next few years we will enter the era of supremely multiplexed spectrographs [e.g., The Dark Energy Spectroscopic Instrument (DESI),<sup>27</sup> the Subaru Prime Focus Spectrograph (PFS),<sup>28</sup> Euclid,<sup>29</sup> the Wide Field Infrared Survey Telescope (WFIRST)<sup>30</sup>], which could dramatically increase the number of galaxies with known spectroscopic redshifts within the local universe. The best strategy to this end is to obtain spectra of bright galaxies with currently unknown redshifts, as is planned as part of the DESI Bright Galaxy Survey (DESI Collaboration et al. 2016). Using the

<sup>27</sup> <https://www.desi.lbl.gov/>

<sup>28</sup> <https://pfs.ipmu.jp/>

<sup>29</sup> <https://sci.esa.int/web/euclid/>

<sup>30</sup> <https://wfIRST.gsfc.nasa.gov/>

same methodology described above, we can estimate the RCF as a function of galaxy brightness (as traced by the PS1  $r$ -band), rather than  $z$  or  $M_{W1}$ . The results from this exercise are summarized in Figure 9.

From Figure 9 it is clear that the vast majority of extremely bright galaxies ( $r_{\text{PS1}} \lesssim 14$  mag) have catalogued redshifts. Figure 9 also shows the RCF for galaxies that are and are not within the SDSS imaging footprint,<sup>31</sup> and a comparison of these two curves highlights the crucial role that SDSS played in terms of identifying galaxies in the local universe and spectroscopically measuring their redshifts. SDSS pushes the completeness of redshift catalogs  $\sim 2$  mag fainter than what is observed outside the SDSS footprint. From our analysis, it is possible to estimate the completeness of existing redshift catalogs as a function of depth, and we find that catalogs are currently  $\sim 90\%$ ,  $50\%$ , and  $10\%$  complete to a depth of  $r_{\text{PS1}} \approx 14.7$  mag,  $16.9$  mag, and  $19.1$  mag respectively. Since the DESI bright galaxy survey is an order-of-magnitude increase over SDSS, Figure 9 makes it clear that the DESI bright galaxy survey will dramatically improve our knowledge of which galaxies reside in the local universe.

## 6. SUMMARY AND CONCLUSIONS

We have presented early results from the ZTF Bright Transient Survey. The BTS utilizes publicly announced discoveries from the ZTF MSIP surveys to spectroscopically classify all extragalactic transients that peak brighter than  $18.5$  mag. Simple filters are applied to the ZTF alert stream with the aim of minimizing false negatives, and we estimate that  $\sim 95\%$  of all  $m_{\text{peak}} \leq 18.5$  mag SNe in the BTS are spectroscopically classified (during the period of the survey in 2018). This effort has resulted in the classification of 761 SNe.

Spectroscopic observations are primarily conducted with SEDM, which is optimized to classify SN with  $m < 19$  mag. For sources that are inaccessible to SEDM, we utilize any available resource in order to obtain a classification. Final BTS classifications utilize spectral template matching, via SNID, and visual inspection of the spectra and (publically available) light curves. Of the 761 classified BTS SNe, the majority are SNe Ia (547), followed by CC SNe, both SNe II (155) and SNe Ib/c (40), with a relatively small number of SLSNe (19). The relative fraction of each of these types of SNe agrees with what has been found in previous studies (Li et al. 2011; Holoiien et al. 2019).

In this early release paper, we have focused on measuring the redshift completeness of local galaxy catalogs by using SNe Ia as a relatively unbiased tracer of galaxies in the local universe. By raw number, we find that less than half of the

<sup>31</sup> This is determined by performing a cross match between BTS host positions and the SDSS imaging catalog. Any hosts with SDSS sources within  $1'$  are assumed to be within the SDSS imaging footprint.

BTS SN host galaxies have known spectroscopic redshifts. In more detail, we find that the RCF falls steeply as a function of redshift, with only  $\sim$ half of all galaxies having known redshifts at  $z \approx 0.05$ , and  $\lesssim 20\%$  of galaxies having known redshifts at  $z \approx 0.1$ . The “missing” galaxies with unknown redshifts have important ramifications when searching for electromagnetic counterparts to multimessenger astronomical events, and suggest that the most complete method to find the EM counterparts for events at  $d \gtrsim 100$  Mpc is to tile the entire error regions associated with gravitational wave (GW) or neutrino event alerts. However, even if this is done the same problem arises again when deciding which candidates among those found by the wide-field searches to follow up spectroscopically. Given the resources available, typically candidates with known distances, that also coincide with the distance constraints in a GW alert, are heavily prioritized for follow-up (e.g., Andreoni et al. 2019; Coughlin et al. 2019). Incorporation of redshifts from the Census of the Local Universe narrowband H $\alpha$  catalog (Cook et al. 2019) may alleviate this problem to some extent in the near future.

The combination of ZTF and SEDM has illustrated the power of focused efforts in the era of very large time-domain surveys. Within the next few years the Large Synoptic Survey Telescope (LSST; Ivezić et al. 2008) will begin full survey operations. LSST will increase the volume of transient discoveries by an order of magnitude relative to on-going surveys in much the same way that PTF, PS1, and others built upon LOSS, and ATLAS and ZTF have built upon those surveys. We have already reached the era where true spectroscopic completeness is impossible for SN surveys, and LSST will greatly exacerbate this problem. Nevertheless, the use of an ultra-low-resolution instrument has allowed us to spectroscopically classify a nearly-complete subset of the discoveries made by ZTF (those with  $m_{\text{peak}} \leq 18.5$  mag). The simple focus of the BTS – classify all the bright transients – has resulted in the largest systematic classification of SNe to date. 761 SNe are included in this early release paper, while the inclusion of 2019 results will eventually bring this number to  $>1800$ . In addition to measuring the RCF, as was done here, our growing sample can be used to: determine volumetric SN rates, measure the expansion of the Universe using low- $z$  SNe Ia, study the demographics of CC SNe, and measure the luminosity function of a wide range of transient phenomena. This, despite the fact that BTS only targets a tiny minority of all transients discovered by ZTF. The BTS demonstrates that a focused triage of an otherwise overwhelming discovery stream can lead to both impactful and novel results.

Based on observations obtained with the Samuel Oschin Telescope 48-inch and the 60-inch Telescope at the Palomar Observatory as part of the Zwicky Transient Facility project. ZTF is supported by the National Science Foundation under Grant No. AST-1440341 and a collaboration including Caltech, IPAC, the Weiz-

mann Institute for Science, the Oskar Klein Center at Stockholm University, the University of Maryland, the University of Washington, Deutsches Elektronen- Synchrotron and Humboldt University, Los Alamos National Laboratories, the TANGO Consortium of Taiwan, the University of Wisconsin at Milwaukee, and Lawrence Berkeley National Laboratories. Operations are conducted by COO, IPAC, and UW.

This work was supported by the GROWTH project funded by the National Science Foundation under PIRE Grant No 1545949. The ZTF forced-photometry service was funded under the Heising-Simons Foundation grant #12540303 (PI: Graham). A.A.M. is funded by the Large Synoptic Survey Telescope Corporation, the Brinson Foundation, and the Moore Foundation in support of the LSSTC Data Science Fellowship Program, he also receives support as a CIERA Fellow by the CIERA Postdoctoral Fellowship Program (Center for Interdisciplinary Exploration and Research in Astrophysics, Northwestern University). M.R. has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 759194 - USNAC). The Oskar Klein Centre is funded by the Swedish Research Council. A.Y.Q.H. is supported by a National Science Foundation Graduate Research Fellowship under Grant No. DGE1144469.

Partially based on observations made with the Nordic Optical Telescope, operated by the Nordic Optical Telescope Scientific Association at the Observatorio del Roque de los Muchachos, La Palma, Spain, of the Instituto de Astrofísica de Canarias. Some of the data presented here were obtained with ALFOSC, which is provided by the Instituto de Astrofísica de Andalucía (IAA) under a joint agreement with the University of Copenhagen and NOTSA.

Some of the data presented herein were obtained at the W. M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California, and NASA; the observatory was made possible by the generous financial support of the W. M. Keck Foundation.

This paper is partly based on observations made with the Italian Telescopio Nazionale Galileo (TNG) operated on the island of La Palma by the Fundación Galileo Galilei of the INAF (Istituto Nazionale di Astrofisica) at the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. Partially based on observations obtained with the Apache Point Observatory 3.5-meter telescope, which is owned and operated by the Astrophysical Research Consortium. Partially based on observations from the LCOGT network. Partially based on public observations collected at the WHT, operated on the island of La Palma by the Isaac Newton Group.

The Liverpool Telescope is operated on the island of La Palma by Liverpool John Moores University in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias with financial support from the UK Science and Technology Facilities Council.

The SED Machine is based upon work supported by the National Science Foundation under Grant No. 1106171.

We thank students Shaney Sze and Ho Ko for assisting with the manual candidate vetting during the summer of 2018.

*Software:* SNID (Blondin & Tonry 2007), astropy (Astropy Collaboration et al. 2013), scipy (Jones et al. 2001), matplotlib (Hunter 2007), pandas (McKinney 2010), emcee (Foreman-Mackey et al. 2013), corner (Foreman-Mackey 2016),

MultinomCI (Signorell & et al. 2019), PYSEDm (Rigault et al. 2019), IRAF (Tody 1986), PyRAF (Science Software Branch at

STScI 2012), pyraf-dbsp (Bellm & Sesar 2016), LPipe (Perley 2019), pyDIS (Davenport et al. 2016).

## APPENDIX

### A. MEASURING THE CONDITIONAL PROBABILITY OF THE RCF

To estimate the conditional probability that a galaxy has a catalogued redshift based on its distance and IR-luminosity, we model the data  $X$  with the Bernoulli distribution

$$X \sim \text{Bern}(p), \quad (\text{A1})$$

where  $p$  is parameterized with a logistic function with dependence on both redshift  $z$  and host galaxy luminosity:

$$p(z, M, \theta) = \frac{1}{1 + \exp(az + bM - c)}, \quad (\text{A2})$$

with host-galaxy absolute magnitude  $M$ , and  $\theta$  representing the model parameters:  $a$ ,  $b$ , and  $c$ , which need to be determined. The precise analytic dependence of  $p$  on  $z$  and  $M$  may not be logistic, however, the purpose of this exercise is to provide a general sense for how the RCF relies on  $z$  and  $M$ . Given that it smoothly transitions over an exponential length scale from 1 to 0, the logistic function works well for this general purpose.

From here it follows that the probability of a host galaxy having a previously catalogued redshift is:

$$Pr(q) = \begin{cases} p(z, M, \theta), & \text{if } q = \text{NED}_z \\ 1 - p(z, M, \theta), & \text{if } q = \text{!NED}_z \end{cases} \quad (\text{A3})$$

and the likelihood of the observations given the data and model parameters is:

$$Pr(q_k | z_k, M_k, \theta) = \prod_{k=1}^K p(z_k, M_k, \theta)^{q_k} (1 - p(z_k, M_k, \theta))^{1-q_k}, \quad (\text{A4})$$

where  $k$  represents the individual observations and  $q_k = 1$  for  $\text{NED}_z$  galaxies and  $q_k = 0$  for  $\text{!NED}_z$  galaxies.

To estimate the model parameters we adopt wide, flat priors, and use the affine-invariant Markov Chain Monte Carlo (MCMC) ensemble sampling technique described by Goodman & Weare (2010), as implemented in the `emcee` software package (Foreman-Mackey et al. 2013). For  $a$  and  $b$  we adopt flat priors bounded between 0 and  $10^6$ . For  $c$  we adopt a flat prior between  $-100$  and  $100$ . We use 25 walkers within the sample, and run the ensemble until it has “converged,” which we define as  $>100$  times longer than the average auto-correlation length of the individual chains from each walker. We find there is a strong covariance between  $b$  and  $c$ , whereas  $a$  is relatively independent, as shown in the corner plot in Figure 10. We will happily make our posterior samples available to readers upon request.

We also constrain the RCF as a function of the host redshift,  $z$ , or host galaxy luminosity, individually. We do this separately from the analysis above, while using the same MCMC procedure with  $p$  in Equations A3 and A4 replaced by

$$p(z, \theta) = \frac{1}{1 + \exp(az - c)}, \quad (\text{A5})$$


for redshift, and

$$p(M, \theta) = \frac{1}{1 + \exp(bM - c)}, \quad (\text{A6})$$

for host galaxy luminosity (where, again, we use absolute magnitude  $M_{W1}$  instead of total luminosity). The results of this procedure are shown in the side panels of Figure 8. In these panels the solid lines show the median value of  $p(z)$ ,  $\text{RCF}(z)$ , and  $p(M)$ ,  $\text{RCF}(M)$ , while the shaded region shows the 90% credible regions for  $p(z)$  and  $p(M)$ .

## REFERENCES

- |  |   |
|--|---|
| Abolfathi, B., Aguado, D. S., Aguilar, G., et al. 2018, <i>ApJS</i> , 235, 42  | Andreoni, I., Goldstein, D. A., Anand, S., et al. 2019, <i>ApJL</i> , 881, L16                    |
| Aguado, D. S., Ahumada, R., Almeida, A., et al. 2019, <i>ApJS</i> , 240, 23  | Angus, C. R., Smith, M., Sullivan, M., et al. 2019, <i>MNRAS</i> , 487, 2215                      |
| Aldering, G., Adam, G., Antilogus, P., et al. 2002, in <i>Proc. SPIE</i> , Vol. 4836, Survey and Other Telescope Technologies and Discoveries, ed. J. A. Tyson & S. Wolff, 61–72 | Arnett, W. D., Bahcall, J. N., Kirshner, R. P., & Woosley, S. E. 1989, <i>ARA&amp;A</i> , 27, 629 |
|  | Astier, P., Guy, J., Regnault, N., et al. 2006, <i>A&amp;A</i> , 447, 31                          |



**Figure 10.** Corner plot showing the posterior distribution of  $a$ ,  $b$ , and  $c$ , used to constrain the joint distribution of the RCF on  $z$  and  $M_{W1}$ ,  $\text{RCF}(z, M_{W1})$ .

- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
- Baade, W., & Zwicky, F. 1934, *Physical Review*, 46, 76
- Bellm, E. C. 2016, *PASP*, 128, 084501
- Bellm, E. C., & Sesar, B. 2016, pyraf-dbsp: Reduction pipeline for the Palomar Double Beam Spectrograph, Astrophysics Source Code Library, , ascl:1602.002
- Bellm, E. C., Kulkarni, S. R., Graham, M. J., et al. 2019a, *PASP*, 131, 018002
- Bellm, E. C., Kulkarni, S. R., Barlow, T., et al. 2019b, *PASP*, 131, 068003
- Ben-Ami, S., Konidaris, N., Quimby, R., et al. 2012, in *Proc. SPIE*, Vol. 8446, Ground-based and Airborne Instrumentation for Astronomy IV, 844686
- Blagorodnova, N., Neill, J. D., Walters, R., et al. 2018, *PASP*, 130, 035003
- Blondin, S., Dessart, L., & Hillier, D. J. 2015, *MNRAS*, 448, 2766
- Blondin, S., & Tonry, J. L. 2007, in *American Institute of Physics Conference Series*, Vol. 924, The Multicolored Landscape of Compact Objects and Their Explosive Origins, ed. T. di Salvo, G. L. Israel, L. Piersant, L. Burderi, G. Matt, A. Tornambe, & M. T. Menna, 312–321
- Branch, D., Dang, L. C., Hall, N., et al. 2006, *PASP*, 118, 560
- Cenko, S. B., Fox, D. B., Moon, D.-S., et al. 2006, *PASP*, 118, 1396
- Chambers, K. C., Magnier, E. A., Metcalfe, N., et al. 2016, arXiv e-prints, arXiv:1612.05560
- Cook, D. O., Kasliwal, M. M., Van Sistine, A., et al. 2019, *ApJ*, 880, 7
- Cortini, G. 2018, Transient Name Server Discovery Report, 2018-393, 1
- Coughlin, M. W., Ahumada, T., Anand, S., et al. 2019, arXiv e-prints, arXiv:1907.12645
- Davenport, J., de Val-Borro, M., & Wilkinson, T. D. 2016, doi:10.5281/zenodo.58753
- De Cia, A., Gal-Yam, A., Rubin, A., et al. 2018, *ApJ*, 860, 100
- DESI Collaboration, Aghamousa, A., Aguilar, J., et al. 2016, arXiv e-prints, arXiv:1611.00036
- Dey, A., Schlegel, D. J., Lang, D., et al. 2019, *AJ*, 157, 168
- Duev, D. A., Mahabal, A., Masci, F. J., et al. 2019, arXiv e-prints, arXiv:1907.11259
- Filippenko, A. V. 1997, *ARA&A*, 35, 309
- Filippenko, A. V., Li, W. D., Treffers, R. R., & Modjaz, M. 2001, in *Astronomical Society of the Pacific Conference Series*, Vol. 246, IAU Colloq. 183: Small Telescope Astronomy on Global Scales, ed. B. Paczynski, W.-P. Chen, & C. Lemme, 121–+
- Fitzpatrick, E. L., & Massa, D. 2007, *ApJ*, 663, 320
- Flesch, E. W. 2015, *PASA*, 32, e010
- Foley, R. J., Li, W., Moore, M., et al. 2006, Central Bureau Electronic Telegrams, 695, 1
- Foley, R. J., Smith, N., Ganeshalingam, M., et al. 2007, *ApJL*, 657, L105
- Foreman-Mackey, D. 2016, *The Journal of Open Source Software*, 24, doi:10.21105/joss.00024. <http://dx.doi.org/10.5281/zenodo.45906>
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, 125, 306
- Fox, O. D., Azalee Bostroem, K., Van Dyk, S. D., et al. 2014, *ApJ*, 790, 17

- Frieman, J. A., Bassett, B., Becker, A., et al. 2008, *AJ*, 135, 338
- Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, *A&A*, 595, A1
- Gal-Yam, A. 2012, *Science*, 337, 927
- Gal-Yam, A., Mazzali, P., Ofek, E. O., et al. 2009, *Nature*, 462, 624
- Galama, T. J., Vreeswijk, P. M., van Paradijs, J., et al. 1998, *Nature*, 395, 670
- Gehrels, N., Cannizzo, J. K., Kanner, J., et al. 2016, *ApJ*, 820, 136
- Goobar, A., & Leibundgut, B. 2011, *Annual Review of Nuclear and Particle Science*, 61, 251
- Goodman, J., & Weare, J. 2010, *Communications in Applied Mathematics and Computational Science*, 5, 65
- Goodman, L. A. 1965, *Technometrics*, 7, 247. <https://www.tandfonline.com/doi/abs/10.1080/00401706.1965.10490252>
- Gorbovskey, E. S., Lipunov, V. M., Kornilov, V. G., et al. 2013, *Astronomy Reports*, 57, 233
- Graham, M. J., Kulkarni, S. R., Bellm, E. C., et al. 2019, *PASP*, 131, 078001
- Grzegorzec, J. 2018, *Transient Name Server Discovery Report*, 2018-582, 1
- Gutiérrez, C. P., Anderson, J. P., Hamuy, M., et al. 2017, *ApJ*, 850, 89
- Helfand, D. J., White, R. L., & Becker, R. H. 2015, *ApJ*, 801, 26
- Hirata, K., Kajita, T., Koshihara, M., et al. 1987, *Physical Review Letters*, 58, 1490
- Ho, A. Y. Q., Phinney, E. S., Ravi, V., et al. 2019, *ApJ*, 871, 73
- Hodgkin, S. T., Wyrzykowski, L., Blagorodnova, N., & Kposov, S. 2013, *Philosophical Transactions of the Royal Society of London Series A*, 371, 20120239
- Holoien, T. W. S., Stanek, K. Z., Kochanek, C. S., et al. 2017a, *MNRAS*, 464, 2672
- Holoien, T. W. S., Brown, J. S., Stanek, K. Z., et al. 2017b, *MNRAS*, 467, 1098
- . 2017c, *MNRAS*, 471, 4966
- Holoien, T. W. S., Brown, J. S., Vallety, P. J., et al. 2019, *MNRAS*, 484, 1899
- Howell, D. A. 2001, *ApJL*, 554, L193
- Hunter, J. D. 2007, *Computing In Science & Engineering*, 9, 90
- Itagaki, K. 2018a, *Transient Name Server Discovery Report*, 2018-1614, 1
- . 2018b, *Transient Name Server Discovery Report*, 2018-1766, 1
- Ivezić, Ž., Tyson, J. A., Acosta, E., et al. 2008, *ArXiv e-prints*, arXiv:0805.2366
- Jarrett, T. H., Cohen, M., Masci, F., et al. 2011, *ApJ*, 735, 112
- Jones, E., Oliphant, T., Peterson, P., et al. 2001, *SciPy: Open source scientific tools for Python*, , [Online; accessed |today;]. <http://www.scipy.org/>
- Kasen, D. 2006, *ApJ*, 649, 939
- Kasliwal, M. M., & Nissanke, S. 2014, *ApJL*, 789, L5
- Kasliwal, M. M., Kulkarni, S. R., Gal-Yam, A., et al. 2012, *ApJ*, 755, 161
- Kasliwal, M. M., Cannella, C., Bagdasaryan, A., et al. 2019, *PASP*, 131, 038003
- Kulkarni, S. R., Perley, D. A., & Miller, A. A. 2018, *ApJ*, 860, 22
- Lang, D., Hogg, D. W., & Mykytyn, D. 2016a, *The Tractor: Probabilistic astronomical source detection and measurement*, , ascl:1604.008
- Lang, D., Hogg, D. W., & Schlegel, D. J. 2016b, *AJ*, 151, 36
- Law, N. M., Kulkarni, S. R., Dekany, R. G., et al. 2009, *PASP*, 121, 1395
- Li, W., Leaman, J., Chornock, R., et al. 2011, *MNRAS*, 412, 1441
- Li, W. D., Filippenko, A. V., Treffers, R. R., et al. 2000, in *American Institute of Physics Conference Series*, Vol. 522, American Institute of Physics Conference Series, ed. S. S. Holt & W. W. Zhang, 103–106
- Ligo Scientific Collaboration, & VIRGO Collaboration. 2019a, *GRB Coordinates Network, Circular Service*, 24168
- . 2019b, *GRB Coordinates Network, Circular Service*, 24442
- . 2019c, *GRB Coordinates Network, Circular Service*, 25606
- . 2019d, *GRB Coordinates Network, Circular Service*, 25707
- Liu, Y.-Q., Modjaz, M., Bianco, F. B., & Graur, O. 2016, *ApJ*, 827, 90
- Lunnan, R., Yan, L., Perley, D. A., et al. 2019, *arXiv e-prints*, arXiv:1910.02968
- Mahabal, A., Rebbapragada, U., Walters, R., et al. 2019, *PASP*, 131, 038002
- Maraston, C., Pforr, J., Henriques, B. M., et al. 2013, *MNRAS*, 435, 2764
- Martin, D. C., Fanson, J., Schiminovich, D., et al. 2005, *ApJL*, 619, L1
- Masci, F. J., Laher, R. R., Rusholme, B., et al. 2019, *PASP*, 131, 018003
- McCray, R. 1993, *ARA&A*, 31, 175
- McKinney, W. 2010, in *Proceedings of the 9th Python in Science Conference*, ed. S. van der Walt & J. Millman, 51 – 56
- Modjaz, M., Liu, Y. Q., Bianco, F. B., & Graur, O. 2016, *ApJ*, 832, 108
- Modjaz, M., Blondin, S., Kirshner, R. P., et al. 2014, *AJ*, 147, 99
- Nordin, J., Brinnel, V., van Santen, J., et al. 2019, *arXiv e-prints*, arXiv:1904.05922
- Norgaard-Nielsen, H. U., Hansen, L., Jorgensen, H. E., Aragon Salamanca, A., & Ellis, R. S. 1989, *Nature*, 339, 523
- Oke, J. B., & Gunn, J. E. 1982, *PASP*, 94, 586
- Oke, J. B., Cohen, J. G., Carr, M., et al. 1995, *PASP*, 107, 375
- Papadogiannakis, S., Dhawan, S., Morosin, R., & Goobar, A. 2019a, *MNRAS*, 485, 2343
- Papadogiannakis, S., Goobar, A., Amanullah, R., et al. 2019b, *MNRAS*, 483, 5045

- Pastorello, A., Smartt, S. J., Mattila, S., et al. 2007, *Nature*, 447, 829
- Pastorello, A., Smartt, S. J., Botticella, M. T., et al. 2010, *ApJL*, 724, L16
- Patat, F., Cappellaro, E., Danziger, J., et al. 2001, *ApJ*, 555, 900
- Patterson, M. T., Bellm, E. C., Rusholme, B., et al. 2019, *PASP*, 131, 018001
- Perley, D. A. 2019, *PASP*, 131, 084503
- Perlmutter, S., Aldering, G., Goldhaber, G., et al. 1999, *ApJ*, 517, 565
- Piascik, A. S., Steele, I. A., Bates, S. D., et al. 2014, in *Proc. SPIE*, Vol. 9147, *Ground-based and Airborne Instrumentation for Astronomy V*, 91478H
- Prentice, S. J., Maguire, K., Smartt, S. J., et al. 2018, *ApJL*, 865, L3
- Quimby, R. M., Aldering, G., Wheeler, J. C., et al. 2007, *ApJL*, 668, L99
- Quimby, R. M., Kulkarni, S. R., Kasliwal, M. M., et al. 2011, *Nature*, 474, 487
- Quimby, R. M., De Cia, A., Gal-Yam, A., et al. 2018, *ApJ*, 855, 2
- Riess, A. G., Filippenko, A. V., Challis, P., et al. 1998, *AJ*, 116, 1009
- Rigault, M., Neill, J. D., Blagorodnova, N., et al. 2019, *A&A*, 627, A115
- Schlafly, E. F., & Finkbeiner, D. P. 2011, *ApJ*, 737, 103
- Schlafly, E. F., Meisner, A. M., & Green, G. M. 2019, *ApJS*, 240, 30
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Schlegel, E. M. 1990, *MNRAS*, 244, 269
- Schmidt, B. P., Kirshner, R. P., Eastman, R. G., et al. 1993, *Nature*, 364, 600
- Science Software Branch at STScI. 2012, PyRAF: Python alternative for IRAF, *Astrophysics Source Code Library*, , , ascl:1207.011
- Secrest, N. J., Dudik, R. P., Dorland, B. N., et al. 2015, *ApJS*, 221, 12
- Shappee, B., Prieto, J., Stanek, K. Z., et al. 2014, in *American Astronomical Society Meeting Abstracts*, Vol. 223, *American Astronomical Society Meeting Abstracts #223*, 236.03
- Signorell, A., & et al. 2019, *DescTools: Tools for Descriptive Statistics*, *r* package version 0.99.29.  
<https://cran.r-project.org/package=DescTools>
- Silverman, J. M., Foley, R. J., Filippenko, A. V., et al. 2012a, *MNRAS*, 425, 1789
- . 2012b, *MNRAS*, 425, 1789
- Smartt, S. J., Valenti, S., Fraser, M., et al. 2015, *A&A*, 579, A40
- Smith, N., Li, W., Foley, R. J., et al. 2007, *ApJ*, 666, 1116
- Soumagnac, M. T., & Ofek, E. O. 2018, *PASP*, 130, 075002
- Sun, P., Lau, A., Liu, J., et al. 2018, *Transient Name Server Discovery Report*, 2018-2007, 1
- Tachibana, Y., & Miller, A. A. 2018, *PASP*, 130, 128001
- Tanaka, Y. 2018, *Transient Name Server Discovery Report*, 2018-469, 1
- Tartaglia, L., Sand, D. J., Valenti, S., et al. 2018, *ApJ*, 853, 62
- Taubenberger, S. 2017, *The Extremes of Thermonuclear Supernovae*, 317
- Tody, D. 1986, in *Proc. SPIE*, Vol. 627, *Instrumentation in astronomy VI*, ed. D. L. Crawford, 733
- Tonry, J. L., Denneau, L., Heinze, A. N., et al. 2018, *PASP*, 130, 064505
- Turatto, M., Cappellaro, E., Danziger, I. J., et al. 1993, *MNRAS*, 262, 128
- Wen, X.-Q., Wu, H., Zhu, Y.-N., et al. 2013, *MNRAS*, 433, 2946
- Williamson, M., Modjaz, M., & Bianco, F. B. 2019, *ApJL*, 880, L22
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, 140, 1868
- Zackay, B., Ofek, E. O., & Gal-Yam, A. 2016, *ApJ*, 830, 27
- Zhang, M., Ding, Y., Liu, S., et al. 2018, *Transient Name Server Discovery Report*, 2018-1393, 1
- Zhang, T.-M., Wang, X.-F., Chen, J.-C., et al. 2015, *Research in Astronomy and Astrophysics*, 15, 215
- Zwicky, F. 1938a, *PASP*, 50, 215
- . 1938b, *ApJ*, 88, 529
- . 1942, *ApJ*, 96, 28