

---

# A Reduction from Reinforcement Learning to No-Regret Online Learning

---

Ching-An Cheng  
Georgia Tech

Remi Tachet des Combes  
Microsoft Research Montreal

Byron Boots  
UW

Geoff Gordon  
Microsoft Research Montreal

## Abstract

We present a reduction from reinforcement learning (RL) to no-regret online learning based on the saddle-point formulation of RL, by which *any* online algorithm with sublinear regret can generate policies with provable performance guarantees. This new perspective decouples the RL problem into two parts: regret minimization and function approximation. The first part admits a standard online-learning analysis, and the second part can be quantified independently of the learning algorithm. Therefore, the proposed reduction can be used as a tool to systematically design new RL algorithms. We demonstrate this idea by devising a simple RL algorithm based on mirror descent and the generative-model oracle. For any  $\gamma$ -discounted tabular RL problem, with probability at least  $1 - \delta$ , it learns an  $\epsilon$ -optimal policy using at most  $\tilde{O}\left(\frac{|\mathcal{S}||\mathcal{A}|\log(\frac{1}{\delta})}{(1-\gamma)^4\epsilon^2}\right)$  samples. Furthermore, this algorithm admits a direct extension to linearly parameterized function approximators for large-scale applications, with computation and sample complexities independent of  $|\mathcal{S}|, |\mathcal{A}|$ , though at the cost of potential approximation bias.

## 1 INTRODUCTION

Reinforcement learning (RL) is a fundamental problem for sequential decision making in unknown environments. One of its core difficulties, however, is the need for algorithms to infer long-term consequences based on limited, noisy, short-term feedback. As a result, designing RL algorithms that are both scalable and provably sample efficient has been challenging.

---

In this work, we revisit the classic linear-program (LP) formulation of RL [1, 2] in an attempt to tackle this long-standing question. We focus on the associated saddle-point problem of the LP (given by Lagrange duality), which has recently gained traction due to its potential for computationally efficient algorithms with theoretical guarantees [3–11]. But in contrast to these previous works based on stochastic approximation, here we consider a reformulation through the lens of online learning, i.e. regret minimization. Since the pioneering work of Gordon [12], Zinkevich [13], online learning has evolved into a ubiquitous tool for systematic design and analysis of iterative algorithms. Therefore, if we can identify a reduction from RL to online learning, we can potentially leverage it to build efficient RL algorithms.

We will show this idea is indeed feasible. We present a reduction by which *any* no-regret online algorithm, after observing  $N$  samples, can find a policy  $\hat{\pi}_N$  in a policy class  $\Pi$  satisfying  $V^{\hat{\pi}_N}(p) \geq V^{\pi^*}(p) - o(1) - \epsilon_\Pi$ , where  $V^\pi(p)$  is the accumulated reward of policy  $\pi$  with respect to some unknown initial state distribution  $p$ ,  $\pi^*$  is the optimal policy, and  $\epsilon_\Pi \geq 0$  is a measure of the expressivity of  $\Pi$  (see Section 4.2 for definition).

Our reduction is built on a refinement of online learning, called Continuous Online Learning (COL), which was proposed to model problems where loss gradients across rounds change continuously with the learner’s decisions [14]. COL has a strong connection to equilibrium problems (EPs) [15, 16], and any monotone EP (including our saddle-point problem of interest) can be framed as no-regret learning in a properly constructed COL problem [14]. Using this idea, our reduction follows naturally by first converting an RL problem to an EP and then the EP to a COL problem.

Framing RL as COL reveals new insights into the relationship between approximate solutions to the saddle-point problem and approximately optimal policies. Importantly, this new perspective shows that the RL problem can be separated into two parts: regret minimization and function approximation. The first part admits standard treatments from the online learning

literature, and the second part can be quantified *independently* of the learning process. For example, one can accelerate learning by adopting optimistic online algorithms [17, 18] that account for the predictability in COL, without worrying about function approximators. Because of these problem-agnostic features, the proposed reduction can be used to systematically design efficient RL algorithms with performance guarantees.

As a demonstration, we design an RL algorithm based on arguably the simplest online learning algorithm: mirror descent. Assuming a generative model<sup>1</sup>, we prove that, for *any* tabular Markov decision process (MDP), with probability at least  $1 - \delta$ , this algorithm learns an  $\epsilon$ -optimal policy for the  $\gamma$ -discounted accumulated reward, using at most  $\tilde{O}\left(\frac{|\mathcal{S}||\mathcal{A}|\log(\frac{1}{\delta})}{(1-\gamma)^4\epsilon^2}\right)$  samples, where  $|\mathcal{S}|, |\mathcal{A}|$  are the sizes of state and action spaces, and  $\gamma$  is the discount rate. Furthermore, thanks to the separation property above, our algorithm admits a natural extension with linearly parameterized function approximators, whose sample and per-round computation complexities are *linear* in the number of parameters and independent of  $|\mathcal{S}|, |\mathcal{A}|$ , though at the cost of policy performance bias due to approximation error.

This sample complexity improves the current best provable rate of the saddle-point RL setup [3–6] by a large factor of  $\frac{|\mathcal{S}|^2}{(1-\gamma)^2}$ , *without* making any assumption on the MDP.<sup>2</sup> This improvement is attributed to our new online-learning-style analysis that uses a cleverly selected comparator in the regret definition. While it is possible to devise a minor modification of the previous stochastic mirror descent algorithms, e.g. [5], achieving the same rate with our new analysis, we remark that our algorithm is considerably simpler and removes a projection required in previous work [3–6].

Finally, we do note that the same sample complexity can also be achieved, e.g., by model-based RL and (phased) Q-learning [19, 20]. However, these methods either have super-linear runtime, with no obvious route for improvement, or could become unstable when using function approximators without further assumption.

## 2 SETUP & PRELIMINARIES

Let  $\mathcal{S}$  and  $\mathcal{A}$  be state and action spaces, which can be discrete or continuous. We consider  $\gamma$ -discounted infinite-horizon problems for  $\gamma \in [0, 1)$ . Our goal is to find a policy  $\pi(a|s)$  that maximizes the discounted average return  $V^\pi(p) := \mathbb{E}_{s \sim p}[V^\pi(s)]$ , where  $p$  is the

<sup>1</sup>In practice, it can be approximated by running a behavior policy with sufficient exploration [19].

<sup>2</sup>[5] has the same sample complexity but requires the MDP to be ergodic under any policy.

initial state distribution,

$$V^\pi(s) := (1 - \gamma)\mathbb{E}_{\xi \sim \rho_\pi(s)}\left[\sum_{t=0}^{\infty} \gamma^t r(s_t, a_t)\right] \quad (1)$$

is the value function of  $\pi$  at state  $s$ ,  $r : \mathcal{S} \times \mathcal{A} \rightarrow [0, 1]$  is the reward function, and  $\rho_\pi(s)$  is the distribution of trajectory  $\xi = s_0, a_0, s_1, \dots$  generated by running  $\pi$  from  $s_0 = s$  in an MDP. We assume that the initial distribution  $p(s_0)$ , the transition  $\mathcal{P}(s'|s, a)$ , and the reward function  $r(s, a)$  in the MDP are unknown but can be queried through a *generative model*, i.e. we can sample  $s_0$  from  $p$ ,  $s'$  from  $\mathcal{P}$ , and  $r(s, a)$  for any  $s \in \mathcal{S}$  and  $a \in \mathcal{A}$ . We remark that the definition of  $V^\pi$  in (1) contains a  $(1 - \gamma)$  factor. We adopt this setup to make writing more compact. We denote the optimal policy as  $\pi^*$  and its value function as  $V^*$  for short.

### 2.1 Duality in RL

Our reduction is based on the linear-program (LP) formulation of RL. We provide a short recap here (please see Appendix A and [21] for details).

To show how  $\max_\pi V^\pi(p)$  can be framed as a LP, let us define the average state distribution under  $\pi$ ,  $d^\pi(s) := (1 - \gamma)\sum_{t=0}^{\infty} \gamma^t d_t^\pi(s)$ , where  $d_t^\pi$  is the state distribution at time  $t$  visited by running  $\pi$  from  $p$  (e.g.  $d_0^\pi = p$ ). By construction,  $d^\pi$  satisfies the stationarity property,

$$d^\pi(s') = (1 - \gamma)p(s') + \gamma\mathbb{E}_{s \sim d^\pi}\mathbb{E}_{a \sim \pi|s}[\mathcal{P}(s'|s, a)]. \quad (2)$$

With  $d^\pi$ , we can write  $V^\pi(p) = \mathbb{E}_{s \sim d^\pi}\mathbb{E}_{a \sim \pi|s}[r(s, a)]$  and our objective  $\max_\pi V^\pi(p)$  equivalently as:

$$\begin{aligned} \max_{\boldsymbol{\mu} \in \mathbb{R}^{|\mathcal{S}||\mathcal{A}|}: \boldsymbol{\mu} \geq \mathbf{0}} \quad & \mathbf{r}^\top \boldsymbol{\mu} \\ \text{s.t.} \quad & (1 - \gamma)\mathbf{p} + \gamma\mathbf{P}^\top \boldsymbol{\mu} = \mathbf{E}^\top \boldsymbol{\mu} \end{aligned} \quad (3)$$

where  $\mathbf{r} \in \mathbb{R}^{|\mathcal{S}||\mathcal{A}|}$ ,  $\mathbf{p} \in \mathbb{R}^{|\mathcal{S}|}$ , and  $\mathbf{P} \in \mathbb{R}^{|\mathcal{S}||\mathcal{A}| \times |\mathcal{S}|}$  are vector forms of  $r$ ,  $p$ , and  $\mathcal{P}$ , respectively, and  $\mathbf{E} = \mathbf{I} \otimes \mathbf{1} \in \mathbb{R}^{|\mathcal{S}||\mathcal{A}| \times |\mathcal{S}|}$  (we use  $|\cdot|$  to denote the cardinality of a set,  $\otimes$  the Kronecker product,  $\mathbf{I} \in \mathbb{R}^{|\mathcal{S}| \times |\mathcal{S}|}$  is the identity, and  $\mathbf{1} \in \mathbb{R}^{|\mathcal{A}|}$  the vector of ones). In (3),  $\mathcal{S}$  and  $\mathcal{A}$  may seem to have finite cardinalities, but the same formulation extends to countable or even continuous spaces (under proper regularity assumptions; see [22]). We adopt this abuse of notation (emphasized by bold-faced symbols) for compactness.

The variable  $\boldsymbol{\mu}$  of the LP in (3) resembles a joint distribution  $d^\pi(s)\pi(a|s)$ . To see this, notice that the constraint in (3) is reminiscent of (2), and implies  $\|\boldsymbol{\mu}\|_1 = 1$ , i.e.  $\boldsymbol{\mu}$  is a probability distribution. Then one can show  $\mu(s, a) = d^\pi(s)\pi(a|s)$  when the constraint is satisfied, which implies that (3) is the same as  $\max_\pi V^\pi(p)$  and its solution  $\boldsymbol{\mu}^*$  corresponds to  $\mu^*(s, a) = d^{\pi^*}(s)\pi^*(a|s)$  of the optimal policy  $\pi^*$ .

As (3) is a LP, it suggests looking at its dual, which turns out to be the classic LP formulation of RL<sup>3</sup>,

$$\begin{aligned} \min_{\mathbf{v} \in \mathbb{R}^{|\mathcal{S}|}} \quad & \mathbf{p}^\top \mathbf{v} \\ \text{s.t.} \quad & (1 - \gamma)\mathbf{r} + \gamma \mathbf{P}\mathbf{v} \leq \mathbf{E}\mathbf{v}. \end{aligned} \quad (4)$$

It can be verified that for all  $\mathbf{p} > 0$ , the solution to (4) satisfies the Bellman equation [23] and therefore is the optimal value function  $\mathbf{v}^*$  (the vector form of  $V^*$ ). We note that, for any policy  $\pi$ ,  $V^\pi$  by definition satisfies a stationarity property

$$V^\pi(s) = \mathbb{E}_{a \sim \pi|s} [(1 - \gamma)r(s, a) + \gamma \mathbb{E}_{s' \sim \mathcal{P}|s, a} [V^\pi(s')]] \quad (5)$$

which can be viewed as a dual equivalent of (2) for  $d^\pi$ . Because, for any  $s \in \mathcal{S}$  and  $a \in \mathcal{A}$ ,  $r(s, a)$  is in  $[0, 1]$ , (5) implies  $V^\pi(s)$  lies in  $[0, 1]$  too.

## 2.2 Toward RL: the Saddle-Point Setup

The LP formulations above require knowing the probabilities  $p$  and  $\mathcal{P}$  and are computationally inefficient. When only generative models are available (as in our setup), one can alternatively exploit the duality relationship between the two LPs in (3) and (4), and frame RL as a saddle-point problem [3]. Let us define

$$\mathbf{a}_\mathbf{v} := \mathbf{r} + \frac{1}{1-\gamma}(\gamma \mathbf{P} - \mathbf{E})\mathbf{v} \quad (6)$$

as the *advantage function* with respect to  $\mathbf{v}$  (where  $\mathbf{v}$  is not necessarily a value function). Then the Lagrangian connecting the two LPs can be written as

$$\mathcal{L}(\mathbf{v}, \boldsymbol{\mu}) := \mathbf{p}^\top \mathbf{v} + \boldsymbol{\mu}^\top \mathbf{a}_\mathbf{v}, \quad (7)$$

which leads to the saddle-point formulation,

$$\min_{\mathbf{v} \in \mathcal{V}} \max_{\boldsymbol{\mu} \in \mathcal{M}} \mathcal{L}(\mathbf{v}, \boldsymbol{\mu}), \quad (8)$$

where the constraints are

$$\mathcal{V} = \{\mathbf{v} \in \mathbb{R}^{|\mathcal{S}|} : \mathbf{v} \geq 0, \|\mathbf{v}\|_\infty \leq 1\} \quad (9)$$

$$\mathcal{M} = \{\boldsymbol{\mu} \in \mathbb{R}^{|\mathcal{S}||\mathcal{A}|} : \boldsymbol{\mu} \geq 0, \|\boldsymbol{\mu}\|_1 = 1\}. \quad (10)$$

The solution to (8) is exactly  $(\mathbf{v}^*, \boldsymbol{\mu}^*)$ , but notice that extra constraints on the norm of  $\boldsymbol{\mu}$  and  $\mathbf{v}$  are introduced in  $\mathcal{V}, \mathcal{M}$ , compared with (3) and (4). This is a common practice, which uses known bound on the solutions of (3) and (4) (discussed above) to make the search spaces  $\mathcal{V}$  and  $\mathcal{M}$  in (8) compact and as small as possible so that optimization converges faster.

Having compact variable sets allows using first-order stochastic methods, such as stochastic mirror descent

<sup>3</sup>Our setup in (4) differs from the classic one in the  $(1 - \gamma)$  factor in the constraint due to the average setup.

and mirror-prox [24, 25], to efficiently solve the problem. These methods only require using the generative model to compute unbiased estimates of the gradients  $\nabla_{\mathbf{v}} \mathcal{L} = \mathbf{b}_\boldsymbol{\mu}$  and  $\nabla_{\boldsymbol{\mu}} \mathcal{L} = \mathbf{a}_\mathbf{v}$ , where we define

$$\mathbf{b}_\boldsymbol{\mu} := \mathbf{p} + \frac{1}{1-\gamma}(\gamma \mathbf{P} - \mathbf{E})^\top \boldsymbol{\mu} \quad (11)$$

as the *balance function* with respect to  $\boldsymbol{\mu}$ .  $\mathbf{b}_\boldsymbol{\mu}$  measures whether  $\boldsymbol{\mu}$  violates the stationarity constraint in (3) and can be viewed as the dual of  $\mathbf{a}_\mathbf{v}$ . When the state or action space is too large, one can resort to function approximators to represent  $\mathbf{v}$  and  $\boldsymbol{\mu}$ , which are often realized by linear basis functions for the sake of analysis [10].

## 2.3 COL and EPs

Finally, we review the COL setup in [14], which we will use to design the reduction from the saddle-point problem in (8) to online learning in the next section.

Recall that an online learning problem describes the iterative interactions between a learner and an opponent. In round  $n$ , the learner chooses a decision  $x_n$  from a decision set  $\mathcal{X}$ , the opponent chooses a per-round loss function  $l_n : \mathcal{X} \rightarrow \mathbb{R}$  based on the learner's decisions, and then information about  $l_n$  (e.g. its gradient  $\nabla l_n(x_n)$ ) is revealed to the learner. The performance of the learner is usually measured in terms of regret with respect to some  $x' \in \mathcal{X}$ ,

$$\text{Regret}_N(x') := \sum_{n=1}^N l_n(x_n) - \sum_{n=1}^N l_n(x').$$

When  $l_n$  is convex and  $\mathcal{X}$  is compact and convex, many no-regret (i.e.  $\text{Regret}_N(x') = o(N)$ ) algorithms are available, such as mirror descent and follow-the-regularized-leader [26–28].

COL is a subclass of online learning problems where the loss sequence changes continuously with respect to the played decisions of the learner [14]. In COL, the opponent is equipped with a bifunction  $f : (x, x') \mapsto f_x(x')$ , where any fixed  $x' \in \mathcal{X}$ ,  $\nabla f_x(x')$  is continuous in  $x \in \mathcal{X}$ . The opponent selects per-round losses based on  $f$ , but the learner does not know  $f$ : in round  $n$ , if the learner chooses  $x_n$ , the opponent sets

$$l_n(x) = f_{x_n}(x), \quad (12)$$

and returns, e.g., a stochastic estimate of  $\nabla l_n(x_n)$  (the regret is still measured in terms of the noise-free  $l_n$ ).

In [14], a natural connection is shown between COL and equilibrium problems (EPs). As EPs include the saddle-point problem of interest, we can use this idea to turn (8) into a COL problem. Recall an EP is defined as follows: Let  $\mathcal{X}$  be compact and  $F : (x, x') \mapsto F(x, x')$  be a bifunction s.t.  $\forall x, x' \in \mathcal{X}$ ,  $F(\cdot, x')$  is

continuous,  $F(x, \cdot)$  is convex, and  $F(x, x) \geq 0$ .<sup>4</sup> The problem  $\text{EP}(\mathcal{X}, F)$  aims to find  $x^* \in \mathcal{X}$  s.t.

$$F(x^*, x) \geq 0, \quad \forall x \in \mathcal{X}. \quad (13)$$

By its definition, a natural residual function to quantify the quality of an approximation solution  $x$  to EP is  $r_{ep}(x) := -\min_{x' \in \mathcal{X}} F(x, x')$  which describes the degree to which (13) is violated at  $x$ . We say a bifunction  $F$  is *monotone* if,  $\forall x, x' \in \mathcal{X}$ ,  $F(x, x') + F(x', x) \leq 0$ , and *skew-symmetric* if the equality holds.

EPs with monotone bifunctions represent general convex problems, including convex optimization problems, saddle-point problems, variational inequalities, etc. For instance, a convex-concave problem  $\min_{y \in \mathcal{Y}} \max_{z \in \mathcal{Z}} \phi(y, z)$  can be cast as  $\text{EP}(\mathcal{X}, F)$  with  $\mathcal{X} = \mathcal{Y} \times \mathcal{Z}$  and the skew-symmetric bifunction [29]

$$F(x, x') := \phi(y', z) - \phi(y, z'), \quad (14)$$

where  $x = (y, z)$  and  $x' = (y', z')$ . In this case,  $r_{ep}(x) = \max_{z' \in \mathcal{Z}} \phi(y, z') - \min_{y' \in \mathcal{Y}} \phi(y', z)$  is the duality gap.

Cheng et al. [14] show that a learner achieves sublinear dynamic regret in COL if and only if the same algorithm can solve  $\text{EP}(\mathcal{X}, F)$  with  $F(x, x') = f_x(x') - f_x(x)$ . Concretely, they show that, given a monotone  $\text{EP}(\mathcal{X}, F)$  with  $F(x, x) = 0$  (which is satisfied by (14)), one can construct a COL problem by setting  $f_{x'}(x) := F(x', x)$ , i.e.  $l_n(x) = F(x_n, x)$ , such that any no-regret algorithm can generate an approximate solution to the EP.

**Proposition 1.** [14] *If  $F$  is skew-symmetric and  $l_n(x) = F(x_n, x)$ , then  $r_{ep}(\hat{x}_N) \leq \frac{1}{N} \text{Regret}_N$ , where  $\text{Regret}_N = \max_{x \in \mathcal{X}} \text{Regret}_N(x)$ , and  $\hat{x}_N = \frac{1}{N} \sum_{n=1}^N x_n$ ; the same guarantee holds also for the best decision in  $\{x_n\}_{n=1}^N$ .*

### 3 AN ONLINE LEARNING VIEW

We present an alternate online-learning perspective on the saddle-point formulation in (8). This analysis paves a way for our reduction in the next section. By reduction, we mean realizing the two steps below:

1. Define a sequence of online losses such that any algorithm with sublinear regret can produce an approximate solution to the saddle-point problem.
2. Convert the approximate solution in the first step to an approximately optimal policy in RL.

<sup>4</sup>We restrict ourselves to this convex and continuous case as it is sufficient for our problem setup.

Methods to achieve these two steps individually are not new. The reduction from convex-concave problems to no-regret online learning is well known [30]. Likewise, the relationship between the approximate solution of (8) and policy performance is also available; this is how the saddle-point formulation [5] works in the first place. So couldn't we just use these existing approaches? We argue that purely combining these two techniques fails to fully capture important structure that resides in RL. While this will be made precise in the later analyses, we highlight the main insights here.

Instead of treating (8) as an *adversarial* two-player online learning problem [30], we adopt the recent reduction to COL [14] reviewed in Section 2.3. The main difference is that the COL approach takes a single-player setup and retains the Lipschitz continuity in the source saddle-point problem. This single-player perspective is in some sense cleaner and, as we will show in Section 4.2, provides a simple setup to analyze effects of function approximators. Additionally, due to continuity, the losses in COL are predictable and therefore make designing fast algorithms possible.

With the help of the COL reformulation, we study the relationship between the approximate solution to (8) and the performance of the associated policy in RL. We are able to establish a tight bound between the residual and the performance gap, resulting in a large improvement of  $\frac{|S|^2}{(1-\gamma)^2}$  in sample complexity compared with the best bounds in the literature of the saddle-point setup, *without* adding extra constraints on  $\mathcal{X}$  and assumptions on the MDP. Overall, this means that *stronger* sample complexity guarantees can be attained by *simpler* algorithms, as we demonstrate in Section 5.

The missing proofs of this section are in Appendix B.

#### 3.1 The COL Formulation of RL

First, let us exercise the above COL idea with the saddle-point formulation of RL in (8). To construct the EP, we can let  $\mathcal{X} = \{x = (\mathbf{v}, \boldsymbol{\mu}) : \mathbf{v} \in \mathcal{V}, \boldsymbol{\mu} \in \mathcal{M}\}$ , which is compact. According to (14), the bifunction  $F$  of the associated  $\text{EP}(\mathcal{X}, F)$  is naturally given as

$$\begin{aligned} F(x, x') &:= \mathcal{L}(\mathbf{v}', \boldsymbol{\mu}) - \mathcal{L}(\mathbf{v}, \boldsymbol{\mu}') \\ &= \mathbf{p}^\top \mathbf{v}' + \boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}'} - \mathbf{p}^\top \mathbf{v} - \boldsymbol{\mu}'^\top \mathbf{a}_{\mathbf{v}} \end{aligned} \quad (15)$$

which is skew-symmetric, and  $x^* := (\mathbf{v}^*, \boldsymbol{\mu}^*)$  is a solution to  $\text{EP}(\mathcal{X}, F)$ . This identification gives us a COL problem with the loss in the  $n$ th round defined as

$$l_n(x) := \mathbf{p}^\top \mathbf{v} + \boldsymbol{\mu}_n^\top \mathbf{a}_{\mathbf{v}} - \mathbf{p}^\top \mathbf{v}_n - \boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}_n} \quad (16)$$

where  $x_n = (\mathbf{v}_n, \boldsymbol{\mu}_n)$ . We see  $l_n$  is a linear loss. Moreover, because of the continuity in  $\mathcal{L}$ , it is predictable, i.e.  $l_n$  can be (partially) inferred from past feedback as the MDP involved in each round is the same.

### 3.2 Policy Performance and Residual

By Proposition 1, any no-regret algorithm, when applied to (16), provides guarantees in terms of the residual function  $r_{ep}(x)$  of the EP. But this is not the end of the story. We also need to relate the learner decision  $x \in \mathcal{X}$  to a policy  $\pi$  in RL and then convert bounds on  $r_{ep}(x)$  back to the policy performance  $V^\pi(p)$ . Here we follow the common rule in the literature and associate each  $x = (\mathbf{v}, \boldsymbol{\mu}) \in \mathcal{X}$  with a policy  $\pi_\mu$  defined as

$$\pi_\mu(a|s) \propto \mu(s, a). \quad (17)$$

In the following, we relate the residual  $r_{ep}(x)$  to the performance gap  $V^*(p) - V^{\pi_\mu}(p)$  through a relative performance measure defined as

$$r_{ep}(x; x') := F(x, x) - F(x, x') = -F(x, x') \quad (18)$$

for  $x, x' \in \mathcal{X}$ , where the last equality follows from the skew-symmetry of  $F$  in (15). Intuitively, we can view  $r_{ep}(x; x')$  as comparing the performance of  $x$  with respect to the comparator  $x'$  under an optimization problem proposed by  $x$ , e.g. we have  $l_n(x_n) - l_n(x') = r_{ep}(x_n; x')$ . And by the definition in (18), it holds that  $r_{ep}(x; x') \leq \max_{x' \in \mathcal{X}} -F(x, x') = r_{ep}(x)$ .

We are looking for inequalities in the form  $V^*(p) - V^{\pi_\mu}(p) \leq \kappa(r_{ep}(x; x'))$  that hold for *all*  $x \in \mathcal{X}$  with some strictly increasing function  $\kappa$  and some  $x' \in \mathcal{X}$ , so we can get *non-asymptotic* performance guarantees once we combine the two steps described at the beginning of this section. For example, by directly applying results of [14] to the COL in (16), we get  $V^*(p) - V^{\hat{\pi}_N}(p) \leq \kappa(\frac{\text{Regret}_N}{N})$ , where  $\hat{\pi}_N$  is the policy associated with the average/best decision in  $\{x_n\}_{1=N}$ .

#### 3.2.1 The Classic Result

Existing approaches (e.g. [4–6]) to the saddle-point point formulation in (8) rely on the relative residual  $r_{ep}(x; x^*)$  with respect to the optimal solution to the problem  $x^*$ , which we restate in our notation.

**Proposition 2.** *For any  $x = (\mathbf{v}, \boldsymbol{\mu}) \in \mathcal{X}$ , if  $\mathbf{E}^\top \boldsymbol{\mu} \geq (1 - \gamma)\mathbf{p}$ ,  $r_{ep}(x; x^*) \geq (1 - \gamma) \min_s p(s) \|\mathbf{v}^* - \mathbf{v}^{\pi_\mu}\|_\infty$ .*

Therefore, although the original saddle-point problem in (8) is framed using  $\mathcal{V}$  and  $\mathcal{M}$ , in practice, an extra constraint, such as  $\mathbf{E}^\top \boldsymbol{\mu} \geq (1 - \gamma)\mathbf{p}$ , is added into  $\mathcal{M}$ , i.e. these algorithms consider instead

$$\mathcal{M}' = \{\boldsymbol{\mu} \in \mathbb{R}^{|\mathcal{S}||\mathcal{A}|} : \boldsymbol{\mu} \in \mathcal{M}, \mathbf{E}^\top \boldsymbol{\mu} \geq (1 - \gamma)\mathbf{p}\}, \quad (19)$$

so that the marginal of the estimate  $\boldsymbol{\mu}$  can have the sufficient coverage required in Proposition 2. This condition is needed to establish non-asymptotic guarantees on the performance of the policy generated by  $\boldsymbol{\mu}$  [3, 5, 6], but it can sometimes be impractical to

realize, e.g., when  $\mathbf{p}$  is unknown. Without it, extra assumptions (like ergodicity [5]) on the MDP are needed.

However, Proposition 2 is undesirable for a number of reasons. First, the bound is quite conservative, as it concerns the uniform error  $\|\mathbf{v}^* - \mathbf{v}^{\pi_\mu}\|_\infty$  whereas the objective in RL is about the gap  $V^*(p) - V^{\pi_\mu}(p) = \mathbf{p}^\top (\mathbf{v}^* - \mathbf{v}^{\pi_\mu})$  with respect to the initial distribution  $p$  (i.e. a weighted error). Second, the constant term  $(1 - \gamma) \min_s p(s)$  can be quite small (e.g. when  $p$  is uniform, it is  $\frac{1-\gamma}{|\mathcal{S}|}$ ) which can significantly amplify the error in the residual. Because a no-regret algorithm typically decreases the residual in  $O(N^{-1/2})$  after seeing  $N$  samples, the factor of  $\frac{1-\gamma}{|\mathcal{S}|}$  earlier would turn into a multiplier of  $\frac{|\mathcal{S}|^2}{(1-\gamma)^2}$  in sample complexity. This makes existing saddle-point approaches sample inefficient in comparison with other RL methods like Q-learning [20]. Lastly, enforcing  $\mathbf{E}^\top \boldsymbol{\mu} \geq (1 - \gamma)\mathbf{p}$  requires knowing  $\mathbf{p}$  (which is unavailable in our setup) and adds extra projection steps during optimization. When  $\mathbf{p}$  is unknown, while it is possible to modify this constraint to use a uniform distribution, this might worsen the constant factor and could introduce bias.

One may conjecture that the bound in Proposition 2 could perhaps be tightened by better analyses. However, we prove this is impossible in general.

**Proposition 3.** *There is a class of MDPs such that, for some  $x \in \mathcal{X}$ , Proposition 2 is an equality.*

We note that Proposition 3 does not hold for *all* MDPs. Indeed, if one makes stronger assumptions on the MDP, such as that the Markov chain induced by *every* policy is ergodic [5], then it is possible to show, for all  $x \in \mathcal{X}$ ,  $r_{ep}(x; x^*) = c \|\mathbf{v}^* - \mathbf{v}^{\pi_\mu}\|_\infty$  for some constant  $c$  independent of  $\gamma$  and  $|\mathcal{S}|$ , when one constrains  $\mathbf{E}^\top \boldsymbol{\mu} \geq (1 - \gamma + \gamma\sqrt{c})\mathbf{p}$ . Nonetheless, this construct still requires adding an undesirable constraint to  $\mathcal{X}$ .

#### 3.2.2 Curse of Covariate Shift

Why does this happen? We can view this issue as a form of *covariate shift*, i.e. a mismatch between distributions. To better understand it, we notice a simple equality, which has often been used implicitly, e.g. in the technical proofs of [5].

**Lemma 1.** *For any  $x = (\mathbf{v}, \boldsymbol{\mu})$ , if  $x' \in \mathcal{X}$  satisfies (2) and (5) (i.e.  $\mathbf{v}'$  and  $\boldsymbol{\mu}'$  are the value function and state-action distribution of policy  $\pi_{\boldsymbol{\mu}'}$ ),  $r_{ep}(x; x') = -\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}'}$ .*

Lemma 1 implies  $r_{ep}(x; x^*) = -\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}^*}$ , which is non-negative. This term is similar to an equality called the performance difference lemma [31, 32].

**Lemma 2.** *Let  $\mathbf{v}^\pi$  and  $\boldsymbol{\mu}^\pi$  denote the value and state-action distribution of some policy  $\pi$ . Then for any function  $\mathbf{v}'$ , it holds that  $\mathbf{p}^\top (\mathbf{v}^\pi - \mathbf{v}') = (\boldsymbol{\mu}^\pi)^\top \mathbf{a}_{\mathbf{v}'}$ . In*

particular, it implies  $V^\pi(p) - V^{\pi'}(p) = (\boldsymbol{\mu}^\pi)^\top \mathbf{a}_{\mathbf{v}^{\pi'}}$ .

From Lemmas 1 and 2, we see that the difference between the residual  $r_{ep}(x; x^*) = -\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}^*}$  and the performance gap  $V^{\pi\mu}(p) - V^{\pi^*}(p) = (\boldsymbol{\mu}^{\pi\mu})^\top \mathbf{a}_{\mathbf{v}^*}$  is due to the mismatch between  $\boldsymbol{\mu}$  and  $\boldsymbol{\mu}^{\pi\mu}$ , or more specifically, the mismatch between the two marginals  $\mathbf{d} = \mathbf{E}^\top \boldsymbol{\mu}$  and  $\mathbf{d}^{\pi\mu} = \mathbf{E}^\top \boldsymbol{\mu}^{\pi\mu}$ . Indeed, when  $\mathbf{d} = \mathbf{d}^{\pi\mu}$ , the residual is equal to the performance gap. However, in general, we do not have control over that difference for the sequence of variables  $\{x_n = (\mathbf{v}_n, \boldsymbol{\mu}_n) \in \mathcal{X}\}$  an algorithm generates. The sufficient condition in Proposition 2 attempts to mitigate the difference, using the fact  $\mathbf{d}^{\pi\mu} = (1 - \gamma)\mathbf{p} + \gamma\mathbf{P}_{\pi\mu}^\top \mathbf{d}^{\pi\mu}$  from (2), where  $\mathbf{P}_{\pi\mu}$  is the transition matrix under  $\pi\mu$ . But the missing half  $\gamma\mathbf{P}_{\pi\mu}^\top \mathbf{d}^{\pi\mu}$  (due to the long-term effects in the MDP) introduces the unavoidable, weak constant  $(1 - \gamma)\min_s p(s)$ , if we want to have a uniform bound on  $\|\mathbf{v}^* - \mathbf{v}^{\pi\mu}\|_\infty$ . The counterexample in Proposition 3 was designed to maximize the effect of covariate shift, so that  $\boldsymbol{\mu}$  fails to capture state-action pairs with high advantage. To break the curse, we must properly weight the gap between  $\mathbf{v}^*$  and  $\mathbf{v}^{\pi\mu}$  instead of relying on the uniform bound on  $\|\mathbf{v}^* - \mathbf{v}^{\pi\mu}\|_\infty$  as before.

## 4 THE REDUCTION

The analyses above reveal both good and bad properties of the saddle-point setup in (8). On the one hand, we showed that approximate solutions to the saddle-point problem in (8) can be obtained by running any no-regret algorithm in the single-player COL problem defined in (16); many efficient algorithms are available from the online learning literature. On the other hand, we also discovered a root difficulty in converting an approximate solution of (8) to an approximately optimal policy in RL (Proposition 2), even after imposing strong conditions like (19). At this point, one may wonder if the formulation based on (8) is fundamentally sample inefficient compared with other approaches to RL, but this is actually not true.

Our main contribution shows that learning a policy through running a no-regret algorithm in the COL problem in (16) is, in fact, as sample efficient in policy performance as other RL techniques, even without the common constraint in (19) or extra assumptions on the MDP like ergodicity imposed in the literature.

**Theorem 1.** *Let  $X_N = \{x_n \in \mathcal{X}\}_{n=1}^N$  be any sequence. Let  $\hat{\pi}_N$  be the policy given by  $\hat{x}_N$  via (17), which is either the average or the best decision in  $X_N$ . Define  $y_N^* := (\mathbf{v}^{\hat{\pi}_N}, \boldsymbol{\mu}^*)$ . Then  $V^{\hat{\pi}_N}(p) \geq V^*(p) - \frac{\text{Regret}_N(y_N^*)}{N}$ .*

Theorem 1 shows that if  $X_N$  has sublinear regret, then

both the average policy and the best policy in  $X_N$  converge to the optimal policy in performance with a rate  $O(\text{Regret}_N(y_N^*)/N)$ . Compared with existing results obtained through Proposition 2, the above result removes the factor  $(1 - \gamma)\min_s p(s)$  and impose *no* assumption on  $X_N$  or the MDP. Indeed Theorem 1 holds for *any* sequence. For example, when  $X_N$  is generated by stochastic feedback of  $l_n$ , Theorem 1 continues to hold, as the regret is defined in terms of  $l_n$ , not of the sampled loss. Stochasticity only affects the regret rate.

In other words, we have shown that when  $\boldsymbol{\mu}$  and  $\mathbf{v}$  can be directly parameterized, an approximately optimal policy for the RL problem can be obtained by running any no-regret online learning algorithm, and that the policy quality is simply dictated by the regret rate. To illustrate, in Section 5 we will prove that simply running mirror descent in this COL produces an RL algorithm that is as sample efficient as other common RL techniques. One can further foresee that algorithms leveraging the continuity in COL—e.g. mirror-prox [25] or PicCoLO [18]—and variance reduction can lead to more sample efficient RL algorithms.

Below we will also demonstrate how to use the fact that COL is *single-player* (see Section 2.3) to cleanly incorporate the effects of using function approximators to model  $\boldsymbol{\mu}$  and  $\mathbf{v}$ . We will present a corollary of Theorem 1, which separates the problem of *learning*  $\boldsymbol{\mu}$  and  $\mathbf{v}$ , and that of *approximating*  $\mathcal{M}$  and  $\mathcal{V}$  with function approximators. The first part is controlled by the rate of regret in online learning, and the second part depends on only the chosen class of function approximators, independently of the learning process. As these properties are agnostic to problem setups and algorithms, our reduction leads to a framework for systematic synthesis of new RL algorithms with performance guarantees. The missing proofs of this section are in Appendix C.

### 4.1 Proof of Theorem 1

The main insight of our reduction is to adopt, in defining  $r_{ep}(x; x')$ , a comparator  $x' \in \mathcal{X}$  based on the output of the algorithm (represented by  $x$ ), instead of the fixed comparator  $x^*$  (the optimal pair of value function and state-action distribution) that has been used conventionally, e.g. in Proposition 2. While this idea seems unnatural from the standard saddle-point or EP perspective, it is possible, because the regret in online learning is measured against the worst-case choice in  $\mathcal{X}$ , which is allowed to be selected in *hindsight*. Specifically, we propose to select the following comparator to directly bound  $V^*(p) - V^{\hat{\pi}_N}(p)$  instead of the conservative measure  $\|V^* - V^{\hat{\pi}_N}\|_\infty$  used before.

**Proposition 4.** *For  $x = (\mathbf{v}, \boldsymbol{\mu}) \in \mathcal{X}$ , define  $y_x^* :=$*

$(\mathbf{v}^{\pi^\mu}, \boldsymbol{\mu}^*) \in \mathcal{X}$ . It holds  $r_{ep}(x; y_x^*) = V^*(p) - V^{\pi^\mu}(p)$ .

To finish the proof, let  $\hat{x}_N$  be either  $\frac{1}{N} \sum_{n=1}^N x_n$  or  $\arg \min_{x \in X_N} r_{ep}(x; y_x^*)$ , and let  $\hat{\pi}_N$  denote the policy given by (17). First,  $V^*(p) - V^{\hat{\pi}_N}(p) = r_{ep}(\hat{x}_N; y_N^*)$  by Proposition 4. Next we follow the proof idea of Proposition 1 in [14]: because  $F$  is skew-symmetric and  $F(y_N^*, \cdot)$  is convex, we have by (18)

$$\begin{aligned} V^*(p) - V^{\hat{\pi}_N}(p) &= r_{ep}(\hat{x}_N; y_N^*) = -F(\hat{x}_N, y_N^*) \\ &= F(y_N^*, \hat{x}_N) \leq \frac{1}{N} \sum_{n=1}^N F(y_N^*, x_n) \\ &= \frac{1}{N} \sum_{n=1}^N -F(x_n, y_N^*) = \frac{1}{N} \text{Regret}_N(y_N^*). \end{aligned}$$

## 4.2 Function Approximators

When the state and action spaces are large or continuous, directly optimizing  $\mathbf{v}$  and  $\boldsymbol{\mu}$  can be impractical. Instead we can consider optimizing over a subset of feasible choices parameterized by function approximators

$$\mathcal{X}_\Theta = \{\mathbf{x}_\theta = (\phi_\theta, \psi_\theta) : \psi_\theta \in \mathcal{M}, \theta \in \Theta\}, \quad (20)$$

where  $\phi_\theta$  and  $\psi_\theta$  are functions parameterized by  $\theta \in \Theta$ , and  $\Theta$  is a parameter set. Because COL is a single-player setup, we can extend the previous idea and Theorem 1 to provide performance bounds in this case by a simple rearrangement (see Appendix C), which is a common trick used in the online imitation learning literature [33–35]. Notice that, in (20), we require only  $\psi_\theta \in \mathcal{M}$ , but not  $\phi_\theta \in \mathcal{V}$ , because for the performance bound in our reduction to hold, we only need the constraint  $\mathcal{M}$  (see Lemma 4 in proof of Proposition 4).

**Corollary 1.** *Let  $X_N = \{x_n \in \mathcal{X}_\Theta\}_{n=1}^N$  be any sequence. Let  $\hat{\pi}_N$  be the policy given either by the average or the best decision in  $X_N$ . It holds that*

$$V^{\hat{\pi}_N}(p) \geq V^*(p) - \frac{\text{Regret}_N(\Theta)}{N} - \epsilon_{\Theta, N}$$

where  $\epsilon_{\Theta, N} = \min_{x_\theta \in \mathcal{X}_\Theta} r_{ep}(\hat{x}_N; y_N^*) - r_{ep}(\hat{x}_N; x_\theta)$  measures the expressiveness of  $X_\Theta$ , and  $\text{Regret}_N(\Theta) := \sum_{n=1}^N l_n(x_n) - \min_{x \in \mathcal{X}_\Theta} \sum_{n=1}^N l_n(x)$ .

We can quantify  $\epsilon_{\Theta, N}$  with the basic Hölder’s inequality.

**Proposition 5.** *Let  $\hat{x}_N = (\hat{\mathbf{v}}_N, \hat{\boldsymbol{\mu}}_N)$ . Under the setup in Corollary 1, regardless of the parameterization, it is true that  $\epsilon_{\Theta, N}$  is no larger than*

$$\begin{aligned} &\min_{(\mathbf{v}_\theta, \boldsymbol{\mu}_\theta) \in \mathcal{X}_\Theta} \frac{\|\boldsymbol{\mu}_\theta - \boldsymbol{\mu}^*\|_1}{1 - \gamma} + \min_{\mathbf{w}: \mathbf{w} \geq 1} \|\mathbf{b} \hat{\boldsymbol{\mu}}_N\|_{1, \mathbf{w}} \|\mathbf{v}_\theta - \mathbf{v}^{\hat{\pi}_N}\|_{\infty, 1/\mathbf{w}} \\ &\leq \min_{(\mathbf{v}_\theta, \boldsymbol{\mu}_\theta) \in \mathcal{X}_\Theta} \frac{1}{1 - \gamma} \left( \|\boldsymbol{\mu}_\theta - \boldsymbol{\mu}^*\|_1 + 2 \|\mathbf{v}_\theta - \mathbf{v}^{\hat{\pi}_N}\|_{\infty} \right). \end{aligned}$$

where the norms are defined as  $\|\mathbf{x}\|_{1, \mathbf{w}} = \sum_i w_i |x_i|$  and  $\|\mathbf{x}\|_{\infty, 1/\mathbf{w}} = \max_i w_i^{-1} |x_i|$ .

---

## Algorithm 1 Mirror descent for RL

---

**Input:**  $\epsilon$  optimality of the  $\gamma$ -average return  
 $\delta$  maximal failure probability  
 generative model of an MDP

**Output:**  $\hat{\pi}_N = \pi^{\hat{\boldsymbol{\mu}}_N}$

- 1:  $x_1 = (\mathbf{v}_1, \boldsymbol{\mu}_1)$  where  $\boldsymbol{\mu}_1$  is uniform and  $\mathbf{v}_1 \in \mathcal{V}$
  - 2: Set  $N = \tilde{\Omega}\left(\frac{|\mathcal{S}||\mathcal{A}| \log(\frac{1}{\delta})}{(1-\gamma)^2 \epsilon^2}\right)$  and  $\eta = (1-\gamma)(|\mathcal{S}||\mathcal{A}|N)^{-1/2}$
  - 3: Set the Bregman divergence as (22)
  - 4: **for**  $n = 1 \dots N - 1$  **do**
  - 5:   Sample  $g_n$  according to (24)
  - 6:   Update to  $x_{n+1}$  according to (21)
  - 7: **end for**
  - 8: Set  $(\hat{\mathbf{v}}_N, \hat{\boldsymbol{\mu}}_N) = \hat{x}_N = \frac{1}{N} \sum_{n=1}^N x_n$
- 

Proposition 5 says  $\epsilon_{\Theta, N}$  depends on how well  $\mathcal{X}_\Theta$  captures the value function of the output policy  $\mathbf{v}^{\hat{\pi}_N}$  and the optimal state-action distribution  $\boldsymbol{\mu}^*$ . We remark that this result is independent of how  $\mathbf{v}^{\hat{\pi}_N}$  is generated. Furthermore, Proposition 5 makes *no* assumption whatsoever on the structure of function approximators. It even allows sharing parameters  $\theta$  between  $\mathbf{v} = \phi_\theta$  and  $\boldsymbol{\mu} = \psi_\theta$ , e.g., they can be a bi-headed neural network, which is common for learning shared feature representations. More precisely, the structure of the function approximator would only affect whether  $l_n((\phi_\theta, \psi_\theta))$  remains a convex function in  $\theta$ , which determines the difficulty of designing algorithms with sublinear regret.

In other words, the proposed COL formulation provides a reduction which dictates the policy performance with two separate factors: 1) the rate of regret  $\text{Regret}_N(\Theta)$  which is controlled by the choice of online learning algorithm; 2) the approximation error  $\epsilon_{\Theta, N}$  which is determined by the choice of function approximators. These two factors can almost be treated independently, except that the choice of function approximators would determine the properties of  $l_n((\phi_\theta, \psi_\theta))$  as a function of  $\theta$ , and the choice of  $\Theta$  needs to ensure (20) is admissible.

## 5 SAMPLE COMPLEXITY OF MIRROR DESCENT

We demonstrate the power of our reduction by applying perhaps the simplest online learning algorithm, mirror descent, to the proposed COL problem in (16) with stochastic feedback (Algorithm 1). For transparency, we discuss the tabular setup. We will show a natural extension to basis functions at the end.

Recall that mirror descent is a first-order algorithm, whose update rule can be written as

$$x_{n+1} = \arg \min_{x \in \mathcal{X}} \langle g_n, x \rangle + \frac{1}{\eta} B_R(x \| x_n) \quad (21)$$

where  $\eta > 0$  is the step size,  $g_n$  is the feedback direction, and  $B_R(x||x') = R(x) - R(x') - \langle \nabla R(x'), x - x' \rangle$  is the Bregman divergence generated by a strictly convex function  $R$ . Based on the geometry of  $\mathcal{X} = \mathcal{V} \times \mathcal{M}$ , we consider a natural Bregman divergence of the form

$$B_R(x'||x) = \frac{1}{2|\mathcal{S}|} \|\mathbf{v}' - \mathbf{v}\|_2^2 + KL(\boldsymbol{\mu}'||\boldsymbol{\mu}) \quad (22)$$

This choice mitigates the effects of dimension (e.g. if we set  $x_1 = (\mathbf{v}_1, \boldsymbol{\mu}_1)$  with  $\boldsymbol{\mu}_1$  being the uniform distribution, it holds  $B_R(x'||x_1) = \tilde{O}(1)$  for any  $x' \in \mathcal{X}$ ).

To define the feedback direction  $g_n$ , we slightly modify the per-round loss  $l_n$  in (16) and consider a new loss

$$h_n(x) := \mathbf{b}_{\boldsymbol{\mu}_n}^\top \mathbf{v} + \boldsymbol{\mu}^\top \left( \frac{1}{1-\gamma} \mathbf{1} - \mathbf{a}_{\mathbf{v}_n} \right) \quad (23)$$

that shifts  $l_n$  by a constant, where  $\mathbf{1}$  is the vector of ones. One can verify that  $l_n(x) - l_n(x') = h_n(x) - h_n(x')$ , for all  $x, x' \in \mathcal{X}$  when  $\boldsymbol{\mu}, \boldsymbol{\mu}'$  in  $x$  and  $x'$  satisfy  $\|\boldsymbol{\mu}\|_1 = \|\boldsymbol{\mu}'\|_1$  (which holds for Algorithm 1). Therefore, using  $h_n$  does not change regret. The reason for using  $h_n$  instead of  $l_n$  is to make  $\nabla_{\boldsymbol{\mu}} h_n((\mathbf{v}, \boldsymbol{\mu}))$  (and its unbiased approximation) a positive vector, so the regret bound can have a better dimension dependency. This is a common trick used in online learning (e.g. EXP3 [36]) for optimizing variables living in a simplex ( $\boldsymbol{\mu}$  here).

We set the first-order feedback  $g_n$  as an unbiased *sampled* estimate of  $\nabla h_n(x_n)$ . In round  $n$ , this is realized by two independent calls of the generative model:

$$g_n = \left[ \begin{array}{c} \tilde{\mathbf{P}}_n + \frac{1}{1-\gamma} (\gamma \tilde{\mathbf{P}}_n - \mathbf{E}_n)^\top \tilde{\boldsymbol{\mu}}_n \\ |\mathcal{S}| |\mathcal{A}| \left( \frac{1}{1-\gamma} \hat{\mathbf{1}}_n - \hat{\mathbf{r}}_n - \frac{1}{1-\gamma} (\gamma \hat{\mathbf{P}}_n - \hat{\mathbf{E}}_n) \mathbf{v}_n \right) \end{array} \right] \quad (24)$$

Let  $g_n = [\mathbf{g}_{n,v}; \mathbf{g}_{n,\mu}]$ . For  $\mathbf{g}_{n,v}$ , we sample  $\mathbf{p}$ , sample  $\boldsymbol{\mu}_n$  to get a state-action pair, and query the transition  $\mathbf{P}$  at the state-action pair sampled from  $\boldsymbol{\mu}_n$ . ( $\tilde{\mathbf{p}}_n$ ,  $\tilde{\mathbf{P}}_n$ , and  $\tilde{\boldsymbol{\mu}}_n$  denote the single-sample estimate of these probabilities.) For  $\mathbf{g}_{n,\mu}$ , we first sample *uniformly* a state-action pair (which explains the factor  $|\mathcal{S}| |\mathcal{A}|$ ), and then query the reward  $\mathbf{r}$  and the transition  $\mathbf{P}$ . ( $\hat{\mathbf{1}}_n$ ,  $\hat{\mathbf{r}}_n$ ,  $\hat{\mathbf{P}}_n$ , and  $\hat{\mathbf{E}}_n$  denote the single-sample estimates.) To emphasize, we use  $\tilde{\cdot}$  and  $\hat{\cdot}$  to distinguish the empirical quantities obtained by these two independent queries. By construction, we have  $\mathbf{g}_{n,\mu} \geq 0$ . It is clear that this direction  $g_n$  is unbiased, i.e.  $\mathbb{E}[g_n] = \nabla h_n(x_n)$ . Moreover, it is extremely sparse and can be computed using  $O(1)$  sample, computational, and memory complexities.

Below we show this algorithm, despite being extremely simple, has strong theoretical guarantees. In other words, we obtain simpler versions of the algorithms proposed in [3, 5, 10] but with improved performance.

**Theorem 2.** *With probability  $1-\delta$ , Algorithm 1 learns an  $\epsilon$ -optimal policy with  $\tilde{O}\left(\frac{|\mathcal{S}| |\mathcal{A}| \log(\frac{1}{\delta})}{(1-\gamma)^2 \epsilon^2}\right)$  samples.*

Note that the above statement makes no assumption on the MDP (except the tabular setup for simplifying analysis). Also, because the definition of value function in (1) is scaled by a factor  $(1-\gamma)$ , the above result translates into a sample complexity in  $\tilde{O}\left(\frac{|\mathcal{S}| |\mathcal{A}| \log(\frac{1}{\delta})}{(1-\gamma)^4 \epsilon^2}\right)$  for the conventional discounted accumulated rewards.

## 5.1 Proof Sketch of Theorem 2

The proof is based on the basic property of mirror descent and martingale concentration. We provide a sketch here; please refer to Appendix D for details. Let  $y_N^* = (\mathbf{v}^{\hat{\pi}_N}, \boldsymbol{\mu}^*)$ . We bound the regret in Theorem 1 by the following rearrangement, where the first equality below is because  $h_n$  is a constant shift from  $l_n$ .

$$\begin{aligned} \text{Regret}_N(y_N^*) &= \sum_{n=1}^N h_n(x_n) - \sum_{n=1}^N h_n(y_N^*) \\ &\leq \left( \sum_{n=1}^N (\nabla h_n(x_n) - g_n)^\top x_n \right) + \left( \max_{x \in \mathcal{X}} \sum_{n=1}^N g_n^\top (x_n - x) \right) \\ &\quad + \left( \sum_{n=1}^N (g_n - \nabla h_n(x_n))^\top y_N^* \right) \end{aligned}$$

We recognize the first term is a martingale, because  $x_n$  does not depend on  $g_n$ . Therefore, we can appeal to a Bernstein-type martingale concentration and prove it is in  $\tilde{O}\left(\frac{\sqrt{N} |\mathcal{S}| |\mathcal{A}| \log(\frac{1}{\delta})}{1-\gamma}\right)$ . For the second term, by treating  $g_n^\top x$  as the per-round loss, we can use standard regret analysis of mirror descent and show a bound in  $\tilde{O}\left(\frac{\sqrt{N} |\mathcal{S}| |\mathcal{A}|}{1-\gamma}\right)$ . For the third term, because  $\mathbf{v}^{\hat{\pi}_N}$  in  $y_N^* = (\mathbf{v}^{\hat{\pi}_N}, \boldsymbol{\mu}^*)$  depends on  $\{g_n\}_{n=1}^N$ , it is *not* a martingale. Nonetheless, we are able to handle it through a union bound and show it is again no more than  $\tilde{O}\left(\frac{\sqrt{N} |\mathcal{S}| |\mathcal{A}| \log(\frac{1}{\delta})}{1-\gamma}\right)$ . Despite the union bound, it does not increase the rate because we only need to handle  $\mathbf{v}^{\hat{\pi}_N}$ , not  $\boldsymbol{\mu}^*$  which induces a martingale. To finish the proof, we substitute this high-probability regret bound into Theorem 1 to obtain the desired claim.

## 5.2 Extension to Function Approximators

The above algorithm assumes the tabular setup for illustration purposes. In Appendix E, we describe a direct extension of Algorithm 1 that uses linearly parameterized function approximators of the form  $x_\theta = (\Phi \boldsymbol{\theta}_v, \Psi \boldsymbol{\theta}_\mu)$ , where columns of bases  $\Phi, \Psi$  belong to  $\mathcal{V}$  and  $\mathcal{M}$ , respectively, and  $(\boldsymbol{\theta}_v, \boldsymbol{\theta}_\mu) \in \Theta$ .

Overall the algorithm stays the same, except the gradient is computed by chain-rule, which can be done in  $O(\dim(\Theta))$  time and space. While this seems worse,



the computational complexity per update actually improves to  $O(\dim(\Theta))$  from the slow  $O(|\mathcal{S}||\mathcal{A}|)$  (required before for the projection in (21)), as now we only optimize in  $\Theta$ . Moreover, we prove that its sample complexity is also better, though at the cost of bias  $\epsilon_{\Theta, N}$  in Corollary 1. Therefore, the algorithm becomes applicable to large-scale or continuous problems.

**Theorem 3.** *Under a proper choice of  $\Theta$  and  $B_R$ , with probability  $1 - \delta$ , Algorithm 1 learns an  $(\epsilon + \epsilon_{\Theta, N})$ -optimal policy with  $\tilde{O}\left(\frac{\dim(\Theta)\log(\frac{1}{\delta})}{(1-\gamma)^2\epsilon^2}\right)$  samples.*

The proof is in Appendix E, and mainly follows Section 5.1. First, we choose some  $\Theta$  to satisfy (20) so we can use Corollary 1 to reduce the problem into regret minimization. To make the sample complexity independent of  $|\mathcal{S}|, |\mathcal{A}|$ , the key is to uniformly sample over the columns of  $\Psi$  (instead of over all states and actions like (24)) when computing unbiased estimates of  $\nabla_{\theta_\mu} h_n((\theta_v, \theta_\mu))$ . The intuition is that we should only focus on the places our basis functions care about (of size  $\dim(\Theta)$ ), instead of wasting efforts to visit all possible combinations (of size  $|\mathcal{S}||\mathcal{A}|$ ).

## 6 CONCLUSION

We propose a reduction from RL to no-regret online learning that provides a systematic way to design new RL algorithms with performance guarantees. Compared with existing approaches, our framework makes no assumption on the MDP and naturally works with function approximators. To illustrate, we design a simple RL algorithm based on mirror descent; it achieves similar sample complexity as other RL techniques, but uses minimal assumptions on the MDP and is scalable to large or continuous problems. This encouraging result evidences the strength of the online learning perspective. As a future work, we believe even faster learning in RL is possible by leveraging control variate for variance reduction and by applying more advanced online techniques [17, 18] that exploit the continuity in COL to predict the future gradients.

### Acknowledgements

This research is partially supported by NVIDIA Graduate Fellowship.

### References

- [1] Alan S Manne et al. Linear programming and sequential decision models. Technical report, Cowles Foundation for Research in Economics, Yale University, 1959.
- [2] Eric V Denardo and Bennett L Fox. Multichain Markov renewal programs. *SIAM Journal on Applied Mathematics*, 16(3):468–487, 1968.
- [3] Mengdi Wang and Yichen Chen. An online primal-dual method for discounted Markov decision processes. In *Conference on Decision and Control*, pages 4516–4521. IEEE, 2016.
- [4] Yichen Chen and Mengdi Wang. Stochastic primal-dual methods and sample complexity of reinforcement learning. *arXiv preprint arXiv:1612.02516*, 2016.
- [5] Mengdi Wang. Randomized linear programming solves the discounted Markov decision problem in nearly-linear running time. *ArXiv e-prints*, 2017.
- [6] Donghwan Lee and Niao He. Stochastic primal-dual Q-learning. *arXiv preprint arXiv:1810.08298*, 2018.
- [7] Mengdi Wang. Primal-dual  $\pi$  learning: Sample complexity and sublinear run time for ergodic Markov decision problems. *arXiv preprint arXiv:1710.06100*, 2017.
- [8] Qihang Lin, Selvaprabu Nadarajah, and Negar Soheili. Revisiting approximate linear programming using a saddle point based reformulation and root finding solution approach. Technical report, working paper, U. of Il. at Chicago and U. of Iowa, 2017.
- [9] Bo Dai, Albert Shaw, Niao He, Lihong Li, and Le Song. Boosting the actor with dual critic. In *International Conference on Learning Representation*, 2018.
- [10] Yichen Chen, Lihong Li, and Mengdi Wang. Scalable bilinear  $\pi$  learning using state and action features. *arXiv preprint arXiv:1804.10328*, 2018.
- [11] Chandrashekar Lakshminarayanan, Shalabh Bhatnagar, and Csaba Szepesvári. A linearly relaxed approximate linear program for Markov decision processes. *IEEE Transactions on Automatic Control*, 63(4):1185–1191, 2018.
- [12] Geoffrey J Gordon. Regret bounds for prediction problems. In *Conference on Learning Theory*, volume 99, pages 29–40, 1999.
- [13] Martin Zinkevich. Online convex programming and generalized infinitesimal gradient ascent. In *International Conference on Machine Learning*, pages 928–936, 2003.
- [14] Ching-An Cheng, Jonathan Lee, Ken Goldberg, and Byron Boots. Online learning with continuous variations: Dynamic regret and reductions. *arXiv preprint arXiv:1902.07286*, 2019.
- [15] Eugen Blum. From optimization and variational inequalities to equilibrium problems. *Math. student*, 63:123–145, 1994.

- [16] M Bianchi and S Schaible. Generalized monotone bifunctions and equilibrium problems. *Journal of Optimization Theory and Applications*, 90(1):31–43, 1996.
- [17] Alexander Rakhlin and Karthik Sridharan. Online learning with predictable sequences. *arXiv preprint arXiv:1208.3728*, 2012.
- [18] Ching-An Cheng, Xinyan Yan, Nathan Ratliff, and Byron Boots. Predictor-corrector policy optimization. In *International Conference on Machine Learning*, 2019.
- [19] Michael J Kearns and Satinder P Singh. Finite-sample convergence rates for q-learning and indirect algorithms. In *Advances in neural information processing systems*, pages 996–1002, 1999.
- [20] Sham Machandranath Kakade et al. *On the sample complexity of reinforcement learning*. PhD thesis, University of London London, England, 2003.
- [21] Martin L Puterman. *Markov decision processes: discrete stochastic dynamic programming*. John Wiley & Sons, 2014.
- [22] Onésimo Hernández-Lerma and Jean B Lasserre. *Discrete-time Markov control processes: basic optimality criteria*, volume 30. Springer Science & Business Media, 2012.
- [23] Richard Bellman. The theory of dynamic programming. *Bulletin of the American Mathematical Society*, 60(6):503–515, 1954.
- [24] Arkadi Nemirovski, Anatoli Juditsky, Guanghui Lan, and Alexander Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on optimization*, 19(4):1574–1609, 2009.
- [25] Anatoli Juditsky, Arkadi Nemirovski, and Claire Tauvel. Solving variational inequalities with stochastic mirror-prox algorithm. *Stochastic Systems*, 1(1):17–58, 2011.
- [26] Nicolo Cesa-Bianchi and Gabor Lugosi. *Prediction, learning, and games*. Cambridge university press, 2006.
- [27] Shai Shalev-Shwartz et al. Online learning and online convex optimization. *Foundations and Trends® in Machine Learning*, 4(2):107–194, 2012.
- [28] Elad Hazan et al. Introduction to online convex optimization. *Foundations and Trends® in Optimization*, 2(3-4):157–325, 2016.
- [29] Alejandro Jofré and Roger J-B Wets. Variational convergence of bifunctions: motivating applications. *SIAM Journal on Optimization*, 24(4):1952–1979, 2014.
- [30] Jacob Abernethy, Peter L Bartlett, and Elad Hazan. Blackwell approachability and no-regret learning are equivalent. In *Annual Conference on Learning Theory*, pages 27–46, 2011.
- [31] Andrew Y Ng, Daishi Harada, and Stuart Russell. Policy invariance under reward transformations: Theory and application to reward shaping. In *International Conference on Machine Learning*, volume 99, pages 278–287, 1999.
- [32] Sham Kakade and John Langford. Approximately optimal approximate reinforcement learning. In *International Conference on Machine Learning*, volume 2, pages 267–274, 2002.
- [33] Stéphane Ross, Geoffrey Gordon, and Drew Bagnell. A reduction of imitation learning and structured prediction to no-regret online learning. In *International Conference on Artificial Intelligence and Statistics*, pages 627–635, 2011.
- [34] Ching-An Cheng and Byron Boots. Convergence of value aggregation for imitation learning. *International Conference on Artificial Intelligence and Statistics*, 2018.
- [35] Ching-An Cheng, Xinyan Yan, Evangelos A Theodorou, and Byron Boots. Accelerating imitation learning with predictive models. In *International Conference on Artificial Intelligence and Statistics*, 2019.
- [36] Elad Hazan. Introduction to online convex optimization. *Foundations and Trends in Optimization*, 2(3-4):157–325, 2016. URL <http://dblp.uni-trier.de/db/journals/ftopt/ftopt2>.
- [37] Colin McDiarmid. Concentration. In *Probabilistic methods for algorithmic discrete mathematics*, pages 195–248. Springer, 1998.

## Appendix

### A Review of RL Setups

We provide an extended review of different formulations of RL for interested readers. First, let us recall the problem setup. Let  $\mathcal{S}$  and  $\mathcal{A}$  be state and action spaces, and let  $\pi(a|s)$  denote a policy. For  $\gamma \in [0, 1)$ , we are interested in solving a  $\gamma$ -discounted infinite-horizon RL problem:

$$\max_{\pi} V^{\pi}(p), \quad \text{s.t.} \quad V^{\pi}(p) := (1 - \gamma) \mathbb{E}_{s_0 \sim p} \mathbb{E}_{\xi \sim \rho_{\pi}(s_0)} \left[ \sum_{t=0}^{\infty} \gamma^t r(s_t, a_t) \right] \quad (25)$$

where  $V^{\pi}(p)$  is the discounted average return,  $r : \mathcal{S} \times \mathcal{A} \rightarrow [0, 1]$  is the reward function,  $\rho_{\pi}(s_0)$  denotes the distribution of trajectory  $\xi = s_0, a_0, s_1, \dots$  generated by running  $\pi$  from state  $s_0$  in a Markov decision process (MDP), and  $p$  is a fixed but unknown initial state distribution.

#### A.1 Coordinate-wise Formulations

**RL in terms of stationary state distribution** Let  $d_t^{\pi}(s)$  denote the state distribution at time  $t$  given by running  $\pi$  starting from  $p$ . We define its  $\gamma$ -weighted mixture as

$$d^{\pi}(s) := (1 - \gamma) \sum_{t=0}^{\infty} \gamma^t d_t^{\pi}(s) \quad (26)$$

We can view  $d^{\pi}$  in (26) as a form of stationary state distribution of  $\pi$ , because it is a valid probability distribution of state and satisfies the stationarity property below,

$$d^{\pi}(s') = (1 - \gamma)p(s') + \gamma \mathbb{E}_{s \sim d^{\pi}} \mathbb{E}_{a \sim \pi|s} [\mathcal{P}(s'|s, a)] \quad (2)$$

where  $\mathcal{P}(s'|s, a)$  is the transition probability of the MDP. The definition in (26) generalizes the concept of stationary distribution of MDP; as  $\gamma \rightarrow 1$ ,  $d^{\pi}$  is known as the limiting average state distribution, which is the same as the stationary distribution of the MDP under  $\pi$ , if one exists. Moreover, with the property in (2),  $d^{\pi}$  summarizes the Markov structure of RL, and allows us to write (25) simply as

$$\max_{\pi} V^{\pi}(p), \quad \text{s.t.} \quad V^{\pi}(p) = \mathbb{E}_{s \sim d^{\pi}} \mathbb{E}_{a \sim \pi|s} [r(s, a)] \quad (27)$$

after commuting the order of expectation and summation. That is, an RL problem aims to maximize the expected reward under the stationary state-action distribution generated by the policy  $\pi$ .

**RL in terms of value function** We can also write (25) in terms of value function. Recall

$$V^{\pi}(s) := (1 - \gamma) \mathbb{E}_{\xi \sim \rho_{\pi}(s_0)|s_0=s} \left[ \sum_{t=0}^{\infty} \gamma^t r(s_t, a_t) \right] \quad (1)$$

is the value function of  $\pi$ . By definition,  $V^{\pi}$  (like  $d^{\pi}$ ) satisfies a stationarity property

$$V^{\pi}(s) = \mathbb{E}_{a \sim \pi|s} \left[ (1 - \gamma)r(s, a) + \gamma \mathbb{E}_{s' \sim \mathcal{P}|s, a} [V^{\pi}(s')] \right] \quad (5)$$

which can be viewed as a dual equivalent of (2). Because  $r$  is in  $[0, 1]$ , (5) implies  $V^{\pi}$  lies in  $[0, 1]$ .

The value function  $V^*$  (a shorthand of  $V_{\pi^*}$ ) of the optimal policy  $\pi^*$  of the RL problem satisfies the so-called Bellman equation [23]:  $V^*(s) = \max_{a \in \mathcal{A}} (1 - \gamma)r(s, a) + \gamma \mathbb{E}_{s' \sim \mathcal{P}|s, a} [V^*(s')]$ , where the optimal policy  $\pi^*$  can be recovered as the arg max. Equivalently, by the definition of max, the Bellman equation amounts to finding the smallest  $V$  such that  $V(s) \geq (1 - \gamma)r(s, a) + \gamma \mathbb{E}_{s' \sim \mathcal{P}|s, a} [V(s')]$ ,  $\forall s \in \mathcal{S}, a \in \mathcal{A}$ . In other words, the RL problem in (25) can be written as

$$\min_V \mathbb{E}_{s \sim p} [V(s)] \quad \text{s.t.} \quad V(s) \geq (1 - \gamma)r(s, a) + \gamma \mathbb{E}_{s' \sim \mathcal{P}|s, a} [V(s')], \quad \forall s \in \mathcal{S}, a \in \mathcal{A} \quad (28)$$

#### A.2 Linear Programming Formulations

We now connect the above two alternate expressions through the classical LP setup of RL [1, 2].

**LP in terms of value function** The classic LP formulation<sup>5</sup> is simply a restatement of (28):

$$\min_{\mathbf{v}} \mathbf{p}^\top \mathbf{v} \quad \text{s.t.} \quad (1 - \gamma)\mathbf{r} + \gamma\mathbf{P}\mathbf{v} \leq \mathbf{E}\mathbf{v} \quad (4)$$

where  $\mathbf{p} \in \mathbb{R}^{|\mathcal{S}|}$ ,  $\mathbf{v} \in \mathbb{R}^{|\mathcal{S}|}$ , and  $\mathbf{r} \in \mathbb{R}^{|\mathcal{S}||\mathcal{A}|}$  are the vector forms of  $p$ ,  $V$ ,  $r$ , respectively,  $\mathbf{P} \in \mathbb{R}^{|\mathcal{S}||\mathcal{A}| \times |\mathcal{S}|}$  is the transition probability<sup>6</sup>, and  $\mathbf{E} = \mathbf{I} \otimes \mathbf{1} \in \mathbb{R}^{|\mathcal{S}||\mathcal{A}| \times |\mathcal{S}|}$  (we use  $|\cdot|$  to denote the cardinality of a set,  $\otimes$  the Kronecker product,  $\mathbf{I} \in \mathbb{R}^{|\mathcal{S}| \times |\mathcal{S}|}$  is the identity, and  $\mathbf{1} \in \mathbb{R}^{|\mathcal{A}|}$  a vector of ones). It is easy to verify that for all  $\mathbf{p} > 0$ , the solution to (4) is the same and equal to  $\mathbf{v}^*$  (the vector form of  $V^*$ ).

**LP in terms of stationary state-action distribution** Define the Lagrangian function

$$\mathcal{L}(\mathbf{v}, \mathbf{f}) := \mathbf{p}^\top \mathbf{v} + \mathbf{f}^\top ((1 - \gamma)\mathbf{r} + \gamma\mathbf{P}\mathbf{v} - \mathbf{E}\mathbf{v}) \quad (29)$$

where  $\mathbf{f} \geq \mathbf{0} \in \mathbb{R}^{|\mathcal{S}||\mathcal{A}|}$  is the Lagrangian multiplier. By Lagrangian duality, the dual problem of (4) is given as  $\max_{\mathbf{f} \geq \mathbf{0}} \min_{\mathbf{v}} \mathcal{L}(\mathbf{v}, \mathbf{f})$ . Or after substituting the optimality condition of  $\mathbf{v}$  and define  $\boldsymbol{\mu} := (1 - \gamma)\mathbf{f}$ , we can write the dual problem as another LP problem

$$\max_{\boldsymbol{\mu} \geq \mathbf{0}} \mathbf{r}^\top \boldsymbol{\mu} \quad \text{s.t.} \quad (1 - \gamma)\mathbf{p} + \gamma\mathbf{P}^\top \boldsymbol{\mu} = \mathbf{E}^\top \boldsymbol{\mu} \quad (3)$$

Note that this problem like (4) is normalized: we have  $\|\boldsymbol{\mu}\|_1 = 1$  because  $\|\mathbf{p}\|_1 = 1$ , and

$$\|\boldsymbol{\mu}\|_1 = \mathbf{1}^\top \mathbf{E}^\top \boldsymbol{\mu} = (1 - \gamma)\mathbf{1}^\top \mathbf{p} + \gamma\mathbf{1}^\top \mathbf{P}^\top \boldsymbol{\mu} = (1 - \gamma)\|\mathbf{p}\|_1 + \gamma\|\boldsymbol{\mu}\|_1$$

where we use the facts that  $\boldsymbol{\mu} \geq \mathbf{0}$  and  $\mathbf{P}$  is a stochastic transition matrix. This means that  $\boldsymbol{\mu}$  is a valid state-action distribution, from which we see that the equality constraint in (3) is simply a vector form (2). Therefore, (3) is the same as (27) if we define the policy  $\pi$  as the conditional distribution based on  $\boldsymbol{\mu}$ .

## B Missing Proofs of Section 3

### B.1 Proof of Lemma 1

**Lemma 1.** For any  $x = (\mathbf{v}, \boldsymbol{\mu})$ , if  $x' \in \mathcal{X}$  satisfies (2) and (5) (i.e.  $\mathbf{v}'$  and  $\boldsymbol{\mu}'$  are the value function and state-action distribution of policy  $\pi_{\boldsymbol{\mu}'}$ ),  $r_{ep}(x; x') = -\boldsymbol{\mu}'^\top \mathbf{a}_{\mathbf{v}'}$ .

*Proof.* First note that  $F(x, x) = 0$ . Then as  $x'$  satisfies stationarity, we can use Lemma 2 below and write

$$\begin{aligned} r_{ep}(x; x') &= F(x, x) - F(x, x') \\ &= -F(x, x') \\ &= -(\mathbf{p}^\top \mathbf{v}' - \mathbf{p}^\top \mathbf{v}) - \boldsymbol{\mu}'^\top \mathbf{a}_{\mathbf{v}'} + \boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}} \quad (\because \text{Definition of } F \text{ in (15)}) \\ &= -\boldsymbol{\mu}'^\top \mathbf{a}_{\mathbf{v}} - \boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}'} + \boldsymbol{\mu}'^\top \mathbf{a}_{\mathbf{v}} \quad (\because \text{Lemma 2}) \\ &= -\boldsymbol{\mu}'^\top \mathbf{a}_{\mathbf{v}'} \end{aligned}$$

□

### B.2 Proof of Lemma 2

**Lemma 2.** Let  $\mathbf{v}^\pi$  and  $\boldsymbol{\mu}^\pi$  denote the value and state-action distribution of some policy  $\pi$ . Then for any function  $\mathbf{v}'$ , it holds that  $\mathbf{p}^\top (\mathbf{v}^\pi - \mathbf{v}') = (\boldsymbol{\mu}^\pi)^\top \mathbf{a}_{\mathbf{v}'}$ . In particular, it implies  $V^\pi(p) - V^{\pi'}(p) = (\boldsymbol{\mu}^\pi)^\top \mathbf{a}_{\mathbf{v}^{\pi'}}$ .

*Proof.* This is the well-known performance difference lemma. The proof is based on the stationary properties in (2) and (5), which can be stated in vector form as

$$(\boldsymbol{\mu}^\pi)^\top \mathbf{E}\mathbf{v}^\pi = (\boldsymbol{\mu}^\pi)^\top ((1 - \gamma)\mathbf{r} + \gamma\mathbf{P}\mathbf{v}^\pi) \quad \text{and} \quad (1 - \gamma)\mathbf{p} + \gamma\mathbf{P}^\top \boldsymbol{\mu}^\pi = \mathbf{E}^\top \boldsymbol{\mu}^\pi$$

---

<sup>5</sup>Our setup in (4) differs from the classic one in the  $(1 - \gamma)$  factor in the constraint to normalize the problem.

<sup>6</sup>We arrange the coordinates in a way such that along the  $|\mathcal{S}||\mathcal{A}|$  indices are contiguous in actions.

The proof is a simple application of these two properties.

$$\begin{aligned}
 \mathbf{p}^\top(\mathbf{v}^\pi - \mathbf{v}') &= \frac{1}{1-\gamma}(\mathbf{E}^\top \boldsymbol{\mu}^\pi - \gamma \mathbf{P}^\top \boldsymbol{\mu}^\pi)^\top(\mathbf{v}^\pi - \mathbf{v}') \\
 &= \frac{1}{1-\gamma}(\boldsymbol{\mu}^\pi)^\top((\mathbf{E} - \gamma \mathbf{P})\mathbf{v}^\pi - (\mathbf{E} - \gamma \mathbf{P})\mathbf{v}') \\
 &= \frac{1}{1-\gamma}(\boldsymbol{\mu}^\pi)^\top((1-\gamma)\mathbf{r} - (\mathbf{E} - \gamma \mathbf{P})\mathbf{v}') = (\boldsymbol{\mu}^\pi)^\top \mathbf{a}_{\mathbf{v}'}
 \end{aligned}$$

where we use the stationarity property of  $\boldsymbol{\mu}^\pi$  in the first equality and that  $\mathbf{v}^\pi$  in the third equality.  $\square$

### B.3 Proof of Proposition 2

**Proposition 2.** For any  $x = (\mathbf{v}, \boldsymbol{\mu}) \in \mathcal{X}$ , if  $\mathbf{E}^\top \boldsymbol{\mu} \geq (1-\gamma)\mathbf{p}$ ,  $r_{ep}(x; x^*) \geq (1-\gamma) \min_s p(s) \|\mathbf{v}^* - \mathbf{v}^{\pi_\mu}\|_\infty$ .

*Proof.* This proof mainly follows the steps in [5] but written in our notation. First Lemma 1 shows  $r_{ep}(x; x^*) = -\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}^*}$ . We then lower bound  $-\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}^*}$  by reversing the proof of the performance difference lemma (Lemma 2).

$$\begin{aligned}
 \boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}^*} &= \frac{1}{1-\gamma} \boldsymbol{\mu}^\top ((1-\gamma)\mathbf{r} - (\mathbf{E} - \gamma \mathbf{P})\mathbf{v}^*) && (\because \text{Definition of } \mathbf{a}_{\mathbf{v}^*}) \\
 &= \frac{1}{1-\gamma} \boldsymbol{\mu}^\top ((\mathbf{E} - \gamma \mathbf{P})\mathbf{v}^{\pi_\mu} - (\mathbf{E} - \gamma \mathbf{P})\mathbf{v}^*) && (\because \text{Stationarity of } \mathbf{v}^{\pi_\mu}) \\
 &= \frac{1}{1-\gamma} \boldsymbol{\mu}^\top (\mathbf{E} - \gamma \mathbf{P})(\mathbf{v}^{\pi_\mu} - \mathbf{v}^*) \\
 &= \frac{1}{1-\gamma} \mathbf{d}^\top (\mathbf{I} - \gamma \mathbf{P}_{\pi_\mu})(\mathbf{v}^{\pi_\mu} - \mathbf{v}^*)
 \end{aligned}$$

where we define  $\mathbf{d} := \mathbf{E}^\top \boldsymbol{\mu}$  and  $\mathbf{P}_{\pi_\mu}$  as the state-transition of running policy  $\pi_\mu$ .

We wish to further upper bound this quantity. To proceed, we appeal to the Bellman equation of the optimal value function  $\mathbf{v}^*$  and the stationarity of  $\mathbf{v}^{\pi_\mu}$ :

$$\mathbf{v}^* \geq (1-\gamma)\mathbf{r}_{\pi_\mu} + \gamma \mathbf{P}_{\pi_\mu} \mathbf{v}^* \quad \text{and} \quad \mathbf{v}^{\pi_\mu} = (1-\gamma)\mathbf{r}_{\pi_\mu} + \gamma \mathbf{P}_{\pi_\mu} \mathbf{v}^{\pi_\mu},$$

which together imply that  $(\mathbf{I} - \gamma \mathbf{P}_{\pi_\mu})(\mathbf{v}^{\pi_\mu} - \mathbf{v}^*) \leq 0$ . We will also use the stationarity of  $\mathbf{d}^{\pi_\mu}$  (the average state distribution of  $\pi_\mu$ ):  $\mathbf{d}^{\pi_\mu} = (1-\gamma)\mathbf{p} + \gamma \mathbf{P}_{\pi_\mu}^\top \mathbf{d}^{\pi_\mu}$ .

Since  $\mathbf{d} \geq (1-\gamma)\mathbf{p}$  in the assumption, we can then write

$$\begin{aligned}
 \boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}^*} &= \frac{1}{1-\gamma} \mathbf{d}^\top (\mathbf{I} - \gamma \mathbf{P}_{\pi_\mu})(\mathbf{v}^{\pi_\mu} - \mathbf{v}^*) \\
 &\leq \mathbf{p}^\top (\mathbf{I} - \gamma \mathbf{P}_{\pi_\mu})(\mathbf{v}^{\pi_\mu} - \mathbf{v}^*) \\
 &\leq -\min_s p(s) \|(\mathbf{I} - \gamma \mathbf{P}_{\pi_\mu})(\mathbf{v}^{\pi_\mu} - \mathbf{v}^*)\|_\infty \\
 &\leq -\min_s p(s) (1-\gamma) \|\mathbf{v}^{\pi_\mu} - \mathbf{v}^*\|_\infty.
 \end{aligned}$$

Finally, flipping the sign of the inequality concludes the proof.  $\square$

### B.4 Proof of Proposition 3

**Proposition 3.** There is a class of MDPs such that, for some  $x \in \mathcal{X}$ , Proposition 2 is an equality.

*Proof.* We show this equality holds for a class of MDPs. For simplicity, let us first consider an MDP with three states 1, 2, 3 and for each state there are three actions (*left*, *right*, *stay*). They correspond to intuitive, deterministic transition dynamics

$$\mathcal{P}(\max\{s-1, 1\} | s, \text{left}) = 1, \quad \mathcal{P}(\min\{s+1, 3\} | s, \text{right}) = 1, \quad \mathcal{P}(s | s, \text{stay}) = 1.$$

We set the reward as  $r(s, \text{right}) = 1$  for  $s = 1, 2, 3$  and zero otherwise. It is easy to see that the optimal policy is  $\pi^*(\text{right}|s) = 1$ , which has value function  $\mathbf{v}^* = [1, 1, 1]^\top$ .

Now consider  $x = (\mathbf{v}, \boldsymbol{\mu}) \in \mathcal{X}$ . To define  $\boldsymbol{\mu}$ , let  $\mu(s, a) = d(s)\pi_{\boldsymbol{\mu}}(a|s)$ . We set

$$\pi_{\boldsymbol{\mu}}(\text{right}|1) = 1, \quad \pi_{\boldsymbol{\mu}}(\text{stay}|2) = 1, \quad \pi_{\boldsymbol{\mu}}(\text{right}|3) = 1$$

That is,  $\pi_{\boldsymbol{\mu}}$  is equal to  $\pi^*$  except when  $s = 2$ . One can verify the value function of this policy is  $\mathbf{v}^{\pi_{\boldsymbol{\mu}}} = [(1 - \gamma), 0, 1]^\top$ .

As far as  $d$  is concerned ( $\mathbf{d} = \mathbf{E}^\top \boldsymbol{\mu}$ ), suppose the initial distribution is uniform, i.e.  $\mathbf{p} = [1/3, 1/3, 1/3]^\top$ , we choose  $d$  as  $\mathbf{d} = (1 - \gamma)\mathbf{p} + \gamma[1, 0, 0]^\top$ , which satisfies the assumption in Proposition 2. Therefore, we have  $\boldsymbol{\mu} \in \mathcal{M}'$  and we will let  $\mathbf{v}$  be some arbitrary point in  $\mathcal{V}$ .

Now we show for this choice  $x = (\mathbf{v}, \boldsymbol{\mu}) \in \mathcal{V} \times \mathcal{M}'$ , the equality in Proposition 2 holds. By Lemma 1, we know  $r_{ep}(x; x') = -\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}^*}$ . Recall the advantage is defined as  $\mathbf{a}_{\mathbf{v}^*} = \mathbf{r} + \frac{1}{1-\gamma}(\gamma\mathbf{P} - \mathbf{E})\mathbf{v}^*$ . Let  $A_{V^*}(s, a)$  denote the functional form of  $\mathbf{a}_{\mathbf{v}^*}$  and define the expected advantage:

$$A_{V^*}(s, \pi_{\boldsymbol{\mu}}) := \mathbb{E}_{a \sim \pi_{\boldsymbol{\mu}}}[A_{V^*}(s, a)].$$

We can verify it has the following values:

$$A_{V^*}(1, \pi_{\boldsymbol{\mu}}) = 0, \quad A_{V^*}(2, \pi_{\boldsymbol{\mu}}) = -1, \quad A_{V^*}(3, \pi_{\boldsymbol{\mu}}) = 0.$$

Thus, the above construction yields

$$r_{ep}(x; x^*) = -\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}^*} = \frac{(1 - \gamma)}{3} = (1 - \gamma) \min_s p(s) \|\mathbf{v}^* - \mathbf{v}^{\pi_{\boldsymbol{\mu}}}\|_\infty$$

One can easily generalize this 3-state MDP to an  $|S|$ -state MDP where states are partitioned into three groups.  $\square$

## C Missing Proofs of Section 4

### C.1 Proof of Proposition 4

**Proposition 4.** For  $x = (\mathbf{v}, \boldsymbol{\mu}) \in \mathcal{X}$ , define  $y_x^* := (\mathbf{v}^{\pi_{\boldsymbol{\mu}}}, \boldsymbol{\mu}^*) \in \mathcal{X}$ . It holds  $r_{ep}(x; y_x^*) = V^*(p) - V^{\pi_{\boldsymbol{\mu}}}(p)$ .

*Proof.* First we generalize Lemma 1.

**Lemma 3.** Let  $x = (\mathbf{v}, \boldsymbol{\mu})$  be arbitrary. Consider  $\tilde{x}' = (\mathbf{v}' + \mathbf{u}', \boldsymbol{\mu}')$ , where  $\mathbf{v}'$  and  $\boldsymbol{\mu}'$  are the value function and state-action distribution of policy  $\pi_{\boldsymbol{\mu}'}$ , and  $\mathbf{u}'$  is arbitrary. It holds that  $r_{ep}(x; \tilde{x}') = -\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}'} - \mathbf{b}_{\boldsymbol{\mu}}^\top \mathbf{u}'$ .

To proceed, we write  $y_x^* = (\mathbf{v}^* + (\mathbf{v}^{\pi_{\boldsymbol{\mu}}} - \mathbf{v}^*), \boldsymbol{\mu}^*)$  and use Lemma 3, which gives  $r_{ep}(x; y_x^*) = -\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}^*} - \mathbf{b}_{\boldsymbol{\mu}}^\top (\mathbf{v}^{\pi_{\boldsymbol{\mu}}} - \mathbf{v}^*)$ . To relate this equality to the policy performance gap, we also need the following equality.

**Lemma 4.** For  $\boldsymbol{\mu} \in \mathcal{M}$ , it holds that  $-\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}^*} = V^*(p) - V^{\pi_{\boldsymbol{\mu}}}(p) + \mathbf{b}_{\boldsymbol{\mu}}^\top (\mathbf{v}^{\pi_{\boldsymbol{\mu}}} - \mathbf{v}^*)$ .

Together they imply the desired equality  $r_{ep}(x; y_x^*) = V^*(p) - V^{\pi_{\boldsymbol{\mu}}}(p)$ .  $\square$

#### C.1.1 Proof of Lemma 3

**Lemma 3.** Let  $x = (\mathbf{v}, \boldsymbol{\mu})$  be arbitrary. Consider  $\tilde{x}' = (\mathbf{v}' + \mathbf{u}', \boldsymbol{\mu}')$ , where  $\mathbf{v}'$  and  $\boldsymbol{\mu}'$  are the value function and state-action distribution of policy  $\pi_{\boldsymbol{\mu}'}$ , and  $\mathbf{u}'$  is arbitrary. It holds that  $r_{ep}(x; \tilde{x}') = -\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}'} - \mathbf{b}_{\boldsymbol{\mu}}^\top \mathbf{u}'$ .

*Proof.* Let  $x' = (\mathbf{v}', \boldsymbol{\mu}')$ . As shorthand, define  $\mathbf{f}' := \mathbf{v}' + \mathbf{u}'$ , and  $\mathbf{L} := \frac{1}{1-\gamma}(\gamma\mathbf{P} - \mathbf{E})$  (i.e. we can write  $\mathbf{a}_{\mathbf{f}'} = \mathbf{r} + \mathbf{L}\mathbf{f}'$ ). Because  $r_{ep}(x; x') = -F(x, x') = -(\mathbf{p}^\top \mathbf{v}' + \boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}'} - \mathbf{p}^\top \mathbf{v} - \boldsymbol{\mu}'^\top \mathbf{a}_{\mathbf{v}'})$ , we can write

$$\begin{aligned} r_{ep}(x; \tilde{x}') &= -\mathbf{p}^\top \mathbf{f}' - \boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{f}'} + \mathbf{p}^\top \mathbf{v} + \boldsymbol{\mu}'^\top \mathbf{a}_{\mathbf{v}'} \\ &= (-\mathbf{p}^\top \mathbf{v}' - \boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}'} + \mathbf{p}^\top \mathbf{v} + \boldsymbol{\mu}'^\top \mathbf{a}_{\mathbf{v}'}) - \mathbf{p}^\top \mathbf{u}' - \boldsymbol{\mu}^\top \mathbf{L}\mathbf{u}' \\ &= r_{ep}(x; x') - \mathbf{p}^\top \mathbf{u}' - \boldsymbol{\mu}^\top \mathbf{L}\mathbf{u}' \\ &= r_{ep}(x; x') - \mathbf{b}_{\boldsymbol{\mu}}^\top \mathbf{u}' \end{aligned}$$

Finally, by Lemma 1, we have also  $r_{ep}(x; x') = -\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}'}$  and therefore the final equality.  $\square$

### C.1.2 Proof of Lemma 4

**Lemma 4.** For  $\boldsymbol{\mu} \in \mathcal{M}$ , it holds that  $-\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}^*} = V^*(p) - V^{\pi_\mu}(p) + \mathbf{b}_\mu^\top (\mathbf{v}^{\pi_\mu} - \mathbf{v}^*)$ .

*Proof.* Following the setup in Lemma 3, we prove the statement by the rearrangement below:

$$\begin{aligned} -\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}'} &= -(\boldsymbol{\mu}^{\pi_\mu})^\top \mathbf{a}_{\mathbf{v}'} + (\boldsymbol{\mu}^{\pi_\mu})^\top \mathbf{a}_{\mathbf{v}'} - \boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}'} \\ &= V^{\pi'}(p) - V^{\pi_\mu}(p) + (\boldsymbol{\mu}^{\pi_\mu} - \boldsymbol{\mu})^\top \mathbf{a}_{\mathbf{v}'} \\ &= \left( V^{\pi'}(p) - V^{\pi_\mu}(p) \right) + (\boldsymbol{\mu}^{\pi_\mu} - \boldsymbol{\mu})^\top \mathbf{r} + (\boldsymbol{\mu}^{\pi_\mu} - \boldsymbol{\mu})^\top \mathbf{L}\mathbf{v}' \end{aligned}$$

where the second equality is due to the performance difference lemma, i.e. Lemma 2, and the last equality uses the definition  $\mathbf{a}_{\mathbf{v}'} = \mathbf{r} + \mathbf{L}\mathbf{v}'$ . For the second term above, let  $\mathbf{r}_{\pi_\mu}$  and  $\mathbf{P}_{\pi_\mu}$  denote the expected reward and transition under  $\pi_\mu$ . Because  $\boldsymbol{\mu} \in \mathcal{M}$ , we can rewrite it as

$$\begin{aligned} (\boldsymbol{\mu}^{\pi_\mu} - \boldsymbol{\mu})^\top \mathbf{r} &= (\mathbf{E}^\top \boldsymbol{\mu}^{\pi_\mu} - \mathbf{E}^\top \boldsymbol{\mu}) \mathbf{r}_{\pi_\mu} \\ &= ((1 - \gamma)\mathbf{p} + \gamma \mathbf{P}^\top \boldsymbol{\mu}^{\pi_\mu} - \mathbf{E}^\top \boldsymbol{\mu}) \mathbf{r}_{\pi_\mu} \\ &= (1 - \gamma)\mathbf{b}_\mu^\top \mathbf{r}_{\pi_\mu} + \gamma(\boldsymbol{\mu}^{\pi_\mu} - \boldsymbol{\mu})^\top \mathbf{P} \mathbf{r}_{\pi_\mu} \\ &= (1 - \gamma)\mathbf{b}_\mu^\top \left( \mathbf{r}_{\pi_\mu} + \gamma \mathbf{P}_{\pi_\mu} \mathbf{r}_{\pi_\mu} + \gamma^2 \mathbf{P}_{\pi_\mu}^2 \mathbf{r}_{\pi_\mu} + \dots \right) \\ &= \mathbf{b}_\mu^\top \mathbf{v}^{\pi_\mu} \end{aligned}$$

where the second equality uses the stationarity of  $\boldsymbol{\mu}^{\pi_\mu}$  given by (2). For the third term, it can be written

$$(\boldsymbol{\mu}^{\pi_\mu} - \boldsymbol{\mu})^\top \mathbf{L}\mathbf{v}' = (-\mathbf{p} - \mathbf{L}^\top \boldsymbol{\mu})^\top \mathbf{v}' = -\mathbf{b}_\mu^\top \mathbf{v}'$$

where the first equality uses stationarity, i.e.  $\mathbf{b}_\mu^{\pi_\mu} = \mathbf{p} + \mathbf{L}^\top \boldsymbol{\mu}^{\pi_\mu} = 0$ . Finally combining the three steps, we have

$$-\boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}'} = V^{\pi'}(p) - V^{\pi_\mu}(p) + \mathbf{b}_\mu^\top (\mathbf{v}^{\pi_\mu} - \mathbf{v}')$$

$\square$

### C.2 Proof of Corollary 1

**Corollary 1.** Let  $X_N = \{x_n \in \mathcal{X}_\theta\}_{n=1}^N$  be any sequence. Let  $\hat{\pi}_N$  be the policy given either by the average or the best decision in  $X_N$ . It holds that

$$V^{\hat{\pi}_N}(p) \geq V^*(p) - \frac{\text{Regret}_N(\Theta)}{N} - \epsilon_{\Theta, N}$$

where  $\epsilon_{\Theta, N} = \min_{x_\theta \in \mathcal{X}_\theta} r_{ep}(\hat{x}_N; y_N^*) - r_{ep}(\hat{x}_N; x_\theta)$  measures the expressiveness of  $X_\theta$ , and  $\text{Regret}_N(\Theta) := \sum_{n=1}^N l_n(x_n) - \min_{x \in \mathcal{X}_\Theta} \sum_{n=1}^N l_n(x)$ .

*Proof.* This can be proved by a simple rearrangement

$$V^*(p) - V^{\hat{\pi}_N}(p) = r_{ep}(\hat{x}_N; y_N^*) - r_{ep}(\hat{x}_N; x_\theta) \leq \epsilon_{\Theta, N} + \frac{\text{Regret}_N(\Theta)}{N}$$

where the first equality is Proposition 4 and the last inequality is due to the skew-symmetry of  $F$ , similar to the proof of Theorem 1.  $\square$

### C.3 Proof of Proposition 5

**Proposition 5.** Let  $\hat{x}_N = (\hat{\mathbf{v}}_N, \hat{\boldsymbol{\mu}}_N)$ . Under the setup in Corollary 1, regardless of the parameterization, it is true that  $\epsilon_{\Theta, N}$  is no larger than

$$\begin{aligned} & \min_{(\mathbf{v}_\theta, \boldsymbol{\mu}_\theta) \in \mathcal{X}_\Theta} \frac{\|\boldsymbol{\mu}_\theta - \boldsymbol{\mu}^*\|_1}{1-\gamma} + \min_{\mathbf{w}: \mathbf{w} \geq 1} \|\mathbf{b}_{\hat{\boldsymbol{\mu}}_N}\|_{1, \mathbf{w}} \|\mathbf{v}_\theta - \mathbf{v}^{\hat{\pi}_N}\|_{\infty, 1/\mathbf{w}} \\ & \leq \min_{(\mathbf{v}_\theta, \boldsymbol{\mu}_\theta) \in \mathcal{X}_\Theta} \frac{1}{1-\gamma} \left( \|\boldsymbol{\mu}_\theta - \boldsymbol{\mu}^*\|_1 + 2\|\mathbf{v}_\theta - \mathbf{v}^{\hat{\pi}_N}\|_\infty \right). \end{aligned}$$

where the norms are defined as  $\|\mathbf{x}\|_{1, \mathbf{w}} = \sum_i w_i |x_i|$  and  $\|\mathbf{x}\|_{\infty, 1/\mathbf{w}} = \max_i w_i^{-1} |x_i|$ .

*Proof.* For shorthand, let us set  $x = (\mathbf{v}, \boldsymbol{\mu}) = \hat{x}_N$  and write also  $\pi_\mu = \hat{\pi}_N$  as the associated policy. Let  $y_x^* = (\mathbf{v}^{\pi_\mu}, \boldsymbol{\mu}^*)$  and similarly let  $x_\theta = (\mathbf{v}_\theta, \boldsymbol{\mu}_\theta) \in \mathcal{X}_\Theta$ . With  $r_{ep}(x; x') = -F(x, x')$  and (15), we can write

$$\begin{aligned} r_{ep}(x; y_x^*) - r_{ep}(x; x_\theta) &= \left( -\mathbf{p}^\top \mathbf{v}^{\pi_\mu} - \boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}^{\pi_\mu}} + \mathbf{p}^\top \mathbf{v} + \boldsymbol{\mu}^{*\top} \mathbf{a}_{\mathbf{v}} \right) - \left( -\mathbf{p}^\top \mathbf{v}_\theta - \boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}_\theta} + \mathbf{p}^\top \mathbf{v} + \boldsymbol{\mu}_\theta^\top \mathbf{a}_{\mathbf{v}} \right) \\ &= \mathbf{p}^\top (\mathbf{v}_\theta - \mathbf{v}^{\pi_\mu}) + (\boldsymbol{\mu}^* - \boldsymbol{\mu}_\theta)^\top \mathbf{a}_{\mathbf{v}} + \boldsymbol{\mu}^\top (\mathbf{a}_{\mathbf{v}_\theta} - \mathbf{a}_{\mathbf{v}^{\pi_\mu}}) \\ &= \mathbf{b}_\mu^\top (\mathbf{v}_\theta - \mathbf{v}^{\pi_\mu}) + (\boldsymbol{\mu}^* - \boldsymbol{\mu}_\theta)^\top \mathbf{a}_{\mathbf{v}} \end{aligned}$$

Next we quantize the size of  $\mathbf{a}_{\mathbf{v}}$  and  $\mathbf{b}_\mu$ .

**Lemma 5.** For  $(\mathbf{v}, \boldsymbol{\mu}) \in \mathcal{X}$ ,  $\|\mathbf{a}_{\mathbf{v}}\|_\infty \leq \frac{1}{1-\gamma}$  and  $\|\mathbf{b}_\mu\|_1 \leq \frac{2}{1-\gamma}$ .

*Proof of Lemma 5.* Let  $\Delta$  denote the set of distributions

$$\begin{aligned} \|\mathbf{a}_{\mathbf{v}}\|_\infty &= \frac{1}{1-\gamma} \|(1-\gamma)\mathbf{r} + \gamma\mathbf{P}\mathbf{v} - \mathbf{E}\mathbf{v}\|_\infty \leq \frac{1}{1-\gamma} \max_{a, b \in [0, 1]} |a - b| \leq \frac{1}{1-\gamma} \\ \|\mathbf{b}_\mu\|_1 &= \frac{1}{1-\gamma} \|(1-\gamma)\mathbf{p} + \gamma\mathbf{P}^\top \boldsymbol{\mu} - \mathbf{E}^\top \boldsymbol{\mu}\|_1 \leq \frac{1}{1-\gamma} \max_{\mathbf{q}, \mathbf{q}' \in \Delta} \|\mathbf{q} - \mathbf{q}'\|_1 \leq \frac{2}{1-\gamma} \end{aligned}$$

□

Therefore, we have preliminary upper bounds:

$$\begin{aligned} (\boldsymbol{\mu}^* - \boldsymbol{\mu}_\theta)^\top \mathbf{a}_{\mathbf{v}} &\leq \|\mathbf{a}_{\mathbf{v}}\|_\infty \|\boldsymbol{\mu}^* - \boldsymbol{\mu}_\theta\|_1 \leq \frac{1}{1-\gamma} \|\boldsymbol{\mu}^* - \boldsymbol{\mu}_\theta\|_1 \\ \mathbf{b}_\mu^\top (\mathbf{v}_\theta - \mathbf{v}^{\pi_\mu}) &\leq \|\mathbf{b}_\mu\|_1 \|\mathbf{v}_\theta - \mathbf{v}^{\pi_\mu}\|_\infty \leq \frac{2}{1-\gamma} \|\mathbf{v}_\theta - \mathbf{v}^{\pi_\mu}\|_\infty \end{aligned}$$

However, the second inequality above can be very conservative, especially when  $\mathbf{b}_\mu \approx 0$  which can be likely when it is close to the end of policy optimization. To this end, we introduce a free vector  $\mathbf{w} \geq 1$ . Define norms  $\|\mathbf{v}\|_{\infty, 1/\mathbf{w}} = \max_i \frac{|v_i|}{w_i}$  and  $\|\boldsymbol{\delta}\|_{1, \mathbf{w}} = \sum_i w_i |\delta_i|$ . Then we can instead have an upper bound

$$\mathbf{b}_\mu^\top (\mathbf{v}_\theta - \mathbf{v}^{\pi_\mu}) \leq \min_{\mathbf{w}: \mathbf{w} \geq 1} \|\mathbf{b}_\mu\|_{1, \mathbf{w}} \|\mathbf{v}_\theta - \mathbf{v}^{\pi_\mu}\|_{\infty, 1/\mathbf{w}}$$

Notice that when  $\mathbf{w} = \mathbf{1}$  the above inequality reduces to  $\mathbf{b}_\mu^\top (\mathbf{v}_\theta - \mathbf{v}^{\pi_\mu}) \leq \|\mathbf{b}_\mu\|_1 \|\mathbf{v}_\theta - \mathbf{v}^{\pi_\mu}\|_\infty$ , which as we showed has an upper bound  $\frac{2}{1-\gamma} \|\mathbf{v}_\theta - \mathbf{v}^{\pi_\mu}\|_\infty$ .

Combining the above upper bounds, we have an upper bound on  $\epsilon_{\Theta, N}$ :

$$\begin{aligned} \epsilon_{\Theta, N} = r_{ep}(x; y_x^*) - r_{ep}(x; x_\theta) &\leq \frac{1}{1-\gamma} \|\boldsymbol{\mu}_\theta - \boldsymbol{\mu}^*\|_1 + \min_{\mathbf{w}: \mathbf{w} \geq 1} \|\mathbf{b}_\mu\|_{1, \mathbf{w}} \|\mathbf{v}_\theta - \mathbf{v}^{\pi_\mu}\|_{\infty, 1/\mathbf{w}} \\ &\leq \frac{1}{1-\gamma} \left( \|\boldsymbol{\mu}_\theta - \boldsymbol{\mu}^*\|_1 + 2\|\mathbf{v}_\theta - \mathbf{v}^{\pi_\mu}\|_\infty \right). \end{aligned}$$

Since it holds for any  $\theta \in \Theta$ , we can minimize the right-hand side over all possible choices. □



## D Proof of Sample Complexity of Mirror Descent

**Theorem 2.** *With probability  $1 - \delta$ , Algorithm 1 learns an  $\epsilon$ -optimal policy with  $\tilde{O}\left(\frac{|\mathcal{S}||\mathcal{A}|\log(\frac{1}{\delta})}{(1-\gamma)^2\epsilon^2}\right)$  samples.*

The proof is a combination of the basic property of mirror descent (Lemma 9) and the martingale concentration. Define  $K = |\mathcal{S}||\mathcal{A}|$  and  $\kappa = \frac{1}{1-\gamma}$  as shorthands. We first slightly modify the per-round loss used to compute the gradient. Recall  $l_n(x) := \mathbf{p}^\top \mathbf{v} + \boldsymbol{\mu}_n^\top \mathbf{a}_\mathbf{v} - \mathbf{p}^\top \mathbf{v}_n - \boldsymbol{\mu}^\top \mathbf{a}_{\mathbf{v}_n}$  and let us consider instead a loss function

$$h_n(x) := \mathbf{b}_{\boldsymbol{\mu}_n}^\top \mathbf{v} + \boldsymbol{\mu}^\top (\kappa \mathbf{1} - \mathbf{a}_{\mathbf{v}_n})$$

which shifts  $l_n$  by a constant in each round. (Note for all the decisions  $(\mathbf{v}_n, \boldsymbol{\mu}_n)$  produced by Algorithm 1  $\boldsymbol{\mu}_n$  always satisfies  $\|\boldsymbol{\mu}_n\|_1 = 1$ ). One can verify that  $l_n(x) - l_n(x') = h_n(x) - h_n(x')$ , for all  $x, x' \in \mathcal{X}$ , when  $\boldsymbol{\mu}, \boldsymbol{\mu}'$  in  $x$  and  $x'$  satisfy  $\|\boldsymbol{\mu}\|_1 = \|\boldsymbol{\mu}'\|_1$  (which holds for Algorithm 1). As the definition of regret is relative, we may work with  $h_n$  in online learning and use it to define the feedback.

The reason for using  $h_n$  instead of  $l_n$  is to make  $\nabla_{\boldsymbol{\mu}} h_n((\mathbf{v}, \boldsymbol{\mu}))$  (and its unbiased approximation) a positive vector (because  $\kappa \geq \|\mathbf{a}_\mathbf{v}\|_\infty$  for any  $\mathbf{v} \in \mathcal{V}$ ), so that the regret bound can have a better dependency on the dimension for learning  $\boldsymbol{\mu}$  that lives in the simplex  $\mathcal{M}$ . This is a common trick used in the online learning, e.g. in EXP3.

To run mirror descent, we set the first-order feedback  $g_n$  received by the learner as an unbiased estimate of  $\nabla h_n(x_n)$ . For round  $n$ , we construct  $g_n$  based on *two* calls of the generative model:

$$g_n = \begin{bmatrix} \mathbf{g}_{n,v} \\ \mathbf{g}_{n,\mu} \end{bmatrix} = \begin{bmatrix} \tilde{\mathbf{p}}_n + \frac{1}{1-\gamma} (\gamma \tilde{\mathbf{P}}_n - \mathbf{E}_n)^\top \tilde{\boldsymbol{\mu}}_n \\ K(\kappa \hat{\mathbf{1}}_n - \hat{\mathbf{r}}_n - \frac{1}{1-\gamma} (\gamma \hat{\mathbf{P}}_n - \hat{\mathbf{E}}_n) \mathbf{v}_n) \end{bmatrix}$$

For  $\mathbf{g}_{n,v}$ , we sample  $\mathbf{p}$ , then sample  $\boldsymbol{\mu}_n$  to get a state-action pair, and finally query the transition dynamics  $\mathbf{P}$  at the state-action pair sampled from  $\boldsymbol{\mu}_n$ . ( $\tilde{\mathbf{p}}_n$ ,  $\tilde{\mathbf{P}}_n$ , and  $\tilde{\boldsymbol{\mu}}_n$  denote the single-sample empirical approximation of these probabilities.) For  $\mathbf{g}_{n,\mu}$ , we first sample *uniformly* a state-action pair (which explains the factor  $K$ ), and then query the reward  $\mathbf{r}$  and the transition dynamics  $\mathbf{P}$ . ( $\hat{\mathbf{1}}_n$ ,  $\hat{\mathbf{r}}_n$ ,  $\hat{\mathbf{P}}_n$ , and  $\hat{\mathbf{E}}_n$  denote the single-sample empirical estimates.) To emphasize, we use  $\tilde{\cdot}$  and  $\hat{\cdot}$  to distinguish the empirical quantities obtained by these two independent queries. By construction, we have  $\mathbf{g}_{n,\mu} \geq 0$ . It is clear that this direction  $g_n$  is unbiased, i.e.  $\mathbb{E}[g_n] = \nabla h_n(x_n)$ . Moreover, it is extremely sparse and can be computed using  $O(1)$  sample, computational, and memory complexities.

Let  $y_N^* = (\mathbf{v}^{\hat{\pi}_N}, \boldsymbol{\mu}^*)$ . We bound the regret by the following rearrangement.

$$\begin{aligned} \text{Regret}_N(y_N^*) &= \sum_{n=1}^N l_n(x_n) - \sum_{n=1}^N l_n(y_N^*) \\ &= \sum_{n=1}^N h_n(x_n) - \sum_{n=1}^N h_n(y_N^*) \\ &= \sum_{n=1}^N \nabla h_n(x_n)^\top (x_n - y_N^*) \\ &= \left( \sum_{n=1}^N (\nabla h_n(x_n) - g_n)^\top x_n \right) + \left( \sum_{n=1}^N g_n^\top (x_n - y_N^*) \right) + \left( \sum_{n=1}^N (g_n - \nabla h_n(x_n))^\top y_N^* \right) \\ &\leq \left( \sum_{n=1}^N (\nabla h_n(x_n) - g_n)^\top x_n \right) + \left( \max_{x \in \mathcal{X}} \sum_{n=1}^N g_n^\top (x_n - x) \right) + \left( \sum_{n=1}^N (g_n - \nabla h_n(x_n))^\top y_N^* \right), \quad (30) \end{aligned}$$

where the third equality comes from  $h_n$  being linear. We recognize the first term is a martingale  $M_N = \sum_{n=1}^N (\nabla h_n(x_n) - g_n)^\top x_n$ , because  $x_n$  does not depend on  $g_n$ . Therefore, we can appeal to standard martingale concentration property. For the second term, it can be upper bounded by standard regret analysis of mirror descent, by treating  $g_n^\top x$  as the per-round loss. For the third term, because  $y_N^* = (\mathbf{v}^{\hat{\pi}_N}, \boldsymbol{\mu}^*)$  depends on  $\{g_n\}_{n=1}^N$ , it is not a martingale. Nonetheless, we will be able to handle it through a union bound. Below, we give details for bounding these three terms.

### D.1 The First Term: Martingale Concentration

For the first term,  $\sum_{n=1}^N (\nabla h_n(x_n) - g_n)^\top x_n$ , we use a martingale concentration property. Specifically, we adopt a Bernstein-type inequality [37, Theorem 3.15]:

**Lemma 6.** [37, Theorem 3.15] *Let  $M_0, \dots, M_N$  be a martingale and let  $F_0 \subseteq F_1 \subseteq \dots \subseteq F_n$  be the filtration such that  $M_n = \mathbb{E}_{|F_n}[M_{n+1}]$ . Suppose there are  $b, \sigma < \infty$  such that for all  $n$ , given  $F_{n-1}$ ,  $M_n - M_{n-1} \leq b$ , and  $\mathbb{V}_{|F_{n-1}}[M_n - M_{n-1}] \leq \sigma^2$  almost surely. Then for any  $\epsilon \geq 0$ ,*

$$P(M_N - M_0 \geq \epsilon) \leq \exp\left(\frac{-\epsilon^2}{2N\sigma^2(1 + \frac{b\epsilon}{3N\sigma^2})}\right).$$

Lemma 6 implies, with probability at least  $1 - \delta$ ,

$$M_N - M_0 \leq \sqrt{2N\sigma^2(1 + o(1)) \log\left(\frac{1}{\delta}\right)},$$

where  $o(1)$  means convergence to 0 as  $N \rightarrow \infty$ .

To apply Lemma 6, we need to provide bounds on the properties of the martingale difference:

$$\begin{aligned} M_n - M_{n-1} &= (\nabla h_n(x_n) - g_n)^\top x_n \\ &= (\kappa \mathbf{1} - \mathbf{a}_{\mathbf{v}_n} - \mathbf{g}_{n,\mu})^\top \boldsymbol{\mu}_n + (\mathbf{b}_{\boldsymbol{\mu}_n} - \mathbf{g}_{n,v})^\top \mathbf{v}_n. \end{aligned}$$

For the first term  $(\kappa \mathbf{1} - \mathbf{a}_{\mathbf{v}_n} - \mathbf{g}_{n,\mu})^\top \boldsymbol{\mu}_n$ , we use the lemma below:

**Lemma 7.** *Let  $\boldsymbol{\mu} \in \mathcal{M}$  be arbitrary, chosen independently from the randomness of  $\mathbf{g}_{n,\mu}$  when  $F_{n-1}$  is given. Then it holds  $|(\kappa \mathbf{1} - \mathbf{a}_{\mathbf{v}_n} - \mathbf{g}_{n,\mu})^\top \boldsymbol{\mu}| \leq \frac{2(1+K)}{1-\gamma}$  and  $\mathbb{V}_{|F_{n-1}}[(\kappa \mathbf{1} - \mathbf{a}_{\mathbf{v}_n} - \mathbf{g}_{n,\mu})^\top \boldsymbol{\mu}] \leq \frac{4K}{(1-\gamma)^2}$ .*

*Proof.* By triangular inequality,

$$|(\kappa \mathbf{1} - \mathbf{a}_{\mathbf{v}_n} - \mathbf{g}_{n,\mu})^\top \boldsymbol{\mu}| \leq |(\kappa \mathbf{1} - \mathbf{a}_{\mathbf{v}_n})^\top \boldsymbol{\mu}| + |\mathbf{g}_{n,\mu}^\top \boldsymbol{\mu}|.$$

For the deterministic part, using Lemma 5 and Hölder's inequality,

$$|(\kappa \mathbf{1} - \mathbf{a}_{\mathbf{v}_n})^\top \boldsymbol{\mu}| \leq \kappa + \|\mathbf{a}_{\mathbf{v}_n}\|_\infty \|\boldsymbol{\mu}\|_1 \leq \frac{2}{1-\gamma}.$$

For the stochastic part, let  $i_n$  be index of the sampled state-action pair and  $j_n$  be the index of the transited state sampled at the pair given by  $i_n$ . With abuse of notation, we will use  $i_n$  to index both  $\mathcal{S} \times \mathcal{A}$  and  $\mathcal{S}$ . With this notation, we may derive

$$\begin{aligned} |\mathbf{g}_{n,\mu}^\top \boldsymbol{\mu}| &= |K \boldsymbol{\mu}^\top (\kappa \hat{\mathbf{1}}_n - \hat{\mathbf{r}}_n - \frac{1}{1-\gamma} (\gamma \hat{\mathbf{P}}_n - \hat{\mathbf{E}}_n) \mathbf{v}_n)| \\ &= K \mu_{i_n} \left| \kappa - r_{i_n} - \frac{\gamma v_{n,j_n} - v_{n,i_n}}{1-\gamma} \right| \\ &\leq \frac{2K \mu_{i_n}}{1-\gamma} \leq \frac{2K}{1-\gamma} \end{aligned}$$

where we use the facts that  $r_{i_n}, v_{n,j_n}, v_{n,i_n} \in [0, 1]$  and  $\mu_{i_n} \leq 1$ .

For  $\mathbb{V}_{|F_{n-1}}[(\kappa \mathbf{1} - \mathbf{a}_{\mathbf{v}_n} - \mathbf{g}_{n,\mu})^\top \boldsymbol{\mu}_n]$ , we can write it as

$$\begin{aligned} \mathbb{V}_{|F_{n-1}}[(\kappa \mathbf{1} - \mathbf{a}_{\mathbf{v}_n} - \mathbf{g}_{n,\mu})^\top \boldsymbol{\mu}] &= \mathbb{V}_{|F_{n-1}}[\mathbf{g}_{n,\mu}^\top \boldsymbol{\mu}] \\ &\leq \mathbb{E}_{|F_{n-1}}[|\mathbf{g}_{n,\mu}^\top \boldsymbol{\mu}_n|^2] \\ &= \sum_{i_n} \frac{1}{K} \mathbb{E}_{j_n|i_n} \left[ K^2 \mu_{i_n}^2 \left( \kappa - r_{i_n} - \frac{\gamma v_{n,j_n} - v_{n,i_n}}{1-\gamma} \right)^2 \right] \\ &\leq \frac{4K}{(1-\gamma)^2} \sum_{i_n} \mu_{i_n}^2 \\ &\leq \frac{4K}{(1-\gamma)^2} \left( \sum_{i_n} \mu_{i_n} \right)^2 \leq \frac{4K}{(1-\gamma)^2} \end{aligned}$$

where in the second inequality we use the fact that  $|\kappa - r_{i_n} - \frac{\gamma v_{n,j_n} - v_{n,i_n}}{1-\gamma}| \leq \frac{2}{1-\gamma}$ .  $\square$

For the second term  $(\mathbf{b}_{\boldsymbol{\mu}_n} - \mathbf{g}_{n,v})^\top \mathbf{v}_n$ , we use the following lemma.

**Lemma 8.** *Let  $\mathbf{v} \in \mathcal{V}$  be arbitrary, chosen independently from the randomness of  $\mathbf{g}_{n,v}$  when  $F_{n-1}$  is given. Then it holds that  $|(\mathbf{b}_{\boldsymbol{\mu}_n} - \mathbf{g}_{n,v})^\top \mathbf{v}| \leq \frac{4}{1-\gamma}$  and  $\mathbb{V}_{|F_{n-1}}[(\mathbf{b}_{\boldsymbol{\mu}_n} - \mathbf{g}_{n,v})^\top \mathbf{v}] \leq \frac{4}{(1-\gamma)^2}$ .*

*Proof.* We appeal to Lemma 5, which shows  $\|\mathbf{b}_{\boldsymbol{\mu}_n}\|_1, \|\mathbf{g}_{n,v}\|_1 \leq \frac{2}{1-\gamma}$ , and derive

$$|(\mathbf{b}_{\boldsymbol{\mu}_n} - \mathbf{g}_{n,v})^\top \mathbf{v}| \leq (\|\mathbf{b}_{\boldsymbol{\mu}_n}\|_1 + \|\mathbf{g}_{n,v}\|_1) \|\mathbf{v}\|_\infty \leq \frac{4}{1-\gamma}.$$

Similarly, for the variance, we can write

$$\mathbb{V}_{|F_{n-1}}[(\mathbf{b}_{\boldsymbol{\mu}_n} - \mathbf{g}_{n,v})^\top \mathbf{v}] = \mathbb{V}_{|F_{n-1}}[\mathbf{g}_{n,v}^\top \mathbf{v}] \leq \mathbb{E}_{|F_{n-1}}[(\mathbf{g}_{n,v}^\top \mathbf{v})^2] \leq \frac{4}{(1-\gamma)^2}. \quad \square$$

Thus, with helps from the two lemmas above, we are able to show

$$M_n - M_{n-1} \leq |(\kappa \mathbf{1} - \mathbf{a}_{\mathbf{v}_n} - \mathbf{g}_{n,\mu})^\top \boldsymbol{\mu}_n| + |(\mathbf{b}_{\boldsymbol{\mu}_n} - \mathbf{g}_{n,v})^\top \mathbf{v}_n| \leq \frac{4 + 2(1+K)}{1-\gamma}$$

as well as (because  $\mathbf{g}_{n,\mu}$  and  $\mathbf{g}_{n,v}$  are computed using independent samples)

$$\mathbb{V}_{|F_{n-1}}[M_n - M_{n-1}] \leq \mathbb{E}_{|F_{n-1}}[|(\kappa \mathbf{1} - \mathbf{a}_{\mathbf{v}_n} - \mathbf{g}_{n,\mu})^\top \boldsymbol{\mu}_n|^2] + \mathbb{E}_{|F_{n-1}}[|(\mathbf{b}_{\boldsymbol{\mu}_n} - \mathbf{g}_{n,v})^\top \mathbf{v}_n|^2] \leq \frac{4(1+K)}{(1-\gamma)^2}$$

Now, since  $M_0 = 0$ , by martingale concentration in Lemma 6, we have

$$P \left( \sum_{n=1}^N (\nabla h_n(x_n) - g_n)^\top x_n > \epsilon \right) \leq \exp \left( \frac{-\epsilon^2}{2N\sigma^2(1 + \frac{b\epsilon}{3N\sigma^2})} \right)$$

with  $b = \frac{6+2K}{1-\gamma}$  and  $\sigma^2 = \frac{4(1+K)}{(1-\gamma)^2}$ . This implies that, with probability at least  $1 - \delta$ , it holds

$$\sum_{n=1}^N (\nabla h_n(x_n) - g_n)^\top x_n \leq \sqrt{N \frac{8(1+K)}{(1-\gamma)^2} (1 + o(1)) \log \left( \frac{1}{\delta} \right)} = \tilde{O} \left( \frac{\sqrt{NK \log \left( \frac{1}{\delta} \right)}}{1-\gamma} \right)$$

## D.2 Static Regret of Mirror Descent

Next we move onto deriving the regret bound of mirror descent with respect to the online loss sequence:

$$\max_{x \in \mathcal{X}} \sum_{n=1}^N g_n^\top (x_n - x)$$

This part is quite standard; nonetheless, we provide complete derivations below.

We first recall a basic property of mirror descent

**Lemma 9.** Let  $\mathcal{X}$  be a convex set. Suppose  $R$  is 1-strongly convex with respect to some norm  $\|\cdot\|$ . Let  $g$  be an arbitrary vector and define, for  $x \in \mathcal{X}$ ,

$$y = \arg \min_{x' \in \mathcal{X}} \langle g, x' \rangle + B_R(x'|x)$$

Then for all  $z \in \mathcal{X}$ ,

$$\langle g, y - z \rangle \leq B_R(z|x) - B_R(z|y) - B_R(y|x) \quad (31)$$

*Proof.* Recall the definition  $B_R(x'|x) = R(x') - R(x) - \langle \nabla R(x), x' - x \rangle$ . The optimality of the proximal map can be written as

$$\langle g + \nabla R(y) - \nabla R(x), y - z \rangle \leq 0, \quad \forall z \in \mathcal{X}.$$

By rearranging the terms, we can rewrite the above inequality in terms of Bregman divergences as follows and derive the first inequality (31):

$$\begin{aligned} \langle g, y - z \rangle &\leq \langle \nabla R(x) - \nabla R(y), y - z \rangle \\ &= B_R(z|x) - B_R(z|y) + \langle \nabla R(x) - \nabla R(y), y \rangle - \langle \nabla R(x), x \rangle + \langle \nabla R(y), y \rangle + R(x) - R(y) \\ &= B_R(z|x) - B_R(z|y) + \langle \nabla R(x), y - x \rangle + R(x) - R(y) \\ &= B_R(z|x) - B_R(z|y) - B_R(y|x), \end{aligned}$$

which concludes the lemma.  $\square$

Let  $x' \in \mathcal{X}$  be arbitrary. Applying this lemma to the  $n$ th iteration of mirror descent in (21), we get

$$\langle g_n, x_{n+1} - x' \rangle \leq \frac{1}{\eta} (B_R(x'|x_n) - B_R(x'|x_{n+1}) - B_R(x_{n+1}|x_n))$$

By a telescoping sum, we then have

$$\sum_{n=1}^N \langle g_n, x_n - x' \rangle \leq \frac{1}{\eta} B_R(x'|x_1) + \sum_{n=1}^N \langle g_n, x_{n+1} - x_n \rangle - \frac{1}{\eta} B_R(x_{n+1}|x_n).$$

We bound the right-hand side as follows. Recall that based on the geometry of  $\mathcal{X} = \mathcal{V} \times \mathcal{M}$ , we considered a natural Bregman divergence of the form:

$$B_R(x'|x) = \frac{1}{2|\mathcal{S}|} \|\mathbf{v}' - \mathbf{v}\|_2^2 + KL(\boldsymbol{\mu}'||\boldsymbol{\mu})$$

Let  $x_1 = (\mathbf{v}_1, \boldsymbol{\mu}_1)$  where  $\boldsymbol{\mu}_1$  is uniform. By this choice, we have:

$$\frac{1}{\eta} B_R(x'|x_1) \leq \frac{1}{\eta} \max_{x \in \mathcal{X}} B_R(x|x_1) \leq \frac{1}{\eta} \left( \frac{1}{2} + \log(K) \right).$$

We now decompose each item in the above sum as:

$$\begin{aligned} \langle g_n, x_{n+1} - x_n \rangle - \frac{1}{\eta} B_R(x_{n+1}|x_n) &= \left( \mathbf{g}_{n,v}^\top (\mathbf{v}_{n+1} - \mathbf{v}_n) - \frac{1}{2\eta|\mathcal{S}|} \|\mathbf{v}_n - \mathbf{v}_{n+1}\|_2^2 \right) \\ &\quad + \left( \mathbf{g}_{n,\mu}^\top (\boldsymbol{\mu}_{n+1} - \boldsymbol{\mu}_n) - \frac{1}{\eta} KL(\boldsymbol{\mu}_{n+1}||\boldsymbol{\mu}_n) \right) \end{aligned}$$

and we upper bound them using the two lemmas below (recall  $\mathbf{g}_{n,\mu} \geq 0$  due to the added  $\kappa \mathbf{1}$  term).

**Lemma 10.** For any vector  $x, y, g$  and scalar  $\eta > 0$ , it holds  $\langle g, x - y \rangle - \frac{1}{2\eta} \|x - y\|_2^2 \leq \frac{\eta \|g\|_2^2}{2}$ .

*Proof.* By Cauchy-Swartz inequality,  $\langle g, x - y \rangle - \frac{1}{2\eta} \|x - y\|_2^2 \leq \|g\|_2 \|x - y\|_2 - \frac{1}{2\eta} \|x - y\|_2^2 \leq \frac{\eta \|g\|_2^2}{2}$ .  $\square$

**Lemma 11.** Suppose  $B_R(x||y) = KL(x||y)$  and  $x, y$  are probability distributions, and  $g \geq 0$  element-wise. Then, for  $\eta > 0$ ,

$$-\frac{1}{\eta}B_R(y||x) + \langle g, x - y \rangle \leq \frac{\eta}{2} \sum_i x_i (g_i)^2 = \frac{\eta}{2} \|g\|_x^2.$$

*Proof.* Let  $\Delta$  denotes the unit simplex.

$$\begin{aligned} -B_R(y||x) + \langle \eta g, x - y \rangle &\leq \langle \eta g, x \rangle + \max_{y' \in \Delta} \langle -\eta g, y' \rangle - B_R(y' || x) \\ &= \langle \eta g, x \rangle + \log \left( \sum_i x_i \exp(-\eta g_i) \right) && (\because \text{convex conjugate of KL divergence}) \\ &\leq \langle \eta g, x \rangle + \log \left( \sum_i x_i \left( 1 - \eta g_i + \frac{1}{2}(\eta g_i)^2 \right) \right) && (\because e^x \leq 1 + x + \frac{1}{2}x^2 \text{ for } x \leq 0) \\ &= \langle \eta g, x \rangle + \log \left( 1 + \sum_i x_i \left( -\eta g_i + \frac{1}{2}(\eta g_i)^2 \right) \right) \\ &\leq \langle \eta g, x \rangle + \sum_i x_i \left( -\eta g_i + \frac{1}{2}(\eta g_i)^2 \right) && (\because \log(x) \leq x - 1) \\ &= \frac{1}{2} \sum_i x_i (\eta g_i)^2 = \frac{\eta^2}{2} \|g\|_x^2. \end{aligned}$$

Finally, dividing both sides by  $\eta$ , we get the desired result.  $\square$

Thus, we have the upper bound  $\langle g_n, x_{n+1} - x_n \rangle - \frac{1}{\eta}B_R(x_{n+1}||x_n) = \frac{\eta|\mathcal{S}|\|\mathbf{g}_{n,v}\|_2^2}{2} + \frac{\eta\|\mathbf{g}_{n,\mu}\|_{\mu_n}^2}{2}$ . Together with the upper bound on  $\frac{1}{\eta}B_R(x' || x_1)$ , it implies that

$$\begin{aligned} \sum_{n=1}^N \langle g_n, x_n - x' \rangle &\leq \frac{1}{\eta}B_R(x' || x_1) + \sum_{n=1}^N \langle g_n, x_{n+1} - x_n \rangle - \frac{1}{\eta}B_R(x_{n+1} || x_n) \\ &\leq \frac{1}{\eta} \left( \frac{1}{2} + \log(K) \right) + \frac{\eta}{2} \sum_{n=1}^N |\mathcal{S}| \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\mu_n}^2. \end{aligned} \quad (32)$$

We can expect, with high probability,  $\sum_{n=1}^N |\mathcal{S}| \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\mu_n}^2$  concentrates toward its expectation, i.e.

$$\sum_{n=1}^N |\mathcal{S}| \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\mu_n}^2 \leq \sum_{n=1}^N \mathbb{E}[|\mathcal{S}| \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\mu_n}^2] + o(N).$$

Below we quantify this relationship using martingale concentration. First we bound the expectation.

**Lemma 12.**  $\mathbb{E}[\|\mathbf{g}_{n,v}\|_2^2] \leq \frac{4}{(1-\gamma)^2}$  and  $\mathbb{E}[\|\mathbf{g}_{n,\mu}\|_{\mu_n}^2] \leq \frac{4K}{(1-\gamma)^2}$ .

*Proof.* For the first statement, using the fact that  $\|\cdot\|_2 \leq \|\cdot\|_1$  and Lemma 5, we can write

$$\mathbb{E}[\|\mathbf{g}_{n,v}\|_2^2] \leq \mathbb{E}[\|\mathbf{g}_{n,v}\|_1^2] = \mathbb{E}[\|\tilde{\mathbf{p}}_n + \frac{1}{1-\gamma}(\gamma\tilde{\mathbf{P}}_n - \mathbf{E}_n)^\top \tilde{\boldsymbol{\mu}}_n\|_1^2] \leq \frac{4}{(1-\gamma)^2}.$$

For the second statement, let  $i_n$  be the index of the sampled state-action pair and  $j_n$  be the index of the transited-to state sampled at the pair given by  $i_n$ . With abuse of notation, we will use  $i_n$  to index both  $\mathcal{S} \times \mathcal{A}$  and  $\mathcal{S}$ .

$$\begin{aligned} \mathbb{E}[\|\mathbf{g}_{n,\mu}\|_{\mu_n}^2] &= \mathbb{E} \left[ \sum_{i_n} \frac{1}{K} \mathbb{E}_{j_n|i_n} \left[ K^2 \mu_{i_n} \left( \kappa - r_{i_n} - \frac{\gamma v_{n,j_n} - v_{n,i_n}}{1-\gamma} \right)^2 \right] \right] \\ &\leq \frac{4K}{(1-\gamma)^2} \mathbb{E} \left[ \sum_{i_n} \mu_{i_n} \right] \leq \frac{4K}{(1-\gamma)^2}. \end{aligned} \quad \square$$

To bound the tail, we resort to the Höfdding-Azuma inequality of martingale [37, Theorem 3.14].

**Lemma 13** (Azuma-Hoeffding). *Let  $M_0, \dots, M_N$  be a martingale and let  $F_0 \subseteq F_1 \subseteq \dots \subseteq F_n$  be the filtration such that  $M_n = \mathbb{E}_{|F_n}[M_{n+1}]$ . Suppose there exists  $b < \infty$  such that for all  $n$ , given  $F_{n-1}$ ,  $|M_n - M_{n-1}| \leq b$ . Then for any  $\epsilon \geq 0$ ,*

$$P(M_N - M_0 \geq \epsilon) \leq \exp\left(\frac{-2\epsilon^2}{Nb^2}\right)$$

To apply Lemma 13, we consider the martingale

$$M_N = \sum_{n=1}^N |\mathcal{S}| \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\mu_n}^2 - \left( \sum_{n=1}^N \mathbb{E}[|\mathcal{S}| \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\mu_n}^2] \right)$$

To bound the change of the size of martingale difference  $|M_n - M_{n-1}|$ , we follow similar steps to Lemma 12.

**Lemma 14.**  $\|\mathbf{g}_{n,v}\|_2^2 \leq \frac{4}{(1-\gamma)^2}$  and  $\|\mathbf{g}_{n,\mu}\|_{\mu_n}^2 \leq \frac{4K^2}{(1-\gamma)^2}$ .

Note  $\|\mathbf{g}_{n,\mu}\|_{\mu}^2$  is  $K$ -factor larger than  $\mathbb{E}[\|\mathbf{g}_{n,\mu}\|_{\mu}^2]$  and  $K \geq 1$ . Therefore, we have

$$|M_n - M_{n-1}| \leq |\mathcal{S}| \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\mu_n}^2 + |\mathcal{S}| \mathbb{E}[\|\mathbf{g}_{n,v}\|_2^2] + \mathbb{E}[\|\mathbf{g}_{n,\mu}\|_{\mu_n}^2] \leq \frac{8(|\mathcal{S}| + K^2)}{(1-\gamma)^2}$$

Combining these results, we have, with probability as least  $1 - \delta$ ,

$$\begin{aligned} \sum_{n=1}^N |\mathcal{S}| \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\mu_n}^2 &\leq \sum_{n=1}^N \mathbb{E}[|\mathcal{S}| \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\mu_n}^2] + \frac{4\sqrt{2}(|\mathcal{S}| + K^2)}{(1-\gamma)^2} \sqrt{N \log\left(\frac{1}{\delta}\right)} \\ &\leq \frac{4(K + |\mathcal{S}|)}{(1-\gamma)^2} N + \frac{4\sqrt{2}(|\mathcal{S}| + K^2)}{(1-\gamma)^2} \sqrt{N \log\left(\frac{1}{\delta}\right)} \end{aligned}$$

Now we suppose we set  $\eta = \frac{1-\gamma}{\sqrt{KN}}$ . From (38), we then have

$$\begin{aligned} \sum_{n=1}^N \langle g_n, x_n - x' \rangle &\leq \frac{1}{\eta} \left( \frac{1}{2} + \log(K) \right) + \frac{\eta}{2} \sum_{n=1}^N |\mathcal{S}| \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\mu_n}^2 \\ &\leq \frac{\sqrt{KN}}{1-\gamma} \left( \frac{1}{2} + \log(K) \right) + \frac{1-\gamma}{\sqrt{KN}} \left( \frac{2(K + |\mathcal{S}|)}{(1-\gamma)^2} N + \frac{2\sqrt{2}(|\mathcal{S}| + K^2)}{(1-\gamma)^2} \sqrt{N \log\left(\frac{1}{\delta}\right)} \right) \\ &\leq \tilde{O} \left( \frac{\sqrt{KN}}{1-\gamma} + \frac{\sqrt{K^3 \log \frac{1}{\delta}}}{1-\gamma} \right). \end{aligned}$$

### D.3 Union Bound

Lastly, we provide an upper bound on the last component:

$$\sum_{n=1}^N (g_n - \nabla h_n(x_n))^\top y_N^*.$$

Because  $y_N^*$  depends on  $g_n$ , this term does not obey martingale concentration like the first component  $\sum_{n=1}^N (\nabla h_n(x_n) - g_n)^\top x_n$  which we analyzed in Appendix D.1 To resolve this issue, we utilize the concept of covering number and derive a union bound.

Recall for a compact set  $\mathcal{Z}$  in a norm space, the covering number  $\mathcal{N}(\mathcal{Z}, \epsilon)$  with  $\epsilon > 0$  is the minimal number of  $\epsilon$ -balls that covers  $\mathcal{Z}$ . That is, there is a set  $\{z_i \in \mathcal{Z}\}_{i=1}^{\mathcal{N}(\mathcal{Z}, \epsilon)}$  such that  $\max_{z \in \mathcal{Z}} \min_{z' \in B(\mathcal{Z}, \epsilon)} \|z - z'\| \leq \epsilon$ . Usually the covering number  $\mathcal{N}(\mathcal{Z}, \epsilon)$  is polynomial in  $\epsilon$  and perhaps exponential in the ambient dimension of  $\mathcal{Z}$ .

The idea of covering number can be used to provide a union bound of concentration over compact sets, which we summarize as a lemma below.

**Lemma 15.** *Let  $f, g$  be two random  $L$ -Lipschitz functions. Suppose for some  $a > 0$  and some fixed  $z \in \mathcal{Z}$  selected independently of  $f, g$ , it holds*

$$P(|f(z) - g(z)| > \epsilon) \leq \exp(-a\epsilon^2)$$

Then it holds that

$$P\left(\sup_{z \in \mathcal{Z}} |f(z) - g(z)| > \epsilon\right) \leq \mathcal{N}\left(\mathcal{Z}, \frac{\epsilon}{4L}\right) \exp\left(\frac{-a\epsilon^2}{4}\right)$$

*Proof.* Let  $\mathcal{C}$  denote a set of covers of size  $\mathcal{N}(\mathcal{Z}, \frac{\epsilon}{4L})$ . Then, for any  $z \in \mathcal{Z}$  which could depend on  $f, g$ ,

$$\begin{aligned} |f(z) - g(z)| &\leq \min_{z' \in \mathcal{C}} |f(z) - f(z')| + |f(z') - g(z')| + |g(z') - g(z)| \\ &\leq \min_{z' \in \mathcal{C}} 2L\|z - z'\| + |f(z') - g(z')| \\ &\leq \frac{\epsilon}{2} + \max_{z' \in \mathcal{C}} |f(z') - g(z')| \end{aligned}$$

Thus,  $\sup_{z \in \mathcal{Z}} |f(z) - g(z)| > \epsilon \implies \max_{z' \in \mathcal{C}} |f(z') - g(z')| > \frac{\epsilon}{2}$ . Therefore, we have the union bound.

$$P\left(\sup_{z \in \mathcal{Z}} |f(z) - \mathbb{E}[f(z)]| > \epsilon\right) \leq \mathcal{N}\left(\mathcal{Z}, \frac{\epsilon}{4L}\right) \exp\left(\frac{-a\epsilon^2}{4}\right). \quad \square$$

We now use Lemma 15 to bound the component  $\sum_{n=1}^N (g_n - \nabla h_n(x_n))^\top y_N^*$ . We recall by definition, for  $x = (\mathbf{v}, \boldsymbol{\mu})$ ,

$$(\nabla h_n(x_n) - g_n)^\top x = (\kappa \mathbf{1} - \mathbf{a}_{\mathbf{v}_n} - \mathbf{g}_{n,\mu})^\top \boldsymbol{\mu} + (\mathbf{b}_{\boldsymbol{\mu}_n} - \mathbf{g}_{n,v})^\top \mathbf{v}$$

Because  $y_N^* = (\mathbf{v}^{\hat{\pi}_N}, \boldsymbol{\mu}^*)$ , we can write the sum of interest as

$$\sum_{n=1}^N (g_n - \nabla h_n(x_n))^\top y_N^* = \sum_{n=1}^N (\mathbf{g}_{n,\mu} - \kappa \mathbf{1} + \mathbf{a}_{\mathbf{v}_n})^\top \boldsymbol{\mu}^* + \sum_{n=1}^N (\mathbf{g}_{n,v} - \mathbf{b}_{\boldsymbol{\mu}_n})^\top \mathbf{v}^{\hat{\pi}_N}$$

For the first term, because  $\boldsymbol{\mu}^*$  is set beforehand by the MDP definition and does not depend on the randomness during learning, it is a martingale and we can apply the steps in Appendix D.1 to show,

$$\sum_{n=1}^N (\mathbf{g}_{n,\mu} - \kappa \mathbf{1} + \mathbf{a}_{\mathbf{v}_n})^\top \boldsymbol{\mu}^* = \bar{O}\left(\frac{\sqrt{NK \log(\frac{1}{\delta})}}{1 - \gamma}\right)$$

For the second term, because  $\mathbf{v}^{\hat{\pi}_N}$  depends on the randomness in the learning process, we need to use a union bound. Following the steps in Appendix D.1, we see that for some fixed  $\mathbf{v} \in \mathcal{V}$ , it holds

$$P\left(\left|\sum_{n=1}^N (\mathbf{g}_{n,v} - \mathbf{b}_{\boldsymbol{\mu}_n})^\top \mathbf{v}\right| > \epsilon\right) \leq \exp\left(-\frac{(1-\gamma)^2}{N} \epsilon^2\right)$$

where some constants were ignored for the sake of conciseness. Note also that it does not have the  $\sqrt{K}$  factor because of Lemma 8. To apply Lemma 15, we need to know the order of covering number of  $\mathcal{V}$ . Since  $\mathcal{V}$  is an  $|\mathcal{S}|$ -dimensional unit cube in the positive orthant, it is straightforward to show  $\mathcal{N}(\mathcal{V}, \epsilon) \leq \max(1, (1/\epsilon)^{|\mathcal{S}|})$  (by simply discretizing evenly in each dimension). Moreover, the functions  $\sum_{n=1}^N \mathbf{g}_{n,v}^\top \mathbf{v}$  and  $\sum_{n=1}^N \mathbf{b}_{\boldsymbol{\mu}_n}^\top \mathbf{v}$  are  $\frac{N}{1-\gamma}$ -Lipschitz in  $\|\cdot\|_\infty$ .

Applying Lemma 15 then gives us:

$$P\left(\sup_{v \in \mathcal{V}} \left|\sum_{n=1}^N (\mathbf{g}_{n,v} - \mathbf{b}_{\boldsymbol{\mu}_n})^\top \mathbf{v}\right| > \epsilon\right) \leq \mathcal{N}\left(\mathcal{V}, \frac{\epsilon(1-\gamma)}{4N}\right) \exp\left(-\frac{(1-\gamma)^2}{4N} \epsilon^2\right).$$

For a given  $\delta$ , we thus want to find the smallest  $\epsilon$  such that:

$$\delta \geq \mathcal{N} \left( \mathcal{V}, \frac{\epsilon(1-\gamma)}{4N} \right) \exp \left( -\frac{(1-\gamma)^2}{4N} \epsilon^2 \right).$$

That is:

$$\log\left(\frac{1}{\delta}\right) \leq \frac{(1-\gamma)^2}{4N} \epsilon^2 + |\mathcal{S}| \min(0, \log(\frac{\epsilon(1-\gamma)}{4N})).$$

Picking  $\epsilon = O \left( \log(N) \frac{\sqrt{N \log(\frac{1}{\delta})}}{1-\gamma} \right) = \tilde{O} \left( \frac{\sqrt{N \log(\frac{1}{\delta})}}{1-\gamma} \right)$  guarantees that the inequality is verified asymptotically.

Combining these two steps, we have shown overall, with probability at least  $1 - \delta$ ,

$$\sum_{n=1}^N (g_n - \nabla h_n(x_n))^\top y_N^* = \tilde{O} \left( \frac{\sqrt{N K \log(\frac{1}{\delta})}}{1-\gamma} \right).$$

#### D.4 Summary

In the previous subsections, we have provided high probability upper bounds for each term in the decomposition

$$\text{Regret}_N(y_N^*) \leq \left( \sum_{n=1}^N (\nabla h_n(x_n) - g_n)^\top x_n \right) + \left( \max_{x \in \mathcal{X}} \sum_{n=1}^N g_n^\top (x_n - x) \right) + \left( \sum_{n=1}^N (g_n - \nabla h_n(x_n))^\top y_N^* \right)$$

implying with probability at least  $1 - \delta$ ,

$$\text{Regret}_N(y_N^*) \leq \tilde{O} \left( \frac{\sqrt{N K \log(\frac{1}{\delta})}}{1-\gamma} \right) + \tilde{O} \left( \frac{\sqrt{K N}}{1-\gamma} + \frac{\sqrt{K^3 \log \frac{1}{\delta}}}{1-\gamma} \right) = \tilde{O} \left( \frac{\sqrt{N |\mathcal{S}| |\mathcal{A}| \log(\frac{1}{\delta})}}{1-\gamma} \right)$$

By Theorem 1, this would imply with probability at least  $1 - \delta$ ,

$$V^{\hat{\pi}_N}(p) \geq V^*(p) - \frac{\text{Regret}_N(y_N^*)}{N} \geq V^*(p) - \tilde{O} \left( \frac{\sqrt{|\mathcal{S}| |\mathcal{A}| \log(\frac{1}{\delta})}}{(1-\gamma)\sqrt{N}} \right)$$

In other words, the sample complexity of mirror descent to obtain an  $\epsilon$  approximately optimal policy (i.e.  $V^*(p) - V^{\hat{\pi}_N}(p) \leq \epsilon$ ) is at most  $\tilde{O} \left( \frac{|\mathcal{S}| |\mathcal{A}| \log(\frac{1}{\delta})}{(1-\gamma)^2 \epsilon^2} \right)$ .

## E Sample Complexity of Mirror Descent with Basis Functions

Here we provide further discussions on the sample complexity of running Algorithm 1 with linearly parameterized function approximators and the proof of Theorem 3.

**Theorem 3.** *Under a proper choice of  $\Theta$  and  $B_R$ , with probability  $1 - \delta$ , Algorithm 1 learns an  $(\epsilon + \epsilon_{\Theta, N})$ -optimal policy with  $\tilde{O} \left( \frac{\dim(\Theta) \log(\frac{1}{\delta})}{(1-\gamma)^2 \epsilon^2} \right)$  samples.*

### E.1 Setup

We suppose that the decision variable is parameterized in the form  $x_\theta = (\Phi \theta_v, \Psi \theta_\mu)$ , where  $\Phi, \Psi$  are given nonlinear basis functions and  $(\theta_v, \theta_\mu) \in \Theta$  are the parameters to learn. For modeling the value function, we suppose each column in  $\Phi$  is a vector (i.e. function) such that its  $\|\cdot\|_\infty$  is less than one. For modeling the state-action distribution, we suppose each column in  $\Psi$  is a state-action distribution from which we can draw samples. This choice implies that every column of  $\Phi$  belongs to  $\mathcal{V}$ , and every column of  $\Psi$  belongs to  $\mathcal{M}$ .



Considering the geometry of  $\Phi$  and  $\Psi$ , we consider a compact and convex parameter set

$$\Theta = \{\theta = (\boldsymbol{\theta}_v, \boldsymbol{\theta}_\mu) : \|\boldsymbol{\theta}_v\|_2 \leq \frac{C_v}{\sqrt{\dim(\boldsymbol{\theta}_v)}}, \boldsymbol{\theta}_\mu \geq 0, \|\boldsymbol{\theta}_\mu\|_1 = 1\}$$

where  $C_v < \infty$ . The constant  $C_v$  acts as a regularization in learning: if it is too small, the bias (captured as  $\epsilon_{\Theta, N}$  in Corollary 1 restated below) becomes larger; if it is too large, the learning becomes slower.

This choice of  $\Theta$  makes sure, for  $\theta = (\boldsymbol{\theta}_v, \boldsymbol{\theta}_\mu) \in \Theta$ ,  $\Psi\boldsymbol{\theta}_\mu \in \mathcal{M}$  and  $\|\Phi\boldsymbol{\theta}_v\|_\infty \leq \|\boldsymbol{\theta}_v\|_1 \leq C_v$ . Therefore, by the above construction, we can verify that the requirement in Corollary 1 is satisfied, i.e. for  $\theta = (\boldsymbol{\theta}_v, \boldsymbol{\theta}_\mu) \in \Theta$ , we have  $(\Phi\boldsymbol{\theta}_v, \Psi\boldsymbol{\theta}_\mu) \in \mathcal{X}_\Theta$ .

**Corollary 1.** *Let  $X_N = \{x_n \in \mathcal{X}_\theta\}_{n=1}^N$  be any sequence. Let  $\hat{\pi}_N$  be the policy given either by the average or the best decision in  $X_N$ . It holds that*

$$V^{\hat{\pi}_N}(p) \geq V^*(p) - \frac{\text{Regret}_N(\Theta)}{N} - \epsilon_{\Theta, N}$$

where  $\epsilon_{\Theta, N} = \min_{x_\theta \in \mathcal{X}_\theta} r_{ep}(\hat{x}_N; y_N^*) - r_{ep}(\hat{x}_N; x_\theta)$  measures the expressiveness of  $X_\theta$ , and  $\text{Regret}_N(\Theta) := \sum_{n=1}^N l_n(x_n) - \min_{x \in \mathcal{X}_\Theta} \sum_{n=1}^N l_n(x)$ .

## E.2 Online Loss and Sampled Gradient

Let  $\theta = (\boldsymbol{\theta}_v, \boldsymbol{\theta}_\mu) \in \Theta$ . In view of the parameterization above, we can identify the online loss in (23) in the parameter space as

$$h_n(\theta) := \mathbf{b}_{\boldsymbol{\mu}_n}^\top \Phi \boldsymbol{\theta}_v + \boldsymbol{\theta}_\mu^\top \Psi^\top \left( \frac{1}{1-\gamma} \mathbf{1} - \mathbf{a}_{\mathbf{v}_n} \right) \quad (33)$$

where we have the natural identification  $x_n = (\mathbf{v}_n, \boldsymbol{\mu}_n) = (\Phi\boldsymbol{\theta}_{v,n}, \Psi\boldsymbol{\theta}_{\mu,n})$  and  $\theta_n = (\boldsymbol{\theta}_{v,n}, \boldsymbol{\theta}_{\mu,n}) \in \Theta$  is the decision made by the online learner in the  $n$ th round. Note that because this extension of Algorithm 1 makes sure  $\|\boldsymbol{\theta}_{\mu,n}\|_1 = 1$  for every iteration, we can still use  $h_n$ . For writing convenience, we will continue to overload  $h_n$  as a function of parameter  $\theta$  in the following analyses.

Mirror descent requires gradient estimates of  $\nabla h_n(\theta_n)$ . Here we construct an unbiased stochastic estimate of  $\nabla h_n(\theta_n)$  as

$$g_n = \begin{bmatrix} \mathbf{g}_{n,v} \\ \mathbf{g}_{n,\mu} \end{bmatrix} = \begin{bmatrix} \Phi^\top (\tilde{\mathbf{p}}_n + \frac{1}{1-\gamma} (\gamma \tilde{\mathbf{P}}_n - \mathbf{E}_n)^\top \tilde{\boldsymbol{\mu}}_n) \\ \dim(\boldsymbol{\theta}_\mu) \hat{\Psi}_n^\top \left( \frac{1}{1-\gamma} \hat{\mathbf{1}}_n - \hat{\mathbf{r}}_n - \frac{1}{1-\gamma} (\gamma \hat{\mathbf{P}}_n - \hat{\mathbf{E}}_n) \mathbf{v}_n \right) \end{bmatrix} \quad (34)$$

using two calls of the generative model (again we overload the symbol  $g_n$  for the analyses in this section):

- The upper part  $\mathbf{g}_{n,v}$  is constructed similarly as before in (24): First we sample the initial state from the initial distribution, the state-action pair using the learned state-action distribution, and then the transition to state at the sampled state-action pair. We evaluate  $\Phi$ 's values at those samples to construct  $\mathbf{g}_{n,v}$ . Thus,  $\mathbf{g}_{n,v}$  would generally be a dense vector of size  $\dim(\boldsymbol{\theta}_v)$  (unless the columns of  $\Phi$  are sparse to begin with).
- The lower part  $\mathbf{g}_{n,\mu}$  is constructed slightly differently. Recall for the tabular version in (24), we uniformly sample over the state and action spaces. Here instead we first sample uniformly a column (i.e. a basis function) in  $\Psi$  and then sample a state-action pair according to the sampled column, which is a distribution by design. Therefore, the multiplier due to uniform sampling in the second row of (34) is now  $\dim(\boldsymbol{\theta}_\mu)$  rather than  $|\mathcal{S}||\mathcal{A}|$  in (24). The matrix  $\hat{\Psi}_n$  is extremely sparse, where only the single sampled entry (the column and the state-action pair) is one and the others are zero. In fact, one can think of the tabular version as simply using basis functions  $\Psi = \mathbf{I}$ , i.e. each column is a delta distribution. Under this identification, the expression in (34) matches the one in (24).

It is straightforward to verify that  $\mathbb{E}[g_n] = \nabla h_n(\theta_n)$  for  $g_n$  in (34).

### E.3 Proof of Theorem 3

We follow the same steps of the analysis of the tabular version. We will highlight the differences/improvement due to using function approximations.

First, we use Corollary 1 in place of Theorem 1. To properly handle the randomness, we revisit its derivation to slightly tighten the statement, which was simplified for the sake of cleaner exposition. Define

$$y_{N,\theta}^* = (\mathbf{v}_{N,\theta}^*, \boldsymbol{\mu}_\theta^*) := \arg \max_{x_\theta \in \mathcal{X}_\theta} r_{ep}(\hat{x}_N; x_\theta).$$

For writing convenience, let us also denote  $\theta_N^* = (\boldsymbol{\theta}_{v,N}^*, \boldsymbol{\theta}_\mu^*) \in \Theta$  as the corresponding parameter of  $y_{N,\theta}^*$ . We remark that  $\boldsymbol{\mu}_\theta^*$  (i.e.  $\boldsymbol{\theta}_\mu^*$ ), which tries to approximate  $\boldsymbol{\mu}^*$ , is fixed before the learning process, whereas  $\mathbf{v}_{N,\theta}^*$  (i.e.  $\boldsymbol{\theta}_{v,N}^*$ ) could depend on the stochasticity in the learning. Using this new notation and the steps in the proof of Corollary 1, we can write

$$\begin{aligned} V^*(p) - V^{\hat{\pi}_N}(p) &= r_{ep}(\hat{x}_N; y_N^*) \\ &= \epsilon_{\Theta,N} + r_{ep}(\hat{x}_N; y_{N,\theta}^*) \leq \epsilon_{\Theta,N} + \frac{\text{Regret}_N(y_{N,\theta}^*)}{N} \end{aligned}$$

where the first equality is Proposition 4, the last inequality follows the proof of Theorem 1, and we recall the definition  $\epsilon_{\Theta,N} = r_{ep}(\hat{x}_N; y_N^*) - r_{ep}(\hat{x}_N; y_{N,\theta}^*)$ .

The rest of the proof is very similar to that of Theorem 1, because linear parameterization does not change the convexity of the loss sequence. Let  $y_N^* = (\mathbf{v}^{\hat{\pi}_N}, \boldsymbol{\mu}^*)$ . We bound the regret by the following rearrangement.

$$\begin{aligned} \text{Regret}_N(y_{N,\theta}^*) &= \sum_{n=1}^N l_n(x_n) - \sum_{n=1}^N l_n(y_{N,\theta}^*) \\ &= \sum_{n=1}^N h_n(\theta_n) - \sum_{n=1}^N h_n(\theta_N^*) \\ &= \sum_{n=1}^N \nabla h_n(\theta_n)^\top (\theta_n - \theta_N^*) \\ &= \left( \sum_{n=1}^N (\nabla h_n(\theta_n) - g_n)^\top \theta_n \right) + \left( \sum_{n=1}^N g_n^\top (\theta_n - \theta_N^*) \right) + \left( \sum_{n=1}^N (g_n - \nabla h_n(\theta_n))^\top \theta_N^* \right) \\ &\leq \left( \sum_{n=1}^N (\nabla h_n(\theta_n) - g_n)^\top \theta_n \right) + \left( \max_{\theta \in \Theta} \sum_{n=1}^N g_n^\top (\theta_n - \theta) \right) + \left( \sum_{n=1}^N (g_n - \nabla h_n(\theta_n))^\top \theta_N^* \right) \end{aligned} \quad (35)$$

where the second equality is due to the identification in (33).

We will solve this online learning problem with mirror descent

$$\theta_{n+1} = \arg \min_{\theta \in \Theta} \langle g_n, \theta \rangle + \frac{1}{\eta} B_R(\theta \| \theta_n) \quad (36)$$

with step size  $\eta > 0$  and a Bregman divergence that is a straightforward extension of (22)

$$B_R(\theta' \| \theta) = \frac{1}{2} \frac{\dim(\theta_v)}{C_v^2} \|\theta'_v - \theta_v\|_2^2 + KL(\boldsymbol{\theta}'_\mu \| \boldsymbol{\theta}_\mu) \quad (37)$$

where the constant  $\frac{\dim(\theta_v)}{C_v^2}$  is chosen to make the size of Bregman divergence dimension-free (at least up to log factors). Below we analyze the size of the three terms in (35) like what we did for Theorem 2.

### E.4 The First Term: Martingale Concentration

The first term is a martingale. We will use this part to highlight the different properties due to using basis functions. The proof follows the steps in Appendix D.1, but now the martingale difference of interest is instead

$$\begin{aligned} M_n - M_{n-1} &= (\nabla h_n(\theta_n) - g_n)^\top \theta_n \\ &= (\Psi^\top (\kappa \mathbf{1} - \mathbf{a}_{v_n}) - \mathbf{g}_{n,\mu})^\top \boldsymbol{\theta}_{\mu,n} + (\Phi^\top \mathbf{b}_{\mu_n} - \mathbf{g}_{n,v})^\top \boldsymbol{\theta}_{v,n} \end{aligned}$$

They now have nicer properties due to the way  $\mathbf{g}_{n,\mu}$  is sampled.

For the first term  $(\Psi^\top(\kappa\mathbf{1} - \mathbf{a}_{\mathbf{v}_n}) - \mathbf{g}_{n,\mu})^\top \boldsymbol{\theta}_\mu$ , we use the lemma below, where we recall the filtration  $F_n$  is naturally defined as  $\{\theta_1, \dots, \theta_n\}$ .

**Lemma 16.** *Let  $\theta = (\theta_v, \boldsymbol{\theta}_\mu) \in \Theta$  be arbitrary that is chosen independent of the randomness of  $\mathbf{g}_{n,\mu}$  when  $F_{n-1}$  is given. Then it holds  $|(\kappa\mathbf{1} - \mathbf{a}_{\mathbf{v}_n} - \mathbf{g}_{n,\mu})^\top \boldsymbol{\theta}| \leq \frac{2(1+\dim(\boldsymbol{\theta}_\mu))}{1-\gamma}$  and  $\mathbb{V}_{|F_{n-1}}[(\kappa\mathbf{1} - \mathbf{a}_{\mathbf{v}_n} - \mathbf{g}_{n,\mu})^\top \boldsymbol{\theta}_n] \leq \frac{4\dim(\boldsymbol{\theta}_\mu)}{(1-\gamma)^2}$ .*

*Proof.* By triangular inequality,

$$|(\Psi^\top(\kappa\mathbf{1} - \mathbf{a}_{\mathbf{v}_n}) - \mathbf{g}_{n,\mu})^\top \boldsymbol{\theta}_\mu| \leq |(\kappa\mathbf{1} - \mathbf{a}_{\mathbf{v}_n})^\top \Psi \boldsymbol{\theta}_\mu| + |\mathbf{g}_{n,\mu}^\top \boldsymbol{\theta}_\mu|$$

For the deterministic part, using Lemma 5 and Hölder's inequality,

$$|(\kappa\mathbf{1} - \mathbf{a}_{\mathbf{v}_n})^\top \Psi \boldsymbol{\theta}_\mu| \leq \kappa + \|\mathbf{a}_{\mathbf{v}_n}\|_\infty \|\Psi \boldsymbol{\theta}_\mu\|_1 \leq \frac{2}{1-\gamma}$$

For the stochastic part, let  $k_n$  denote the sampled column index,  $i_n$  be index of the sampled state-action pair using the column of  $k_n$ , and  $j_n$  be the index of the transited state sampled at the pair given by  $i_n$ . With abuse of notation, we will use  $i_n$  to index both  $\mathcal{S} \times \mathcal{A}$  and  $\mathcal{S}$ . Let  $\boldsymbol{\mu} = \Psi \boldsymbol{\theta}_\mu$ . With this notation, we may derive

$$\begin{aligned} |\mathbf{g}_{n,\mu}^\top \boldsymbol{\theta}_\mu| &= |\dim(\boldsymbol{\theta}_\mu) \boldsymbol{\theta}_\mu^\top \hat{\Psi}_n^\top (\kappa \hat{\mathbf{1}}_n - \hat{\mathbf{r}}_n - \frac{1}{1-\gamma} (\gamma \hat{\mathbf{P}}_n - \hat{\mathbf{E}}_n) \mathbf{v}_n)| \\ &= \dim(\boldsymbol{\theta}_\mu) \theta_{\mu,k_n} |\kappa - r_{i_n} - \frac{\gamma v_{n,j_n} - v_{n,i_n}}{1-\gamma}| \\ &\leq \frac{2\dim(\boldsymbol{\theta}_\mu) \theta_{\mu,k_n}}{1-\gamma} \leq \frac{2\dim(\boldsymbol{\theta}_\mu)}{1-\gamma} \end{aligned}$$

where we use the facts  $r_{i_n}, v_{n,j_n}, v_{n,i_n} \in [0, 1]$  and  $\theta_{\mu,k_n} \leq 1$ .

Let  $\psi_\mu^{(k)}$  denote the  $k$ th column of  $\Psi$ . For  $\mathbb{V}_{|F_{n-1}}[(\kappa\mathbf{1} - \mathbf{a}_{\mathbf{v}_n} - \mathbf{g}_{n,\mu})^\top \boldsymbol{\theta}_n]$ , we can write it as

$$\begin{aligned} \mathbb{V}_{|F_{n-1}}[(\Psi^\top(\kappa\mathbf{1} - \mathbf{a}_{\mathbf{v}_n}) - \mathbf{g}_{n,\mu})^\top \boldsymbol{\theta}_\mu] &= \mathbb{V}_{|F_{n-1}}[\mathbf{g}_{n,\mu}^\top \boldsymbol{\theta}_\mu] \\ &\leq \mathbb{E}_{|F_{n-1}}[|\mathbf{g}_{n,\mu}^\top \boldsymbol{\theta}_\mu|^2] \\ &= \sum_{k_n} \frac{1}{\dim(\boldsymbol{\theta}_\mu)} \sum_{i_n} \psi_{\mu,i_n}^{(k_n)} \mathbb{E}_{j_n|i_n} \left[ \dim(\boldsymbol{\theta}_\mu)^2 \theta_{\mu,k_n}^2 \left( \kappa - r_{i_n} - \frac{\gamma v_{n,j_n} - v_{n,i_n}}{1-\gamma} \right)^2 \right] \\ &\leq \frac{4\dim(\boldsymbol{\theta}_\mu)}{(1-\gamma)^2} \sum_{k_n} \theta_{\mu,k_n}^2 \sum_{i_n} \psi_{\mu,i_n}^{(k_n)} \\ &\leq \frac{4\dim(\boldsymbol{\theta}_\mu)}{(1-\gamma)^2} \left( \sum_{k_n} \theta_{\mu,k_n} \right)^2 \leq \frac{4\dim(\boldsymbol{\theta}_\mu)}{(1-\gamma)^2} \end{aligned}$$

where in the second inequality we use the fact that  $|\kappa - r_{i_n} - \frac{\gamma v_{n,j_n} - v_{n,i_n}}{1-\gamma}| \leq \frac{2}{1-\gamma}$ .  $\square$

For the second term  $(\Phi^\top \mathbf{b}_{\boldsymbol{\mu}_n} - \mathbf{g}_{n,v})^\top \boldsymbol{\theta}_{v,n}$ , we use this lemma.

**Lemma 17.** *Let  $\boldsymbol{\theta} \in \mathcal{V}$  be arbitrary that is chosen independent of the randomness of  $\mathbf{g}_{n,v}$  when  $F_{n-1}$  is given. Then it holds  $|(\Phi^\top \mathbf{b}_{\boldsymbol{\mu}_n} - \mathbf{g}_{n,v})^\top \boldsymbol{\theta}| \leq \frac{4C_v}{1-\gamma}$  and  $\mathbb{V}_{|F_{n-1}}[(\Phi^\top \mathbf{b}_{\boldsymbol{\mu}_n} - \mathbf{g}_{n,v})^\top \boldsymbol{\theta}] \leq \frac{4C_v^2}{(1-\gamma)^2}$ .*

*Proof.* We appeal to Lemma 5, which shows  $\|\mathbf{b}_{\boldsymbol{\mu}_n}\|_1 \leq \frac{2}{1-\gamma}$  and

$$\|\tilde{\mathbf{p}}_n + \frac{1}{1-\gamma} (\gamma \tilde{\mathbf{P}}_n - \mathbf{E}_n)^\top \tilde{\boldsymbol{\mu}}_n\|_1 \leq \frac{2}{1-\gamma}$$

Therefore, overall we can derive

$$|(\Phi^\top \mathbf{b}_{\boldsymbol{\mu}_n} - \mathbf{g}_{n,v})^\top \boldsymbol{\theta}| \leq \left( \|\mathbf{b}_{\boldsymbol{\mu}_n}\|_1 + \|\tilde{\mathbf{p}}_n + \frac{1}{1-\gamma} (\gamma \tilde{\mathbf{P}}_n - \mathbf{E}_n)^\top \tilde{\boldsymbol{\mu}}_n\|_1 \right) \|\Phi \boldsymbol{\theta}_v\|_\infty \leq \frac{4C_v}{1-\gamma}$$

where we use again each column in  $\Phi$  has  $\|\cdot\|_\infty$  less than one, and  $\|\cdot\|_\infty \leq \|\cdot\|_2$ . Similarly, for the variance, we can write

$$\mathbb{V}_{|F_{n-1}}[(\Phi^\top \mathbf{b}_{\mu_n} - \mathbf{g}_{n,v})^\top \theta] = \mathbb{V}_{|F_{n-1}}[\mathbf{g}_{n,v}^\top \theta] \leq \mathbb{E}_{|F_{n-1}}[(\mathbf{g}_{n,v}^\top \theta)^2] \leq \frac{4C_v^2}{(1-\gamma)^2} \quad \square$$

From the above two lemmas, we see the main difference from the what we had in Appendix D.1 for the tabular case is that, the martingale difference now scales in  $O\left(\frac{C_v + \dim(\theta_\mu)}{1-\gamma}\right)$  instead of  $O\left(\frac{|\mathcal{S}||\mathcal{A}|}{1-\gamma}\right)$ , and its variance scales in  $O\left(\frac{C_v^2 + \dim(\theta_\mu)}{(1-\gamma)^2}\right)$  instead of  $O\left(\frac{|\mathcal{S}||\mathcal{A}|^2}{(1-\gamma)^2}\right)$ . We note the constant  $C_v$  is universal, independent of the problem size.

Following the similar steps in Appendix D.1, these new results imply that

$$P\left(\sum_{n=1}^N (\nabla h_n(\theta_n) - g_n)^\top \theta_n > \epsilon\right) \leq \exp\left(\frac{-\epsilon^2}{2N\sigma^2\left(1 + \frac{b\epsilon}{3N\sigma^2}\right)}\right)$$

with  $b = O\left(\frac{C_v + \dim(\theta_\mu)}{1-\gamma}\right)$  and  $O\left(\frac{C_v^2 + \dim(\theta_\mu)}{(1-\gamma)^2}\right)$ . This implies that, with probability at least  $1 - \delta$ , it hold

$$\sum_{n=1}^N (\nabla h_n(\theta_n) - g_n)^\top \theta_n = \tilde{O}\left(\frac{\sqrt{N(C_v^2 + \dim(\theta_\mu)) \log\left(\frac{1}{\delta}\right)}}{1-\gamma}\right)$$

## E.5 Static Regret of Mirror Descent

Again the steps here are very similar to those in Appendix D.2. We concern bounding the static regret.

$$\max_{\theta \in \Theta} \sum_{n=1}^N g_n^\top (\theta_n - \theta)$$

From Appendix D.2, we recall this can be achieved by the mirror descent's optimality condition. The below inequality is true, for any  $\theta' \in \Theta$ :

$$\sum_{n=1}^N \langle g_n, \theta_n - \theta' \rangle \leq \frac{1}{\eta} B_R(\theta' | \theta_1) + \sum_{n=1}^N \langle g_n, \theta_{n+1} - \theta_n \rangle - \frac{1}{\eta} B_R(\theta_{n+1} | \theta_n)$$

Based on our choice of Bregman divergence given in (37), i.e.

$$B_R(\theta' | \theta) = \frac{1}{2} \frac{\dim(\theta_v)}{C_v^2} \|\theta'_v - \theta_v\|_2^2 + KL(\theta'_\mu | \theta_\mu), \quad (37)$$

we have  $\frac{1}{\eta} B_R(\theta' | \theta_1) \leq \frac{\tilde{O}(1)}{\eta}$ . For each  $\langle g_n, \theta_{n+1} - \theta_n \rangle - \frac{1}{\eta} B_R(\theta_{n+1} | \theta_n)$ , we will use again the two basic lemmas we proved in Appendix D.2.

**Lemma 10.** *For any vector  $x, y, g$  and scalar  $\eta > 0$ , it holds  $\langle g, x - y \rangle - \frac{1}{2\eta} \|x - y\|_2^2 \leq \frac{\eta \|g\|_2^2}{2}$ .*

**Lemma 11.** *Suppose  $B_R(x|y) = KL(x|y)$  and  $x, y$  are probability distributions, and  $g \geq 0$  element-wise. Then, for  $\eta > 0$ ,*

$$-\frac{1}{\eta} B_R(y|x) + \langle g, x - y \rangle \leq \frac{\eta}{2} \sum_i x_i (g_i)^2 = \frac{\eta}{2} \|g\|_x^2.$$

Thus, we have the upper bound

$$\langle g_n, \theta_{n+1} - \theta_n \rangle - \frac{1}{\eta} B_R(\theta_{n+1} | \theta_n) = \frac{C_v^2}{\dim(\theta_v)} \frac{\eta \|\mathbf{g}_{n,v}\|_2^2}{2} + \frac{\eta \|\mathbf{g}_{n,\mu}\|_{\theta_{\mu,n}}^2}{2}$$

Together with the upper bound on  $\frac{1}{\eta}B_R(x'|x_1)$ , it implies that

$$\begin{aligned} \sum_{n=1}^N \langle g_n, x_n - x' \rangle &\leq \frac{1}{\eta}B_R(x'|x_1) + \sum_{n=1}^N \langle g_n, x_{n+1} - x_n \rangle - \frac{1}{\eta}B_R(x_{n+1}|x_n) \\ &\leq \frac{\tilde{O}(1)}{\eta} + \frac{\eta}{2} \sum_{n=1}^N \frac{C_v^2}{\dim(\theta_v)} \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\theta_{\mu,n}}^2 \end{aligned} \quad (38)$$

We can expect, with high probability,  $\sum_{n=1}^N \frac{C_v^2}{\dim(\theta_v)} \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\theta_{\mu,n}}^2$  concentrates toward its expectation, i.e.

$$\sum_{n=1}^N \frac{C_v^2}{\dim(\theta_v)} \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\theta_{\mu,n}}^2 \leq \sum_{n=1}^N \mathbb{E} \left[ \frac{C_v^2}{\dim(\theta_v)} \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\theta_{\mu,n}}^2 \right] + o(N)$$

To bound the right-hand side, we will use the upper bounds below, which largely follow the proof of Lemma 16 and Lemma 17.

**Lemma 18.**  $\mathbb{E}[\|\mathbf{g}_{n,v}\|_2^2] \leq \frac{4\dim(\theta_v)}{(1-\gamma)^2}$  and  $\mathbb{E}[\|\mathbf{g}_{n,\mu}\|_{\theta_{\mu,n}}^2] \leq \frac{4\dim(\theta_\mu)}{(1-\gamma)^2}$ .

**Lemma 19.**  $\|\mathbf{g}_{n,v}\|_2^2 \leq \frac{4\dim(\theta_v)}{(1-\gamma)^2}$  and  $\|\mathbf{g}_{n,\mu}\|_{\theta_{\mu,n}}^2 \leq \frac{4\dim(\theta_\mu)}{(1-\gamma)^2}$ .

By Azuma-Hoeffding's inequality in Lemma 13,

$$\begin{aligned} \sum_{n=1}^N \frac{C_v^2}{\dim(\theta_v)} \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\theta_{\mu,n}}^2 &\leq \sum_{n=1}^N \mathbb{E} \left[ \frac{C_v^2}{\dim(\theta_v)} \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\theta_{\mu,n}}^2 \right] + O \left( \frac{C_v^2 + \dim(\theta_\mu)^2}{(1-\gamma)^2} \sqrt{N \log \left( \frac{1}{\delta} \right)} \right) \\ &\leq O \left( \frac{C_v^2 + \dim(\theta_\mu)}{(1-\gamma)^2} N \right) + O \left( \frac{C_v^2 + \dim(\theta_\mu)^2}{(1-\gamma)^2} \sqrt{N \log \left( \frac{1}{\delta} \right)} \right) \end{aligned}$$

Now we suppose we set  $\eta = O \left( \frac{1-\gamma}{\sqrt{N(C_v^2 + \dim(\theta_\mu))}} \right)$ . We have

$$\sum_{n=1}^N \langle g_n, \theta_n - \theta' \rangle \leq \frac{\tilde{O}(1)}{\eta} + \frac{\eta}{2} \sum_{n=1}^N \frac{C_v^2}{\dim(\theta_v)} \|\mathbf{g}_{n,v}\|_2^2 + \|\mathbf{g}_{n,\mu}\|_{\theta_{\mu,n}}^2 \leq \tilde{O} \left( \frac{\sqrt{(C_v^2 + \dim(\theta_\mu))N}}{1-\gamma} \right)$$

### E.5.1 Union Bound

Lastly we use an union bound to handle the term

$$\sum_{n=1}^N (g_n - \nabla h_n(\theta_n))^\top \theta_N^*$$

We follow the steps in Appendix D.3: we will use again the fact that  $\theta_N^* = (\theta_{v,N}^*, \theta_\mu^*) \in \Theta$ , so we can handle the part with  $\theta_\mu^*$  using the standard martingale concentration, and the part with  $\theta_{v,N}^*$  using the union bound.

Using the previous analyses, we see can first show that the martingale due to the part  $\theta_\mu^*$  concentrates in  $\tilde{O} \left( \frac{\sqrt{N \dim(\theta_\mu) \log(\frac{1}{\delta})}}{1-\gamma} \right)$ . Likewise, using the union bound, we can show the martingale due to the part  $\theta_{v,N}^*$  concentrates in  $\tilde{O} \left( \frac{\sqrt{N C_v^2 \log(\frac{N}{\delta})}}{1-\gamma} \right)$  where  $\mathcal{N}$  some proper the covering number of the set  $\left\{ \theta_v : \|\theta_v\|_2 \leq \frac{C_v}{\sqrt{\dim(\theta_v)}} \right\}$ . Because  $\log \mathcal{N} = O(\dim(\theta_v))$  for an Euclidean ball. We can combine the two bounds and show together

$$\sum_{n=1}^N (g_n - \nabla h_n(\theta_n))^\top \theta_N^* = \tilde{O} \left( \frac{\sqrt{N(C_v^2 \dim(\theta_v) + \dim(\theta_\mu)) \log(\frac{1}{\delta})}}{1-\gamma} \right)$$

### E.5.2 Summary

Combining the results of the three parts above, we have, with probability  $1 - \delta$ ,

$$\begin{aligned}
 & \text{Regret}_N(y_{N,\theta}^*) \\
 & \leq \left( \sum_{n=1}^N (\nabla h_n(\theta_n) - g_n)^\top \theta_n \right) + \left( \max_{\theta \in \Theta} \sum_{n=1}^N g_n^\top (\theta_n - \theta) \right) + \left( \sum_{n=1}^N (g_n - \nabla h_n(\theta_n))^\top \theta_n^* \right) \\
 & = \tilde{O} \left( \frac{\sqrt{N(\dim(\boldsymbol{\theta}_\mu) + C_v^2) \log(\frac{1}{\delta})}}{1 - \gamma} \right) + \tilde{O} \left( \frac{\sqrt{(C_v^2 + \dim(\boldsymbol{\theta}_\mu))N}}{1 - \gamma} \right) + \tilde{O} \left( \frac{\sqrt{N(C_v^2 \dim(\boldsymbol{\theta}_v) + \dim(\boldsymbol{\theta}_\mu)) \log(\frac{1}{\delta})}}{1 - \gamma} \right) \\
 & = \tilde{O} \left( \frac{\sqrt{N \dim(\Theta) \log(\frac{1}{\delta})}}{1 - \gamma} \right)
 \end{aligned}$$

where the last step is due to  $C_v$  is a universal constant. Or equivalently, the above bounds means a sample complexity in  $\tilde{O} \left( \frac{\dim(\Theta) \log(\frac{1}{\delta})}{(1-\gamma)^2 \epsilon^2} \right)$ . Finally, we recall the policy performance has a bias  $\epsilon_{\Theta,N}$  in Corollary 1 due to using function approximators. Considering this effect, we have the final statement.