

# Enhanced UV radiation and dense clumps in Mrk 231's molecular outflow

Claudia Cicone <sup>\*1,2</sup>, Roberto Maiolino<sup>3,4</sup>, Susanne Aalto<sup>5</sup>, Sebastien Muller<sup>5</sup>, and Chiara Feruglio<sup>6</sup>

<sup>1</sup> Institute of Theoretical Astrophysics, University of Oslo, P.O. Box 1029, Blindern, 0315 Oslo, Norway

<sup>2</sup> INAF-Osservatorio Astronomico di Brera, via Brera 28, 20121, Milan, Italy

<sup>3</sup> Cavendish Laboratory, University of Cambridge, 19 J. J. Thomson Ave, Cambridge CB3 0HE, UK

<sup>4</sup> Kavli Institute for Cosmology, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

<sup>5</sup> Department of Space, Earth and Environment, Chalmers University of Technology, Onsala Observatory, SE-439 92 Onsala, Sweden

<sup>6</sup> INAF - Osservatorio Astronomico di Trieste, via G.B. Tiepolo 11, 34143 Trieste, Italy

Received 27-Sep-2019 / Accepted: 25-Nov-2019

## ABSTRACT

We present interferometric observations of the CN(1-0) line emission in Mrk 231 and combine them with previous observations of CO and other H<sub>2</sub> gas tracers to study the physical properties of the massive molecular outflow. We find a strong boost of the CN/CO(1-0) line luminosity ratio in the outflow, which is unprecedented compared to any other known Galactic or extragalactic source. For the dense gas phase in the outflow traced by the HCN and CN emissions, we infer  $X_{\text{CN}} \equiv [\text{CN}]/[\text{H}_2] > X_{\text{HCN}}$  by at least a factor of three, with H<sub>2</sub> gas densities of  $n_{\text{H}_2} \sim 10^{5-6} \text{ cm}^{-3}$ . In addition, for the first time, we resolve narrow spectral features in the HCN(1-0) and HCO<sup>+</sup>(1-0) high-velocity line wings tracing the dense phase of the outflow. The velocity dispersions of these spectral features,  $\sigma_v \sim 7 - 20 \text{ km s}^{-1}$ , are consistent with those of massive extragalactic giant molecular clouds detected in nearby starburst nuclei. The H<sub>2</sub> gas masses inferred from the HCN data are quite high,  $M_{\text{mol}} \sim 0.3 - 5 \times 10^8 M_{\odot}$ . Our results suggest that massive, denser molecular gas complexes survive embedded into the more diffuse H<sub>2</sub> phase of the outflow, and that the chemistry of such outflowing dense clouds is affected by enhanced UV radiation.

**Key words.** galaxies:active — galaxies: evolution — galaxies: individual (Mrk 231) — galaxies: interactions — galaxies: ISM — submillimeter: ISM

## 1. Introduction

Mrk 231 is a late-stage merger, classified as ultra-luminous infrared galaxy (ULIRG) ( $L_{\text{IR}}(8 - 1000\mu\text{m}) = 1.3 \times 10^{46} \text{ erg s}^{-1}$ ), likely on its path to becoming an elliptical galaxy (Lipari et al. 2005). At least ~30% of the IR luminosity is due to dust heated by an active galactic nucleus (AGN) (Nardini et al. (2010), but Veilleux et al. (2009) estimate a higher AGN contribution). Despite its weak emission in the hard X-ray band (Teng et al. 2014; Piconcelli et al. 2013; Braitto et al. 2004), the elevated AGN bolometric luminosity ( $L_{\text{AGN}} \sim 8 \times 10^{45} \text{ erg s}^{-1}$ , Leighly et al. 2014) makes Mrk 231 the closest quasar known. However, most of the IR luminosity traces a young starburst (age < 250 Myr, Davies et al. 2007), forming stars at a rate of  $\text{SFR} \gtrsim 200 M_{\odot} \text{ yr}^{-1}$  (Taylor et al. 1999), fed by  $\gtrsim 10^9 M_{\odot}$  of molecular (H<sub>2</sub>) gas distributed in a kpc-size disk (Downes & Solomon 1998).

The multi-wavelength and multi-scale emission of Mrk 231 is overwhelmingly affected by *explosive* phenomena, which are a manifestation of feedback from the star formation and quasar activity, as predicted by theoretical models (Chevalier & Clegg 1985; Silk & Rees 1998; Murray et al. 2005; Faucher-Giguère & Quataert 2012; Zubovas & King 2012, 2014; Nims et al. 2015; Thompson et al. 2016; Ishibashi et al. 2018) and ‘observed’ in hydrodynamical simulations (Di Matteo et al. 2005; Sijacki et al. 2007; Costa et al. 2015, 2018; Biernacki & Teyssier 2018; Richings & Faucher-Giguère 2018; Schneider et al. 2018). On nuclear scales, Feruglio et al. (2015) reported observations of a

possible X-ray ultra fast wind with  $v = 20,000 \text{ km s}^{-1}$ . Broad absorption lines (BALs) were detected in both low and high ionisation species, classifying Mrk 231 as a FeLoBAL quasar (Lipari et al. 2009). The high-velocity material responsible for the BALs is believed to dominate even the Ly $\alpha$ , CIV, and CIII] emission lines, all showing spectral profiles blueshifted by  $-3500 \text{ km s}^{-1}$  (Veilleux et al. 2013). Besides the BAL winds, which are confined within  $r < 10 \text{ pc}$ , Mrk 231 hosts one of the best-studied *galaxy-scale* outflows, revealed for the first time in OH absorption by Fischer et al. (2010) and in CO emission by Feruglio et al. (2010). Using these earlier observations it was estimated that the outflow entrains  $\sim 3 \times 10^8 M_{\odot}$  of H<sub>2</sub> gas and extends out to at least  $r \sim 1 \text{ kpc}$  into the interstellar medium (ISM) (Cicone et al. 2012). Blue-shifted NaID (Rupke & Veilleux 2011; Lipari et al. 2009) and H<sub>1</sub> 21cm (Morganti et al. 2016; Teng et al. 2013) absorption features have also been detected, and they trace the H<sub>1</sub> gas component of the outflow.

At present, most galaxy-scale molecular outflows have been studied exclusively through low-J CO transitions (Cicone et al. 2014; Pereira-Santaella et al. 2018; Fluetsch et al. 2019). Such lines, thanks to their low critical density and high abundance of CO in the ISM (provided CO is sufficiently shielded), are ubiquitous tracers of H<sub>2</sub> gas, including the diffuse envelopes of molecular clouds that do not host active star formation. On the other hand, transitions from species such as HCN, HCO<sup>+</sup>, and HNC, because of their higher critical density, should arise mainly from the denser regions of molecular complexes associated with star formation sites (Gao & Solomon 2004). There-

\* claudia.cicone@astro.uio.no

fore, these molecules (including CN, see below) are commonly referred to as ‘dense gas’ tracers, and we will adopt this terminology throughout the text. However, we note that a significant portion of the emission from such tracers may still be associated with gas that has a density much lower than the nominal critical density ( $n_{\text{crit}}$ ) of the corresponding molecular transition. Indeed,  $n_{\text{crit}}$  is calculated analytically using the optically thin approximation and so it does not account for radiative trapping (Scoville & Solomon 1974), which can lower the volume density needed to excite the line by more than one order of magnitude with respect to  $n_{\text{crit}}$ . To deal with line trapping, Shirley (2015) propose to use the ‘effective excitation density’ ( $n_{\text{eff}}$ ), empirically defined as the gas density that produces a detectable integrated line flux arbitrarily set equal to  $\int T_R dv = 1 \text{ K km s}^{-1}$ , where  $T_R$  is the radiation temperature<sup>1</sup>. Furthermore, the  $n_{\text{crit}}$  computation in general assumes molecular hydrogen ( $\text{H}_2$ ) as the only collisional partner, hence neglecting collisions with electrons and atomic hydrogen. Collisions with electrons may become relevant in a medium where a large fraction of Carbon is ionised but Hydrogen is in the molecular phase, leading to the excitation of molecules such as HCN even in lower density  $\text{H}_2$  gas ( $n_{\text{H}_2} \sim 10^3 \text{ cm}^{-3}$ , e.g. Goldsmith & Kauffmann (2017); Kauffmann et al. (2017)).

The first detection of HCN,  $\text{HCO}^+$ , and HNC in the outflow of Mrk 231 was reported by Aalto et al. (2012a). Up to now, this outflow has been studied in eight different species, four detected in absorption by *Herschel* (e.g. OH,  $^{18}\text{OH}$ ,  $\text{H}_2\text{O}$ ,  $\text{OH}^+$ , Fischer et al. 2010; Sturm et al. 2011; González-Alfonso et al. 2014, 2018), and four imaged in emission using (sub-)millimetre interferometry (e.g. CO, HCN, HNC,  $\text{HCO}^+$ , see Feruglio et al. 2010; Cicone et al. 2012; Aalto et al. 2012b, 2015; Alatalo 2015; Feruglio et al. 2015; Lindberg et al. 2016).

In this paper we report the detection of exceptionally bright Cyanide Radical (CN)  $N = 1 - 0$  emission arising from the outflow of Mrk 231. CN is a tracer of high density molecular gas (for  $T_k = 100 \rightarrow 10 \text{ K}$ ,  $n_{\text{crit}} \simeq 6.4 \times 10^4 \rightarrow 4.1 \times 10^5 \text{ cm}^{-3}$  and  $n_{\text{eff}} \simeq 4.6 \times 10^3 \rightarrow 3.8 \times 10^4 \text{ cm}^{-3}$ , assuming a reference column density of  $N_{\text{CN}} = 10^{14} \text{ cm}^{-2}$ , Shirley (2015)). The production of CN is aided by the presence of moderate UV fields, either via the reaction  $\text{CH} + \text{N} \rightarrow \text{CN} + \text{H}$ , since CH is formed rapidly in a medium where both  $\text{C}^+$  and  $\text{H}_2$  are abundant (Fuente et al. 1993; Meier et al. 2015), or via the photodissociation of HCN:  $\text{HCN} + \gamma_{\text{UV}} \rightarrow \text{CN} + \text{H}$  (Boger & Sternberg 2005). For these reasons, CN is expected to be enhanced in photon dominated regions (PDRs), i.e. portions of the clouds where the chemistry is dominated by the interactions with UV photons (e.g. Aalto et al. 2002; Rodríguez-Franco et al. 1998), but possibly also in the presence of X-rays and/or cosmic rays (Meijerink et al. 2007; García-Burillo et al. 2010; Harada et al. 2015). Observations of Galactic star forming regions have shown that the mere presence of dense molecular gas is not itself a prior for enhanced CN emission. Instead, CN tends to be brighter on the surfaces of molecular clouds exposed to interstellar UV radiation (PDRs), but also inside molecular clouds embedding young massive stars (Yamagishi et al. 2018). Notably, CN remains very sensitive to variations in star formation activity even when its emission is averaged across kpc-scale regions in extragalactic sources (Watanabe et al. 2014).

In this paper we also present the first detection of spectrally-resolved features in the broad wings of the HCN(1-0) and  $\text{HCO}^+(1-0)$  emission lines. Their velocity dispersions are con-

sistent with individual massive molecular clouds entrained in the outflow. We discuss their properties and compare them with known Galactic and extragalactic giant molecular clouds (GMCs).

The paper is organised as follows. The observations are described in § 2. In § 3.1 we show the CN(1-0) data and combine them with previous (sub-)millimetre observations of other molecular lines to study the physical properties of the outflowing  $\text{H}_2$  gas. In § 3.2 we present and discuss new observational evidence for an intrinsic clumpiness of the broad spectral wings of the HCN(1-0) and  $\text{HCO}^+(1-0)$  lines. Our results are discussed in § 4 and summarised in § 5. Throughout the paper we adopt a standard  $\Lambda\text{CDM}$  cosmological model with  $H_0 = 67.8 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ,  $\Omega_\Lambda = 0.692$ ,  $\Omega_M = 0.308$  (Planck Collaboration et al. 2016). At the distance of Mrk 231 (redshift  $z = 0.04217$ , luminosity distance  $D_L = 192.4 \text{ Mpc}$ ), the physical scale is  $0.859 \text{ kpc arcsec}^{-1}$ . Uncertainties correspond to  $1\sigma$  statistical errors.

## 2. Observations

The  $N = 1 - 0$   $J = 3/2 - 1/2$  and  $J = 1/2 - 1/2$  spin-doublet transition of CN (hereafter, CN(1-0)) consists of several hyperfine lines (five in the brighter  $J = 3/2 - 1/2$  group and four in the  $J = 1/2 - 1/2$  group, e.g. Skatrud et al. (1983)), with a frequency separation that is much smaller than the molecular line width of Mrk 231 (maximum separation between the  $J = 3/2 - 1/2$  hyperfine levels,  $\Delta\nu^{\text{max}} = 32.3 \text{ MHz} = 85 \text{ km s}^{-1}$ ). The CN(1-0) doublet was included within the 3.6 GHz-wide spectral bandwidth of the IRAM Plateau de Bure Interferometer (PdBI, now Northern Extended Millimeter Array) observations targeting also the CO(1-0) transition and described in Cicone et al. (2012). However, the CN(1-0) data are shown and analysed for the first time in this paper. The phase centre of these observations is  $\text{RA}(J2000) = 12:56:14.200$ ;  $\text{Dec}(J2000) = +56:52:25.0$ . The other main technical parameters such as rest and observed frequency, date, array configuration, synthesised beam, and rms noise are summarised in Table 1. For data reduction and analysis we used the CLIC and MAPPING packages within the GILDAS software<sup>2</sup>. The 3mm continuum adjacent to the CO(1-0) and CN(1-0) lines was estimated between  $109.3037 < \nu_{\text{obs}}[\text{GHz}] < 109.8779$  and  $111.2470 < \nu_{\text{obs}}[\text{GHz}] < 111.8603$ , and subtracted from the  $uv$  visibility data prior to imaging and cleaning.

In this paper we also use archival IRAM PdBI observations of other molecular gas tracers that have been detected in the outflow of Mrk 231, presented in previous publications. Technical details and relevant references for these archival datasets are also reported in Table 1.

## 3. Results

### 3.1. CN and high density tracers in the outflow

The continuum-subtracted maps of the total CN(1-0) line emission and of the blue- and red-shifted CN(1-0) components are presented in Figure 1. For the line wings, we selected velocities that are not affected by blending of the two CN(1-0) spingroups. Besides the well-known central concentration of  $\text{H}_2$  gas embedding the molecular disk already studied by Downes & Solomon (1998), we identify a northern component, detected in CN at the  $12\sigma$  level, which is spatially extended and offset by  $\sim 3 - 7 \text{ kpc}$  with respect to the nucleus of Mrk 231 (left panel of Fig. 1).

<sup>2</sup> <http://www.iram.fr/IRAMFR/GILDAS>

<sup>1</sup> We refer to Shirley (2015) for further details on this definition, but we note that there are different nuances of  $n_{\text{eff}}$  in the literature, see also Leroy et al. (2017).

Table 1: IRAM PdBI observations of Mrk 231 used in this paper:

Line(s)	$\nu_{\text{rest}}$ [GHz]	$\nu_{\text{obs}}$ [GHz]	Date	Conf.	Clean beam ["'x'"]	rms (100 km s <sup>-1</sup> ) [mJy beam <sup>-1</sup> ]	Ref.	Proj. ID
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
CN(1-0)	113.4910 <sup>†</sup>	108.8987 <sup>†</sup>	2010	6C	2.8 × 2.6	0.47	This work	UA26
CO(1-0)	115.2712	110.6069	2009-2010	6C/5D	3.2 × 2.8	0.25	Cicone et al. (2012)	UA26, T02F
CO(2-1)	230.5380	221.2096	2010	6C/5D	1.3 × 1.0	0.79	Cicone et al. (2012)	UB26
CO(3-2)	345.7960	331.8038	2012	6D	1.5 × 1.3	2.1	Feruglio et al. (2015)	V087
HCN(1-0) <sup>‡</sup>	88.6319	85.0455	2011	6B	1.4 × 1.2	0.37	Aalto et al. (2012a)	U-D
HNC(2-1)	177.2610	170.0884	2012	5D	2.9 × 2.0	0.60	Lindberg et al. (2016)	W028
HCN(3-2)	265.8862	255.1275	2012-2013	6A/6B	0.4 × 0.3	0.54	Aalto et al. (2015)	V086, WA85

Notes: (1) Molecular transition; (2) rest frequency; (3) observed (sky) frequency; (4) observing dates; (5) configuration of the interferometer; (6) Synthesised beam size achieved after cleaning with natural weighting by using a support encompassing the  $3\sigma$  contour levels; (7)  $1\sigma$  rms per beam calculated in channels of  $\Delta\nu = 100 \text{ km s}^{-1}$ ; (8) relevant references; (9) identification code of the observing programme. <sup>†</sup> Central frequency of the  $J = 3/2 - 1/2$  spingroup (i.e. the brightest component of the CN(1-0) doublet); <sup>‡</sup> These observations include also the HCO<sup>+</sup>(1-0), HNC(1-0), and HC<sub>3</sub>N(10-9) transitions (see Fig. 2).




Fig. 1: IRAM PdBI continuum-subtracted maps of the CN(1-0) line emission. From left to right: (i) total emission integrated between  $-800 < v [\text{km s}^{-1}] < 1600$ , where velocities are calculated with respect to the  $J = 3/2 - 1/2$  spingroup (Table 1); (ii) blue wing of the  $J = 3/2 - 1/2$  line ( $-500 < v [\text{km s}^{-1}] < -300$ ); (iii) red wing, integrated between  $108.372 < \nu_{\text{obs}} [\text{GHz}] < 108.463$ , corresponding to velocities  $300 < v [\text{km s}^{-1}] < 550$  with respect to the central frequency of the fainter  $J = 1/2 - 1/2$  spingroup (where the red line wing is not affected by blending). All maps are corrected for the primary beam (45.6'' FWHM) and show the central  $25'' \times 25''$  region. The synthesised beam is reported at the bottom-left corner of the maps. Negative contours (dashed lines) correspond to  $(-6\sigma, -3\sigma, -2\sigma)$ , while positive contours (solid lines) correspond to  $(2\sigma, 3\sigma, 6\sigma, 9\sigma, n \times 12\sigma)$ , with  $n = 1, 2$ , etc where  $\sigma$  is the rms of the cleaned datacube (see Table 1).

Table 2: Line fluxes resulting from the simultaneous fit to the total integrated spectra:

Transition	Narrow core <sup>(*)</sup> [Jy km s <sup>-1</sup> ]	Blue Wing <sup>(†)</sup> [Jy km s <sup>-1</sup> ]	Red Wing <sup>(‡)</sup> [Jy km s <sup>-1</sup> ]	Total line [Jy km s <sup>-1</sup> ]
CO(1-0)	71.0 ± 0.6	9.2 ± 0.7	2.6 ± 0.2	83.0 ± 1.0
CN(1-0) $J = 3/2 - 1/2$	9.2 ± 0.4	4.1 ± 0.5	1.5 ± 0.3	14.8 ± 0.8
CN(1-0) $J = 1/2 - 1/2$	5.0 ± 0.3	2.2 ± 0.3	0.8 ± 0.2	8.1 ± 0.5
HNC(1-0)	4.3 ± 0.3	1.1 ± 0.4	0.5 ± 0.2	5.8 ± 0.5
HCO <sup>+</sup> (1-0)	6.4 ± 0.3	0.4 ± 0.4	0.7 ± 0.3	7.5 ± 0.6
HCN(1-0)	6.9 ± 0.3	2.9 ± 0.4	0.7 ± 0.3	10.6 ± 0.6
CO(2-1)	182.0 ± 1.7	23 ± 2	9.9 ± 1.1	215 ± 3
HNC(2-1)	22.0 ± 0.5	8.1 ± 0.6	3.0 ± 0.3	33.2 ± 0.8
CO(3-2)	545 ± 4	51 ± 5	40 ± 4	637 ± 8
HCN(3-2)	20.1 ± 1.0	12.5 ± 1.4	2.9 ± 0.8	35.4 ± 1.9
HC <sub>3</sub> N(10-9)				1.5 ± 0.5

Notes: <sup>(\*)</sup> Narrow core best-fit kinematics:  $v = 18.0 \pm 0.2 \text{ km s}^{-1}$ ;  $\sigma_v = 82.3 \pm 0.3 \text{ km s}^{-1}$ . <sup>(†)</sup> Blue wing best-fit kinematics:  $v = -135 \pm 16 \text{ km s}^{-1}$ ;  $\sigma_v = 168 \pm 8 \text{ km s}^{-1}$ . <sup>(‡)</sup> Red wing best-fit kinematics:  $v = 437 \pm 7 \text{ km s}^{-1}$ ;  $\sigma_v = 128 \pm 7 \text{ km s}^{-1}$ . The HC<sub>3</sub>N(10-9) line is modelled by a single Gaussian, with unconstrained width and velocity, shifted by  $v = -990 \pm 50 \text{ km s}^{-1}$  with respect to the HNC(1-0) line, with  $\sigma_v = 180 \pm 50 \text{ km s}^{-1}$  and  $S^{\text{peak}} = 3.4 \pm 0.8 \text{ mJy}$ .




Fig. 2: Total spectra - and simultaneous fit - of the CO(1-0), CN(1-0), HNC(1-0), HCO<sup>+</sup>(1-0), HCN(1-0), CO(2-1), HCN(2-1), CO(3-2), and HCN(3-2) molecular lines in Mrk 231. Overlaid on the data, we show the individual Gaussians employed in the simultaneous fit (up to three components for each line, red dashed curves) and the total best-fit curve (blue solid lines). Only the fainter HC<sub>3</sub>N(10-9) line was fitted by a single Gaussian with unconstrained velocity and width. The rms values per spectral channel are: 1.7 mJy per  $\delta v = 14$  km s<sup>-1</sup> (CO(1-0)); 2.3 mJy per  $\delta v = 43$  km s<sup>-1</sup> (CN(1-0)); 1.7 mJy per  $\delta v = 40$  km s<sup>-1</sup> (HNC(1-0)); 1.4 mJy per  $\delta v = 70$  km s<sup>-1</sup> (HCO<sup>+</sup> and HCN(1-0)); 8 mJy per  $\delta v = 26$  km s<sup>-1</sup> (CO(2-1)); 1.2 mJy per  $\delta v = 60$  km s<sup>-1</sup> (HCN(2-1)); 26 mJy per  $\delta v = 18$  km s<sup>-1</sup> (CO(3-2)); 5 mJy per  $\delta v = 47$  km s<sup>-1</sup> (HCN(3-2)).

To our knowledge, this is the first detection of such an extended molecular gas structure outside the main disk of Mrk 231. In Appendix A we show the CN(1-0) and CO(1-0) spectra extracted from a region encompassing the northern CN feature. The properties of this structure, i.e. its spectral line profile, spatial extent, and CN/CO(1-0) line luminosity ratio, suggest that it may represent an  $> 5$  kpc-size extension of the molecular outflow. Further supporting such interpretation, we note that previous CO and HCN observations have already evidenced blue-shifted and red-shifted features extending by  $\sim 2$  kpc north of the nucleus of Mrk 231 (Cicone et al. 2012; Aalto et al. 2012a; Feruglio et al. 2015), i.e. in a region lying between the newly-discovered northern CN structure and the well-known central kpc component of the outflow.

The total CN(1-0) spectrum of Mrk 231 is presented in Fig 2, together with the CO(1-0), HNC(1-0), HCO<sup>+</sup>(1-0), HCN(1-0), CO(2-1), HCN(2-1), CO(3-2), and HCN(3-2) line spectra. The sizes of the emitting regions differ slightly among different transitions, due to a combination of (i) different beam sizes, since higher resolution data (e.g. taken with configurations A and B of the IRAM PdBI, see Table 1) have lower sensitivity to large-scale structures because of interferometric filtering, and (ii) physical conditions, where higher excitation lines tend to be more compact than CO(1-0), as already noted for the outflow of Mrk 231 by Cicone et al. (2012). For this reason, we employed slightly different aperture sizes for different tracers, chosen to maximise the total emission detected in the corresponding primary beam-corrected datacubes. More specifically, the spec-

tra shown in Fig 2 were extracted from circular apertures with diameters of 10'' for CO(1-0), CN(1-0), CO(2-1), and HCN(2-1); 5'' for CO(3-2); 4'' for HCN(1-0), HCO<sup>+</sup>(1-0), and HNC(1-0); and 3'' for HCN(3-2). We note however that adopting the same 10''-size aperture for all lines would not change our results, but it would just result in slightly noisier HCN(1-0), HNC(1-0), HCO<sup>+</sup>(1-0), and HCN(3-2) spectra.

The striking similarity between the spectral profiles of all these different molecular transitions allows us to perform a simultaneous fit, whose results are overlaid on the data in Fig 2. The fit employs three Gaussian components, which are tied to have the same velocity and width for each line. Such simultaneous fitting procedure allows us to robustly deblend the broad wings of the two CN spin-groups. The amplitude ratio between the  $J = 1/2 - 1/2$  and  $J = 3/2 - 1/2$  components is fixed for each Gaussian, and constrained by the fit to be  $0.55 \pm 0.03$ , which is close to the theoretical optically thin value (0.5), as already noted by Henkel et al. (2014) using lower S/N single-dish CN(1-0) observations of Mrk 231. All other best-fit values are listed in Table 2.

Figure 2 shows that the broad wings are significantly more prominent in the dense gas tracers (e.g. CN, HCN) than in (low-J) CO transitions. For example, the CN(1-0) and HCN(1-0) wings-to-core flux ratios are higher respectively by a factor of four and three compared to CO(1-0).

The enhancement of high density tracers in Mrk 231's outflow is further explored in Figure 3, where we plotted, separately for the three Gaussian components fitted to Mrk 231's lines,




Fig. 3: CN/CO(1-0) versus HNC/CO(1-0) (*top*), HCO<sup>+</sup>/CO(1-0) (*middle*), and HCN/CO(1-0) (*bottom*) line luminosity ratios for the different line components of Mrk 231 (narrow core, blue wing, red wing, and total emission), including literature samples of nearby galaxies (grey circles) and GMCs (grey stars). Measurements corresponding to extragalactic molecular outflows (NGC 253 and NGC 3256, see Table B.1) are highlighted with purple circles. The CN(1-0) luminosity corresponds to the sum of the two spingroup components. When only the brightest  $J = 3/2 - 1/2$  spingroup line flux is available in the literature, we extrapolated the total CN(1-0) flux by multiplying this value by 1.5. The dashed line indicates the 1:1 relation. All errorbars include a 10% calibration uncertainty, which was added in quadrature to the statistical error ( $1\sigma$ ). Upper limits are shown at the  $3\sigma$  level. Sources with  $L'_{\text{CN}(1-0)}/L'_{\text{CO}(1-0)} > 0.2$  are labelled in the plot.




Fig. 4: CN/HCN(1-0) versus HNC/HCN(1-0) line luminosity ratios for the different line components of Mrk 231, compared to a sample of local galaxies and Galactic GMCs from the literature. See also caption of Fig. 3.

the CN/CO ( $J = 1 - 0$ ) against the HNC/CO, HCO<sup>+</sup>/CO, and HCN/CO ( $J = 1 - 0$ ) line luminosity ratios (i.e. the ratios between the line luminosities  $L'_{\text{line}}$  in units of  $[\text{K km s}^{-1} \text{ pc}^2]$ , calculated using the definition by Solomon & Vanden Bout (2005), which are equivalent to brightness temperature ratios). For comparison, Figure 3 includes also a sample of nearby galaxies as well as Galactic and extragalactic GMCs drawn from the literature (all literature values are reported in Table B.1 in Appendix B). Most of the literature data points correspond to integrated line measurements, without differentiating between disk and outflow components. The only exceptions are the galaxies NGC 253 and NGC 3256, for which molecular line fluxes have been measured at different locations by Meier et al. (2015) and Harada et al. (2018), including a few positions coincident with their molecular outflows (highlighted with purple circles in Fig 3).

It is evident from Fig. 3 that  $L'_{\text{CN}(1-0)}/L'_{\text{CO}(1-0)}$  is significantly boosted in both the blue- and red-shifted sides of Mrk 231's outflow, compared to the value measured in the narrow core of the lines, which should trace mainly the molecular disk. An enhancement with respect to the core is measured also in  $L'_{\text{HNC}(1-0)}/L'_{\text{CO}(1-0)}$  and  $L'_{\text{HCN}(1-0)}/L'_{\text{CO}(1-0)}$  for both line wings, and in  $L'_{\text{HCO}^+(1-0)}/L'_{\text{CO}(1-0)}$  for the red wing (but not for the blue wing). However, among all line ratios explored here,  $L'_{\text{CN}(1-0)}/L'_{\text{CO}(1-0)}$  shows the largest enhancement in the line wings relative to the core.

Furthermore, the  $L'_{\text{CN}(1-0)}/L'_{\text{CO}(1-0)}$  values of  $0.70 \pm 0.13$  and  $0.9 \pm 0.2$  measured respectively in the blue and red wings are significantly higher than the corresponding  $L'_{\text{HNC}(1-0)}/L'_{\text{CO}(1-0)}$  ( $0.19 \pm 0.07$  and  $0.28 \pm 0.16$  for the blue and red wings),  $L'_{\text{HCO}^+(1-0)}/L'_{\text{CO}(1-0)}$  ( $0.46 \pm 0.19$  for the red wing) and  $L'_{\text{HCN}(1-0)}/L'_{\text{CO}(1-0)}$  ( $0.54 \pm 0.12$  and  $0.46 \pm 0.19$  for the blue and red wings) values. Hence, CN shows the highest luminosity advantage in the line wings with respect to CO. Quite strikingly, the red wing of Mrk 231 is as bright in CN(1-0) as in CO(1-0) emission. Such high CN/CO line luminosity ratios are unprecedented if compared to any other extragalactic or Galactic measurement available in the literature. In particular, neither NGC 253 nor

NGC 3256 show an enhancement of CN (or other dense gas tracers) in the outflow comparable to that seen in Mrk 231, suggesting that the physical state of the molecular gas in outflow can differ significantly from source to source. The CN(1-0) line emission is exceptionally bright in the outflow of Mrk 231, not only compared to Mrk231's own molecular disk (traced by the 'core' component shown in Fig 3), but also compared to any other known astrophysical environment.

However, while Mrk 231's outflow shows an unprecedented enhancement of high density tracers such as CN(1-0) compared to the CO(1-0) line emission (which should be a good proxy of the total H<sub>2</sub> gas mass), the line luminosity ratios computed between different dense gas tracers (e.g. CN/HCN, HNC/HCN) are not uncommon compared to other known astrophysical sources. Indeed, Figure 4 shows that the  $L'_{\text{CN}(1-0)}/L'_{\text{HCN}(1-0)}$  and  $L'_{\text{HNC}(1-0)}/L'_{\text{HCN}(1-0)}$  values measured in the wings and in the core of Mrk 231's lines are formally consistent with each other (although  $L'_{\text{CN}(1-0)}/L'_{\text{HCN}(1-0)}$  is slightly higher in the red wing than in the other components). Furthermore, both line luminosity ratios fall within the range of values measured in the literature sample.

### 3.2. Spectrally-resolved clumps in the HCN and HCO<sup>+</sup> wings

The broad wings of the HCN(1-0) and HCO<sup>+</sup>(1-0) lines, first detected by Aalto et al. (2012a), show several narrow substructures when examined at high spectral resolution. These features are clearly visible in the spectrum extracted from the central beam (containing  $\sim 70\%$  of the total HCN(1-0) flux), presented in Figure 5. To analyse the properties of these spectral clumps, we performed a simultaneous fit to the HCN(1-0) and HCO<sup>+</sup>(1-0) transitions by using multiple Gaussian components, whose results are overlaid on the data in Fig 5. The fit confirms that clumps labelled as 'A1', 'A2', and 'A3' are detected in both the HCN and HCO<sup>+</sup> lines (with the same velocity and dispersion), whereas the HCN clumps labelled as 'B1', 'B2', and 'B3' do not have a counterpart in HCO<sup>+</sup>. Only one feature, labelled as 'C1', is detected in HCO<sup>+</sup> but not in HCN. As discussed later on, all these clumps are detected at high significance. Such differences between those species were already pointed out by Lindberg et al. (2016) based on the analysis of Mrk 231's spectral line profiles. In particular, they noted the presence of HCN and HCO<sup>+</sup> features at different velocities in the line wings. Moreover, their radiative transfer modelling of the HCN and HCO<sup>+</sup> line emission is inconsistent when using the same outflow properties for both species: while a clumpy outflow model could reproduce the HCN emission, the same model would lead to an unrealistically high outflow mass for HCO<sup>+</sup>. These facts suggest that the HCN and HCO<sup>+</sup> line emissions arise in regions with somehow different physical/chemical properties, hereby a clumpy and chemically differentiated outflow. One possible origin of this difference could be related to the presence of shock fronts in the outflow, with HCN tracing the shocked regions between the outflow and the surrounding gas, and high-velocity HCO<sup>+</sup> tracing pre-shock gas, closer to the nucleus.

The best-fit Gaussian parameters of the clumps are reported in Table 3. The velocity dispersions of  $\sigma_v \sim 7 - 20 \text{ km s}^{-1}$  are consistent with individual GMCs. These  $\sigma_v$  values are larger than typical Milky Way GMCs ( $\langle\sigma_v\rangle \sim 2 \text{ km s}^{-1}$ ) but smaller than those measured for clouds populating the dense nuclear starburst of NGC 253 ( $\langle\sigma_v\rangle \sim 22 \text{ km s}^{-1}$ , Leroy et al. 2015). By assuming HCN-to-H<sub>2</sub> conversion factors in the range  $3.1 - 24 M_{\odot} (\text{K km s}^{-1} \text{ pc}^2)^{-1}$  (e.g. Gao & Solomon 2004; Aalto

et al. 2015; Leroy et al. 2015), the HCN luminosities of the clumps would correspond to molecular gas masses of  $M_{\text{mol}} \sim 0.3 - 4.9 \times 10^8 M_{\odot}$  (see Table 3). Such masses are one order of magnitude larger than typical GMCs in the MW and M33 (Rosolowsky et al. 2003), but roughly comparable (in the lower-mass end limit) to the more massive GMCs populating the dense environment within the nuclear starburst of NGC 253 (Leroy et al. 2015).

The spectral clumps appear less significant in spectra extracted from apertures larger than the central beam. Only features A1, A2, B2 and B3 are clearly detected also in the spectrum extracted from a 4''-size aperture, with best-fit  $v$  and  $\sigma_v$  values consistent with those measured within the central beam (Table 3). The lower significance of the spectral clumps in large apertures is due to the fact that they are spatially unresolved at the resolution of the HCN(1-0) data ( $\sim 1 \text{ kpc}$ , see Table 1). Hence, their flux is maximised in the central beam, but it is washed out in larger apertures because of the superposition with additional outflow components.

We checked the significance of the HCN(1-0) clumps in the  $uv$  visibility data, by analysing the  $uv$  amplitude as a function of  $uv$  radius ( $uv$  plots). The interferometric images of the HCN(1-0) clumps and the corresponding  $uv$  plots are shown in Fig. 6. Both the maps and the  $uv$  plots, with their flat trend of  $uv$  visibilities as a function of  $uv$  radius, confirm that the HCN clumps are spatially unresolved in our observations. A fit to the  $uv$  plots using a point source model, whose results are also listed in Table 3, shows that the HCN clumps (A1-A3, B1-B3) are individually detected at a high significance, ranging from  $S/N \sim 4.8 - 7.4$ . Furthermore, as shown in Table 3, the integrated fluxes derived from the  $uv$  fit are overall consistent with those obtained from the simultaneous spectral fit shown in Figure 5. Since the  $uv$  visibility data contain the *total* flux within a certain velocity range (chosen as  $v_c \pm 2\sigma_v$ , where  $v_c$  and  $\sigma_v$  are the central velocity and dispersion of each spectral clump, listed in Table 3), while the spectrum in Fig. 5 samples only the central beam, such flux consistency between the central beam and the  $uv$  data suggests that the clumps dominate the HCN flux within the velocity ranges sampled by their narrow line profile. In other words, at the velocity of the clumps, there is little additional flux in more spatially extended and diffuse components.

We searched for the presence of clumps also in the other molecular gas tracers, by inspecting the spectra extracted from the central beam. The HNC(1-0) and CN(1-0) line wings show clumpy features similar to HCN(1-0), but the lower S/N of these datasets and, in the case of CN, the blending of the two spin-groups, do not allow us to reliably fit any of these features. In the CO(1-0) transition, the line wings are intrinsically much smoother compared to the high density tracers, even when studied at the highest available spectral resolution. The only discernible structure in the CO(1-0) line wings is a red-shifted feature with best-fit Gaussian parameters:  $v = 459 \pm 10 \text{ km s}^{-1}$ ,  $\sigma_v = 39 \pm 17 \text{ km s}^{-1}$  and  $S_{\text{CO}}^{\text{peak}} = 1.7 \pm 0.6 \text{ mJy}$ , which does not correspond to any of the clumps revealed in HCN or HCO<sup>+</sup>.

Our results support a scenario where the molecular outflow entrains high-velocity, dense molecular gas clouds, dominating the emission in the HCN and HCO<sup>+</sup> line wings, which are embedded in a more diffuse phase that contributes most to the observed CO(1-0) line wings. A similar conclusion was independently reached by Cicone et al. (2018) for the dual AGN and LIRG NGC 6240, based on a multi-tracer analysis of the extended molecular outflow.




Fig. 5: Spectral fit to the HCN(1-0) and HCO<sup>+</sup>(1-0) line wings showing the presence of several spectrally-resolved narrow substructures (here often indicated as ‘clumps’). The spectrum was extracted from the cleaned datacube before continuum-subtraction and using a beam-size aperture centred at RA(J2000)=12:56:14.216 and Dec(J2000)=56:52:25.186. The rms is 1.8 mJy per  $\delta v = 6.9 \text{ km s}^{-1}$  spectral channel. The clumps’ solutions that were simultaneously fitted to both transitions are plotted in magenta, whereas features detected only in HCN (HCO<sup>+</sup>) are shown in cyan (orange). The best-fit Gaussian parameters are reported in Table 3.

Table 3: Analysis of the HCN(1-0) and HCO<sup>+</sup>(1-0) outflowing clumps

ID	Spectral fit (Gaussian)						$uv$ visibility fit (point source model)				
	$v$ [km s <sup>-1</sup> ]	$\sigma_v$ [km s <sup>-1</sup> ]	$S_{\text{peak}}^{\text{HCO}^+}$ [mJy]	$S_{\text{peak}}^{\text{HCN}}$ [mJy]	S/N (HCN)	$\int S^{\text{HCN}} dv$ [Jy km s <sup>-1</sup> ]	$M_{\text{mol}}^{\ddagger}$ [10 <sup>8</sup> M <sub>⊙</sub> ]	$\delta\text{RA}$ [″]	$\delta\text{Dec}$ [″]	S/N	$\int S^{\text{HCN}} dv$ [Jy km s <sup>-1</sup> ]
A1	-465 ± 2	9 ± 2	2.4 ± 0.8	4.4 ± 0.9	4.9	0.10 ± 0.03	0.5-3.6	-0.05 ± 0.13	0.17 ± 0.12	4.8	0.11 ± 0.02
A2	-406 ± 2	7 ± 2	1.6 ± 0.9	3.9 ± 1.0	3.9	0.07 ± 0.03	0.3-2.4	0.03 ± 0.15	0.09 ± 0.13	4.4	0.08 ± 0.02
A3	313 ± 3	11 ± 3	2.2 ± 0.8	3.6 ± 0.8	4.5	0.10 ± 0.03	0.4-3.4	0.11 ± 0.10	-0.23 ± 0.09	6.2	0.15 ± 0.03
B1	-665 ± 4	14 ± 4	< 2.1 <sup>†</sup>	3.0 ± 0.7	4.3	0.11 ± 0.04	0.5-3.7	-0.28 ± 0.10	-0.16 ± 0.09	6.6	0.17 ± 0.03
B2	-349 ± 6	20 ± 6	< 2.1 <sup>†</sup>	2.8 ± 0.7	4.0	0.14 ± 0.05	0.6-4.9	-0.12 ± 0.09	0.12 ± 0.08	7.1	0.24 ± 0.03
B3	526 ± 3	11 ± 3	< 2.7 <sup>†</sup>	4.1 ± 0.9	4.6	0.11 ± 0.04	0.5-3.9	-0.11 ± 0.09	0.10 ± 0.08	7.4	0.16 ± 0.02
C1	-297 ± 6	13 ± 6	2.1 ± 0.8	< 2.4 <sup>†</sup>	< 3	< 0.07 <sup>†</sup>	< 2.0 <sup>†</sup>				

<sup>†</sup>  $3\sigma$  upper limit; <sup>‡</sup> Range of dense molecular gas masses estimated from the HCN(1-0) line luminosity following  $M_{\text{mol}} [M_{\odot}] = \alpha_{\text{HCN}} L'_{\text{HCN}(1-0)} [\text{K km s}^{-1} \text{ pc}^2]$ , assuming  $\alpha_{\text{HCN}} \in [3.1, 24] M_{\odot} (\text{K km s}^{-1} \text{ pc}^2)^{-1}$ , where the lower  $\alpha_{\text{HCN}}$  value has been proposed by García-Burillo et al. (2012) for (U)LIRGs, whereas the higher  $\alpha_{\text{HCN}}$  has been measured by Leroy et al. (2015) for individual GMCs in the starburst region of NGC 253. We note that both Gao & Solomon (2004), based on integrated observations of local galaxies and on the assumption that HCN traces very dense gas, and Shimajiri et al. (2017), based on observations of Galactic GMCs (see also Kauffmann et al. (2017)) propose an intermediate value of  $\alpha_{\text{HCN}} = 10 M_{\odot} (\text{K km s}^{-1} \text{ pc}^2)^{-1}$ .

## 4. Discussion

### 4.1. Enhanced CN/HCN abundance ratios

We aim at understanding the origin of the enhanced CN/CO(1-0) and HCN/CO(1-0) line luminosity ratios measured in the outflow of Mrk 231 (Figure 3). This can be due to a combination of (i) higher fractional abundances of such tracers ( $X_{\text{HCN}}$ ,  $X_{\text{CN}}$ , i.e. the abundance ratios of these species over the total H<sub>2</sub> abundance) and/or (ii) a substantial entrainment of dense gas with  $n > 10^4 \text{ cm}^{-3}$ , which would increase the excitation of these lines (Aalto et al. 2012b, 2015). A high abundance of HCN may be the signature of shocks (e.g. see Aalto et al. 2012b and references therein), whereas a high  $X_{\text{CN}}$  should trace gas exposed to a strong UV radiation field (PDR-like behaviour) from massive stars. García-Burillo et al. (2010) proposed that CN emis-

sion is amplified in the vicinity of an AGN due to the X-ray radiation field, although there is still no consensus on this result, since many AGN hosts show the opposite behaviour, with  $L'_{\text{CN}}/L'_{\text{CO}}$  values that are lower by a factor of 2-3 compared to the average ratios measured in pure starbursts (Wilson 2018; Pérez-Beaupuits et al. 2007). The prominent CN line wings of Mrk 231 may indicate that the outflow is bathed in a strong UV radiation field, either due to star formation in the galaxy disk or to new stars forming within the outflow itself. The latter is a scenario proposed by various recent models and simulations (Zubovas et al. 2013; Zubovas & King 2014; Ishibashi & Fabian 2012; Ishibashi et al. 2013; El-Badry et al. 2016; Wang & Loeb 2018; Decataldo et al. 2019). Star formation in the outflow could have major implications, since such stars would have highly radial orbit and could, for instance, contribute to the formation of




Fig. 6: Interferometric maps and corresponding  $uv$  visibility plots of the HCN(1-0) clumps (top rows: A1, A2, A3; bottom rows: B1, B2, and B3, see also labels on the top-left corner of the maps). The maps show the continuum-subtracted emission integrated between  $v_c - 2\sigma_v < v < v_c + 2\sigma_v$ , where  $v_c$  and  $\sigma_v$  are the best-fit central velocity and velocity dispersion obtained from the spectral line fitting (Table 3). Each map is  $10'' \times 10''$ , and the synthesised beam is  $1.45'' \times 1.21''$ . Contours correspond to  $(-3\sigma, -2\sigma, 2\sigma, 3\sigma, 4\sigma, 5\sigma, 6\sigma)$ . Below each map we report the corresponding  $uv$  plot, binned in intervals of 30m. The red solid and dashed lines represent the best-fit value and its associated uncertainty obtained using a point source model (Table 3).

the spheroidal component of galaxies. Observational confirmations of this new mode of star formation have been presented recently (Maiolino et al. 2017; Gallagher et al. 2019; Rodríguez del Pino et al. 2019). Within the context of our findings, young massive stars formed inside the outflow would provide a much

stronger (internal) UV radiation field with respect to the illumination from stars in the disk or even from the AGN.



By using the RADEX<sup>3</sup> dense cloud models developed by Aalto et al. (2015) to reproduce the HCN(3-2)/(1-0) line luminosity ratios in Mrk 231's outflow, we can attempt to find a combination of  $X_{\text{HCN}}$ ,  $X_{\text{CN}}$ ,  $T_{\text{kin}}$  and  $n_{\text{H}_2}$  solutions that can fit also the CN/HCN and CN spin doublet line ratios (Table 2). We assume that the HCN and CN line emissions arise from the same dense cloud population, while the low-J CO line emission is due to a different, more diffuse phase of the outflow. We recall that in these models (see also Aalto et al. 2015), the dense clouds can be either self gravitating, virialized clouds, i.e. implying that their internal velocity dispersion ( $\Delta v_{\text{sg}}$ ) is locked to their mass ( $M_{\text{vir}}$ ) and size ( $R$ ) through  $\Delta v_{\text{sg}} = (GM_{\text{vir}}/R)^{1/2}$ , or unbound clouds, for which  $\Delta v \gg \Delta v_{\text{sg}}$ . We explore CN and HCN abundances in the range between  $10^{-8}$  and  $10^{-6}$ . We find that, depending on whether the clouds are self gravitating or unbound, the models produce very different values for the absolute CN and HCN abundances, hence  $X_{\text{CN}}$  and  $X_{\text{HCN}}$  remain quantitatively unconstrained for the outflow with current data. However, all possible solutions that fit the observed line ratios consistently require  $X_{\text{CN}} > X_{\text{HCN}}$ , with an CN abundance that is at least a factor of three higher than the HCN abundance. Gas densities for such outflow phase (traced by the CN and HCN emissions) are  $n_{\text{H}_2} \sim 10^5 - 10^6 \text{ cm}^{-3}$ , with temperatures not much higher than  $T_{\text{kin}} \sim 20 \text{ K}$ . Since CN is a well-known PDR tracer (see also § 1), these results strongly suggest that the whole dense cloud population in outflow is affected by UV radiation. We should mention that high CN abundances may be due also to cosmic rays (e.g. see work done on the Galactic centre by Harada et al. (2015)), which are known to permeate Mrk 231's outflow as inferred by González-Alfonso et al. (2018) based on the OH<sup>+</sup> enhancement. However, it is not clear whether a cosmic ray chemistry would also explain  $X_{\text{CN}} > X_{\text{HCN}}$ .

In the literature (see Appendix B), there are only two other extragalactic molecular outflows with available CN(1-0) observations (NGC 3256 and NGC 253, see also Table B.1), and neither of them shows the exceptionally high  $L'_{\text{CN}(1-0)}/L'_{\text{CO}(1-0)}$  values that we measure in Mrk 231. However, in NGC 3256, Sakamoto et al. (2014) and Harada et al. (2018) find an enhancement of CN and HCN at the outflow position compared to other locations within the galaxy merger. Different from CN, there are quite a few extragalactic outflows with HCN and/or HCO<sup>+</sup> observations. In Arp 220, observations by Barcos-Muñoz et al. (2018) imply  $L'_{\text{HCN}(1-0)}/L'_{\text{CO}(1-0)} \sim 0.9$  and  $\sim 7$  for the blue- and red-shifted outflows, which are significantly higher than the corresponding luminosity ratio of  $0.20 \pm 0.03$  obtained by using the HCN and CO emissions integrated across the whole galaxy (Table B.1). In IC 5063, the high HCO<sup>+</sup>(4-3)/CO(2-1) ratio measured in the outflow by Oosterloo et al. (2017) points to an additional dense component with  $n \gtrsim 10^6 \text{ cm}^{-3}$  that is not accounted for by the low-J CO emission (which is likely optically thin, e.g. Dasyra et al. (2016)). Aladro et al. (2018) detected the blue-shifted component of Mrk 273's molecular outflow in HCN(3-2) emission, and its location is consistent with the CO(1-0) component imaged by Cicone et al. (2014). Among other extragalactic outflows detected in dense molecular gas tracers we list NGC 1068 (HCN(4-3) and HCO<sup>+</sup>(4-3) emission, García-Burillo et al. 2014), NGC 1266 (CS(2-1) and HCN(1-0) emis-

sion, Alatalo et al. 2015), and IC 860 (CS(7-6) absorption, Aalto et al. 2019).

Interestingly, even though the molecular streamers embedded in the starburst-driven outflows of NGC 253 and M 82 have been detected in dense gas tracers, the latter do not show a significant enhancement compared to CO emission. In NGC 253, the CN emission is boosted (with respect to both CO and C<sup>17</sup>O) in the central starburst, but not in the streamers (Meier et al. 2015). The HCN/CO luminosity ratio is instead similar between starburst and outflow (Walter et al. 2017). The outflow of M82 was detected in HCO<sup>+</sup>(1-0) and HCN(1-0) emission (Salas et al. 2014), but the mass traced by these molecules is a small fraction ( $\gtrsim 2\%$ ) of the total (CO-based) mass of the outflow. Hence, the molecular gas filaments embedded in the starburst-driven outflows of NGC 253 and M82 appear to differ in their physical properties from Mrk 231's outflow.

The exceptional boost of CN and HCN in Mrk 231's outflow may hold clues to explain the enhancement of such dense gas tracers in integrated observations of this ULIRG. Indeed, although the total flux is dominated by the narrow component of the molecular lines (as also reflected by the line ratios in Fig. 3, where the integrated data point lies closer to the core than to the wings) we note that the CN/CO and HCN/CO luminosity ratios measured in the narrow core are quite high compared to the nearby galaxy population and to local GMCs, although not as extreme as those measured in the wings. We cannot rule out that the low-projected velocity component of the outflow is contaminating significantly also the narrow emission<sup>4</sup>. In this case, since such component would share the same CN and HCN enhancement of the high velocity wings, it would also boost the line ratios measured for the narrow core. In the hypothesis that Mrk 231-like outflows have a chemical impact on the global ISM of their host galaxies, they would resemble scaled-up versions of the so-called 'chemically active' protostellar outflows revealed in young stellar objects (YSOs), which are characterised by enhanced abundances of shock sensitive tracers such as SiO (Tafalla & Hacar 2013) or even CN and HCO<sup>+</sup> (Bachiller et al. 2001).

#### 4.2. Dense clumps embedded in the outflow

The discovery of narrow features in the HCN and HCO<sup>+</sup> line wings suggests that the outflow embeds coherent dense molecular gas complexes, with velocity dispersions similar to extragalactic GMCs detected in dense environments. Furthermore, our data show that, while such clumps dominate the high-velocity emission from the high density tracers, they are not present in the wings of low-J CO lines. The HCN clouds embedded in Mrk 231's outflow are massive, with inferred  $M_{\text{mol}}$  values in the range  $\sim 0.3 - 4.9 \times 10^8 M_{\odot}$  (Table 3), and have large velocity dispersions of  $\sim 7 - 20 \text{ km s}^{-1}$ . Such unusually high masses may result from an observational bias, where multiple outflowing clouds with overlapping line-of-sight velocities contribute to the narrow sub-structures that we identify in the (beam-averaged) HCN spectra in Fig. 5. However, it is also possible that we are witnessing the survival of only the biggest and/or the most massive molecular clouds in outflow, which are less affected by photoevaporation (Decataldo et al. 2019) and hydrodynamical instabilities (Gronke & Oh 2018; Armillotta et al. 2017).

<sup>3</sup> RADEX is a publicly available non-local thermodynamic equilibrium radiative transfer code, which uses as input parameters the molecular gas column density ( $N_{\text{H}_2}$ ), the line width ( $\Delta v$ ), the gas temperature ( $T_{\text{kin}}$ ), the background temperature (set equal to the Cosmic Microwave Background temperature at  $z = 0$ , i.e.  $T_{\text{bg}} = 2.73 \text{ K}$ ), and the gas volume density (van der Tak et al. 2007)

<sup>4</sup> Only very high S/N, spatially resolved observations would allow us to quantify such contribution, by de-blending the low-velocity outflow emission from the rotating disk.

These physical properties, i.e. masses and velocity dispersions a factor of ten larger than GMCs in the Galactic disk, seem to be more typical of molecular clouds forming in dense environments, such as those residing in the nuclear starburst of NGC 253. The latter, identified by Leroy et al. (2015) through HCN-to-CO intensity peaks (one of which may be even entrained in the outflow), are characterised by higher surface and volume densities, shorter free fall time-scales and so higher star formation efficiencies compared to typical disk GMCs. That the HCN clumps detected in Mrk 231's outflow correspond to star forming complexes is a very interesting prospect, especially in the context of the new scenarios predicting the formation of stars inside outflows. Indeed, dense and clumpy gas are conditions that typically result into rapid gravitational collapse and star formation (Decataldo et al. 2019); hence our finding provides further support for this new mode of star formation.

We note that, if the clouds' mass distribution in outflow differs strongly from that of typical GMCs, as suggested by our results, it would result in a different initial mass function (IMF) of the stars that would form within such environment (Hopkins 2012), and possibly explain other exotic properties of (U)LIRGs such as their high  $^{18}\text{O}$  abundance (which favours a top-heavy IMF, e.g. Brown & Wilson 2019). However, a valid objection is that, even if the high-velocity clumps trace individual clouds, we know that, at least in the Milky Way, not all GMCs are active stellar nurseries. Indeed, even within GMCs, HCN emission can be quite widespread and up to  $\sim 50\%$  of it can be associated with extended regions with  $n_{\text{H}_2} \lesssim 10^3 \text{ cm}^{-3}$ , i.e. two orders of magnitude below the critical density for collisions with  $\text{H}_2$  molecules (Pety et al. 2017; Kauffmann et al. 2017; Harada et al. 2019). In these regions, HCN may be excited by electron collisions (Goldsmith & Kauffmann 2017). Additional data are required to settle this issue.

## 5. Summary and Conclusions

We have reported the detection of an exceptional enhancement of CN(1-0) line emission in the molecular outflow of Mrk 231. More specifically, we measured  $L'_{\text{CN}(1-0)}/L'_{\text{CO}(1-0)}$  values of  $0.70 \pm 0.13$  (blue-shifted wing) and  $0.9 \pm 0.2$  (red-shifted wing), which are not only much higher than the value measured in the narrow core component ( $0.21 \pm 0.03$ ), but also higher than any other known Galactic or extragalactic source. The CN/CO(1-0) line luminosity ratios computed for the line wings are also significantly higher than the respective HCN/CO(1-0),  $\text{HCO}^+$ /CO(1-0) and HNC/CO(1-0) line luminosity ratios, indicating that, among the dense gas tracers explored so far in Mrk 231's outflow, CN has the highest luminosity advantage with respect to CO.

We have found that, in order to reproduce these ratios, the dense gas cloud population entrained in the outflow must be characterised by an CN abundance that is at least a factor of three higher than the HCN abundance, by gas densities of  $n_{\text{H}_2} \sim 10^{5-6} \text{ cm}^{-3}$ , and kinetic temperatures not much higher than  $T_{\text{kin}} \sim 20 \text{ K}$ . This result can be interpreted (although not uniquely) in terms of an outflow chemistry that is heavily affected by UV radiation, since CN emission has been found to be amplified mainly in PDRs. A population of young massive stars formed inside the outflow, as predicted by recent models, would naturally provide an explanation for such strong UV radiation field.

In the literature there are only two other extragalactic molecular outflows with available CN measurements, NGC 253 and

NGC 3256, and neither of them shows an CN enhancement similar to Mrk 231. Although we are still limited by statistics, we may conclude that the outflow of Mrk 231 is rather exceptional in its chemical and physical properties. In the future, using deeper and higher spatial resolution observations, it would be interesting to explore whether the outflow component is responsible for the exceptionally strong emission from dense  $\text{H}_2$  gas tracers observed in the global ISM of this ULIRG.

Additionally, we have shown that the HCN(1-0) and  $\text{HCO}^+(1-0)$  line wings are spectrally resolved into several narrow sub-structures with velocity dispersions of  $\sigma_v \sim 7 - 20 \text{ km s}^{-1}$  and inferred masses of  $M_{\text{mol}} \sim 0.3 - 4.9 \times 10^8 M_{\odot}$ , which we interpret as individual dense clumps entrained in Mrk 231's outflow. These properties are consistent with the most massive extragalactic GMCs populating the nuclear starburst of NGC 253. Three of the outflowing clumps are detected simultaneously in both the HCN and  $\text{HCO}^+(1-0)$  transitions, while additional three (one) are detected only in HCN (or  $\text{HCO}^+$ ), supporting the hypothesis of a chemically differentiated outflow. An analysis of the  $uv$  visibility data indicates that these narrow HCN outflow features are spatially unresolved at the resolution of our data ( $\sim 1 \text{ kpc}$ ), but, at their line-of-sight velocity, there is little additional HCN flux in more spatially extended and diffuse components. The CO spectral line wings are instead intrinsically much smoother, implying that low-J CO emission is dominated by a more diffuse component of the outflow.

This discovery strongly suggests that the outflow of Mrk 231 embeds coherent molecular gas cloud complexes, similar to GMCs, but overall more massive (and with different chemical properties). This possibly indicates that we are witnessing the survival of only the biggest molecular clouds that are less affected by photo-evaporation and hydrodynamical instabilities, i.e. the processes that we expect to be at work in energetic multi-phase outflows and that may quickly destroy most of the coldest and densest gas. If these clumps correspond to star forming complexes, as predicted by recent theoretical models, the stars that would form from gravitational collapse of these massive clouds would probably follow a different IMF. However, this intriguing hypothesis requires follow-up observations to confirm the star forming nature of such clumps and better constrain their masses.

Finally, our interferometric maps have revealed a new molecular gas structure in Mrk 231, offset to the North by  $5''$ . Its high CN/CO line luminosity ratio and broad blue-shifted line profiles suggest that this northern component may trace an additional extension of the outflow, reaching out to  $r > 5 \text{ kpc}$  from the nucleus.

*Acknowledgements.* The interpretation of our data, especially for what concerns the ISM chemistry in the presence of strong UV fields, benefited greatly from extensive discussions with the late Malcolm Walmsley. We are deeply grateful to him, not only for his help with this paper, but also for the several insightful and enlightening discussions that some of us had with him on a broad range of topics in Astrophysics. CC thanks Jens Kauffmann for providing the Orion data, Nanase Harada and Kazushi Sakamoto for providing non-published measurements of NGC 3256, Paola Severgnini for insightful discussions about the multi-wavelength emission of Mrk 231, and Paola Caselli for valuable explanations on the properties of CN. This work is based on observations carried out under project numbers UA26, T02F, UB26, V087, U-D, W028, V026, WA85 with the IRAM NOEMA Interferometer. IRAM is supported by INSU/CNRS (France), MPG (Germany) and IGN (Spain). This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 664931. RM acknowledges ERC Advanced Grant 695671 'QUENCH' and support by the Science and Technology Facilities Council (STFC).

## References

- Aalto, S., García-Burillo, S., Müller, S., et al. 2015, *A&A*, 574, A85
- Aalto, S., García-Burillo, S., Müller, S., et al. 2012a, *A&A*, 537, A44
- Aalto, S., García-Burillo, S., Müller, S., et al. 2012b, *A&A*, 537, A44
- Aalto, S., Müller, S., König, S., et al. 2019, arXiv e-prints, arXiv:1905.07275
- Aalto, S., Polatidis, A. G., Hüttemeister, S., & Curran, S. J. 2002, *A&A*, 381, 783
- Aladro, R., König, S., Aalto, S., et al. 2018, *A&A*, 617, A20
- Aladro, R., Martín, S., Riquelme, D., et al. 2015, *A&A*, 579, A101
- Alatalo, K. 2015, *ApJ*, 801, L17
- Alatalo, K., Lacy, M., Lanz, L., et al. 2015, *ApJ*, 798, 31
- Armillotta, L., Fraternali, F., Werk, J. K., Prochaska, J. X., & Marinacci, F. 2017, *MNRAS*, 470, 114
- Baan, W. A., Henkel, C., Loenen, A. F., Baudry, A., & Wiklind, T. 2008, *A&A*, 477, 747
- Bachiller, R., Pérez Gutiérrez, M., Kumar, M. S. N., & Tafalla, M. 2001, *A&A*, 372, 899
- Barcos-Muñoz, L., Aalto, S., Thompson, T. A., et al. 2018, *ApJ*, 853, L28
- Biernacki, P. & Teyssier, R. 2018, *MNRAS*, 475, 5688
- Boger, G. I. & Sternberg, A. 2005, *ApJ*, 632, 302
- Braito, V., Della Ceca, R., Piconcelli, E., et al. 2004, *A&A*, 420, 79
- Brown, T. & Wilson, C. D. 2019, *ApJ*, 879, 17
- Chevalier, R. A. & Clegg, A. W. 1985, *Nature*, 317, 44
- Cicone, C., Feruglio, C., Maiolino, R., et al. 2012, *A&A*, 543, A99
- Cicone, C., Maiolino, R., Sturm, E., et al. 2014, *A&A*, 562, A21
- Cicone, C., Severgnini, P., Papadopoulos, P. P., et al. 2018, *ApJ*, 863, 143
- Costa, T., Rosdahl, J., Sijacki, D., & Haehnelt, M. G. 2018, *MNRAS*, 479, 2079
- Costa, T., Sijacki, D., & Haehnelt, M. G. 2015, *MNRAS*, 448, L30
- Costagliola, F., Aalto, S., Rodríguez, M. I., et al. 2011, *Astronomy and Astrophysics*, 528, A30
- Dasyra, K. M., Combes, F., Oosterloo, T., et al. 2016, *A&A*, 595, L7
- Davies, R. I., Müller Sánchez, F., Genzel, R., et al. 2007, *ApJ*, 671, 1388
- Decataldo, D., Pallottini, A., Ferrara, A., Vallini, L., & Gallerani, S. 2019, *MNRAS*, 487, 3377
- Di Matteo, T., Springel, V., & Hernquist, L. 2005, *Nature*, 433, 604
- Downes, D. & Solomon, P. M. 1998, *ApJ*, 507, 615
- El-Badry, K., Wetzel, A., Geha, M., et al. 2016, *ApJ*, 820, 131
- Faucher-Giguère, C.-A. & Quataert, E. 2012, *MNRAS*, 425, 605
- Feruglio, C., Fiore, F., Carniani, S., et al. 2015, *A&A*, 583, A99
- Feruglio, C., Maiolino, R., Piconcelli, E., et al. 2010, *A&A*, 518, L155+
- Fischer, J., Sturm, E., González-Alfonso, E., et al. 2010, *A&A*, 518, L41+
- Fluetsch, A., Maiolino, R., Carniani, S., et al. 2019, *MNRAS*, 483, 4586
- Fuente, A., Martín-Pintado, J., Cernicharo, J., & Bachiller, R. 1993, *A&A*, 276, 473
- Gallagher, R., Maiolino, R., Belfiore, F., et al. 2019, *MNRAS*, 485, 3409
- Gao, Y. & Solomon, P. M. 2004, *The Astrophysical Journal Supplement Series*, 152, 63
- García-Burillo, S., Combes, F., & Usero, A. e. a. 2014, *A&A*, 567, A125
- García-Burillo, S., Usero, A., Alonso-Herrero, A., et al. 2012, *A&A*, 539, A8
- García-Burillo, S., Usero, A., Fuente, A., et al. 2010, *A&A*, 519, A2
- Goldsmith, P. F. & Kauffmann, J. 2017, *ApJ*, 841, 25
- González-Alfonso, E., Fischer, J., Bruderer, S., et al. 2018, *ApJ*, 857, 66
- González-Alfonso, E., Fischer, J., Graciá-Carpio, J., et al. 2014, *A&A*, 561, A27
- Gronke, M. & Oh, S. P. 2018, *MNRAS*, 480, L111
- Harada, N., Nishimura, Y., Watanabe, Y., et al. 2019, *ApJ*, 871, 238
- Harada, N., Riquelme, D., Viti, S., et al. 2015, *A&A*, 584, A102
- Harada, N., Sakamoto, K., Martín, S., et al. 2018, *ApJ*, 855, 49
- Henkel, C., Asiri, H., Ao, Y., et al. 2014, *A&A*, 565, A3
- Hopkins, P. F. 2012, *MNRAS*, 423, 2037
- Imanishi, M., Nakanishi, K., & Izumi, T. 2017, *ApJ*, 849, 29
- Ishibashi, W. & Fabian, A. C. 2012, *MNRAS*, 427, 2998
- Ishibashi, W., Fabian, A. C., & Canning, R. E. A. 2013, *MNRAS*, 431, 2350
- Ishibashi, W., Fabian, A. C., & Maiolino, R. 2018, *MNRAS*, 476, 512
- Kauffmann, J., Goldsmith, P. F., Melnick, G., et al. 2017, *A&A*, 605, L5
- Leighly, K. M., Terndrup, D. M., Baron, E., et al. 2014, *ApJ*, 788, 123
- Leroy, A. K., Bolatto, A. D., Ostriker, E. C., et al. 2015, *ApJ*, 801, 25
- Leroy, A. K., Usero, A., Schrubba, A., et al. 2017, *ApJ*, 835, 217
- Lindberg, J. E., Aalto, S., Müller, S., et al. 2016, *A&A*, 587, A15
- Lipari, S., Sanchez, S. F., Bergmann, M., et al. 2009, *MNRAS*, 392, 1295
- Lipari, S., Terlevich, R., Zheng, W., et al. 2005, *MNRAS*, 360, 416
- Maiolino, R., Russell, H. R., Fabian, A. C., et al. 2017, *Nature*, 544, 202
- Meier, D. S., Walter, F., Bolatto, A. D., et al. 2015, *ApJ*, 801, 63
- Meijerink, R., Spaans, M., & Israel, F. P. 2007, *A&A*, 461, 793
- Morganti, R., Veilleux, S., Oosterloo, T., Teng, S. H., & Rupke, D. 2016, *A&A*, 593, A30
- Murray, N., Quataert, E., & Thompson, T. A. 2005, *ApJ*, 618, 569
- Nardini, E., Risaliti, G., Watabe, Y., Salvati, M., & Sani, E. 2010, *MNRAS*, 405, 2505
- Nims, J., Quataert, E., & Faucher-Giguère, C.-A. 2015, *MNRAS*, 447, 3612
- Nishimura, Y., Shimonishi, T., Watanabe, Y., et al. 2016, *ApJ*, 818, 161
- Oosterloo, T., Raymond Oonk, J. B., Morganti, R., et al. 2017, *A&A*, 608, A38
- Papadopoulos, P. P., van der Werf, P. P., Xilouris, E. M., et al. 2012, *MNRAS*, 426, 2601
- Pereira-Santaella, M., Colina, L., García-Burillo, S., et al. 2018, *A&A*, 616, A171
- Pérez-Beaupuits, J. P., Aalto, S., & Gerebro, H. 2007, *A&A*, 476, 177
- Pety, J., Guzmán, V. V., Orkisz, J. H., et al. 2017, *A&A*, 599, A98
- Piconcelli, E., Miniutti, G., Ranalli, P., et al. 2013, *MNRAS*, 428, 1185
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016, *A&A*, 594, A13
- Privon, G. C., Herrero-Illana, R., Evans, A. S., et al. 2015, *ApJ*, 814, 39
- Richings, A. J. & Faucher-Giguère, C.-A. 2018, *MNRAS*, 474, 3673
- Rodríguez del Pino, B., Arribas, S., Piqueras López, J., Villar-Martín, M., & Colina, L. 2019, *MNRAS*, 486, 344
- Rodríguez-Franco, A., Martín-Pintado, J., & Fuente, A. 1998, *A&A*, 329, 1097
- Rosolowsky, E., Engargiola, G., Plambeck, R., & Blitz, L. 2003, *ApJ*, 599, 258
- Rupke, D. S. N. & Veilleux, S. 2011, *ApJ*, 729, L27+
- Sakamoto, K., Aalto, S., Combes, F., Evans, A., & Peck, A. 2014, *ApJ*, 797, 90
- Salas, P., Galaz, G., Salter, D., et al. 2014, *ApJ*, 797, 134
- Schneider, E. E., Robertson, B. E., & Thompson, T. A. 2018, *ApJ*, 862, 56
- Scoville, N. Z. & Solomon, P. M. 1974, *ApJ*, 187, L67
- Shimajiri, Y., André, P., Braine, J., et al. 2017, *A&A*, 604, A74
- Shirley, Y. L. 2015, *PASP*, 127, 299
- Sijacki, D., Springel, V., Di Matteo, T., & Hernquist, L. 2007, *MNRAS*, 380, 877
- Silk, J. & Rees, M. J. 1998, *A&A*, 331, L1
- Skatrud, D. D., De Lucia, F. C., Blake, G. A., & Sastry, K. V. L. N. 1983, *Journal of Molecular Spectroscopy*, 99, 35
- Solomon, P. M. & Vanden Bout, P. A. 2005, *ARA&A*, 43, 677
- Sturm, E., González-Alfonso, E., Veilleux, S., et al. 2011, *ApJL*, 733, L16
- Tafalla, M. & Hacar, A. 2013, *A&A*, 552, L9
- Takano, S., Nakajima, T., & Kohn, K. 2019, *PASJ*, 65
- Taylor, G. B., Silver, C. S., Ulvestad, J. S., & Carilli, C. L. 1999, *The Astrophysical Journal*, 519, 185
- Teng, S. H., Brandt, W. N., Harrison, F. A., et al. 2014, *ApJ*, 785, 19
- Teng, S. H., Veilleux, S., & Baker, A. J. 2013, *ApJ*, 765, 95
- Thompson, T. A., Quataert, E., Zhang, D., & Weinberg, D. H. 2016, *MNRAS*, 455, 1830
- van der Tak, F. F. S., Black, J. H., Schöier, F. L., Jansen, D. J., & van Dishoeck, E. F. 2007, *Astronomy and Astrophysics*, 468, 627
- Veilleux, S., Meléndez, M., Sturm, E., Graciá-Carpio, J., & Fischer, J. e. a. 2013, *ApJ*, 776, 27
- Veilleux, S., Rupke, D. S. N., Kim, D. C., et al. 2009, *ApJS*, 182, 628
- Walter, F., Bolatto, A. D., Leroy, A. K., et al. 2017, *ApJ*, 835, 265
- Wang, X. & Loeb, A. 2018, *New A*, 61, 95
- Watanabe, Y., Nishimura, Y., Harada, N., et al. 2017, *ApJ*, 845, 116
- Watanabe, Y., Sakai, N., Sorai, K., & Yamamoto, S. 2014, *ApJ*, 788, 4
- Wilson, C. D. 2018, *MNRAS*, 477, 2926
- Yamagishi, M., Nishimura, A., Fujita, S., et al. 2018, *ApJS*, 235, 9
- Zubovas, K. & King, A. 2012, *ApJ*, 745, L34
- Zubovas, K. & King, A. R. 2014, *MNRAS*, 439, 400
- Zubovas, K., Nayakshin, S., King, A., & Wilkinson, M. 2013, *MNRAS*, 433, 3079




Fig. A.1: CO(1-0) and CN(1-0) emission line spectra extracted from a circular aperture with  $7''$  radius centred at RA(J2000)=12:56:14.207, Dec(J2000)=56.52.30.994, sampling the northern feature of CN emission visible in the left panel of Fig. 1.

## Appendix A: A possible northern extension of the molecular outflow

The interferometric map of the total CN(1-0) line emission shown in the left panel of Figure 1 presents an extended feature, offset to the North by  $\sim 5''$  with respect to Mrk 231's molecular disk. As noted in the § 1, Mrk 231 hosts multiple extranuclear shells and bubbles that are associated with feedback processes (e.g. Lípari et al. 2005; Lipari et al. 2009; Morganti et al. 2016). However, most of the structures identified so far are located *South* of the nucleus, whereas such northern CN emission does not overlap with any other known optical or radio continuum component.

To further study the origin of this emission, we show in Figure A.1 the CO(1-0) and CN(1-0) line spectra extracted from a  $7''$ -radius aperture, which samples the full extent of this northern CN component down to the  $2\sigma$  level. The fluxes measured by integrating the spectra are  $13.0 \pm 0.8 \text{ Jy km s}^{-1}$  and  $6 \pm 2 \text{ Jy km s}^{-1}$ , respectively for CO(1-0) and CN(1-0). Most of the emission is blue-shifted in both tracers, with prominent broad blue-shifted components. The  $L'_{\text{CN}(1-0)}/L'_{\text{CO}(1-0)}$  ratio is  $0.47 \pm 0.16$ , which is intermediate between the values measured in the narrow core of Mrk 231's integrated line emission and in the broad wings (see Fig. 3).

Hence, overall, the CN/CO line luminosity ratio and the broad and blueshifted line profiles suggest that the northern structure may trace an additional, extended ( $r > 5 \text{ kpc}$ ) component of the molecular outflow. The nature of such emission needs to be investigated and confirmed with dedicated follow-up observations. However, the outflow hypothesis is also corroborated by the detection, both in molecular (Aalto et al. 2012b; Cicone et al. 2012; Feruglio et al. 2015), and in neutral atomic gas tracers (Rupke & Veilleux 2011), of high velocity gas up to  $\sim 2 - 3 \text{ kpc}$  North of the molecular disk, in a region bridging between the newly discovered CN-bright northern component and the well-known kpc-scale central molecular outflow.

## Appendix B: Literature sample with CN(1-0) observations

Table B.1: Literature Sample

Name (1)	CO(1-0) (2)	CN(1-0) (3)	HCN(1-0) (4)	HNC(1-0) (5)	HCO <sup>+</sup> (1-0) (6)	Type (7)	Units (8)	Ref (9)
Arp220	470.7 ± 0.9	44.3 ± 0.9	54.9 ± 1.2	38.6 ± 1.7	22.9 ± 1.0	I	Jy km/s	a
IC694	283.5	13 ± 3 (*)	11.8 ± 1.8	4.9 ± 1.2	19.2 ± 1.9	I	Jy km/s	b
NGC3690	290 ± 30	61 ± 12 (*)	14.	---	11.1	I	Jy km/s	b,c
NGC1614	201.6 ± 0.9	11.5 ± 1.1	4.7 ± 1.7	1.4 ± 1.4	9.0 ± 1.8	I	Jy km/s	d
NGC2146	1150 ± 30	49 ± 12 (*)	< 18.7	7.5 ± 1.4	19.7 ± 3.3	I	Jy km/s	b,c
NGC2623	161	13.4	7.8 ± 0.7	---	4.0 ± 0.7	I	Jy km/s	e,f
NGC6240	327 ± 3	34.0 ± 1.7	16.7 ± 0.9	3.3 ± 0.8	27.1 ± 1.0	I	Jy km/s	d,g
NGC3256 <sup>N</sup>	1168 ± 3	280 ± 3	197 ± 1.0	110.2 ± 0.9	205.3 ± 1.0	N	K km/s	h,i,j
NGC3256 <sup>S</sup>	1344 ± 3	73 ± 3	57.0 ± 1.0	28.1 ± 0.9	87.9 ± 1.0	N	K km/s	h,i,j
NGC3256 <sup>C</sup>	1037 ± 2	95 ± 3	93.2 ± 1.0	389.0 ± 0.9	139.5 ± 0.9	OF	K km/s	h,i,j
NGC3256 <sup>TNE</sup>	487 ± 2	25 ± 3	37.3 ± 1.0	16.4 ± 0.9	71.9 ± 1.0	TT	K km/s	h,i,j
NGC3256 <sup>TSE</sup>	749 ± 2	25 ± 3	36.6 ± 1.0	13.8 ± 0.9	70.5 ± 1.0	TT	K km/s	h,i,j
NGC3256 <sup>TSW</sup>	990 ± 2	23 ± 3	51.9 ± 1.0	17.5 ± 0.9	84.5 ± 1.0	TT	K km/s	h,i,j
NGC3256 <sup>OS</sup>	537 ± 3	33 ± 3	40.4 ± 1.0	12.3 ± 0.9	64.6 ± 1.0	OF	K km/s	h,i,j
NGC3256 <sup>SW</sup>	1317 ± 2	67 ± 3	76.7 ± 1.0	33.3 ± 0.9	124.1 ± 1.0	N	K km/s	h,i,j
NGC7130	257 ± 14	20 ± 4 (*)	16.5	10.0 ± 1.2	---	I	Jy km/s	b,c
NGC1808	1900 ± 140	154 ± 4 (*)	107 ± 5	45 ± 11	59 ± 4	I	Jy km/s	b
NGC660	659.0 ± 1.8	64.6 ± 1.6	28.7 ± 1.2	15.0 ± 1.2	29.9 ± 1.2	I	Jy km/s	d
NGC1068 <sup>CND</sup>	155.0 ± 1.0	46.1 ± 0.5	26.4 ± 0.6	5.6 ± 0.4	13.3 ± 0.7	N	K km/s	k
05414+5840	411.1	17.4 ± 0. (*)	15 ± 3	< 3.8	9 ± 3	I	Jy km/s	b
12243-0036	165.4	14.1 ± 2.7 (*)	13.0 ± 1.6	13.4 ± 1.6	9.7 ± 1.6	I	Jy km/s	b
M83	1121 ± 14	94.1 ± 0.7	50.5 ± 0.6	20.4 ± 0.4	46.1 ± 0.7	I	Jy km/s	a
15107+0724	118.1	11.4 (*)	13 ± 4	7 ± 4	7 ± 2	I	Jy km/s	b
17208-0014	171.	14.5	11.8 ± 0.8	9.1 ± 1.0	9.1 ± 0.8	I	Jy km/s	d,f
NGC253 <sup>tot</sup>	870. ± 3	150.0 ± 1.4	70.0 ± 1.0	30.0 ± 1.0	59.0 ± 1.0	I	K km/s	k
NGC253 <sup>SB</sup>	16000 ± 700	2540 ± 80	2420 ± 110	---	2090 ± 100	N	K km/s	l
NGC253 <sup>r1</sup>	720 ± 70	33 ± 3	24 ± 2	---	27 ± 3	OF	K km/s	l
NGC253 <sup>r10</sup>	1920 ± 190	112 ± 9	179 ± 18	---	153 ± 15	OF	K km/s	l
NGC253 <sup>r2</sup>	1920 ± 190.	76 ± 6	119 ± 12	---	76 ± 8	OF	K km/s	l
IC342 <sup>tot</sup>	160.0 ± 0.4	20.4 ± 0.4	11.60 ± 0.10	4.5 ± 0.10	8.4 ± 0.10	N	K km/s	k
M82	2658 ± 14.	213 ± 2	117 ± 2	52.0 ± 1.1	167 ± 4	I	Jy km/s	a
NGC4945	9886.6 ± 28.4	1134 ± 30 (*)	420 ± 8	225 ± 4	450 ± 4	I	Jy km/s	b
AM2055-425	49	1.5 ± 0.3 (*)	---	---	---	I	Jy km/s	m
AM2246-490	34	2.4 ± 0.3 (*)	---	---	---	I	Jy km/s	m
VV114	613	6.0 (*)	---	---	---	I	Jy km/s	m
AM1300-233	89	2.6 ± 0.3 (*)	---	---	---	I	Jy km/s	m
NGC1377	45 ± 5	0.48 ± 0.12 (*)	---	---	---	I	Jy km/s	m
IC860	73.1	4.6	6.7 ± 0.7	3.6 ± 0.6	4.2 ± 0.6	I	Jy km/s	d,f
NGC3079	1090 ± 50	32 ± 4	25.0 ± 1.3	6.6 ± 0.9	28.1 ± 1.4	I	Jy km/s	d
NGC4194	171.3 ± 1.0	10.0 ± 1.2	2.8 ± 0.4	1.5 ± 0.4	3.7 ± 0.4	I	Jy km/s	d
NGC7771	464.0 ± 1.4	23 ± 2	20.3 ± 0.9	12.0 ± 1.0	18.5 ± 0.9	I	Jy km/s	d
NGC3556	251.2 ± 1.8	6.0 ± 0.8	3.2 ± 0.5	1.0 ± 0.5	5.1 ± 0.5	I	Jy km/s	d
NGC7674	121.1 ± 1.2	9.6 ± 1.5	< 1.1	< 1.1	< 1.1	I	Jy km/s	d
UGC2866	427.8 ± 1.7	16.0 ± 1.2	8.8 ± 0.7	4.5 ± 0.8	12.9 ± 0.7	I	Jy km/s	d
UGC5101	96.9 ± 1.8	22.1 ± 1.6	6.6 ± 0.9	5.4 ± 1.1	2.4 ± 1.0	I	Jy km/s	d
M51	249 ± 8	38.0 ± 0.4	26.7 ± 0.5	8.9 ± 0.3	13.2 ± 0.3	I	Jy km/s	a
M51_P1	68.6 ± 0.3	3.2 ± 0.08	2.74 ± 0.09	0.96 ± 0.06	2.04 ± 0.08	I	K km/s	n
M51_P2	54.1 ± 0.2	1. ± 0.3	1.70 ± 0.10	0.63 ± 0.07	1.38 ± 0.09	I	K km/s	n
NGC7469	288 ± 7	29 ± 2	11.7 ± 0.6	5.4 ± 0.6	14.3 ± 0.7	I	Jy km/s	a
IIIZw35	70.6	8.5	6.5 ± 0.9	---	3.0 ± 0.6	I	Jy km/s	e,f
05189-2524	25.7	6.7	4.4 ± 0.8	2.3 ± 0.7	3.3 ± 0.8	I	Jy km/s	f,o
09111-1007 <sup>w</sup>	51	6	3.6 ± 0.5	< 1.5	5.0 ± 1.0	I	Jy km/s	f,o
10173+0828	24.2	2	2.6 ± 0.7	---	1.7 ± 0.7	I	Jy km/s	e,f
12224-0624	22.6	3.1	< 3.3	< 4.0	< 4.0	I	Jy km/s	f,o
NGC4418	117	15.1	12.9 ± 0.9	6.0 ± 0.8	7.6 ± 0.8	I	Jy km/s	d,f
13120-5453	234	39.2	---	---	---	I	Jy km/s	f
14378-3651	24.7	3	---	---	---	I	Jy km/s	f

Follows on next page

Name (1)	CO(1-0) (2)	CN(1-0) (3)	HCN(1-0) (4)	HNC(1-0) (5)	HCO <sup>+</sup> (1-0) (6)	Type (7)	Units (8)	Ref (9)
20551-4250	54.9	3	1.4 ± 0.1	0.37 ± 0.04	2.20 ± 0.10	I	Jy km/s	f, p
NGC7479	270	9 <sup>(*)</sup>	< 20.2	---	---	I	Jy km/s	f, q
W51GMC	172 ± 4	16.0 ± 1.2	5.5 ± 0.2	2.4 ± 0.3	4.5 ± 0.2	C	K km/s	r
W51GMCA	1617 ± 11	124 ± 7	62.0 ± 1.0	40. ± 1.0	54. ± 2	C	K km/s	r
W51GMCB	748 ± 6	55 ± 4	30. ± 0.6	16.3 ± 0.6	23.4 ± 0.6	C	K km/s	r
W51GMCC	369 ± 6	23 ± 2	11.8 ± 0.5	5.6 ± 0.6	9.3 ± 0.5	C	K km/s	r
W51GMCD	186 ± 6	16 ± 2	5.2 ± 0.3	2.2 ± 0.3	4.5 ± 0.3	C	K km/s	r
W51GMCE	77 ± 3	13 ± 1.4	2.1 ± 0.3	0.6 ± 0.3	1.8 ± 0.2	C	K km/s	r
W51e1/e2	1375 ± 6	154 ± 3	77.2 ± 0.5	50.0 ± 0.4	75.2 ± 0.5	C	K km/s	r
OrionB	60.43	1.16 <sup>(*)</sup>	1.54	0.445	1.63	C	K km/s	s
LMC_N79	16.80 ± 0.06	0.68 ± 0.09 <sup>(*)</sup>	1.51 ± 0.03	0.45 ± 0.03	2.797 ± 0.019	C	K km/s	t
LMC_N44C	25.58 ± 0.05	0.51 ± 0.05 <sup>(*)</sup>	1.192 ± 0.017	0.383 ± 0.013	2.164 ± 0.015	C	K km/s	t
LMC_N113	46.96 ± 0.07	0.90 ± 0.12 <sup>(*)</sup>	3.15 ± 0.05	1.39 ± 0.03	4.29 ± 0.04	C	K km/s	t
OrionA	3.0 ± 1.2 × 10 <sup>4</sup>	140 ± 600 <sup>(*)</sup>	1000 ± 400	---	---	C	K km/s pc <sup>2</sup>	u

**Notes.** Col.(1): source name. Cols.(2-6): integrated line fluxes of the corresponding transitions. Col.(7): source type. I= integrated observation; N= nucleus; OF=outflow region; TT: tidal tail; C = giant molecular cloud. Col.(8): units of the fluxes reported in Cols.(2-6); Col.(9): references: (a) Aladro et al. (2015); (b) Baan et al. (2008); (c) Aalto et al. (2002); (d) Costagliola et al. (2011); (e) García-Burillo et al. (2012); (f) Lutz et al. 2019 (in prep); (g) Papadopoulos et al. (2012); (h) Harada et al. (2018); (i) Sakamoto et al. (2014); (j) Harada, Sakamoto (Private Comm.); (k) Takano et al. (2019); (l) Meier et al. (2015); (m) Wilson (2018); (n) Watanabe et al. (2014); (o) Privon et al. (2015); (p) Imanishi et al. (2017); (q) Gao & Solomon (2004); (r) Watanabe et al. (2017); (s) Pety et al. (2017); (t) Nishimura et al. (2016); (u) Kauffmann et al. (2017) and J. Kauffmann (priv. comm.); (\*) In these cases only the flux of the CN(1-0)  $J = 3/2 - 1/2$  spingroup is available, and it has been multiplied by a factor of 1.5 to evaluate the total CN(1-0) flux.