SOLVING EQUATION SYSTEMS IN ω -CATEGORICAL ALGEBRAS

MANUEL BODIRSKY AND THOMAS QUINN-GREGSON

ABSTRACT. We study the computational complexity of deciding whether a given set of term equalities and inequalities has a solution in an ω -categorical algebra \mathfrak{A} . There are ω -categorical groups where this problem is undecidable. We show that if \mathfrak{A} is an ω -categorical semilattice or an abelian group, then the problem is in P or NP-hard. The hard cases are precisely those where $\operatorname{Pol}(\mathfrak{A}, \neq)$ has a uniformly continuous minor-preserving map to the clone of projections on a two-element set. The results provide information about algebras \mathfrak{A} such that $\operatorname{Pol}(\mathfrak{A}, \neq)$ does not satisfy this condition, and they are of independent interest in universal algebra. In our proofs we rely on the Barto-Pinsker theorem about the existence of pseudo-Siggers polymorphisms. To the best of our knowledge, this is the first time that the pseudo-Siggers identity has been used to prove a complexity dichotomy.

1. Introduction

The problem of deciding whether a given system of linear equations has a solution in \mathbb{Z}_p is one of the central computational problems that can be solved in polynomial time, for example by Gaussian elimination. The problem can also be rephrased as follows: fix the structure $(\mathbb{Z}_p; +, 0, 1, \dots, p-1)$ where + is the binary addition operation and $0, 1, \dots, p-1$ are constants; the problem is then to decide whether a given conjunction of atomic formulas in the signature of this structure is satisfiable in this structure. Analogous computational problems can be formulated for other algebraic structures \mathfrak{A} instead of $(\mathbb{Z}_p; +, 0, \dots, p-1)$, and have been studied systematically in the special cases of groups [26], monoids [36], and semigroups [30].

An even more general class of computational problems is the class of constraint satisfaction problems (CSPs); here we fix a structure \mathfrak{A} with a finite signature τ , and the task is to decide whether a given conjunction of atomic τ -formulas is satisfiable in \mathfrak{A} . This problem, denoted by CSP(\mathfrak{A}), is typically introduced only for relational signatures; the restriction to relational signatures is not severe, because we may replace each operation f of arity kin \mathfrak{A} by the k+1-ary relation $R_f := \{(x_1, \ldots, x_k, x_0) \mid x_0 = f(x_1, \ldots, x_k)\}$. Then every atomic formula over \mathfrak{A} can be translated into a finite set of atomic formulas in the new signature to obtain a satisfiability-equivalent instance in the new signature. We might have to introduce some additional variables to eliminate nested terms in atomic formulas, but the overall reduction changes the size of the input only by a linear factor.

It has been conjectured by Feder and Vardi [25] that CSPs for fixed structures \mathfrak{A} with a finite domain have a *complexity dichotomy* in the sense that they are either in P or

Both authors have received funding from the Deutsche Forschungsgemeinschaft (DFG) and from the European Research Council (Grant Agreement no. 681988, CSP-Infinity).

NP-complete. The dichotomy conjecture has been confirmed recently, independently by Bulatov [20] and by Zhuk [46]. This achievement has been made possible because of an important link between constraint satisfaction and central topics in universal algebra; see, e.g., the survey articles in [31].

There are many famous computational problems that can be phrased as solving equation systems over algebraic structures \mathfrak{A} with an infinite domain; for example, $CSP(\mathfrak{A})$ for the structure $\mathfrak{A}=(\mathbb{Z};+,\cdot,1)$ is Hilbert's tenth problem, and known to be undecidable [35], whereas the problem can be solved in polynomial time for $\mathfrak{A}=(\mathbb{Z};+,1)$ (see, e.g., [42]). In full generality, the mentioned connection between constraint satisfaction and universal algebra breaks down (see the survey article [13]). However, if the structure \mathfrak{A} is ω -categorical, i.e., if all countably infinite models of the first-order theory of $\mathfrak A$ are isomorphic, then the universal-algebraic approach is still applicable [5, 14]. Note that when studying the CSP of infinite-domain structures $\mathfrak A$ we still require the signature of $\mathfrak A$ to be finite. In particular, we no longer have constants for every element in the domain. If the signature contains no constants at all, then solving equation systems becomes trivial for many algebraic structures: for instance for monoids with unit element 1, we might satisfy all the equations by setting all variables to 1. The natural signature for studying the problem of solving equations over infinite domains is to additionally allow inequalities in the input, i.e., atomic formulas of the form $s \neq t$ where s and t are terms. In this article we study problems of the form $CSP(\mathfrak{A},\neq)$ where \mathfrak{A} is a finite-signature algebra¹. For example, for a given monoid \mathfrak{A} , the problem $CSP(\mathfrak{A},\neq)$ is non-trivial in general since we may no longer map all the variables in the input to 1.

1.1. **Applications.** If $CSP(\mathfrak{A}, \neq)$ can be solved in polynomial time, then various other interesting computational problems can be solved in polynomial time, too. Let \mathfrak{A} be an algebra with a finite signature τ and a (finite or infinite) domain A. The *Identity Checking Problem (for* \mathfrak{A}) is the problem of deciding whether for given τ -terms s, t over the variables x_1, \ldots, x_n the identity $s \approx t$ is valid in \mathfrak{A} , i.e., whether

(1.1)
$$\mathfrak{A} \models \forall x_1, \dots, x_n \colon s(x_1, \dots, x_n) = t(x_1, \dots, x_n).$$

Note that this is the case if and only if there are no elements $a_1, \ldots, a_n \in A$ such that $\mathfrak{A} \models s(a_1, \ldots, a_n) \neq t(a_1, \ldots, a_n)$; by introducing additional variables and equations we can translate this into an instance of $CSP(\mathfrak{A}, \neq)$ which is unsatisfiable if and only if (1.1) holds. Hence, if $CSP(\mathfrak{A}, \neq)$ is in NP, then the Identity Checking Problem for \mathfrak{A} is in CoNP, and if $CSP(\mathfrak{A}, \neq)$ is in P, then the Identity Checking Problem for \mathfrak{A} is in P, too.

In the so-called Entailment Problem (for \mathfrak{A}) we are given a finite set of equations $s_1 = t_1, \ldots, s_m = t_m$ and another equation $s_0 = t_0$ over a common set of variables V, and the question is whether every assignment $V \to A$ that satisfies $s_1 = t_1 \wedge \cdots \wedge s_m = t_m$ also satisfies $s_0 = t_0$. Note that this is the case if and only if the formula $s_1 = t_1 \wedge \cdots \wedge s_m = t_m \wedge s_0 \neq t_0$ is unsatisfiable, so again the problem reduces in polynomial time to the complement of $CSP(\mathfrak{A}, \neq)$.

¹An algebra is simply a structure with a purely functional signature; see [27] for basic terminology.

Finally, there is a strong link between $\mathrm{CSP}(\mathfrak{A},\neq)$ and the problem $\mathrm{CSP}(\mathfrak{A},a_1,\ldots,a_n)$, where $a_1,\ldots,a_n\in A$ are constants, if the algebra \mathfrak{A} is model-complete. The notion of model-completeness is a central concept from model theory and can be seen as a weak form of quantifier elimination: \mathfrak{A} is model-complete if every first-order sentence is equivalent to an existential sentence over \mathfrak{A} . It follows from results in [7,12] that if \mathfrak{A} is model-complete, then for all $a_1,\ldots,a_n\in A$ the problem $\mathrm{CSP}(\mathfrak{A},\neq)$ and the problem $\mathrm{CSP}(\mathfrak{A},\neq,a_1,\ldots,a_n)$ are polynomial-time equivalent; in particular, there is a polynomial-time reduction from $\mathrm{CSP}(\mathfrak{A},a_1,\ldots,a_n)$ to $\mathrm{CSP}(\mathfrak{A},\neq)$. Conversely, we will see that if \mathfrak{A} satisfies an additional assumption, namely \mathfrak{A} that there is an embedding $\mathfrak{A}^2 \hookrightarrow \mathfrak{A}$ (i.e., an isomorphism between $\mathfrak{A}^2 = \mathfrak{A} \times \mathfrak{A}$ and a substructure of \mathfrak{A}), then there are $a_1,\ldots,a_n\in A$ such that there is a polynomial-time reduction from $\mathrm{CSP}(\mathfrak{A},\neq)$ to $\mathrm{CSP}(\mathfrak{A},a_1,\ldots,a_n)$ (Proposition 3.8).

- 1.2. **Results.** We initiate the study the computational complexity of $CSP(\mathfrak{A}, \neq)$ for ω -categorical algebras \mathfrak{A} . We first observe that there are ω -categorical groups \mathfrak{A} such that $CSP(\mathfrak{A}, \neq)$ is undecidable (Section 5.2). For abelian ω -categorical groups, however, we show that $CSP(\mathfrak{A}, \neq)$ is in P or NP-complete (Theorem 5.16). Recall that if P and NP are distinct then there are also problems in NP that are of intermediate complexity, i.e., neither in P nor NP-hard ([32]). We also show a P versus NP-hard complexity dichotomy for ω -categorical semilattices (Theorem 6.5).
- 1.3. Outline. In our proofs we rely on recent universal-algebraic results for ω -categorical structures, in particular from [2,4,5], so we start by giving a self-contained introduction to the universal-algebraic approach in Section 2. Universal-algebraic concepts are also needed to precisely state the border between the NP-hard and the polynomial cases in our results. In Section 3 we specialise the universal-algebraic approach to structures of the form (\mathfrak{A}, \neq) where \mathfrak{A} is an algebra, and in Section 4 we specialise further to monoids. Section 5 contains our classification for ω -categorical abelian groups. Finally, Section 6 contains our classification for ω -categorical semilattices. We close with a discussion and some open problems in Section 7.

2. The Universal-Algebraic Approach

The universal-algebraic approach is based on the following concept from universal algebra. An operation $f: A^k \to A$ preserves a relation $R \subseteq A^m$ if for all $t_1, \ldots, t_k \in R$ the m-tuple $f(t_1, \ldots, t_k)$ obtained from applying f componentwise is also contained in R. Note that if $g: A^m \to A$ is an operation, then f preserves the $graph R_g$ of g, defined as

$$R_g := \{(a_1, \dots, a_m, g(a_1, \dots, a_m)) \mid a_1, \dots, a_m \in A\},\$$

if and only if f commutes with g, i.e., for all $a_{1,1}, \ldots, a_{n,m} \in A$

$$f(g(a_{1,1},\ldots,a_{1,m}),\ldots,g(a_{n,1},\ldots,a_{n,m}))$$

= $g(f(a_{1,1},\ldots,a_{n,1}),\ldots,f(a_{1,m},\ldots,a_{n,m})).$

²This property is equivalent to a property that is often referred to as *convexity* in the theoretical computer science literature [37] and will play an important role in this article.

In this case we say that f preserves g. An operation f is a polymorphism of a structure \mathfrak{A} if f preserves all relations and all operations of \mathfrak{A} . Note that the projections $\pi_i^k \colon A^k \to A$ defined by $\pi_i^k(a_1,\ldots,a_k) := a_i$ is a polymorphism of every structure with domain A. We also would like to mention that similarly as the set of all automorphisms of \mathfrak{A} forms a group, the set of all polymorphisms of \mathfrak{A} , denoted by $\operatorname{Pol}(\mathfrak{A})$, forms a clone, i.e., the set of polymorphisms is closed under composition and contain the projections. The clone of projections on a two-element set will be denoted by \mathscr{P} . A map between two clones is called minor-preserving if it maps operations to operations of the same arity, and if

$$\xi(f(p_1,\ldots,p_n))=\xi(f)(p_1,\ldots,p_n)$$

for all n-ary operations f and projections p_1, \ldots, p_n of the same arity m. In the introduction we have mentioned that $CSP(\mathfrak{A})$ is for every finite structure \mathfrak{A} with finite relational signature in P or NP-complete; using polymorphisms, the border between the two cases can be stated as follows, combining results from [4, 20, 22, 43, 46].

Theorem 2.1. Let $\mathfrak A$ be a structure with finite domain and finite relational signature. Then either

• \mathfrak{A} has a polymorphism $s \colon A^6 \to A$ which is Siggers, i.e., satisfies

$$s(x, y, x, z, y, z) \approx s(y, x, z, x, z, y)$$

in this case, $CSP(\mathfrak{A})$ is in P, or

• $Pol(\mathfrak{A})$ has a minor-preserving map to \mathscr{P} ; in this case, $CSP(\mathfrak{A})$ is NP-complete.

The fact that $CSP(\mathfrak{A})$ is NP-hard if \mathfrak{A} does not have a Siggers polymorphism [43] was already known before the break-through result from [20,46]. The equivalence of the existence of a Siggers polymorphism and of the non-existence of a minor-preserving map to \mathscr{P} is from [4]. For general ω -categorical structures, the equivalence is no longer valid [2], but we still have the following hardness condition.

Theorem 2.2 ([4]). Let \mathfrak{A} be an ω -categorical structure with a finite relational⁴ signature. If $\operatorname{Pol}(\mathfrak{A})$ has a uniformly continuous³ minor-preserving map to \mathscr{P} then $\operatorname{CSP}(\mathfrak{A})$ is NP-hard.

To apply this hardness condition, we need the following terminology from [7]. An ω -categorical structure $\mathfrak A$ is called a *core* if every endomorphism of $\mathfrak A$ (i.e., every homomorphism from $\mathfrak A$ to $\mathfrak A$) is an embedding. Two structures $\mathfrak A$ and $\mathfrak B$ are called *homomorphically equivalent* if there is a homomorphism from $\mathfrak A$ to $\mathfrak B$ and vice versa. Clearly, two structures that are homomorphically equivalent have the same CSP.

Theorem 2.3 ([7,12]). Every ω -categorical relational structure \mathfrak{B} is homomorphically equivalent to a model-complete core structure \mathfrak{C} , which is unique up to isomorphism, and again ω -categorical, and which will be called the model-complete core of \mathfrak{C} .

³In our setting, $\xi \colon \mathscr{C} \to \mathscr{P}$ is uniformly continuous if and only if there exists a finite set $F \subseteq C$ such that if $f, g \in \mathscr{C}$ agree on F, then $\xi(f) = \xi(g)$.

A first-order formula ϕ is called *primitive positive* if it is of the form

$$\exists \bar{x}(\phi_1(\bar{x}) \wedge \cdots \wedge \phi_n(\bar{x})),$$

where ϕ_1, \ldots, ϕ_n are atomic formulas. Every primitive positive relation of a relational structure \mathfrak{B} is preserved by the polymorphisms of \mathfrak{B} . If \mathfrak{B} is ω -categorical then conversely every relation left invariant by polymorphisms of \mathfrak{B} is primitive positive definable [14].

The following is implied by results in [4].

Proposition 2.4. Let \mathfrak{B} be an ω -categorical relational⁴ structure.

- If $\mathfrak C$ is homomorphically equivalent to $\mathfrak B$ then there is a uniformly continuous minor-preserving map from $\operatorname{Pol}(\mathfrak B)$ to $\operatorname{Pol}(\mathfrak C)$.
- If \mathfrak{C} is the model-complete core of \mathfrak{B} and $c_1, \ldots, c_n \in C$, then $\operatorname{Pol}(\mathfrak{C})$ (and $\operatorname{Pol}(\mathfrak{B})$) has a uniformly continuous minor-preserving map to $\operatorname{Pol}(\mathfrak{C}, c_1, \ldots, c_n)$.
- If $\mathfrak A$ is a substructure of $\mathfrak B$ whose domain is primitive positive definable in $\mathfrak B$, then there is a uniformly continuous minor-preserving map from $\operatorname{Pol}(\mathfrak A)$ to $\operatorname{Pol}(\mathfrak A)$.

Hence, if $\operatorname{Pol}(\mathfrak{C}, c_1, \ldots, c_n)$ or $\operatorname{Pol}(\mathfrak{A})$ in Proposition 2.4 has a uniformly continuous minor-preserving map to \mathscr{P} , then $\operatorname{CSP}(\mathfrak{B})$ is NP-hard by Theorem 2.2, because the composition of uniformly continuous minor-preserving maps is uniformly continuous and minor-preserving. For model-complete cores, we will use the following result.

Theorem 2.5 (Barto and Pinsker [5]). Let \mathfrak{C} be an ω -categorical relational⁴ structure which is a model-complete core. Then at least one of the following holds.

• \mathfrak{C} has a pseudo-Siggers polymorphism, i.e., a polymorphism $s: C^6 \to C$ and endomorphisms $e_1, e_2: C \to C$ satisfying

$$e_1(s(x,y,x,z,y,z)) \approx e_2(s(y,x,z,x,z,y)).$$

• $Pol(\mathfrak{C})$ has a uniformly continuous minor-preserving map to \mathscr{P} .

In this article we will show how to use the pseudo-Siggers identity to obtain structural information about \mathfrak{C} if \mathfrak{C} is of the form (\mathfrak{A}, \neq) where \mathfrak{A} is an ω -categorical semilattice or abelian group.

Remark 2.6. In many situations, the two items in Theorem 2.5 are mutually exclusive; two general conditions that imply this have been presented in [2]. However, these conditions do not cover our setting, not even in the special case of semilattices. Abelian groups are covered, but this requires an extra argument that will be given in Section 5.5.

3. Algebras

An algebra \mathfrak{A} is a structure with domain A and with a purely functional signature. The n-ary polymorphisms of \mathfrak{A} are precisely the (algebra) homomorphisms $g:A^n\to A$. In this section we make some observations that are relevant for the universal-algebraic approach to the CSP of structures of the form (\mathfrak{A}, \neq) .

⁴It will be explained in Remark 3.5 that the result also holds for general structures that also might contain operations.

Conventions. We write $\omega = \{0, 1, 2, \dots\}$ for the set of natural numbers including zero. The equality symbol is always allowed in first-order formulas.

- 3.1. Homogeneity. An important source of ω -categorical structures comes from Fraïssé-amalgamation. The age of a τ -structure is the class of all finitely generated τ -structures that embed into the structure. A structure is called homogeneous if every isomorphism between finitely generated substructures extends to an automorphism. Let τ be a countable signature and let \mathcal{K} be a class of finitely generated τ -structures which is closed under subalgebras, has the joint embedding property and the amalgamation property, and contains countably many isomorphism types of structures. Then there exists a countable homogeneous τ -structure $\mathscr F$ whose age is $\mathcal K$ (Theorem 6.1.2. in [28]). A structure $\mathfrak A$ is called uniformly locally finite if there exists a function $f:\omega\to\omega$ such that every substructure of $\mathfrak A$ generated by n elements has at most f(n) elements. If $\mathfrak A$ is ω -categorical then $\mathfrak A$ must be uniformly locally finite. Conversely, every homogeneous uniformly locally finite structure is ω -categorical ([28], Corollary 6.2).
- 3.2. Model companions. Let \mathfrak{A} and \mathfrak{B} be algebras with the same signature. Note that homomorphisms between (\mathfrak{A}, \neq) and (\mathfrak{B}, \neq) must be embeddings (which is not true in general if \mathfrak{A} and \mathfrak{B} are arbitrary structures). It follows that structures of the form (\mathfrak{A}, \neq) must be cores. Two structures \mathfrak{A} and \mathfrak{B} are called *companions* if they satisfy the same universal first-order sentences (for instance, if \mathfrak{A} is a semilattice, so is every companion of \mathfrak{A}). Note that in this case, \mathfrak{A} and \mathfrak{B} have the same age. The implication from (1) to (2) in the following lemma can be shown by a compactness argument (see, e.g., [7]); it is straightforward to prove the other implications in cyclic order.

Lemma 3.1. Let \mathfrak{A} and \mathfrak{B} be ω -categorical algebras. Then the following are equivalent.

- (1) \mathfrak{A} and \mathfrak{B} are companions;
- (2) $\mathfrak{A} \hookrightarrow \mathfrak{B}$ and $\mathfrak{B} \hookrightarrow \mathfrak{A}$;
- (3) (\mathfrak{A}, \neq) and (\mathfrak{B}, \neq) are homomorphically equivalent;
- (4) $CSP(\mathfrak{A}, \neq)$ and $CSP(\mathfrak{B}, \neq)$ are the same computational problem;
- (5) $Age(\mathfrak{A}) = Age(\mathfrak{B})$.

A structure \mathfrak{B} is called a *model companion of* \mathfrak{A} if \mathfrak{A} and \mathfrak{B} are companions and \mathfrak{B} is model-complete. Every ω -categorical structure has a model companion [40], which is unique up to isomorphism and ω -categorical (see, e.g., [27]). For illustration, we present an example of an ω -categorical algebra and its model companion.

Example 3.2. For $a, b \in \mathbb{Q}$ we write [a, b] for $\{x \in \mathbb{Q} \mid a \leq x \leq b\}$ and min for the binary operation that returns the minimum of its two arguments. Then $([0, 1]; \min)$ and $(\mathbb{Q}; \min)$ are companions. Since $(\mathbb{Q}; \min)$ is model-complete, it is the model companion of $([0, 1]; \min)$.

Unfortunately, several of the results that we cited in Section 2 were originally only formulated for relational signatures. But it is not difficult to see that they also hold for structures that might involve operations, as we will see in the following. The definition of model-complete cores for general ω -categorical structures \mathfrak{B} is the same as the one we

gave for the relational case: a structure \mathfrak{C} is a model-complete core of \mathfrak{B} if \mathfrak{C} and \mathfrak{B} are homomorphically equivalent and \mathfrak{C} is a model-complete core.

Let \mathfrak{B} be a structure. We write \mathfrak{B}^* for the relational structure obtained from \mathfrak{B} by replacing each operation g in \mathfrak{B} of arity k by a relation symbol R_g of arity k+1 that denotes in \mathfrak{B}^* the graph of the operation $g^{\mathfrak{B}}$.

Remark 3.3. There are homogeneous algebras \mathfrak{A} such that \mathfrak{A}^* is not homogeneous: for example, consider the group $\mathfrak{A} := \mathbb{Z}_2 \times \mathbb{Z}_3$ generated by an element a of order 2 and an element b of order 3. Then it is easy to verify that \mathfrak{A} is homogeneous in the signature $\{\cdot\}$ of semigroups, but in \mathfrak{A}^* the substructures induced by $\{a\}$ and $\{b\}$ are isomorphic, and no automorphism of \mathfrak{A}^* maps a to b.

Lemma 3.4. Let \mathfrak{B} be an ω -categorical structure. Then \mathfrak{B} has a model-complete core \mathfrak{C} , which is unique up to isomorphism and again ω -categorical. Moreover, \mathfrak{C}^* is the model-complete core of \mathfrak{B}^* .

Proof. By Theorem 2.3, the relational structure \mathfrak{B}^* has a model-complete core \mathfrak{C}' which is ω -categorical. Since \mathfrak{C}' and \mathfrak{B}^* are homomorphically equivalent, there are homomorphisms $h \colon \mathfrak{B}^* \to \mathfrak{C}'$ and $i \colon \mathfrak{C}' \to \mathfrak{B}^*$. For each k-ary function symbol g from the signature τ of \mathfrak{B} , the relation denoted by R_g in \mathfrak{C}' is the graph of a k-ary operation on C'. Indeed⁵ let $\phi(x_1, \ldots, x_n)$ be the formula $\exists z \colon R_g(x_1, \ldots, x_n, z)$ and let $u_1, \ldots, u_n \in C'$. Then there exists $z \in B$ such that $g^{\mathfrak{B}}(i(u_1), \ldots, i(u_n)) = z$. Hence, $(i(u_1), \ldots, i(u_n), z) \in R_g^{\mathfrak{B}^*}$ and thus $(h \circ i(u_1), \ldots, h \circ i(u_n), h(z)) \in R_g^{\mathfrak{C}'}$, so $\mathfrak{C}' \models \phi(h \circ i(u_1), \ldots, h \circ i(u_n))$ and thus $\mathfrak{C}' \models \phi(u_1, \ldots, u_n)$. Moreover, if $(u_1, \ldots, u_n, a), (u_1, \ldots, u_n, b) \in R_g^{\mathfrak{C}'}$ then $(i(u_1), \ldots, i(u_n), i(a)), (i(u_1), \ldots, i(u_n), i(b)) \in R_g^{\mathfrak{B}^*}$, so i(a) = i(b) since $R_g^{\mathfrak{B}^*}$ is the graph of the function $g^{\mathfrak{B}}$. Thus, a = b because i is injective.

Let \mathfrak{C} be the τ -structure with the same domain and relations as \mathfrak{C}' and such that every operation symbol $g \in \tau$ denotes the operation whose graph is $R_g^{\mathfrak{C}'}$. Clearly, \mathfrak{C}^* equals \mathfrak{C}' . We prove that \mathfrak{C} is a model-complete core of \mathfrak{B} : the maps h and i are homomorphisms from \mathfrak{B} to \mathfrak{C} and from \mathfrak{C} to \mathfrak{B} , respectively, showing that \mathfrak{B} and \mathfrak{C} are homomorphically equivalent. Every endomorphism of \mathfrak{C} is an endomorphism of \mathfrak{C}' , and hence preserves all first-order formulas over \mathfrak{C}' and also preserves all first-order formulas over \mathfrak{C} . So \mathfrak{C} is a model-complete core.

If \mathfrak{D} is a model-complete core that is homomorphically equivalent with \mathfrak{B} , then \mathfrak{D}^* is homomorphically equivalent to \mathfrak{C}^* , and hence \mathfrak{D}^* and \mathfrak{C}^* are isomorphic. It follows that \mathfrak{D} and \mathfrak{C} are isomorphic, showing the uniqueness of \mathfrak{C} up to isomorphism.

As in the relational case, because of the uniqueness of the model-complete core up to isomorphism we call \mathfrak{C} the model-complete core of \mathfrak{B} .

Remark 3.5. Similarly as in the proof of Lemma 3.4 it can be shown that the assumption in Theorem 2.2, Proposition 2.4, and Theorem 2.5 that the structures are relational can be dropped.

⁵This would not be true for arbitrary structures \mathfrak{C}' that are homomorphically equivalent to \mathfrak{B}^* , but we will use model-completeness.

Corollary 3.6. Let \mathfrak{A} be an ω -categorial algebra and \mathfrak{C} its model companion. Then (\mathfrak{C}, \neq) is the model-complete core of (\mathfrak{A}, \neq) .

Proof. The structure (\mathfrak{C}, \neq) is a model-complete core and homomorphically equivalent to (\mathfrak{A}, \neq) by Lemma 3.1. So the model-complete core of (\mathfrak{A}, \neq) must be isomorphic to (\mathfrak{C}, \neq) .

3.3. Square embeddings. In theoretical computer science [37], a first-order τ -theory T is called *convex* if for every finite set of atomic τ -formulas S the set

$$T \cup S \cup \{x_1 \neq y_1, \dots, x_m \neq y_m\}$$

is satisfiable if and only if $T \cup S \cup \{x_i \neq y_i\}$ is satisfiable for each $i \in \{1, ..., m\}$. If \mathfrak{A} is a structure, we write $Th(\mathfrak{A})$ for the first-order theory of \mathfrak{A} , i.e., for the set of all first-order sentences that hold in \mathfrak{A} . If $T = Th(\mathfrak{A})$ for some structure \mathfrak{A} , then an alternative terminology [19] for convexity is that \neq is 1-independent from \mathfrak{A} .

Proposition 3.7. Let \mathfrak{A} be an ω -categorical algebra. Then the following are equivalent.

- (1) Th(\mathfrak{A}) is convex;
- (2) A has a binary injective polymorphism;
- (3) $\mathfrak{A}^2 \hookrightarrow \mathfrak{A}$;
- (4) $\mathfrak{A}^k \hookrightarrow \mathfrak{A}$ for all $k \in \omega$;
- (5) Age(\mathfrak{A}) is closed under finite direct products.

Proof. The equivalence of (1) and (2) is shown for relational ω -categorical structures [8], and the same proof also works for ω -categorical structures with functions. The implication from (2) to (3) holds because \mathfrak{A} is an algebra. The implications from (3) to (4) and from (4) to (5) are clear. For the implication from (5) to (1), suppose that $\operatorname{Th}(\mathfrak{A}) \cup S \cup \{x_i \neq y_i\}$ is satisfiable for each $i \in \{1, \ldots, m\}$. Let \mathfrak{A}_i be the substructure of \mathfrak{A} induced by the variables of S and $\{x_i, y_i\}$. Then by assumption $\mathfrak{A}_1 \times \cdots \times \mathfrak{A}_m$ is a substructure of \mathfrak{A} and witnesses that $\operatorname{Th}(\mathfrak{A}) \cup S \cup \{x_1 \neq y_1, \ldots, x_m \neq y_m\}$ is satisfiable. \square

We present a pair of applications of square embeddings in the context of equation solving.

Proposition 3.8. Let \mathfrak{A} be a model-complete ω -categorical structure with finite signature τ such that $\mathfrak{A}^2 \hookrightarrow \mathfrak{A}$. Then there are finitely many $a_1, \ldots, a_n \in A$ such that $\mathrm{CSP}(\mathfrak{A}, \neq)$ is polynomial-time equivalent to $\mathrm{CSP}(\mathfrak{A}, a_1, \ldots, a_n)$.

Proof. We have already mentioned in the introduction that if \mathfrak{B} is a model-complete ω -categorical structure then for all $a_1, \ldots, a_n \in B$ there is a polynomial-time reduction from $CSP(\mathfrak{B}, a_1, \ldots, a_n)$ to $CSP(\mathfrak{B})$; see [7,11]. For the converse reduction, note that for every conjunction of atomic τ -formulas ϕ the formula

$$\phi \wedge s_1 \neq t_1 \wedge \cdots \wedge s_m \neq t_m$$

is satisfiable in \mathfrak{A} if and only if $\phi \wedge s_i \neq t_i$ is satisfiable in \mathfrak{A} for each $i \in \{1, ..., m\}$, because Th(\mathfrak{A}) is convex. By introducing new variables and new identities in ϕ , we may assume that each of the conjuncts $s_i \neq t_i$ is in fact of the form $x_i \neq y_i$ for variables x_i and y_i . To test whether $\phi \wedge x_i \neq y_i$ is satisfiable, we pick representatives $a_1, ..., a_l$ for each orbit of

pairs in Aut(\mathfrak{A}). Note that $\phi \wedge x_i \neq y_i$ is satisfiable if and only if $\phi \wedge x_i = a \wedge y_i = a'$ is satisfiable in \mathfrak{A} for some orbit representatives $a, a' \in \{a_1, \ldots, a_l\}$. Hence, $CSP(\mathfrak{A}, \neq)$ can be reduced to $CSP(\mathfrak{A}, a_1, \ldots, a_l)$. The reduction is in AC_0 , and in particular in Logspace and Ptime.

Proposition 3.9. Let \mathfrak{A} and \mathfrak{B} be ω -categorical algebras with the same signature τ such that $\mathfrak{A}^2 \hookrightarrow \mathfrak{A}$. If $CSP(\mathfrak{A}, \neq)$ is in P then there is a polynomial-time reduction from $CSP(\mathfrak{A} \times \mathfrak{B}, \neq)$ to $CSP(\mathfrak{B}, \neq)$.

Proof. Let ϕ be a conjunction of atomic τ -formula and consider formula

$$\Phi := \phi \wedge x_1 \neq y_1 \wedge \cdots \wedge x_m \neq y_m$$

with variables V, where we again assume without loss of generality that $x_i, y_i \in V$. We claim that Φ is satisfiable in $\mathfrak{A} \times \mathfrak{B}$ if and only if there exists a partition $\{1, 2, \ldots, m\} = I \cup J$ such that $\Phi_I = \phi \wedge \bigwedge_{i \in I} x_i \neq y_i$ is satisfiable in \mathfrak{A} and $\Phi_J = \phi \wedge \bigwedge_{j \in J} x_i \neq y_i$ is satisfiable in \mathfrak{B} .

Let $f: V \to A \times B$ be an assignment satisfying Φ . Let $f_A: V \to A$ be given by $f_A(v) = a$ if and only if f(v) = (a, b) for some $b \in B$; dually define f_B . Hence $f(v) = (f_A(v), f_B(v))$. Then f_A and f_B satisfies ϕ , and if $f(x_i) \neq f(y_i)$ then either $f_A(x_i) \neq f_A(y_i)$ or $f_B(x_i) \neq f_B(y_i)$ (or both). Letting $I = \{k: f_A(x_i) \neq f_A(y_i)\}$ and $J = \{k: f_A(x_i) = f_A(y_i)\}$ and $f_B(x_i) \neq f_B(y_i)\}$ we obtain desired partition of $\{1, 2, ..., m\}$.

Conversely, let V_I and V_J be the variables of Φ_I and Φ_J , respectively. Let $g_A \colon V_I \to A$ and $g_B \colon V_J \to B$ be assignments satisfying Φ_I and Φ_J , respectively. Fix $a \in A$ and $b \in B$. Expand g_A to a map $g'_A \colon V \to A$ by letting $g'_A(x) = a$ for any $x \in V \setminus V_I$; dually obtain g'_B . Then the map $V \to A \times B$ given by $v \mapsto (g'_A(v), g'_B(v))$ is an assignment satisfying Φ . This finishes the proof of our claim.

This gives rise to the following method for determining the satisfiability of Φ . For each $1 \leq i \leq m$ we check (in polynomial-time) if $\Phi_i = \phi \wedge x_i \neq y_i$ is satisfiable in A. Let K be the set of $i \in \{1, \ldots, m\}$ for which Φ_i is not satisfied in A. By the claim and Proposition 3.7 we have that Φ is satisfiable in $\mathfrak{A} \times \mathfrak{B}$ if and only if $\phi \wedge \bigwedge_{k \in K} x_k \neq y_k$ is satisfiable in \mathfrak{B} .

The following lemma shows that the property to have a square embedding implies the existence of a pseudo-Siggers polymorphism in an important situation.

Lemma 3.10. Let \mathfrak{A} be an ω -categorical algebra. If \mathfrak{A}^2 is isomorphic to \mathfrak{A} , then \mathfrak{A} has a pseudo-Siggers polymorphism.

Proof. Clearly there exists an isomorphism $g: \mathfrak{A}^6 \to \mathfrak{A}$. Let $\alpha: A \to A$ be the map defined as follows. For $a \in A$, let $(a_1, \ldots, a_6) \in A^6$ be such that $g(a_1, \ldots, a_6) = a$. Define $\alpha(a) := g(a_2, a_1, a_4, a_3, a_6, a_5)$; then α is an automorphism of \mathfrak{A} , because g is an isomorphism, and for all $x, y, z \in A$ we have that $\alpha(g(x, y, x, z, y, z)) = g(y, x, z, x, z, y)$, so g is a pseudo-Siggers polymorphism.

3.4. Pseudo-Siggers polymorphisms. If \mathfrak{A} is an ω -categorical algebra, then the existence of a pseudo-Siggers polymorphism of (\mathfrak{A}, \neq) has an interesting consequence, which is in fact equivalent if the algebra \mathfrak{A} is even homogeneous.

Lemma 3.11. Let \mathfrak{A} be an ω -categorical algebra. If $s \in \operatorname{Pol}^{(6)}(\mathfrak{A}, \neq)$ is a pseudo-Siggers operation then for all $x, y, z, u, v, w \in A$

$$s(x, y, x, z, y, z) = s(u, v, u, w, v, w)$$

$$\Leftrightarrow s(y, x, z, x, z, y) = s(v, u, w, u, w, v).$$

If \mathfrak{A} is homogeneous, the converse implication holds as well.

Proof. Let s be a pseudo-Siggers operation, i.e., there are $e_1, e_2 \in \text{End}(\mathfrak{A}, \neq)$ such that $e_1(s(x, y, x, z, y, z)) = e_2(s(y, x, z, x, z, y))$. Now observe that

$$s(x, y, x, z, y, z) = s(u, v, u, w, v, w)$$

$$\Leftrightarrow e_1(s(x, y, x, z, y, z)) = e_1(s(u, v, u, w, v, w)) \qquad \text{(since } e_1 \text{ is injective)}$$

$$\Leftrightarrow e_2(s(y, x, z, x, z, y)) = e_2(s(v, u, w, u, w, v)) \qquad \text{(by assumption)}$$

$$\Leftrightarrow s(y, x, z, x, z, y) = s(v, u, w, u, w, v) \qquad \text{(since } e_2 \text{ is injective)}.$$

Conversely, suppose that s satisfies (3.1). By the lift lemma (Lemma 3 in [17]) it suffices to show that for every finite $F \subseteq A$ there exists $\alpha \in \operatorname{Aut}(\mathfrak{A})$ such that $s(x,y,x,z,y,z) = \alpha s(y,x,z,x,z,y)$ for all $x,y,z \in F$. Since \mathfrak{A} is homogeneous, it suffices to verify that for every $k \in \omega$ and $a_1, a_2, a_3 \in F^k$ the k-tuples $s(a_1, a_2, a_1, a_3, a_2, a_3)$ and $s(a_2, a_1, a_3, a_1, a_3, a_2)$ satisfy the same atomic formulas in the language of (\mathfrak{A}, \neq) . So let r, t be terms in the language of \mathfrak{A} such that $r(s(a_1, a_2, a_1, a_3, a_2, a_3)) = t(s(a_1, a_2, a_1, a_3, a_2, a_3))$. Then

$$s(r(a_1), r(a_2), r(a_1), r(a_3), r(a_2), r(a_3)) = r(s(a_1, a_2, a_1, a_3, a_2, a_3))$$

$$= t(s(a_1, a_2, a_1, a_3, a_2, a_3))$$

$$= s(t(a_1), t(a_2), t(a_1), t(a_3), t(a_2), t(a_3))$$

and therefore the assumption implies that

$$s(r(a_2), r(a_1), r(a_3), r(a_1), r(a_3), r(a_2)) = s(t(a_2), t(a_1), t(a_3), t(a_1), t(a_3), t(a_2)),$$

which in turn implies that $r(s(a_2, a_1, a_3, a_1, a_3, a_2)) = t(s(a_2, a_1, a_3, a_1, a_3, a_2))$. Symmetrically, one can show that every atomic formula that holds on $s(a_2, a_1, a_3, a_1, a_3, a_2)$ also holds on $s(a_1, a_2, a_1, a_3, a_2, a_3)$.

We would like to point out that in later sections, whenever we use the assumption that \mathfrak{A} has a pseudo-Siggers polymorphism, we do this by using Property (3.1), and we are not aware of an ω -categorical algebra where the converse of Lemma 3.11 does not hold.

4. Monoids

Let $\mathfrak{M} = (M; \cdot, 1)$ be a monoid. Polymorphisms $f: M^n \to M$ of \mathfrak{M} have the particularly pleasing property that they decompose in the following sense: for any $x_1, \ldots, x_n \in M$ we have

$$(4.1) f(x_1, x_2, \dots, x_n) = f(x_1, 1, \dots, 1) \cdot f(1, x_2, 1, \dots, 1) \cdots f(1, 1, \dots, 1, x_n).$$

For $I \subseteq \{1, ..., n\}$ we write $x_I^{(n)}$ for the *n*-tuple whose *i*-th component is x if $i \in I$ and 1 otherwise.

Definition 4.1. Let $f \in \operatorname{Pol}^{(n)}(\mathfrak{M})$. Let $f_I \colon M \to M$ be the operation given by

$$f_I(x) := f(x_I^{(n)}).$$

Remark 4.2. The unary constant operation $x \mapsto 1$ is an endomorphism of every monoid \mathfrak{M} . It follows that for every $I \subseteq \{1, \ldots, n\}$ and every $f \in \operatorname{Pol}^{(n)}(\mathfrak{M})$ the operation f_I is an endomorphism of \mathfrak{M} . Every polymorphism of (\mathfrak{M}, \neq) must preserve $M \setminus \{1\}$.

Note that f_I is not necessarily a self-embedding of $(M; \cdot, 1)$. For example, the projection f(x, y) = x is a polymorphism of $(M; \cdot, 1)$, and $f_{\{2\}}$ is constant. However, there must be a subset I of $\{1, \ldots, n\}$ such that f_I is an embedding.

Proposition 4.3. Let $(M; \cdot, 1, \neq)$ be a monoid and $f: M^n \to M$ a polymorphism of $(M; \cdot, 1)$. Then for any partition $I_1 \cup I_2 \cup \cdots \cup I_k$ of $\{1, \ldots, n\}$ there exists $j \in \{1, \ldots, k\}$ such that f_{I_j} is a self-embedding of $(M; \cdot, 1)$.

Proof. Let $I_1 \cup I_2 \cup \cdots \cup I_k$ be a partition of $\{1, \ldots, n\}$. Suppose for contradiction that for every $j \in \{1, \ldots, k\}$ there exist distinct $x_j, y_j \in M$ such that $f_{I_j}(x_j) = f_{I_j}(y_j)$. Let \bar{x} be the *n*-tuple such that $\bar{x}_i = x_j$ if $i \in I_j$, and similarly let \bar{y} be the *n*-tuple such that $\bar{y}_i = y_j$ if $i \in I_j$. Then

$$f(\bar{x}) = f((x_1)_{I_1}^{(n)} \cdots (x_k)_{I_k}^{(n)}) = f((x_1)_{I_1}^{(n)}) \cdots f((x_k)_{I_k}^{(n)})$$

$$= f_{I_1}(x_1) \cdots f_{I_k}(x_k)$$

$$= f_{I_1}(y_1) \cdots f_{I_k}(y_k) = f((y_1)_{I_1}^{(n)} \cdots (y_k)_{I_k}^{(n)}) = f(\bar{y})$$

showing that f does not preserve \neq , a contradiction.

Proposition 4.4. A monoid $(M; \cdot, 1, \neq)$ has a pseudo-Siggers polymorphism if and only if there are a polymorphism $s \colon M^6 \to M$ and self-embeddings a, b of $(M; \cdot, 1)$ such that

$$a(s_{\{i,j\}}(x)) = b(s_{\{i,j\}}(x))$$

for all $x \in M$ and $\{i, j\} \in \{\{1, 3\}, \{2, 5\}, \{4, 6\}\}.$

Proof. For any $x, y, z \in M$ we have

$$a(s(x,y,x,z,y,z)) = a(s_{\{1,3\}}(x)s_{\{2,5\}}(y)s_{\{4,6\}}(z)) = a(s_{\{1,3\}}(x))a(s_{\{2,5\}}(y))a(s_{\{4,6\}}(z))$$
 to which the result follows. \Box

5. Groups

Let \mathfrak{G} be an ω -categorical group. There are homogeneous ω -categorical groups such that $CSP(\mathfrak{G}, \neq)$ is undecidable (Section 5.2). However, we are able to classify the complexity of $CSP(\mathfrak{G}, \neq)$ if \mathfrak{G} is additionally abelian; in this case, $CSP(\mathfrak{G}, \neq)$ is in P or NP-complete (Section 5.5). Some of the structural results we obtain not only hold for abelian groups, but for general groups with a pseudo-Siggers polymorphism, and they will be presented in Section 5.1.

The order of an element $g \in G$ is the cardinality of the subgroup of \mathfrak{G} generated by g. We say that \mathfrak{G} is a torsion group if every element of \mathfrak{G} is of finite order. Since an

 ω -categorical group is uniformly locally finite, it must be of *finite exponent* [38], i.e., there exists $n \in \omega$ such that $g^n = 1$ for every $g \in G$; the minimum such n is called the *exponent* of \mathfrak{G} . This follows from the well-known fact that ω -categorical structures \mathfrak{B} are uniformly locally finite (Corollary 7.3.2 in [27]), that is, there exists a function $f: \omega \to \omega$ such that for every $n \in \omega$ each substructure of \mathfrak{B} generated by n elements has at most f(n) elements. In particular, every ω -categorical group must be a torsion group.

Remark 5.1. Note that the identity element 1 of \mathfrak{G} has the quantifier-free definition $x \cdot x = x$ (in the language $\{\cdot\}$ of semigroups), and we may therefore assume that there is a constant symbol for 1 in the signature. Similarly, the inverse function has a quantifier-free definition, and we assume that the signature contains a unary function symbol for taking inverses.

5.1. **Pseudo-Siggers Groups.** An *involution* is an element of \mathfrak{G} of order 2. An element $x \in G$ is *central* if xg = gx for all $g \in G$. Clearly, an involution generates a normal subgroup if and only if it is central. Since every group is in particular a monoid, we may use Proposition 4.3 and obtain the following.

Proposition 5.2. Let \mathfrak{G} be an ω -categorical group such that (\mathfrak{G}, \neq) has a pseudo-Siggers polymorphism f. Then

- $\mathfrak{G}^2 \hookrightarrow \mathfrak{G}$, or
- There is a central involution $i \in G$ such that $\mathfrak{G} \times \mathfrak{G}/\langle i \rangle \hookrightarrow \mathfrak{G}$.

Proof. By Proposition 4.3 we may assume without loss of generality that $f_{\{1,3\}}$ is an embedding. Let g be the binary polymorphism of (\mathfrak{G}, \neq) given by

$$g(x,y) := f_{\{1,3\}}(x) \cdot f_{\{2,4,5,6\}}(y) = f(x,1,x,1,1,1) \cdot f(1,y,1,y,y,y) = f(x,y,x,y,y,y).$$

Claim. Im(g) is bi-embeddable with $\mathfrak{G} \times \text{Im}(f_{\{2,4,5,6\}})$. Suppose that there exist $x,y \in G$ such that g(x,1) = g(1,y). Then x = 1 or y = 1 since g preserves \neq . So $\text{Im}(f_{\{1,3\}}) \cap \text{Im}(f_{\{2,4,5,6\}}) = \{1\}$. Moreover, the subgroups $\text{Im}(f_{\{1,3\}})$ and $\text{Im}(f_{\{2,4,5,6\}})$ of \mathfrak{G} are commuting, since for all $x,y \in G$

$$f_{\{1,3\}}(x)f_{\{2,4,5,6\}}(y) = f(x,y,x,y,y,y) = f_{\{2,4,5,6\}}(y)f_{\{1,3\}}(x).$$

It follows that Im(g) is generated by $\text{Im}(f_{\{1,3\}}) \cup \text{Im}(f_{\{2,4,5,6\}})$, and bi-embeddable with $\mathfrak{G} \times \text{Im}(f_{\{2,4,5,6\}})$, as required.

If $f_{\{2,4,5,6\}}$ is an embedding, then $g \colon \mathfrak{G}^2 \hookrightarrow \mathfrak{G}$ and we are done, so suppose that this is not the case. Then there exists an $i \in G \setminus \{1\}$ such that $f_{\{2,4,5,6\}}(i) = 1$. So we have f(1,i,1,i,i) = f(1,1,1,1,1,1) and Equation 3.1 implies that f(i,1,i,1,i,i) = f(1,1,1,1,1,1). Hence, $f(i,i,i,i,i^2,i^2) = 1$, and we must have $i^2 = 1$ because otherwise f would not preserve $G \setminus \{1\}$. Note also that if $x \in G$ is such that f(1,x,1,x,x,x,x) = 1 then x = i or x = 1: otherwise, f(1,x,1,x,x,x,x) = 1 = f(i,1,i,1,i,i) in contradiction to the assumption that f preserves \neq . So the kernel of $f_{\{2,4,5,6\}}$ is $\{1,i\} = \langle i \rangle$ and $(x,y\langle i \rangle) \mapsto g(x,y)$ is an embedding $\mathfrak{G} \times \mathfrak{G}/\langle i \rangle \hookrightarrow \mathfrak{G}$.

Let \mathfrak{H} and \mathfrak{K} be ω -categorical groups. We say that \mathfrak{H} and \mathfrak{K} are of relatively prime exponent if the exponents of \mathfrak{H} and \mathfrak{K} are co-prime.

Lemma 5.3. Let \mathfrak{H} and \mathfrak{K} be ω -categorical groups of relatively prime exponent. If the structure $(\mathfrak{H} \times \mathfrak{K}, \neq)$ has a pseudo-Siggers polymorphism, then (\mathfrak{H}, \neq) and (\mathfrak{K}, \neq) have pseudo-Siggers polymorphisms, too.

Proof. Let $h \in \omega$ be the exponent of \mathfrak{H} and $k \in \omega$ the exponent of \mathfrak{K} , and let $\mathfrak{G} := \mathfrak{H} \times \mathfrak{K}$. Suppose that (\mathfrak{G}, \neq) has a pseudo-Siggers polymorphism s. Then \mathfrak{H} is isomorphic to the subgroup $\mathfrak{H} \times \{1^{\mathfrak{K}}\}$ of \mathfrak{G} , which is precisely the set of all elements of G that satisfy $x^h = 1$; hence, $H \times \{1^{\mathfrak{K}}\}$ is primitive positive definable in \mathfrak{G} , and the restriction of s to this set is a pseudo-Siggers polymorphism of $(\mathfrak{H} \times \{1^{\mathfrak{K}}\}, \neq)$. Therefore, (\mathfrak{H}, \neq) has a pseudo-Siggers polymorphism. \square

- 5.2. Undecidable ω -categorical groups. Saracino and Wood [39] showed that there are 2^{ω} non-isomorphic homogeneous ω -categorical groups. Homogeneous ω -categorical structures have quantifier elimination [28] and in particular they are model-complete. Hence, if two homogeneous ω -categorical structures are companions, then they must be isomorphic [28]. Recall from Lemma 3.1 that ω -categorical algebras $\mathfrak A$ and $\mathfrak B$ are companions if and only if $\mathrm{CSP}(\mathfrak A,\neq)$ and $\mathrm{CSP}(\mathfrak B,\neq)$ are the same computational problem. Since there are only countably many Turing machines, it follows that there are ω -categorical homogeneous groups $\mathfrak A$ such that $\mathrm{CSP}(\mathfrak A,\neq)$ is undecidable.
- 5.3. The abelian case. We now consider ω -categorical abelian groups \mathfrak{G} . Our main results are a characterisation of the case that (\mathfrak{G}, \neq) has a pseudo-Siggers polymorphism and a full complexity classification for $\mathrm{CSP}(\mathfrak{G}, \neq)$. It is standard to then use additive notation; so the identity element is from now on denoted by 0 and the group composition by +.

Recall that a p-group is a group whose elements have orders that are powers of a fixed prime p. For example, the cyclic group \mathbb{Z}_{p^n} of order p^n is a p-group. A subgroup of \mathfrak{G} which is a p-group is also called a p-subgroup. We write $\bigoplus_{i\in I}G_i$ for the direct sum of the G_i , i.e., for the subgroup of $\prod_{i\in I}G_i$ containing all elements that are 1 at all but finitely many indices. If $G_i = G$ for all $i \in I$ then we also write $G^{(I)}$ instead of $\bigoplus_{i\in I}G$. Finite direct products $\mathfrak{G}_1 \times \cdots \times \mathfrak{G}_k$ coincide with finite direct sums $\mathfrak{G}_1 \oplus \cdots \oplus \mathfrak{G}_k$ and we use the latter notation in this section. As mentioned earlier, an ω -categorical group \mathfrak{G} must be a torsion group.

Theorem 5.4 (Theorem 1 in [29]). Every abelian torsion group \mathfrak{G} is the direct sum of its p-subgroups.

Recall that every ω -categorical group must be of finite exponent.

Corollary 5.5. Every abelian group \mathfrak{G} of finite exponent is a finite direct sum of its p-subgroups.

As a consequence of Corollary 5.5 and Lemma 5.3 we need to consider abelian p-groups. We first recall another basic fact from the theory of abelian groups.

Theorem 5.6 (Theorem 6 in [29]). Every abelian group of finite exponent is a direct sum of cyclic groups.

A group is called *trivial* if it only consists of the identity element 1, and *non-trivial* otherwise.

Corollary 5.7. Let \mathfrak{G} be a non-trivial countable ω -categorical p-group. Then there exists $k \in \omega$ such that

(5.1)
$$\mathfrak{G} = \mathbb{Z}_{p^{n_1}}^{(s_1)} \oplus \cdots \oplus \mathbb{Z}_{p^{n_k}}^{(s_k)}$$

where $n_1, \ldots, n_k \in \{1, 2, 3, \ldots\}$ and $s_1, \ldots, s_k \in \{1, 2, \ldots\} \cup \{\omega\}$.

Proof. Theorem 5.6 shows that $\mathfrak{G} = \bigoplus_{i \in \omega} \mathbb{Z}_{c_i}$ for some integer sequence $(c_i)_{i \in \omega}$ with $c_i \geq 2$ for all $i \in \omega$. Theorem 5.4 implies that each c_i is of the form $c_i = p^{n_i}$ for some $n_i \in \{1, 2, 3, \ldots\}$. So we may write $\mathfrak{G} = \bigoplus_{i \in \omega} \mathbb{Z}_{p^{n_i}}^{(s_i)}$ where $s_i \in \omega \cup \{\omega\}$. As \mathfrak{G} has finite exponent we have $s_i = 0$ for all but finitely many $i \in \omega$.

Understanding p-groups up to bi-embeddability will be useful. For groups of the form (5.1) we define

$$m_{\mathfrak{G}} := \begin{cases} 0 & \text{if } s_i \in \omega \text{ for every } i \in \{1, \dots, k\} \\ \max\{n : s_n = \omega\} & \text{otherwise.} \end{cases}$$

Note that \mathfrak{G} is finite if and only if $m_{\mathfrak{G}} = 0$. The following is a consequence of a more general result about bi-embeddability of abelian p-groups [6] (Corollary 5.4); see Remark 4.12 in [23].

Lemma 5.8. Let $k, \ell \in \omega$ and $s_n, t_n \in \omega \cup \{\omega\}$. Then $\mathfrak{G} = \bigoplus_{n \in \{1, \dots, k\}} \mathbb{Z}_{p^n}^{(s_n)}$ and $\mathfrak{H} = \bigoplus_{n \in \{1, \dots, \ell\}} \mathbb{Z}_{p^n}^{(t_n)}$ are bi-embeddable if and only if $m_{\mathfrak{G}} = m_{\mathfrak{H}}$ and $s_n = t_n$ for all $n \geq m_{\mathfrak{G}}$.

We apply this lemma to the two possibilities that arise in Proposition 5.2 and start with the easier situation where $\mathfrak{G}^2 \hookrightarrow \mathfrak{G}$.

Lemma 5.9. Let \mathfrak{G} be an ω -categorical abelian p-group. Then $\mathfrak{G}^2 \hookrightarrow \mathfrak{G}$ if and only if \mathfrak{G} is bi-embeddable with $\mathbb{Z}_{p^n}^{(\omega)}$ for some $n \in \omega$.

Proof. The statement is trivial if \mathfrak{G} is trivial. Otherwise, by Corollary 5.7, \mathfrak{G} can be written as $\mathfrak{G} = \bigoplus_{n \in \{1,\dots,k\}} \mathbb{Z}_{p^n}^{(s_n)}$. Then $s_n = 2s_n$ for every $n > m_{\mathfrak{G}}$ by Lemma 5.8. Hence, $s_n = 0$ for every $n \geq m_{\mathfrak{G}}$, and again by Lemma 6.14 we conclude that \mathfrak{G} is bi-embeddable with $\mathbb{Z}_{p^m_{\mathfrak{G}}}^{(\omega)}$. The converse is immediate.

We now treat the other possibility that arises in Proposition 5.2 and which involves the quotient $\mathfrak{G}/\langle x \rangle$ for a central involution x of \mathfrak{G} . We first have to recall how this quotient looks like if \mathfrak{G} is a countable abelian ω -categorical 2-group and hence of the form as described in Corollary 5.7.

Lemma 5.10. Let $\mathfrak{G} = \mathbb{Z}_{2^1}^{(s_1)} \oplus \cdots \oplus \mathbb{Z}_{2^k}^{(s_k)}$ be an abelian ω -categorical 2-group and let $x \in G$ be a central involution. Then $G/\langle x \rangle$ is isomorphic to

$$\mathbb{Z}_{2^{1}}^{(s_{1})} \oplus \cdots \oplus \mathbb{Z}_{2^{i-2}}^{(s_{i-2})} \oplus \mathbb{Z}_{2^{i-1}}^{(s_{i-1}+1)} \oplus \mathbb{Z}_{2^{i}}^{(s_{i}-1)} \oplus \mathbb{Z}_{2^{i+1}}^{(s_{i+1})} \oplus \cdots \oplus \mathbb{Z}_{2^{k}}^{(s_{k})}$$

for some $i \in \{1, \ldots, k\}$ (where $s_0 = 0$ if i = 1).

Proof. Let $g_1, g_2, \dots \in G$ be such that $\mathfrak{G} = \bigoplus_{i \in \omega} \langle g_i \rangle$.

Claim. x is a power of some generator g_i . Suppose without loss of generality that $x = (m_1g_1, \ldots, m_rg_r, 0, \ldots)$ for $r \in \omega$ and $m_1, \ldots, m_r \in \omega$ powers of two. Let $t \in \{1, \ldots, r\}$ be such that m_t is minimal among m_1, \ldots, m_r , and let $a_i := m_i/m_t$ for $i \in \{1, \ldots, r\}$. Let $y = (a_1g_1, \ldots, a_rg_r, 0, 0, \ldots)$ be so that $x = y^{m_t}$. Since a_1, \ldots, a_r have greatest common divisor 1, we may use the following lemma (Lemma II.3.b in [41]): if g_1, \ldots, g_r are generators of an abelian group \mathfrak{H} , and if a_1, \ldots, a_r are integers with greatest common divisor 1, then the element $a_1g_1 + a_2g_2 + \cdots + a_rg_r$ is one of a set of r generators of \mathfrak{H} . So there are $h_2, \ldots, h_r \in G$ such that \mathfrak{G} can be written as

$$\mathfrak{G} = \langle y \rangle \oplus \bigoplus_{i \in \{2, \dots, r\}} \langle h_i \rangle \oplus \bigoplus_{i \in \{r, r+1, \dots\}} \langle g_i \rangle$$

Since $\mathbb{Z}_{2^s}/\mathbb{Z}_2$ is isomorphic to $\mathbb{Z}_{2^{s-1}}$, we get that $\mathbb{Z}_{2^1}^{(s_1)} \oplus \cdots \oplus \mathbb{Z}_{2^k}^{(s_k)}/\langle x \rangle$ is of the form as described in the statement.

Lemma 5.11. A non-trivial ω -categorical abelian 2-group \mathfrak{G} is bi-embeddable with $\mathfrak{G} \oplus \mathfrak{G}/\langle x \rangle$ for some involution $x \in G$ if and only if

- $\mathfrak{G}^2 \hookrightarrow \mathfrak{G}$, or
- \mathfrak{G} is bi-embeddable with $\mathbb{Z}_{2^n}^{(\omega)} \oplus \mathbb{Z}_{2^{n+1}}$ for some $n \in \omega$.

Proof. By Corollary 5.7, the 2-group \mathfrak{G} can be written as $\bigoplus_{n\in\{1,\ldots,k\}} \mathbb{Z}_{2^n}^{(s_n)}$ for $s_k > 0$. First suppose that \mathfrak{G} is bi-embeddable with $\mathfrak{G} \oplus \mathfrak{G}/\langle x \rangle$ for some involution $x \in G$. If $\mathfrak{G}^2 \hookrightarrow \mathfrak{G}$ then we are done. Otherwise, by Lemma 5.9 the group \mathfrak{G} is not bi-embeddable with $\mathbb{Z}_{p^n}^{(\omega)}$ for some $n \in \omega$. Corollary 5.7 then implies that $s_k < \omega$. By Lemma 5.10 there exists a unique $i \in \{1,\ldots,k\}$ such that $G/\langle x \rangle$ is isomorphic to

$$\mathbb{Z}_{2}^{(s_{1})} \oplus \cdots \oplus \mathbb{Z}_{2^{i-2}}^{(s_{i-2})} \oplus \mathbb{Z}_{2^{i-1}}^{(s_{i-1}+1)} \oplus \mathbb{Z}_{2^{i}}^{(s_{i}-1)} \oplus \mathbb{Z}_{2^{i+1}}^{(s_{i+1})} \oplus \cdots \oplus \mathbb{Z}_{2^{k}}^{(s_{k})}.$$

Since \mathfrak{G} is bi-embeddable with $\mathfrak{G} \oplus \mathfrak{G}/\langle x \rangle$ Lemma 5.8 implies that either $s_k = s_k(s_k - 1)$ if k = i or $s_k = 2s_k$ otherwise. Since s_k is finite and non-zero (as \mathfrak{G} contains an involution, it must be non-trivial) we conclude that k = i and $s_k = 1$. If s_{k-1} is finite then $s_{k-1} = 2s_{k-1} + 1$ by Lemma 5.8, a contradiction. So $s_{k-1} = \omega$, and by Lemma 5.8 \mathfrak{G} is bi-embeddable with $\mathbb{Z}_{2^{k-1}}^{(\omega)} \oplus \mathbb{Z}_{2^k}$, as required.

Conversely, if $\mathfrak{G}^2 \hookrightarrow \mathfrak{G}$ then by Lemma 5.9 \mathfrak{G} is bi-embeddable with $\mathbb{Z}_{2^n}^{(\omega)}$ for some $n \in \omega$, and in fact $n \geq 1$ since \mathfrak{G} is non-trivial. Hence, there exists an involution $x \in G$ and $\mathfrak{G}/\langle x \rangle$ is biembeddable with $\mathbb{Z}_{2^{n-1}} \oplus \mathbb{Z}_{2^n}^{(\omega)}$ by Lemma 5.10, which is bi-embeddable with $\mathbb{Z}_{2^n}^{(\omega)}$ by Lemma 5.8. We conclude that \mathfrak{G} is bi-embeddable with $\mathfrak{G} \oplus \mathfrak{G}/\langle x \rangle$. If \mathfrak{G} is bi-embeddable with $\mathbb{Z}_{2^n}^{(\omega)} \oplus \mathbb{Z}_{2^{n+1}}$ for some $n \in \omega$, let x be an involution generated by an element of order 2^{n+1} . Then $\mathfrak{G}/\langle x \rangle$ is bi-embeddable with $\mathbb{Z}_{2^n}^{(\omega)} \oplus \mathbb{Z}_{2^n}$, which is isomorphic to $\mathbb{Z}_{2^n}^{(\omega)}$ and hence $\mathfrak{G} \oplus \mathfrak{G}/\langle x \rangle$ is bi-embeddable with $\mathbb{Z}_{2^n}^{(\omega)} \oplus \mathbb{Z}_{2^{n+1}} \oplus \mathbb{Z}_{2^n}^{(\omega)}$ and hence with \mathfrak{G} .

Proposition 5.12. Let \mathfrak{G} be an ω -categorical abelian group such that (\mathfrak{G}, \neq) has a pseudo-Siggers polymorphism. Then \mathfrak{G} is bi-embeddable with $\mathbb{Z}_m^{(\omega)}$ or with $\mathbb{Z}_m^{(\omega)} \oplus \mathbb{Z}_{2m}$ for some $m \geq 1$. *Proof.* If \mathfrak{G} is trivial then it is bi-embeddable with $\mathbb{Z}_1^{(\omega)}$ and we are done. Otherwise, by Corollary 5.5 the group \mathfrak{G} is $\mathfrak{G}_{p_1} \oplus \cdots \oplus \mathfrak{G}_{p_r}$, where $r \geq 1, p_1, \ldots, p_r$ are primes, and \mathfrak{G}_{p_i} for $i \in \{1, \ldots, r\}$ is a non-trivial p_i -subgroup of \mathfrak{G} . By Lemma 5.3, for each $p \in \{p_1, \ldots, p_r\}$ the structure (\mathfrak{G}_p, \neq) has a pseudo-Siggers polymorphism.

By Proposition 5.2 we have $\mathfrak{G}_p^2 \hookrightarrow \mathfrak{G}_p$ or \mathfrak{G}_p is bi-embeddable with $\mathfrak{G}_p \oplus \mathfrak{G}_p / \langle x \rangle$ for some involution $x \in G$ and p = 2. In the first case, \mathfrak{G}_p is bi-embeddable with $\mathbb{Z}_{p^n}^{(\omega)}$ for some $n \in \omega$ by Lemma 5.9. In the latter case, \mathfrak{G}_p is bi-embeddable with $\mathbb{Z}_{2^n}^{(\omega)} \oplus \mathbb{Z}_{2^{n+1}}$ or with $\mathbb{Z}_{2^n}^{(\omega)}$ for some $n \in \omega$ by Lemma 5.11. So we deduce that there are $n_1, \ldots, n_r \in \{1, 2, 3, \ldots\}$ such that \mathfrak{G} is bi-embeddable with

- $\mathbb{Z}_{p_1^{n_1}}^{(\omega)} \oplus \cdots \oplus \mathbb{Z}_{p_r^{n_r}}^{(\omega)}$ or with
- $\mathbb{Z}_{p_1^{n_1}}^{(\omega)} \oplus \mathbb{Z}_{p_1^{n_1+1}} \oplus \mathbb{Z}_{p_2^{n_2}}^{(\omega)} \oplus \cdots \oplus \mathbb{Z}_{p_r^{n_r}}^{(\omega)}$ where $p_1 = 2$ and $p_i > 2$ for every $i \in \{2, \ldots, r\}$.

Let $m := p_1^{n_1} p_2^{n_2} \cdots p_r^{n_r}$. In case (1), the group \mathfrak{G} is isomorphic to $\mathbb{Z}_m^{(\omega)}$. In case (2), we have

$$\mathbb{Z}_{p_1^{n_1}}^{(\omega)} \oplus \mathbb{Z}_{p_1^{n_1+1}} \oplus \mathbb{Z}_{p_2^{n_2}}^{(\omega)} \oplus \cdots \oplus \mathbb{Z}_{p_r^{n_r}}^{(\omega)} \simeq \mathbb{Z}_{p_1^{n_1}}^{(\omega)} \oplus \cdots \oplus \mathbb{Z}_{p_r^{n_r}}^{(\omega)} \oplus \mathbb{Z}_{2^{n_1+1}} \oplus \mathbb{Z}_{p_2^{n_2}} \oplus \cdots \oplus \mathbb{Z}_{p_r^{n_r}}^{(\omega)} \oplus \mathbb{Z}_{2m}.$$

5.4. Polynomial-time tractable abelian groups. Let $n \in \omega$. In this section we present a polynomial-time algorithm for $\mathrm{CSP}(\mathbb{Z}_{2n} \oplus \mathbb{Z}_n^{(\omega)}, \neq)$. Continuing with the additive notation, we let $\mathbb{Z}_k = \{0, 1, \ldots, k-1\}$ for each $k \in \omega$. So let Φ be a conjunction of atomic formulas of the form x = y + z and of the form $x \neq y$ over a finite set of variables V. We would like to test whether Φ is satisfiable in $\mathbb{Z}_{2n} \oplus \mathbb{Z}_n^{(\omega)}$ for some fixed $n \geq 2$.

Remark 5.13. Note that over \mathbb{Z}_2 every disequality $x \neq y$ can be translated into an equality x = y + 1 and hence satisfiability of the entire system can be solved in polynomial time with Gaussian elimination. The same trick does not work for solvability in \mathbb{Z}_n if $n \geq 3$, and indeed satisfiability of disequalities over \mathbb{Z}_n is NP-complete for $n \geq 3$.

Linear equation systems over \mathbb{Z}_k , for any $k \in \omega$, can be solved in polynomial time [26]. We need this algorithm for equation systems over \mathbb{Z}_{2n} . Alternatively, we can use a more general algorithm of Bulatov and Dalmau for constraints preserved by a Maltsev operation [21]. Let $m: \mathbb{Z}_{2n}^3 \to \mathbb{Z}_{2n}$ be the Maltsev operation given by $(x, y, z) \mapsto x - y + z$. Observe that m is idempotent, preserves the graph of addition, and also preserves the following relation

$$R := \{(x_1, x_2) \in \mathbb{Z}_{2n}^2 \mid x_1 - x_2 = n\}.$$

To see this, let $(x_1, x_2), (y_1, y_2), (z_1, z_2) \in R$. Then

$$m(x_1, y_1, z_1) - m(x_2, y_2, z_2) = x_1 - x_2 - (y_1 - y_2) + z_1 - z_2 = n - n + n = n.$$

The algorithm. Let Φ_e be all the conjuncts in Φ that are equations, and let Φ_d be all the conjuncts in Φ that are disequalities. Our algorithm is the following.

- (1) Test for each disequality $x \neq y$ in Φ_d with Gaussian elimination whether Φ_e implies x = y in \mathbb{Z}_n . Let Φ_d^* be the set of all disequalities where this is the case.
- (2) For each inequality $x \neq y$ in Φ_d^* , add the constraint R(x,y) to Φ_e , and solve the resulting instance of the CSP over \mathbb{Z}_{2n} with the Bulatov-Dalmau algorithm for Maltsev constraints. The algorithm accepts if and only if the Maltsev instance is satisfiable.

Theorem 5.14. Let $n \geq 1$. Then the algorithm presented above solves $CSP(\mathbb{Z}_{2n} \oplus \mathbb{Z}_n^{(\omega)}, \neq)$ in polynomial time.

Proof. It is clear that the algorithm has a polynomial running time. To prove the correctness of this algorithm, first suppose that Φ has a solution $s: V \to \mathbb{Z}_{2n} \oplus \mathbb{Z}_n^{(\omega)}$. Let $x \neq y$ be a disequality from Φ_d^* , and let $s(x) = (s_1(x), s_2(x), \dots)$ and $s(y) = (s_1(y), s_2(y), \dots)$. By the definition of Φ_d^* , we must have that $s(x)_i = s(y)_i$ for $i \in \{1, 2, \dots\}$ and $s(x)_1 = s(y)_1 \pmod{n}$. Since $s(x) \neq s(y)$ we must have $s(x)_1 \neq s(y)_1$, and hence $s(x)_1 - s(y)_1 = n$ and $(s(x)_1, s(y)_1) \in R$. Therefore, if the Bulatov-Dalmau algorithm rejects, then our algorithm correctly rejects the input.

Conversely, suppose that the algorithm accepts. Hence, the input to the Maltsev constraints has a solution $r: V \to \mathbb{Z}_{2n}$. Let ϕ_1, \ldots, ϕ_k be the disequalities in $\Phi_d \setminus \Phi_d^*$. For each $i \leq k$ there exists a solution t_i to Φ_e over \mathbb{Z}_n such that t_i satisfies ϕ_i . We then construct a solution $s: V \to \mathbb{Z}_{2n} \oplus \mathbb{Z}_n^k$ of Φ as follows:

$$s(x) := (r(x), t_1(x), \dots, t_k(x))$$

which may naturally be viewed as a solution $s: V \to \mathbb{Z}_{2n} \oplus \mathbb{Z}_n^{(\omega)}$. The map s satisfies Φ_e since each of r, t_1, \ldots, t_k does. Moreover, s satisfies Φ_d^* since r does. Finally, s satisfies ϕ_i for $i \in \{1, \ldots, k\}$ since t_i does.

5.5. The classification. We combine the results obtained in the previous sections to prove our complexity dichotomy for ω -categorical abelian groups (Theorem 5.16). The border is given by the existence of a pseudo-Siggers polymorphism of the model companion of (\mathfrak{G}, \neq) . Then we strengthen the statement by providing an exact characterisation of those ω -categorical abelian groups \mathfrak{G} such that (\mathfrak{G}, \neq) has a pseudo-Siggers polymorphism (Theorem 5.17). We finally prove that for abelian groups the two cases in Theorem 2.5 are disjoint, which provides yet another equivalent characterisation of the complexity border in terms of uniformly continuous minor-preserving maps to \mathscr{P} .

Proposition 5.15. If \mathfrak{G} is an ω -categorical abelian group then $CSP(\mathfrak{G}, \neq)$ is in NP.

Proof. By Theorem 5.4 the group \mathfrak{G} can be written as $\mathfrak{G}_{p_1} \oplus \cdots \oplus \mathfrak{G}_{p_r}$, where $r \in \omega$ and p_1, \ldots, p_r are distinct primes, and \mathfrak{G}_{p_i} for $i \in \{1, \ldots, r\}$ is a non-trivial p_i -subgroup of \mathfrak{G} . By Lemma 5.8 each \mathfrak{G}_{p_i} is bi-embeddable with $\mathbb{Z}_{p_i^{n_i}}^{(\omega)} \oplus \mathfrak{H}_i$ for some $n_i \in \omega \cup \{0\}$ and finite abelian group \mathfrak{H}_i . Hence \mathfrak{G} is bi-embeddable with $\mathbb{Z}_n^{(\omega)} \oplus \mathfrak{H}_i$, where $n = p_1^{n_1} p_2^{n_2} \cdots p_r^{n_r}$

and $\mathfrak{H} = \mathfrak{H}_1 \oplus \mathfrak{H}_2 \oplus \cdots \oplus \mathfrak{H}_r$. Since $\mathrm{CSP}(\mathbb{Z}_n^{(\omega)}, \neq)$ is in P and $(\mathbb{Z}_n^{(\omega)})^2 \hookrightarrow \mathbb{Z}_n^{(\omega)}$, it follows from Proposition 3.9 that there is a polynomial-time reduction from $\mathrm{CSP}(\mathbb{Z}_n^{(\omega)} \oplus \mathfrak{H}, \neq)$ to $\mathrm{CSP}(\mathfrak{H}, \neq)$, and hence from $\mathrm{CSP}(\mathfrak{G}, \neq)$ to $\mathrm{CSP}(\mathfrak{H}, \neq)$. Since \mathfrak{H} is finite, the result follows.

Theorem 5.16. Let \mathfrak{G} be an ω -categorical abelian group and let \mathfrak{H} be its model companion. If (\mathfrak{H}, \neq) has a pseudo-Siggers polymorphism, then $CSP(\mathfrak{G}, \neq)$ is in P. Otherwise, $CSP(\mathfrak{G}, \neq)$ is NP-complete.

Proof. If (\mathfrak{H}, \neq) does not have a pseudo-Siggers polymorphism, then $\mathrm{CSP}(\mathfrak{H}, \neq)$ and therefore also $\mathrm{CSP}(\mathfrak{G}, \neq)$ are NP-hard by Theorem 2.5, and thus NP-complete by Proposition 5.15. Otherwise, Proposition 5.12 implies that \mathfrak{H} is bi-embeddable with $\mathbb{Z}_n^{(\omega)}$ or with $\mathbb{Z}_n^{(\omega)} \oplus \mathbb{Z}_{2n}$ for some $n \in \omega$. In this case the polynomial-time tractability of $\mathrm{CSP}(\mathfrak{H}, \neq)$ and therefore also of $\mathrm{CSP}(\mathfrak{G}, \neq)$ follows from Theorem 5.14.

The border between polynomial-time tractable and NP-hard cases can be described mathematically in several equivalent ways.

Theorem 5.17. Let \mathfrak{G} be an ω -categorical abelian group. Then the following are equivalent.

- (1) (\mathfrak{G}, \neq) has a pseudo-Siggers polymorphism;
- (2) \mathfrak{G} is bi-embeddable with either $\mathbb{Z}_n^{(\omega)}$ or $\mathbb{Z}_n^{(\omega)} \oplus \mathbb{Z}_{2n}$ for some $n \geq 1$;
- (3) the model-complete core of (\mathfrak{G}, \neq) has a pseudo-Siggers polymorphism.

Proof. The implication from (1) to (2) is Proposition 5.12.

To prove the implication from (2) to (3) it suffices to prove that $\mathbb{Z}_n^{(\omega)}$ and $\mathbb{Z}_n^{(\omega)} \oplus \mathbb{Z}_{2n}$ are model-complete and have a pseudo-Siggers polymorphism. Since $(\mathbb{Z}_n^{(\omega)})^2$ is isomorphic to $\mathbb{Z}_n^{(\omega)}$, the structure $(\mathbb{Z}_n^{(\omega)}, \neq)$ has a pseudo-Siggers polymorphism by Lemma 3.10. Moreover, it is well-known and easy to see that $\mathbb{Z}_n^{(\omega)}$ is homogeneous, and therefore model-complete.

Now let $\mathfrak{G} = \mathbb{Z}_n^{(\omega)} \oplus \mathbb{Z}_{2n}$. Let $(a_i)_{i \in \omega}$ be a sequence of elements of G of order n and let b be an element of G of order 2n such that $\mathfrak{G} = \bigoplus_{i \in \omega} \langle a_i \rangle \oplus \langle b \rangle$. We construct a pseudo-Siggers polymorphism s of \mathfrak{G} as follows. Let $g: \left(\bigoplus_{i \in \omega} \langle a_i \rangle\right)^6 \to \bigoplus_{i \in \omega \setminus \{1, \dots, 6\}} \langle a_i \rangle$ be an isomorphism. Note that the map $h: \left(\bigoplus_{i \in \omega} \langle a_i \rangle\right)^6 \to \bigoplus_{i \in \omega \setminus \{1, \dots, 6\}} \langle a_i \rangle$ given by $h(x_1, x_2, x_3, x_4, x_5, x_6) := g(x_2, x_1, x_4, x_3, x_6, x_5)$ is an isomorphism, too, and hence there exists an automorphism α of $\bigoplus_{i \in \omega \setminus \{1, \dots, 6\}} \langle a_i \rangle$ such that $\alpha(h(x_1, \dots, x_6)) = g(x_1, \dots, x_6)$. Note that for all $x, y, z \in \bigoplus_{i \in \omega} \langle a_i \rangle$ we have

$$g(x,y,x,z,y,z) = \alpha(h(x,y,x,z,y,z)) = \alpha(g(y,x,z,x,z,y)).$$

Extend g to a homomorphism $f: \mathfrak{G}^6 \to \mathfrak{G}$ by defining

$$f_{\{k\}}(b) := \begin{cases} a_k + b & \text{if } k \in \{1, 4, 5\} \\ a_k + 2b & \text{if } k \in \{2, 3, 6\} \end{cases}.$$

This fully determines f by (4.1) and because \mathfrak{G} is generated by b and $(a_i)_{i\in\omega}$. We claim that f preserves \neq . Indeed, suppose that $f(x_1,\ldots,x_6)=f(y_1,\ldots,y_6)$. Write x_i as (c_i,r_ib)

and y_i as $(d_i, s_i b)$ where c_i and d_i are elements of $\bigoplus_{i \in \omega} \langle a_i \rangle$ and $r_i, s_i \in \{0, \ldots, 2n-1\}$. From the definition of f we then obtain that $g(c_1, \ldots, c_6) = g(d_1, \ldots, d_6)$, which implies that $c_1 = d_1, \ldots, c_6 = d_6$ since g is injective. So $f(r_1 b, \ldots, r_6 b) = f(s_1 b, \ldots, s_6 b)$, and it suffices to show that $r_i b = s_i b$ for some $i \in \{1, \ldots, 6\}$. Let $r := r_1 + 2r_2 + 2r_3 + r_4 + r_5 + 2r_6$ and $s := s_1 + 2s_2 + 2s_3 + s_4 + s_5 + 2s_6$. Then

$$f(r_1b, \dots, r_6b) = (r_1a_1 + \dots + r_6a_6) + rb = (r_1a_1 + \dots + r_6a_6) + sb = f(s_1b, \dots, s_6b)$$

forces $r-s=0 \mod 2n$ and $r_i=s_i \mod n$ for every $i\in\{1,\ldots,6\}$, say $r_i=s_i+k_in$. Then as $2r_i=2s_i \mod 2n$ we have

$$r - (2r_2 + 2r_3 + 2r_6) = r_1 + r_4 + r_5 = s_1 + s_4 + s_5 = s - (2s_2 + 2s_3 + 2s_6) \mod 2n.$$

Hence $(k_1 + k_4 + k_5)n = 0 \mod 2n$, so $k_1 + k_4 + k_5$ is even. One of k_1, k_4 , and k_5 must be even, say $k_1 = 2t$ is even (the other two cases can be shown analogously). Then $r_1b = (s_1 + k_1n)b = (s_1 + 2nt)b = s_1b$ since 2nb = 0 and we are done.

Extend α by setting $\alpha(b) := b$ and

$$\alpha(a_1, a_2, a_3, a_4, a_5, a_6) := (a_2, a_1, a_4, a_3, a_6, a_5);$$

again, this determines α on all of G. Then α witnesses that f is a pseudo-Siggers polymorphism: for each $r \in \omega$ we have

$$\alpha f(rb, 1, rb, 1, 1, 1) = \alpha (a_1 + rb + a_3 + 2rb) = \alpha (a_1 + a_3 + 3rb)$$
$$= a_2 + a_4 + 3rb = a_2 + 2rb + a_4 + rb = f(1, rb, 1, rb, 1, 1)$$

and it follows that $\alpha f_{\{1,3\}} = f_{\{2,4\}}$. Similarly, we get $\alpha f_{\{2,5\}} = f_{\{2,5\}}$ and $\alpha f_{\{4,6\}} = f_{\{4,6\}}$, and hence f is pseudo-Siggers by Proposition 4.4.

Finally, the implication from (3) to (1) is a well-known general fact for ω -categorical structures that follows from the lift lemma presented in [17] (see [5]): if the model-complete core of an ω -categorial structure \mathfrak{B} has a pseudo-Siggers polymorphism, then so has \mathfrak{B} . \square

Item (3) in Theorem 5.17 is the condition from the first infinite-domain tractability conjecture (see [5]). We already know from Theorem 5.16, Theorem 5.17, and from Theorem 2.2 that if \mathfrak{G} is an ω -categorical abelian group such that (\mathfrak{G}, \neq) has a pseudo-Siggers polymorphism, then $\operatorname{Pol}(\mathfrak{G}, \neq)$ cannot have a uniformly continuous minor-preserving map to \mathscr{P} , unless P=NP. It is surprisingly difficult to verify this also without the complexity-theoretic assumption; however, by bounding the orbit growth⁶ of (\mathfrak{G}, \neq) this follows from a result of [1], as we will see below.

Proposition 5.18. Let \mathfrak{G} be an ω -categorical abelian group. Then (\mathfrak{G}, \neq) has a pseudo-Siggers polymorphism if and only if $\operatorname{Pol}(\mathfrak{G}, \neq)$ has no uniformly continuous minorpreserving map to \mathscr{P} .

Proof. Let \mathfrak{H} be the model companion of \mathfrak{G} . First suppose that (\mathfrak{G}, \neq) has no uniformly continuous minor-preserving map to \mathscr{P} . Then neither has (\mathfrak{H}, \neq) , by Proposition 2.4, and so (\mathfrak{H}, \neq) has a pseudo-Siggers polymorphism by Theorem 2.5. Hence, (\mathfrak{G}, \neq) has a pseudo-Siggers polymorphism by Theorem 5.17.

⁶We thank Michael Kompatscher for discussing the growth rate of $\mathbb{Z}_2^{(\omega)}$.

Now suppose that (\mathfrak{G}, \neq) has a uniformly continuous minor-preserving map to \mathscr{P} . It is known that if \mathfrak{B} is an ω -categorical structure with a uniformly continuous minor-preserving map to \mathscr{P} and if the number of orbits of n-tuples of $\operatorname{Aut}(\mathfrak{B})$ grows slower than doubly exponentially, then \mathfrak{B} cannot have a pseudo-Siggers polymorphism [2]. Steitz [44] proved that for every ω -categorical ω -stable structure \mathfrak{B} there exists $m \in \omega$ such that

the number of orbits of n-tuples of $\operatorname{Aut}(\mathfrak{B})$ is smaller than 2^{mn^2} . The groups $\mathbb{Z}_n^{(\omega)}$ and $\mathbb{Z}_n^{(\omega)} \oplus \mathbb{Z}_{2n}$ are for every $n \in \omega$ totally categorical: the models of cardinality κ are clearly isomorphic to $\mathbb{Z}_n^{(\kappa)}$ or to $\mathbb{Z}_n^{(\kappa)} \oplus \mathbb{Z}_{2n}$, respectively, and hence in particular ω -stable (see [45]).

6. Semilattices

A semilattice is an algebra $(S; \land)$ where \land is a binary operation that is associative, commutative, and idempotent. For $a, b \in A$ we define $a \leq b$ iff a = ab = ba. Clearly, $(A; \leq)$ is a partial order and polymorphisms of $(A; \land)$ are monotone with respect to \leq . Note that semilattices with a greatest element \bot are special monoids (where the greatest element takes the role of 1). As in the case of monoids, we often omit the symbol \land and write ab instead of $a \land b$.

Example 6.1. Let \mathfrak{P}_n be the Boolean algebra with the atoms $\{1,\ldots,n\}$. Clearly, the $\{\wedge\}$ -reduct \mathfrak{S}_n of \mathfrak{P}_n is a semilattice. It is well-known that every finite semilattice $(S;\wedge)$ has the following embedding e into \mathfrak{S}_n , for n:=|S|, which we recall here for the convenience of the reader: if b is any bijection between S and $\{1,\ldots,n\}$, define

$$e(x) := \bigvee_{y \le x} b(y).$$

This map is injective: if $e(x_1) \leq e(x_2)$ then in particular $b(x_1) \leq e(x_2)$ and since $b(x_1)$ is an atom we must have $b(x_1) \leq b(y)$ for some $y \leq x_2$. Since b(y) is an atom we must have $x_1 = y$ and hence $x_1 \leq x_2$. Together with the symmetric argument we obtain that $e(x_1) = e(x_2)$ implies that $x_1 = x_2$.

Moreover, e preserves \wedge : let $x_1, x_2, x_3 \in S$ be such that $x_1 \wedge x_2 = x_3$. Let a be an atom of \mathfrak{S}_n and $c \in S$ be such that b(c) = a. Then

$$a \leq e(x_1) \wedge e(x_2)$$
 if and only if $a \leq e(x_1)$ and $a \leq e(x_2)$ if and only if $c \leq x_1$ and $c \leq x_2$ if and only if $c \leq (x_1 \wedge x_2) = x_3$ if and only if $a \leq e(x_3)$.

This shows that $e(x_1) \wedge e(x_2) = e(x_3)$ and concludes the proof.

Example 6.2. The class \mathcal{L} of all finite semilattices forms an amalgamation class. To see this, suppose that \mathfrak{B}_1 and \mathfrak{B}_2 are two finite semilattices such that $B_1 \cap B_2$ is the domain of a subsemilattice \mathfrak{A} of both \mathfrak{B}_1 and \mathfrak{B}_2 . We have to prove that there exists a finite semilattice

 \triangle

 \mathfrak{C} and embeddings $e_1 \colon \mathfrak{B}_1 \hookrightarrow \mathfrak{C}$ and $e_2 \colon \mathfrak{B}_2 \hookrightarrow \mathfrak{C}$ such that $e_1(a) = e_2(a)$ for all $a \in A$. Note that the poset $(A; \leq)$ induced by \mathfrak{A} is a subposet of the posets $(B_1; \leq)$ and $(B_2; \leq)$ induced by \mathfrak{B}_1 and \mathfrak{B}_2 , respectively. Let \mathfrak{C} be the Dedekind-McNeille completion [33] of the poset amalgam of $(B_1; \leq)$ and $(B_2; \leq)$. Then \mathfrak{C} has the required properties.

We write \mathfrak{U} for the Fraïssé-limit of \mathcal{L} , i.e., for the up to isomorphism unique countable universal homogeneous semilattice, studied e.g. in [34] where a finite axiomatisation of its first-order theory is presented. Note that \mathfrak{U} is ω -categorical, because it is uniformly locally finite: in the subalgebra of \mathfrak{U} generated by u_1, \ldots, u_n there are precisely the elements of the form $\bigwedge_{u \in V}$ for a subset V of $\{u_1, \ldots, u_n\}$, and hence their number is bounded by 2^n . Also note that \mathcal{L} is closed under taking finite direct products, and it follows from Proposition 3.7 that $\mathfrak{U}^2 \hookrightarrow \mathfrak{U}$.

 $CSP(\mathfrak{U}, \neq)$ might be viewed as a special case of Horn-Horn set constraints [10], which might be solved in polynomial time. We do not want to introduce Horn-Horn sets constraints here, but for the convenience show how to derive a polynomial-time algorithm for $CSP(\mathfrak{U}, \neq)$ from Fact 24 in [10], which implies the following.

Lemma 6.3. Let $\phi = x \neq y \land \psi$ by a primitive positive formula of (\mathfrak{U}, \neq) where ψ does not contain formulas involving \neq . Then ϕ is satisfiable if and only if either $\psi \land x = 0 \land y = 1$ or $\psi \land x = 1 \land y = 0$ is satisfiable in $(\{0,1\}; \land, 0, 1)$.

Proposition 6.4. $CSP(\mathfrak{U}, \neq)$ can be solved in polynomial time.

Proof. Since $\mathfrak{U}^2 \hookrightarrow \mathfrak{U}$ (Proposition 3.7) it suffices to consider the situation that ϕ is of the form $x \neq y \land \psi$ where ψ does not contain formulas involving \neq . By the previous lemma ϕ is satisfiable if and only if either $\psi \land x = 0 \land y = 1$ or $\psi \land x = 1 \land y = 0$ is satisfiable in $(\{0,1\}; \land, 0, 1)$. However $CSP(\{0,1\}; \land, 0, 1)$ can be solved in polynomial time (see, e.g., [30]), to which the result follows.

In this section we prove the following dichotomy result.

Theorem 6.5. Let \mathfrak{S} be a countable ω -categorical semilattice. Then either

- (1) there is a uniformly continuous minor-preserving from $\operatorname{Pol}(\mathfrak{S}, \neq)$ to the clone of projections, in which case $\operatorname{CSP}(\mathfrak{S}, \neq)$ is NP-hard, or
- (2) the model-companion \mathfrak{C} of \mathfrak{S} is isomorphic to \mathfrak{U} . In this case $\operatorname{Pol}(\mathfrak{S}, \neq)$ is in P.

We first prove that the two cases are indeed disjoint. The identities that appear in the next proposition have been discovered by Jakub Rydval in a different context [18]; note that these identities have *height one* and hence are preserved by all minor-preserving maps [4].

Proposition 6.6. There are $f, g_1, \ldots, g_4 \in \text{Pol}(\mathfrak{U}, \neq)$ such that for all $x, y \in U$

$$g_1(y, x, x) = f(x, y, x, x),$$
 $g_2(y, x, x) = f(y, x, x, x),$
 $g_1(x, y, x) = f(x, x, y, x),$ $g_2(x, y, x) = f(x, x, y, x),$
 $g_1(x, x, y) = f(x, x, x, y),$ $g_2(x, x, y) = f(x, x, x, y),$

```
g_3(y, x, x) = f(y, x, x, x), g_4(y, x, x) = f(y, x, x, x),

g_3(x, y, x) = f(x, y, x, x), g_4(x, y, x) = f(x, y, x, x),

g_3(x, x, y) = f(x, x, x, y), g_4(x, x, y) = f(x, x, y, x).
```

Proof. Let Φ be the following (infinite) set of atomic $\{\land, \neq\}$ -formulas. The variables of these formulas consist of the elements of semilattices $(W_0; \land)$, $(W_1; \land)$, ..., $(W_4; \land)$ such that there is an isomorphism α_0 from \mathfrak{U}^4 to $(W_0; \land)$ and an isomorphism α_i from \mathfrak{U}^3 to $(W_i; \land)$ for each $i \in \{1, ..., 4\}$. The formulas in Φ come from three groups.

- For $i \in \{0, ..., 4\}$ and $x, y, z \in W_i$ the set Φ contains the atomic formula $(x \wedge y) = z$ if and only if $(x \wedge z) = z$ holds in $(W_i; \wedge)$.
- For $i \in \{0, ..., 4\}$ and distinct $x_1, x_2 \in W_i$ the set Φ contains $x_1 \neq x_2$.
- Finally, whenever $g_i(x_1, x_2, x_3) = f(y_1, y_2, y_3, y_4)$ is an identity from the statement then the set Φ contains the atomic formula x = y where x is the variable $\alpha_i(x_1, x_2, x_3) \in W_i$ and y is the variable $\alpha_0(y_1, \ldots, y_4) \in W_0$.

Note that any satisfying assignment of Φ in \mathfrak{U} restricted to the elements of W_i defines a polymorphism of \mathfrak{U} because of the atomic formulas of the first group; these polymorphisms will be injective because of the atomic formulas of the second group; and jointly they satisfy the identities given in the statement because of the atomic formulas of the third group.

Since $\mathfrak U$ is ω -categorical, it suffices to show that every finite subset ϕ of formulas in Φ is satisfiable. If ϕ contains no atomic formulas of the form $x \neq y$ then ϕ is trivially satisfiable by mapping all variables to the same element of $\mathfrak U$. Since $\mathfrak U^2 \hookrightarrow \mathfrak U$ it suffices by Proposition 3.7 to consider the situation that ϕ is of the form $x \neq y \land \psi$ where ψ does not contain formulas involving \neq . By Lemma 6.3 it suffices to show that either $\psi \land x = 0 \land y = 1$ or or $\psi \land x = 1 \land y = 0$ is satisfiable in $(\{0,1\}; \land, 0, 1)$. We construct a solution to $\psi \land x = 1 \land y = 0$. Suppose that x, which must be assigned to 1, is of the form $\alpha_1(y, x, x)$. If the variable $x_0 := \alpha_0(x, y, x, x)$ is present in ϕ , assign it to 1, too. If the variable $x_4 := \alpha_4(x, x, y)$ is present in ϕ , assign it to 1, too. Note that assigning all other variables to 0 is a satisfying solution to ϕ .

Before we prove Theorem 6.5 we consider a special case that will be used in the proof. Every linear order $(A; \leq)$ gives rise to a semilattice $(A; \wedge)$, by defining $a \wedge b := a$ iff $a \leq b$. Semilattices that arise from linear orders in this way are called *linear*. A semilattice $(S; \wedge)$ is called *semilinear* if for every $x \in S$ the subsemilattice induced by $\{y \in S \mid y \leq x\}$ is linear.

Lemma 6.7. Let $(S; \wedge)$ be a countable ω -categorical semilinear semilattice with |S| > 1. Then $Pol(S; \wedge, \neq)$ has a uniformly continuous minor-preserving map to \mathscr{P} .

Proof. If |S| = 2 then the statement follows from the proof of Schaefers classification: $(S; \wedge, \neq)$ is neither preserved by min, max, or constant operations because these operations do not preserve \neq , and neither preserved by minority and majority because these operations do not preserve \wedge . If |S| > 2 is finite then the statement holds because of the well-known fact that in this case all polymorphisms of $(S; \neq)$ only depend on one argument. Otherwise, $(S; \wedge)$ is countably infinite, and so is its model companion $(C; \wedge)$. First consider the case that $(C; \wedge)$ is linear. Every ω -categorical infinite linear order embeds all finite linear

orders, and thus $(C; \wedge)$ and $(\mathbb{Q}; \min)$ have the same age. Since both structures are model-complete, they have the same first-order theory, and by the ω -categoricity of $(\mathbb{Q}; \min)$ they are isomorphic. Let

$$R := \{ (x, y, z) \in \mathbb{Q}^3 \mid x = y < z \lor x = z < y \}.$$

The primitive positive formula $\psi(x, y, z)$ given by

$$\min(y, z) = x \land y \neq z$$

defines R in $(\mathbb{Q}; \min, \neq)$. It follows from Theorem 51 in [15] in combination with Theorem 28 in [16] that $\operatorname{Pol}(\mathbb{Q}; R)$ has a (uniformly) continuous homomorphism to \mathscr{P} and hence the same holds for $\operatorname{Pol}(\mathbb{Q}; \min, \neq)$. It follows in particular that there exists a uniformly continuous minor-preserving map from $\operatorname{Pol}(\mathbb{Q}; \min, \neq)$ to \mathscr{P} (see [4]).

To prove the general case, choose an element $x \in C$ such that $T := \{y \in C \mid y \leq x\}$ contains more than one element; such an element x must exist because C has more than one element. Then the subsemilattice of $(C; \land)$ with domain T is linear, and we have proved above that $\operatorname{Pol}(T; \land, \neq)$ has a uniformly continuous minor-preserving map to \mathscr{P} . Since T is primitive positive definable in an expansion of the model-complete core $(C; \land, \neq)$, it follows that $\operatorname{Pol}(C; \land, \neq)$ has a uniformly continuous minor-preserving map to \mathscr{P} , too (Proposition 2.4).

Proof of Theorem 6.5. If there is a uniformly continuous minor-preserving map from $\operatorname{Pol}(\mathfrak{S},\neq)$ to \mathscr{P} , then the NP-hardness of $\operatorname{CSP}(\mathfrak{S},\neq)$ follows from Theorem 2.2. In this case, the model-companion of \mathfrak{S} cannot be isomorphic to \mathfrak{U} : otherwise, there would be a minor-preserving map from \mathfrak{U} via $\operatorname{Pol}(\mathfrak{S},\neq)$ to \mathscr{P} , and such a map preserves the identities from Proposition 6.6 (see [4]). Clearly, these identities cannot be satisfied by projections on a two-element set, so we reached a contradiction.

Now suppose that there is no uniformly continuous minor-preserving map from $\operatorname{Pol}(\mathfrak{S},\neq)$ to \mathscr{P} . We will prove that for every $n \in \omega$, \mathfrak{S}_n embeds into every infinite ω -categorical semilattice \mathfrak{T} such that $\operatorname{Pol}(\mathfrak{T},\neq)$ has no uniformly continuous minor-preserving map to \mathscr{P} . The proof is by induction on $n \in \omega$. The statement is clear for n=1, and for n=2 the statement follows from Lemma 6.7. Now suppose inductively that $n \geq 2$ is such that \mathfrak{S}_n embeds into every ω -categorical semilattice \mathfrak{T} such that $\operatorname{Pol}(\mathfrak{T},\neq)$ has no uniformly continuous minor-preserving map to \mathscr{P} . We would like to show that \mathfrak{S}_{n+1} embeds into \mathfrak{T} , too. We may assume that \mathfrak{T} is model-complete; otherwise, let \mathfrak{C} be the model-companion of \mathfrak{T} , which exists and is again an ω -categorical countable semilattice (Lemma 3.4). Moreover, there is a uniformly continuous minor-preserving map from $\operatorname{Pol}(\mathfrak{T},\neq)$ to $\operatorname{Pol}(\mathfrak{C},\neq)$, and hence there is no uniformly continuous minor-preserving map from $\operatorname{Pol}(\mathfrak{C},\neq)$ to \mathscr{P} . We may therefore replace \mathfrak{T} by \mathfrak{C} .

Claim 1. For all $a, b \in T$ with a < b the subsemilattice of \mathfrak{T} induced by the interval $[a, b] := \{x \in T \mid a \leq x \leq b\}$ embeds \mathfrak{S}_n . Clearly, the structure $(\mathfrak{T}, \neq, a, b)$ is a model-complete core, too. Proposition 2.4 thus implies that $(\mathfrak{T}, \neq, a, b)$ has no uniformly continuous minor-preserving map to \mathscr{P} . The set [a, b] is primitive positive definable in (\mathfrak{T}, a, b) , and $([a, b]; \wedge)$ is an ω -categorical subsemilattice of \mathfrak{T} . So there is a uniformly continuous

minor-preserving map from $\operatorname{Pol}(\mathfrak{T},\neq,a,b)$ to $\operatorname{Pol}([a,b];\wedge,\neq)$ and hence $\operatorname{Pol}([a,b];\wedge,\neq)$ cannot have a uniformly continuous minor-preserving map to \mathscr{P} . The inductive assumption then implies that there exists an embedding $e \colon \mathfrak{S}_n \hookrightarrow ([a,b];\wedge)$.

Claim 2. There exist $a, b, c \in T$ such that a < b < c and

(6.1)
$$f(b, a, b, a, a, a) \neq f(c, a, c, a, a, a)$$

(6.2) or
$$f(a, b, a, a, b, a) \neq f(a, c, a, a, c, a)$$
.

Suppose otherwise. By Claim 1, there exists an embedding $e: \mathfrak{S}_2 \to \mathfrak{T}$.


Let $y, z \in T$ be the images of the atoms of \mathfrak{S}_2 under e, and let $w := e(\mathbf{1})$ and $x := e(\mathbf{0})$.

Again by Claim 1, there exists an embedding e' of \mathfrak{S}_2 into [y, w].

Let u and v be the images of the atoms of \mathfrak{S}_2 under e';

we may then replace y by uv and then x by yz;

in the resulting constellation we have uv = y and yz = x; see the figure to the right.



Since we assumed that (6.1) neither holds for (a, b, c) = (x, y, w) nor for (a, b, c) = (x, z, w), we have

$$f(y, x, y, x, x, x) = f(w, x, w, x, x, x) = f(z, x, z, x, x, x)$$

and so

$$f(x, x, x, x, x, x) = f(y, x, y, x, x, x) f(z, x, z, x, x, x) = f(y, x, y, x, x, x).$$

A similar argument applied to the image of e' and (6.2) shows that f(y, y, y, y, y, y) = f(y, w, y, y, w, y). Hence,

$$\begin{split} f(y,z,y,x,z,x) &= f(y,w,y,y,w,y) f(y,z,y,z,z,z) \\ &= f(y,y,y,y,y) f(y,z,y,z,z,z) \\ &= f(y,x,y,x,x,x) = f(x,x,x,x,x,x,x) \end{split}$$

and so by (3.1)

$$f(y, z, y, x, z, x) = f(x, x, x, x, x, x, x) = f(z, y, x, y, x, z)$$

contradicting that f preserves \neq .

Suppose that $f(b, a, b, a, a, a) \neq f(c, a, c, a, a, a)$; the other case from Claim 2 can be shown similarly. Let $g: \{b, c\} \times [a, b] \to T$ be given by

$$g(x,y) := f(x,y,x,y,y,y).$$

We show that g is injective. Suppose first that g(b,y) = g(b,y') for $y,y' \in [a,b]$. Then by (3.1) we have f(y,b,y,b,y,y) = f(y',b,y',b,y',y'). Since yb = y and y'b = y' we have

$$f(y,y,y,y,y,y) = g(b,y)f(y,b,y,b,y,y) = g(b,y')f(y',b,y',b,y',y') = f(y',y',y',y',y',y',y')$$

and so y=y' as f preserves \neq . A similar argument shows that g(c,y)=g(c,y') forces y=y'. Finally, if g(b,x)=g(c,x) then multiplying both sides by f(c,a,c,a,a,a) we obtain

f(b, a, b, a, a, a) = f(c, a, c, a, a, a), contradicting our hypothesis. This shows that $\mathfrak{S}_1 \times \mathfrak{S}_n$ embeds into \mathfrak{T} . Since $\mathfrak{S}_1 \times \mathfrak{S}_n \simeq \mathfrak{S}_{n+1}$ we have that \mathfrak{S}_{n+1} embeds into \mathfrak{T} , which concludes the induction.

We obtain that \mathfrak{S} and \mathfrak{C} have the same age as \mathfrak{U} . Since \mathfrak{C} and \mathfrak{U} are model-complete, they must be isomorphic. The polynomial-time tractability of $CSP(\mathfrak{S}, \neq)$ then follows from Proposition 6.4.

Corollary 6.8. Let \mathfrak{S} be a countable ω -categorical semilattice. Then $\mathrm{CSP}(\mathfrak{S}, \neq)$ is in P or NP-hard.

7. CONCLUSION AND FUTURE WORK

In previous work about CSPs for ω -categorical structures \mathfrak{A} , algebras were used to analyse the polymorphism clones of \mathfrak{A} (and these algebras are oligomorphic, but never ω -categorical). In this article, in contrast, the structure \mathfrak{A} itself is assumed to be an algebra, expanded by the disequality relation. Our result underlines the importance of

- uniformly continuous minor-preserving maps $\operatorname{Pol}(\mathfrak{A}, \neq) \to \mathscr{P}$ as a tool for proving hardness (in Section 6), and
- pseudo-Siggers polymorphisms of model-complete cores to obtain structural (and subsequently algorithmic) results.

We are not aware of any previous result that would use the Siggers (or pseudo-Siggers) identity directly, even for algebras over finite domains. For example, the result that every structure with a finite domain that has a Siggers polymorphism also has a cyclic polymorphism departs from the (a priori) weaker assumption that the structure has a Taylor polymorphism [3]. We close with some open problems.

- Let \mathfrak{A} be an ω -categorical model-complete algebra. Suppose that $\mathfrak{A}^2 \hookrightarrow \mathfrak{A}$. Does then (\mathfrak{A}, \neq) have a binary pseudo-symmetric polymorphism? The converse is false (a counterexample can be found in the class of ω -categorical model-complete algebras with a single unary function symbol). The forward implication is true if \mathfrak{A}^2 is isomorphic to \mathfrak{A} [9].
- Let \mathfrak{L} be an ω -categorical lattice. Does $\mathrm{CSP}(\mathfrak{L}, \neq)$ satisfy a complexity dichotomy P versus NP-hard? We may assume that \mathfrak{L} is model-complete. Using similar techniques as in Section 6 it is possible to show that if $\mathrm{Pol}(\mathfrak{L}, \neq)$ does not have uniformly continuous minor-preserving maps to \mathscr{P} , then \mathfrak{L} must have a square embedding $\mathfrak{L}^2 \hookrightarrow \mathfrak{L}$. So we may assume that the age of \mathfrak{L} is closed under taking products and thus contains the class of all distributive lattices. The countable homogeneous universal distributive lattice has a uniformly continuous continuous minor-preserving maps to \mathscr{P} (this has essentially been observed in [10]). Are there any other examples of model-complete ω -categorical lattices with $\mathfrak{L}^2 \hookrightarrow \mathfrak{L}$? Note that the class of all finite lattices forms an amalgamation class, but the countable homogeneous lattice \mathfrak{L} with this age is not ω -categorical: this can be seen from the fact that every finite lattice can be embedded in a finite lattice with three

generators [24] (page 224). Hence, there exists an infinite number of inequivalent formulas with three variables over \mathfrak{L} .

REFERENCES

- [1] Libor Barto, Michael Kompatscher, Miroslav Olšák, Trung Van Pham, and Michael Pinsker, *The equivalence of two dichotomy conjectures for infinite domain constraint satisfaction problems*, Proceedings of the 32nd annual ACM/IEEE symposium on logic in computer science lics'17, 2017. Preprint arXiv:1612.07551.
- [2] ______, Equations in oligomorphic clones and the constraint satisfaction problem for ω-categorical structures, Journal of Mathematical Logic 19 (2019), no. 2, #1950010. An extended abstract appeared at the Proceedings of the 32nd Annual ACM/IEEE Symposium on Logic in Computer Science LICS'17.
- [3] Libor Barto and Marcin Kozik, Absorbing subalgebras, cyclic terms and the constraint satisfaction problem, Logical Methods in Computer Science 8/1 (2012), no. 07, 1–26.
- [4] Libor Barto, Jakub Opršal, and Michael Pinsker, *The wonderland of reflections*, Israel Journal of Mathematics **223** (2018), no. 1, 363–398.
- [5] Libor Barto and Michael Pinsker, *The algebraic dichotomy conjecture for infinite domain constraint satisfaction problems*, Proceedings of the 31th Annual IEEE Symposium on Logic in Computer Science LICS'16, 2016, pp. 615–622. Preprint arXiv:1602.04353.
- [6] Jon Barwise and Paul Eklof, *Infinitary properties of abelian torsion groups*, Annals of Mathematical Logic 2 (1970), no. 1, 25–68.
- [7] Manuel Bodirsky, Cores of countably categorical structures, Logical Methods in Computer Science (LMCS) 3 (2007), no. 1, 1–16.
- [8] ______, Complexity classification in infinite-domain constraint satisfaction, 2012. Mémoire d'habilitation à diriger des recherches, Université Diderot Paris 7. Available at arXiv:1201.0856.
- [9] Manuel Bodirsky, Hubie Chen, Jan Kára, and Timo von Oertzen, Maximal infinite-valued constraint languages, Theoretical Computer Science (TCS) 410 (2009), 1684–1693. A preliminary version appeared at ICALP'07.
- [10] Manuel Bodirsky and Martin Hils, *Tractable set constraints*, Journal of Artificial Intelligence Research 45 (2012), 731–759.
- [11] Manuel Bodirsky, Martin Hils, and Barnaby Martin, On the scope of the universal-algebraic approach to constraint satisfaction, Proceedings of the symposium on logic in computer science (LICS), 2010July, pp. 90–99.
- [12] ______, On the scope of the universal-algebraic approach to constraint satisfaction, Logical Methods in Computer Science (LMCS) 8 (2012), no. 3. An extended abstract that announced some of the results appeared in the proceedings of Logic in Computer Science (LICS'10).
- [13] Manuel Bodirsky and Marcello Mamino, Constraint satisfaction problems over numeric domains, The constraint satisfaction problem: Complexity and approximability, 2017, pp. 79–111.
- [14] Manuel Bodirsky and Jaroslav Nešetřil, Constraint satisfaction with countable homogeneous templates, Journal of Logic and Computation 16 (2006), no. 3, 359–373.
- [15] Manuel Bodirsky and Michael Pinsker, Reducts of Ramsey structures, AMS Contemporary Mathematics, vol. 558 (Model Theoretic Methods in Finite Combinatorics) (2011), 489–519.
- [16] _____, Topological Birkhoff, Transactions of the American Mathematical Society **367** (2015), 2527–2549.
- [17] _____, Canonical functions: a proof via topological dynamics, 2016. Preprint arXiv:1610.09660.
- [18] Manuel Bodirsky and Jakub Rydval, Temporal constraint satisfaction problems in fixed point logic, 2019. Preprint.
- [19] M. Broxvall, P. Jonsson, and J. Renz, *Disjunctions, independence, refinements*, Artificial Intelligence **140** (2002), no. 1/2, 153–173.

- [20] Andrei A. Bulatov, A dichotomy theorem for nonuniform CSPs, 58th IEEE annual symposium on foundations of computer science, FOCS 2017, berkeley, ca, usa, october 15-17, 2017, 2017, pp. 319–330.
- [21] Andrei A. Bulatov and Víctor Dalmau, A simple algorithm for Mal'tsev constraints, SIAM Journal on Computing 36 (2006), no. 1, 16–27.
- [22] Andrei A. Bulatov, Andrei A. Krokhin, and Peter G. Jeavons, Classifying the complexity of constraints using finite algebras, SIAM Journal on Computing 34 (2005), 720–742.
- [23] Filippo Calderoni and Simon Thomas, The bi-embeddability relation for countable abelian groups, Transactions of the AMS **371** (2019), 2237–2254.
- [24] Richard A. Dean, Component subsets of the free lattice on n generators, Proceedings of the American Mathematical Society 7 (1956), no. 2, 220–226.
- [25] Tomás Feder and Moshe Y. Vardi, The computational structure of monotone monadic SNP and constraint satisfaction: a study through Datalog and group theory, SIAM Journal on Computing 28 (1999), 57–104.
- [26] M. Goldmann and A Russell, *The complexity of solving equations over finite groups*, Information and Computation 178 (2002), no. 1, 253–262.
- [27] Wilfrid Hodges, *Model theory*, Cambridge University Press, 1993.
- [28] _____, A shorter model theory, Cambridge University Press, Cambridge, 1997.
- [29] Irvin Kaplansky, Infinite abelian groups, The University of Michigan Press, Ann Arbor, 1969.
- [30] Ondrej Klíma, Pascal Tesson, and Denis Thérien, Dichotomies in the complexity of solving systems of equations over finite semigroups, Theory Comput. Syst. 40 (2007), no. 3, 263–297.
- [31] Andrei A. Krokhin and Stanislav Zivny (eds.), *The constraint satisfaction problem: Complexity and approximability*, Dagstuhl Follow-Ups, vol. 7, Schloss Dagstuhl Leibniz-Zentrum fuer Informatik, 2017.
- [32] Richard E. Ladner, On the structure of polynomial time reducibility, Journal of the ACM 22 (1975), no. 1, 155–171.
- [33] H. M. MacNeille, Partially ordered sets, Transactions of the AMS 42 (1937), no. 3, 416–460.
- [34] John K. Truss Manfred Droste Dietrich Kuske, On homogeneous semilattices and their automorphism groups, Order 16 (1999), 31–56.
- [35] Yuri V. Matiyasevich, Hilbert's tenth problem, MIT Press, Cambridge, Massachusetts, 1993.
- [36] Cristopher Moore, Pascal Tesson, and Denis Thérien, Satisfiability of systems of equations over finite monoids, Mathematical foundations of computer science 2001, 26th international symposium, MFCS 2001 marianske lazne, czech republic, august 27-31, 2001, proceedings, 2001, pp. 537–547.
- [37] Greg Nelson and Derek C. Oppen, Fast decision procedures based on congruence closure, Journal of the ACM 27 (1980), no. 2, 356–364.
- [38] J. G. Rosenstein, ℵ₀-categoricity of groups, Journal of Algebra **25** (1973), no. 3, 435–467.
- [39] D. Saracino and C. Wood, QE nil-2 groups of exponent 4, Journal of Algebra 76 (1982), 337–352.
- [40] Dan Saracino, Model companions for ℵ₀-categorical theories, Proceedings of the AMS 39 (1973), 591–598.
- [41] Eugene Schenkman, Group theory, Krieger Pub Co, 1975.
- [42] Alexander Schrijver, *Theory of linear and integer programming*, Wiley Interscience Series in Discrete Mathematics and Optimization, 1998.
- [43] Mark H. Siggers, A strong Mal'cev condition for varieties omitting the unary type, Algebra Universalis **64** (2010), no. 1, 15–20.
- [44] Philip W. Steitz, Upper bounds for growth in the ryll-nardzewski function of an ω -categorical, ω -stable theory, Israel Journal of Mathematics 77 (1992), no. 3, 335–343.
- [45] Katrin Tent and Martin Ziegler, A course in model theory, Lecture Notes in Logic, Cambridge University Press, 2012.

[46] Dmitriy Zhuk, A proof of CSP dichotomy conjecture, 58th IEEE annual symposium on foundations of computer science, FOCS 2017, berkeley, ca, usa, october 15-17, 2017, 2017, pp. 331–342. https://arxiv.org/abs/1704.01914.