

Singular Vectors From Singular Values

Weiwei Xu* Michael K. Ng[†]

Abstract

In the recent paper [1], Denton et al. provided the eigenvector-eigenvalue identity for Hermitian matrices, and a survey was also given for such identity in the literature. The main aim of this paper is to present the identity related to singular vectors and singular values of a general matrix.

Keywords: Singular values, singular vectors, matrix, submatrices and identity

Mathematics Subject Classification: 15A18

1 Introduction

In the recent paper [1], Denton et al. provided the eigenvector-eigenvalue identity for Hermitian matrices, and a survey was also given for such identity in the literature. The main aim of this paper is to present the identity related to singular vectors and singular values of a general matrix. Indeed, this topic has been studied by Thompson [2]. In particular, the interlacing inequalities for singular values of submatrices are derived in [2].

Let A be a $m \times n$ matrix with singular values $\sigma_i(A)$ and normed left singular vectors u_i and right singular vectors v_l . A^H denotes the conjugate transpose of A . The elements of each left singular vector are denoted u_{ij} and right singular vector v_{ls} . Let \bar{A}_j be $(n-1) \times n$ matrix after deleting the j -th row of A and $\sigma_k(\bar{A}_j)$ be singular values of \bar{A}_j . Let \hat{A}_s be $n \times (n-1)$

*Nanjing University of Information Science & Technology, China (Email: wwx19840904@sina.com). Research supported in part by Natural Science Foundation of China under Grant No. 11971243 and Natural Science Foundation of Jiangsu Province under grant BK20181405.

[†]Department of Mathematics, The University of Hong Kong, Pokfulam, Hong Kong (Email: mng@maths.hku.hk). Research supported in part by the HKRGC GRF 12306616, 12200317, 12300218 and 12300519, and HKU Grant 104005583.

matrix after deleting the s -th column of A and $\sigma_t(\hat{A}_s)$ be singular values of \hat{A}_s .

Lemma 1[1] *Let A be a $n \times n$ Hermitian matrix with distinct eigenvalues $\lambda_i(A)$ and normed eigenvectors v_i . The elements of each eigenvector are denoted $v_{i,j}$. Let M_j be the $(n-1) \times (n-1)$ submatrix of A that results from deleting the j -th column and the j -th row, with eigenvalues $\lambda_k(M_j)$. Then*

$$|v_{ij}|^2 = \frac{\prod_{k=1}^{n-1} (\lambda_i(A) - \lambda_k(M_j))}{\prod_{k=1, k \neq i}^n (\lambda_i(A) - \lambda_k(A))}, \quad 1 \leq i, j \leq n.$$

Lemma 2[1] *Let A be a $n \times n$ Hermitian matrix with eigenvalues $\lambda_i(A)$ and normed eigenvectors v_i . The elements of each eigenvector are denoted $v_{i,j}$. Let M_j be the $(n-1) \times (n-1)$ submatrix of A that results from deleting the j -th column and the j -th row, with eigenvalues $\lambda_k(M_j)$. Then*

$$|v_{ij}|^2 \prod_{k=1, k \neq i}^n (\lambda_i(A) - \lambda_k(A)) = \prod_{k=1}^{n-1} (\lambda_i(A) - \lambda_k(M_j)), \quad 1 \leq i, j \leq n.$$

In this paper, we establish the following theorem. We first present singular vectors from distinct singular values of $n \times n$ matrix A .

Theorem 1 *Let $A = U^H \Sigma V \in \mathbf{C}^{n \times n}$ with distinct singular values $\sigma_i(A)$ and $U = (u_{ij}), 1 \leq i, j \leq n$ be left singular vector matrix of A and $V = (v_{ls}), 1 \leq l, s \leq n$ be right singular vector matrix of A . Let \bar{A}_j be $(n-1) \times n$ matrix after deleting the j -th row of A and $\sigma_k(\bar{A}_j)$ be singular values of \bar{A}_j . Let \hat{A}_s be $n \times (n-1)$ matrix after deleting the s -th column of A and $\sigma_t(\hat{A}_s)$ be singular values of \hat{A}_s . Then*

$$|u_{ij}|^2 = \frac{\prod_{k=1}^{n-1} (\sigma_i^2(A) - \sigma_k^2(\bar{A}_j))}{\prod_{k=1, k \neq i}^n (\sigma_i^2(A) - \sigma_k^2(A))}, \quad 1 \leq i, j \leq n,$$

$$|v_{ls}|^2 = \frac{\prod_{t=1}^{n-1} (\sigma_l^2(A) - \sigma_t^2(\hat{A}_s))}{\prod_{t=1, t \neq l}^n (\sigma_l^2(A) - \sigma_t^2(A))}, \quad 1 \leq l, s \leq n.$$

Then we deduce singular vectors from singular values of $m \times n$ matrix A .

Theorem 2 Let $A = U^H \Sigma V \in \mathbf{C}^{m \times n}$ with singular values $\sigma_1(A) \geq \sigma_2(A) \geq \dots \geq \sigma_r(A) > 0, \sigma_{r+1}(A) = \dots = \sigma_{\min\{m,n\}}(A) = \dots = \sigma_{\max\{m,n\}}(A) = 0$ and $U = (u_{ij}), 1 \leq i, j \leq m$ be left singular vector matrix of A and $V = (v_{ls}), 1 \leq l, s \leq n$ be right singular vector matrix of A . Let \bar{A}_j be $(m-1) \times n$ matrix after deleting the j -th row of A and $\sigma_k(\bar{A}_j)$ be singular values of \bar{A}_j . Let \hat{A}_s be $m \times (n-1)$ matrix after deleting the s -th column of A and $\sigma_t(\hat{A}_s)$ be singular values of \hat{A}_s . Then

$$|u_{ij}|^2 \prod_{k=1, k \neq i}^m (\sigma_i^2(A) - \sigma_k^2(A)) = \prod_{k=1}^{m-1} (\sigma_i^2(A) - \sigma_k^2(\bar{A}_j)), \quad 1 \leq i, j \leq m,$$

$$|v_{ls}|^2 \prod_{t=1, t \neq l}^n (\sigma_l^2(A) - \sigma_t^2(A)) = \prod_{t=1}^{n-1} (\sigma_l^2(A) - \sigma_t^2(\hat{A}_s)), \quad 1 \leq l, s \leq n.$$

We note that the identities in Theorems 1 and 2 are not given in the previous results. The organization of this paper is given as follows. In Section 2, we present the proofs of the two main theorems. The concluding remarks are given in Section 3.

2 The Proof

In this section, we give the proofs of Theorems 1 and 2.

Proof of Theorem 1. Let singular value decompositions of A be $A = U^H \Sigma V$,

where $U \in \mathbf{U}^n, V \in \mathbf{U}^n$. (i) Let $A = \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{pmatrix}$ with $a_1, \dots, a_n \in \mathbf{C}^{1 \times n}$. Since

$$AA^H = U^H \Sigma \Sigma^H U,$$

by Lemma 1 we have

$$|u_{ij}|^2 \prod_{k=1, k \neq i}^n (\lambda_i(AA^H) - \lambda_k(AA^H)) = \prod_{k=1}^{n-1} (\lambda_i(AA^H) - \lambda_k(S_j)), \quad 1 \leq i, j \leq n, \quad (1)$$

where S_j is the $(n-1) \times (n-1)$ submatrix of AA^H that results from deleting the j th column and the j th row, with eigenvalues $\lambda_k(S_j)$. It is easy to see

that

$$S_j = \begin{pmatrix} a_1 a_1^H & \cdots & a_1 a_{j-1}^H & a_1 a_{j+1}^H & \cdots & a_1 a_n^H \\ a_2 a_1^H & \cdots & a_2 a_{j-1}^H & a_2 a_{j+1}^H & \cdots & a_2 a_n^H \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{j-1} a_1^H & \cdots & a_{j-1} a_{j-1}^H & a_{j-1} a_{j+1}^H & \cdots & a_{j-1} a_n^H \\ a_{j+1} a_1^H & \cdots & a_{j+1} a_{j-1}^H & a_{j+1} a_{j+1}^H & \cdots & a_{j+1} a_n^H \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ a_n a_1^H & \cdots & a_n a_{j-1}^H & a_n a_{j+1}^H & \cdots & a_n a_n^H \end{pmatrix}.$$

Let \bar{A}_j be $(n-1) \times n$ matrix after deleting the j -th row of A and $\sigma_k(\bar{A}_j)$ be singular values of \bar{A}_j , then

$$\bar{A}_j = \begin{pmatrix} a_1 \\ \vdots \\ a_{j-1} \\ a_{j+1} \\ \vdots \\ a_n \end{pmatrix}$$

and

$$\sigma_k^2(\bar{A}_j) = \lambda_k^2(\bar{A}_j \bar{A}_j^H). \quad (2)$$

Since A has distinct singular values, by (1) and (2) we have

$$|u_{ij}|^2 = \frac{\prod_{k=1}^{n-1} (\sigma_i^2(A) - \sigma_k^2(\bar{A}_j))}{\prod_{k=1, k \neq i}^n (\sigma_i^2(A) - \sigma_k^2(A))}, \quad 1 \leq i, j \leq n.$$

(ii) Let $A = (b_1, \dots, b_n)$ with $b_1, \dots, b_n \in \mathbf{C}^n$. Since

$$A^H A = V^H \Sigma^H \Sigma V,$$

by Lemma 1 we have

$$|v_{ij}|^2 \prod_{k=1, k \neq i}^n (\lambda_i(A^H A) - \lambda_k(A^H A)) = \prod_{k=1}^{n-1} (\lambda_i(A^H A) - \lambda_k(H_j)), \quad 1 \leq i, j \leq n, \quad (3)$$

where H_j is the $(n-1) \times (n-1)$ submatrix of $A^H A$ that results from deleting the j th column and the j th row, with eigenvalues $\lambda_k(H_j)$. It is easy to see that

$$H_j = \begin{pmatrix} b_1^H b_1 & \cdots & b_1^H b_{j-1} & b_1^H b_{j+1} & \cdots & b_1^H b_n \\ b_2^H b_1 & \cdots & b_2^H b_{j-1} & b_2^H b_{j+1} & \cdots & b_2^H b_n \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ b_{j-1}^H b_1 & \cdots & b_{j-1}^H b_{j-1} & b_{j-1}^H b_{j+1} & \cdots & b_{j-1}^H b_n \\ b_{j+1}^H b_1 & \cdots & b_{j+1}^H b_{j-1} & b_{j+1}^H b_{j+1} & \cdots & b_{j+1}^H b_n \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ b_n^H b_1 & \cdots & b_n^H b_{j-1} & b_n^H b_{j+1} & \cdots & b_n^H b_n \end{pmatrix}.$$

Let \hat{A}_j be $n \times (n-1)$ matrix after deleting the j -th row of A and $\sigma_k(\hat{A}_j)$ be singular values of \hat{A}_j , then

$$\hat{A}_j = (b_1, \dots, b_{j-1}, b_{j+1}, \dots, b_n)$$

and

$$\sigma_k^2(\hat{A}_j) = \lambda_k^2(\hat{A}_j^H \hat{A}_j). \quad (4)$$

Since A has distinct singular values, by (3) and (4) we have

$$|v_{ls}|^2 = \frac{\prod_{t=1}^{n-1} (\sigma_t^2(A) - \sigma_t^2(\hat{A}_s))}{\prod_{t=1, t \neq l}^n (\sigma_t^2(A) - \sigma_t^2(A))}, \quad 1 \leq l, s \leq n.$$

This completes the proof.

By using the similar trick, we can show the results of Theorem 2.

Proof of Theorem 2. Let singular value decompositions of A be $A = U^H \Sigma V$,

where $U \in \mathbf{U}^m, V \in \mathbf{U}^n$. (i) Let $A = \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_m \end{pmatrix}$ with $a_1, \dots, a_m \in \mathbf{C}^{1 \times n}$.

Since

$$AA^H = U^H \Sigma \Sigma^H U,$$

by Lemma 1 we have

$$|u_{ij}|^2 \prod_{k=1, k \neq i}^m (\lambda_i(AA^H) - \lambda_k(AA^H)) = \prod_{k=1}^{m-1} (\lambda_i(AA^H) - \lambda_k(\tilde{S}_j)), \quad 1 \leq i, j \leq m, \quad (5)$$

where \tilde{S}_j is the $(m-1) \times (m-1)$ submatrix of AA^H that results from deleting the j th column and the j th row, with eigenvalues $\lambda_k(\tilde{S}_j)$. It is easy to see that

$$\tilde{S}_j = \begin{pmatrix} a_1 a_1^H & \cdots & a_1 a_{j-1}^H & a_1 a_{j+1}^H & \cdots & a_1 a_m^H \\ a_2 a_1^H & \cdots & a_2 a_{j-1}^H & a_2 a_{j+1}^H & \cdots & a_2 a_m^H \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{j-1} a_1^H & \cdots & a_{j-1} a_{j-1}^H & a_{j-1} a_{j+1}^H & \cdots & a_{j-1} a_m^H \\ a_{j+1} a_1^H & \cdots & a_{j+1} a_{j-1}^H & a_{j+1} a_{j+1}^H & \cdots & a_{j+1} a_m^H \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ a_m a_1^H & \cdots & a_m a_{j-1}^H & a_m a_{j+1}^H & \cdots & a_m a_m^H \end{pmatrix}.$$

Let \tilde{A}_j be $(m-1) \times n$ matrix after deleting the j -th row of A and $\sigma_k(\tilde{A}_j)$ be singular values of \tilde{A}_j , then

$$\tilde{A}_j = \begin{pmatrix} a_1 \\ \vdots \\ a_{j-1} \\ a_{j+1} \\ \vdots \\ a_m \end{pmatrix}$$

and

$$\sigma_k^2(\tilde{A}_j) = \lambda_k^2(\tilde{A}_j \tilde{A}_j^H). \quad (6)$$

By (5) and (6) we have

$$|u_{ij}|^2 \prod_{k=1, k \neq i}^m (\sigma_i^2(A) - \sigma_k^2(A)) = \prod_{k=1}^{m-1} (\sigma_i^2(A) - \sigma_k^2(\tilde{A}_j)), \quad 1 \leq i, j \leq m.$$

(ii) Let $A = (b_1, \dots, b_n)$ with $b_1, \dots, b_n \in \mathbf{C}^m$. Since

$$A^H A = V^H \Sigma^H \Sigma V,$$

by Lemma 1 we have

$$|v_{ij}|^2 \prod_{k=1, k \neq i}^n (\lambda_i(A^H A) - \lambda_k(A^H A)) = \prod_{k=1}^{n-1} (\lambda_i(A^H A) - \lambda_k(H_j)), \quad 1 \leq i, j \leq n, \quad (7)$$

where H_j is the $(n-1) \times (n-1)$ submatrix of $A^H A$ that results from deleting the j th column and the j th row, with eigenvalues $\lambda_k(H_j)$. It is easy to see that

$$H_j = \begin{pmatrix} b_1^H b_1 & \cdots & b_1^H b_{j-1} & b_1^H b_{j+1} & \cdots & b_1^H b_n \\ b_2^H b_1 & \cdots & b_2^H b_{j-1} & b_2^H b_{j+1} & \cdots & b_2^H b_n \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ b_{j-1}^H b_1 & \cdots & b_{j-1}^H b_{j-1} & b_{j-1}^H b_{j+1} & \cdots & b_{j-1}^H b_n \\ b_{j+1}^H b_1 & \cdots & b_{j+1}^H b_{j-1} & b_{j+1}^H b_{j+1} & \cdots & b_{j+1}^H b_n \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ b_n^H b_1 & \cdots & b_n^H b_{j-1} & b_n^H b_{j+1} & \cdots & b_n^H b_n \end{pmatrix}.$$

Let \hat{A}_j be $n \times (n-1)$ matrix after deleting the j -th row of A and $\sigma_k(\hat{A}_j)$ be singular values of \hat{A}_j , then

$$\hat{A}_j = (b_1, \dots, b_{j-1}, b_{j+1}, \dots, b_n)$$

and

$$\sigma_k^2(\hat{A}_j) = \lambda_k^2(\hat{A}_j^H \hat{A}_j). \quad (8)$$

By (7) and (8) we have

$$|v_{ls}|^2 = \frac{\prod_{t=1}^{n-1} (\sigma_t^2(A) - \sigma_t^2(\hat{A}_s))}{\prod_{t=1, t \neq l}^n (\sigma_t^2(A) - \sigma_t^2(A))}, \quad 1 \leq l, s \leq n.$$

This completes the proof.

3 Concluding Remarks

In [1], Denton et al. revisited the identity of eigenvalue-eigenvector for Hermitian matrices. This identity has been discovered by many researchers in the literature. In this paper, we derived similar identity for singular vectors and singular values of general matrices. However, we find these results have not been studied previously. As a future research work, it may be useful to apply this new identity formula to matrix perturbation problems and the calculation of arbitrary singular values and singular vector instead of all singular values and singular vectors.

References

- [1] P. Denton, S. Parke, T. Tao and X. Zhang, Eigenvectors from Eigenvalues: a Survey of a Basic Identity in Linear Algebra, arXiv:1908.03795.
- [2] R. Thompson, Principal Submatrices IX: Interlacing Inequalities for Singular Values of Submatrices, *Linear Algebra and its Applications*, V5 (1972), pp. 1-12.