

## WILLMORE SPHERES IN THE 3-SPHERE REVISITED

SEBASTIAN HELLER

ABSTRACT. Bryant [1] classified all Willmore spheres in 3-space to be given by minimal surfaces in  $\mathbb{R}^3$  with embedded planar ends. This note provides new explicit formulas for genus 0 minimal surfaces in  $\mathbb{R}^3$  with  $2k + 1$  embedded planar ends for all  $k \geq 4$ . Peng and Xiao claimed these examples to exist in [6], but in the same paper they also claimed the existence of a minimal surface with 7 embedded planar ends, which was falsified by Bryant [2].

## 1. SURFACES

Let  $\phi: \hat{\Sigma} \rightarrow S^3$  be a compact, smooth, conformally parametrised, and immersed surface such that for a suitable chosen point  $p \in S^3$ , with its stereographic projection  $\pi_p: S^3 \setminus \{p\} \rightarrow \mathbb{R}^3$ , the composition

$$f = \pi_p \circ \phi: \Sigma = \hat{\Sigma} \setminus \phi^{-1}\{p\} \rightarrow \mathbb{R}^3$$

is a minimal surface in  $\mathbb{R}^3$ . We call such minimal surfaces  $f$  minimal surfaces with embedded planar ends. It was shown by Bryant [1] that all Willmore spheres in the 3-sphere are of this type. Conversely, every  $\phi$  as above is a Willmore sphere. By definition, (immersed) Willmore surfaces  $\phi: \hat{\Sigma} \rightarrow S^3$  are the critical points for the Willmore functional

$$\mathcal{W}(\phi) = \int_{\hat{\Sigma}} (H^2 - K + 1) dA, \quad (1)$$

where 1 is the sectional curvature of the round 3-sphere,  $H$  is the mean curvature,  $dA$  and  $K$  are the area form and the curvature of the induced metric of  $\phi$ . The Willmore energy of a Willmore sphere is given by  $4\pi(n - 1)$ , with  $n$  being the number of ends of  $f = \pi_p \circ \phi$ .

**Remark.** *Peng and Xiao claim the existence of a minimal surface with 7 embedded planar ends in [6] and remark that the existence for  $n = 2k + 1 \geq 9$  follows by a long but straight forward computation. Though the  $n = 7$  case was falsified by Bryant [2], we show that the surfaces predicted in [6] do exist for  $n \geq 9$  by giving a simple and explicit parametrization.*

Let  $X$  be a Riemann surface, and  $g: X \rightarrow \mathbb{K}^r$  be a smooth map, where  $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$ , and  $r \in \mathbb{N}$ . We denote by

$$dg = \partial g + \bar{\partial} g$$

the decomposition of the differential  $dg$  of  $g$  into its complex linear part  $\partial g$  and its complex antilinear part  $\bar{\partial} g$ . For a minimal surface  $f: \Sigma \rightarrow \mathbb{R}^3$  there exists a holomorphic line bundle  $S \rightarrow \Sigma$  with  $S^2 = K_{\Sigma}$  and two holomorphic sections  $s_1, s_2 \in H^0(\Sigma, S)$  such that

$$\partial f = (s_1^2 + s_2^2, i s_1^2 - i s_2^2, -2i s_1 s_2). \quad (2)$$

This is called the Weierstrass representation, and the two spinors  $(s_1, s_2)$  are the Weierstrass data of  $f$ . In the case a minimal surface  $f: \Sigma \rightarrow \mathbb{R}^3$  with embedded planar ends its Weierstrass data are meromorphic spinors on  $\hat{\Sigma}$  with first order poles at  $\phi^{-1}(p)$ , see for example [5] and the references therein. Moreover,  $\partial f$  has no residues at the embedded planar ends.

1.1. **Existing examples in the literature.** Peng and Xiao [6] considered the following ansatz

$$s_1 = \frac{z(z^k - c)}{(z^k - 1)(z^k - \lambda)} \sqrt{dz} \quad \text{and} \quad s_2 = \frac{(z^k - a)(z^k - b)}{z(z^k - 1)(z^k - \lambda)} \sqrt{dz}$$

for the Weierstrass data of a minimal surface of the  $(2k + 1)$ -punctured sphere

$$\Sigma = \mathbb{C}P^1 \setminus \{z \in \mathbb{C} \mid z(z^k - 1)(z^k - \lambda) = 0\},$$

where  $a, b, c, \lambda \in \mathbb{C}$  are pairwise distinct and satisfy the following algebraic condition:

$$0 = \text{res}_p s_1^2 = \text{res}_p s_1 s_2 = \text{res}_p s_2^2 \tag{3}$$

holds at every point  $p \in \{z \in \mathbb{C} \mid z(z^k - 1)(z^k - \lambda) = 0\}$ . Peng and Xiao [6] claim that a solution  $(a, b, c, \lambda)$  always exist implying via (2) the existence of immersed minimal surfaces with  $2k + 1$  ends.

Appropriate solutions of 3 for  $k \in \{4, 5, 6\}$  and explicit formula for the corresponding minimal surface  $f$  can be computed easily. For example, for  $k = 4$ ,

$$a = 10 - 4\sqrt{7} + \sqrt{\frac{635}{3} - 80\sqrt{7}}, \quad b = 10 + 4\sqrt{7} + \sqrt{\frac{635}{3} + 80\sqrt{7}}, \quad c = -3 - 4\sqrt{\frac{3}{5}}, \quad \lambda = -31 - 8\sqrt{15}$$

solve Equation (3). We obtain a minimal surface  $f = \Re(F)$  with 9 embedded planar ends, where

$$F: \mathbb{C}P^1 \setminus \{z \in \mathbb{C} \mid z(z^k - 1)(z^k - \lambda) = 0\} \longrightarrow \mathbb{C}^3$$

is given by

$$F(z) = \frac{1}{45z(-1+z^4)(31+8\sqrt{15}+z^4)} \begin{pmatrix} 6\sqrt{-1}z^2(-15 - 4\sqrt{15} + 5z^4) \\ 5(31 + 8\sqrt{15}) - 3z^4(153 + 40\sqrt{15} + 15z^4) \\ \sqrt{-1}(5(31 + 8\sqrt{15}) - 3z^4(147 + 40\sqrt{15} + 15z^4)) \end{pmatrix}.$$

## 2. CURVES

The following description of genus 0 minimal surfaces with embedded planar ends is due to Bryant [2], see also [3]. Consider a genus 0 minimal surface  $f: \mathbb{C}P^1 \setminus \{p_1, \dots, p_n\} \rightarrow \mathbb{R}^3$  with  $n$  embedded planar ends. As  $\partial f$  has no residues at the ends  $p_1, \dots, p_n$ , the surface  $f$  is the real part of a meromorphic map  $F: \mathbb{C}P^1 \rightarrow \mathbb{C}^3$  with simple poles at  $p_1, \dots, p_n$ . Since  $f$  is conformally parametrised  $F$  is a null curve, i.e., with respect to the standard symmetric inner product  $(\cdot, \cdot)$  on  $\mathbb{C}^3$  we have  $(\partial F, \partial F) = 0$ . Consider  $\mathbb{C}^5$  with the inner product

$$\langle \cdot, \cdot \rangle = -e_0^* \otimes e_4^* - e_4^* \otimes e_0^* + e_1^* \otimes e_1^* + e_2^* \otimes e_2^* + e_3^* \otimes e_3^*,$$

the 3-quadric

$$\mathcal{Q}^3 = P\{v \in \mathbb{C}^5 \setminus \{0\} \mid \langle v, v \rangle = 0\}$$

and the conformal embedding

$$\Psi: \mathbb{C}^3 \rightarrow \mathcal{Q}^3; \quad (z_1, z_2, z_3) \mapsto \left[\frac{1}{2}(z_1^2 + z_2^2 + z_3^2), z_1, z_2, z_3, 1\right].$$

For a minimal sphere  $f = \Re(F)$  with  $n$  embedded planar ends,  $\Psi \circ F: \mathbb{C}P^1 \rightarrow \mathcal{Q}^3$  is an unbranched rational curve of degree  $n$ . Moreover,  $\Psi \circ F$  is again a null curve, i.e., for every local holomorphic lift  $\hat{\Psi}$  of  $\Psi \circ F$  the condition  $\langle \hat{\Psi}, \hat{\Psi} \rangle = 0 = \langle \partial \hat{\Psi}, \partial \hat{\Psi} \rangle$  holds. Conversely, every (nondegenerate) unbranched null curve gives rise to a minimal surface with embedded planar ends by reversing the above construction.

Let  $V = \mathbb{C}^4$  be equipped with the 2-form  $\Omega = dx_1 \wedge dx_2 + dx_3 \wedge dx_4$ . Consider the 5-dimensional space  $W = \{\eta \in \Lambda^2 V \mid \Omega(\eta) = 0\}$  equipped with the non-degenerated symmetric inner product  $\frac{1}{2}\Omega \wedge \Omega$ , and the corresponding 3-quadric  $\mathcal{Q}_\Omega$  of null lines in  $PW$ . Identifying  $(W, \frac{1}{2}\Omega \wedge \Omega) \cong (\mathbb{C}^5, \langle \cdot, \cdot \rangle)$  yields  $\mathcal{Q}_\Omega \cong \mathcal{Q}^3$ . A holomorphic curve  $\psi: \hat{\Sigma} \rightarrow \mathbb{C}P^3$  is a contact curve if  $\Omega(\hat{\psi} \wedge \partial \hat{\psi}) = 0$  holds for every local holomorphic lift  $\hat{\psi}$  of  $\psi$ . Then, with respect to a local holomorphic coordinate  $z$ , the map  $z \mapsto \hat{\psi} \wedge \frac{\partial \hat{\psi}}{\partial z}$  gives a local lift of a well-defined holomorphic curve

$$\psi^2: \hat{\Sigma} \rightarrow \mathcal{Q}_\Omega \subset PW,$$

the second associated curve of  $\psi$ . It is a null curve with respect to  $\frac{1}{2}\Omega \wedge \Omega$ . A curve into a projective space is called nondegenerate if it is not contained in any hyperplane. The Klein correspondence (see [3]) states

that every nondegenerate null curve is given by a nondegenerate contact curve in  $\mathbb{C}P^3$ . For  $\hat{\Sigma} = \mathbb{C}P^1$  and  $\psi$  of degree  $d$  its second associated curve  $\psi^2$  is unbranched if and only if the total branch order of  $\psi$  is  $d - 3$ . This is a direct consequence of the Plücker relations applied to the duality between  $\psi$  and its third associated curve, see [3]. Hence, to construct genus 0 minimal surfaces with  $2k + 1$  embedded planar ends, we have to construct rational contact curves of degree  $2k$  with total branch order  $2k - 3$ .

**2.1. Rational contact curves of degree  $2k$  with total branch order  $2k - 3$ .** For  $k \in \mathbb{N} \setminus \{3\}$  consider the map  $\psi: \mathbb{C}P^1 \rightarrow \mathbb{C}P^3$  defined via the lift

$$\hat{\psi}(z) = \begin{pmatrix} z^3 \\ \frac{1}{6}(-6 + 13k - 9k^2 + 2k^3 + \frac{12(-3+2k)}{-3+k}z^k + 3k(-3 + 2k)z^k - 6z^{2k}) \\ \frac{1}{2}(-3 + 2k)z(-2 + k + 2z^k) \\ z^2(-1 + k + z^k) \end{pmatrix}. \quad (4)$$

This is a nondegenerate rational curve of degree  $2k$  if  $k \in \mathbb{N}^{>3}$ . It can be directly verified that it is a contact curve, i.e.,  $\Omega(\hat{\psi} \wedge \frac{\partial \hat{\psi}}{\partial z}) = 0$ . For  $k \in \mathbb{N}^{>3}$  its branch points are at the  $k$ -th roots of unity and at  $z = \infty$ . The branch order at the roots of unity is 1, and at  $z = \infty$  the branch order is  $k - 3$ . Hence, the total branch order is  $2k - 3$ . The construction fails for  $k \leq 3$ : for  $k = 1$ , the curve is of degree 2 and branched at  $z = 0$  and  $z = 1$ . Thus, its second associate curve cannot be unbranched. For  $k = 2$ , the degree of the curve is  $2 \neq 4$ . For  $k = 3$  and  $k = 0$ , the formula (4) gives a point in  $\mathbb{C}P^3$ . For  $k = 4$  we obtain the first valid example

$$\hat{\psi}(z) = (z^3, 5 + 20z^4 - z^8, 5z + 5z^5, 3z^2 + z^6).$$

Together with the examples of unbranched null curves of even degree  $d \geq 4$  and the nonexistence results of Bryant in [2], this shows:

**Theorem.** *There exist a Willmore sphere  $\phi_n: \mathbb{C}P^1 \rightarrow S^3$  with Willmore energy  $4\pi(n - 1)$  if and only if  $n \in \mathbb{N} \setminus \{2, 3, 5, 7\}$ .*

For completeness, we state the formula for the genus 0 minimal surfaces  $f = \Re(F)$  with  $(2k + 1)$  embedded planar ends corresponding to the contact curve (4). Note that these surfaces are only determined up to Goursat transformations [4], see also [2]. We leave it as an exercise for the interested reader to verify that we have reobtained the surfaces of Peng and Xiao [6] up to a Goursat transformation and reparametrisation. The meromorphic map  $F: \mathbb{C}P^1 \rightarrow \mathbb{C}^3$  is given by

$$F(z) = \begin{pmatrix} \frac{-12\sqrt{-1}(-3+k)(-3+2k)z^2(2-3k+k^2-2z^k)}{4z((-3+k)(-1+k)^2(-3+2k)-6(-1+k)(-3+2k)z^k-3(-3+k)z^{2k})} \\ \frac{(-1+k)(-12(3-2k)^2(-2+k)z^k-12(-3+k)(-3+2k)z^{2k}+(-3+k)((6-7k+2k^2)^2-12z^4))}{4z((-3+k)(-1+k)^2(-3+2k)-6(-1+k)(-3+2k)z^k-3(-3+k)z^{2k})} \\ \frac{\sqrt{-1}(-1+k)(12(3-2k)^2(-2+k)z^k+12(-3+k)(-3+2k)z^{2k}+(-3+k)((6-7k+2k^2)^2-12z^4))}{4z((-3+k)(-1+k)^2(-3+2k)-6(-1+k)(-3+2k)z^k-3(-3+k)z^{2k})} \end{pmatrix}.$$

## REFERENCES

- [1] R. L. Bryant, A duality theorem for Willmore surfaces, *J. Diff. Geo.* 20 (1984), no. 1, 23–53.
- [2] R. L. Bryant, Surfaces in conformal geometry, *Proc. of symposia in pure math.*, 48 (1988), 227–240.
- [3] R. L. Bryant, Notes on projective, contact, and null curves, *arxiv:1905.06117*.
- [4] E. Goursat, Sur un mode de transformation des surfaces minima. *Acta Math.*, Vol 11, 135–186, 1887.
- [5] R. Kusner and N. Schmitt, The Spinor Representation of Minimal Surfaces, *arxiv.org/dg-ga/9512003*.
- [6] C.K. Peng and L. Xiao, Willmore surfaces and minimal surfaces with flat ends, *Geometry and Topology of Submanifolds X*, 259–265 (2000).

INSTITUTE OF DIFFERENTIAL GEOMETRY, LEIBNIZ UNIVERSITÄT HANNOVER, WELFENGARTEN 1, 30167 HANNOVER

*E-mail address:* seb.heller@gmail.com