# Modular and fractional *L*-intersecting families of vector spaces

Rogers Mathew <sup>\*1</sup>, Tapas Kumar Mishra<sup>2</sup>, Ritabrata Ray<sup>3</sup>, and Shashank Srivastava<sup>4</sup>

<sup>1</sup>Department of Computer Science and Engineering, Indian Institute of Technology Hyderabad, India.

rogers@iith.ac.in

<sup>2</sup>Department of Computer Science and Engineering, National Institute of Technology Rourkela, India.

mishrat@nitrkl.ac.in

<sup>3</sup>Department of Electrical & Computer Engineering, Cornell University, Ithaka, NY 14853, USA.

rayritabrata96@gmail.com

<sup>4</sup> Toyota Technological Institute at Chicago, Chicago, IL 60637, USA. shashanks@ttic.edu

#### Abstract

In the first part of this paper, we prove the following theorem which is the q-analogue of a generalized modular Ray-Chaudhuri-Wilson Theorem shown in [Alon, Babai, Suzuki, J. Combin. Theory Series A, 1991]. It is also a generalization of the main theorem in [Frankl and Graham, European J. Combin. 1985] under certain circumstances.

• Let V be a vector space of dimension n over a finite field of size q. Let  $K = \{k_1, \ldots, k_r\}, L = \{\mu_1, \ldots, \mu_s\}$  be two disjoint subsets of  $\{0, 1, \ldots, b-1\}$  with  $k_1 < \cdots < k_r$ . Let  $\mathcal{F} = \{V_1, V_2, \ldots, V_m\}$  be a family of subspaces of V such that  $\forall i \in [m], \dim(V_i) \equiv k_t \pmod{b}$ , for some  $k_t \in K$ ; for every distinct  $i, j \in [m], \dim(V_i \cap V_j) \equiv \mu_t \pmod{b}$ , for some  $\mu_t \in L$ . Moreover, it is given that neither of the following two conditions hold:

(i) q + 1 is a power of 2, and b = 2

(ii) q = 2, b = 6

Then,

$$\begin{aligned} |\mathcal{F}| &\leq \begin{cases} N(n,s,r,q), & \text{if } (s+k_r \leq n \text{ and } r(s-r+1) \leq b-1) \text{ or } (s < k_1+r) \\ N(n,s,r,q) + \sum_{t \in [r]} {n \brack k_t}_q, & \text{otherwise} \end{cases} \\ \text{, where } N(n,s,r,q) &:= {n \brack s}_q + {n \brack s-1}_q + \dots + {n \brack s-r+1}_q. \end{aligned}$$

In the second part of this paper, we prove q-analogues of results on a recent notion called *fractional L-intersecting family* of sets for families of subspaces of a given vector space over a finite field of size q. We use the above theorem to obtain a general upper bound to the cardinality of such families. We give an improvement to this general upper bound in certain special cases.

<sup>\*</sup>This author was supported by a grant from the Science and Engineering Research Board, Department of Science and Technology, Govt. of India (project number: MTR/2019/000550).

## 1 Introduction

Let [n] be the set of all natural numbers from 1 to n. A family  $\mathcal{F}$  of subsets of [n] is called intersecting if every set in  $\mathcal{F}$  has a non-empty intersection with every other set in  $\mathcal{F}$ . One of the earliest studies on intersecting families dates back to the famous Erdős-Ko-Rado Theorem [Erdos et al., 1961] about maximal uniform intersecting families. Ray-Chaudhuri and Wilson [Ray-Chaudhuri and Wilson, 1975] introduced the notion of *L*-intersecting families. Let  $L = \{l_1, \ldots, l_s\}$  be a set of non-negative integers. A family  $\mathcal{F}$  of subsets of [n] is said to be *L*intersecting if for every distinct  $F_i, F_j$  in  $\mathcal{F}, |F_i \cap F_j| \in L$ . The Ray-Chaudhuri-Wilson Theorem states that if  $\mathcal{F}$  is *t*-uniform (that is, every set in  $\mathcal{F}$  is *t*-sized), then  $|\mathcal{F}| \leq {n \choose s}$ . This bound is tight as shown by the set of all *s*-sized subsets of [n] with  $L = \{0, \ldots, s-1\}$ . Frankl-Wilson Theorem

[Frankl and Wilson, 1981a] extends this to non-uniform families by showing that  $|\mathcal{F}| \leq \sum_{i=0}^{n} {n \choose i}$ ,

where  $\mathcal{F}$  is any family of subsets of [n] that is *L*-intersecting. The collection of all the subsets of [n] of size at most *s* with  $L = \{0, \ldots s - 1\}$  is a tight example to this bound. The first proofs of these theorems were based on the technique of higher incidence matrices. Alon, Babai, and Suzuki in [Alon et al., 1991] generalized the Frankl-Wilson Theorem using a proof that operated on spaces of multilinear polynomials. They showed that if the sizes of the sets in  $\mathcal{F}$  belong to  $K = \{k_1, \ldots, k_r\}$  with each  $k_i > s - r$ , then  $|\mathcal{F}| \leq {n \choose s} + \cdots + {n \choose s - r + 1}$ . A modular version of the Ray-Chaudhuri-Wilson Theorem was shown in [Frankl and Wilson, 1981b]. This result was generalized in [Alon et al., 1991]. See [Liu and Yang, 2014] for a survey on *L*-intersecting families.

Researchers have also been working on similar intersection theorems for subspaces of a given vector space over a finite field. Hsieh [Hsieh, 1975], and Deza and Frankl [Deza and Frankl, 1983] showed Erdős-Ko-Rado type theorems for subspaces. Let V be a vector space of dimension n over a finite field of size q. The number of d-dimensional subspaces of V is given by the q-binomial coefficient  $\begin{bmatrix} n \\ d \end{bmatrix}_q = \frac{(q^n-1)(q^{n-1}-1)\cdots(q^{n-d+1}-1)}{(q^{d-1}-1)\cdots(q^{-1})}$ . The following theorem which is a q-analog of the Ray-Chaudhuri-Wilson Theorem by considering families of subspaces instead of subsets is due to

[Frankl and Graham, 1985].

**Theorem 1.1.** [Theorem 1.1 in [Frankl and Graham, 1985]] Let V be a vector space over of dimension n over a finite field of size q. Let  $\mathcal{F} = \{V_1, V_2, \ldots, V_m\}$  be a family of subspaces of V such that  $\dim(V_i) = k, \forall i \in [m]$ . Let  $0 \le \mu_1 < \mu_2 < \cdots < \mu_s < b$  be integers such that  $k \not\equiv \mu_t \pmod{b}$ , for any t. For every  $1 \le i < j \le m$ ,  $\dim(V_i \cap V_j) \equiv \mu_t \pmod{b}$ , for some t. Then,

$$|\mathcal{F}| \le \begin{bmatrix} n \\ s \end{bmatrix}_q$$

except possibly for  $q = 2, b = 6, s \in \{3, 4\}$ .

**Example 1** (Remark 3.2 in [Frankl and Graham, 1985]). Let n = k + s. Let  $\mathcal{F}$  be the family of all the k-dimensional subspaces of V, where V is an n-dimensional vector space over a finite field of size q. Observe that, for any two distinct  $V_i, V_j \in \mathcal{F}, k - s \leq \dim(V_i \cap V_j) \leq k - 1$ . This is a tight example for Theorem 1.1.

Alon et al. in [Alon et al., 1991] proved a generalization of the non-modular version of the above theorem. This result was subsequently strengthened in [Liu et al., 2018].

Our paper is divided into two parts. In Part A of the paper, we prove the following theorem which is a generalization of Theorem 1.1 due to Frankl and Graham under certain circumstances.

It is also the q-analogue of a generalized modular Ray-Chaudhuri-Wilson Theorem shown in [Alon et al., 1991]. We assume that  $\begin{bmatrix} a \\ b \end{bmatrix}_q = 0$ , when b < 0 or b > a. Let

$$N(n,s,r,q) := \begin{bmatrix} n \\ s \end{bmatrix}_q + \begin{bmatrix} n \\ s-1 \end{bmatrix}_q + \dots + \begin{bmatrix} n \\ s-r+1 \end{bmatrix}_q$$

**Theorem 1.2.** Let V be a vector space of dimension n over a finite field of size q. Let  $K = \{k_1, \ldots, k_r\}, L = \{\mu_1, \ldots, \mu_s\}$  be two disjoint subsets of  $\{0, 1, \ldots, b-1\}$  with  $k_1 < \cdots < k_r$ . Let  $\mathcal{F} = \{V_1, V_2, \ldots, V_m\}$  be a family of subspaces of V such that  $\forall i \in [m], \dim(V_i) \equiv k_t \pmod{b}$ , for some  $k_t \in K$ ; for every distinct  $i, j \in [m], \dim(V_i \cap V_j) \equiv \mu_t \pmod{b}$ , for some  $\mu_t \in L$ . Moreover, it is given that neither of the following two conditions hold:

- (i) q + 1 is a power of 2, and b = 2
- (ii) q = 2, b = 6

Then,

$$|\mathcal{F}| \leq \begin{cases} N(n, s, r, q), & \text{if } (s + k_r \leq n \text{ and } r(s - r + 1) \leq b - 1) \text{ or } (s < k_1 + r) \\ N(n, s, r, q) + \sum_{t \in [r]} {n \brack k_t}_q, & \text{otherwise.} \end{cases}$$

In Part B, we study a notion of fractional *L*-intersecting families which was introduced in [Balachandran et al., 2019]. We say a family  $\mathcal{F} = \{F_1, F_2, \ldots, F_m\}$  of subsets of [n] is a *fractional L*-intersecting family, where *L* is a set of irreducible fractions between 0 and 1, if  $\forall i, j \in [m], i \neq j$ ,  $\frac{|F_i \cap F_j|}{|F_i|} \in L$  or  $\frac{|F_i \cap F_j|}{|F_j|} \in L$ . In this paper, we extend this notion from subsets to subspaces of a vector space over a finite field.

**Definition 1.3.** Let  $L = \{\frac{a_1}{b_1}, \dots, \frac{a_s}{b_s}\}$  be a set of positive irreducible fractions, where every  $\frac{a_i}{b_i} < 1$ . Let  $\mathcal{F} = \{V_1, \dots, V_m\}$  be a family of subspaces of a vector space V over a finite field. We say  $\mathcal{F}$  is a fractional L-intersecting family of subspaces if for every two distinct  $i, j \in [m]$ ,  $\frac{\dim(V_i \cap V_j)}{\dim(V_i)} \in L$  or  $\frac{\dim(V_i \cap V_j)}{\dim(V_j)} \in L$ .

When every subspace in  $\mathcal{F}$  is of dimension exactly k, it is an L'-intersecting family where  $L' = \{\frac{a_1k}{b_1}, \ldots, \frac{a_sk}{b_s}\}$ . Applying Theorem 1.1, we get  $|\mathcal{F}| \leq \begin{bmatrix}n\\s\end{bmatrix}_q$ . A tight example to this is the collection of all k-dimensional subspaces of V with  $L = \{\frac{0}{k}, \ldots, \frac{k-1}{k}\}$ . However, the problem of bounding the cardinality of a fractional L-intersecting family of subspaces becomes more interesting when  $\mathcal{F}$  contains subspaces of various dimensions. In Part B of this paper, we obtain upper bounds for the cardinality of a fractional L-intersecting family of subspaces that are q-analogs of the results in [Balachandran et al., 2019]. With the help of Theorem 1.2 that we prove in Part A, we obtain the following result in Part B.

**Theorem 1.4.** Let  $L = \{\frac{a_1}{b_1}, \frac{a_2}{b_2}, \dots, \frac{a_s}{b_s}\}$  be a collection of positive irreducible fractions, where every  $\frac{a_i}{b_i} < 1$ . Let  $\mathcal{F}$  be a fractional L-intersecting family of subspaces of a vector space V of dimension n over a finite field of size q. Let  $t = \max_{i \in [s]} b_i$ ,  $g(t, n) = \frac{2(2t+\ln n)}{\ln(2t+\ln n)}$ , and  $h(t, n) = \min(g(t, n), \frac{\ln n}{\ln t})$ . Then,

$$|\mathcal{F}| \le 2g(t,n)h(t,n)\ln(g(t,n)) \begin{bmatrix} n\\ s \end{bmatrix}_q + h(t,n)\sum_{i=1}^{s-1} \begin{bmatrix} n\\ i \end{bmatrix}_q.$$

Further, if  $2g(t, n) \ln(g(t, n)) \le n + 2$ , then

$$|\mathcal{F}| \le 2g(t,n)h(t,n)\ln(g(t,n)) \begin{bmatrix} n\\s \end{bmatrix}_q.$$

**Example 2.** Let s be a constant,  $L = \{\frac{0}{s}, \frac{1}{s}, \dots, \frac{s-1}{s}\}$ , and  $\mathcal{F}$  be the family of all the s-sized subspaces of V. Clearly,  $\mathcal{F}$  is a fractional L-intersecting family showing that the bound in Theorem 1.4 is asymptotically tight up to a multiplicative factor of  $\frac{\ln^2 n}{\ln \ln n}$ .

We improve the bound obtained in Theorem 1.4 for the special case when  $L = \{\frac{a}{b}\}$ , where b is a prime.

**Theorem 1.5.** Let  $L = \{\frac{a}{b}\}$ , where  $\frac{a}{b}$  is a positive irreducible fraction less than 1 and b is a prime. Let  $\mathcal{F}$  be a fractional *L*-intersecting family of subspaces of a vector space V of dimension n over a finite field of size q. Then, we have  $|\mathcal{F}| \leq (b-1)({n \brack 1}_q + 1) \lceil \frac{\ln n}{\ln b} \rceil + 2$ .

**Example 3.** Let  $L = \{\frac{1}{2}\}$ . Let V be a vector space of dimension n over a finite field of size q. Let  $\{v_1, v_2, \ldots, v_n\}$  be a basis of V. Let  $V' := span(\{v_2, \ldots, v_n\})$  be an (n-1)-dimensional subspace of V. Let  $\mathcal{F}$  be the set of all  $\begin{bmatrix} n-1\\1 \end{bmatrix}_q$  2-dimensional subspaces of V each of which is obtained by a span of  $v_1$  and each of the  $\begin{bmatrix} n-1\\1 \end{bmatrix}_q$  1-dimensional subspaces of V'. This example

shows that when b and q are constants, the bound in Theorem 1.5 is asymptotically tight up to a multiplicative factor of  $\ln n$ .

## PART A: Generalized modular RW Theorem for subspaces

As mentioned before, in this part we prove Theorem 1.2. The approach followed here is similar to the approach used in proving Theorem 1.5, a generalized modular Ray-Chaudhuri-Wilson Theorem for subsets, in [Alon et al., 1991]. We start by stating the Zsigmondy's Theorem which will be used in the proof of Theorem 1.2.

**Theorem 1.6** ([Zsigmondy, 1892]). For any  $q, b \in \mathbb{N}$ , there exists a prime p such that  $q^b \equiv 1 \pmod{p}$ ,  $q^i \not\equiv 1 \pmod{p} \quad \forall i, \ 0 < i < b$ , except when (i) q + 1 is a power of 2, b = 2, or (ii) q = 2, b = 6.

#### 1.1 Notations used in PART A

Unless defined explicitly, in the rest of Part A, the symbols  $K = \{k_1, \ldots, k_r\}$ ,  $r, L = \{\mu_1, \ldots, \mu_s\}$ ,  $s, q, V, \mathcal{F}, n, b, m, \text{ and } V_1, \ldots, V_m$  are defined as they are defined in Theorem 1.2. We shall use  $U \subseteq V$  to denote that U is a subspace of V. Using Zsigmondy's Theorem, we find a prime p so that  $q^i \not\equiv 1 \pmod{p}$  for 0 < i < b and  $q^b \equiv 1 \pmod{p}$ . This is possible except in the two cases specified in Theorem 1.6. We ignore these two cases from now on in the rest of Part A.

#### 1.2 Möbius inversion over the subspace poset

Consider the partial order defined on the set of subspaces of the vector space V over a finite field of size q under the 'containment' relation. Let  $\alpha$  be a function from the set of subspaces of V to  $\mathbb{F}_p$ . A function  $\beta$  from the set of subspaces of V to  $\mathbb{F}_p$  is the zeta transform of  $\alpha$  if  $\forall W \subseteq V, \ \beta(W) = \sum_{U \subseteq W} \alpha(U)$ . Then, applying the *Möbius inversion formula* we get  $\forall W \subseteq V$ ,

 $\alpha(W) = \sum_{U \subseteq W} \mu(U, W) \beta(U)$ , where  $\alpha$  is called the *Möbius transform* of  $\beta$  and  $\mu(U, W)$  is the *Möbius function* for the subspace poset. In the proposition below, we show that the *Möbius function* for the subspace poset is defined as

$$\mu(X,Y) = \begin{cases} {}^{(-1)^d q^{\binom{d}{2}}}, \text{ if } X \subseteq Y \\ 0, & \text{otherwise} \end{cases}$$

,  $\forall X, Y \subseteq V$  with d = dim(Y) - dim(X).

**Proposition 1.7.** Let  $\alpha$  and  $\beta$  be functions from the set of subspaces of V to  $\mathbb{F}_p$ . Then,  $\forall W \subseteq V$ ,

$$\beta(W) = \sum_{U \subseteq W} \alpha(U) \iff \alpha(W) = \sum_{\substack{U \subseteq W \\ d = \dim(W) - \dim(U)}} (-1)^d q^{\frac{d(d-1)}{2}} \beta(U)$$

Proof. Proof in Appendix A.

**Definition 1.8.** Given two subspaces U and W of the vector space V, we define their union space  $U \cup W$  as the span of union of sets of vectors in U and W.

**Proposition 1.9.** Let  $\alpha$  and  $\beta$  be functions as defined in Proposition 1.7. Then,  $\forall W, Y$  such that  $W \subseteq Y \subseteq V$ ,

$$\sum_{\substack{T: \ W \subseteq T \subseteq Y \\ d = \dim(Y) - \dim(T)}} (-1)^d q^{\frac{d(d-1)}{2}} \beta(T) = \sum_{U: \ U \cup W = Y} \alpha(U)$$

Proof. Proof in Appendix B

**Corollary 1.10.** For any non-negative integer g, the following are equivalent for functions  $\alpha$  and  $\beta$  defined in Proposition 1.7:

(i)  $\alpha(U) = 0, \forall U \subseteq V \text{ with } dim(U) \ge g.$ 

(ii) 
$$\sum_{\substack{W \subseteq T \subseteq Y \\ d = \dim(Y) - \dim(T)}} (-1)^d q^{\frac{d(d-1)}{2}} \beta(T) = 0, \forall W, Y \subseteq V \text{ with } \dim(Y) - \dim(W) \ge g$$

**Definition 1.11.** Let  $H = \{h_1, h_2, \ldots, h_t\}$  be a subset of  $\{0, 1, \ldots, n\}$  where  $h_1 < h_2 < \cdots < h_t$ . We say H has a gap of size  $\geq g$  if either  $h_1 \geq g - 1$ ,  $n - h_t \geq g - 1$ , or  $h_{i+1} - h_i \geq g$  for some  $i \in [t-1]$ .

**Lemma 1.12.** Let  $\alpha$  and  $\beta$  be functions as in Proposition 1.7. Let  $H \subseteq \{0, 1, ..., n\}$  be a set of integers and g an integer,  $0 \le g \le n$ . Moreover, we have the following conditions:

- (i)  $\forall U \subseteq V$ , we have  $\alpha(U) = 0$  whenever  $dim(U) \ge g$ .
- (ii)  $\forall T \subseteq V$ , we have  $\beta(T) = 0$  whenever  $dim(T) \notin H$ .
- (iii) H has a gap  $\geq g + 1$ .

Then,  $\alpha = \beta = 0$ .

*Proof.* Let  $H = \{h_1, h_2, \ldots, h_{|H|}\}$ . Suppose the gap is between  $h_i$  and  $h_{i-1}$ , or between -1 and  $h_1$ , then we may assume  $\exists i$  such that  $h_i \in H$  and  $h_i - j \notin H$  for  $1 \leq j \leq g$  and  $h_i - g \geq 0$ . Choose any two subspaces, say U and W, of V of dimensions  $h_i$  and  $h_i - g$ , respectively. Since  $\dim(U) \geq g, \alpha(U) = 0$ . We know from Corollary 1.10 that

$$\sum_{\substack{W \subseteq T \subseteq U\\ cdim(U) - dim(T)}} (-1)^d q^{\frac{d(d-1)}{2}} \beta(T) = 0$$

But whenever  $dim(T) < h_i$ , it lies between  $h_i - g$  and  $h_i - 1$ , and hence  $\beta(T) = 0$ . Then,

$$\sum_{\substack{W \subseteq T \subseteq U\\ d = dim(\overline{U}) - dim(T)}} (-1)^d q^{\frac{d(d-1)}{2}} \beta(T) = \beta(U) = 0$$

Since our choice of U was arbitrary, we may conclude that  $\beta(U) = 0, \forall U \subseteq V$  with  $dim(U) = h_i$ . Thus, we can remove  $h_i$  from the set H, and then use the same procedure to further reduce the size of H till it is an empty set. If H is empty,  $\beta(U) = 0, \forall U \subseteq V$ , giving  $\alpha(U) = \beta(U) = 0$  as required.

Now suppose the gap was between  $h_{|H|}$  and n + 1. In this case, we take U of dimension  $h_{|H|}$  and W of dimension  $h_{|H|} + g$  to show that  $\beta(U) = 0$ , and remove  $h_{|H|}$  from H. Note that removing a number from the set H can never reduce the gap.

## **1.3** Defining functions $f^{x,y}$ and $g^{x,y}$

d=

Consider all the subspaces of the vector space V. We can impose an ordering on the subspaces of same dimension, and use the natural ordering across dimensions, so that every subspace can be uniquely represented by a pair of integers  $\langle d, e \rangle$ , indicating that it is the  $e^{th}$  subspace of dimension d,  $0 \leq d \leq n$ ,  $1 \leq e \leq \begin{bmatrix} n \\ d \end{bmatrix}_q$ . Let us call that subspace  $V_{d,e}$ . Let S be the number of subspaces of V of dimension at most s, that is,  $S = \sum_{t=0}^{s} \begin{bmatrix} n \\ t \end{bmatrix}_q$ . Let each subspace  $V_{d,e}$  of dimension at most s be represented as a 0-1 containment vector  $v_{d,e}$  of S entries, each entry of the vector denoting whether a particular subspace of dimension  $\leq s$  is contained in  $V_{d,e}$  or not.

$$v_{d,e}^{x,y} = \begin{cases} 1, \text{ if } V_{x,y} \text{ is a subspace of } V_{d,e} \\ 0, \text{ otherwise} \end{cases}$$

The vector  $v_{d,e}$  is made up of  $v_{d,e}^{x,y}$  values for  $0 \le x \le s, 1 \le y \le {n \brack x_{q}}$ , making it a vector of size S. For  $0 \le x \le s, 1 \le y \le {n \brack x_{q}}$  we define functions  $f^{x,y} : \mathbb{F}_{2}^{S} \to \mathbb{F}_{p}$  as

$$f^{x,y}(v) = f^{x,y}(v^{0,1}, v^{1,1}, \dots, v^{1, {n \brack 1}_q}, \dots, v^{s,1}, \dots, v^{s, {n \brack s}_q}) := v^{x,y}.$$

For  $0 \le x \le s - r, 1 \le y \le {n \brack x_q}$ , we define functions  $g^{x,y} : \mathbb{F}_2^S \to \mathbb{F}_p$  as

$$g^{x,y}(v) = f^{x,y}(v) \prod_{t \in [r]} (\sum_{j=1}^{\binom{n}{1}_q} v^{1,j} - \binom{k_t}{1}_q)$$

Let  $\Omega$  denote  $\mathbb{F}_2^S$ . The functions  $f^{x,y}$  and  $g^{x,y}$  reside in the space  $\mathbb{F}_p^{\Omega}$ .

#### **1.4** Swallowing trick: linear independence of functions $f^{x,y}$ and $g^{x,y}$

**Lemma 1.13.** Let  $s + k_r \leq n$  and  $r(s - r + 1) \leq b - 1$ . The functions  $g^{x,y}$ ,  $0 \leq x \leq s - r$ ,  $1 \leq y \leq {n \brack x_q}$ , are linearly independent in the function space  $\mathbb{F}_p^{\Omega}$  over  $\mathbb{F}_p$ .

*Proof.* We wish to show that the only solution to  $\sum_{\substack{0 \le x \le s-r \\ 1 \le y \le \binom{n}{x}_g}} \alpha^{x,y} g^{x,y} = 0$  is the trivial solution

 $\alpha^{x,y} = 0, \ \forall x, y.$  We define function  $\alpha$  from the set of all subspaces of V to  $\mathbb{F}_p$  as:

$$\alpha(V_{d,e}) = \begin{cases} \alpha^{d,e}, \text{ if } 0 \le d \le s - r \\ 0, \text{ if } d > s - r \end{cases}$$

We show that functions  $\alpha$  and  $\beta(U) := \sum_{T \subseteq U} \alpha(T)$  satisfy the conditions of Lemma 1.12, thereby implying  $\alpha(U) = 0$ ,  $\forall U \subseteq V$ , including  $\alpha(V_{d,e}) = \alpha^{d,e} = 0$  for  $0 \leq d \leq s - r$ , which will in turn imply that the functions  $g^{x,y}$  above are linearly independent.

Let  $H = \{x : 0 \le x \le n, x \equiv k_t \pmod{b}, t \in [r]\}$ . We claim that H has a gap of size at least s - r + 2. Suppose  $n \ge b + k_1$ . Then, since  $r(s - r + 1) \le b - 1$ , the gap is between  $k_1$  and  $b + k_1$ . Suppose  $(s + k_r \le) n < b + k_1$ . Then, the gap is right above  $k_r$ . This proves the claim. We now need to show that for  $T \subseteq V$ ,  $\beta(T) = 0$  whenever  $\dim(T) \notin H$ , or whenever  $\dim(T) \not\equiv k_t \pmod{b}$ , for any  $t \in [r]$ . Suppose  $v_T$  is the S-sized containment vector for T. When  $\dim(T) \not\equiv k_t \pmod{b}$  for any  $t \in [r]$ , it follows from the property of the prime p given by Theorem 1.6 that  $\sum_{1 \le j \le [n] \atop 1} v_T^{1,j} - [k_t \atop 1]_q \neq 0$  in  $\mathbb{F}_p$ , for every  $t \in [r]$ .

$$\beta(T) = \sum_{U \subseteq T} \alpha(U) = \sum_{\substack{\dim(U) \leq s-r \\ U \subseteq T}} \alpha(U) = \sum_{\substack{0 \leq d \leq s-r \\ 1 \leq e \leq \lfloor n \\ d \rfloor_{c}}} \alpha(V^{d,e}) f^{d,e}(v_T)$$

Since  $\sum_{1 \le j \le \begin{bmatrix} n \\ 1 \end{bmatrix}_q} v_T^{1,j} - \begin{bmatrix} k_t \\ 1 \end{bmatrix}_q \ne 0$  in  $\mathbb{F}_p$  for every  $t \in [r], f^{d,e}(v_T) = c(T)g^{d,e}(v_T)$  where  $c(T) \ne 0$ .

Then,

$$\beta(T) = c(T) \sum_{\substack{0 \le d \le s - r \\ 1 \le e \le {n \brack d_q}}} \alpha(V^{d,e}) g^{d,e}(v_T) = c(T) \sum_{\substack{0 \le d \le s - r \\ 1 \le e \le {n \brack d_q}}} \alpha^{d,e} g^{d,e}(v_T) = c(T) \cdot 0 = 0.$$

Since the set H and the functions  $\alpha$  and  $\beta$  satisfy the conditions of Lemma 1.12, we have  $\alpha = 0$ . This proves the lemma.

Recall that we are given a family  $\mathcal{F} = \{V_1, V_2, \dots, V_m\}$  of subspaces of V such that  $\forall i \in [m]$ , dim $(V_i) \equiv k_t \pmod{b}$ , for some  $k_t \in K$ . Further, dim $(V_i \cap V_j) \equiv \mu_t \pmod{b}$ , for some  $\mu_t \in L$ and K and L are disjoint subsets of  $\{0, 1, \dots, b-1\}$ . Let  $v_i$  be the containment vector of size S corresponding to subspace  $V_i \in \mathcal{F}$ . We define the following functions from  $\mathbb{F}_2^S \to \mathbb{F}_p$ .

$$g^{i}(v) = g^{i}(v^{0,1}, v^{1,1}, \dots, v^{1, {n \brack 1_{q}}}, \dots, v^{s,1}, \dots, v^{s, {n \choose s_{q}}})$$
$$:= \prod_{j=1}^{s} \left( \sum_{1 \le y \le {n \brack 1_{q}}} \left( v^{1,y}_{i} v^{1,y} \right) - {\mu_{j} \choose 1}_{q} \right)$$

Let  $v = v_j$ . Then,  $\sum_{1 \le y \le {n \brack 1}_q} (v_i^{1,y} v^{1,y})$  counts the number of 1-dimensional subspaces common to  $V_i$  and  $V_j$ . That is,  $\sum_{1 \le y \le {n \brack 1}_q} v_i^{1,y} v^{1,y} = \begin{bmatrix} \dim(V_i \cap V_j) \\ 1 \end{bmatrix}_q$ . In  $\mathbb{F}_p$ ,  $\begin{bmatrix} \dim(V_i \cap V_j) \\ 1 \end{bmatrix}_q \neq \begin{bmatrix} \mu_t \\ 1 \end{bmatrix}_q$  for any  $1 \le t \le s$ , if i = j, and  $\begin{bmatrix} \dim(V_i \cap V_j) \\ 1 \end{bmatrix}_q = \begin{bmatrix} \mu_t \\ 1 \end{bmatrix}_q$  for some  $1 \le t \le s$  if  $i \ne j$ . Accordingly,  $g^i(v_j) = \begin{cases} 0, & i \ne j \\ \ne 0, & i = j \end{cases}$ .

**Lemma 1.14** (Swallowing trick 1). Let  $s + k_r \leq n$  and  $r(s - r + 1) \leq b - 1$ . The collection of functions  $g^i$ ,  $1 \leq i \leq m$  together with the functions  $g^{x,y}$ ,  $0 \leq x \leq s - r$ ,  $1 \leq y \leq \begin{bmatrix} n \\ x \end{bmatrix}_q$  are linearly independent in  $\mathbb{F}_p^{\Omega}$  over  $\mathbb{F}_p$ .

*Proof.* Let

$$\sum_{1 \le i \le m} \alpha^i g^i + \sum_{\substack{0 \le x \le s - r \\ 1 \le y \le \begin{bmatrix} n \\ x \end{bmatrix}_q}} \alpha^{x,y} g^{x,y} = 0 \tag{1}$$

We know that  $g^i(v_j) = 0$  whenever  $i \neq j$ , and  $g^{x,y}(v_i) = 0, 1 \leq i \leq m$ . The latter holds because  $\dim(V_i) \equiv k_t \pmod{b}$ , say equal to  $bl + k_t$ , for some  $t \in [r]$ . Number of 1-dimensional subspaces in  $V_i$  will then be  $\begin{bmatrix} bl + k_t \\ 1 \end{bmatrix}_q$  which is equal to  $\begin{bmatrix} k_t \\ 1 \end{bmatrix}_q$  in  $\mathbb{F}_p$ . Suppose we evaluate L.H.S. of Equation (1) on  $v_1$ , then all terms except the first one vanish. This gives us  $\alpha^1 = 0$ , and reduces the relation by one term from left. Next, we put  $v = v_2$  to get  $\alpha^2 = 0$ , and so on. Finally, all  $\alpha^i$  terms are zero, and we are left only with functions  $g^{x,y}$ . These  $\alpha^{x,y}$  values are zero from Lemma 1.13. Therefore, we have shown that (1) implies that  $\alpha^i = 0, 1 \leq i \leq m$  and  $\alpha^{x,y} = 0, 0 \leq x \leq s-r, 1 \leq y \leq \begin{bmatrix} n \\ x \end{bmatrix}_q$ , and hence the given functions are linearly independent.  $\Box$ 

## **1.5 Proof of Theorem 1.2:** $(s + k_r \le n \text{ and } r(s - r + 1) \le b - 1)$

**Lemma 1.15.** The collection of functions  $f^{x,y}$ ,  $0 \le x \le s$ ,  $1 \le y \le {n \brack x}_q$ , spans all the functions  $g^{x,y}$ ,  $0 \le x \le s - r$ ,  $1 \le y \le {n \brack x}_q$  as well as the functions  $g^i$ ,  $1 \le i \le m$ .

#### *Proof.* Proof in Appendix C

This means that the above functions  $g^{x,y}$  and  $g^i$  belong to the span of functions  $f^{x,y}$  which is a function space of dimension at most S. From Lemma 1.14, we know that  $g^{x,y}$  and  $g^i$  are together linearly independent. Thus,

$$\sum_{j=0}^{s-r} {n \brack j}_q + m \le S = \sum_{j=0}^{s} {n \brack j}_q.$$
$$\Rightarrow |\mathcal{F}| = m \le {n \brack s}_q + {n \brack s-1}_q + \dots + {n \brack s-r+1}_q.$$

We thus have the following theorem.

**Theorem 1.16.** Let V be a vector space of dimension n over a finite field of size q. Let  $K = \{k_1, \ldots, k_r\}, L = \{\mu_1, \ldots, \mu_s\}$  be two disjoint subsets of  $\{0, 1, \ldots, b-1\}$  with  $k_1 < \cdots < k_r$ . Assume  $s + k_r \leq n$  and  $r(s - r + 1) \leq b - 1$ . Let  $\mathcal{F} = \{V_1, V_2, \ldots, V_m\}$  be a family of subspaces of V such that  $\forall i \in [m], \dim(V_i) \equiv k_t \pmod{b}$ , for some  $k_t \in K$ ; for every distinct  $i, j \in [m], \dim(V_i \cap V_j) \equiv \mu_t \pmod{b}$ , for some  $\mu_t \in L$ . Moreover, it is given that neither of the following two conditions hold:

(i) q + 1 is a power of 2, and b = 2

(ii) 
$$q = 2, b = 6$$

Then,  $|\mathcal{F}| \leq N(n, s, r, q)$ .

#### **1.6** Proof of Theorem 1.2: unrestricted

We start with the following lemma which claims that the functions  $g^{x,y}, 0 \le x \le s-r, 1 \le y \le {n \brack x}_q$ , and  $\forall t \in [r], x \not\equiv k_t \pmod{b}$  are linearly independent over  $\mathbb{F}_p$ .

**Lemma 1.17.** The collection of functions  $\{g^{x,y} \mid 0 \le x \le s-r, 1 \le y \le \begin{bmatrix} n \\ x \end{bmatrix}_q$ , and  $\forall t \in [r], x \ne k_t \pmod{b}$  are linearly independent in the function space  $\mathbb{F}_p^{\Omega}$  over  $\mathbb{F}_p$ .

*Proof.* Recall that  $g^{x,y}(v) = f^{x,y}(v) \prod_{t \in [r]} \left(\sum_{j=1}^{\binom{n}{1}_q} v^{1,j} - \begin{bmatrix} k_t \\ 1 \end{bmatrix}_q \right)$ . Assume, for the sake of contradiction,  $\sum_{\substack{0 \le x \le s-r \\ x \not\equiv k_t \pmod{p}, \forall t \in [r]}} \alpha^{x,y} g^{x,y} = 0$  with at least one  $\alpha^{x,y}$  as non-zero. Let  $\langle x_0, y_0 \rangle$  be the first

subspace, based on the ordering of subspaces defined in Section 1.3, such that  $\alpha^{x_0,y_0}$  is non-zero. Evaluating both sides on  $v_{x_0,y_0}$ , we see that all  $f^{x,y}$  (and therefore  $g^{x,y}$ ) with  $\langle x, y \rangle$  higher in the ordering than  $\langle x_0, y_0 \rangle$  will vanish (due to the virtue of our ordering), and so we get  $\alpha^{x_0,y_0} = 0$  which is a contradiction. Here we have crucially used the fact that by ignoring  $x \equiv k_t \pmod{p}$  cases, for any  $t \in [r]$ , we make sure that  $v_{x_0,y_0}$  used above always has  $x_0 \not\equiv k_t \pmod{b}$  and  $\begin{bmatrix} n \\ 1 \end{bmatrix}$ 

therefore 
$$\left(\sum_{j=1}^{\lfloor 1 \rfloor_q} v_{x_0,y_0}^{1,j} - \begin{bmatrix} k_t \\ 1 \end{bmatrix}_q\right) \not\equiv 0 \pmod{p}, \ \forall t \in [r].$$

**Lemma 1.18** (Swallowing trick 2). The collection of functions  $g^i$ ,  $1 \le i \le m$  together with the functions  $g^{x,y}$ ,  $0 \le x \le s - r, x \ne k_t \pmod{b}$ ,  $\forall t \in [r], 1 \le y \le \begin{bmatrix} n \\ x \end{bmatrix}_q$  are linearly independent in  $\mathbb{F}_p^{\Omega}$  over  $\mathbb{F}_p$ .

*Proof.* Proof is similar to the proof of Lemma 1.14.

Since s < b, for any  $0 \le x \le s - r$  and for any  $t \in [r]$ ,  $x \ne k_t \pmod{b}$  is equivalent to  $x \ne k_t$ . Combining Lemmas 1.17, 1.18 and 1.15, we have

$$\sum_{\substack{0 \le j \le s-r, \\ j \ne k_t, t \in [r]}} {n \brack j}_q + m \le \sum_{j=0}^s {n \brack j}_q$$

This implies,

$$|\mathcal{F}| = m \leq = \begin{cases} N(n, s, r, q), \text{ if } s < k_1 + r\\ N(n, s, r, q) + \sum_{t \in [r]} \begin{bmatrix} n\\k_t \end{bmatrix}_q, \text{ otherwise} \end{cases}$$

We thus have the following theorem which combined with Theorem 1.16 yields Theorem 1.2.

**Theorem 1.19.** Let V be a vector space of dimension n over a finite field of size q. Let  $K = \{k_1, \ldots, k_r\}, L = \{\mu_1, \ldots, \mu_s\}$  be two disjoint subsets of  $\{0, 1, \ldots, b-1\}$  with  $k_1 < \cdots < k_r$ . Let  $\mathcal{F} = \{V_1, V_2, \ldots, V_m\}$  be a family of subspaces of V such that  $\forall i \in [m], \dim(V_i) \equiv k_t \pmod{b}$ , for some  $k_t \in K$ ; for every distinct  $i, j \in [m], \dim(V_i \cap V_j) \equiv \mu_t \pmod{b}$ , for some  $\mu_t \in L$ . Moreover, it is given that neither of the following two conditions hold:

(i) q+1 is a power of 2, and b=2

(ii) 
$$q = 2, b = 6$$

Then,

$$|\mathcal{F}| \leq \left\{egin{array}{l} N(n,s,r,q), & ext{if } (s < k_1 + r) \ N(n,s,r,q) + \sum\limits_{t \in [r]} {n \brack k_t}_q, & ext{otherwise} \end{array}
ight.$$

## **PART B:** Fractional L-intersecting families of subspaces

Let  $L = \{\frac{a_1}{b_1}, \dots, \frac{a_s}{b_s}\}$  be a collection of positive irreducible fractions, each strictly less than 1. Let V be a vector space of dimension n over a finite field of size q. Let  $\mathcal{F}$  be a family of subspaces of V. Recall that, we call  $\mathcal{F}$  a fractional L-intersecting family of subspaces if  $\forall A, B \in \mathcal{F}$ ,  $dim(A \cap B) \in \{\frac{a_i}{b_i}dim(A), \frac{a_i}{b_i}dim(B)\}$ , for some  $\frac{a_i}{b_i} \in L$ . In Section 2, we prove a general upper bound for the size of a fractional L-intersecting family using Theorem 1.2 proved in Part A. In Section 3, we improve this upper bound for the special case when  $L = \{\frac{a}{b}\}$  is a singleton set with b being a prime number.

## 2 A general upper bound

The key idea we use here is to split the fractional L intersecting family  $\mathcal{F}$  into subfamilies and then use Theorem 1.2 to bound each of them.

**Lemma 2.1.** Let  $L = \{\frac{a_1}{b_1}, \frac{a_2}{b_2}, \dots, \frac{a_s}{b_s}\}$ , where every  $\frac{a_i}{b_i}$  is a irreducible fraction in the open interval (0, 1). Let  $\mathcal{F} = \{V_1, \dots, V_m\}$  be a fractional *L*-intersecting family of subspaces of a vector space *V* of dimension *n* over a finite field of size *q*. Let  $\mathcal{F}_k^p$  denote subspaces in  $\mathcal{F}$  whose dimensions

leave a remainder  $k \pmod{p}$ , where p is a prime number. That is,  $\mathcal{F}_k^p := \{W \in \mathcal{F} \mid dim(W) \equiv k \pmod{p}\}$ . Further, let k > 0, and  $p > \max(b_1, b_2, \ldots, b_s)$ . Then,

$$|\mathcal{F}_{k}^{p}| \leq \begin{cases} \begin{bmatrix} n \\ s \end{bmatrix}_{q}, \text{ if } (2p \leq n+2) \text{ or } (s < k+1) \\ \begin{bmatrix} n \\ s \end{bmatrix}_{q} + \begin{bmatrix} n \\ k \end{bmatrix}_{q}, \text{ otherwise.} \end{cases}$$

Proof. Apply Theorem 1.2 with family  $\mathcal{F}$  replaced by  $\mathcal{F}_k^p$ ,  $K = \{k\}$ , r = 1, b replaced by p, and each  $\mu_i$  replaced by  $\frac{a_i}{b_i}k$ . Let  $s' (\leq s)$  be the number of distinct  $\mu_i$ 's. Notice that k > 0, and  $p > b_i > a_i$  ensure that  $k \not\equiv \frac{a_i}{b_i}k \pmod{p}$ . Moreover, since  $s' \leq p - 1$  and  $k \leq p - 1$ , we have  $s' + k \leq n$  if  $2p \leq n+2$ . Since  $p > b_i$  and every  $b_i \geq 2$ , we have p > 2. This avoids the bad cases (i) and (ii) of Theorem 1.2. Thus, we satisfy the premise of Theorem 1.2 and the conclusion follows.

Suppose  $2p \le n+2$ . The above lemma immediately gives us a bound of  $|\mathcal{F}| \le |\mathcal{F}_0^p| + (p-1) \begin{bmatrix} n \\ s \end{bmatrix}_q$ . But it could be that most subspaces belong to  $\mathcal{F}_0^p$ . To overcome this problem, we instead choose a set of primes P such that no subspace can belong to  $\mathcal{F}_0^p$  for every  $p \in P$ . A natural choice is to take just enough primes in increasing order so that the product of these primes exceeds n, because then any subspace with dimension divisible by all primes in P will have a dimension greater than n, which is not possible.

**Lemma 2.2.** Let  $L = \{\frac{a_1}{b_1}, \frac{a_2}{b_2}, \dots, \frac{a_s}{b_s}\}$ , where every  $\frac{a_i}{b_i}$  is an irreducible fraction in the open interval (0, 1). Let  $\mathcal{F} = \{V_1, \dots, V_m\}$  be a fractional *L*-intersecting family of subspaces of a vector space *V* of dimension *n* over a finite field of size *q*. Let  $t := \max(b_1, b_2, \dots, b_s)$  and  $g(t, n) := \frac{2(2t+\ln n)}{\ln(2t+\ln n)}$ . Suppose  $2g(t, n) \ln(g(t, n)) \leq n+2$ . Then,

$$|\mathcal{F}| \le 2g^2(t,n)\ln(g(t,n)) \begin{bmatrix} n\\s \end{bmatrix}_q$$

*Proof.* For some  $\beta$  to be chosen later, choose P to be the set  $\{p_{\alpha+1}, p_{\alpha+2}, \ldots, p_{\beta}\}$  where  $p_l$  denotes the  $l^{th}$  prime number and  $p_{\alpha} \leq t < p_{\alpha+1} < p_{\alpha+2} < \cdots < p_{\beta}$ . Let l# denote the product of all primes less than or equal to l. Thus,  $p_l\#$  which is known as the *primorial function*, is the product of the first l primes. It is known that  $p_l\# = e^{(1+o(1))l \ln l}$  and  $l\# = e^{(1+o(1))l}$ . We require the following condition for the set P:

$$\frac{p_{\beta}\#}{t\#} > n$$

Using the bounds for  $p_l \#$  and l # discussed above, we find that it is sufficient to choose  $\beta \geq \frac{2(2t+\ln n)}{\ln(2t+\ln n)} := g(t,n)$ . Let  $\beta = g(t,n)$ . From the Prime Number Theorem, it follows that  $p_\beta$  (and so  $p_{\alpha+1}, p_{\alpha+2}, \ldots, p_{\beta-1}$  as well) is at most  $2g(t,n) \ln(g(t,n))$ . We are given that  $2p \leq 2p_\beta \leq n+2$ , for every  $p \in P$ . We apply Lemma 2.1 with  $p = p_{\alpha+1}$  to get

$$|\mathcal{F}| \le |\mathcal{F}_0^{p_{\alpha+1}}| + (p_{\alpha+1} - 1) \begin{bmatrix} n \\ s \end{bmatrix}_q$$

Next, apply Lemma 2.1 on  $\mathcal{F}_0^{p_{\alpha+1}}$  with  $p = p_{\alpha+2}$  and so on. As argued above, no subspace is left uncovered after we reach  $p_{\beta}$ . This means,

$$\begin{aligned} |\mathcal{F}| &\leq (p_{\alpha+1} + p_{\alpha+2} + \dots + p_{\beta} - (\beta - \alpha)) \begin{bmatrix} n \\ s \end{bmatrix}_{q} \\ &< (\beta - \alpha) p_{\beta} \begin{bmatrix} n \\ s \end{bmatrix}_{q} \\ &< \beta p_{\beta} \begin{bmatrix} n \\ s \end{bmatrix}_{q} \\ &\leq 2g^{2}(t, n) \ln(g(t, n)) \begin{bmatrix} n \\ s \end{bmatrix}_{q} \end{aligned}$$

**Lemma 2.3.** Let  $L = \{\frac{a_1}{b_1}, \frac{a_2}{b_2}, \dots, \frac{a_s}{b_s}\}$ , where every  $\frac{a_i}{b_i}$  is an irreducible fraction in the open interval (0, 1). Let  $\mathcal{F} = \{V_1, \dots, V_m\}$  be a fractional *L*-intersecting family of subspaces of a vector space *V* of dimension *n* over a finite field of size *q*. Let  $t := \max(b_1, b_2, \dots, b_s)$  and  $g(t, n) := \frac{2(2t+\ln n)}{\ln(2t+\ln n)}$ . Then,

$$|\mathcal{F}| \le 2g^2(t,n)\ln(g(t,n)) \begin{bmatrix} n\\ s \end{bmatrix}_q + g(t,n)\sum_{i=1}^{s-1} \begin{bmatrix} n\\ i \end{bmatrix}_q$$

*Proof.* Let  $P = \{p_{\alpha+1}, p_{\alpha+2}, \ldots, p_{\beta}\}$ , where  $\beta = g(t, n)$  and  $p_{\beta} \leq 2g(t, n) \ln(g(t, n))$ . The proof is similar to the proof of Lemma 2.2. We apply Lemma 2.1 with  $p = p_{\alpha+1}$  to show that

$$|\mathcal{F}| \le |\mathcal{F}_0^{p_{\alpha+1}}| + (p_{\alpha+1} - 1) \begin{bmatrix} n \\ s \end{bmatrix}_q + \sum_{i=1}^{s-1} \begin{bmatrix} n \\ i \end{bmatrix}_q$$

Next, we apply Lemma 2.1 on  $\mathcal{F}_0^{p_{\alpha+1}}$  with  $p = p_{\alpha+2}$  and so on as shown in the proof of Lemma 2.2 to get the desired bound.

$$\begin{aligned} |\mathcal{F}| &\leq (p_{\alpha+1} + p_{\alpha+2} + \dots + p_{\beta} - (\beta - \alpha)) \begin{bmatrix} n \\ s \end{bmatrix}_{q} + (\beta - \alpha) \sum_{i=1}^{s-1} \begin{bmatrix} n \\ i \end{bmatrix}_{q} \\ &\leq (\beta - \alpha) \Big( p_{\beta} \begin{bmatrix} n \\ s \end{bmatrix}_{q} + \sum_{i=1}^{s-1} \begin{bmatrix} n \\ i \end{bmatrix}_{q} \Big) \\ &\leq \beta \Big( p_{\beta} \begin{bmatrix} n \\ s \end{bmatrix}_{q} + \sum_{i=1}^{s-1} \begin{bmatrix} n \\ i \end{bmatrix}_{q} \Big) \\ &\leq 2g^{2}(t, n) \ln(g(t, n)) \begin{bmatrix} n \\ s \end{bmatrix}_{q} + g(t, n) \sum_{i=1}^{s-1} \begin{bmatrix} n \\ s \end{bmatrix}_{q} \end{aligned}$$

Since  $p_{\alpha+1} > t$ , we have  $p_{\alpha+1}p_{\alpha+2}\cdots p_{\beta} > t^{\beta-\alpha}$ . This implies that, if  $t^{\beta-\alpha} \ge n$ , then the product of the primes in P will be greater than n as desired. Substituting  $\beta - \alpha$  with  $\frac{\ln n}{\ln t}$  (and  $p_{\beta}$  with  $2g(t,n)\ln(g(t,n))$ ) in the second inequality above, we get another upper bound of

 $|\mathcal{F}| \leq 2g(t,n) \frac{\ln(n)\ln(g(t,n))}{\ln t} {n \choose s}_q + \frac{\ln n}{\ln t} \sum_{i=1}^{s-1} {n \choose i}_q.$  We can do a similar substitution for  $\beta - \alpha$  in the calculations done at the end of the proof of Lemma 2.2 to get a similar bound.

Combining all the results in this section, we get Theorem 1.4

## 3 An improved bound for singleton L

In this section, we improve the upper bound for the size of a fractional *L*-intersecting family obtained in Theorem 1.4 for the special case  $L = \{\frac{a}{b}\}$ , where *b* is a constant prime. Before we give the proof, below we restate the the statement of Theorem 1.5.

**Statement of Theorem 1.5**: Let  $L = \{\frac{a}{b}\}$ , where  $\frac{a}{b}$  is a positive irreducible fraction less than 1 and b is a prime. Let  $\mathcal{F}$  be a fractional L-intersecting family of subspaces of a vector space V of dimension n over a finite field of size q. Then, we have  $|\mathcal{F}| \leq (b-1)(\binom{n}{1} + 1)\lceil \frac{\ln n}{\ln b} \rceil + 2$ .

*Proof.* We assume that all the subspaces in the family except possibly one subspace, say W, have a dimension divisible by b. Otherwise,  $\mathcal{F}$  cannot satify the property of a fractional  $\frac{a}{b}$ -intersecting family. Let us ignore W in the discussion to follow. For any subspace  $V_i$  that is not the zero subspace, let k be the largest power of b that divides  $dim(V_i)$ . Then,  $dim(V_i) = rb^{k+1} + jb^k$ , for some  $1 \leq j < b, r \geq 0$ . Consider the subfamily,

$$\mathcal{F}^{j,k} = \{ V_i : b^k | dim(V_i), b^{k+1} \not | dim(V_i), dim(V_i) = rb^{k+1} + jb^k \text{ for some } r \ge 0, j \in [b] \}$$

The subfamily  $\mathcal{F}^{j,k}$ ,  $1 \leq k \leq \lceil \frac{\ln n}{\ln b} \rceil$ ,  $1 \leq j < b$ , cover each and every subspace (except the zero subspace and the subspace W) of  $\mathcal{F}$  exactly once. We will show that  $|\mathcal{F}^{j,k}| \leq {n \brack 1}_q + 1$ , which when multiplied with the number of values j and k can take will immediately imply the theorem.

Let  $m^{j,k} = |\mathcal{F}^{j,k}|$ . Let  $M^{j,k}$  be an  $m^{j,k} \times \begin{bmatrix} n \\ 1 \end{bmatrix}_q 0$ -1 matrix whose rows correspond to the

subspaces of  $\mathcal{F}^{j,k}$  in any given order, whose columns correspond to the 1-dimensional subspaces of V in any given order, and the  $(i-l)^{th}$  entry is 1 if and only if the  $i^{th}$  subspace of  $\mathcal{F}^{j,k}$  contains the  $l^{th}$  1-dimensional subspace. Let  $N^{j,k} = M^{j,k} \cdot (M^{j,k})^T$ . Any diagonal entry  $N_{i,i}^{j,k}$  is the number of 1-dimensional subspaces in the  $i^{th}$  subspace in  $\mathcal{F}^{j,k}$ , and an off-diagonal entry  $N_{i,l}^{j,k}$ is number of 1-dimensional subspaces common to the  $i^{th}$  and  $l^{th}$  subspaces of  $\mathcal{F}^{j,k}$ .

$$\begin{split} N_{i,i}^{j,k} &= \begin{bmatrix} r_1 b^{k+1} + j b^k \\ 1 \end{bmatrix}_q = \begin{bmatrix} b^{k-1} \\ 1 \end{bmatrix}_q \begin{bmatrix} r_1 b^2 + j b \\ 1 \end{bmatrix}_{q^{b^{k-1}}},\\ N_{i,l}^{j,k} &= \begin{bmatrix} r_2 a b^k + j a b^{k-1} \\ 1 \end{bmatrix}_q = \begin{bmatrix} b^{k-1} \\ 1 \end{bmatrix}_q \begin{bmatrix} r_2 a b + j a \\ 1 \end{bmatrix}_{q^{b^{k-1}}}, \end{split}$$

for some  $r_1, r_2$  (may be different for different values of i, l). Let  $P^{j,k}$  be the matrix obtained by dividing each entry of  $N^{j,k}$  by  $\begin{bmatrix} b^{k-1} \\ 1 \end{bmatrix}_{c}$ .

$$\det(N^{j,k}) = \begin{bmatrix} b^{k-1} \\ 1 \end{bmatrix}_q^{m^{j,k}} \det(P^{j,k})$$

We will show that  $\det(P^{j,k})$  is non-zero, thereby implying  $\det(N^{j,k})$  is also non-zero. Consider  $\det(P^{j,k}) \pmod{\begin{bmatrix} b \\ 1 \end{bmatrix}}_{c^{b^{k-1}}}$ .

$$\begin{split} P_{i,i}^{j,k} &\equiv \begin{bmatrix} r_1b^2 + jb \\ 1 \end{bmatrix}_{q^{b^{k-1}}} \pmod{\begin{bmatrix} b \\ 1 \end{bmatrix}_{q^{b^{k-1}}}} \equiv \begin{pmatrix} \mod \begin{bmatrix} b \\ 1 \end{bmatrix}_{q^{b^{k-1}}} \pmod{\begin{bmatrix} b \\ 1 \end{bmatrix}_{q^{b^{k-1}}}} \\ P_{i,l}^{j,k} &\equiv \begin{bmatrix} r_2ab + ja \\ 1 \end{bmatrix}_{q^{b^{k-1}}} \pmod{\begin{bmatrix} b \\ 1 \end{bmatrix}_{q^{b^{k-1}}}} \equiv \begin{bmatrix} r_3 \\ 1 \end{bmatrix}_{q^{b^{k-1}}} \pmod{\begin{bmatrix} b \\ 1 \end{bmatrix}_{q^{b^{k-1}}}}, \end{split}$$

where  $r_3 = ja \mod b$  and  $1 \le r_3 \le b-1$  (since j < b, a < b, and b is a prime, we have  $1 \le r_3 \le b-1$ ). We know that the determinant of an  $r \times r$  matrix where diagonal entries are 0 and off-diagonal entries are all 1 is  $(-1)^{r-1}(r-1)$ .

$$\det(P^{j,k}) \equiv {\binom{r_3}{1}}_{q^{b^{k-1}}}^{m^{j,k}} (-1)^{m^{j,k}-1} (m^{j,k}-1) \pmod{\binom{b}{1}}_{q^{b^{k-1}}}^{p^{j,k}}$$

Let  $Q^{j,k}$  be the matrix formed by taking all but the last row and the last column of  $P^{j,k}$ .

$$\det(Q^{j,k}) \equiv {\binom{r_3}{1}}_{q^{b^{k-1}}} {}^{m^{j,k}-1} (-1)^{m^{j,k}-2} (m^{j,k}-2) \pmod{\binom{b}{1}}_{q^{b^{k-1}}}$$

We will now show that one of  $\det(P^{j,k})$  or  $\det(Q^{j,k})$  is non-zero (mod  $\begin{bmatrix} b\\1 \end{bmatrix}_{q^{b^{k-1}}}$ ) and therefore non-zero in  $\mathbb{R}$ . First, we show that  $\begin{bmatrix} r_3\\1 \end{bmatrix}_{q^{b^{k-1}}}^{m^{j,k}}$  is not divisible by  $\begin{bmatrix} b\\1 \end{bmatrix}_{q^{b^{k-1}}}$ . Suppose  $s_3 \equiv r_3^{-1}$ (mod b).

$$\begin{bmatrix} r_3 \\ 1 \end{bmatrix}_{q^{b^{k-1}}}^{m^{j,k}} \begin{bmatrix} s_3 \\ 1 \end{bmatrix}_{q^{r_3b^{k-1}}}^{m^{j,k}} = \begin{bmatrix} r_3s_3 \\ 1 \end{bmatrix}_{q^{b^{k-1}}}^{m^{j,k}}$$
$$\begin{bmatrix} r_3s_3 \\ 1 \end{bmatrix}_{q^{b^{k-1}}}^{m^{j,k}} \equiv \begin{bmatrix} 1 \\ 1 \end{bmatrix}_{q^{b^{k-1}}}^{m^{j,k}} \pmod{\begin{bmatrix} b \\ 1 \end{bmatrix}_{q^{b^{k-1}}}} \equiv 1 \pmod{\begin{bmatrix} b \\ 1 \end{bmatrix}_{q^{b^{k-1}}}}$$

Therefore,  $\begin{bmatrix} r_3 \\ 1 \end{bmatrix}_{q^{b^{k-1}}}^{m^{j,k}}$  is invertible modulo  $\begin{bmatrix} b \\ 1 \end{bmatrix}_{q^{b^{k-1}}}$ , and hence the former is not divisible by the latter. Suppose  $\begin{bmatrix} r_3 \\ 1 \end{bmatrix}_{q^{b^{k-1}}}^{m^{j,k}} (-1)^{m^{j,k}-1} (m^{j,k}-1)$  is divible by  $\begin{bmatrix} b \\ 1 \end{bmatrix}_{q^{b^{k-1}}}$ . We may ignore  $(-1)^{m^{j,k}-1}$  for divisibility purpose. Then, there must be a product of prime powers that is equal to  $(m^{j,k}-1)$  multiplied by  $\begin{bmatrix} r_3 \\ 1 \end{bmatrix}_{q^{b^{k-1}}}^{m^{j,k}}$  such that this product is divisible by  $\begin{bmatrix} b \\ 1 \end{bmatrix}_{q^{b^{k-1}}}$ . Observe that,  $\begin{bmatrix} r_3 \\ 1 \end{bmatrix}_{q^{b^{k-1}}}^{m^{j,k-1}}$  has only lesser powers of the same primes, and  $m^{j,k}-1$  and  $m^{j,k}-2$  cannot have any prime in common. So, the product  $\begin{bmatrix} r_3 \\ 1 \end{bmatrix}_{q^{b^{k-1}}}^{m^{j,k-1}} (m^{j,k}-2)$  cannot be divisible by  $\begin{bmatrix} b \\ 1 \end{bmatrix}_{q^{b^{k-1}}}$ , which is what we wanted to prove.

Therefore, either  $P^{j,k}$  or  $Q^{j,k}$  is a full rank matrix, or  $rank(P^{j,k}) \ge m^{j,k} - 1$ . Being a non-zero multiple of  $P^{j,k}$ ,  $rank(N^{j,k}) \ge m^{j,k} - 1$ . But we know that  $rank(AB) \le \min(rank(A), rank(B))$ , for any two matrices A, B.

$$\begin{split} m^{j,k} - 1 &\leq rank(N^{j,k}) \leq \min(rank(M^{j,k}), rank((M^{j,k})^T)) \\ &= rank(M^{j,k}) \\ &\leq \begin{bmatrix} n \\ 1 \end{bmatrix}_a \end{split}$$

Or,  $m^{j,k} \leq \begin{bmatrix} n \\ 1 \end{bmatrix}_q + 1$ , as required. It follows that,

$$|\mathcal{F}| = m \le 2 + \sum_{\substack{1 \le k \le \lceil \frac{\ln n}{\ln b} \rceil \\ 1 \le j < b}} m^{j,k} \le (b-1) \left( \binom{n}{1}_q + 1 \right) \left\lceil \frac{\ln n}{\ln b} \right\rceil + 2.$$

## 4 Concluding remarks

In Theorem 1.2, for  $|\mathcal{F}|$  to be at most N(n, s, r, q), one of the necessary conditions is  $r(s-r+1) \leq b-1$ . When r = 1, this condition is always true as  $L \subseteq \{0, 1, \ldots, b-1\}$ . However, when  $r \geq 2$ , it is not the case. Would it be possible to get the same upper bound for  $|\mathcal{F}|$  without having to satisfy such a strong necessary condition? Another interesting question concerning Theorem 1.2 is regarding its tightness. From Example 1, we know that Theorem 1.2 is tight when r = 1. However, since Theorem 1.2 requires the sets K and L to be disjoint it is not possible to extend the construction in Example 1 to obtain a tight example for the case  $r \geq 2$ . Further, we know of no other tight example for this case. Therefore, we are not clear whether Theorem 1.2 is tight when  $r \geq 2$ .

We believe that the upper bounds given by Theorems 1.4 and 1.5 are not tight. Proving tight upper bounds in both the scenarios is a question that is obviously interesting. One possible approach to try would be to answer the following simpler question. Consider the case when  $L = \{\frac{1}{2}\}$ . We call such a family a *bisection-closed family* of subspaces. Let  $\mathcal{F}$  be a bisection closed family of subspaces of a vector space V of dimension n over a finite field of size q. From Theorem 1.5, we know that  $|\mathcal{F}| \leq ({n \brack 1}_q + 1) \log_2 n + 2$ . We believe that  $|\mathcal{F}| \leq c {n \brack 1}_q$ , where c is a constant. Example 3 gives a 'trivial' bisection-closed family of size  ${n-1 \brack 1}_q$  where every subspace contains the vector  $v_1$ . It would be interesting to look for non-trivial examples of large

bisection-closed families.

## 5 References

- [Alon et al., 1991] Alon, N., Babai, L., and Suzuki, H. (1991). Multilinear polynomials and frankl-ray-chaudhuri-wilson type intersection theorems. Journal of Combinatorial Theory, Series A, 58(2):165 – 180.
- [Balachandran et al., 2019] Balachandran, N., Mathew, R., and Mishra, T. K. (2019). Fractional L-intersecting families. The Electronic Journal of Combinatorics, 26(2):40.

- [Deza and Frankl, 1983] Deza, M. and Frankl, P. (1983). Erdös-ko-rado theorem—22 years later. SIAM Journal on Algebraic Discrete Methods, 4(4):419–431.
- [Erdos et al., 1961] Erdos, P., Ko, C., and Rado, R. (1961). Intersection theorems for systems of finite sets. The Quarterly Journal of Mathematics, 12:313–320.
- [Frankl and Graham, 1985] Frankl, P. and Graham, R. (1985). Intersection theorems for vector spaces. *European Journal of Combinatorics*, 6(2):183 187.
- [Frankl and Wilson, 1981a] Frankl, P. and Wilson, R. M. (1981a). Intersection theorems with geometric consequences. *Combinatorica*, 1(4):357–368.
- [Frankl and Wilson, 1981b] Frankl, P. and Wilson, R. M. (1981b). Intersection theorems with geometric consequences. *Combinatorica*, 1(4):357–368.
- [Hsieh, 1975] Hsieh, W. (1975). Intersection theorems for systems of finite vector spaces. Discrete Mathematics, 12(1):1–16.
- [Liu and Yang, 2014] Liu, J. and Yang, W. (2014). Set systems with restricted k-wise lintersections modulo a prime number. *European Journal of Combinatorics*, 36:707–719.
- [Liu et al., 2018] Liu, J. Q., Zhang, S. G., and Xiao, J. M. (2018). A common generalization to theorems on set systems with l-intersections. Acta Mathematica Sinica, English Series, 34(7):1087–1100.
- [Ray-Chaudhuri and Wilson, 1975] Ray-Chaudhuri, D. K. and Wilson, R. M. (1975). On tdesigns. Osaka J. Math., 12(3):737–744.
- [Zsigmondy, 1892] Zsigmondy, K. (1892). Zur theorie der potenzreste. Monatshefte f
  ür Mathematik und Physik, 3(1):265–284.

## A Proof of Proposition 1.7

*Proof.* (Forward direction)  $\beta(W) = \sum_{U \subseteq W} \alpha(U)$  is given.

$$\sum_{\substack{U \subseteq W \\ d = dim(W) - dim(U)}} (-1)^d q^{\frac{d(d-1)}{2}} \beta(U) = \sum_{\substack{U \subseteq W \\ d = dim(W) - dim(U)}} \left( (-1)^d q^{\frac{d(d-1)}{2}} \sum_{U' \subseteq U} \alpha(U') \right)$$
$$= \sum_{\substack{U' \subseteq W \\ U' \subseteq W}} \left( \alpha(U') \left( \sum_{\substack{U' \subseteq U \subseteq W \\ d = dim(W) - dim(U)}} (-1)^d q^{\frac{d(d-1)}{2}} \right) \right)$$
$$= \sum_{\substack{U' \subseteq W \\ U' \subseteq W}} \left( \alpha(U') \left( \sum_{\substack{d = 0 \\ d = 0}} \left( \left( \frac{dim(W) - dim(U')}{d} \right)_q (-1)^d q^{\frac{d(d-1)}{2}} \right) \right) \right)$$

We know that  $\sum_{d=0}^{n} {n \brack d}_{q} (-1)^{d} q^{\frac{d(d-1)}{2}} = 0$  unless n = 0, and 1 otherwise. Therefore, only dim(W) - dim(U') = 0 or U' = W term will survive in the summation above, giving,

$$\sum_{\substack{U\subseteq W\\ d=\dim(W)-\dim(U)}} (-1)^d q^{\frac{d(d-1)}{2}} \beta(U) = \alpha(W)$$

(Reverse direction)  $\alpha(W) = \sum_{\substack{U \subseteq W \\ d = dim(W) - dim(U)}} (-1)^d q^{\frac{d(d-1)}{2}} \beta(U)$  is given.

$$\begin{split} \sum_{U \subseteq W} \alpha(U) &= \sum_{U \subseteq W} \left( \sum_{\substack{U' \subseteq U \\ d = dim(U) - dim(U')}} (-1)^d q^{\frac{d(d-1)}{2}} \beta(U') \right) \\ &= \sum_{U' \subseteq W} \left( \beta(U') \sum_{\substack{U' \subseteq U \subseteq W \\ d = dim(U) - dim(U')}} (-1)^d q^{\frac{d(d-1)}{2}} \right) \\ &= \sum_{U' \subseteq W} \left( \beta(U') \sum_{\substack{d = 0 \\ d = 0}}^{\dim(W) - dim(U')} \left( (-1)^d q^{\frac{d(d-1)}{2}} \left[ \frac{dim(W) - dim(U')}{d} \right]_q \right) \right) \end{split}$$

The only term that will survive in the above summation is the term when dim(W) - dim(U') = 0 or U' = W. This proves  $\sum_{U \subseteq W} \alpha(U) = \beta(W)$ .

## B Proof of Proposition 1.9

*Proof.* We know that  $\beta(W) = \sum_{U \subseteq W} \alpha(U)$ .

$$\sum_{\substack{T: \ W \subseteq T \subseteq Y\\ d=dim(Y)-dim(T)}} (-1)^d q^{\frac{d(d-1)}{2}} \beta(T) = \sum_{\substack{T: \ W \subseteq T \subseteq Y\\ d=dim(Y)-dim(T)}} \left( (-1)^d q^{\frac{d(d-1)}{2}} \sum_{U \subseteq T} \alpha(U) \right)$$
$$= \sum_{U \subseteq Y} \left( \alpha(U) \sum_{\substack{T: U \subseteq T,\\ W \subseteq T \subseteq Y,\\ d=dim(Y)-dim(T)}} (-1)^d q^{\frac{d(d-1)}{2}} \right)$$

Let W' be the span of union (of vectors) of U and W. From our notation above,  $W' = U \cup W$ . As T is a superspace of both U and W, T contains all the vectors of U and W. Thus,  $W' \subseteq T$ .

$$\sum_{\substack{T: \ W \subseteq T \subseteq Y \\ d = \dim(Y) - \dim(T)}} (-1)^d q^{\frac{d(d-1)}{2}} \beta(T) = \sum_{U \subseteq Y} \left( \alpha(U) \sum_{\substack{T: W \cup U \subseteq T \subseteq Y \\ d = \dim(Y) - \dim(T)}} (-1)^d q^{\frac{d(d-1)}{2}} \right)$$
$$= \sum_{U \subseteq Y} \left( \alpha(U) \sum_{d=0}^{\dim(Y) - \dim(W')} \left( (-1)^d q^{\frac{d(d-1)}{2}} \left[ \frac{\dim(Y) - \dim(W')}{d} \right]_q \right) \right)$$

Again, the only non-zero term in the second summation above is when dim(Y) - dim(W') = 0, or Y = W'.

$$\sum_{\substack{T: \ W \subseteq T \subseteq Y\\ d=dim(Y)-dim(T)}} (-1)^d q^{\frac{d(d-1)}{2}} \beta(T) = \sum_{U: U \cup W = Y} \alpha(U)$$

## C Proof of Lemma 1.15

Proof. Let  $v \in \mathbb{F}_2^S$ . The key observation here is that the product  $f^{x,y}(v)f^{1,z}(v), 0 \le x \le s-1, 1 \le y \le \begin{bmatrix}n\\x\end{bmatrix}_q, 1 \le z \le \begin{bmatrix}n\\1\end{bmatrix}_q$  may be replaced by the function  $f^{x',w}(v)$ , where  $x \le x' \le x+1, 1 \le w \le \begin{bmatrix}n\\x'\end{bmatrix}_q$ . If  $V_{1,z} \subseteq V_{x,y}$ , it is trivial that  $f^{x,y}(v)f^{1,z}(v) = f^{x,y}(v)$ , since  $f^{x,y}(v) = 1$  only if  $f^{1,z}(v) = 1$ . If  $V_{1,z} \not\subseteq V_{x,y}$ , we let  $V_{x',w}$  be the span of union of vectors of  $V_{1,z}$  and  $V_{x,y}$ . Suppose, a vector space U contains both  $V_{1,z}$  and  $V_{x,y}$ . Then, it is clear that it must contain the span of their union as well. Similarly, a vector space U that does not contain either  $V_{1,z}$  or  $V_{x,y}$ , cannot contain  $V_{x',w}$ . Thus,  $f^{x,y}(v)f^{1,z}(v) = f^{x',w}(v)$ . To see why x' = x + 1 (in case  $V_{1,z} \not\subseteq V_{x,y}$ ), the space  $V_{x',w}$  may be obtained by taking any (non-zero) vector of  $V_{1,z}$  and introducing it into the basis of  $V_{x,y}$ . The space spanned by this extended basis is exactly  $V_{x',w}$  by definition, and the size of basis has increased by exactly 1.

By induction, it follows that,

$$f^{1,y_1}(v)f^{1,y_2}(v)\cdots f^{1,y_l}(v) = f^{x,y}(v)$$

for some x, y where,  $1 \le x \le l, 1 \le y \le {n \brack x}_q$ . That is, a product of l functions of the form  $f^{1,y}$  may be replaced by a single function  $f^{x,y}$  where x is at most l.

Now consider functions

$$\begin{split} g^{i}(v) &= g^{i}(v^{0,1}, v^{1,1}, \cdots, v^{1, {[n] \atop l]_{q}}}, \cdots, v^{s,1}, \cdots, v^{s, {[n] \atop s]_{q}}}) \\ &= \prod_{j=1}^{s} \left( \sum_{1 \le y \le {[n] \atop l]_{q}}} \left( v^{1,y}_{i} v^{1,y} \right) - {[\mu_{j} \atop 1]_{q}} \right) \\ &= \prod_{j=1}^{s} \left( \sum_{1 \le y \le {[n] \atop l]_{q}}} \left( v^{1,y}_{i} f^{1,y}_{i}(v) \right) - {[\mu_{j} \atop 1]_{q}} \right) \end{split}$$

Since the functions  $f^{x,y}$  only take 0/1 values, we can reduce any exponent of 2 or more on the function after expanding the product to 1. Moreover, the terms will all be products of the form  $f^{1,y_1}f^{1,y_2}\cdots f^{1,y_l}(v), 1 \leq l \leq s$ . These are replaced according to the observation above by single function of the form  $f^{x,y}(v)$ , and thus the set of function  $f^{x,y}, 0 \le x \le s, 1 \le y \le {n \brack x}_q$ span all functions  $g^i(v)$ . Note that  $f^{0,1}(v)$  is the constant function 1. Similarly, for  $0 \le x \le s - r, 1 \le y \le \begin{bmatrix} n \\ x \end{bmatrix}_q$ ,

$$g^{x,y}(v) = f^{x,y}(v) \prod_{t \in [r]} \left( \sum_{j=1}^{\binom{n}{1}_{q}} v^{1,j} - \binom{k_{t}}{1}_{q} \right)$$
  
$$= f^{x,y}(v) \prod_{t \in [r]} \left( \sum_{j=1}^{\binom{n}{1}_{q}} f^{1,j}(v) - \binom{k_{t}}{1}_{q} \right)$$
  
$$= f^{x,y}(v) \left( \sum_{\substack{r \\ x'=0}}^{r} \sum_{\substack{y'=1 \\ y'=1}}^{\binom{n}{2}} c_{x',y'} f^{x',y'}(v) \right)$$
  
$$= \sum_{x'=0}^{s} \sum_{\substack{y'=1 \\ y'=1}}^{\binom{n}{2}} c_{x',y'} f^{x',y'}(v)$$
  
$$(c_{x',y'} \text{ are constants})$$

Thus, the set of function  $f^{x,y}, 0 \le x \le s, 1 \le y \le {n \brack x}_q$  span all functions  $g^{x,y}(v), 0 \le x \le x \le x \le x$  $s-r, 1 \le y \le \begin{bmatrix} n \\ x \end{bmatrix}_{q}.$