

A parsimonious family of multivariate Poisson-lognormal distributions for clustering multivariate count data

Sanjeena Subedi* Ryan Browne †

Abstract

Multivariate count data are commonly encountered through high-throughput sequencing technologies in bioinformatics, text mining, or in sports analytics. Although the Poisson distribution seems a natural fit to these count data, its multivariate extension is computationally expensive. In most cases mutual independence among the variables is assumed, however this fails to take into account the correlation among the variables usually observed in the data. Recently, mixtures of multivariate Poisson-lognormal (MPLN) models have been used to analyze such multivariate count measurements with a dependence structure. In the MPLN model, each count is modeled using an independent Poisson distribution conditional on a latent multivariate Gaussian variable. Due to this hierarchical structure, the MPLN model can account for over-dispersion as opposed to the traditional Poisson distribution and allows for correlation between the variables. Rather than relying on a Monte Carlo-based estimation framework which is computationally inefficient, a fast variational-EM based framework

*Department of Mathematical Sciences, Binghamton University, State University of New York, 4400 Vestal Parkway East, Binghamton, NY, USA 13902. e: sdang@binghamton.edu

†Department of Statistics and Actuarial Science, University of Waterloo, 200 University Avenue West, Waterloo, ON, Canada. e: ryan.browne@uwaterloo.ca

is used here for parameter estimation. Further, a parsimonious family of mixtures of Poisson-lognormal distributions are proposed by decomposing the covariance matrix and imposing constraints on these decompositions. Utility of such models is shown using simulated and benchmark datasets.

Keywords: BIC, clustering, count data, MCLUST, mixture models, model-based clustering, MPLN, variational approximations, Variational EM algorithm.

1 Introduction

With the emergence of next generation sequencing technologies that provide a fast and cost-effective data generation platform, multivariate count data are becoming ubiquitous in bioinformatics. Although there has been a big explosion in the approaches for data generation and a dramatic reduction in cost and time, efficiently analyzing these complex biological data sets still remains a challenge. Cluster analysis is widely used in bioinformatics for identification and analysis of population heterogeneity. Clustering allows us to summarize data into homogenous groups or clusters, each with its unique characteristics. de Souto et al. (2008) did a comparative analysis of several clustering techniques on 35 different cancer gene expression data sets and concluded that a mixture model-based clustering approach showed superior performance in terms of recovering the true structure of the data as opposed to the traditional distance-based approaches such as k -means and hierarchical methods. Model-based clustering, which utilizes mixture models, has been increasingly used in the last two decades (Gollini and Murphy, 2014; Subedi and McNicholas, 2014; Bouveyron and Brunet, 2012; Subedi et al., 2015; Browne and McNicholas, 2015; Kosmidis and Karlis, 2016; Dang et al., 2015; Vrbik and McNicholas, 2014; Tortora et al., 2019; Subedi and McNicholas, 2020). A finite mixture model assumes that the population consists of a finite mixture

of subpopulations or components, which can be represented by a parametric model. If d -dimensional random vector \mathbf{Y} arises from a G component finite mixture model, its density can be written as

$$f(\mathbf{y} \mid \boldsymbol{\vartheta}) = \sum_{g=1}^G \pi_g f(\mathbf{y} \mid \boldsymbol{\varphi}_g),$$

where π_g for $g = 1, \dots, G$ is the mixing proportion such that $\pi_g > 0$ and $\sum_g \pi_g = 1$, $f(\cdot)$ is the component specific probability density function or the probability mass function with parameters $\boldsymbol{\varphi}_g$, and $\boldsymbol{\vartheta}$ contains all the unknown parameters for the finite mixture. Depending on the nature of the data, an appropriate $f(\cdot)$ is used.

A number of initial works on univariate analysis for a count variable from next generation sequencing data focused on a Poisson model (Marioni et al., 2008; Bullard et al., 2010; Witten et al., 2010). However, a Poisson distribution also imposes a mean-variance relationship such that mean and variance must be equal and most modern biological data typically exhibits over-dispersion (i.e., the observed variation is larger than what is predicted by the Poisson model, see Anders and Huber (2010)). A commonly used approach to incorporate this additional variability in univariate case is to model the mean parameter of Poisson distribution as a random variable and impose a distribution. Several different distributions has been proposed: the inverse-Gaussian distribution yields a Poisson-inverse Gaussian distribution (Holla, 1967), the inverse-Gamma distribution yields a Poisson-inverse Gamma distribution Willmot (1993), the lognormal distribution yields a Poisson-lognormal distribution (Bulmer, 1974), and the gamma distribution yields a gamma-Poisson distribution (i.e., the negative binomial distribution) (Greenwood and Yule, 1920; Collings and Margolin, 1985). See Karlis and Xekalaki (2005) for a detailed list of other possible priors. Another approach is to scale the mean parameter by a random variable and impose a beta

prior on that scaling random variable resulting in a Poisson-Beta distribution (Gurland, 1958). Amongst these, the most popular approach for statistical methods pertaining next generation sequence data is the gamma-Poisson distribution also known as the negative binomial distribution (Robinson et al., 2010; Yu et al., 2013; Love et al., 2014; Dong et al., 2016). In a gamma-Poisson distribution, λ is modeled as a random variable from the Gamma distribution such that $Y_i | \lambda_i \sim \text{Poisson}(\lambda)$ and $\lambda_i \sim \text{Gamma}(\alpha, \beta)$. The resulting marginal distribution of Y_i will then be a negative binomial distribution (Greenwood and Yule, 1920; Collings and Margolin, 1985) with mean $\mathbb{E}(Y) = \alpha\beta$, dispersion parameter $\phi = \frac{1}{\alpha}$, and variance $\mathbb{V}(Y) = \mathbb{E}(Y) + \mathbb{E}(Y)^2\phi > \mathbb{E}(Y)$. Note that this is a univariate distribution.

While the multivariate extensions of these distributions seem a natural choice to modelling multivariate count data, these models are computationally expensive and/or they impose restrictions on the covariance structure. For example, in addition to the computational burden (Brijs et al., 2004), the multivariate Poisson distribution by (Teicher, 1954; Campbell, 1934) can only allow for positive correlation among the variables Karlis and Meligkotsidou (2007); Inouye et al. (2017). Several different formulation of multivariate version of the Poisson-beta distribution was studied by Sarabia and Gómez-Déniz (2011); these formulations are either computationally intensive especially for high dimensional data, have restrictions on the covariance structure or the parameters in the marginal distributions are not free (i.e., they share the hyperparameters). Hence, independence between variables is assumed for multivariate analysis in most cases (Rau et al., 2015; Si et al., 2014; Papastamoulis et al., 2016), which fails to take into account the correlation between variables.

Silva et al. (2019) developed a mixtures of multivariate Poisson-lognormal (MPLN; Aitchison and Ho, 1989) distributions to cluster multivariate count measurements. In MPLN

models, the counts are modeled using a hierarchical structure such that

$$Y_{ij} | \theta_{ij} \sim \text{Poisson}(e^{\theta_{ij}}) \text{ and } \boldsymbol{\theta}_i = (\theta_{i1}, \dots, \theta_{id}) = (\log \lambda_{i1}, \dots, \log \lambda_{id}) \sim \mathcal{N}_d(\boldsymbol{\mu}, \boldsymbol{\Sigma}),$$

where $\mathcal{N}_d(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ denotes a d -dimensional Gaussian distribution with mean $\boldsymbol{\mu}$ and covariance $\boldsymbol{\Sigma}$. Due to this hierarchical structure, the MPLN model can account for over-dispersion as opposed to the traditional Poisson distribution and allows for correlation. An attractive property of MPLN distribution is that it can support both positive and negative correlation. However, the posterior of the latent variables do not have a closed form and therefore, these models came with heavy computation cost as they relied on Bayesian MCMC-based approaches, discussed further in Section 2.1. Bayesian MCMC-based approaches for models with latent variables can be computationally intensive (Blei et al., 2017) and the computational time is further compounded in model-based clustering where the expected value of the latent variable needs to be computed several times (further detailed in Section 2.2). In this paper, we propose a computationally efficient framework for clustering using mixtures of multivariate Poisson-lognormal distributions using variational Gaussian approximations. We also impose constraints on the covariance matrices of latent variable using an eigen-decomposition which results in a parsimonious family of models. Section 2.3 provides the variational EM estimation algorithm, Section 3 provides results on simulations with comparisons to a popular Poisson-based mixture approach and results on a real data set. Section concludes with some conclusions and future directions.

2 Methodology

2.1 Multivariate Poisson-lognormal distribution

In multivariate Poisson-lognormal (MPLN; Aitchison and Ho, 1989) distribution, the counts are modeled using a hierarchical structure such that

$$Y_{ij} \mid \theta_{ij} \sim \text{Poisson}(e^{\theta_{ij}}) \quad \text{and} \quad \boldsymbol{\theta}_i \sim \mathcal{N}_d(\boldsymbol{\mu}, \boldsymbol{\Sigma}),$$

where $\boldsymbol{\theta}_i = (\theta_{i1}, \dots, \theta_{id})$ and $\mathcal{N}_d(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ denotes a d -dimensional Gaussian distribution with mean $\boldsymbol{\mu}$ and covariance $\boldsymbol{\Sigma}$. The mean and expected value of the \mathbf{Y} (Aitchison and Ho, 1989; Georgescu et al., 2014) can be found using the law of iterative expectation,

$$\begin{aligned} \mathbb{E}(Y_j) &= \mathbb{E}[\mathbb{E}(Y_j \mid \theta_j)] &&= e^{\mu_j + 1/2 \times \Sigma_{jj}}, \\ \mathbb{V}(Y_j) &= \mathbb{E}[\mathbb{V}(Y_j \mid \theta_j)] + \mathbb{V}[\mathbb{E}(Y_j \mid \theta_j)] &&= \mathbb{E}(Y_j) + \mathbb{E}(Y_j)^2(e^{\Sigma_{jj}} - 1), \\ \text{cov}(Y_j, Y_k) &= \mathbb{E}[\text{cov}(Y_j, Y_k \mid \theta_j, \theta_k)] + \text{cov}[\mathbb{E}(Y_j \mid \theta_j), \mathbb{E}(Y_k \mid \theta_k)] &&= \mathbb{E}(Y_j) \mathbb{E}(Y_k)(e^{\Sigma_{jk}} - 1), \end{aligned}$$

where Y_j is the j^{th} entry of \mathbf{Y} , μ_j is the j^{th} entry of $\boldsymbol{\mu}$, Σ_{jk} is the entry in the j^{th} row and k^{th} column of the matrix $\boldsymbol{\Sigma}$ and $j \neq k$. Due to this hierarchical structure, the MPLN model can account for over-dispersion and also allows for correlation.

The marginal probability density of \mathbf{Y} can be written as:

$$f_{\mathbf{Y}}(\mathbf{y}) = \int_{\mathbb{R}^d} \left[\prod_{j=1}^d p(y_j \mid \theta_j) \right] \phi_d(\boldsymbol{\theta} \mid \boldsymbol{\mu}, \boldsymbol{\Sigma}) d\boldsymbol{\theta},$$

where θ_j and y_j are the j^{th} element of $\boldsymbol{\theta}$ and \mathbf{y} respectively, $p(\cdot)$ is the probability mass function of the Poisson distribution with mean $\lambda_j = e^{\theta_j}$ and $\phi_d(\cdot)$ is the probability density function of d -dimensional Gaussian distribution with mean $\boldsymbol{\mu}$ and covariance $\boldsymbol{\Sigma}$. Note that

the marginal distribution of Y involves multiple integrals and cannot be further simplified. Hence, Aitchison and Ho (1989) suggested that the maximum likelihood estimation of the parameters requires a mix of Newton-Raphson and steepest ascent methods. An alternate approach is to use an expectation-maximization (EM) algorithm (Dempster et al., 1977), which is an iterative approach for maximizing the likelihood when the data are incomplete or are treated as incomplete. In an MPLN distribution, the observed variables are the counts \mathbf{Y} and the missing data are the latent variables $\boldsymbol{\theta}$. The EM-algorithm comprises of two iterative steps: an E-step and an M-step. In E-step, the expected value of the complete data (i.e. observed and missing data) log-likelihood given the observed data and current parameter estimate (i.e., updates from previous iteration) is computed which is then maximized in the subsequent M-step. These step are repeated until convergence to obtain the maximum likelihood estimate of the parameters. In order to compute the expected value of the complete data log-likelihood, $\mathbb{E}(\boldsymbol{\theta} \mid \mathbf{y})$ and $\mathbb{E}(\boldsymbol{\theta}\boldsymbol{\theta}' \mid \mathbf{y})$ need to be computed. The conditional distribution of $f_{\boldsymbol{\theta}|\mathbf{Y}}(\boldsymbol{\theta} \mid \mathbf{y}, \boldsymbol{\mu}, \boldsymbol{\Sigma})$ is given by

$$f_{\boldsymbol{\theta}|\mathbf{Y}}(\boldsymbol{\theta} \mid \mathbf{y}, \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{\left[\prod_{j=1}^d p(y_j \mid \theta_j) \right] \phi_d(\boldsymbol{\theta} \mid \boldsymbol{\mu}, \boldsymbol{\Sigma})}{f_{\mathbf{Y}}(\mathbf{y} \mid \boldsymbol{\mu}, \boldsymbol{\Sigma})}.$$

However, as mentioned above, the marginal distribution of Y involves multiple integrals and cannot be further simplified. Georgescu et al. (2014) proposed a maximum likelihood estimation of the parameters for MPLN distribution in the Monte Carlo EM (MCEM) framework (Wei and Tanner, 1990); they utilized a Monte Carlo EM algorithm using importance sampling to find the empirical expected value of the complete data log-likelihood. Chagneau et al. (2011) utilized an MCMC-based approach for parameter estimation. However, it comes with a heavy computational overhead, further exacerbated as the dimensionality and sample size rise (Chagneau et al., 2011).

2.2 Mixtures of Multivariate Poisson-lognormal distributions

Silva et al. (2019) proposed a mixture of multivariate Poisson-lognormal distributions to cluster multivariate count measurements. A G -component mixture of MPLN distributions can be written as

$$f(\mathbf{y} \mid \boldsymbol{\vartheta}) = \sum_{g=1}^G \pi_g f_{\mathbf{Y}}(\mathbf{y} \mid \boldsymbol{\mu}_g, \boldsymbol{\Sigma}_g) = \sum_{g=1}^G \pi_g \int_{\mathbb{R}^d} \left[\prod_{j=1}^d p(y_{ij} \mid \theta_{ij}) \right] \phi_d(\boldsymbol{\theta}_i \mid \boldsymbol{\mu}_g, \boldsymbol{\Sigma}_g) d\boldsymbol{\theta}_i,$$

where $\boldsymbol{\vartheta}$ denotes all model parameters and $f_{\mathbf{Y}}(\mathbf{y} \mid \boldsymbol{\mu}_g, \boldsymbol{\Sigma}_g)$ denotes the distribution of the g^{th} component with parameters $\boldsymbol{\mu}_g$ and $\boldsymbol{\Sigma}_g$. In model-based clustering, an additional component membership indicator variable \mathbf{Z} is introduced which is assumed to be unknown and $Z_{ig} = 1$ if the observation i^{th} belongs to group g and $Z_{ig} = 0$ otherwise. Hence, the complete data now comprises of observed expression levels \mathbf{y} , underlying latent variable $\boldsymbol{\theta}$, and unknown group membership \mathbf{z} . Therefore, the complete-data log-likelihood is

$$l_c(\boldsymbol{\mu}, \boldsymbol{\Sigma} \mid \mathbf{y}, \boldsymbol{\theta}, \mathbf{z}) = \sum_{g=1}^G \sum_{i=1}^n z_{ig} \left[\log \phi_d(\boldsymbol{\theta}_i \mid \boldsymbol{\mu}_g, \boldsymbol{\Sigma}_g) + \sum_{j=1}^d \log p(y_{ij} \mid \theta_{ij}) \right].$$

In order to utilize the EM framework for parameter estimation for the mixtures of MPLN distributions, finding the expected value of the complete-data log-likelihood requires the conditional expectations $\mathbb{E}(Z_{ig}\boldsymbol{\theta}_i \mid \mathbf{y}_i, \boldsymbol{\mu}_g, \boldsymbol{\Sigma}_g)$ and $\mathbb{E}(Z_{ig}\boldsymbol{\theta}_i\boldsymbol{\theta}_i' \mid \mathbf{y}_i, \boldsymbol{\mu}_g, \boldsymbol{\Sigma}_g)$. Silva et al. (2019) utilized an MCMC-EM algorithm using **Stan** (Hoffman and Gelman, 2014; Stan Development Team, 2015) to compute the required conditional expectations. **Stan** uses the No-U-Turn Sampler (NUTS; Hoffman and Gelman, 2014), an adaptive variant of the Hamiltonian Monte Carlo (HMC; Neal et al., 2011) that requires no tuning parameters and can efficiently sample from a posterior distribution when the parameters are correlated. However, in a clustering context, even such efficient sampling techniques like NUTS comes with heavy computational

burden as MCMC-EM needs to be conducted at every iteration of MCMC-EM after updating the model parameters and the component indicator variable. Additionally, Bayesian techniques based on MCMC sampling algorithms also comes with increased computational overhead and possible difficulty in determining convergence as the complexity of the model increases. When dealing with such complex models, convergence can also be quite slow. This crucial aspect is partly why applications of such models have been limited to small dimensions (Georgescu et al., 2014). Additionally, when the true number of groups, i.e., the number of components of the mixture model, is unknown, the MCMC-EM algorithm must be employed for every possible number of components and the optimal number of components is chosen in conjunction with a model selection criterion which might entail extreme computational cost.

2.3 Variational approximation of mixtures of MPLN distributions

Variational approximation (Wainwright et al., 2008) is an approximate inference technique which has been very popular in machine learning. It presents an alternative parameter estimation framework for MPLN distribution by using a computationally convenient approximating density in place of a more complex but ‘true’ posterior density. Using computationally convenient Gaussian densities, complex posterior distributions are approximated by minimizing the Kullback-Leibler (KL) divergence between the true and the approximating densities and therefore, reducing the computational overhead. Several studies have shown that it delivers accurate approximations (Archambeau et al., 2007; Challis and Barber, 2013; Arridge et al., 2018). Khan et al. (2013) developed variational inference framework for several non-conjugate latent Gaussian models. Arridge et al. (2018) also developed a variational Gaussian approximation (VGA) to the posterior distribution arising from the univariate Poisson model with a Gaussian prior and derived an explicit expression for the lower bound, and

showed the existence and uniqueness of the optimal Gaussian approximation.

Here, we propose a variational Gaussian approximation to the marginal distribution of the observed variable i.e. $f(\mathbf{y})$ in the mixtures of MPLN distributions. Suppose, we have an approximating density $q(\boldsymbol{\theta})$, the marginal log-likelihood can be written as

$$\begin{aligned}\log f(\mathbf{y}) &= \int_{\mathbb{R}^d} \log f(\mathbf{y})q(\boldsymbol{\theta})d\boldsymbol{\theta} \\ &= \int_{\mathbb{R}^d} \log \frac{f(\mathbf{y}, \boldsymbol{\theta})/q(\boldsymbol{\theta})}{f(\boldsymbol{\theta} | \mathbf{y})/q(\boldsymbol{\theta})}q(\boldsymbol{\theta})d\boldsymbol{\theta} \\ &= \int_{\mathbb{R}^d} [\log f(\mathbf{y}, \boldsymbol{\theta}) - \log q(\boldsymbol{\theta})] q(\boldsymbol{\theta})d\boldsymbol{\theta} + D_{KL}(q||f) \\ &= F(q, \mathbf{y}) + D_{KL}(q||f),\end{aligned}$$

where $D_{KL}(q||f) = \int_{\mathbb{R}^d} q(\boldsymbol{\theta}) \log \frac{q(\boldsymbol{\theta})}{f(\boldsymbol{\theta}|\mathbf{y})}d\boldsymbol{\theta}$ is the Kullback-Leibler (KL) divergence between $f(\boldsymbol{\theta} | \mathbf{y})$ and approximating distribution $q(\boldsymbol{\theta})$, and $F(q, \mathbf{y}) = \int_{\mathbb{R}^d} [\log f(\mathbf{y}, \boldsymbol{\theta}) - \log q(\boldsymbol{\theta})] q(\boldsymbol{\theta})d\boldsymbol{\theta}$ is our evidence lower bound (ELBO). To minimize the KL divergence, we maximize our ELBO. In VGA, $q(\boldsymbol{\theta})$ is assumed to be a Gaussian distribution. Assuming $q(\boldsymbol{\theta}) = \mathcal{N}(\mathbf{m}, \mathbf{S})$, the lower bound for the log $f(\mathbf{y})$ becomes

$$\begin{aligned}F(q, \mathbf{y}) &= -\frac{1}{2}(\mathbf{m} - \boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{m} - \boldsymbol{\mu}) - \frac{1}{2} \text{tr}(\boldsymbol{\Sigma}^{-1}\mathbf{S}) + \frac{1}{2} \log |\mathbf{S}| - \frac{1}{2} \log |\boldsymbol{\Sigma}| - \frac{d}{2} + \mathbf{m}'\mathbf{y} \\ &\quad - \sum_{j=1}^d \left(e^{m_j + \frac{1}{2}S_{jj}} + \log(y_j!) \right).\end{aligned}$$

This lower bound is strictly jointly concave with respect to the mean (\mathbf{m}) and variance (\mathbf{S}) of the approximating distribution and hence, similar to Arridge et al. (2018), parameter estimation can be obtained via Newton's method and fixed-point method. This alleviates the need of the use of MCMC-based approach to get samples from the posterior distribution of the latent variable $\boldsymbol{\theta}$ and the computational burden that comes along with such

approaches. Additionally, the EM framework using variational Gaussian approximation algorithm is monotonic as opposed to MCMC-based EM. A similar variational framework for MPLN distribution was recently proposed by Chiquet et al. (2019) for network models.

The complete data log-likelihood of the mixtures of MPLN distributions can be written as:

$$\begin{aligned} l_c(\boldsymbol{\vartheta} | \mathbf{y}) &= \sum_{g=1}^G \sum_{i=1}^n z_{ig} \log \pi_g + \sum_{g=1}^G \sum_{i=1}^n z_{ig} \log f(\mathbf{y}_i | \boldsymbol{\mu}_g, \boldsymbol{\Sigma}_g) \\ &= \sum_{g=1}^G \sum_{i=1}^n z_{ig} \log \pi_g + \sum_{g=1}^G \sum_{i=1}^n z_{ig} \left[\int_{\mathbb{R}^d} [\log f(\mathbf{y}, \boldsymbol{\theta}_{ig}) - \log q(\boldsymbol{\theta}_{ig})] q(\boldsymbol{\theta}_{ig}) d\boldsymbol{\theta}_{ig} + D_{KL}(q_{ig} \| f_{ig}) \right] \end{aligned}$$

where $D_{KL}(q_{ig} \| f_{ig}) = \int_{\mathbb{R}^d} q(\boldsymbol{\theta}_{ig}) \log \frac{q(\boldsymbol{\theta}_{ig})}{f(\boldsymbol{\theta}_i | \mathbf{y}_i, Z_{ig}=1)} d\boldsymbol{\theta}_{ig}$ is the Kullback-Leibler (KL) divergence between $f(\boldsymbol{\theta}_i | \mathbf{y}_i, Z_{ig} = 1)$ and approximating distribution $q(\boldsymbol{\theta}_{ig})$. Assuming $q(\boldsymbol{\theta}_{ig}) = \mathcal{N}(\mathbf{m}_{ig}, \mathbf{S}_{ig})$, the log-likelihood becomes:

$$l_c(\boldsymbol{\vartheta} | \mathbf{y}) = \sum_{g=1}^G \sum_{i=1}^n z_{ig} \log \pi_g + \sum_{g=1}^G \sum_{i=1}^n z_{ig} [F(q_{ig}, \mathbf{y}_i) + D_{KL}(q_{ig} \| f_{ig})],$$

where the ELBO for each observation \mathbf{y}_i is

$$\begin{aligned} F(q_{ig}, \mathbf{y}_i) &= \frac{1}{2} \log |\mathbf{S}_{ig}| - \frac{1}{2} (\mathbf{m}_{ig} - \boldsymbol{\mu}_g)' \boldsymbol{\Sigma}_g^{-1} (\mathbf{m}_{ig} - \boldsymbol{\mu}_g) - \text{tr}(\boldsymbol{\Sigma}_g^{-1} \mathbf{S}_{ig}) - \frac{1}{2} \log |\boldsymbol{\Sigma}_g| - \frac{d}{2} + \mathbf{m}'_{ig} \mathbf{y}_i \\ &\quad - \sum_{j=1}^d \left(e^{m_{igj} + \frac{1}{2} S_{ig,jj}} + \log(y_{ij}!) \right), \end{aligned}$$

where m_{igj} is the j^{th} element of the \mathbf{m}_{ig} and $S_{ig,jj}$ is the j^{th} diagonal element of the matrix \mathbf{S}_{ig} .

The variational parameters that maximize the ELBO will minimize the KL divergence between the true posterior and the approximating density.

Parameter estimation can be done in an iterative EM-type approach such that the following steps are iterated.

1. Conditional on the variational parameters \mathbf{m}_{ig} , \mathbf{S}_{ig} and on $\boldsymbol{\mu}_g$ and $\boldsymbol{\Sigma}_g$, the $\mathbb{E}(Z_{ig})$ is computed. Given $\boldsymbol{\mu}_g$ and $\boldsymbol{\Sigma}_g$,

$$\mathbb{E}(Z_{ig} | \mathbf{y}_i) = \frac{\pi_g f(\mathbf{y} | \boldsymbol{\mu}_g, \boldsymbol{\Sigma}_g)}{\sum_{h=1}^G \pi_h f(\mathbf{y} | \boldsymbol{\mu}_h, \boldsymbol{\Sigma}_h)}.$$

Note that this involves the marginal distribution of Y which is difficult to compute. Hence, we use an approximation of $\mathbb{E}(Z_{ig})$ where we replace the marginal density of the exponent of ELBO such that

$$\hat{Z}_{ig} \stackrel{\text{def}}{=} \frac{\pi_g \exp [F(q_{ig}, \mathbf{y}_i)]}{\sum_{h=1}^G \pi_h \exp [F(q_{ih}, \mathbf{y}_i)]}.$$

This approximation is computationally convenient and a similar framework was utilized by Tang et al. (2015); Gollini and Murphy (2014). This approximation works well in simulation studies and real data analysis.

2. Given \hat{Z}_{ig} , variational parameters \mathbf{m}_{ig} and \mathbf{S}_{ig} is updated conditional on $\boldsymbol{\mu}_g$ and $\boldsymbol{\Sigma}_g$. The lower bound is strictly jointly concave with respect to \mathbf{m}_{ig} and \mathbf{S}_{ig} of the approximating distribution. Therefore, we can update \mathbf{m}_{ig} and \mathbf{S}_{ig} as following:

- (a) fixed-point method for updating \mathbf{S}_{ig} is

$$\mathbf{S}_{ig}^{(t+1)} = \left\{ \boldsymbol{\Sigma}_g^{-1} + \mathbf{I} \odot \mathbf{exp} \left[\mathbf{m}_{ig}^{(t)} + \frac{1}{2} \text{diag} \left(\mathbf{S}_{ig}^{(t)} \right) \right] \mathbf{1}'_d \right\}^{-1}$$

where the vector function $\mathbf{exp}[\mathbf{a}] = (e^{a_1}, \dots, e^{a_d})'$ is a vector of exponential each element of the d -dimensional vector \mathbf{a} , $\text{diag}(\mathbf{S}) = (\mathbf{S}_{11} \dots, \mathbf{S}_{dd})$ puts the diagonal elements of the $d \times d$ matrix \mathbf{S} into a d -dimensional vector, \odot the Hadmard

product and $\mathbf{1}_d$ is a d -dimensional vector of ones ;

(b) Newton's method to update \mathbf{m}_{ig} is

$$\mathbf{m}_{ig}^{(t+1)} = \mathbf{m}_{ig}^{(t)} - \mathbf{S}_{ig}^{(t+1)} \left\{ \exp \left[\mathbf{m}_{ig}^{(t)} + \frac{1}{2} \text{diag} \left(\mathbf{S}_{ig}^{(t+1)} \right) \right] + \Sigma_g^{-1} \left(\mathbf{m}_{ig}^{(t)} - \boldsymbol{\mu}_g \right) - \mathbf{y}_i \right\}; \quad \text{and}$$

3. given \hat{z}_{ig} and the variational parameters \mathbf{m}_{ig} and \mathbf{S}_{ig} , the parameters $\boldsymbol{\pi}$, $\boldsymbol{\mu}_g$ and Σ_g .

$$\begin{aligned} \pi_g &= \frac{\sum_{i=1}^n \hat{Z}_{ig}}{n}, \\ \boldsymbol{\mu}_g &= \frac{\sum_{i=1}^n \hat{Z}_{ig} \mathbf{m}_{ig}}{\sum_{i=1}^n \hat{Z}_{ig}}, \quad \text{and} \\ \Sigma_g &= \frac{\sum_{i=1}^n \hat{Z}_{ig} (\mathbf{m}_{ig} - \boldsymbol{\mu}_g) (\mathbf{m}_{ig} - \boldsymbol{\mu}_g)'}{\sum_{i=1}^n \hat{Z}_{ig}} - \frac{\sum_{i=1}^n \hat{Z}_{ig} \mathbf{S}_{ig}}{\sum_{i=1}^n \hat{Z}_{ig}}. \end{aligned}$$

2.4 Parsimonious family of models

For the mixtures of MPLN distribution, the number of free parameters in the covariance matrices of the latent variable is $Gd(d+1)/2$, i.e., the number of parameters increases quadratically with d . Banfield and Raftery (1993) proposed an eigen-decomposition of the covariance matrices of mixtures of Gaussian distributions such that $\Sigma_g = \lambda_g \mathbf{D}_g \mathbf{A}_g \mathbf{D}_g'$ where \mathbf{D}_g is the matrix of eigenvectors and \mathbf{A}_g is a diagonal matrix proportional to the eigenvalues of Σ_g , such that $|\mathbf{A}_g| = 1$, and λ_g is the associated constant of proportionality. For a Gaussian distribution, this decomposition results in a geometric interpretation of the constraints such that λ_g controls the cluster volume, \mathbf{A}_g controls the cluster shape, and \mathbf{D}_g controls the cluster orientation. Celeux and Govaert (1995) imposed constraints on these parameters of the covariance matrix to be equal or different among groups resulting in a family of 14 models known as Gaussian parsimonious clustering models (GPCM). Parameter estimation in an EM framework for 12 of these 14 GPCM family and MM framework of

Browne and McNicholas (2014) for the remaining two models is available in the R package `mclust` (Scrucca et al., 2016). Here, we propose an eigen-decomposition of the component-specific covariance matrices of the latent variable $\boldsymbol{\theta}$ to introduce parsimony. The geometric interpretation does not hold for observed data and therefore, we only focused on the following subset of the GPCM family of model given in Table 1.

Table 1: Geometric interpretation of the subset of the models considered in our study obtained via eigen-decomposition of $\boldsymbol{\Sigma}_g$.

| Model (Covariance) | Volume | Shape | Orientation | Parameters |
|---|----------|-----------|-------------|----------------------|
| EII (λI) | Equal | Spherical | - | 1 |
| VII ($\lambda_g I$) | Variable | Spherical | - | G |
| E EI ($\lambda \mathbf{A}$) | Equal | Equal | Ax-Alg | d |
| V VI ($\lambda_g \mathbf{A}_g$) | Variable | Variable | Ax-Alg | dG |
| EEE ($\lambda \mathbf{D} \mathbf{A} \mathbf{D}^T$) | Equal | Equal | Equal | $d(d+1)/2$ |
| V VE ($\lambda_g \mathbf{D} \mathbf{A}_g \mathbf{D}^T$) | Variable | Variable | Equal | $d(d+1)/2 + (G-1)d$ |
| E EV ($\lambda \mathbf{D}_g \mathbf{A} \mathbf{D}_g^T$) | Equal | Equal | Variable | $Gd(d+1)/2 - (G-1)d$ |
| V VV ($\lambda_g \mathbf{D}_g \mathbf{A}_g \mathbf{D}_g^T$) | Variable | Variable | Variable | $Gd(d+1)/2$ |

Note that within a component $\text{cov}(Y_j, Y_k) = \mathbb{E}(Y_j) \mathbb{E}(Y_k)(e^{\Sigma_{jk}} - 1)$, therefore, independence among the latent variable $\boldsymbol{\theta}$ will result in independence among the observed variable \mathbf{Y} . We focused on EII, VII, EEI, and VVI to recover diagonal covariance structure when variables are independent. Additionally, to introduce some parsimony compared to VVV model, we also focused some additional non-diagonal constraints such as EEE, VVE and EEV. Given \mathbf{m}_{ig} and \mathbf{S}_{ig} and using the sample covariance matrix of \mathbf{m}_g as $\hat{\boldsymbol{\Sigma}}_{\text{sample}} = \frac{\sum_{i=1}^n \hat{Z}_{ig} (\mathbf{m}_{ig} - \boldsymbol{\mu}_g)(\mathbf{m}_{ig} - \boldsymbol{\mu}_g)'}{\sum_{i=1}^n \hat{Z}_{ig}} - \frac{\sum_{i=1}^n \hat{Z}_{ig} \mathbf{S}_{ig}}{\sum_{i=1}^n \hat{Z}_{ig}}$ the parameter estimates are analogous to their Gaussian counterpart. See Celeux and Govaert (1995); Browne and McNicholas (2014) for details.

2.5 Initialization, Model Selection, and Performance assessment

EM algorithm relies heavily on the initial values (Biernacki et al., 2003) and hence, good starting values are crucial. While initialization of the clusters using k -means is one of the widely used approach for symmetric distributions, it may not be appropriate for our over-dispersed count datasets as k -means initialization is equivalent to fitting a Gaussian mixture models with spherical clusters which will not be ideal for these over-dispersed multivariate counts. Therefore, we utilized small EM initialization for our analysis. For each G , the initialization “EII” model of consist of using 20 different random partitions of datasets into G clusters and running 20 iterations of our variational EM algorithm. The parameters associated with the largest log-likelihood among the 20 random partitions was then used to initialize all eight models which is then run until convergence. Convergence of the algorithm for these models is determined using a modified Aitken acceleration criterion. where at iteration m . Convergence is based on Aitken acceleration (Aitken, 1926) which is considered to satisfied when $l_{\infty}^{(m+1)} - l_{\infty}^{(m)}$ is positive and smaller than some ϵ , where $l^{(m)}$ is the value of the log-likelihood and $l_{\infty}^{(m+1)}$ is an asymptotic estimate of the log-likelihood given by

$$l_{\infty}^{(m+1)} = l^{(m)} + \frac{l^{(m+1)} - l^{(m-1)}}{1 - a^{(m)}} \quad \text{where} \quad a^{(m)} = \frac{l^{(m+1)} - l^{(m)}}{l^{(m)} - l^{(m-1)}}$$

(Böhning et al., 1994). Here, we used $\epsilon = 0.001$.

Typically, the number of component is unknown in cluster analysis. In our case, the covariance structure that provides the best fit is also unknown. Hence, we fit the model with all eight covariance structure to a range of G and the Bayesian information criteria (BIC) (Schwarz et al., 1978) is used to assess the goodness of fit of each fitted model and select the number of G and corresponding to the corresponding covariance structure that has the best

fitness to the data. The BIC of a fitted model is computed by

$$\text{BIC}(\hat{\boldsymbol{\vartheta}}) = -2\ell(\hat{\boldsymbol{\vartheta}}) + \psi \log(n),$$

where $\ell(\boldsymbol{\vartheta})$ is the observed data log-likelihood function here, n is the sample size, and ψ is the number of free parameters in the model. Note that here we compute approximation of BIC using *ELBO* (Chen et al., 2018). We will refer to the approximation of BIC as BIC in the paper. Additionally, the variational parameters can be regarded as proxies for the latent variables and therefore, we do not penalize for variational parameters. The model that has the smallest BIC value is selected and the number of underlying groups is therefore estimated. Performance assessment clustering performance was done in using the Adjusted Rand Index (ARI; Hubert and Arabie, 1985): 1 indicates perfect agreement between the true and predicted classification and the expected value of the ARI under random classification is zero. A value < 0 indicate a classification that is worse than would be expected under random assignment. For simulation studies where we know the true class labels, we also compared the performance of our proposed model to a Poisson mixture model implemented in the R package `HTSCluster` (Rau et al., 2015). Note that it is a clustering algorithm designed for clustering high-throughput transcriptome sequencing (HTS) data and therefore has a built-in normalization option which could not be turned off. Therefore, we set the normalization option to “TMM” for our analysis.

3 Analyses

3.1 Simulation Study 1

We generated 100 three dimensional datasets of size $N = 2000$ with three components with mixing proportions $\boldsymbol{\pi} = (0.2, 0.5, 0.3)$, known mean and a completely unconstrained VVV covariance structure (see Table 2 for the values used to generate the data). We ran our model for $G = 1 : 4$ and all eight covariance structures. In 100 out of the 100 datasets, a three component VVV model was selected with an average ARI of 0.99 and standard deviation (sd) of 0.003. Summary of the parameter estimates is provided in Table 2.

Table 2: Summary of the average and standard errors (SE) of the estimated parameters from the 100 datasets where a $G = 3$ component VVV model was selected in Simulation Study 1.

| | True Parameters | Average of Estimated Parameters | Standard errors |
|----------------------|-----------------|---------------------------------|--------------------|
| $\boldsymbol{\mu}_1$ | (6, 3, 3) | (6.00, 3.00, 3.00) | (0.03, 0.04, 0.04) |
| $\boldsymbol{\mu}_2$ | (3, 5, 3) | (3.00, 5.00, 3.00) | (0.02, 0.02, 0.02) |
| $\boldsymbol{\mu}_3$ | (5, 3, 5) | (5.00, 3.00, 5.00) | (0.02, 0.03, 0.02) |

$$\boldsymbol{\Sigma}_1 = \begin{bmatrix} 0.30 & 0.15 & 0.20 \\ 0.15 & 0.40 & 0.30 \\ 0.20 & 0.30 & 0.40 \end{bmatrix}, \quad \hat{\boldsymbol{\Sigma}}_1 = \begin{bmatrix} 0.30 & 0.15 & 0.20 \\ 0.15 & 0.40 & 0.30 \\ 0.20 & 0.30 & 0.40 \end{bmatrix}, \quad \text{SE}(\hat{\boldsymbol{\Sigma}}_1) = \begin{bmatrix} 0.02 & 0.02 & 0.02 \\ 0.02 & 0.03 & 0.03 \\ 0.02 & 0.03 & 0.03 \end{bmatrix}$$

$$\boldsymbol{\Sigma}_2 = \begin{bmatrix} 0.30 & 0.15 & 0.20 \\ 0.15 & 0.40 & 0.30 \\ 0.20 & 0.30 & 0.40 \end{bmatrix}, \quad \hat{\boldsymbol{\Sigma}}_2 = \begin{bmatrix} 0.30 & 0.15 & 0.20 \\ 0.15 & 0.40 & 0.30 \\ 0.20 & 0.30 & 0.40 \end{bmatrix}, \quad \text{SE}(\hat{\boldsymbol{\Sigma}}_2) = \begin{bmatrix} 0.02 & 0.01 & 0.01 \\ 0.01 & 0.02 & 0.02 \\ 0.01 & 0.02 & 0.02 \end{bmatrix}$$

$$\boldsymbol{\Sigma}_3 = \begin{bmatrix} 0.20 & -0.15 & -0.10 \\ -0.15 & 0.40 & -0.10 \\ -0.10 & -0.10 & 0.20 \end{bmatrix}, \quad \hat{\boldsymbol{\Sigma}}_3 = \begin{bmatrix} 0.20 & -0.15 & -0.10 \\ -0.15 & 0.39 & -0.10 \\ -0.10 & -0.10 & 0.20 \end{bmatrix}, \quad \text{SE}(\hat{\boldsymbol{\Sigma}}_3) = \begin{bmatrix} 0.01 & 0.01 & 0.01 \\ 0.01 & 0.02 & 0.01 \\ 0.01 & 0.01 & 0.01 \end{bmatrix}$$

Similarly, we also ran the Poisson mixture model in the R package `HTScluster` for $G =$

1 : 4 and selected the best model with BIC. In 100 out of the 100 datasets, it overestimated the number of components resulting in a four component model was selected with an average ARI of 0.75 and a standard deviation of 0.11.

3.2 Simulation Study 2

Here, we generated 100 six dimensional datasets of size $N = 500$ with two components with mixing proportions $\boldsymbol{\pi} = (0.59, 0.41)$, known mean and a completely constrained EII (λ) covariance structure (see Table 3 for the values used to generate the data). We ran our model for $G = 1 : 3$ and all eight covariance structures. In 99 out of 100 datasets, a two component EII model was selected with an average ARI of 1.00 and a standard deviation of 0.00 and in one of the 100 dataset, a two component VII model was selected. Note, that VII is still imposes an isotropic (spherical) structure on $\boldsymbol{\Sigma}$ but it allows $\boldsymbol{\Sigma}$ to be vary among groups. In our case, with $G = 2$, it just added one additional parameter so the penalty for selecting a slightly complicated model was very small. Summary of the parameter estimates of the 99 out of the 100 datasets where the correct model is selected is provided in 3.

Table 3: Summary of the average and standard errors (SE) of the estimated parameters from the 99 out the 100 datasets where a $G = 2$ component EII model was selected in Simulation Study 2.

| g | $\boldsymbol{\mu}_g$ | | $\hat{\boldsymbol{\mu}}_g$ |
|------|--------------------------|---------|---------------------------------------|
| 1 | (5, 6, 5, 5, 5, 6) | Average | (5.00, 6.00, 5.01, 5.00, 5.01, 6.01) |
| | | SE | (0.05, 0.06, 0.05, 0.05, 0.05, 0.06) |
| 2 | (2.5, 3, 2.5, 3, 3, 2.5) | Average | (2.49, 3.01, 2.50, 3.00, 3.00, 2.49) |
| | | SE | (0.07, 0.07, 0.07, 0.07, 0.07, 0.07) |
| g | λ | | $\hat{\lambda}$ |
| 1, 2 | 1 | Average | 0.99 |
| | | SE | 0.03 |

Similarly, we also ran the Poisson mixture model in the R package `HTSCluster` for $G = 1 :$

3 and selected the best model with BIC. In 100 out of the 100 datasets, it overestimated the number of components resulting in a three component model was selected with an average ARI of 0.04 and a standard deviation of 0.02.

3.3 Simulation studies on data generated from other models

3.3.1 Simulation Study 3: Datasets generated using mixtures of negative binomial distributions

Here, we generated 100 six dimensional datasets of size $N = 2000$ from a two component mixtures of independent negative binomial distributions with mixing proportions $\boldsymbol{\pi} = (0.79, 0.21)$. We ran all our eight models for $G = 1, \dots, 4$ and model selection was done using BIC. In 100 out of 100 datasets, a two EII component model was selected with perfect classification. Note that, it always selected a diagonal covariance structure where the covariance is 0. Table 4 shows that our model was able to recover the true mean and variance of the data generated from the mixtures of negative binomial distributions.

Similarly, we also ran the Poisson mixture model in the R package `HTSCluster` for $G = 1 : 4$ and selected the best model with BIC. In 100 out of the 100 datasets, it overestimated the number of components resulting in a four component model was selected with an average ARI of 0.29 and a standard deviation of 0.01.

3.3.2 Simulation Study 4: Datasets generated using mixtures of independent Poisson distributions

Here, we used 100 four dimensional datasets generated using a mixtures of independent Poisson distributions with mixing proportions $\boldsymbol{\pi} = (0.59, 0.41)$. We ran all our eight models for $G = 1, \dots, 4$ and model selection was done using BIC. In 100 out of the 100 datasets, a

Table 4: Summary of the average and standard errors (SE) of the estimated means for the 100 datasets in Simulation Study 3 where the data was generated from a mixture of independent negative binomial distributions.

| g=1 | |
|--------------------------------|--|
| True Mean | (1000, 500, 1000, 500, 1000 , 500) |
| Average of Estimated Means | (1000.28, 500.24, 999.84, 500.09, 999.75, 500.25) |
| SE of Estimated Means | (2.66, 1.50, 2.31, 1.43, 2.91, 1.36) |
| True Variance | (11000, 3000, 11000, 3000, 11000, 3000) |
| Average of Estimated Variances | (11045.99, 3012.67, 11036.90, 30011.06, 11034.85, 3012.86) |
| SE of Estimated Variances | (147.40, 37.25, 148.18, 37.31, 152.14, 39.56) |
| g=2 | |
| True Mean | (500, 1000, 500, 1000, 500, 500) |
| Average of Estimated Means | (500.10, 999.81, 500.00, 1000.53, 499.91, 499.75) |
| SE of Estimated Means | (2.44, 5.38, 5.37, 2.71, 2.39, 2.68) |
| True Variance | (3000, 11000, 3000,11000, 3000, 3000) |
| Average of Estimated Variances | (3011.23, 11036.40, 3010.06, 11051.85, 3009.10, 3007.34) |
| SE of Estimated Variances | (43.19, 182.23, 45.03, 193.22, 43.65, 43.70) |

two component EII model was selected with perfect classification for all datasets. Note that here, the diagonal structure for Σ was always selected as the datasets were generated from mixtures of independent Poisson distributions. Table 5 shows that our model was able to recover the true mean the data generated from the mixtures of Poisson distributions very well but it overestimated the variable slightly which would be expected as MPLN imposes a structure such that $\mathbb{V}(Y) > \mathbb{E}(Y)$ but for Poisson distribution $\mathbb{V}(Y) = \mathbb{E}(Y)$.

Similarly, we also ran the Poisson mixture model in the R package `HTSCluster` for $G = 1 : 4$ and selected the best model with BIC. In 100 out of the 100 datasets, it selected the correct $G = 2$ model with an ARI of 1.

Table 5: Summary of the average and standard errors (SE) of the estimated means for the 100 datasets in Simulation Study 4 where the data was generated using a mixtures of independent Poisson distributions.

| g=1 | |
|--------------------------------|---------------------------------------|
| True Mean | (1000, 1500, 1500, 1000) |
| Average of Estimated Means | (999.98, 1500.16, 1500.27, 1000.09) |
| SE of Estimated Means | (1.31, 1.84, 1.77, 1.40) |
| True Variance | (1000, 1500, 1500, 1000) |
| Average of Estimated Variances | (1027.97, 1563.15, 1563.27, 1028.09) |
| SE of Estimated Variances | (6.21, 14.37, 14.28, 6.52) |
| g=2 | |
| True Mean | (1000, 1000, 1000, 1500) |
| Average of Estimated Means | (999.94, 1000.04, 999.96, 1499.97) |
| SE of Estimated Means | (1.39, 1.32, 1.38, 1.79) |
| True Variance | (1000, 1000, 1000, 1500) |
| Average of Estimated Variances | (1027.93, 1028.03, 1027.95, 1562.95) |
| SE of Estimated Variances | (6.50, 6.13, 6.57, 14.51) |

3.4 Real data analysis

3.5 Chronic Kidney Disease Data Set

Here, we used the Chronic Kidney Disease (CKD) Dataset available via UCI Machine Learning repository. The dataset comprises of 25 attributes regarding various blood measurements and various categorical measurements pertaining disease status on 400 individuals. Here, we focus on three discrete count measurements: **bgr**: blood glucose random; **wbcc**: white blood cell count; and **pcv**packed cell volume with the aim of classifying the disease status of the patient for CKD. Information on the status of CKD is available in the variable **class**. Before the analysis, observations with missing values on our 4 variables (**bgr**, **wbcc**, **pcv** and **class**) were removed resulting in 261 observations. We ran our algorithm for $G = 1, \dots, 4$ and BIC selected a two component "VVI" model. As seen in Table 6, the first cluster

comprised of mostly CKD patients and the second cluster comprised mostly of non-CKD patients.

Table 6: Cross tabulation of the clusters obtained by mixtures of MPLN and HTScluster against the CKD status.

| | Mix. MPLN | | HTScluster | | | |
|---------|-----------|-----|------------|----|----|----|
| | 1 | 2 | 1 | 2 | 3 | 4 |
| CKD | 95 | 29 | 28 | 18 | 29 | 49 |
| Not CKD | 0 | 135 | 24 | 0 | 16 | 95 |

Visualization of the true cluster structure and predicted cluster structure are shown in Figure 1. We also ran HTScluster on the dataset for $G = 1, \dots, 4$ and it selected a four component model. However, the cluster structure appears to be random (see Figure 1 and Table 6).




Figure 1: Scatter plot matrix of the chronic kidney dataset where the colors/symbols represents the CKD status on the left, colors/symbols representing the cluster structure recovered by MPLN in the middle, and colors/symbols representing the cluster structure recovered by HTScluster on the right.

4 Conclusions

Here, we propose an efficient framework for parameter estimation for clustering using mixtures of multivariate Poisson-lognormal distributions that utilizes a variational Gaussian

approximation. The variational Gaussian approximation provides a fast and deterministic framework for estimating the model parameters and alleviates the computational overhead due to MCMC-EM. Through eigen-decomposition of the component’s covariance matrices of the latent variable and imposition of constraints, a family of parsimonious models is proposed. Through simulation studies, we show that the proposed models provide competitive performance, the parameters were very close to the true parameters (when the correct model was chosen), and close to perfect classification was obtained using the model selected by BIC. Additionally, we also generate data from mixtures of univariate Poisson distributions and mixtures of negative binomial distributions and show that our proposed model can recover the true mean of the models fairly well. When datasets were generated from a Poisson model which assumes $\mathbb{V}(Y) = \mathbb{E}(Y)$, the variance was overestimated slightly as MPLN imposes the structure that $\mathbb{V}(Y) > \mathbb{E}(Y)$. In the case where the datasets were generated from a mixture of univariate negative binomial distributions, the variance estimation was much closer as both MPLN and negative binomial impose the structure that $\mathbb{V}(Y) > \mathbb{E}(Y)$. Additionally, when data are generated from mixtures of univariate distributions, our approach selects a model with a diagonal covariance structure where variables are independent.

While our proposed work is motivated by modern biological data such as RNA-seq data, here we only focus on a general framework that is applicable to any multivariate count data. For biological data, extensions of these approaches may require some structure that needs to be imposed on the datasets to normalize for library specific factors such as library size and transcript specific factors such as gene length. Although the use of variational Gaussian approximations alleviates some of the challenges of MCMC-EM, the algorithm can still be prone to computational issues encountered by any traditional EM algorithm such as convergence to local maxima, singularities, etc. Furthermore, a systematic evaluation for the speed and accuracy of variational EM against the MCMC-EM would be very valuable and

is ongoing. In our work, BIC seems to perform fairly well, however, some future work will focus on investigation of different model-selection criteria for MPLN mixture models and for discrete data in general.

References

- Aitchison, J. and C. Ho (1989). The multivariate Poisson-log normal distribution. *Biometrika* 76(4), 643–653.
- Aitken, A. C. (1926). A series formula for the roots of algebraic and transcendental equations. *Proceedings of the Royal Society of Edinburgh* 45, 14–22.
- Anders, S. and W. Huber (2010). Differential expression analysis for sequence count data. *Nature Precedings*, 1–1.
- Archambeau, C., D. Cornford, M. Opper, and J. Shawe-Taylor (2007). Gaussian process approximations of stochastic differential equations. *Journal of Machine Learning Research* 1, 1–16.
- Arridge, S. R., K. Ito, B. Jin, and C. Zhang (2018). Variational gaussian approximation for poisson data. *Inverse Problems* 34(2), 025005.
- Banfield, J. D. and A. E. Raftery (1993). Model-based Gaussian and non-Gaussian clustering. *Biometrics* 49(3), 803–821.
- Biernacki, C., G. Celeux, and G. Govaert (2003). Choosing starting values for the em algorithm for getting the highest likelihood in multivariate gaussian mixture models. *Computational Statistics & Data Analysis* 41(3-4), 561–575.

- Blei, D. M., A. Kucukelbir, and J. D. McAuliffe (2017). Variational inference: A review for statisticians. *Journal of the American statistical Association* 112(518), 859–877.
- Böhning, D., E. Dietz, R. Schaub, P. Schlattmann, and B. Lindsay (1994). The distribution of the likelihood ratio for mixtures of densities from the one-parameter exponential family. *Annals of the Institute of Statistical Mathematics* 46, 373–388.
- Bouveyron, C. and C. Brunet (2012). Simultaneous model-based clustering and visualization in the fisher discriminative subspace. *Statistics and Computing* 22(1), 301–324.
- Brijs, T., D. Karlis, G. Swinnen, K. Vanhoof, G. Wets, and P. Manchanda (2004). A multivariate poisson mixture model for marketing applications. *Statistica Neerlandica* 58(3), 322–348.
- Browne, R. P. and P. D. McNicholas (2014). Estimating common principal components in high dimensions. *Advances in Data Analysis and Classification* 8(2), 217–226.
- Browne, R. P. and P. D. McNicholas (2015). A mixture of generalized hyperbolic distributions. *Canadian Journal of Statistics* 43(2), 176–198.
- Bullard, J. H., E. Purdom, K. D. Hansen, and S. Dudoit (2010). Evaluation of statistical methods for normalization and differential expression in mrna-seq experiments. *BMC Bioinformatics* 11(1), 94.
- Bulmer, M. (1974). On fitting the poisson lognormal distribution to species-abundance data. *Biometrics*, 101–110.
- Campbell, J. (1934). The poisson correlation function. *Proceedings of the Edinburgh Mathematical Society* 4(1), 18–26.

- Celeux, G. and G. Govaert (1995). Gaussian parsimonious clustering models. *Pattern Recognition* 28, 781–793.
- Chagneau, P., F. Mortier, N. Picard, and J.-N. Bacro (2011). A hierarchical bayesian model for spatial prediction of multivariate non-gaussian random fields. *Biometrics* 67(1), 97–105.
- Challis, E. and D. Barber (2013). Gaussian kullback-leibler approximate inference. *The Journal of Machine Learning Research* 14(1), 2239–2286.
- Chen, Y.-C., Y. S. Wang, E. A. Erosheva, et al. (2018). On the use of bootstrap with variational inference: Theory, interpretation, and a two-sample test example. *The Annals of Applied Statistics* 12(2), 846–876.
- Chiquet, J., M. Mariadassou, and S. Robin (2019). Variational inference for sparse network reconstruction from count data. In *Proceedings of the 36th International Conference on Machine Learning*, Volume 97 of *Proceedings of Machine Learning Research*.
- Collings, B. J. and B. H. Margolin (1985). Testing goodness of fit for the poisson assumption when observations are not identically distributed. *Journal of the American Statistical Association* 80(390), 411–418.
- Dang, U. J., R. P. Browne, and P. D. McNicholas (2015). Mixtures of multivariate power exponential distributions. *Biometrics* 71(4), 1081–1089.
- de Souto, M. C., I. G. Costa, D. S. de Araujo, T. B. Ludermir, and A. Schliep (2008). Clustering cancer gene expression data: a comparative study. *BMC Bioinformatics* 9(1), 1.

- Dempster, A. P., N. M. Laird, and D. B. Rubin (1977). Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society: Series B* 39(1), 1–38.
- Dong, K., H. Zhao, T. Tong, and X. Wan (2016). Nbllda: negative binomial linear discriminant analysis for rna-seq data. *BMC Bioinformatics* 17(1), 369.
- Georgescu, V., N. Desassis, S. Soubeyrand, A. Kretzschmar, and R. Senoussi (2014). An automated MCEM algorithm for hierarchical models with multivariate and multitype response variables. *Communications in Statistics-Theory and Methods* 43(17), 3698–3719.
- Gollini, I. and T. B. Murphy (2014). Mixture of latent trait analyzers for model-based clustering of categorical data. *Statistics and Computing* 24(4), 569–588.
- Greenwood, M. and G. U. Yule (1920). An inquiry into the nature of frequency distributions representative of multiple happenings with particular reference to the occurrence of multiple attacks of disease or of repeated accidents. *Journal of the Royal statistical society* 83(2), 255–279.
- Gurland, J. (1958). A generalized class of contagious distributions. *Biometrics* 14(2), 229–249.
- Hoffman, M. D. and A. Gelman (2014). The no-u-turn sampler: adaptively setting path lengths in hamiltonian monte carlo. *Journal of Machine Learning Research* 15(1), 1593–1623.
- Holla, M. (1967). On a poisson-inverse gaussian distribution. *Metrika* 11(1), 115–121.
- Hubert, L. and P. Arabie (1985). Comparing partitions. *Journal of Classification* 2, 193–218.

- Inouye, D. I., E. Yang, G. I. Allen, and P. Ravikumar (2017). A review of multivariate distributions for count data derived from the poisson distribution. *Wiley Interdisciplinary Reviews: Computational Statistics* 9(3), e1398.
- Karlis, D. and L. Meligkotsidou (2007). Finite mixtures of multivariate poisson distributions with application. *Journal of Statistical Planning and Inference* 137(6), 1942–1960.
- Karlis, D. and E. Xekalaki (2005). Mixed poisson distributions. *International Statistical Review* 73(1), 35–58.
- Khan, M. E., A. Aravkin, M. Friedlander, and M. Seeger (2013). Fast dual variational inference for non-conjugate latent gaussian models. In *International Conference on Machine Learning*, pp. 951–959.
- Kosmidis, I. and D. Karlis (2016). Model-based clustering using copulas with applications. *Statistics and Computing* 26(5), 1079–1099.
- Love, M. I., W. Huber, and S. Anders (2014). Moderated estimation of fold change and dispersion for rna-seq data with deseq2. *Genome Biology* 15(12), 550.
- Marioni, J. C., C. E. Mason, S. M. Mane, M. Stephens, and Y. Gilad (2008). Rna-seq: an assessment of technical reproducibility and comparison with gene expression arrays. *Genome Research* 18(9), 1509–1517.
- Neal, R. M. et al. (2011). Mcmc using hamiltonian dynamics. *Handbook of Markov Chain Monte Carlo* 2(11).
- Papastamoulis, P., M.-L. Martin-Magniette, and C. Maugis-Rabusseau (2016). On the estimation of mixtures of Poisson regression models with large number of components. *Computational Statistics & Data Analysis* 93, 97–106.

- Rau, A., C. Maugis-Rabusseau, M.-L. Martin-Magniette, and G. Celeux (2015). Co-expression analysis of high-throughput transcriptome sequencing data with Poisson mixture models. *Bioinformatics* 31(9), 1420–1427.
- Robinson, M. D., D. J. McCarthy, and G. K. Smyth (2010). edgeR: a bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26(1), 139–140.
- Sarabia, J. M. and E. Gómez-Déniz (2011). Multivariate poisson-beta distributions with applications. *Communications in Statistics-Theory and Methods* 40(6), 1093–1108.
- Schwarz, G. et al. (1978). Estimating the dimension of a model. *The Annals of Statistics* 6(2), 461–464.
- Scrucca, L., M. Fop, T. B. Murphy, and A. E. Raftery (2016). mclust 5: clustering, classification and density estimation using Gaussian finite mixture models. *The R Journal* 8(1), 205–233.
- Si, Y., P. Liu, P. Li, and T. P. Brutnell (2014). Model-based clustering for RNA-seq data. *Bioinformatics* 30(2), 197–205.
- Silva, A., S. J. Rothstein, P. D. McNicholas, and S. Subedi (2019). A multivariate poisson-log normal mixture model for clustering transcriptome sequencing data. *BMC Bioinformatics* 20(1), 394.
- Stan Development Team (2015). *Stan: A C++ library for probability and sampling (Version 2.8.0)*.
- Subedi, S. and P. D. McNicholas (2014). Variational bayes approximations for clustering

- via mixtures of normal inverse gaussian distributions. *Advances in Data Analysis and Classification* 8(2), 167–193.
- Subedi, S. and P. D. McNicholas (2020). A variational approximations-dic rubric for parameter estimation and mixture model selection within a family setting. *Journal of Classification*, 1–20.
- Subedi, S., A. Punzo, S. Ingrassia, and P. D. McNicholas (2015). Cluser-weighed t -factor analyzers for robus model-based clusering and dimension reducion. *Statistical Methods & Applications* 24(4), 623–649.
- Tang, Y., R. P. Browne, and P. D. McNicholas (2015). Model based clustering of high-dimensional binary data. *Computational Statistics & Data Analysis* 87, 84–101.
- Teicher, H. (1954). On the multivariate poisson distribution. *Scandinavian Actuarial Journal* 1954(1), 1–9.
- Tortora, C., B. C. Franczak, R. P. Browne, and P. D. McNicholas (2019). A mixture of coalesced generalized hyperbolic distributions. *Journal of Classification* 36(1), 26–57.
- Vrbik, I. and P. D. McNicholas (2014). Parsimonious skew mixture models for model-based clustering and classification. *Computational Statistics and Data Analysis* 71, 196–210.
- Wainwright, M. J., M. I. Jordan, et al. (2008). Graphical models, exponential families, and variational inference. *Foundations and Trends® in Machine Learning* 1(1–2), 1–305.
- Wei, G. C. and M. A. Tanner (1990). A Monte Carlo implementation of the EM algorithm and the poor man’s data augmentation algorithms. *Journal of the American Statistical Association* 85(411), 699–704.

- Willmot, G. E. (1993). On recursive evaluation of mixed poisson probabilities and related quantities. *Scandinavian Actuarial Journal* 1993(2), 114–133.
- Witten, D., R. Tibshirani, S. G. Gu, A. Fire, and W.-O. Lui (2010). Ultra-high throughput sequencing-based small rna discovery and discrete statistical biomarker analysis in a collection of cervical tumours and matched controls. *BMC Biology* 8(1), 58.
- Yu, D., W. Huber, and O. Vitek (2013). Shrinkage estimation of dispersion in negative binomial models for rna-seq experiments with small sample size. *Bioinformatics* 29(10), 1275–1282.