STRONG UNIQUENESS POLYNOMIALS: THE COMPLEX CASE

TA THI HOAI AN, JULIE TZU-YUEH WANG, AND PIT-MANN WONG

ABSTRACT. The theory of strong uniqueness polynomials, satisfying the separation condition (first introduced by Fujimoto [4]), for complex meromorphic functions is quite complete. We construct examples of strong uniqueness polynomials which do not necessary satisfy the separation condition by constructing regular 1-forms of Wronskian type, a method introduced in [2]. We also use this method to produce a much easier proof in establishing the necessary and sufficient conditions for a polynomial, satisfying the separation condition, to be a strong uniqueness polynomials for meromorphic functions and rational functions.

1. Introduction

Recall that a polynomial P defined over C is said to be an uniqueness polynomial for meromorphic (respectively, rational) functions if it satisfies the condition P(f) =P(g) for non-constant meromorphic functions f, g implies that $f \equiv g$; P is said to be a strong uniqueness polynomial if it satisfies the condition P(f) = cP(g) for non-constant meromorphic (respectively, rational) functions f, g and some nonzero constant c implies that c=1 and $f\equiv g$. A polynomial P is said to separate the roots of its derivative P' if $P(\alpha) \neq P(\beta)$ for any distinct roots α and β of P'. For simplicity, we shall refer to this simply as the separation condition. A fairly complete picture of strong uniqueness polynomials for meromorphic functions (resp. rational functions) satisfying the separation condition is now known due to the works of Fujimoto ([4], [5]), and An and Wang [1] (resp. Khoai and An [6], and Wang [8]). As it turns out the separation condition though sufficient is not necessary. The first result (see Theorem 1) of this article is to construct examples of strong uniqueness polynomials not satisfying the separation condition using the method of constructing regular 1-forms of Wronskian type introduced in [2]. For polynomials satisfying the separation condition, the method of [2] also allows us to give a much easier proof of the necessary and sufficient conditions of strong uniqueness for rational functions (cf. [8]), and meromorphic functions (cf. [1]). The arguments in [8] and [1] using the truncated second main theorem for rational functions and meromorphic functions are no longer needed using the method of this article. The method also avoids some of the rather technical arguments of quadratic transformation used in [8] and [1]. The main results are as follows.

Theorem 1. Let $P(X) = a_n X^n + \sum_{i=0}^m a_i X^i$ $(0 \le m < n, a_i \in \mathbb{C} \text{ and } a_n, a_m \ne 0)$ be a polynomial of degree n. Let $I = \{i \mid a_i \ne 0\}, l = \min\{i \mid i \in I\}$ and $J = \{i - l \mid i \in I\}$. Then the following statements are valid.

- (i) If $n-m \geq 3$ then P is a strong uniqueness polynomial for rational functions if and only if the greatest common divisor of the indices in I is 1 and the greatest common divisor of the indices in J is also 1.
- (ii) If $n-m \geq 4$ then P is a strong uniqueness polynomial for meromorphic functions if and only if the greatest common divisor of the indices in I is 1 and the greatest common divisor of the indices in J is also 1.

In the Theorem above it is possible that $0 \in J$ and we use the convention that 0 is divisible by all integers.

Remark 1. If $n-m \ge 3$ and the greatest common divisor of the indices in J is 1 then $\#I \ge 3$. For if #I = 2 then $I = \{n, m\}$. Hence l = m and $J = \{n - m, 0\}$. By our convention, the greatest common divisor of the indices in J is $n - m \ge 3$. Thus P cannot be a strong uniqueness polynomial.

Remark 2. Theorem 1 can be stated for more general polynomials. Let $P(X) = X^n + a_{n-1}X^{n-1} + a_{n-2}X^{n-2} + \cdots + a_1X + a_0$ be a polynomial of degree n defined over \mathbf{C} , and let $P_0(X) = P(X - \frac{a_{n-1}}{n}) = X^n + b_{n-2}X^{n-2} + b_{n-3}X^{n-3} + \cdots + b_1X + b_0$. If

(A)
$$a_{n-2} = \frac{n-1}{2n} a_{n-1}^2$$

then $b_{n-2}=0$; if

(B)
$$a_{n-2} = \frac{n-1}{2n} a_{n-1}^2$$
 and $a_{n-3} = \frac{(n-1)(n-2)}{6n^2} a_{n-1}^3$

then $b_{n-2} = b_{n-3} = 0$. Thus $P_0(X)$ is a polynomial for which Theorem 1 is applicable. It is clear that P is a strong uniqueness polynomial (for rational functions or meromorphic functions) if and only if P_0 is a strong uniqueness polynomial.

The following concept plays an important role in the strong uniqueness Theorems for polynomials satisfying the separation condition.

Definition. A subset S of \mathbf{C} is said to be *affine rigid* if no non-trivial affine transformation of \mathbf{C} preserves S.

Theorem 2. Let P(X) be a polynomial of degree n over \mathbb{C} , and $P'(X) = \lambda(X - \alpha_1)^{m_1} \dots (X - \alpha_l)^{m_l}$ where λ is a nonzero constant and $\alpha_i \neq \alpha_j$ for $1 \leq i \neq j \leq l$.

Suppose that P(X) satisfies the separation condition, i.e., $P(\alpha_i) \neq P(\alpha_j)$ if $i \neq j$. Then

- (i) P(X) is a uniqueness polynomial for rational functions if and only if $l \geq 3$, or l = 2 and $\min\{m_1, m_2\} \geq 2$.
- (ii) P(X) is a strong uniqueness polynomial for rational functions if and only if the set of zeros of P is affinely rigid and one of the following conditions is satisfied: (a) $l=2, \min\{m_1, m_2\} \geq 2$, or (b) $l\geq 3$, except when n=4, $m_1=m_2=m_3=1$ and

$$\frac{P(\alpha_1)}{P(\alpha_2)} = \frac{P(\alpha_2)}{P(\alpha_3)} = \frac{P(\alpha_3)}{P(\alpha_1)} = w, \quad where \ w^2 + w + 1 = 0.$$

Theorem 3. Let P(X) be a polynomial of degree n over \mathbb{C} , and $P'(X) = \lambda(X - \alpha_1)^{m_1} \dots (X - \alpha_l)^{m_l}$ where λ is a nonzero constant and $\alpha_i \neq \alpha_j$ for $1 \leq i \neq j \leq l$. Suppose that P(X) satisfies the separation condition, i.e., $P(\alpha_i) \neq P(\alpha_j)$ if $i \neq j$ and that the set of zeros of P is affinely rigid. Then

- (i) P(X) is a uniqueness polynomial for meromorphic functions if and only if one of the following conditions is satisfied: (a) l≥ 3, except when n = 4, m₁ = m₂ = m₃ = 1; or (b) l = 2 and min{m₁, m₂} ≥ 2 except when n = 5, m₁ = m₂ = 2.
- (ii) P(X) is a strong uniqueness polynomial for meromorphic functions if and only if the set of zeros of P is affinely rigid and one of the following conditions is satisfied: (a) l≥ 3, except when n = 4, m₁ = m₂ = m₃ = 1; or
 (b) l = 2 and min{m₁, m₂} ≥ 2 except when n = 5, m₁ = m₂ = 2.

For polynomials of the special type $(X - \alpha)^n + a(X - \alpha)^m + b$, we have the following complete characterization.

Corollary 1. Let $P(X) = (X - \alpha)^n + a(X - \alpha)^m + b$ be a polynomial of degree n and $1 \le m \le n - 1$. Then

- (i) P(X) is a uniqueness polynomial for rational functions if and only if $n \ge 4$, $n-m \ge 2$, gcd(n,m) = 1 and $a \ne 0$;
- (ii) P(X) is a strong uniqueness polynomial for rational functions if and only if $n \ge 4$, $n m \ge 2$, gcd(n, m) = 1, $a \ne 0$, and $b \ne 0$;
- (iii) P(X) is a uniqueness polynomial for meromorphic functions if and only if $n \geq 5$, $n m \geq 2$, gcd(n, m) = 1 and $a \neq 0$;
- (iv) P(X) is a strong uniqueness polynomial for meromorphic functions if and only if $n \ge 5$, $n m \ge 2$, gcd(n, m) = 1, $a \ne 0$, and $b \ne 0$.

The idea, as usual, is to show that the following curves, associated to the polynomial P, in $\mathbf{P}^2(\mathbf{C})$ is Brody hyperbolic (in our case this is equivalent to Kobayashi hyperbolic because the curves under consideration are compact):

$$C = [F(X, Y, Z) = 0], C_c = [F_c(X, Y, Z) = 0], c \neq 0, 1$$

where F(X, Y, Z) is the homogenization of the polynomial:

$$\frac{P(X) - P(Y)}{X - Y}$$

and $F_c(X, Y, Z)$ is the homogenization of the polynomial:

$$P(X) - cP(Y), \ c \neq 0, 1.$$

Remark. The set of zeros of P(X) is affinely rigid if and only if F(X, Y, Z) and each $F_c(X, Y, Z)$, $c \neq 0$, 1, have no linear factors

A projective curve is Brody hyperbolic if and only if the genus of the curve is at least 2. We also say that a projective curve C is algebraically hyperbolic if every algebraic map $f: \mathbb{C} \to C$ is constant. It is well-known that a projective curve is algebraically hyperbolic if and only if the genus of the curve is at least 1. For general P the singularities of C and C_c can be complicated which makes it difficult to use the classical genus formula. Moreover, one needs irreducibility of the curves in order to apply the genus formula and, unfortunately irreducibility is usually very difficult to verify even for the special type of polynomials that we are using. For these reasons we adopt the approach in [2] by constructing sufficiently many explicit non-trivial regular 1-forms of Wronskian type on these curves under the assumptions of Theorem 1. The main advantage of using Wronskian type 1-forms is that it is only necessary to show that there is no linear factor (component). The reason being that a curve is Brody (resp. algebraic) hyperbolic if and only each of its components is Brody (resp. algebraic) hyperbolic. A regular 1-form of Wronskian is non-trivial on a component if and only if the component is nonlinear and the existence of q linearly independent regular 1-form(s) on a component implies that the component must be of genus at least g. We shall introduce the notion of regular 1-forms of Wronskian type in section 2 and show via examples how these forms may be constructed. This procedure will then be applied to the curves C and C_c in section 3.

2. REGULAR 1-FORMS OF WRONSKIAN TYPE

In this section we deal with the practical problem of computing the genus of a curve in $\mathbf{P}^2(\mathbf{C})$. For a smooth curve this is easily computed via the well-known genus formula g = (n-1)(n-2)/2 where n is the degree of the smooth curve.

Note that (n-1)(n-2)/2 is the number of distinct monomials of degree n-3 in z_0 , z_1 and z_2 . There is also a genus formula for irreducible singular curves in terms of the Milnor number and the number of local branches at each of the singular point. It is usually quite a chore in computing these invariants when the singularity is complicated; moreover, it is usually extremely difficult to check their irreducibility condition. On the other hand, in Nevanlinna Theory a priori knowledge of irreducibility is usually not necessary. The process, based on the Second Main Theorem, will automatically break down if the curve has any component of genus one. The reason being that, in the Second Main Theorem there is a ramification term which comes from the Wronskian of a map into projective space. For this reason we shall develop a procedure of computing genus, based on the Wronskian, without a priori knowledge of irreducibility. The main idea is as follows. Observe that

$$\frac{\begin{vmatrix} z_i & z_j \\ dz_i & dz_j \end{vmatrix}}{z_j^2} = \frac{z_i}{z_j} \begin{vmatrix} 1 & 1 \\ \frac{dz_i}{z_i} & \frac{dz_j}{z_j} \end{vmatrix} = d(\frac{z_i}{z_j}), \ i \neq j$$

being the differential of a well-defined rational function is a well-defined rational 1-form on $\mathbf{P}^2(\mathbf{C})$ with homogeneous coordinates z_0, z_1 and z_2 . Denote

$$W(z_i, z_j) := \begin{vmatrix} z_i & z_j \\ dz_i & dz_j \end{vmatrix} = z_i dz_j - z_j dz_i.$$

Thus, for any rational function ϕ on $\mathbf{P}^2(\mathbf{C})$ then, for $i \neq j$:

$$\phi \frac{W(z_i, z_j)}{z_j^2}, \ W(z_i, z_j) = z_i z_j \begin{vmatrix} 1 & 1 \\ \frac{dz_i}{z_i} & \frac{dz_j}{z_j} \end{vmatrix}$$

is a well-defined rational 1-form on $\mathbf{P}^2(\mathbf{C})$. Equivalently, for any homogeneous polynomials R and S such that deg $S = \deg R + 2$ then

(2.1)
$$\frac{R}{S}W(z_i, z_j) = \phi \frac{W(z_i, z_j)}{z_i^2}, \ \phi = \frac{z_2^2 R}{S}$$

is a well-defined rational 1-form on $\mathbf{P}^2(\mathbf{C})$.

Definition 2.1. Let $C \subset \mathbf{P}^2(\mathbf{C})$ be an algebraic curve. A 1-form on C is said to be regular if it is the restriction (more precisely, the pull-back) of a rational 1-form on $\mathbf{P}^2(\mathbf{C})$ such that the pole set of ω does not intersect C. A 1-form is said to be of Wronskian type if it is of the form (2.1) above.

To see how a regular 1-form of Wronskian type may be constructed we start by dealing with non-singular curves where the idea is most transparent and then extend this to the singular case in the next section. Let $P(Z_0, Z_1, Z_2)$ be a homogeneous polynomial of degree n and let

$$C = \{ [z_0, z_1, z_2] \in \mathbf{P}^2(\mathbf{C}) \mid P(z_0, z_1, z_2) = 0 \}.$$

Then, by Euler's Theorem, for $[z_0, z_1, z_2] \in C$, we have

$$z_0 \frac{\partial P}{\partial z_0}(z_0, z_1, z_2) + z_1 \frac{\partial P}{\partial z_1}(z_0, z_1, z_2) + z_2 \frac{\partial P}{\partial z_2}(z_0, z_1, z_2) = 0.$$

The (Zariski) tangent space of C is defined by the equation $P(z_0, z_1, z_2) = 0$ and

$$dz_0 \frac{\partial P}{\partial z_0}(z_0, z_1, z_2) + dz_1 \frac{\partial P}{\partial z_1}(z_0, z_1, z_2) + dz_2 \frac{\partial P}{\partial z_2}(z_0, z_1, z_2) = 0.$$

These may be expressed as

$$z_{0} \frac{\partial P}{\partial z_{0}}(z_{0}, z_{1}, z_{2}) + z_{1} \frac{\partial P}{\partial z_{1}}(z_{0}, z_{1}, z_{2}) = -z_{2} \frac{\partial P}{\partial z_{2}}(z_{0}, z_{1}, z_{2}),$$

$$dz_{0} \frac{\partial P}{\partial z_{0}}(z_{0}, z_{1}, z_{2}) + dz_{1} \frac{\partial P}{\partial z_{1}}(z_{0}, z_{1}, z_{2}) = -dz_{2} \frac{\partial P}{\partial z_{2}}(z_{0}, z_{1}, z_{2}).$$

Then by Cramer's rule, we have, on C

$$\frac{\partial P}{\partial z_0} = \frac{W(z_1, z_2)}{W(z_0, z_1)} \frac{\partial P}{\partial z_2}, \quad \frac{\partial P}{\partial z_1} = \frac{W(z_2, z_0)}{W(z_0, z_1)} \frac{\partial P}{\partial z_2}$$

provided that $W(z_0, z_1) \not\equiv 0$ on any component of C, i.e., the defining homogeneous polynomial of C has no linear factor of the form $az_0 + bz_1$. Thus

(2.2)
$$\frac{W(z_1, z_2)}{\frac{\partial P}{\partial z_0}(z_0, z_1, z_2)} = \frac{W(z_2, z_0)}{\frac{\partial P}{\partial z_1}(z_0, z_1, z_2)} = \frac{W(z_0, z_1)}{\frac{\partial P}{\partial z_2}(z_0, z_1, z_2)}$$

is a globally well-defined rational 1-form on any component of $\pi^{-1}(C) \subset \mathbb{C}^3 \setminus \{0\}$ where $(\pi : \mathbb{C}^3 \setminus \{0\} \to \mathbb{P}^2(\mathbb{C}))$ is the Hopf fibration); provided that, of course, the expressions make sense, i.e. the denominators are not identically zero when restrict to a component of $\pi^{-1}(C)$. For our purpose, we also require that the form given by (2.2) is not identically trivial when restrict to a component of $\pi^{-1}(C)$. This is equivalent to the condition that the Wronskians in the formula above are not identically zero, i.e., the defining homogeneous polynomial of C has no linear factor of the form $az_i + bz_j$ where $a, b \in \mathbb{C}$, $0 \le i, j \le 2$ and $i \ne j$. If

$$[P=0] \cap \left[\frac{\partial P}{\partial z_0} = 0\right] \cap \left[\frac{\partial P}{\partial z_1} = 0\right] \cap \left[\frac{\partial P}{\partial z_2} = 0\right] = \emptyset$$

(i.e., C is smooth) then, at any point, one of the expression in (2.2) is regular at the point, hence so are the other expressions. This means that

(2.3)
$$\eta = \frac{\begin{vmatrix} z_1 & z_2 \\ dz_1 & dz_2 \end{vmatrix}}{\frac{\partial P}{\partial z_0}}$$

is regular on $\pi^{-1}(C)$ (note that the form η is not well-defined on C unless n=3, see (2.1)). The form $(n=\deg P)$

(2.4)
$$\omega = \frac{\begin{vmatrix} z_1 & z_2 \\ dz_1 & dz_2 \end{vmatrix}}{z_0^2} \frac{z_0^{n-1}}{\frac{\partial P}{\partial z_0}} = \frac{\begin{vmatrix} z_1 & z_2 \\ dz_1 & dz_2 \end{vmatrix}}{\frac{\partial P}{\partial z_0}} z_0^{n-3} = z_0^{n-3} \eta$$

is a well-defined (again by (2.1)) rational 1-form on C. Moreover, as η is regular on C, the 1-form ω is also regular if $n \geq 3$. If n = 3 then $\omega = \eta$ and if $n \geq 4$ then ω is regular and vanishes along the ample divisor $[z_0^{n-3} = 0] \cap C$. Thus for any homogeneous polynomial $Q = Q(z_0, z_1, z_2)$ of degree n - 3, the 1-form

$$\frac{Q}{z_0^{n-3}}\omega = Q\eta$$

is regular on C and vanishes along [Q=0]. Note that the dimension of the vector space of homogeneous polynomials of degree n-3 (a basis is given by all possible monomials) is

$$\frac{(n-1)(n-2)}{2}$$
 = genus of C .

We summarize these in the following Proposition:

Proposition 2.2. Let $C = \{[z_0, z_1, z_2] \in \mathbf{P}^2(\mathbf{C}) | P(z_0, z_1, z_2) = 0\}$ be a non-singular curve of degree $n \geq 3$. If n = 3 then the space of regular 1-forms on C is $\{c\eta | c \in \mathbf{C}\}$ where η is defined by (2.2). If $n \geq 4$ let

$$\{Q_i | Q_i \text{ is a monomial of degree } n-3, 1 \le i \le (n-1)(n-2)/2\}$$

be a basis of homogeneous polynomials of degree n-3 then

$$\{\omega_i = Q_i \eta \mid 1 \le i \le \frac{(n-1)(n-2)}{2}\}$$

is a basis of the space of regular 1-forms on C.

Next we extend the construction to some examples of singular curves.

Example 1. Let $P_{m,n}(z_0, z_1, z_2) = z_0^n + z_1^m z_2^{n-m} + z_2^n = 0, n \ge m \ge 1$. If n = m then the curve $C = [P_{m,n}(z_0, z_1, z_2) = 0]$ is non-singular and so by Proposition 2.2, if n = 3 then all holomorphic 1-forms are constant multiples of

$$\eta = \frac{\begin{vmatrix} z_1 & z_2 \\ dz_1 & dz_2 \end{vmatrix}}{z_0^2}.$$

If n = 4 then

$$\{\frac{z_0\begin{vmatrix}z_1&z_2\\dz_1&dz_2\end{vmatrix}}{z_0^2},\frac{z_1\begin{vmatrix}z_1&z_2\\dz_1&dz_2\end{vmatrix}}{z_0^2},\frac{z_2\begin{vmatrix}z_1&z_2\\dz_1&dz_2\end{vmatrix}}{z_0^2}\}$$

is a basis of holomorphic 1-forms on $C_{4,4}$.

Consider now the case $n > m \ge 1$ then

$$\begin{split} \partial P_{m,n}/\partial z_0 &= n z_0^{n-1} = 0, \\ \partial P_{m,n}/\partial z_1 &= m z_1^{m-1} z_2^{n-m} = 0, \\ \partial P_{m,n}/\partial z_2 &= (m-n) z_1^m z_2^{n-m-1} + n z_2^{n-1} = 0. \end{split}$$

If m+1=n the curve $C_{m,n}=\{[z_0,z_1,z_2]\in \mathbf{P}^2(\mathbf{C})\,|\,P_{m,n}(z_0,z_1,z_2)=0\}$ is still smooth. If $n=m+2\geq 3$ then $C_{m,n}$ is singular with a unique singular point at [0,1,0]. Proposition 2.2 does not apply to singular curves but the procedure of the construction of holomorphic forms can be modified as follows. The identities (2.2) is now of the form:

$$\frac{W(z_1, z_2)}{nz_0^{n-1}} = \frac{W(z_2, z_0)}{(n-2)z_1^{n-3}z_2^2} = \frac{W(z_0, z_1)}{z_2(2z_1^{n-2} + nz_2^{n-2})}$$

where the denominators now have common zero. Instead of taking

$$\eta = \frac{\begin{vmatrix} z_2 & z_0 \\ dz_2 & dz_0 \end{vmatrix}}{\frac{\partial P}{\partial z_1}} = \frac{\begin{vmatrix} z_2 & z_0 \\ dz_2 & dz_0 \end{vmatrix}}{z_1^{n-3} z_2^2}$$

(as in the smooth case) which is not regular, we take

$$\eta = \frac{\begin{vmatrix} z_2 & z_0 \\ dz_2 & dz_0 \end{vmatrix}}{z_1^{n-3} z_2} = (n-2) \frac{\begin{vmatrix} z_0 & z_1 \\ dz_0 & dz_1 \end{vmatrix}}{2z_1^{n-2} + nz_2^{n-2}}$$

which is regular on $\pi^{-1}(C_{m,n})$ because the common zero of the denominators are given by the equation $[z_1 = z_2 = 0]$, i.e., the point [1,0,0] which is not on $C_{m,n}$. Hence

$$\omega = \frac{\begin{vmatrix} z_2 & z_0 \\ dz_2 & dz_0 \end{vmatrix}}{z_0^2} \frac{z_0^2 z_0^{n-4} z_2}{z_1^{n-3} z_0^2} = \frac{\begin{vmatrix} z_2 & z_0 \\ dz_2 & dz_0 \end{vmatrix}}{z_1^{n-3} z_2} z_0^{n-4} = z_0^{n-4} \eta$$

is globally well-defined on $C_{m,n}$, regular and vanishing along $(n-4)[z_0=0]$. This implies that if n=4 then $\omega=\eta$ is a global regular 1-form on $C_{m,4}$. If n=5 then $\omega=z_0\eta$ is globally regular and vanishes along the ample divisor $[z_0=0]$. Indeed we see that

$$\{\omega = \frac{z_0}{z_1^2 z_2} \begin{vmatrix} z_2 & z_0 \\ dz_2 & dz_0 \end{vmatrix}, \frac{z_1}{z_0}\omega = \frac{z_1}{z_1^2 z_2} \begin{vmatrix} z_2 & z_0 \\ dz_2 & dz_0 \end{vmatrix}, \frac{z_2}{z_0}\omega = \frac{z_2}{z_1^2 z_2} \begin{vmatrix} z_2 & z_0 \\ dz_2 & dz_0 \end{vmatrix}\}$$

are linearly independent holomorphic 1-forms on $C_{3,5}$ hence the genus of $C_{3,5}$ is $\geq 3 = \frac{(5-1)(5-2)}{2} - 3$.

More generally if $n = m + k, k \ge 3$ then

$$\frac{W(z_1, z_2)}{nz_0^{n-1}} = \frac{W(z_2, z_0)}{(n-k)z_1^{n-k-1}z_2^k} = \frac{W(z_0, z_1)}{z_2^{k-1}(kz_1^{n-k} + nz_2^{n-k})}$$

and

$$\eta = \frac{\begin{vmatrix} z_2 & z_0 \\ dz_2 & dz_0 \end{vmatrix}}{z_1^{n-k-1} z_2}$$

is regular on $\pi^{-1}(C_{m,n})$ hence

$$\omega = \frac{\begin{vmatrix} z_2 & z_0 \\ dz_2 & dz_0 \end{vmatrix}}{z_0^2} \frac{z_0^2 z_0^{n-k}}{z_1^{n-k-1} z_2} = \frac{\begin{vmatrix} z_2 & z_0 \\ dz_2 & dz_0 \end{vmatrix}}{z_1^{n-k-1} z_2} z_0^{n-k-2} = z_0^{n-k-2} \eta = z_0^{m-2} \eta$$

is globally well-defined on $C_{m,n}$, regular and vanishing along $(n-k-2)[z_0=0]$ if $m=n-k\geq 2$. Let $\{Q_1,...,Q_{m(m-1)/2}\}$ be a basis of monomials of degree m-2 in $\{z_0,z_1,z_2\}$ then

$${Q_i \eta \mid i = 1, ..., m(m-1)/2}$$

are linearly independent global regular 1-forms on $C_{m,m+k}$, $m \ge 2, k \ge 2$. Thus the genus of $C_{m,m+k} \ge m(m-1)/2$ for all $m \ge 2, k \ge 2$.

The procedure of this section will be applied in the next section to deal with the situation of uniqueness polynomials.

3. The case
$$P(X) = P(Y)$$

Let C be a plane curve (not necessarily irreducible) defined by a homogeneous polynomial R(X, Y, Z) = 0 over \mathbf{K} and let \mathfrak{p} be a point of C. A holomorphic map

(3.1)
$$\phi = (\phi_0, \phi_1, \phi_2) : \Delta_{\epsilon} = \{ t \in \mathbf{K} \mid |t| < \epsilon \} \to C, \quad \varphi(0) = \mathfrak{p}$$

is referred to as a holomorphic parameterization of C at \mathfrak{p} . Local holomorphic parameterization exists for sufficiently small ϵ . A rational function Q on the curve C is represented by A/B where A and B are homogeneous polynomials in X,Y,Z such that $B|_C$ is not identically zero. Thus $Q \circ \phi$ is a well-defined meromorphic function on Δ_{ϵ} with Laurent expansion

$$Q \circ \phi(t) = \sum_{i=m}^{\infty} a_i t^i, \qquad a_m \neq 0.$$

The order of $Q \circ \phi$ at t = 0 is by definition m and shall be denoted by

$$\operatorname{ord}_{\mathfrak{n},\phi}Q = \operatorname{ord}_{t=0}Q(\phi(t)).$$

The function $Q \circ \phi$ is holomorphic if and only if $m \geq 0$. The rational function Q is regular at \mathfrak{p} if and only if $Q \circ \phi$ is holomorphic for *all* local holomorphic parameterizations of C at \mathfrak{p} .

Let P(X) be a polynomial of degree n:

$$P(X) = X^{n} + a_{m}X^{m} + a_{m-1}X^{m-1} + \dots + a_{1}X + a_{0}, a_{m} \neq 0$$

defined over C. We have

$$P'(X) = n(X - \alpha_1)^{m_1}...(X - \alpha_l)^{m_l}$$

where $\alpha_i \neq \alpha_j$ for $i \neq j$ and $m_i \geq 1$. Thus $X = \alpha_i$ is a root of order $m_i + 1$ of $P(X) - P(\alpha_i)$ hence:

(3.3)
$$P(X) - P(\alpha_i) = \sum_{j=m_i+1}^n b_{i,j} (X - \alpha_i)^j, \ b_{i,m_i+1} \neq 0, b_{i,n} \neq 0.$$

A polynomial P is said to separate the roots of P' if

(3.4)
$$P(\alpha_i) \neq P(\alpha_j) \text{ for all } 1 \leq i \neq j \leq l.$$

Let F(X,Y,Z) be the homogenization of the polynomial

$$\frac{P(X) - P(Y)}{X - Y} = \sum_{k=1}^{n} a_k \sum_{j=0}^{k-1} X^{k-1-j} Y^j$$

i.e., $F(X, Y, Z) = Z^n \{P(X/Z) - P(Y/Z)\}/(X - Y)$.

Remark. X - Y is not a factor of F(X, Y, Z) since $F(X, X, 1) = P'(X) \not\equiv 0$. Y - aZ or X - aZ, $a \in k$, is not a factor of F(X, Y, Z) either since $P(Y) \not\equiv P(a)$ and $P(X) \not\equiv P(a)$.

For each $1 \leq i \leq l$ we may, by (3.3), express the polynomial F(X, Y, Z) as a polynomial in $(X - \alpha_i Z)$ and $(Y - \alpha_i Z)$:

(3.5)
$$F(X,Y,Z) = \sum_{j=m_i+1}^{n} \left[b_{i,j} \frac{(X - \alpha_i Z)^j - (Y - \alpha_i Z)^j}{X - Y}\right] Z^{n-j},$$

 $b_{i,m_i+1} \neq 0, b_{i,n} \neq 0$. It is clear that the points $(\alpha_i, \alpha_i, 1) \in C = [F(X, Y, Z) = 0], 1 \leq i \leq l$. On the other hand, the separation condition (3.4) implies that $(\alpha_i, \alpha_j, 1) \notin C$ if $i \neq j$. Denote by $P'(X, Z) = Z^{n-1}P'(X/Z)$ and $P'(Y, Z) = Z^{n-1}P'(Y/Z)$ the homogenization of the polynomials P'(X) and P'(Y) respectively, then

(3.6)
$$P'(X,Z) = n \prod_{i=1}^{l} (X - \alpha_i Z), \ P'(Y,Z) = n \prod_{i=1}^{l} (Y - \alpha_i Z).$$

By the remark above it is clear that P'(X, Z) and P'(Y, Z) are not identically zero on any component of C. Differentiating the polynomial F(X, Y, Z) yields:

(3.7)
$$\begin{cases} \frac{\partial F}{\partial X}(X, Y, Z) = \frac{P'(X, Z) - F(X, Y, Z)}{X - Y}, \\ \frac{\partial F}{\partial Y}(X, Y, Z) = \frac{-P'(Y, Z) + F(X, Y, Z)}{X - Y}, \\ \frac{\partial F}{\partial Z}(X, Y, Z) = (n - m)a_m Z^{n - m - 1} \left(\sum_{i=0}^{m - 1} X^{m - i} Y^i + Z H_{m - 2}\right) \end{cases}$$

where $H_{m-2}(X,Y,Z)$ is a homogenous polynomial of degree m-2. Let

$$W(X,Y) = \begin{vmatrix} X & Y \\ dX & dY \end{vmatrix}, \ W(Y,Z) = \begin{vmatrix} Y & Z \\ dY & dZ \end{vmatrix}, \ W(Z,X) = \begin{vmatrix} Z & X \\ dZ & dX \end{vmatrix}$$

be the Wronskians which are regular 1-forms on \mathbb{C}^3 then (see (2.2) or [2]):

(3.8)
$$\gamma := \frac{W(X,Y)}{\frac{\partial F}{\partial Z}} = \frac{W(Y,Z)}{\frac{\partial F}{\partial X}} = \frac{W(Z,X)}{\frac{\partial F}{\partial Y}}$$

is a well-defined non-trivial rational 1-form on (any component of) $\pi^{-1}(C)$ (π : $\mathbf{C}^3 \setminus \{0\} \to \mathbf{P}^2$ is the usual fibration and C = [F(X,Y,Z) = 0] is a curve in \mathbf{P}^2). It is well-defined and non-trivial because, by (3.6) and (3.8), the restriction of $\partial F/\partial X$ to C is

(3.9)
$$\frac{P'(X,Z)}{X-Y} = n \frac{\prod (X - \alpha_i Z)}{X-Y}.$$

By the remark after (3.4), $X - \alpha_i Z$, X - Y and W(Y, Z) are not identically zero on any component of C. Moreover, for any homogeneous polynomials A(X, Y, Z) and B(X, Y, Z) with deg $B = \deg A + 2$, the rational 1-forms

$$R(X,Y,Z)W(X,Y), R(X,Y,Z)W(Y,Z), R(X,Y,Z)W(Z,X)$$

with R(X,Y,Z) = A(X,Y,Z)/B(X,Y,Z) are globally well-defined on \mathbf{P}^2 (see section 2 or [2]).

The next result provides sufficient conditions for the hyperbolicity of the curve C for a class of polynomials which does not necessary satisfy the separation condition.

Proposition 3.1. Let $P(X) = X^n + a_m X^m + a_{m-1} X^{m-1} + \cdots + a_1 X + a_0$, $a_m \neq 0$, be a polynomial of degree n. Assume that the curve C = [F(X,Y,Z) = 0] has no linear component. Then C is algebraically hyperbolic if $n - m \geq 3$ and is Brody hyperbolic if $n - m \geq 4$.

Proof. By (3.7) the rational 1-form γ defined by (3.8) satisfies the condition

$$\gamma = \frac{(X - Y)W(X, Z)}{P'(Y, Z)}
= \frac{(X - Y)W(Y, Z)}{P'(X, Z)}
= \frac{W(X, Y)}{(n - m)a_m Z^{n-m-1} (X^{m-1} + X^{m-2}Y + \dots + Y^m + ZH_{m-2})}$$

on $\pi^{-1}(C)$ where P'(X,Z) and P'(Y,Z) are given by (3.6). If γ is trivial on an irreducible component of $\pi^{-1}(C)$ then F(X,Y,1) has a linear factor contradicting the assumption that F(X,Y,Z) has no linear factor. Thus γ is non-trivial

on any component of C. Let L(X,Y,Z) be any linear form and $K(X,Y,Z) = Z^{n-m-4}(X^{m-1} + X^{m-2}Y + \cdots + Y^{m-1} + ZH_{m-2})$, then the rational 1-form

(3.11)
$$\omega := L(X, Y, Z)K(X, Y, Z)\gamma = \frac{L(X, Y, Z)}{(n - m)a_m Z^3}W(X, Y)$$

is well-defined on \mathbf{P}^2 because the denominator (of the coefficient of W(X,Y)) is two degrees higher than the numerator (see the remark before the Proposition). For the same reason, the rational 1-form

(3.12)
$$\theta := G(X, Y, Z)\gamma = \frac{1}{(n-m)a_m Z^2} W(X, Y)$$

(where $G(X,Y,Z) = Z^{n-m-3}(X^{m-1} + X^{m-2}Y + \cdots + Y^{m-1} + ZH_{m-2})$) is well-defined on \mathbf{P}^2 . It is clear that the pull-back of ω to the curve C is non-trivial on each component of C. We claim that it is also regular. From the definition (3.11) it is clear that the only possible poles of ω are the points $[Z=0] \cap C$. On the other hand, as a form on C, we see via (3.10) that

$$\omega = \frac{(X - Y)L(X, Y, Z)K(X, Y, Z)W(Y, Z)}{P'(X, Z)}$$
$$= \frac{(X - Y)L(X, Y, Z)K(X, Y, Z)W(X, Z)}{P'(Y, Z)}.$$

If Z=0 then, since $n-m-4\geq 0$, the denominator P'(X,Z) in the expression above is reduced to nX^{n-1} (resp. P'(Y,Z) is reduced to nY^{n-1}). Thus, if ω has a pole at a point with Z=0 then we must have X=Y=0 as well which, of course, is impossible in \mathbf{P}^2 . We conclude from this that ω is a regular 1-form on C. Choosing L(X,Y,Z)=X,Y and Z respectively we obtain 3 regular 1-forms on C:

$$\frac{XW(X,Y)}{(n-m)a_mZ^3}, \frac{YW(X,Y)}{(n-m)a_mZ^3}, \frac{W(X,Y)}{(n-m)a_mZ^2}$$

which are linearly independent on each component of C. Thus the genus of each irreducible component of C is not less than 3. By Picard's theorem, this shows that C is Brody hyperbolic. Respectively, for the case $n-m \geq 3$ we have to use θ (as defined in (3.12)) which cannot be further modified (except by constants) we get only one regular 1-form non-trivial on any component, hence we can only conclude that the genus of each irreducible component of C is not less than one. This, however, still implies that C is algebraically hyperbolic as there is no non-constant algebraic map from \mathbf{C} into an elliptic curve.

The condition that C has no linear component is satisfied if we assume that the zero set of P is affine rigid (See [8]). The method in the proof of the preceding Proposition can also be used to treat the case of a polynomial P(X) satisfying the separation condition. First we need a technical Lemma.

Lemma 3.2. Let $\mathfrak{p}_i = (\alpha_i, \alpha_i, 1)$ and ϕ be a local holomorphic parameterization of C = [F(X, Y, Z) = 0] at \mathfrak{p}_i . Then $ord_{\mathfrak{p}_i, \phi}(X - \alpha_i) = ord_{\mathfrak{p}_i, \phi}(Y - \alpha_i) \leq ord_{\mathfrak{p}_i, \phi}(X - Y)$.

Proof. From expression of the curve C at \mathfrak{p}_i , via (3.5), it is easy to see that $\operatorname{ord}_{t=0}(X(\phi(t)) - \alpha_i) = \operatorname{ord}_{t=0}(Y(\phi(t)) - \alpha_i)$ and

$$\operatorname{ord}_{t=0}(X - Y)(\phi(t)) = \operatorname{ord}_{t=0}\{X(\phi(t)) - \alpha_i - (Y(\phi(t)) - \alpha_i)\}$$

$$\geq \min\{\operatorname{ord}_{t=0}(X(\phi(t)) - \alpha_i), \operatorname{ord}_{t=0}(Y(\phi(t)) - \alpha_i)\}$$

$$= \operatorname{ord}_{t=0}(X(\phi(t)) - \alpha_i)$$

as claimed. \Box

Proposition 3.3. Let P(X) be a polynomial satisfying the separation condition (3.4). Denote $\{\alpha_i, 1 \leq i \leq l\}$ be the distinct roots of P'(X) with respective multiplicities $\{m_i, 1 \leq i \leq l\}$. Then

- (i) C is algebraically hyperbolic if $l \geq 3$, or l = 2 and $\min\{m_1, m_2\} \geq 2$;
- (ii) C is Brody hyperbolic if either of the following holds (a) $l \ge 4$, (b) l = 3 and $\max\{m_1, m_2, m_3\} > 1$, (c) l = 2, $\min\{m_1, m_2\} \ge 2$ and $\max\{m_1, m_2\} \ge 3$.

Proof. By (3.10),

$$\gamma = \frac{(X-Y)W(X,Z)}{P'(Y,Z)} = \frac{(X-Y)W(Y,Z)}{P'(X,Z)}$$

on $\pi^{-1}(C)$ where C = [F(X, Y, Z) = 0]. Canceling out the common factor X - Y, we get, via (3.6), the following rational 1-form:

$$\eta = \frac{W(Y,Z)}{(X - \alpha_1 Z)^{m_1}...(X - \alpha_l Z)^{m_l}} = \frac{W(X,Z)}{(Y - \alpha_1 Z)^{m_1}...(Y - \alpha_l Z)^{m_l}}$$

well-defined on $\pi^{-1}(C)$. Without loss of generality, we may assume that $m_1 \ge m_2 \ge m_i$, for $2 \le i \le l$. Suppose that $l \ge 3$, or l = 2 and $\min\{m_1, m_2\} \ge 2$. In either cases we have $n \ge 4$. The rational 1-forms:

$$\omega_0 := (X - Y)^{n-3} \eta = \frac{W(Y, Z)(X - Y)^{n-3}}{(X - \alpha_1 Z)^{m_1} ... (X - \alpha_l Z)^{m_l}},$$

$$\omega_1 := (X - \alpha_1 Z)(X - Y)^{n-4} \eta = \frac{W(Y, Z)(X - Y)^{n-4}}{(X - \alpha_1 Z)^{m_1 - 1} (X - \alpha_2 Z)^{m_2} ... (X - \alpha_l Z)^{m_l}},$$

$$\omega_2 := (X - \alpha_2 Z)(X - Y)^{n-4} \eta = \frac{W(Y, Z)(X - Y)^{n-4}}{(X - \alpha_1 Z)^{m_1} (X - \alpha_2 Z)^{m_2 - 1} ... (X - \alpha_l Z)^{m_l}},$$

are well-defined on the curve C (because each of the denominator of the coefficients of W(Y, Z) is of two degree higher than the numerator). By the remark after (3.4) it is clear that ω_i are non-trivial on any irreducible component of C. Observe also that ω_i does not have any pole along [Z = 0] (otherwise we have X = Y = 0

as well). On the finite part of C (i.e., Z=1) the separation condition (3.4) and the condition that $n \geq 4$ imply that the only possible poles of ω_i are the points $\mathfrak{p}_j = (\alpha_j, \alpha_j, 1), j = 1, ..., l$. Lemma 3.2 implies that, for any local parameterization ϕ of C at $\mathfrak{p}_j, 1 \leq j \leq l$ ($\phi(0) = \mathfrak{p}_j$),

$$\operatorname{ord}_{\mathfrak{p}_{j},\phi}\omega_{0} = (n-3)\operatorname{ord}_{\mathfrak{p}_{j},\phi}(X-Y) - m_{j}\operatorname{ord}_{\mathfrak{p}_{j},\phi}(X-\alpha_{j})$$

$$\geq (n-3-m_{j})\operatorname{ord}_{\mathfrak{p}_{j},\phi}(X-\alpha_{j}).$$

Since $m_1 \ge m_j$ for $1 \le j \le l$ and $m_1 + ... + m_l = \deg P' = n - 1$, we infer that

$$\operatorname{ord}_{\mathfrak{p}_{j},\phi}\omega_{0} \geq (n-3-m_{1})\operatorname{ord}_{\mathfrak{p}_{j},\phi}(X-\alpha_{j}) \geq \{(\sum_{i=2}^{l}m_{i})-2\}\operatorname{ord}_{\mathfrak{p}_{j},\phi}(X-\alpha_{j})$$

which is non-negative if $l \geq 3$ or l = 2 and $\min\{m_1, m_2\} \geq 2$. This implies part (i) of the Proposition.

Similarly, we get via Lemma 3.2:

$$\operatorname{ord}_{\mathfrak{p}_{1},\phi} \omega_{1} = (n-4)\operatorname{ord}_{\mathfrak{p}_{1},\phi} (X-Y) - (m_{1}-1)\operatorname{ord}_{\mathfrak{p}_{1},\phi} (X-\alpha_{1})$$

$$\geq (n-3-m_{1})\operatorname{ord}_{\mathfrak{p}_{1},\phi} (X-\alpha_{1})$$

$$\geq \{(\sum_{i=2}^{l} m_{i}) - 2\}\operatorname{ord}_{\mathfrak{p}_{1},\phi} (X-\alpha_{1}) \geq 0$$

provided that $l \geq 3$ or l = 2 and $\min\{m_1, m_2\} \geq 2$; and for $2 \leq j \leq l$:

$$\operatorname{ord}_{\mathfrak{p}_{j},\phi} \omega_{1} \geq (n - 4 - m_{j})\operatorname{ord}_{\mathfrak{p}_{j},\phi} (X - \alpha_{j})$$

$$\geq \{m_{1} + (\sum_{2 \leq i \neq j \leq l}^{l} m_{i}) - 3\}\operatorname{ord}_{\mathfrak{p}_{j},\phi} (X - \alpha_{j})$$

which is non-negative if $l \geq 4$, or l = 3 and $m_1 \geq 2$, or l = 2 and $m_1 \geq 3$. Thus, under the hypothesis of (ii), ω_1 is also regular on C.

Similarly, we get:

$$\operatorname{ord}_{\mathfrak{p}_{2},\phi} \omega_{2} = (n-4)\operatorname{ord}_{\mathfrak{p}_{2},\phi} (X-Y) - (m_{2}-1)\operatorname{ord}_{\mathfrak{p}_{2},\phi} (X-\alpha_{2})$$

$$\geq (n-3-m_{2})\operatorname{ord}_{\mathfrak{p}_{2},\phi} (X-\alpha_{2})$$

$$\geq \{(\sum_{i=2}^{l} m_{i}) + m_{1} - 2\}\operatorname{ord}_{\mathfrak{p}_{2},\phi} (X-\alpha_{2}) \geq 0$$

provided that $l \geq 3$ or l = 2 and $m_1 \geq 2$;

$$\operatorname{ord}_{\mathfrak{p}_{1},\phi}\omega_{2} \geq (n-4-m_{1})\operatorname{ord}_{\mathfrak{p}_{1},\phi}(X-\alpha_{1})$$
$$\geq \{(\sum_{i=2}^{l}m_{i})-3\}\operatorname{ord}_{\mathfrak{p}_{1},\phi}(X-\alpha_{1})$$

which is non-negative if $l \ge 4$, or l = 3 and $m_2 + m_3 \ge 3$, or l = 2 and $m_2 \ge 3$; and if $l \ge 3$, for $3 \le j \le l$:

$$\operatorname{ord}_{\mathfrak{p}_{j},\phi} \omega_{2} \geq (n - 4 - m_{j})\operatorname{ord}_{\mathfrak{p}_{j},\phi} (X - \alpha_{j})$$

$$\geq \{m_{1} + (\sum_{2 \leq i \neq j \leq l}^{l} m_{i}) - 3\}\operatorname{ord}_{\mathfrak{p}_{j},\phi} (X - \alpha_{j})$$

which is non-negative if $l \geq 4$, or l = 3 and $m_1 \geq 2$. Thus, under the hypothesis of (ii), ω_2 is also regular on C except when (a) l = 3 and $m_2 = m_3 = 1$; (b) l = 2 and $m_2 = 2$.

If $l \geq 4$; l = 3, $m_1 \geq 2$ and $m_2 + m_3 \geq 3$; or l = 2 and $m_2 \geq 3$, ω_1 and ω_2 are both regular and we claim that they are linearly independent on C. For any constants a and b,

$$a\omega_1 + b\omega_2 = (X - Y)^{n-4} (a(X - \alpha_1 Z) + b(X - \alpha_2 Z))\eta,$$

hence it is not identically zero on any component of C because X - Y and $\alpha X + \beta Z$ are not a linear factor of F(X, Y, Z).

If l=3, $m_1 \geq 2$ and $m_2+m_3=2$; or l=2 and $m_2=2$, ω_0 and ω_1 are both regular and we claim that they are linearly independent on C. We note that under these assumptions, $n=m_1+3$ and after a linear transformation we may assume that $\alpha_1=0$. Hence P(X) can be written as

(3.13)
$$P(X) = P(0) + b_0 X^{m_1+1} + b_1 X^{m_1+2} + X^{m_1+3}$$

with $b_0 \neq 0$. Moreover, if $b_1 = 0$, then m_1 is even; otherwise $\pm \alpha_2$ with $\alpha_2^2 = -b_0(m_1+1)/(m_1+3)$ are the other two solutions of P'(X) = 0 and $P(\alpha_2) = P(-\alpha_2)$ which contradicts the separation condition. For any constants a and b,

$$a\omega_0 + b\omega_1 = (X - Y)^{n-4}(a(X - Y) + bX)\eta,$$

hence it is not identically zero on any component of C if a(X - Y) + bX is not a linear factor of F(X,Y,Z). Since X - Y and X is not a factor of F(X,Y,Z), we may assume that a = 1 and $b \neq 0$. Write (X - Y) + bX = (b + 1)X - Y. If it is a linear factor of F(X,Y,Z), then P(X) = P((b+1)X), and by (3.13) we get, $(b+1)^{m_1+1} = (b+1)^{m_1+3} = 1$ and $(b+1)^{m_1+2} = 1$ if $b_1 \neq 0$ which is always the case if m_1 is odd as noted above. Therefore if m_1 is odd then b = 0 since it is the only solution to satisfy $(b+1)^{m_1+1} = (b+1)^{m_1+2} = (b+1)^{m_1+3} = 1$. If m_1 is even, b = 0 is clearly also the only solution to satisfy $(b+1)^{m_1+1} = (b+1)^{m_1+3} = 1$. Hence we conclude that b = 0 which leads to a contradiction. Thus we have constructed two regular 1-forms which are linearly independent on each component of C. This implies that C is Brody hyperbolic.

4. The case
$$P(X) = cP(Y), c \neq 0, 1$$

In this section we shall deal with the curves $C_c = [F_c(X,Y,Z) = 0]$ where $F_c(X,Y,Z)$ is the homogenization of the polynomial P(X) - cP(Y), for $c \neq 0,1$. As in the preceding section let $P'(X) = \lambda(X - \alpha_1)^{m_1}...(X - \alpha_l)^{m_l}, m_i > 0, \lambda \neq 0$ and $\alpha_i \neq \alpha_j$, if $i \neq j$. We have, by direct calculation

(4.1)
$$\begin{cases} \frac{\partial F_c}{\partial X}(X, Y, Z) = P'(X, Z) = \lambda (X - \alpha_1 Z)^{m_1} ... (X - \alpha_l Z)^{m_l}, \\ \frac{\partial F_c}{\partial Y}(X, Y, Z) = -cP'(Y, Z) = -c\lambda (Y - \alpha_1 Z)^{m_1} ... (Y - \alpha_l Z)^{m_l}, \\ \frac{\partial F_c}{\partial Z}(X, Y, Z) = (n - m)a_m Z^{n-m-1} (X^m - cY^m + ZH_{m-1}) \end{cases}$$

on the curve $C_c = [F_c(X, Y, Z) = 0]$ and where P'(X, Z), P'(Y, Z) are as given in (3.6). We have the following analogue of Proposition 3.1 (and with essentially the same proof).

Proposition 4.1. Let $P(X) = X^n + a_m X^m + \cdots + a_0, a_m \neq 0$ be a polynomial of degree n. Suppose that the curve $C_c = [F_c(X, Y, Z) = 0]$ has no linear component then C_c is algebraically hyperbolic if $n - m \geq 3$ and Brody hyperbolic if $n - m \geq 4$.

Proof. As in Proposition 3.1, we have

(4.2)
$$\gamma := \frac{W(X,Y)}{\frac{\partial F_c}{\partial Z}} = \frac{W(Y,Z)}{\frac{\partial F_c}{\partial Y}} = \frac{W(Z,X)}{\frac{\partial F_c}{\partial Y}}$$

on $\pi^{-1}(C_c)$. By (4.1) we also have

$$\gamma = \frac{W(X,Y)}{(n-m)a_m Z^{n-m-1}(X^m - cY^m + ZH_{m-1})} = \frac{W(Y,Z)}{P'(X,Z)} = \frac{W(X,Z)}{-cP'(Y,Z)}.$$

If $n - m \ge 3$, take $G(X, Y, Z) = Z^{n-m-3}(X^m - cY^m + ZH_{m-1})$. Then

$$\theta := G(X, Y, Z)\gamma = \frac{1}{a_m(n-m)Z^2}W(X, Y)$$

is a well-defined rational 1-form on $C_c \subset \mathbf{P}^2$. By construction the only possibility singularity of θ is along Z = 0. However, as in the proof of Proposition 3.1, we conclude that a pole of θ along Z = 0 implies that X = Y = 0 as well. Obviously this is impossible hence θ is regular on C_c .

If $n-m \ge 4$, take $K(X,Y,Z) = Z^{n-m-4}(X^m-cY^m+ZH_{m-1})$. Then as in the proof of Proposition 3.1, for any linear form L(X,Y,Z)

$$\omega := L(X,Y,Z)K(X,Y,Z)\gamma = \frac{L(X,Y,Z)}{a_m(n-m)Z^3}W(X,Y)$$

is well-defined and regular on C_c . By taking L(X, Y, Z) = X, Y, Z respectively we get 3 regular 1-forms linearly independent on each component of C_c

$$\frac{XW(X,Y)}{a_m(n-m)Z^3}, \ \frac{YW(X,Y)}{a_m(n-m)Z^3}, \frac{W(X,Y)}{a_m(n-m)Z^2}.$$

The Proposition follows immediately from this.

We can say more if P satisfies the separation condition. The form γ defined in (4.2) may be expressed, via (4.1), on $\pi^{-1}(C_c)$ as:

$$(4.3) \qquad \gamma = \frac{W(Y,Z)}{\lambda(X - \alpha_1 Z)^{m_1} \dots (X - \alpha_l Z)^{m_l}} \equiv \frac{W(X,Z)}{-c\lambda(Y - \alpha_1 Z)^{m_1} \dots (Y - \alpha_l Z)^{m_l}}.$$

As observed previously there is no pole along $[Z = 0] \cap \pi^{-1}(C_c)$ hence the only possible poles with $Z \neq 0$ are $\pi^{-1}(\alpha_i, \alpha_j, 1)$ satisfying $P(\alpha_i) = cP(\alpha_j)$ (this is equivalent to the condition that $(\alpha_i, \alpha_j, 1) \in C_c$). The separation condition (3.4) implies that there is at most one j satisfying this condition. From now on, we write $j = \tau(i)$ if $(\alpha_i, \alpha_j, 1) \in C_c$. Since $c \neq 0, 1$ we have

(4.4)
$$\tau(i) \neq i \text{ and } \tau(i) \neq \tau(j) \text{ if } i \neq j.$$

We first establish a technique lemma which will be used through this section.

Lemma 4.2. Let P be a polynomial satisfying (3.4). Suppose that $P(\alpha_i) = cP(\alpha_{\tau(i)})$. Let $u \ge \max\{m_i, m_{\tau(i)}\}$ and H_j , $1 \le j \le u$, be linear forms in X, Y, Z such that $H_j(\alpha_i, \alpha_{\tau(i)}, 1) = 0$. Then

$$\eta := \frac{W(Y,Z) \prod_{j=1}^{u} H_j}{(X - \alpha_i Z)^{m_i}}$$

is regular on $\pi^{-1}(C_c)$.

Proof. For simplicity of notation, assume that i=1 and $\tau(1)=l$; and let $\mathfrak{p}_1=(\alpha_1,\alpha_l,1)$. It suffices to check the regularity of η along $\pi^{-1}(\mathfrak{p}_1)$. Since $H_j(\alpha_1,\alpha_l,1)=0$, $H_j=a(X-\alpha_1Z)+b(Y-\alpha_lZ)$ for some $a,b\in \mathbb{C}$. Therefore,

$$\operatorname{ord}_{\mathfrak{p}_1,\phi} H_i \ge \min\{\operatorname{ord}_{\mathfrak{p}_1,\phi}(X - \alpha_1 Z), \operatorname{ord}_{\mathfrak{p}_1,\phi}(Y - \alpha_l Z)\},$$

for any local parametrization ϕ of C_c at \mathfrak{p}_1 . If $\operatorname{ord}_{\mathfrak{p}_1,\phi}(X-\alpha_1 Z) \leq \operatorname{ord}_{\mathfrak{p}_1,\phi}(Y-\alpha_l Z)$, then $\operatorname{ord}_{\mathfrak{p}_1,\phi}H_j = \operatorname{ord}_{\mathfrak{p}_1,\phi}(X-\alpha_1 Z)$. Then it is clear that η is regular on \mathfrak{p}_1 since $u \geq m_1$.

If $\operatorname{ord}_{\mathfrak{p}_1,\phi}(X-\alpha_1 Z) > \operatorname{ord}_{\mathfrak{p}_1,\phi}(Y-\alpha_l Z)$, then $\operatorname{ord}_{\mathfrak{p}_1,\phi}H_j = \operatorname{ord}_{\mathfrak{p}_1,\phi}(Y-\alpha_l Z)$. By (4.3) we have

$$\frac{W(Y,Z)(Y-\alpha_{l}Z)^{u}}{(X-\alpha_{1}Z)^{m_{1}}} \equiv \frac{W(Z,X)(Y-\alpha_{l}Z)^{u-m_{l}}(X-\alpha_{2}Z)^{m_{2}}\dots(X-\alpha_{l}Z)^{m_{l}}}{-c(Y-\alpha_{2}Z)^{m_{2}}\dots(Y-\alpha_{l-1}Z)^{m_{l-1}}}.$$

Therefore, η is regular on $\pi^{-1}(\mathfrak{p}_1)$ since $u \geq m_l$.

The following result (the case in which $\tau(i)$ exists for all i) was first established in [8] and [1] using the truncated version of the Second Main Theorem of Nevanlinna Theory; we include a simpler proof below.

Lemma 4.3. Let P be a polynomial satisfying the separation condition (3.4) and assume that for each $1 \le i \le l$ there exists $\tau(i)$ such that $(\alpha_i, \alpha_{\tau(i)}, 1) \in C_c = [F_c(X, Y, Z) = 0]$, i.e., $P(\alpha_i) = cP(\alpha_{\tau(i)})$ and F_c is the homogenization of P(X) - cP(Y), $c \ne 0, 1$. If, in addition, the curve C_c has no linear component then

- (i) C_c is algebraically hyperbolic if $|m_i m_{\tau(i)}| \geq 2$ for some $1 \leq i \leq l$;
- (ii) C_c is Brody hyperbolic if $|m_i m_{\tau(i)}| \geq 3$ for some $1 \leq i \leq l$.

Proof. The condition in (i) implies that there is some i such that $m_i \geq 3$. There is no loss of generality in assuming that i = 1. The rational 1-form

(4.5)
$$\omega := \frac{(Y - \alpha_{\tau(1)}Z)^{m_1 - 2}}{(X - \alpha_1 Z)^{m_1}} W(Y, Z)$$

is well-defined on C_c . By (4.3) ω may be expressed as:

$$\omega = \lambda (Y - \alpha_{\tau(1)} Z)^{m_1 - 2} \prod_{i=2}^{l} (X - \alpha_i Z)^{m_i} \gamma$$

$$= \frac{(Y - \alpha_{\tau(1)} Z)^{m_1 - m_{\tau(1)} - 2} \prod_{i=2}^{l} (X - \alpha_i Z)^{m_i}}{\prod_{i=2}^{l} (Y - \alpha_{\tau(i)} Z)^{m_{\tau(i)}}} W(X, Z)$$

where γ is defined in (4.2). The first expression implies that the only possible poles of ω on C_c are contained in $[X=\alpha_1]\cap C_c$ while the second expression implies that the only possible poles of ω on C_c are contained in $[Y=\alpha_{\tau(i)}]\cap C_c, i\geq 2$, provided that $m_1-m_{\tau(1)}\geq 2$. This shows that ω is regular on C_c because none of the points $\{(\alpha_1,\alpha_{\tau(i)},1)\mid 2\leq i\leq l\}$ is in C_c . Furthermore, since none of the linear functions $Y-\alpha_{\tau(1)}Z, X-\alpha_{\tau(j)}Z, j\geq 2$ is a factor of F_c , we conclude that ω is non-trivial on any component of C_c . This establishes the first assertion if $m_1-m_{\tau(1)}\geq 2$. A similar argument applied to

$$\omega = \frac{W(X,Z)(X - \alpha_1 Z)^{m_{\tau(1)} - 2}}{(Y - \alpha_{\tau(1)} Z)^{m_{\tau(1)}}} = -c\lambda (X - \alpha_1 Z)^{m_{\tau(1)} - 2} \prod_{i=2}^{l} (Y - \alpha_{\tau(i)} Z)^{m_{\tau(i)}} \gamma$$

establishes the first assertion if $m_{\tau(1)} - m_1 \ge 2$.

For assertion (ii) there is some i such that $m_i \geq 4$. There is no loss of generality in assuming that i = 1. If $m_1 - m_{\tau(1)} \geq 3$ (resp. $m_{\tau(1)} - m_1 \geq 3$) then an argument similar to the one given above shows that

$$\omega_1 := X \frac{W(Y, Z)(Y - \alpha_{\tau(1)}Z)^{m_1 - 3}}{(X - \alpha_1 Z)^{m_1}} \text{ and } \omega_2 := Y \frac{W(Y, Z)(Y - \alpha_{\tau(1)}Z)^{m_1 - 3}}{(X - \alpha_1 Z)^{m_1}}$$

(resp. take

$$\omega_1 := X \frac{W(X,Z)(X-\alpha_1 Z)^{m_{\tau(1)}-3}}{(Y-\alpha_{\tau(1)} Z)^{m_{\tau(1)}}} \ \ \text{and} \ \ \omega_2 := Y \frac{W(X,Z)(X-\alpha_1 Z)^{m_{\tau(1)}-3}}{(Y-\alpha_{\tau(1)} Z)^{m_{\tau(1)}}})$$

are well-defined regular 1-forms non-trivial and linearly independent on every component of C_c .

If the zero set of P(X) is affine rigid then the condition that C_c has no linear component is satisfied for all $c \neq 0, 1$.

In what follows let L_{ij} , $1 \le i \ne j \le l$, be the linear form defined by

(4.6)
$$L_{ij} := (Y - \alpha_{\tau(j)}Z) - \frac{\alpha_{\tau(i)} - \alpha_{\tau(j)}}{\alpha_i - \alpha_j}(X - \alpha_j Z)$$

provided that $\tau(i)$ and $\tau(j)$ exist, i.e., $P(\alpha_i) = cP(\alpha_{\tau(i)})$ and $P(\alpha_j) = cP(\alpha_{\tau(j)})$. Observe that L_{ij} may also be expressed as

(4.7)
$$L_{ij} = (Y - \alpha_{\tau(i)}Z) - \frac{\alpha_{\tau(i)} - \alpha_{\tau(j)}}{\alpha_i - \alpha_j}(X - \alpha_i Z).$$

Assuming that C_c has no linear component then L_{ij} is not identically zero on any component of C_c . At each point $\mathfrak{p}_i = (\alpha_i, \alpha_{\tau(i)}, 1) \in C_c$ and each local parameterization at \mathfrak{p}_i , we infer from (4.7) that

$$(4.8) \operatorname{ord}_{\mathfrak{p}_{i},\phi} L_{ij} \ge \min\{\operatorname{ord}_{\mathfrak{p}_{i},\phi}(X - \alpha_{i}Z), \operatorname{ord}_{\mathfrak{p}_{i},\phi}(Y - \alpha_{\tau(i)}Z)\},$$

and, analogously (from (4.6)) that for each local parameterization ϕ at \mathfrak{p}_i

$$(4.9) \operatorname{ord}_{\mathfrak{p}_{i},\phi} L_{ij} \ge \min \{ \operatorname{ord}_{\mathfrak{p}_{i},\phi} (X - \alpha_{j} Z), \operatorname{ord}_{\mathfrak{p}_{i},\phi} (Y - \alpha_{\tau(i)} Z) \}.$$

By (3.3) we have the following expansion of P(X):

$$P(X) = P(\alpha_i) + \sum_{j=m_i+1}^{n} b_{i,j} (X - \alpha_i Z)^j$$

where $b_{i,m_i+1} \neq 0$ and $b_{i,n} \neq 0$. If $\mathfrak{p}_i = (\alpha_i, \alpha_{\tau(i)}, 1) \in C_c$, then $F_c(X, Y, Z)$ can be expressed in terms of $X - \alpha_i Z$ and $Y - \alpha_{\tau(i)} Z$ as (compare (3.5))

$$F_c(X, Y, Z) = \sum_{j=m_i+1}^n b_{i,j} (X - \alpha_i Z)^j - c \sum_{j=m_{\tau(i)}+1}^n b_{\tau(i),j} (Y - \alpha_{\tau(i)} Z)^j.$$

Let ϕ be a local parameterization of C_c at \mathfrak{p}_i , we see from this expression that

$$(4.10) (m_i + 1) \operatorname{ord}_{\mathfrak{p}_i, \phi}(X - \alpha_i Z) = (m_{\tau(i)} + 1) \operatorname{ord}_{\mathfrak{p}_i, \phi}(Y - \alpha_{\tau(i)} Z).$$

The following Lemma is convenient in establishing the regularity of certain rational forms to be constructed in the proof of Proposition 4.7.

Lemma 4.4. Assume that $L_{ij}(see~(4.6)), i \neq j$, is defined. If $m_{\tau(i)} \leq m_i$ then for any local parameterization ϕ of C_c at the point $\mathfrak{p}_i = (\alpha_i, \alpha_{\tau(i)}, 1)$, we have

$$ord_{\mathfrak{p}_{i},\phi}(X - \alpha_{i}Z) \leq ord_{\mathfrak{p}_{i},\phi}(Y - \alpha_{\tau(i)}Z)$$

and $ord_{\mathfrak{p}_i,\phi}L_{ij}=ord_{\mathfrak{p}_i,\phi}(X-\alpha_i Z),$ consequently, $ord_{\mathfrak{p}_i,\phi}L_{ij}/(X-\alpha_i Z)\geq 0.$

Proof. The assumption together with (4.10) imply that

$$\operatorname{ord}_{\mathfrak{p}_i,\phi}(X - \alpha_i Z) \le \operatorname{ord}_{\mathfrak{p}_i,\phi}(Y - \alpha_{\tau(i)} Z)$$

and (4.8) implies that $\operatorname{ord}_{\mathfrak{p}_i,\phi}L_{ij} = \operatorname{ord}_{\mathfrak{p}_i,\phi}(X - \alpha_i Z)$.

Lemma 4.5. Let P be a polynomial satisfying the separation condition (3.4). If $l \geq 2$ and the curve C_c has no linear component then

- (i) it is algebraically hyperbolic if either of the following conditions holds,
 - (a) there exists an index i_0 such that $m_{i_0} \geq 2$ and $(\alpha_{i_0}, \alpha_j, 1) \notin C_c$ for $1 \leq j \leq l$;
 - (b) there exist indices i_1 and i_2 such that $m_{i_1} = m_{i_2} = 1$ and $(\alpha_{i_k}, \alpha_j, 1), \notin$ C_c for $1 \le j \le l$ and k = 1, 2;
- (ii) it is Brody hyperbolic if either of the following conditions holds,
 - (a) there exists an index i_0 such that $m_{i_0} \geq 3$ and $(\alpha_{i_0}, \alpha_j, 1) \notin C_c$ for $1 \leq j \leq l$;
 - (b) there exists indices i_0 and i_1 such that $m_{i_1} \ge m_{i_0} = 2$ and $(\alpha_{i_0}, \alpha_j, 1) \notin C_c$ for $1 \le j \le l$;
 - (c) there exist indices i_1 and i_2 such that $m_{i_1} + m_{i_2} = 3$ and $(\alpha_{i_k}, \alpha_j, 1), \notin$ C_c for $1 \le j \le l$ and k = 1, 2;
 - (d) $l \geq 3$ and there exist indices i_1 and i_2 such that $m_{i_1} = m_{i_2} = 1$ and $(\alpha_{i_k}, \alpha_j, 1), \notin C_c$ for $1 \leq j \leq l$ and k = 1, 2.

Proof. From (4.3) we see that if $(\alpha_{i_0}, \alpha_j, 1) \notin C_c$ for all $1 \leq j \leq l$, then γ (as defined by (4.3)) is regular along $\pi^{-1}(C_c) \cap [X - \alpha_{i_0}Z = 0]$. The rational 1-form

$$\eta = \frac{W(Y,Z)}{(X - \alpha_{i_0}Z)^2} = \lambda (X - \alpha_{i_0}Z)^{m_{i_0}-2} \prod_{1 \le i \ne i_0 \le l} (X - \alpha_i Z)^{m_i} \gamma$$

is well-defined on C_c and, as $m_{i_0} \geq 2$, it is also regular on C_c . This proves the assertion (ia). Analogously, if (ib) is satisfied then γ is regular along $\pi^{-1}(C_c) \cap [X - \alpha_{i_1}Z = 0]$ and also along $\pi^{-1}(C_c) \cap [X - \alpha_{i_2}Z = 0]$ hence the 1-form

$$\eta = \frac{W(Y,Z)}{(X - \alpha_{i_1}Z)(X - \alpha_{i_2}Z)} = \lambda \prod_{1 \le i \ne i_1, i_2 \le l} (X - \alpha_i Z)^{m_i} \gamma$$

is well-defined and regular on C_c . This completes the proof of (ib).

Similarly, if $(\alpha_{i_0}, \alpha_j, 1) \notin C_c$ for all $1 \leq j \leq l$, and $m_{i_0} \geq 3$, then

$$\eta_1 = \frac{XW(Y,Z)}{(X - \alpha_{i_0}Z)^3} = \lambda X(X - \alpha_{i_0}Z)^{m_{i_0}-3} \prod_{1 \le i \ne i_0 \le l} (X - \alpha_i Z)^{m_i} \gamma,$$

$$\eta_2 = \frac{YW(Y,Z)}{(X - \alpha_{i_0}Z)^3} = \lambda Y(X - \alpha_{i_0}Z)^{m_{i_0}-3} \prod_{1 \le i \ne i_0 \le l} (X - \alpha_i Z)^{m_i} \gamma$$

are two linearly independent regular 1-forms on C_c . This proves the assertion (iia). If $m_{i_1} \geq m_{i_0} = 2$, then there exists an index $i_k \neq i_0$ such that $m_{i_k} = \max_{1 \leq i \leq l} m_i$. Suppose that $(\alpha_{i_0}, \alpha_j, 1) \notin C_c$ for $1 \leq j \leq l$. Let

$$\eta_1 = \frac{W(Y,Z)}{(X - \alpha_{i_0} Z)^2};$$

$$\eta_2 = \frac{W(Y,Z)}{(X - \alpha_{i_0} Z)(X - \alpha_{i_k} Z)} \quad \text{if } (\alpha_{i_k}, \alpha_j, 1) \not\in C_c \text{ for } 1 \le j \le l;$$

and

$$\eta_2 = \frac{(Y - \alpha_{\tau(i_k)}Z)W(Y,Z)}{(X - \alpha_{i_0}Z)^2(X - \alpha_{i_k}Z)} \quad \text{if } (\alpha_{i_k},\alpha_{\tau(i_k)},1) \in C_c.$$

Then η_1 and η_2 are two linearly independent regular 1-forms on C_c . We note the regularity of the second η_2 is due to Lemma 4.4 since $m_{i_k} = \max_{1 \leq i \leq l} m_i$ and $\operatorname{ord}_{\mathfrak{p}_j,\phi}(X - \alpha_{i_k}Z) \leq \operatorname{ord}_{\mathfrak{p}_j,\phi}(Y - \alpha_{\tau(i_k)}Z)$ from (4.10). This completes the proof of (iib). For (iic) we may assume that $m_{i_1} = 2$ and $m_{i_2} = 1$ since $m_{i_1} = 3$ or $m_{i_2} = 3$ is covered by (iia). Then

$$\eta_1 = \frac{W(Y, Z)}{(X - \alpha_{i_1} Z)^2},
\eta_2 = \frac{W(Y, Z)}{(X - \alpha_{i_1} Z)(X - \alpha_{i_2} Z)}$$

are two linearly independent regular 1-forms on C_c .

If $m_{i_1} = m_{i_2} = 1$ and $l \ge 3$, then there exists an index i_t different from i_1 and i_2 such that $m_{i_t} = \max_{1 \le i \le l} m_i$. Then, similarly

$$\eta_1 = \frac{W(Y, Z)}{(X - \alpha_{i_1} Z)(X - \alpha_{i_2} Z)};$$

$$\eta_2 = \frac{W(Y, Z)}{(X - \alpha_{i_1} Z)(X - \alpha_{i_1} Z)} \quad \text{if } (\alpha_{i_t}, \alpha_j, 1), \notin C_c \text{ for } 1 \le j \le l;$$

and

$$\eta_2 = \frac{(Y - \alpha_{\tau(i_t)}Z)W(Y,Z)}{(X - \alpha_{i_1}Z)(X - \alpha_{i_2}Z)(X - \alpha_{i_t}Z)} \quad \text{if } (\alpha_{i_t}, \alpha_{\tau(i_t)}, 1) \in C_c.$$

are two linearly independent regular 1-forms on C_c . This proves (iid).

Remark 4.6. The preceding Lemma implies that (under the assumption that the polynomial P satisfies condition (3.4)), in deciding whether C_c is algebraically hyperbolic, we may assume that for each $1 \le i \le l$ there exists another index $\tau(i)$ such that $(\alpha_i, \alpha_{\tau(i)}, 1) \in C_c$ for all but one index i and, in which case, $m_i = 1$.

Proposition 4.7. Let P be a polynomial of degree $n \geq 4$ satisfying the separation condition (3.4) and assume that the curve $C_c = [F_c(X, Y, Z) = 0], c \neq 0, 1$, has no linear component. Rearrange α_i so that $m_1 \geq m_2 \cdots \geq m_l$. Then

(i) C_c is algebraically hyperbolic if (a) $l \geq 2$ and $m_2 \geq 2$, or (b) $l \geq 3$ and $m_2 = 1$ except when $l = 3, m_1 = m_2 = m_3 = 1$ with

$$\frac{P(\alpha_1)}{P(\alpha_2)} = \frac{P(\alpha_2)}{P(\alpha_3)} = \frac{P(\alpha_3)}{P(\alpha_1)} = c \text{ or } \frac{1}{c};$$

(ii) C_c is Brody hyperbolic if (a) $l \geq 2$ and $m_2 \geq 2$ except when l = 2 and $m_1 = m_2 = 2$, or (b) $l \geq 3$ and $m_2 = 1$ except when l = 3 and $m_1 = m_2 = m_3 = 1$.

Proof. For case (ia) we have $m_1 \geq m_2 \geq 2$, and hence $m_1 + m_2 - 2 \geq m_1 \geq m_i$, i = 1, ..., l. By Lemma 4.5 and Remark 4.6 we may assume that $\tau(1)$ and $\tau(2)$ exist such that $\mathfrak{p}_i = (\alpha_i, \alpha_{\tau(i)}, 1) \in C_c$ for i = 1, 2. Thus L_{12} is defined. The rational 1-form

(4.11)
$$\omega_1 := \frac{L_{12}^{m_1 + m_2 - 2}}{(X - \alpha_1 Z)^{m_1} (X - \alpha_2 Z)^{m_2}} W(Y, Z)$$

is well-defined (the denominator of the coefficient of the Wronskian is two degree higher than the numerator) on C_c . We claim that ω_1 is regular. It suffices to check regularity at \mathfrak{p}_i , i=1,2. To check ω_1 is regular at \mathfrak{p}_1 , it suffices to check the rational 1-form

$$\eta = \frac{L_{12}^{m_1 + m_2 - 2}}{(X - \alpha_1 Z)^{m_1}} W(Y, Z)$$

is regular at $\pi^{-1}(\mathfrak{p}_1)$. Since $m_1 + m_2 - 2 \ge m_1 \ge m_i$, i = 1, ..., l, the later assertion is an implication of Lemma 4.2. The regularity of ω_1 at \mathfrak{p}_2 can be checked similarly. Thus ω_1 is regular on C_c and (ia) of the Proposition is established.

Next we consider the case (iia). Since $m_1 \geq m_2 \geq 2$, by Lemma 4.5 we may assume that $\tau(1)$ and $\tau(2)$ exist such that $\mathfrak{p}_i = (\alpha_i, \alpha_{\tau(i)}, 1) \in C_c$ for i = 1, 2. Thus L_{12} is defined. Moreover, if $m_2 = 2$, then by Lemma 4.3 we only need to consider when $m_1 \leq 4$. By the preceding case we already have a regular 1-form ω_1 , defined by (4.11), on C_c . We look for another regular 1-form ω_2 on C_c linearly independent to ω_1 . If $m_2 \geq 2$ and $m_1 \geq 3$, the rational 1-form

$$\omega_2 := \begin{cases} \frac{W(Y,Z)L_{12}^{m_1+m_2-3}(X-\alpha_1Z)}{(X-\alpha_1Z)^{m_1}(X-\alpha_2Z)^{m_2}}, & \text{if} \quad m_2 \ge 3; \\ \frac{W(Y,Z)L_{12}^{m_1-1}(X-\alpha_2Z)}{(X-\alpha_1Z)^{m_1}(X-\alpha_2Z)^2}, & \text{if} \quad m_2 = 2, \text{ and } 3 \le m_1 \le 4. \end{cases}$$

by construction, is well-defined on C_c . Moreover, it is clear that ω_1 and ω_2 are linearly independent on C_c since C_c has no linear component. We claim that ω_2 is actually regular on C_c .

For the case $m_2 \geq 3$ we have $m_1 + m_2 - 3 \geq m_1 \geq m_i$ for all i. Similar to the previous proof of the regularity of ω_1 , we see that ω_2 is regular on C_c by Lemma 4.2. For the case $m_2 = 2$ and $3 \leq m_1 \leq 4$, in the numerator of ω_2 there are $m_1(\geq m_i)$ linear forms vanishing at \mathfrak{p}_2 which implies, by Lemma 4.2, that ω_2 is regular at \mathfrak{p}_2 .

We now check the regularity of ω_2 at \mathfrak{p}_1 . When $m_2 \geq 3$, $m_1 + m_2 - 3 \geq m_1 \geq m_i$ and hence ω_2 is regular at \mathfrak{p}_1 by Lemma 4.2. We now consider when $m_2 = 2$ and $m_1 \geq 3$. We first see that by (4.10),

$$(m_1+1)\operatorname{ord}_{\mathfrak{p}_1,\phi}(X-\alpha_1 Z)=(m_{\tau(1)}+1)\operatorname{ord}_{\mathfrak{p}_1,\phi}(Y-\alpha_{\tau(1)} Z).$$

Since $m_2 = 2$, $m_{\tau(1)} = 2$ or 1. Thus, we infer that $\operatorname{ord}_{\mathfrak{p}_1,\phi} L_{12} = \operatorname{ord}_{\mathfrak{p}_1,\phi} (X - \alpha_1 Z) < \operatorname{ord}_{\mathfrak{p}_1,\phi} (Y - \alpha_{\tau(1)} Z)$. On the other hand, as $Z \equiv 1$ on a neighborhood of \mathfrak{p}_1 ,

$$\operatorname{ord}_{\mathfrak{p}_1,\phi}W(Y,Z) = \operatorname{ord}_{\mathfrak{p}_1,\phi}dY \ge \operatorname{ord}_{\mathfrak{p}_1,\phi}(Y - \alpha_{\tau(1)}Z) - 1.$$

Then

$$\operatorname{ord}_{\mathfrak{p}_{1},\phi}\omega_{2} = \operatorname{ord}_{\mathfrak{p}_{1},\phi}W(Y,Z) + (m_{1} - 1)\operatorname{ord}_{\mathfrak{p}_{1},\phi}L_{12} - m_{1}\operatorname{ord}_{\mathfrak{p}_{1},\phi}(X - \alpha_{1}Z)$$

$$\geq \operatorname{ord}_{\mathfrak{p}_{1},\phi}(Y - \alpha_{\tau(1)}Z) - 1 - \operatorname{ord}_{\mathfrak{p}_{1},\phi}(X - \alpha_{1}Z)$$

$$\geq \frac{m_{1} + 1}{m_{\tau(1)} + 1}\operatorname{ord}_{\mathfrak{p}_{1},\phi}(X - \alpha_{\tau(1)}Z) - 1 - \operatorname{ord}_{\mathfrak{p}_{1},\phi}(X - \alpha_{1}Z)$$

$$\geq \frac{m_{1} - m_{\tau(1)}}{m_{\tau(1)} + 1}\operatorname{ord}_{\mathfrak{p}_{1},\phi}(X - \alpha_{1}Z) - 1.$$

$$(4.12)$$

If $m_{\tau(1)}=1$, then $(m_1-m_{\tau(1)})/(m_{\tau(1)}+1)\geq 1$ and thus (4.12) is non-negative. If $m_{\tau(1)}=2$, then $(m_1-m_{\tau(1)})/(m_{\tau(1)}+1)=1/3$ or 2/3. Since in this case, $(m_1+1)\mathrm{ord}_{\mathfrak{p}_1,\phi}(X-\alpha_1Z)=3\mathrm{ord}_{\mathfrak{p}_1,\phi}(Y-\alpha_{\tau(1)}Z)$ with $m_1+1=4$ or 5, we infer that $\mathrm{ord}_{\mathfrak{p}_1,\phi}(X-\alpha_1Z)\geq 3$. Hence, (4.12) is also non-negative. This completes the proof for this case. Next we treat the case $m_2=2,m_1=2$ and $l\geq 3$. Recall that we may assumed that L_{12} is defined (as $m_1=m_2=2$). If there exists an index $\tau(3)$ such that $(\alpha_3,\alpha_{\tau(3)},1)\in C_c$ hence L_{23} and L_{31} are defined. The rational 1-form

$$\omega_2 = \frac{W(Y, Z)L_{12}L_{23}L_{31}}{(X - \alpha_1 Z)^2(X - \alpha_2 Z)^2(X - \alpha_3 Z)}$$

is then defined. We have $m_i \leq 2$ for all i, and for each \mathfrak{p}_i , i=1,2,3, there are two linear forms in the numerator of ω_2 vanishing at it. We infer from Lemma 4.2 that ω_2 is regular. To check that ω_1 and ω_2 are linearly independent is equivalent to show that the quadratic form $aL_{12}(X-\alpha_3Z)+bL_{23}L_{31},\ a,b\in \mathbf{C}$ is not a factor of F_c . Note that we may assume this quadratic form is irreducible since F_c has no linear factor. We have shown for the case (ia) that there is a regular 1-form in this case. Therefore, F_c cannot have any quadratic factor, and hence ω_1 and ω_2 are linearly independent. If there does not exist an index $\tau(3)$ such that $(\alpha_3, \alpha_{\tau(3)}, 1) \in C_c$ then we take

$$\omega_2 = \frac{W(Y, Z)L_{12}^2}{(X - \alpha_1 Z)^2 (X - \alpha_2 Z)(X - \alpha_3 Z)}$$

and it can be verified similarly via Lemma 4.2 that ω_1 and ω_2 are regular and linearly independent on any component of C_c . This completes the proof for the case $m_1 = m_2 = 2$ and $l \geq 3$.

It remains to deal with the case $m_2 = 1$ and $l \ge 3$ ((ib) and (iib)). In this case we have $m_i = 1$ for all $2 \le i \le l$. We separate the proof into two cases: (1) $m_1 \ge 2$ and (2) $m_1 = 1$. First we treat the case $m_1 \ge 2$. If there does not exist $\tau(1)$ such

that $(\alpha_1, \alpha_{\tau(1)}, 1) \in C_c$ then we may take

$$\omega_1 = \frac{W(Y, Z)}{(X - \alpha_1 Z)^2}$$

which is a regular 1-form on C_c . By Lemma 4.5, we may assume that for each $i \geq 2$ there exists an index $\tau(i)$ such that $(\alpha_i, \alpha_{\tau(i)}, 1) \in C_c$. Since $l \geq 3$ and $m_i = 1$ if $i \geq 2$, $m_{\tau(i)} = 1$ for some $i \geq 2$. Then we may take

$$\omega_2 = \frac{W(Y, Z)(Y - \alpha_{\tau(i)}Z)}{(X - \alpha_1 Z)^2 (X - \alpha_i Z)}$$

which is a regular 1-form on C_c (by (4.10)) and linearly independent to ω_1 . Thus we may assume that there exists $\tau(1)$ such that $\mathfrak{p}_1 = (\alpha_1, \alpha_{\tau(1)}, 1) \in C_c$. Let L be a linear form vanishing at \mathfrak{p}_1 . We first claim that

$$(4.13) \operatorname{ord}_{\mathfrak{p}_1,\phi}(W(Y,Z)L) \ge 2\operatorname{ord}_{\mathfrak{p}_1,\phi}(X - \alpha_1 Z).$$

Since $m_i = 1$ for $i \geq 2$, we have $m_{\tau(1)} = 1$, and

$$2\operatorname{ord}_{\mathfrak{p}_1,\phi}(Y - \alpha_{\tau(1)}Z) = (m_1 + 1)\operatorname{ord}_{\mathfrak{p}_1,\phi}(X - \alpha_1Z)$$

where $m_1 = 2$ or 3. Hence $\operatorname{ord}_{\mathfrak{p}_1,\phi}L = \operatorname{ord}_{\mathfrak{p}_1,\phi}(X - \alpha_1 Z)$ and if $m_1 = 2$ then $\operatorname{ord}_{\mathfrak{p}_1,\phi}(X - \alpha_1 Z) \geq 2$. Thus we have:

$$\operatorname{ord}_{\mathfrak{p}_{1},\phi}W(Y,Z) + \operatorname{ord}_{\mathfrak{p}_{1},\phi}L - 2\operatorname{ord}_{\mathfrak{p}_{1},\phi}(X - \alpha_{1}Z)$$

$$\geq \operatorname{ord}_{\mathfrak{p}_{1},\phi}(Y - \alpha_{\tau(1)}) - \operatorname{ord}_{\mathfrak{p}_{1},\phi}(X - \alpha_{1}) - 1$$

$$\geq \frac{m_{1} - 1}{2}\operatorname{ord}_{\mathfrak{p}_{1},\phi}(X - \alpha_{1}) - 1 \geq 0.$$

Next we claim if there exist one index $i_0 \geq 2$ such that $(\alpha_{i_0}, \alpha_j, 1) \notin C_c$ for all $1 \leq j \leq l$, then C_c is Brody hyperbolic. By (4.13)

$$\omega_{1} = \frac{W(Y, Z)(X - \alpha_{1}Z)}{(X - \alpha_{1}Z)^{2}(X - \alpha_{i_{0}}Z)},$$

$$\omega_{2} = \frac{W(Y, Z)(Y - \alpha_{\tau(1)}Z)}{(X - \alpha_{1}Z)^{2}(X - \alpha_{i_{0}}Z)}$$

are two regular, linearly independent 1-forms. Therefore we may assume that for each i there exists $\tau(i)$ such that $\mathfrak{p}_i = (\alpha_i, \alpha_{\tau(i)}, 1) \in C_c$. Since $i \geq 3$, and $m_i = 1$ for $i \geq 2$, then $m_{\tau(i_0)} = 1$ for some $i_0 \geq 2$. Assume that $i_0 = 2$. Then

$$\omega_1 := \frac{W(Y, Z)L_{12}}{(X - \alpha_1 Z)^2 (X - \alpha_2 Z)}$$

is regular at \mathfrak{p}_1 by (4.13), and is regular at \mathfrak{p}_2 by Lemma 4.2. Therefore, ω_1 is regular on C_c . Similarly,

$$\omega_2 := \frac{W(Y, Z)L_{13}L_{23}}{(X - \alpha_1 Z)^2 (X - \alpha_2 Z)(X - \alpha_3 Z)}$$

is regular at \mathfrak{p}_1 by (4.13), and is regular at \mathfrak{p}_2 and \mathfrak{p}_3 (since $m_i \leq 2$, for $i \geq 1$) by Lemma 4.2. This shows that ω_2 is regular on C_c . To show ω_1 and ω_2 are linearly independent, one can use the previous argument that F_c has no quadratic factor as ω_1 exists. Finally, we consider the case when there is no such index, i.e., for each i there exists an index $\tau(i)$ such that $(\alpha_i, \alpha_{\tau(i)}, 1) \in C_c$. If $l \geq 4$, then

$$\omega_1 := \frac{W(Y, Z)L_{12}L_{34}}{(X - \alpha_1 Z)(X - \alpha_2 Z)(X - \alpha_3 Z)(X - \alpha_4 Z)},$$
$$\omega_2 := \frac{W(Y, Z)L_{13}L_{24}}{(X - \alpha_1 Z)(X - \alpha_2 Z)(X - \alpha_3 Z)(X - \alpha_4 Z)}$$

are well-defined, linearly independent and regular on C_c , similarly. If l=3, then this case L_{12}, L_{13} and L_{23} are defined. There are only two possibilities: (I) $\tau(1) = 2, \tau(2) = 3, \tau(3) = 1$ or (II) $\tau(1) = 3, \tau(3) = 2, \tau(2) = 1$. For (I) we have

$$\frac{P(\alpha_1)}{P(\alpha_2)} = \frac{P(\alpha_2)}{P(\alpha_3)} = \frac{P(\alpha_3)}{P(\alpha_1)} = c$$

and for (II)

$$\frac{P(\alpha_1)}{P(\alpha_3)} = \frac{P(\alpha_3)}{P(\alpha_2)} = \frac{P(\alpha_2)}{P(\alpha_1)} = c.$$

This last identity is equivalent to

$$\frac{P(\alpha_1)}{P(\alpha_2)} = \frac{P(\alpha_2)}{P(\alpha_3)} = \frac{P(\alpha_3)}{P(\alpha_1)} = \frac{1}{c}.$$

Remark. The conditions $P(\alpha_1)/P(\alpha_2) = P(\alpha_2)/P(\alpha_3) = P(\alpha_3)/P(\alpha_1) = c$ imply that $P(\alpha_1) = cP(\alpha_2) = c^2P(\alpha_3) = c^3P(\alpha_1)$ or $P(\alpha_1) = c^{-1}P(\alpha_2) = c^{-2}P(\alpha_3) = c^{-3}P(\alpha_1)$ hence $c^3 = 1$ or $1/c^3 = 1$. Analogously the condition that

$$\frac{P(\alpha_1)}{P(\alpha_2)} = \frac{P(\alpha_2)}{P(\alpha_3)} = \frac{P(\alpha_3)}{P(\alpha_1)} = \frac{1}{c}.$$

imply that $1/c^2 + 1/c + 1 = 0$, i.e., c and 1/c are the two solutions of the equation $w^2 + w + 1$.

5. Proof of the Results

Proof of Theorem 1. From Proposition 3.1 and Proposition 4.1, we see that the assumption of affine rigidity on the set of zeros of P(X) assures that when $n-m \geq 4$ the regular 1-forms

$$\frac{XW(X,Y)}{a_{m}(n-m)Z^{3}},\;\frac{YW(X,Y)}{a_{m}(n-m)Z^{3}},\frac{W(X,Y)}{a_{m}(n-m)Z^{2}}$$

are not identically zero on any component of C and C_c . Clearly, X and Y are not linear factors of F(X,Y,Z) or $F_c(X,Y,Z)$, and W(X,Y) only vanishes identically on linear components such as aX - bY = 0. Therefore, it remains to show that

F(X,Y,Z) and $F_c(X,Y,Z)$, $c \neq 0,1$ has no linear factor of the form aX - bY if and only if the greatest common divisor of the non-zero indices in I is 1 and the greatest common divisor of the non-zero indices in J is also 1.

Let $I^* = I \setminus \{0\}$ and $J^* = J \setminus \{0\}$. Since 0 is divisible by all integers, the greatest common divisor of I (resp. J) equals the greatest common divisor of I^* (resp. J^*). Therefore we first assume that the greatest common divisor of the non-zero indices in I^* is 1 and the greatest common divisor of the non-zero indices in J^* is also 1. Recall from the remark of the theorem that if $n - m \geq 3$ and the greatest common divisor of the non-zero indices in J is 1, then $\#I \geq 3$. Then, $\#I^*$ and $\#J^*$ are at least 2. Under these assumptions, we want to show that F(X,Y,Z) and $F_c(X,Y,Z)$ has no linear factor of the form aX - bY.

Clearly, F(X,Y,Z) and $F_c(X,Y,Z)$ having no linear factor of the form aX-bY is the same as saying F(X,Y,1) and $F_c(X,Y,1)$ having no linear factor of the form aX-bY. It is also clear that, neither X nor Y is a linear factor of F(X,Y,1) nor $F_c(X,Y,1)$. Hence, we may assume that a=1 and $b\neq 0$. Observe that, as $F_c(X,X,1)=(1-c)P(X)\equiv 0,\ (c\neq 1)$ and $F(X,X,1)=P'(X)\not\equiv 0,\ X-Y$ is not a factor of $F_c(X,Y,1)$, nor F(X,Y,1). Hence $b\neq 1$. The condition that X-bY, $b\neq 1$, is a factor of F(X,Y,1) is equivalent to the condition that $F(bY,Y,1)\equiv 0$. Since

$$(X - Y)F(X, Y, 1) = P(X) - P(Y) = \sum_{i \in I^*} a_i (X^i - Y^i),$$

$$(bY - Y)F(bY, Y, 1) = \sum_{i \in I^*} a_i (b^i - 1)Y^i \equiv 0.$$

Hence, $b^i = 1$ for all $i \in I^*$. Since $\#I^* \ge 2$ and the greatest common divisor of indices in I^* is 1, we can find integers n_i , $i \in I^*$ such that $\sum_{i \in I^*} i n_i = 1$. Therefore, $b = \prod_{i \in I^*} b^{in_i} = 1$ which contradicts our assumption on $b \ne 1$.

Suppose that X-bY is a factor of $F_c(X,Y,1), c \neq 0, 1$, then $P(bY)-cP(Y) \equiv 0$. Therefore

$$\sum_{i \in I} a_i (b^i - c) Y^i \equiv 0.$$

This implies that $b^i=c$ for all $i\in I$. Therefore $b^{i-l}=1$ for all $i\in I$, where $l=\min\{i\mid i\in I\}$. This is equivalent to saying that $b^j=1$ for all $j\in J^*$. Since $\#J^*\geq 2$ and the greatest common divisor of indices in J^* is 1, similarly, we get b=1 which is impossible.

Conversely, suppose that the greatest common divisor of the indices in I^* is r > 1. Then $i = rc_i$ for each $i \in I^*$. Then

$$F(X, Y, 1) = \sum_{i \in I^*} a_i \frac{X^{rc_i} - Y^{rc_i}}{X - Y}.$$

Clearly, $(X^r - Y^r)/X - Y$ is a factor of F(X, Y, 1). In particularly, let b be a primitive r-root of unity. Then X - bY is a linear factor of F(X, Y, 1).

Similarly, suppose that the greatest common divisor of the non-zero indices in J^* is r > 1. Then $i - l = \alpha_i r$ for all $i \in I$ and $i \neq l = \min\{i \mid i \in I\}$. Let b be a primitive r-th root of unity, and take $c = b^l$. We consider first that $b^l \neq 1$, i.e. $c \neq 1$. Then for $i \in I$ and $i \neq l$, $b^{i-l} = b^{\alpha_i r} = 1$, and $b^i = b^{i-l}b^l = c$. Therefore,

$$P(X) - cP(Y) = \sum_{i \in I} a_i (X^i - cY^i) = \sum_{i \in I} a_i (X^i - b^i Y^i).$$

Clearly, X - bY is a linear factor of $F_c(X, Y, Z)$.

If $b^l = 1$, i.e. c = 1, then the same procedure shows that X - bY is a linear factor of P(X) - P(Y) which equals (X - Y)F(X, Y, 1). Since $b \neq 1$, X - bY is a linear factor of F(X, Y, 1).

Proof of Theorem 2 and Theorem 3. The sufficient conditions has been proved in Proposition 3.3 and Proposition 4.7. For the converse part, we only need to consider when l=1; l=2 and $\min\{m_1,m_2\}=1$; l=2 and $m_1=m_2=2$; l=3 and $m_1=m_2=m_3=1$. If l=1, then $P(X)=(X-\alpha_1)^n+a$, $n\geq 2$, $a\in \mathbb{C}$. Therefore, P(X) is not a uniqueness polynomial for rational functions since $P(f+\alpha_1)=P(\xi f+\alpha_1)$ for any rational function f and any n-th roots of unity ξ . Therefore, P is not a uniqueness or strong uniqueness polynomials for rational functions or meromorphic functions.

We recall from [8] that if P satisfies the separation condition, then $\{(\alpha_i, \alpha_i, 1) | 1 \le i \le l, m_i \ge 2\}$ are the only multiple points of C and each $(\alpha_i, \alpha_i, 1)$ is ordinary and has multiplicity m_i ; C_c has at most l multiple points $\{(\alpha_i, \alpha_{\tau(i)}, 1) | 1 \le i \le l, P(\alpha_i) = cP(\alpha_{\tau(i)})\}$ and each $(\alpha_i, \alpha_{\tau(i)}, 1)$ has multiplicity $\min\{m_i, m_{\tau(i)}\} + 1$ and is ordinary if $m_i = m_{\tau(i)}$.

If l=2 and $\min\{m_1, m_2\} = 1$, we may assume that $m_2 = 1$. If $m_1 = 1$, then the curve C is smooth, and hence is irreducible. If $m_1 \geq 2$, then F(X,Y,Z) = 0 has only one singular point $\mathfrak{q}_1 = (\alpha_1, \alpha_1, 1)$ which has multiplicity m_1 . We may assume that F(X,Y,Z) has no linear factor, otherwise P is not a uniqueness polynomial for rational functions or meromorphic functions. If F(X,Y,Z) has a proper irreducible homogeneous factor $H \in \mathbf{C}[X,Y,Z]$, then F = HG for $G \in \mathbf{C}[X,Y,Z]$ and $\deg H \geq 2$ by assumption. Let $m_1^G(\leq \deg G)$ and m_1^H be the multiplicity of \mathfrak{q}_1 in G = 0 and H = 0 respectively. We note that since H is irreducible and $\deg H \geq 2$, $m_1^H < \deg H$. We have $m_1^G + m_1^H = m_1$ and $\deg G + \deg H = \deg F = n - 1 = m_1 + 1$. On the other hand, by Bézout's theorem, we have $m_1^H m_1^G = \deg H \deg G$. Then $m_1^H = \deg H$ and $m_1^G = \deg G$ which leads to a contradiction. Therefore, F is

irreducible and has genus zero by the genus formula. This shows that P is not a uniqueness polynomials or strong uniqueness for rational functions or meromorphic functions.

If l=2 and $m_1=m_2=2$, then n=5 and C has two multiple points $\mathfrak{q}_1=(\alpha_1,\alpha_1,1)$ and $\mathfrak{q}_2=(\alpha_2,\alpha_2,1)$ which are ordinary and has multiplicity 2. One can check similarly via Bézout's theorem that C is irreducible. By the genus formula we see that the genus of C is one. Hence P is not a uniqueness polynomial or strong uniqueness polynomial for meromorphic functions.

If l = 3 and $m_1 = m_2 = m_3 = 1$, then n = 4 and C is a smooth curve (thus irreducible) of genus one. Hence P is not a uniqueness polynomial or strong uniqueness polynomial for meromorphic functions.

Finally, if l = 3, $m_1 = m_2 = m_3 = 1$, and

$$\frac{P(\alpha_1)}{P(\alpha_2)} = \frac{P(\alpha_2)}{P(\alpha_3)} = \frac{P(\alpha_3)}{P(\alpha_1)} = w,$$

for w satisfying $w^2 + w + 1 = 0$. Then F_w has 3 multiple points which are ordinary and each has multiplicity 2. One can check similarly that C_w is irreducible and has genus zero. Therefore P(X) is not a strong uniqueness polynomial for rational functions. This completes the proof.

Proof of Corollary 1. After a linear transformation, we may assume that $P(X) = X^n + aX^m + b$. If $\gcd(m,n) = d > 1$ or a = 0 then $P(X) = P(\xi_d X)$ where ξ_d is a d-primitive roots of unity. Therefore, P(X) is not a uniqueness polynomial for rational functions or meromorphic functions in this case. If n - m = 1, then P'(X) = 0 has two distinct roots and the non-zero root has multiplicity one in P(X). Then by Theorem 2, P(X) is not a uniqueness polynomial for rational functions or meromorphic functions in this case. If b = 0 and $n - m \ge 2$, then $P(\xi_{n-m}X) = \xi_{n-m}^m P(X)$ where ξ_{n-m} is a (n-m)-primitive roots of unity. This implies that $Y - \xi_{n-m}X$ is a linear factor of $F_{\xi_{n-m}}^{-m}(X,Y,Z)$. Therefore, P(X) is not a strong uniqueness polynomial for rational functions or meromorphic functions if b = 0 and $n - m \ge 2$.

From now, we assume that $a \neq 0$, gcd(m, n) = 1, and $n - m \geq 2$. We first claim that P(X) satisfies the separation condition. Since

$$P'(X) = nX^{m-1}(X^{n-m} + \frac{ma}{n}) = nX^{m-1} \prod_{i=0}^{n-m-1} (X - \xi_{n-m}^{i}\alpha)$$

where α satisfies

$$\alpha^{n-m} = \frac{-ma}{n}.$$

Then P(0) = b and

$$\begin{split} P(\xi_{n-m}^i\alpha) &= (\xi_{n-m}^i\alpha)^m(\alpha^{n-m} + a) + b \\ &= \frac{(n-m)a\alpha^m}{n}\xi_{n-m}^{im} + b. \end{split}$$

Clearly, $P(0) \neq P(\xi_{n-m}^i \alpha)$. Since $\gcd(n,m) = 1$, m is also relatively prime to n-m. Therefore, ξ_{n-m}^m is also a (n-m) primitive root of unity and hence $\xi_{n-m}^{mi} \neq \xi_{n-m}^{mj}$ if $0 \leq i \neq j \leq n-m-1$. Therefore, $P(\xi_{n-m}^i \alpha) \neq P(\xi_{n-m}^j \alpha)$ if $0 \leq i \neq j \leq m-n-1$. This concludes that P(X) satisfies the separation condition. Secondly, when $b \neq 0$ we claim the zero set of P(X) is affine rigid, equivalently, F(X,Y,Z) and each $F_c(X,Y,Z)$, $c \neq 0,1$ have no linear factor. It is clear that F(X,Y,Z) is irreducible if $\gcd(n,m)=1$. Suppose that $\nu X - \lambda Y - \mu Z$ is a factor of $F_c(X,Y,Z)$. It's clear that $F_c(X,Y,Z)$ has no factor of the type $\lambda Y + \mu Z$ or $\nu X - \mu Z$. Therefore, we assume that $\nu = 1$ and $\lambda \neq 0$. Then

$$0 \equiv P(\lambda Y + \mu) - cP(Y)$$

= $(\lambda Y + \mu Z)^n - cY^n + a[(\lambda Y + \mu Z)^m - cY^m] + b(1 - c).$

If $\mu=0$, then it implies b(1-c)=0 which is impossible if $b\neq 0$. If $\mu\neq 0$, since $n-m\geq 2$, comparing the terms of degree n-1 in the equation implies that $\lambda=0$ which is impossible. Therefore, S is affinely rigid under the assumption. Thirdly, since P'(X) has n-1 distinct zeros if m=1; or n-m+1 zeros, one with multiplicity m-1 and the others has multiplicity one if $m\geq 1$. By Theorem 2 it is a uniqueness polynomial for rational functions if and only if $n\geq 4$; and by Theorem 3 is a (strong) uniqueness polynomial for meromorphic functions if and only if $n\geq 5$. This completes the proof of (i), (iii) and (iv). For the same reason, P(X) is a strong uniqueness polynomial for rational functions if $n\geq 5$. It is now remains to check the case when n=4 in more details. Since $n-m\geq 2$ and $\gcd(n,m)=1$, we only need to consider when n=4 and m=1. Then $P(X)=X^4+aX+b$ and

$$P'(X) = 4X^3 + aX = 4(X - \alpha)(X - w\alpha)(X - w^2\alpha)$$

with $\alpha^3 = -a/4$ and $w^2 + w + 1 = 0$. From Theorem 2, P(X) is not a strong uniqueness polynomials for rational functions in this case if and only if there is a permutation τ of $\{1,2,3\}$ with $\tau(i) \neq i$ such that

(5.1)
$$\frac{P(\alpha_1)}{P(\alpha_{\tau(1)})} = \frac{P(\alpha_2)}{P(\alpha_{\tau(2)})} = \frac{P(\alpha_3)}{P(\alpha_{\tau(3)})} = w$$

where α_1 , α_2 , α_3 are solutions of P'(X). For the first one, we have

$$P(\alpha) = \frac{3}{4}a\alpha + b, \quad P(w\alpha) = \frac{3}{4}aw\alpha + b, \quad P(w^2\alpha) = \frac{3}{4}aw^2\alpha + b.$$

Since $b \neq 0$, it is easy to see that $P(\alpha) \neq wP(w\alpha)$ and $P(\alpha) = wP(w^2\alpha)$. Therefore it is impossible for P(X) to satisfy (5.1). Hence, P(X) is a strong uniqueness polynomial for rational functions in this case. This concludes the proof for (ii). \square

References

- T. T. H. An, J. T.-Y. Wang, Uniqueness polynomials for complex meromorphic functions, Inter. J. Math. 13 No 10 (2002), 1095-1115
- [2] T. T. H. An, J. T.-Y. Wang, And P.-M. Wong, Unique range sets and uniqueness polynomials in positive characteristic, Acta Arith. to appear.
- [3] W. CHERRY, AND J. T.-Y. WANG, Uniqueness polynomials for entire functions, Inter. J. Math 13 No 3 (2002), 323-332.
- [4] H. Fujimoto, On uniqueness of meromorphic functions sharing finite sets, Amer. J. Math. 122, (2000), 1175–1203.
- [5] H. Fujimoto, On uniqueness polynomials for meromorphic functions, Nagoya Math. J., to appear
- [6] H. H. KHOAI, T. T. H. AN, On uniqueness polynomials and bi-URS for p-adic meromorphic function, J. Number theory 87, (2001), 211–221.
- [7] B. SHIFFMAN, Uniqueness of entire and meromorphic functions sharing finite sets, Complex Variables Theory Appl. 43, (2001), no. 3-4, 433-449.
- [8] J. T.-Y. Wang, Uniqueness polynomials and bi-unique range sets for rational functions and non-Archimedean meromorphic functions, Acta Arith. 104, (2002), 183–200.

Institute of Mathematics, Academia Sinica, Nankang, Taipei 11529, Taiwan, R.O.C. $E\text{-}mail\ address$: tthan@math.sinica.edu.tw

Institute of Mathematics, Academia Sinica, Nankang, Taipei 11529, Taiwan, R.O.C. $E\text{-}mail\ address:}$ jwang@math.sinica.edu.tw

Department of Mathematics, University of Notre Dame, Notre Dame, IN 46556, U.S.A.

E-mail address: wong.2@nd.edu