

# Viscosity solution of a Delta Greek nonlinear Black-Scholes equation

*Rui M.P. Almeida\**, *Teófilo D. Chihaluca†* and *José C.M. Duque‡*

University of Beira Interior  
Center of Mathematics and Applications  
Covilhã, Portugal

May 5, 2020

## Abstract

In this paper, a class of nonlinear option pricing models involving transaction costs is considered. The diffusion coefficient of the nonlinear parabolic equation for the price  $V$  is assumed to be a linear function of the option's underlying asset price and the Gamma Greek  $V_{xx}$ . The main aim of this work is to study the governing PDE of the Delta Greek. The existence of viscosity solutions is proved using the vanishing viscosity method. Regularizing the equation by adding a small perturbation to the initial problem, a sequence of approximate solutions  $u^\varepsilon$  is constructed and then the method of weak limits is applied to prove the convergence of the sequence to the viscosity solution of the Delta equation. The approximate problems constructed are shown to have good regularity, which allows the use of efficient and robust numerical methods.

**Keywords** Delta equation; transaction costs; convergence; vanishing viscosity method.

## 1 Introduction

In financial mathematics, the Black-Scholes model is frequently used for pricing derivatives by means of the reversed-time parabolic partial differential equation (1) ([10, 19, 22]).

$$0 = V_\tau + \frac{1}{2}\sigma^2 S^2 V_{SS} + (r - q)SV_S - rV, \quad S > 0, \quad \tau \in ]0, T[. \quad (1)$$

---

\*ralmeida@ubi.pt

†teofilo.chihaluca@ubi.pt

‡jduque@ubi.pt

In (1)  $V$  is the option value,  $S$  the underlying asset price,  $\tau$  the time,  $T$  the expiry date,  $\sigma$  the volatility,  $r$  the riskless interest rate and  $q$  the dividend rate. Equation (1) has an analytical solution (see [5]) and to obtain approximate solutions there are also several reliable numerical methods, such as the binomial method, the Monte Carlo method, finite difference and finite element methods. For further details we refer the reader to the survey book [14].

The Black-Scholes equation (1) is very effective in an idealistic market, for example, without transaction costs. However transaction costs may arise, for example, when trading securities. Although they are small in general, they can lead to an increase in the option price, in which case the Black-Scholes pricing methodology is no longer valid since perfect hedging is impossible. Consequently, different models have been proposed to modify equation (1) in order to accommodate transaction costs, such as those in [16, 2, 4, 13]. In these models, the constant volatility is replaced by a modified volatility function which can depend on time, on the asset price, on the option value and its derivatives. The resulting model is a nonlinear equation in nondivergence form.

For the general nonlinear Black-Scholes equation, an explicit solution is unknown and the numerical techniques available are far less than for the linear model. Barles and Soner proved in [4] the existence of viscosity solutions for their model and made some simulations using finite differences but no convergence analysis was performed. However, it is known that explicit schemes have the disadvantage that restrictive conditions on the discretization parameters (for instance, the ratio of the time and space step) are needed to obtain stable, convergent schemes. Moreover, the convergence order is only one in time and two in space. Pooley et al. [18] numerically studied the convergence of some finite difference schemes applied to a nonlinear Black-Scholes equation and presented some examples where non-monotone discretization schemes (such as standard Crank-Nicolson time stepping) can converge to incorrect solutions, or lead to instability. In [9], the authors combine high-order compact difference scheme techniques to construct numerical solutions of the transformed non linear equation using the transformation  $x = \ln(S/K)$  with frozen values of the nonlinear volatility term to make the formulation linear. This transformation transforms the spatial domain  $[0, \infty]$  to  $[-\infty, \infty]$  and, in computations, this infinite domain has to be truncated, which essentially omits the degeneracy of the equation at  $S = 0$ . They show that the finite difference solution converges locally uniformly to the unique viscosity solution of the equation. Ankudinova and Ehrhardt [1] used a Crank-Nicolson method combined with a high order compact difference scheme to construct a numerical scheme for the linearized Black-Scholes equation. Company, Jodar and Pintos in [7] proposed a semi-discretization technique which approximates the nonlinear equation with a system of ordinary differential equations and solved the system using the backward Euler scheme. In order to make the high order scheme work, a smoothing technique for the payoff condition is used, which essentially changes the nature of the pricing problem. In [17], a method based on an upwind finite difference scheme for the spatial discretization and on a fully implicit time-stepping scheme is developed. The authors prove that the approximate solution converges unconditionally to the

viscosity solution of the equation.

The partial derivatives of the solution, the Greeks, are considered of major importance in finance, see for example [19]. In particular, the first spatial derivative, referred to as the Delta Greek in finance, is the key for the hedging process in time, that is portfolio projection against market movements as it follows from the Black-Scholes hedging arguments. There are a few papers concerning the calculation of the Delta Greek directly. Radoslav Valkov in [20] presented a convergence analysis of a positivity-preserving fitted finite volume element method (FVEM) for a generalized Black-Scholes equation transformed on finite interval, degenerating on both boundary points. He first formulated the FVEM as a Petrov-Galerkin finite element method using a spatial discretization, previously proposed by the author in [21]. The Garding coercivity of the corresponding discrete bilinear form was established. He obtained stability and error bounds for the solution of the fully-discrete system. Analysis of the impact of the finite domain transformation on the numerical solution of the original problem was given. Recently, Koleva and Vulkov ([12]) constructed and analysed monotone and sign-preserving finite difference schemes for the Delta equation. They proposed some Newton and Picard iterative procedures for solving the non-linear systems of algebraic equations.

The theory of continuous viscosity solutions for fully nonlinear second-order elliptic and parabolic equations has been introduced by Crandall-Lions for the Hamilton-Jacobi equations and is well established nowadays. For more details on this theory we refer the reader to [8] and references therein. However, the Delta equations has discontinuous initial condition, so isn't to expect continuous solution and the conventional theories of viscosity solutions do not apply. There are several definitions of discontinuous viscosity solutions, but most of them are rather ad hoc. In [6] we can see a survey of the latest development on the uniqueness and regularity of the discontinuous solutions of the Hamilton-Jacobi equation, with discontinuous initial data that are continuous outside a set of measure zero. They proved that the discontinuous solutions of the problem is unique when the initial condition is everywhere continuous. They also clarified the connections among the discontinuous solutions from different notions.

Usually, a high order method requires that the solution to the PDE is sufficiently smooth in order to achieve the expected order of convergence. However, it is known that the non-linear Black-Scholes equation generally does not have classic smooth solutions, but only viscosity solutions. Therefore, a numerical solution to the nonlinear Black-Scholes equation by a high order numerical scheme is not necessarily more accurate than that from a first-order discretization scheme, mainly due to the non-smoothness of the given data and the exact solution. Due to the Greeks being relevant for the quantitative analysis, reliable numerical methods are required for the pricing of options which not only provide a good approximation for the price, but also for its derivatives. In this work we study a simplification of the nonlinear Black-Scholes equation proposed by Barles and Soner ([4]) given focus to the corresponding Delta equation. We prove the existence and uniqueness of possibly discontinuous viscosity solutions for the Delta equation. We first regularize the equation by adding a small per-

turbation parameter and then apply the method of weak limits to prove the convergence of the classical solutions of the regularized problem to the viscosity discontinuous solution of the Delta equation. The main goal of this work is to create a basis to establish the convergence of high order robust numerical methods since the regularized problem has good smooth solutions.

The remainder of this paper is organized as follows. In Section 2, the problem is described. In Section 3, we define the regularized approximate problem as a general nonlinear equation. Section 4 is dedicated to obtain some à priori estimatives. In Section 5 we prove the convergence and, finally, in Section 5, we draw some comments.

## 2 The nonlinear model

In 1998 Barles and Soner [4] developed a complex model that modifies Equation (1) and accommodates transaction costs. Following the Hedges and Neuberger utility function approach in [11], they proposed the volatility function

$$\tilde{\sigma} = \sigma^2 \left( 1 + \Psi(e^{r(T-\tau)} a^2 S^2 V_{SS}) \right), \quad (2)$$

where  $\sigma$  is the historical volatility,  $a = \frac{k}{\sqrt{\epsilon}}$  and the function  $\Psi(A)$  is the solution to the following nonlinear differential equation (ODE)

$$\Psi'(A) = \frac{\Psi(A) + 1}{2\sqrt{A\Psi(A)} - A}, \quad A \neq 0, \quad (3)$$

with the initial condition

$$\Psi(0) = 0. \quad (4)$$

In this way Equation (1) becomes the following non-linear Black-Scholes equation with Barles and Soner's model,

$$0 = V_\tau + \frac{1}{2} \sigma^2 \left( 1 + \Psi(e^{r(T-\tau)} a^2 S^2 V_{SS}) \right) S^2 V_{SS} + (r-q)SV_S - rV, \quad S > 0, \tau \in ]0, T[, \quad (5)$$

which we will consider for European options, where  $S$  pays out continuous dividend  $qSdt$  with time step  $dt$ .

A European call option allows the buyer to buy an asset of value  $S$  for a value  $K$  on maturity date  $T$ , while an European put option allows the holder to sell an asset of value  $S$  for a value  $K$  on maturity date  $T$ . For the sake of simplicity, we will only consider the call option. Since the option can only be exercised on maturity, we complement Equation (5) with the following conditions, in order

to avoid arbitrariness:

$$V(S, T) = \max\{S - K, 0\}, \quad \text{when } S \geq 0; \quad (6)$$

$$\lim_{S \rightarrow \infty} \frac{V(S, \tau)}{S - Ke^{-r(T-\tau)}} = 1, \quad \text{for } \tau \in [0, T]; \quad (7)$$

$$V(0, \tau) = 0, \quad \text{for } \tau \in [0, T]; \quad (8)$$

$$\lim_{S \rightarrow \infty} V_S(S, \tau) = 1, \quad \text{for } \tau \in [0, T]. \quad (9)$$

Barles and Soner proved the existence of a viscosity solution for the European option with volatility given by (2). Their numerical results indicate an economically significant price difference between the standard Black-Scholes model and the non-linear model with transaction costs.

Although an explicit solution for Problem (3)-(4), is unknown, the following Lemmas provide some useful information about its behaviour.

**Lemma 1** ([4]). *A straightforward analysis of the ordinary differential equation (3) reveals that*

$$\lim_{A \rightarrow \infty} \frac{\Psi(A)}{A} = 1 \quad \text{and} \quad \lim_{A \rightarrow -\infty} \Psi(A) = -1. \quad (10)$$

**Lemma 2** ([4]). *If*

$$A \geq 0 \quad \text{then} \quad \Psi(A) \geq 0$$

*and if*

$$A \leq 0 \quad \text{then} \quad -1 < \Psi(A) \leq 0.$$

**Lemma 3** ([4]). *The differential  $\Psi'(A) \geq 0$ .*

**Lemma 4** ([7]). *If  $A \geq 0$  then  $\Psi(A) \leq C_1A + C_2$  with  $C_1 \approx 1.1$  and  $C_2 \approx 2.62$ .*




Figure 1: Numerical simulation of  $\Psi$ .

A numerical solution for Problem (3)-(4) is represented in Figure 1. By the lemmas above and figure 1 is reasonable to simplify the model considering  $\Psi(A) = A$  for  $A > 0$ . So, in this work, we study the Black-Scholes Equation (5),

in which the volatility is assumed to be a function of the underlying asset  $S$ , and time  $\tau$  and the Gamma of option (the Greek Gamma is a second derivative  $V_{SS}$ ) that is,

$$0 = V_\tau + \frac{1}{2}\sigma^2 \left(1 + e^{r(T-\tau)}a^2S^2V_{SS}\right) S^2V_{SS} + (r-q)SV_S - rV, \quad S > 0, \tau \in ]0, T[. \quad (11)$$

It is known that  $V_{SS} \geq 0$  for European Puts and Calls in the absence of transaction costs. In case  $V_{SS} < 0$ , Problem (11) is ill-posed and without solution for general pay-off functions (see [16, 2] for similar cases). Therefore we will henceforth assume that  $V_{SS} > 0$ .

## 2.1 Delta equation

Differentiating (11) with respect to  $S$ , the resulting equation is

$$V_{S\tau} + \left(\frac{1}{2}\sigma^2 \left(1 + e^{r(T-\tau)}a^2S^2V_{SS}\right) S^2V_{SS}\right)_S + (r-q)SV_{SS} - qV_S = 0. \quad (12)$$

Differentiating (6), we have

$$V_S(S, T) = \begin{cases} 0, & S < K \\ 1, & S > K \end{cases}. \quad (13)$$

Considering  $x = S$ ,  $t = T - \tau$  and  $u(x, t) = V_S(S, T - \tau)$ , we obtain the parabolic partial differential equation in divergence form,

$$u_t = \left(\frac{1}{2}\sigma^2(1 + e^{rt}a^2x^2u_x)x^2u_x\right)_x + (r-q)xu_x - qu, \quad x > 0, \quad t \in ]0, T]. \quad (14)$$

which we will call Delta equation. We will consider the corresponding initial condition

$$u(x, 0) = u_0(x) = \begin{cases} 0, & 0 < x < K \\ 0.5, & x = K \\ 1, & x > K \end{cases} \quad (15)$$

By (9), the boundary conditions are

$$u(0, t) = g_1(t) = 0, \quad t \in [0, T] \quad (16)$$

and

$$\lim_{x \rightarrow \infty} u(x, t) = 1, \quad t \in [0, T]. \quad (17)$$

Note that condition  $u(0, t) = 0$  is necessary in order to guarantee compatibility between the equation and the initial data. It is not possible to define  $u_0$  in  $x = K$  by the Black-Scholes model, since  $V(S, T)$  is not differentiable at  $S = K$ . However, as we shall see below, this value is not essential, so we will consider the value  $u_0(K) = 0.5$  for practical purposes.

**Proposition 5.** *If function  $V(S, \tau)$  is a solution of Problem (5)-(9), then  $u(x, t) = V_S(S, T - \tau)$  is a solution of problem (14)-(17). Conversely if  $u(x, t)$  is a solution of Problem (14)-(17), then*

$$V(S, \tau) = \int_0^S u(x, T - \tau) dx \quad (18)$$

*is a solution of problem (5)-(9).*

*Proof.* The first statement was proved above. Suppose now that  $V(x, \tau)$  is given by (18). Then making the change of variable  $t = T - \tau$ , it follows that

$$\begin{aligned} V_\tau(S, \tau) &= \int_0^S u_\tau(x, T - \tau) dx = \int_0^S -u_t(x, t) dx = \\ &= - \int_0^S \left( \frac{1}{2} \sigma^2 (1 + e^{rt} a^2 x^2 u_x) x^2 u_x \right)_x + (r - q) x u_x - q u dx. \end{aligned}$$

Using the integration by parts, we obtain

$$V_\tau(S, \tau) = -\frac{1}{2} \sigma^2 (1 + e^{rt} a^2 S^2 u_S(S, t)) S^2 u_S(S, t) - (r - q) S u(S, t) + q \int_0^S u(x, t) dx$$

Differentiating (18) we have  $V_S(S, \tau) = u(S, T - \tau)$  and  $V_{SS}(S, \tau) = u_S(S, T - \tau)$ . Substituting in the last equation, we obtain (5). In order to obtain (6), we just need to calculate de integral

$$V(S, T) = \int_0^S u(x, 0) dx = \begin{cases} 0 & , S < K \\ 0 & , S = K \\ S - K & , S > K \end{cases} .$$

As we mentioned earlier any real value of  $u(K, 0)$  verifies (6). Condition (7) is obtained applying the L'Hôpital's rule, that is,

$$\lim_{S \rightarrow \infty} \frac{V(S, \tau)}{S - K e^{-r(T-\tau)}} = \lim_{S \rightarrow \infty} \frac{\int_0^S u(x, T - \tau) dx}{S - K e^{-r(T-\tau)}} = \lim_{S \rightarrow \infty} \frac{u(S, T - \tau)}{1} = 1$$

by (17). Condition (8) is obvious. Finally, condition (9) is the same as condition (17).  $\square$

Since it is difficult to deal with an infinite spatial domain, we will consider the spatial domain  $]0, b[$  with  $b$  sufficiently far from  $K$  and substitute equation (17) by

$$u(b, t) = 1, \quad t \in [0, T]. \quad (19)$$

There are two main problems concerning the Delta function that we need to deal with. First, the initial condition is not smooth, it has a discontinuity at  $x = K$  and secondly the diffusion term becomes zero at  $x = 0$ . So we expect that the Delta equation has only a viscosity solution. For ease of reading, we recall the notion of viscosity solution in the next subsection.

## 2.2 Viscosity solution

Consider the following general problem

$$\begin{cases} u_t + F(x, t, u, u_x, u_{xx}) = 0, & (x, t) \in Q_T = \Omega \times ]0, T], \\ u(x, t) = g(x, t), & (x, t) \in \partial\Omega \times [0, T[ \\ u(x, 0) = u_0(x), & x \in \bar{\Omega} \end{cases} \quad (20)$$

where  $F : \mathbb{R}^5 \rightarrow \mathbb{R}$  is a continuous function and  $g, u_0$  are given functions. To extend the technique of viscosity solutions to functions that need not to be continuous, we define

$$u^*(x, t) = \limsup_{r \rightarrow 0} \{u(y, \tau) : |(y, \tau) - (x, t)| \leq r, (y, \tau) \in Q_T\} \quad (21)$$

and

$$u_*(x, t) = \liminf_{r \rightarrow 0} \{u(y, \tau) : |(y, \tau) - (x, t)| \leq r, (y, \tau) \in Q_T\} \quad (22)$$

which are called the upper and lower semicontinuous envelopes of  $u$  respectively.

**Definition 6.** Let  $u : \bar{Q}_T \rightarrow \mathbb{R}$  be a bounded function and  $\phi : \bar{Q}_T \rightarrow \mathbb{R}$  be any  $C^{2,1}(\bar{Q}_T)$  function.

- The function  $u$  is called a viscosity subsolution of (20) if, whenever  $(x_0, t_0)$  is a maximum of  $u^* - \phi$ , the following conditions are satisfied:

$$\phi_t(x_0, t_0) + F(x_0, t_0, u^*(x_0, t_0), \phi_x(x_0, t_0), \phi_{xx}(x_0, t_0)) \leq 0, \quad (23)$$

when  $(x_0, t_0) \in Q_T$ ,

$u^*(x_0, t_0) - g(x_0, t_0) \leq 0$  or (23), when  $(x_0, t_0) \in \partial\Omega \times [0, T[$ ,

and

$u^*(x_0, t_0) - u_0^*(x_0) \leq 0$  or (23), when  $x_0 \in \bar{\Omega}, t_0 = 0$ .

- The function  $u$  is called a viscosity supersolution of (20) if, whenever  $(x_0, t_0)$  is a minimum of  $u_* - \phi$ , the following conditions are satisfied:

$$\phi_t(x_0, t_0) + F(x_0, t_0, u_*(x_0, t_0), \phi_x(x_0, t_0), \phi_{xx}(x_0, t_0)) \geq 0, \quad (24)$$

when  $(x_0, t_0) \in Q_T$ ,

$u_*(x_0, t_0) - g(x_0, t_0) \geq 0$  or (24), when  $(x_0, t_0) \in \partial\Omega \times [0, T[$ ,

and

$u_*(x_0, t_0) - u_0^*(x_0) \geq 0$  or (24), when  $x_0 \in \bar{\Omega}, t_0 = 0$ .

- A bounded function  $u$  which is both a viscosity subsolution and a supersolution is called a (non necessarily continuous) viscosity solution.

**Remark 7.** The definition of viscosity solution is an absolute one. This means that if  $u$  is a viscosity subsolution in  $\Omega$ , then it is also a subsolution in  $\Omega'$ , where  $\Omega' \subset \Omega$ .



**Remark 8.** *In the definition of viscosity solution, local maximum can be replaced by global maximum and also by strict local or global maximum. Moreover  $C^{2,1}$  functions can be substituted by smooth functions. Also we can assume that the local maximum is zero. Similar remarks apply to supersolutions.*

To prove the existence of viscosity solutions to Problem (14)-(16) and (19) we will use the vanishing viscosity method to construct a sequence of approximate solutions  $u^\varepsilon$ , we will then apply the method of weak limits to prove the convergence of the sequence to the viscosity solution of the Delta equation. The advantage of the weak limit method is that it allows passing to the limits with only uniform  $L^\infty$  estimates on  $u^\varepsilon$ .

Problem (14)-(16) and (19) can be written as (20) with  $\Omega = ]0, b[$ ,

$$\begin{aligned} F(x, t, u, u_x, u_{xx}) = & - (0.5x^2\sigma^2 (1 + 2e^{rt}a^2x^2u_x)) u_{xx} \\ & - 2\sigma^2e^{rt}a^2x^3u_x^2 - (r - q + \sigma^2)xu_x + qu \end{aligned} \quad (25)$$

with  $g(0, t) = 1$ ,  $g(b, t) = 1$  and  $u_0$  defined by (15).

### 3 Regularization

In order to overcome the difficulties, in dealing with the Delta Equation, raised above, we propose a small perturbation of the initial problem by considering a new approximate problem. We seek a function  $u^\varepsilon(x, t)$  that satisfies the equation

$$u_t^\varepsilon = ((a_0^\varepsilon(t, x, u_x^\varepsilon)u_x^\varepsilon)_x + (r - q)xu_x^\varepsilon - qu^\varepsilon, \quad (x, t) \in Q_T. \quad (26)$$

with  $a_0^\varepsilon = 0.5x^2\sigma^2(1 + e^{rt}a^2x^2u_x^\varepsilon) + \varepsilon$ ,  $\varepsilon > 0$  small, boundary conditions (16), (19) and initial condition

$$u^\varepsilon(x, 0) = \begin{cases} u_0(x), & 0 \leq x < K - \varepsilon \\ H_5(x), & K - \varepsilon \leq x \leq K + \varepsilon \\ u_0(x), & K + \varepsilon < x \leq b \end{cases} \quad (27)$$

where  $H_5(x)$  is the Hermite polynomial of degree 5 that satisfies  $H_5(K - \varepsilon) = u_0(K - \varepsilon)$ ,  $H_5(K + \varepsilon) = u_0(K + \varepsilon)$ ,  $H_5'(K - \varepsilon) = H_5'(K + \varepsilon) = H_5''(K - \varepsilon) = H_5''(K + \varepsilon) = 0$ .

In order to define a weak solution, we multiply (26) by  $\psi$  and integrate with respect to  $x$ :

$$\int_0^b u_t^\varepsilon \psi \, dx - \int_0^b (a_0^\varepsilon u_x^\varepsilon)_x \psi \, dx - \int_0^b (r - q) x u_x^\varepsilon \psi \, dx + \int_0^b q u^\varepsilon \psi \, dx = 0.$$

Then, integrating by parts and assuming that  $\psi(0) = \psi(b) = 0$ , we obtain

$$\int_0^b u_t^\varepsilon \psi \, dx + \int_0^b a_0^\varepsilon u_x^\varepsilon \psi_x \, dx - \int_0^b (r - q) x u_x^\varepsilon \psi \, dx + \int_0^b q u^\varepsilon \psi \, dx = 0. \quad (28)$$

Note that for Equation (28) to make sense, we must have  $u^\varepsilon$ ,  $u_t^\varepsilon$  and  $u_x^\varepsilon \in L^2(0, b)$ , for  $t \in ]0, T]$ . Taking into consideration conditions (16), (19) we choose the test function space to be

$$V_0 = \{ \psi, \psi_x \in L^2(0, b) : \psi(0) = \psi(b) = 0 \},$$

and for the space solution we consider

$$V = \{ u, u_t, u_x \in L^2(0, b) : u(0, t) = 0, u(b, t) = 1, \text{ for all } t \in [0, T] \}.$$

**Definition 9** (Weak solution). *A function  $u^\varepsilon \in V$  is said to be a weak solution of problem (26)-(27) if it satisfies (27) and (28) for all  $\psi \in V_0$ , and  $t \in ]0, T[$ . Relation (28) must be understood as an equality in  $D'(0, T)$ .*

We are interested in obtaining classical solutions using the Leray-Schauder existence theory. So following [15], we need to obtain à priori estimates for all possible weak solutions of Equation (26).

## 4 À priori estimates

This section is entirely devote to obtaining the desired estimates. The main tools required are integral calculations, imbedding inequalities and Gronwall's Lemmas. Since we have the conditions

$$\psi(0) = \psi(b) = 0, \quad u^\varepsilon(0, t) = 0 \quad \text{and} \quad u^\varepsilon(b, t) = 1,$$

we are not able to considerer  $\psi = u^\varepsilon$  in (28), then we will introduce a new function  $v$  defined by

$$v(x, t) = u^\varepsilon(x, t) - \frac{x}{b}.$$

It has the properties

$$u^\varepsilon(x, t) = v(x, t) + \frac{x}{b},$$

$$u_t^\varepsilon(x, t) = v_t(x, t),$$

$$u_x^\varepsilon(x, t) = v_x(x, t) + \frac{1}{b} \quad \text{and} \quad u_{xx}^\varepsilon(x, t) = v_{xx}(x, t).$$

According to (26) the new function satisfies

$$v_t = \left( a_0^\varepsilon \left( v_x + \frac{1}{b} \right) \left( v_x + \frac{1}{b} \right) \right)_x + (r - q)x \left( v_x + \frac{1}{b} \right) - q \left( v + \frac{x}{b} \right), \quad (29)$$

that is,

$$v_t = ((a_1 + a_2 + \varepsilon)v_x)_x + (r - q)xv_x - qv + f, \quad (x, t) \in Q_T \quad (30)$$

with

$$a_1 = 0.5x^2\sigma^2 \left( 1 + e^{rt}a^2x^2 \left( v_x + \frac{1}{b} \right) \right),$$

$$a_2 = \frac{0.5\sigma^2 e^{rt} a^2 x^4}{b},$$

$$f = \frac{(\sigma^2 + r + 2q)x}{b} + \frac{2x^3 \sigma^2 e^{rt} a^2}{b^2}.$$

The boundary and initial conditions for this new problem are

$$v(0, t) = 0, \quad v(b, t) = 0, \quad (31)$$

$$v(x, 0) = v_0(x) = \begin{cases} -\frac{x}{b}, & 0 \leq x < k - \varepsilon \\ H_5(x) - \frac{x}{b}, & k - \varepsilon \leq x \leq k + \varepsilon \\ 1 - \frac{x}{b}, & k + \varepsilon < x \leq b. \end{cases} \quad (32)$$

The definition of weak solution to Problem (30)-(32) is similar to Definition 9.

**Remark 10.** Assumption  $u_x \geq 0$  is equivalent to  $v_x(x, t) + \frac{1}{b} \geq 0$ , and then  $a_1 \geq 0$ ,  $(x, t) \in Q_T$ .

First we prove that the initial condition is smooth.

**Lemma 11.** Let  $v_0(x)$  be as defined in (32). Then, for  $x \in [0, b]$ ,

$$|v_0(x)| \leq C, \quad (33)$$

$$|v_0'(x)| \leq C\varepsilon^{-1}, \quad \text{and} \quad (34)$$

$$|v_0''(x)| \leq C\varepsilon^{-2}, \quad (35)$$

where  $C$  does not depend on  $\varepsilon$ .

*Proof.* Regarding (32) we only need to prove the results for  $H_5(x)$  were  $H_5(x)$  is as defined in (27). Applying the divided difference method, we can write

$$H_5(x) = \frac{1}{8\varepsilon^3}(x - K + \varepsilon)^3 - \frac{3}{16\varepsilon^4}(x - K + \varepsilon)^3(x - K - \varepsilon) \\ + \frac{3}{16\varepsilon^5}(x - K + \varepsilon)^3(x - K - \varepsilon)^2 \quad x \in [K - \varepsilon, K + \varepsilon].$$

Using the triangular inequality, we have

$$|H_5(x)| \leq \frac{1}{8\varepsilon^3}|x - K + \varepsilon|^3 + \frac{3}{16\varepsilon^4}|x - K + \varepsilon|^3|x - K - \varepsilon| + \frac{3}{16\varepsilon^5}|x - K + \varepsilon|^3|x - K - \varepsilon|^2.$$

It is clear that  $|x - K + \varepsilon| \leq 2\varepsilon$  and  $|x - K - \varepsilon| \leq 2\varepsilon$  for  $x \in [K - \varepsilon, K + \varepsilon]$ , and so

$$|H_5(x)| \leq 10,$$

which prove (33). Differentiating  $H_5(x)$ , we obtain

$$H_5'(x) = \frac{3}{8\varepsilon^3}(x - K + \varepsilon)^2 - \frac{9}{16\varepsilon^4}(x - K + \varepsilon)^2(x - K - \varepsilon) - \frac{3}{16\varepsilon^4}(x - K + \varepsilon)^3 \\ + \frac{9}{16\varepsilon^5}(x - K + \varepsilon)^2(x - K - \varepsilon)^2 + \frac{6}{16\varepsilon^5}(x - K + \varepsilon)^3(x - K - \varepsilon)$$

Using the triangular inequality we have

$$|H'_5(x)| \leq \frac{45}{2}\varepsilon^{-1}$$

and so (34) is proved. The same technique permits us to prove (35) by estimating

$$|H''_5(x)| \leq \frac{81}{2}\varepsilon^{-2}.$$

□

**Theorem 12.** *Let  $v$  be a weak solution of (30)-(32) that satisfies Remark 10. Then*

$$\|v\|_{L^\infty(0,T;L^2(0,b))} \leq C \quad (36)$$

$$\text{and } \|v_x\|_{L^2(0,T;L^2(0,b))} \leq C\varepsilon^{-\frac{1}{2}} \quad (37)$$

where  $C = C(a, \sigma, r, q, b, T)$  does not depend on  $\varepsilon$ .

*Proof.* Considering  $\psi = v$  in the definition of a weak solution, we obtain

$$\int_0^b v_t v \, dx + \int_0^b (a_1 + a_2 + \varepsilon)v_x^2 \, dx = (r - q) \int_0^b x v_x v \, dx - q \int_0^b v^2 \, dx + \int_0^b f v \, dx.$$

Integrating by parts the first term on the right hand side we obtain

$$\frac{1}{2} \frac{d}{dt} \int_0^b v^2 \, dx + \int_0^b (a_1 + a_2 + \varepsilon)v_x^2 \, dx = -\frac{r+q}{2} \int_0^b v^2 \, dx - q \int_0^b v^2 \, dx + \int_0^b f v \, dx.$$

By Cauchy's inequality,

$$\frac{1}{2} \frac{d}{dt} \|v\|_{L^2(0,b)}^2 + \varepsilon \|v_x\|_{L^2(0,b)}^2 \leq \frac{1}{2} \|f\|_{L^2(0,b)}^2 + \frac{1}{2} \|v\|_{L^2(0,b)}^2.$$

Integrating with respect to  $t$ ,

$$\|v(x, t)\|_{L^2(0,b)}^2 + 2\varepsilon \int_0^t \|v_x\|_{L^2(0,b)}^2 \, dt \leq \int_0^T \|f\|_{L^2(0,b)}^2 \, dt + \int_0^T \|v\|_{L^2(0,b)}^2 \, dt + \|v(x, 0)\|_{L^2(0,b)}^2.$$

Taking into account that  $v_0 \in L^2(0, b)$  and  $\int_0^T \|f\|_{L^2(0,b)}^2 \, dt < C$  and applying Gronwall's inequality, we prove (36), and then (37) follows easily. □

The next theorem shows that  $v$  is uniformly bounded in  $L^\infty$ .

**Theorem 13.** *Let  $v$  be a weak solution of (30)-(32) that satisfies Remark 10. Then*

$$\|v\|_{L^\infty(0,T,L^\infty(0,b))} \leq C \quad (38)$$

where  $C = C(a, \sigma, r, q, b, T)$  does not depend on  $\varepsilon$ .

*Proof.* Multiplying (30) by  $v^{2k-1}$ ,  $k \geq 1$ , and integrating with respect to  $x$ , we obtain

$$\begin{aligned} \int_0^b v_t v^{2k-1} dx + (2k-1) \int_0^b (a_1 + a_2 + \varepsilon) v^{2k-2} v_x^2 dx &= (r-q) \int_0^b x v_x v^{2k-1} dx \\ &\quad - q \int_0^b v^{2k} dx + \int_0^b f v^{2k-1} dx. \end{aligned}$$

Applying integration by parts to the first term on the right-hand side, we have

$$\begin{aligned} \frac{1}{2k} \frac{d}{dt} \int_0^b v^{2k} dx + (2k-1) \int_0^b (a_1 + a_2 + \varepsilon) v^{2k-2} v_x^2 dx \\ = -\frac{r + (2k-1)q}{2k} \int_0^b v^{2k} dx + \int_0^b f v^{2k-1} dx. \end{aligned}$$

By Remark 10, we can ignore the second term on the left-hand side and then, applying the Hölder's inequality, we arrive at

$$\|v\|_{L^{2k}(0,b)}^{2k-1} \frac{d}{dt} \|v\|_{L^{2k}(0,b)} \leq \|f\|_{L^{2k}(0,b)} \|v\|_{L^{2k}(0,b)}^{2k-1}.$$

Cancelling the term  $\|v\|_{L^{2k}(0,b)}^{2k-1}$  we have

$$\frac{d}{dt} \|v\|_{L^{2k}(0,b)} \leq \|f\|_{L^{2k}(0,b)}.$$

Integrating with respect to  $t$  it results in

$$\|v\|_{L^{2k}(0,b)} \leq \|v_0\|_{L^{2k}(0,b)} + \int_0^T \|f\|_{L^{2k}(0,b)} dt.$$

Since  $v_0 \in L^\infty(0,b)$  and  $\int_0^T \|f\|_{L^\infty(0,b)} dt < C$ , taking  $k \rightarrow \infty$ , (38) follows.  $\square$

Equation (30) can be rewritten as

$$v_t = (a_1 + a_3 + \varepsilon) v_{xx} + a_4 v_x^2 + a_5 v_x - qv + f \quad (39)$$

with

$$a_3 = 0.5x^4 \sigma^2 e^{rt} a^2 \left(v_x + \frac{1}{b}\right) \geq 0 \text{ by Remark 10,}$$

$$a_4 = 2x^3 \sigma^2 e^{rt} a^2 \text{ and}$$

$$a_5 = \frac{4x^3 \sigma^2 e^{rt} a^2}{b} + (r - q + \sigma^2)x.$$

**Theorem 14.** *If  $v$  is a weak solution of (30)-(32) that satisfies Remark 10, then*

$$\|v_x\|_{L^\infty(0,T;L^2(0,b))} \leq C\varepsilon^{-\frac{3}{2}} \quad (40)$$

$$\text{and } \|v_{xx}\|_{L^2(0,T;L^2(0,b))} \leq C\varepsilon^{-2} \quad (41)$$

where  $C = C(a, \sigma, r, q, b, T)$  does not depend on  $\varepsilon$ .

*Proof.* Multiplying (39) by  $v_{xx}$ , and integrating with respect to  $x$ , we obtain

$$\begin{aligned} \int_0^b v_t v_{xx} dx &= \int_0^b (a_1 + a_3 + \varepsilon) v_{xx}^2 dx + \int_0^b a_4 v_x^2 v_{xx} dx \\ &\quad + \int_0^b a_5 v_x v_{xx} dx - q \int_0^b v v_{xx} dx + \int_0^b f v_{xx} dx \\ &\Leftrightarrow [v_x v_t]_0^b - \int_0^b v_x v_{xt} dx - \int_0^b (a_1 + a_3 + \varepsilon) v_{xx}^2 dx = \frac{1}{3} [v_x^3 a_4]_0^b \\ &\quad - \int_0^b \frac{v_x^3}{3} (a_4)_x dx + \frac{1}{2} [v_x^2 a_5]_0^b - \int_0^b \frac{v_x^2}{2} (a_5)_x dx - q \int_0^b v v_{xx} dx + \int_0^b f v_{xx} dx, \end{aligned}$$

where we used integration by parts. By Hölder's inequality we arrive at

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_0^b v_x^2 dx + \varepsilon \int_0^b v_{xx}^2 dx &= C v_x^3(b, t) + C \int_0^b v_x^3 dx + C v_x^2(b, t) \\ + C \int_0^b v_x^2 dx + \frac{\varepsilon}{3} \int_0^b v_{xx}^2 dx + C \varepsilon^{-1} \int_0^b v^2 dx &+ \frac{\varepsilon}{3} \int_0^b v_{xx}^2 dx + C \varepsilon^{-1} \int_0^b f^2 dx. \end{aligned}$$

According to (9) it is reasonable to consider  $v_x(b, t) = 0$ , for  $b$  sufficiently large. Now we use the Gagliardo-Nirenberg interpolation inequalities to eliminate the terms with  $v_x^3$  and thus we obtain

$$\frac{1}{2} \frac{d}{dt} \int_0^b v_x^2 dx + \frac{\varepsilon}{3} \int_0^b v_{xx}^2 dx \leq C \varepsilon^{-1} + \frac{\varepsilon}{4} \int_0^b v_{xx}^2 dx + C \varepsilon^{-3} \int_0^b v^6 dx.$$

Integrating with respect to  $t$  and recalling the previous theorems, the result follows.  $\square$

**Corollary 15.** *If  $u$  is a weak solution of (26) that satisfies  $u_x^\varepsilon > 0$ , then*

$$\int_0^T \|u_x^\varepsilon\|_{L^\infty(0,b)}^2 dt \leq C \varepsilon^{-4}$$

where  $C = C(a, \sigma, r, q, b, T)$  does not depend on  $\varepsilon$ .

*Proof.* By the interpolation inequality and Hölder's inequality, we have

$$\begin{aligned} \|u_x^\varepsilon\|_{L^\infty(0,b)} &\leq C \|u_{xx}^\varepsilon\|_{L^2(0,b)}^{\frac{1}{2}} \|u_x^\varepsilon\|_{L^2(0,b)}^{\frac{1}{2}} \\ \|u_x^\varepsilon\|_{L^\infty(0,b)}^2 &\leq C \|u_{xx}^\varepsilon\|_{L^2(0,b)}^2 + C \|u_x^\varepsilon\|_{L^2(0,b)}^2 \end{aligned}$$

Integrating with respect to  $t$  and using the previous Theorem the result follows.  $\square$

**Theorem 16.** *Let  $v$  be a weak solution of (30)-(32) that satisfies Remark 10. Then*

$$\|v_t\|_{L^2(0,T;L^2(0,b))} \leq C\varepsilon^{-1}$$

where  $C = C(a, \sigma, r, q, b, T)$  do not depends on  $\varepsilon$ .

*Proof.* Multiplying (30) by  $v_t$ , and integrating with respect to  $x$ , we obtain

$$\begin{aligned} \int_0^b v_t^2 dx + \int_0^b (a_1 + a_2 + \varepsilon) v_x v_{xt} dx &= (r - q) \int_0^b x v_x v_t dx - q \int_0^b v v_t dx \\ &\quad + \int_0^b f v_t dx \\ \Leftrightarrow \int_0^b v_t^2 dx + \frac{1}{2} \int_0^b (a_1 + a_2 + \varepsilon) (v_x^2)_t dx &\leq C \int_0^b v_x^2 dx + \frac{1}{4} \int_0^b v_t^2 dx \\ &\quad - \frac{q}{2} \frac{d}{dt} \int_0^b v^2 dx - C \int_0^b f^2 dx + \frac{1}{4} \int_0^b v_t^2 dx. \end{aligned}$$

Concerning the second term on the left hand side we can write

$$\begin{aligned} \frac{1}{2} \int_0^b (a_1 + a_2 + \varepsilon) (v_x^2)_t dx &= \frac{1}{2} \frac{d}{dt} \int_0^b (a_1 + a_2 + \varepsilon) v_x^2 dx \\ &\quad - \frac{1}{2} \int_0^b (a_1 + a_2 + \varepsilon)_t v_x^2 dx. \end{aligned}$$

Since

$$\begin{aligned} (a_2)_t &= \frac{0.5rx^4\sigma^2 e^{rt} a^2}{b} = ra_2, \\ (a_1)_t &= 0.5rx^4\sigma^2 e^{rt} a^2 v_x + 0.5x^4\sigma^2 e^{rt} a^2 v_{xt} + \frac{0.5rx^4\sigma^2 e^{rt} a^2}{b} \\ &= rba_2 v_x + ba_2 v_{xt} + ra_2, \end{aligned}$$

we have

$$\begin{aligned} \frac{1}{2} \int_0^b v_t^2 dx + \frac{1}{2} \frac{d}{dt} \int_0^b (a_1 + a_2 + \varepsilon) v_x^2 dx + \frac{q}{2} \frac{d}{dt} \int_0^b v^2 dx \leq \frac{1}{2} \int_0^b b r a_2 v_x^3 dx \\ + \int_0^b r a_2 v_x^2 dx + \frac{1}{2} \int_0^b b a_2 v_{xt} v_x^2 dx + C \int_0^b v_x^2 dx + C \int_0^b f^2 dx. \end{aligned}$$

Taking into account that

$$\frac{1}{2} \int_0^b b a_2 v_{xt} v_x^2 dx = \frac{1}{6} \int_0^b b a_2 (v_x^3)_t dx = \frac{1}{6} \frac{d}{dt} \int_0^b b a_2 v_x^3 dx - \frac{1}{6} \int_0^b b r a_2 v_x^3 dx,$$

we arrive at

$$\begin{aligned} \frac{1}{2} \int_0^b v_t^2 dx + \frac{1}{2} \frac{d}{dt} \int_0^b (a_1 + a_2 - \frac{1}{3} b a_2 v_x + \varepsilon) v_x^2 dx + \frac{q}{2} \frac{d}{dt} \int_0^b v^2 dx \leq \\ \frac{1}{3} \int_0^b b r a_2 v_x^3 dx + \int_0^b r a_2 v_x^2 dx + C \int_0^b v_x^2 dx + C \int_0^b f^2 dx. \end{aligned}$$

Define  $a_6 = 0.5x^2\sigma^2(1 + x^2a^2e^{rt}(\frac{2}{3}v_x + \frac{1}{b})) \geq 0$  by Remark 10. If we apply the Gagliardo-Nirenberg inequalities, the last equation becomes

$$\begin{aligned} \frac{1}{2} \int_0^b v_t^2 dx + \frac{1}{2} \frac{d}{dt} \int_0^b (a_6 + a_2 + \varepsilon) v_x^2 dx + \frac{q}{2} \frac{d}{dt} \int_0^b v^2 dx \leq C \int_0^b v_{xx}^2 dx \\ + C \int_0^b v^6 dx + C \int_0^b v_x^2 dx + C \int_0^b f^2 dx. \end{aligned}$$

Integrating with respect to  $t$  we have

$$\begin{aligned} \int_0^t \int_0^b v_t^2 dx dt + \int_0^b (a_2 + a_6 + \varepsilon) v_x^2 dx + q \int_0^b v^2 dx \leq C \int_0^T \int_0^b v_{xx}^2 dx dt \\ + C \int_0^T \int_0^b v^6 dx dt + C \int_0^T \int_0^b f^2 dx dt + C \int_0^T \int_0^b v^2 dx dt + q \int_0^b v_0^2 dx \\ + \int_0^b (a_2 + a_6 + \varepsilon) v_x^2(x, 0) dx. \end{aligned}$$

Applying the previous theorem the required result follows.  $\square$

Writing equation (26) as

$$\begin{aligned} u_t^\varepsilon = (0.5x^2\sigma^2 + \sigma^2e^{rt}a^2x^4u_x^\varepsilon + \varepsilon) u_{xx}^\varepsilon + 2\sigma^2a^2e^{rt}x^3(u_x^\varepsilon)^2 \\ + (r - q + \sigma^2)xu_x^\varepsilon - qu^\varepsilon, \end{aligned} \quad (42)$$



and differentiating with respect to  $x$ , we obtain

$$\begin{aligned} u_{tx}^\varepsilon &= \left( (0.5x^2\sigma^2 + \sigma^2 e^{rt} a^2 x^4 u_x^\varepsilon + \varepsilon) u_{xx}^\varepsilon \right)_x + 4\sigma^2 e^{rt} a^2 x^3 u_x^\varepsilon u_{xx}^\varepsilon \\ &\quad + (r - q + \sigma^2) x u_{xx}^\varepsilon + 6\sigma^2 e^{rt} a^2 x^2 (u_x^\varepsilon)^2 + (r - 2q + \sigma^2) u_x^\varepsilon. \end{aligned}$$

Setting  $w = u_x^\varepsilon$ , we arrive at the equation

$$\begin{aligned} w_t &= \left( (0.5x^2\sigma^2 + \sigma^2 e^{rt} a^2 x^4 w + \varepsilon) w_x \right)_x + 4\sigma^2 e^{rt} a^2 x^3 w w_x \\ &\quad + (r - q + \sigma^2) x w_x + 6\sigma^2 e^{rt} a^2 x^2 w^2 + (\sigma^2 - 2q + r) w \end{aligned} \quad (43)$$

with initial and boundary conditions

$$w(x, 0) = v_0'(x), \quad w(0, t) = w(b, t) = 0. \quad (44)$$

**Theorem 17.** *Let  $w$  be the weak solution of (43)-(44) then*

$$\|w\|_{L^\infty(0,T,L^\infty(0,b))} \leq C\varepsilon^{-4} \quad (45)$$

where  $C = C(a, \sigma, r, q, b, T)$  does not depend on  $\varepsilon$ .

*Proof.* The proof is similar to that of Theorem 13. Multiplying (43) by  $w^{2k-1}$ ,  $k \geq 1$ , and integrating with respect to  $x$ , we obtain

$$\begin{aligned} &\int_0^b w_t w^{2k-1} dx + (2k-1) \int_0^b (0.5x^2\sigma^2 + \sigma^2 e^{rt} a^2 x^4 w + \varepsilon) w_x^2 w^{2k-2} dx \\ &= \int_0^b 4\sigma^2 e^{rt} a^2 x^3 w_x w^{2k} dx + \int_0^b (r - q + \sigma^2) x w_x w^{2k-1} dx \\ &\quad + \int_0^b 6\sigma^2 e^{rt} a^2 x^2 w^{2k+1} dx + \int_0^b (\sigma^2 + r - 2q) w^{2k} dx. \end{aligned}$$

Applying integration by parts to the right hand side, we get

$$\begin{aligned} &\frac{1}{2k} \frac{d}{dt} \int_0^b w^{2k} dx + (2k-1) \int_0^b (0.5x^2\sigma^2 + \sigma^2 e^{rt} a^2 x^4 w + \varepsilon) w_x^2 w^{2k-2} dx \\ &= - \int_0^b \frac{12}{2k+1} \sigma^2 e^{rt} a^2 x^2 w^{2k+1} dx - \int_0^b \frac{r - q + \sigma^2}{2k} w^{2k} dx \\ &\quad + \int_0^b 6\sigma^2 e^{rt} a^2 x^2 w^{2k+1} dx + \int_0^b (\sigma^2 + r - 2q) w^{2k} dx. \end{aligned}$$

Taking into account the bounds of the coefficients, we have

$$\frac{1}{2k} \frac{d}{dt} \int_0^b w^{2k} dx + (2k-1)\varepsilon \int_0^b w_x^2 w^{2k-2} dx \leq C \int_0^b w^{2k+1} dx + C \int_0^b w^{2k} dx.$$

Ignoring the second term, because it is nonnegative by Remark 10, and using a Sobolev embedding, we obtain

$$\|w\|_{L^{2k}}^{2k-1} \frac{d}{dt} \|w\|_{L^{2k}} \leq C \|w\|_{L^\infty}^2 b^{\frac{2k-1}{(2k-1)(2k)}} \|w\|_{L^{2k}}^{2k-1} + C \|w\|_{L^{2k}}^{2k}.$$

Therefore

$$\frac{d}{dt} \|w\|_{L^{2k}} \leq C \|w\|_{L^\infty}^2 b^{\frac{1}{2k}} + C \|w\|_{L^{2k}}.$$

Applying Gronwall's inequality and Corollary 15,

$$\begin{aligned} \|w\|_{L^{2k}} &\leq C \|w(x, 0)\|_{L^{2k}} + C b^{\frac{1}{2k}} \int_0^T \|w\|_{L^\infty}^2 dt \\ &\leq C \|w(x, 0)\|_{L^{2k}} + C b^{\frac{1}{2k}} \varepsilon^{-4}. \end{aligned}$$

Finally taking  $k \rightarrow \infty$  and attending to (34), the required follows.  $\square$

This result finishes this section.

## 5 Convergence

In this section we prove the convergence of the approximate solutions  $u^\varepsilon$  of the viscosity solution  $u$  of (14)-(16) and (19). The idea is that the so-called upper weak limit  $\bar{u}$  and the lower weak limit  $\underline{u}$  are, respectively, a viscosity subsolution and supersolution of (20). On, the one hand, we always have  $\underline{u} \leq \bar{u}$  in  $Q_T$ , and on the other hand, the comparison principle implies that  $\bar{u} \leq \underline{u}$  a.e. in  $Q_T$ . Finally, it is easy to see that this equality implies the local  $L^\infty$  convergence of  $u^\varepsilon$  to the function  $u = \underline{u} = \bar{u}$  as  $\varepsilon \rightarrow 0$ , which turns out to be a unique bounded viscosity solution of (25).

With the estimates in the last section in hand we can now prove the existence and uniqueness of a classical solution  $u^\varepsilon$ .

**Theorem 18.** *Problem (26)-(27) has a unique classical solution*

$$u^\varepsilon \in C^{2+\alpha, 1+\alpha/2}(\bar{Q}_T). \quad (46)$$

*Proof.* Attending to (38) and (45) we are able to consider Equation (26) as a linear equation with bounded coefficients. So, applying Theorem 5.1 from chapter VI in [15], we obtain that  $u_x^\varepsilon \in C^{\alpha, \alpha/2}(\bar{Q}_T)$   $0 < \alpha < 1$ . Which allows us to prove that the coefficients are Hölder continuous. Using the Leray-Schauder theory, namely Theorem 5.2 from chapter IV in [15], we conclude that, there exists a unique classical solution of (26) satisfying (46).  $\square$

Taking into account conditions (21) and (22), we have

$$\begin{aligned}\bar{u}(x, t) &= \lim_{r \rightarrow 0} \sup \{u^\varepsilon(y, \tau) : \|(y, \tau) - (x, t)\| \leq r, \varepsilon \leq r, (y, \tau) \in Q_T\} \quad (47) \\ &= \lim_{\varepsilon \rightarrow 0} \sup^* u^\varepsilon(t, x) < \infty\end{aligned}$$

$$\begin{aligned}\underline{u}(x, t) &= \lim_{r \rightarrow 0} \inf \{u^\varepsilon(y, \tau) : \|(y, \tau) - (x, t)\| \geq r, \varepsilon \leq r, (y, \tau) \in Q_T\} \quad (48) \\ &= \lim_{\varepsilon \rightarrow 0} \inf_* u^\varepsilon(t, x) > -\infty\end{aligned}$$

Since  $u^\varepsilon$  is the classical solution of (26)-(27), it is also viscosity solution for each  $\varepsilon > 0$ . Applying Proposition 4.7 in Chapter V of [3] with appropriate adaptations to parabolic equations we can prove the next result.

**Theorem 19.** *Let  $u^\varepsilon(x, t)$  be the classical solution of (26)-(27) then  $\bar{u}$  and  $\underline{u}$  defined by (47) and (48) are sub and super viscous solutions of (25), respectively.*

In order to apply the comparison theorem, we require the next two lemmas.

**Lemma 20.** *The operator  $F$  defined in (25) is proper, that is*

$$F(x, t, s, p, X) \leq F(x, t, s, p, Y) \quad \text{whenever} \quad X \geq Y \quad (49)$$

and

$$F(x, t, r, p, X) \leq F(x, t, s, p, X) \quad \text{whenever} \quad r \leq s \quad (50)$$

where  $x, t, r, s, p, X, Y \in \mathbb{R}$ .

*Proof.* By the definition of  $F$  and the assumptions on the problem, we have

$$F(x, t, s, p, X) - F(x, t, s, p, Y) = \underbrace{(0.5x^2\sigma^2(1 + 2e^{rt}a^2x^2p))}_{\geq 0} (Y - X) \leq 0$$

whenever  $X \geq Y$

and

$$F(x, t, r, p, X) - F(x, t, s, p, X) = q(r - s) < 0 \quad \text{whenever} \quad r \leq s.$$

□

**Lemma 21.** *Let  $F$  be defined by (25). Then there exists  $\gamma > 0$  such that*

$$\gamma(r - s) \leq F(x, t, r, p, X) - F(x, t, s, p, X) \text{ for } r \geq s \text{ and } x, t, p, X \in \mathbb{R},$$

and there is a function  $\omega : [0, \infty] \rightarrow [0, \infty]$  that satisfies  $\omega(0^+) = 0$  such that

$$F(y, t, s, \alpha(x - y), Y) - F(x, t, s, \alpha(x - y), X) \leq \omega(\alpha|x - y|^2 + |x - y|)$$

for  $x, y \in ]0, b[, t \in ]0, T[$  fixed,  $X, Y \in \mathbb{R}$  and  $\alpha$  given by

$$-3\alpha \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \leq \begin{bmatrix} X & 0 \\ 0 & -Y \end{bmatrix} \leq 3\alpha \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}. \quad (51)$$

*Proof.* The first condition was proved in the previous lemma with  $\gamma = q$ . with respect to the second condition, we have

$$\begin{aligned} F(y, t, s, \alpha(x-y), Y) - F(x, t, s, \alpha(x-y), X) &= 0.5\sigma^2(x^2X - y^2Y) + \\ &+ \sigma^2 e^{rt} a^2 \alpha(x-y)(x^4X - y^4Y) + (r-q)\alpha(x-y)^2 + \\ &+ 2\sigma^2 e^{rt} a^2 \alpha^2(x-y)^2(x^3 - y^3). \end{aligned}$$

Using the estimates given by (51) and some power inequalities, we have

$$F(y, t, s, \alpha(x-y), Y) - F(x, t, s, \alpha(x-y), X) \leq C(|x-y| + \alpha|x-y|^2).$$

The result now follows, taking  $\omega(s) = Cs$ . □

Taking into account Theorem 19 and Theorem 8.2 in [8] we now have the following theorem.

**Theorem 22.** *Let  $\bar{u}$  and  $\underline{u}$  be defined by (47) and (48) respectively then  $\bar{u} \leq \underline{u}$  a.e in  $Q_T$ .*

By definition  $\underline{u} \leq \bar{u}$  in  $Q_T$  and by the previous theorem  $\underline{u} \geq \bar{u}$  a.e. in  $Q_T$  let us consider  $u = \underline{u} = \bar{u}$ . Using the parabolic analogue of Lemma 1.9 in chapter V of [3] we conclude that, when  $\varepsilon \rightarrow 0^+$ ,  $u^\varepsilon$  converges in  $L^\infty$  to the viscosity solution  $u = \underline{u} = \bar{u}$  of (25).

## 6 Final comments

In this paper we analyzed a nonlinear generalization of the Black- Scholes equations that arises when options are priced under variable transaction costs for buying and selling underlying assets. The mathematical model is represented by a fully nonlinear parabolic equation with the diffusion coefficient depending linearly on the second derivative of the option price. We proved the existence of not necessarily continuous viscosity solutions. The vanishing viscosity method used provides a way to determine approximate numerical solutions. The à priori estimates obtained are useful when we wish to prove analytically the convergence and convergence order of certain numerical methods. Indeed this is part of our future work.

## Acknowledgements

This work was partially supported by the research projects: Grant N. UID/MAT/00212/2019 - financed by FEDER through the - Programa Operacional Factores de Competitividade, FCT - Fundação para a Ciência e a Tecnologia and Grant BID/ICI-FC/Santander Universidades-UBI/2016.

## References

- [1] J. Ankudinova, M. Ehrhardt, On the numerical solution of nonlinear Black-Scholes equations, *Computers and Mathematics with Applications* 56 (3) (2008) 799 – 812, *mathematical Models in Life Sciences and Engineering*.
- [2] M. Avellaneda, A. Paras., Dynamic hedging portfolios for derivative securities in the presence of large transaction costs., *Applied Mathematical Finance* 1 (1994) 165–193.
- [3] M. Bardi, I. Capuzzo-Dolcetta, *Optimal Control and Viscosity Solutions of Hamilton-Jacobi-Bellman Equations*, *Modern Birkhäuser Classics*, Birkhäuser Boston, 2008.
- [4] G. Barles, H. M. Soner, Option pricing with transaction costs and a nonlinear Black-Scholes equation, *Finance Stoch.* 2 (4) (1998) 369–397.
- [5] F. Black, M. Scholes, The pricing of options and corporate liabilities, *J. Polit. Econ.* 81 (3) (1973) 637–654.
- [6] G.-Q. Chen, B. Su, Discontinuous solutions of Hamilton-Jacobi equations: Existence, uniqueness, and regularity, in: T. Y. Hou, E. Tadmor (Eds.), *Hyperbolic Problems: Theory, Numerics, Applications*, Springer Berlin Heidelberg, Berlin, Heidelberg, 2003, pp. 443–453.
- [7] R. Company, L. Jódar, J.-R. Pintos, A numerical method for European option pricing with transaction costs nonlinear equation, *Math. Comput. Modelling* 50 (5-6) (2009) 910–920.
- [8] M. G. Crandall, H. Ishii, P.-L. Lions, User’s guide to viscosity solutions of second order partial differential equations, *Bulletin of the American mathematical society* 27 (1) (1992) 1–67.
- [9] B. Düring, M. Fournié, A. Jüngel, Convergence of a high-order compact finite difference scheme for a nonlinear Black-Scholes equation, *ESAIM: Mathematical Modelling and Numerical Analysis - Modélisation Mathématique et Analyse Numérique* 38 (2) (2004) 359–369.
- [10] M. Ehrhardt, *Nonl. Models in Math. Finance: Research Trends in Option Pricing*, Nova Science Publishers, 2009.
- [11] S. Hodges, A. Neuberger., Optimal replication of contingent claims under transaction costs, *Review of Futures Markets* 8 (1989) 222–239.
- [12] M. N. Koleva, L. G. Vulkov, Fast computational approach to the Delta Greek of non-linear Black-Scholes equations, *Journal of Computational and Applied Mathematics* 340 (2018) 508 – 522.
- [13] M. Kratka, No mystery behind the smile, *Risk* 9 (1998) 67–71.

- [14] Y. Kwok, *Mathematical models of financial derivatives*, Springer finance, Springer, 1998.
- [15] O. Ladyzhenskaja, V. Solonnikov, N. Ural'tseva, *Linear and Quasi-linear Equations of Parabolic Type*, American Mathematical Society, translations of mathematical monographs, American Mathematical Society, 1968.
- [16] H. E. Leland, Option pricing and replication with transactions costs, *The Journal of Finance* 40 (5) (1985) 1283–1301.
- [17] D. C. Lesmana, S. Wang, An upwind finite difference method for a non-linear Black-Scholes equation governing European option valuation under transaction costs, *Appl. Math. Comput.* 219 (16) (2013) 8811–8828.
- [18] D. M. Pooley, P. A. Forsyth, K. R. Vetzal, Numerical convergence properties of option pricing pdes with uncertain volatility, *IMA Journal of Numerical Analysis* 23 (2) (2003) 241–267.
- [19] D. Ševcovic, B. Stehliková, K. Mikula, *Analytical and numerical methods for pricing financial derivatives*, Nova Science Publ. ISBN (2011) 978–1.
- [20] R. Valkov, Convergence of a finite volume element method for a generalized Black-Scholes equation transformed on finite interval, *Numerical Algorithms* 68 (1) (2015) 61–80.
- [21] R. Valkov, Predictor-corrector balance method for the worst-case 1D option pricing, *Computational Methods in Applied Mathematics* 16 (1) (2015) 175–86.
- [22] P. Wilmott, *Derivatives. The Theory and Practice of Financial Engineering*, John Wiley and Sons, 1999.