

Localisation for the torsion function and the strong Hardy inequality

M. van den Berg
School of Mathematics, University of Bristol
Fry Building, Woodland Road
Bristol BS8 1UG, United Kingdom
mamvdb@bristol.ac.uk

T. Kappeler
Institut für Mathematik, Universität Zürich
Winterthurerstrasse 190, CH-8057 Zürich, Switzerland
thomas.kappeler@math.uzh.ch

23 February 2020

Abstract

Two-sided bounds for the efficiency of the torsion function are obtained in terms of the square of the distance to the boundary function under the hypothesis that the Dirichlet Laplacian satisfies a strong Hardy inequality. Localisation properties of the torsion function are obtained under that hypothesis. An example is analysed in detail.

Mathematics Subject Classification (2000): 35J25, 35P99.

Keywords: Torsion function, Dirichlet boundary condition, localisation, efficiency, Hardy inequality.

1 Introduction and main results

Let Ω be an open set in \mathbb{R}^m , $m \geq 1$, with finite Lebesgue measure $|\Omega|$, $0 < |\Omega| < \infty$, and with boundary $\partial\Omega$. The torsion function for Ω is the unique solution of

$$-\Delta v = 1, \quad v \in H_0^1(\Omega),$$

and is denoted by v_Ω . The function v_Ω is non-negative and satisfies,

$$\lambda_1(\Omega)^{-1} \leq \|v_\Omega\|_\infty \leq (4 + 3m \log 2)\lambda_1(\Omega)^{-1}, \quad (1)$$

where

$$\lambda_1(\Omega) = \inf_{\varphi \in H_0^1(\Omega) \setminus \{0\}} \frac{\|\nabla \varphi\|_2^2}{\|\varphi\|_2^2} \quad (2)$$

is the first eigenvalue of the Dirichlet Laplacian and where $\|\cdot\|_p$ denotes the standard L^p norm, $1 \leq p \leq \infty$. The m -dependent constant in the right-hand side of (1) has subsequently been improved ([17], [24]). We denote the sharp constant by \mathfrak{c}_m ,

$$\mathfrak{c}_m = \sup\{\lambda_1(\Omega)\|v_\Omega\|_\infty : \Omega \text{ open in } \mathbb{R}^m, 0 < |\Omega| < \infty\}. \quad (3)$$

The torsion function and its L^1 norm, the torsional rigidity, play key roles in different parts of analysis. For example the torsional rigidity of a cross section of a beam appears in the computation of the angular change when a beam of a given length and a given modulus of rigidity is exposed to a twisting moment ([4],[22]). It also arises in the definition of gamma convergence [10] and in the study of minimal submanifolds [20]. Moreover, $v_\Omega(x)$ equals the expected lifetime of Brownian motion in Ω starting at $x \in \Omega$. This immediately implies the non-negativity of v_Ω , and that for open sets Ω_1, Ω_2 ,

$$\Omega_1 \subset \Omega_2 \Rightarrow v_{\Omega_1}(x) \leq v_{\Omega_2}(x), \forall x \in \Omega_1. \quad (4)$$

This in turn implies that the torsional rigidity $\|v_\Omega\|_1$ is monotone increasing in Ω . The torsion function has been studied extensively, and numerous works have been written on this subject. We just mention the paper [9], and the references therein.

The torsion function is defined for an open subset $\Omega \subset \mathbb{R}^m$ of infinite measure, provided the bottom of the spectrum of the Dirichlet Laplacian, denoted by $\lambda_1(\Omega)$, is bounded away from 0. Indeed, by considering an increasing sequence of sets $\Omega \cap B(0; k)$, $k \in \mathbb{N}$, where $B(p; R) = \{x \in \mathbb{R}^m : |p - x| < R\}$ denotes the open ball with centre p and radius R , one obtains a weak solution of

$$-\Delta v_\Omega = 1, \quad v|_{\partial\Omega} = 0. \quad (5)$$

This solution is non-negative, satisfies both (4) and (1), and satisfies the probabilistic interpretation mentioned above. See Theorem 1 in [7], Theorem 5.3 in [5], and [10] for further details.

We remark that the torsion function has been defined in greater generality. Instead of the Dirichlet Laplacian one considers a positive, self-adjoint Schrödinger operator $L := -\Delta + V$, where V is bounded, measurable, and non-negative (or even more generally, a positive, self-adjoint, elliptic operator of second order) acting in $L^2(\Omega)$ with Dirichlet or Neumann boundary conditions or on closed manifolds. In the latter situation one requires potentials V such that the bottom of the spectrum of the Laplacian is bounded away from 0. We denote by v_L the torsion function associated to L . It was discovered in recent papers [2], [3], and references therein, that under appropriate conditions, the reciprocal v_L^{-1} of the torsion function v_L can be used for approximating eigenvalues and eigenfunctions of the Schrödinger operator L . The phenomenon of

localisation of eigenfunctions of Schrödinger operators is a prominent and very active research area and has important applications in the applied sciences. The literature is extensive. See for example the review paper of [19]. Explicit estimates on the localisation of the eigenfunctions in terms of bounds on their exponential decay away from the subdomains, where they concentrate, were obtained in [3].

In this paper we consider the κ -localisation, the localisation, and the efficiency of the torsion function of the Dirichlet Laplacian on an open set Ω in \mathbb{R}^m with $0 < |\Omega| < \infty$ – see Definition 2 and Definition 3 below. These notions have been first introduced for Dirichlet eigenfunctions of Schrödinger operators and go back to [19] and respectively, [21], [23]. The notion of efficiency, defined as the mean to max ratio, can be viewed as a rough measure of localisation.

In [6], we studied the efficiency and localisation for the torsion function of a Schrödinger operator $-\Delta + V$ acting in $L^2(\Omega)$ with Dirichlet boundary condition. Among other results it was shown, in Theorem 4, that under appropriate conditions localisation for the torsion function implies localisation for the first Dirichlet eigenfunction. The converse does not hold. Consider for example a sequence of ellipsoids (Ω_n) with semi-axes of length 1 and n respectively. We see by Theorem 1(iii) below that (v_{Ω_n}) does not localise. On the other hand it was shown in Example 10 in [8] that the corresponding sequence of first Dirichlet eigenfunctions localises. Unlike the torsion function the first Dirichlet eigenfunction is not monotone on set inclusion, and this in general complicates its analysis.

In this paper, we continue this study. Our results are obtained under the hypothesis that the Dirichlet Laplacian satisfies the strong Hardy inequality, defined as follows.

Definition 1. *The Dirichlet Laplacian $-\Delta$ acting in $L^2(\Omega)$ satisfies the strong Hardy inequality, with constant $c_\Omega \in (0, \infty)$, if*

$$\|\nabla w\|_2^2 \geq \frac{1}{c_\Omega} \int_\Omega \frac{w^2}{d_\Omega^2}, \quad \forall w \in C_c^\infty(\Omega), \quad (6)$$

where d_Ω is the distance to the boundary function,

$$d_\Omega(x) = \inf\{|x - y| : y \in \mathbb{R}^m \setminus \Omega\}, \quad x \in \Omega.$$

Both the validity and applications of inequalities like (6) to spectral theory and partial differential equations have been investigated in depth. See for example [1], [11], [12], [13] and [14]. In particular it was shown in [1, p. 208], that for any proper simply connected open subset Ω in \mathbb{R}^2 , $c_\Omega = 16$.

Our first result in Theorem 1, stated below, states that the efficiency of the torsion function of the Dirichlet Laplacian on an open set Ω of \mathbb{R}^m with $0 < |\Omega| < \infty$ can be bounded from above and below by the efficiency of the square of the distance to the boundary function whereas item (ii) and (iii) of Theorem 1 give sufficient conditions for the localisation of the torsion function to hold and respectively, not to hold.

In Theorem 2(i)–(iii) we analyse an example illustrating the use of Theorem 1. In Theorem 2(iv) we give an example of κ -localisation with $0 < \kappa < 1$.

We now define the notion of efficiency and localisation in precise terms. The efficiency of v_Ω , introduced in [15], is defined as follows.

Definition 2. Let Ω be an open set in \mathbb{R}^m with $0 < |\Omega| < \infty$. The efficiency, or mean to max ratio, of v_Ω is

$$\Phi(\Omega) = \frac{\|v_\Omega\|_1}{|\Omega|\|v_\Omega\|_\infty}.$$

Let (Ω_n) be a sequence of open sets in \mathbb{R}^m with $0 < |\Omega_n| < \infty$, $n \in \mathbb{N}$. We say that (v_{Ω_n}) has vanishing efficiency if $\lim_{n \rightarrow \infty} \Phi(\Omega_n) = 0$.

The following notion of localisation for the torsion function has been motivated by the one for eigenfunctions in [19], and has been used previously in [6].

Definition 3. For any sequence of open sets (Ω_n) in \mathbb{R}^m with $0 < |\Omega_n| < \infty$, $n \in \mathbb{N}$, let

$$\mathfrak{A}((\Omega_n)) = \left\{ (A_n) : (\forall n \in \mathbb{N})(A_n \subset \Omega_n, A_n \text{ measurable}), \lim_{n \rightarrow \infty} \frac{|A_n|}{|\Omega_n|} = 0 \right\}, \quad (7)$$

and

$$\kappa = \sup \left\{ \limsup_{n \rightarrow \infty} \frac{\int_{A_n} v_{\Omega_n}}{\|v_{\Omega_n}\|_1} : (A_n) \in \mathfrak{A}((\Omega_n)) \right\}. \quad (8)$$

We say that if (i) $0 < \kappa < 1$ then (v_{Ω_n}) κ -localises, (ii) $\kappa = 1$ then (v_{Ω_n}) localises, (iii) $\kappa = 0$ then (v_{Ω_n}) does not localise.

Before stating our main results we review some basic facts. We note that

$$B(x; d_\Omega(x)) \subset \Omega, \quad (9)$$

and $|B(x; d_\Omega(x))| \leq |\Omega|$. Hence d_Ω is bounded from above and

$$\|d_\Omega\|_\infty \leq (\omega_m^{-1}|\Omega|)^{1/m},$$

where $\omega_m = |B(0; 1)|$. Since the torsion function is pointwise increasing with respect to the domain,(4), it follows by (9) that,

$$v_\Omega(x) \geq v_{B(x; d_\Omega(x))}(x) = \frac{d_\Omega(x)^2}{2m}, \quad \forall x \in \Omega, \quad (10)$$

where we have used that

$$v_{B(p; R)}(x) = \frac{1}{2m}(R^2 - |x - p|^2), \quad \forall x \in B(p; R).$$

By (10),

$$\frac{1}{2m}\|d_\Omega^2\|_1 \leq \|v_\Omega\|_1. \quad (11)$$

Under the additional assumption that Ω satisfies (6), it was shown in [7, Theorem 2] that

$$\|v_\Omega\|_1 \leq c_\Omega\|d_\Omega^2\|_1. \quad (12)$$

Furthermore d_Ω is uniformly continuous. Hence for any $\eta \in \mathbb{R}^+$, the subset $\{d_\Omega \geq \eta\}$ is relatively closed in Ω , and hence measurable.

Throughout we denote for $\nu \geq 0$, by j_ν the first positive zero of the Bessel function J_ν . For the ball $B(p; 1) \subset \mathbb{R}^m$, $m \geq 2$, we have $\lambda_1(B(p; 1)) = j_{(m-2)/2}^2$. Our main results are as follows.

Theorem 1. (i) If Ω is an open subset of \mathbb{R}^m with $0 < |\Omega| < \infty$, and which satisfies (6) with the strong Hardy constant c_Ω , then

$$(2m\mathfrak{c}_m c_\Omega)^{-1} \frac{\|d_\Omega^2\|_1}{|\Omega|\|d_\Omega^2\|_\infty} \leq \Phi(\Omega) \leq c_\Omega j_{(m-2)/2}^2 \frac{\|d_\Omega^2\|_1}{|\Omega|\|d_\Omega^2\|_\infty}, \quad (13)$$

where \mathfrak{c}_m is the sharp constant in (3).

(ii) Let (Ω_n) be a sequence of open sets in \mathbb{R}^m with $0 < |\Omega_n| < \infty$, $n \in \mathbb{N}$, and which satisfies (6) with strong Hardy constants c_{Ω_n} . Suppose

$$c = \sup\{c_{\Omega_n} : n \in \mathbb{N}\} < \infty. \quad (14)$$

If (η_n) is a sequence of strictly positive real numbers such that

$$\lim_{n \rightarrow \infty} \frac{|\{d_{\Omega_n} \geq \eta_n\}|}{|\Omega_n|} = 0, \quad (15)$$

and

$$\lim_{n \rightarrow \infty} \frac{\eta_n^2 |\Omega_n|}{\int_{\{d_{\Omega_n} \geq \eta_n\}} d_{\Omega_n}^2} = 0, \quad (16)$$

then (v_{Ω_n}) localises along the sequence $A_n = \{x \in \Omega_n : d_{\Omega_n} \geq \eta_n\}$.

(iii) Let (Ω_n) be a sequence of open sets in \mathbb{R}^m with $0 < |\Omega_n| < \infty$, $n \in \mathbb{N}$, which satisfies (6) with strong Hardy constants c_{Ω_n} . Suppose (14) holds. If any sequence (A_n) of measurable sets, $A_n \subset \Omega_n$, $n \in \mathbb{N}$, with

$$\lim_{n \rightarrow \infty} \frac{|A_n|}{|\Omega_n|} = 0,$$

satisfies

$$\lim_{n \rightarrow \infty} \frac{\int_{A_n} d_{\Omega_n}^2}{\int_{\Omega_n} d_{\Omega_n}^2} = 0,$$

then (v_{Ω_n}) does not localise.

Theorem 1(i) can be interpreted as follows. Given an open subset Ω of \mathbb{R}^m with $0 < |\Omega| < \infty$, we define the efficiency of d_Ω^2 as

$$D(\Omega) = \frac{\|d_\Omega^2\|_1}{|\Omega|\|d_\Omega^2\|_\infty}.$$

Theorem 1(i) asserts that under condition (6), the efficiencies of v_Ω and d_Ω^2 are comparable.

Let (Ω_n) be a sequence of open sets in \mathbb{R}^m with $0 < |\Omega_n| < \infty$. We say that $(d_{\Omega_n}^2)$ localises if

$$\sup \left\{ \limsup_{n \rightarrow \infty} \frac{\int_{A_n} d_{\Omega_n}^2}{\|d_{\Omega_n}^2\|_1} : (A_n) \in \mathfrak{A}((\Omega_n)) \right\} = 1,$$

where $\mathfrak{A}((\Omega_n))$ is given by (7). Theorem 1(iii) asserts that for any sequence (Ω_n) satisfying the conditions of Theorem 1(iii) the following holds: if $(d_{\Omega_n}^2)$ does not localise, then neither does (v_{Ω_n}) .

In Theorem 2 below we analyse the torsion function for a sequence of open simply connected sets $(\Omega_{\varepsilon_n, n})_{n \geq 4} \subset \mathbb{R}^2$ with $0 < \varepsilon_n < 1$. The construction is as follows. Let Q be the open unit square in \mathbb{R}^2 with vertices $(0, 0)$, $(1, 0)$, $(0, 1)$, $(1, 1)$. Let $n \in \mathbb{N}$, and for any given $n \in \mathbb{N}$ let L_1, \dots, L_{n-1} be the closed line segments of lengths $1 - \varepsilon_n$ with endpoints $(\frac{1}{n}, 0)$, $(\frac{2}{n}, 0)$, \dots , $(\frac{n-1}{n}, 0)$ pointing in the direction $(0, 1)$. Let

$$\Omega_{\varepsilon_n, n} = Q \setminus \cup_{j=1}^{n-1} L_j.$$




Figure 1: $\Omega_{\frac{1}{16}, 16}$

Theorem 2. Let $0 < \alpha < 1$, $c > 0$,

$$\varepsilon_n = cn^{-\alpha},$$

and

$$N_{\alpha, c} = \min\{n \in \mathbb{N} : n^\alpha \geq 2c, cn^{1-\alpha} \geq 2\}.$$

(Note that $N_{\alpha, c} \geq 4$.)

(i) If $n \geq N_{\alpha, c}$, then

$$\frac{1}{3072c_2}(cn^{-\alpha} + c^{-2}n^{2\alpha-2}) \leq \Phi(\Omega_{\varepsilon_n, n}) \leq \frac{64j_0^2}{3}(cn^{-\alpha} + c^{-2}n^{2\alpha-2}), \quad (17)$$

where c_2 is given by (3).

(ii) If $0 < \alpha < \frac{2}{3}$, then $(v_{\Omega_{\varepsilon_n, n}})$ localises along the sequence

$$A_n = \{x \in \Omega_{\varepsilon_n, n} : d_{\Omega_{\varepsilon_n, n}}(x) \geq \frac{1}{2n}\}. \quad (18)$$

(iii) If $\frac{2}{3} < \alpha < 1$, then $(v_{\Omega_{\varepsilon_n, n}})$ is not localising.

(iv) If $\alpha = \frac{2}{3}$, then $(v_{\Omega_{\varepsilon_n, n}})$ is κ_c -localising along (18) with

$$\kappa_c = \frac{c^3}{1 + c^3}. \quad (19)$$

The paper is organised as follows. The proofs of Theorem 1 and Theorem 2(i)–(iii) are given in Section 2. The proof of Theorem 2(iv) involves tools from Brownian motion and is deferred to Section 3.

2 Proofs of Theorem 1 and Theorem 2(i)–(iii)

Proof of Theorem 1(i). Since $|\Omega| < \infty$, the inradius $\|d_\Omega\|_\infty$ is finite, and by (2) and (6),

$$\begin{aligned} \lambda_1(\Omega) &\geq c_\Omega^{-1} \inf_{\varphi \in H_0^1(\Omega), \|\varphi\|_2=1} \int_\Omega \frac{\varphi^2}{d_\Omega^2} \\ &\geq c_\Omega^{-1} \|d_\Omega\|_\infty^{-2} \inf_{\varphi \in H_0^1(\Omega), \|\varphi\|_2=1} \|\varphi\|_2^2 \\ &= c_\Omega^{-1} \|d_\Omega\|_\infty^{-2}. \end{aligned} \quad (20)$$

By (3) and (20),

$$\|v_\Omega\|_\infty \leq \mathfrak{c}_m \lambda_1(\Omega)^{-1} \leq \mathfrak{c}_m c_\Omega \|d_\Omega\|_\infty^2. \quad (21)$$

The lower bound in (13) follows from (21) and the lower bound in (11).

Since Ω has inradius $\|d_\Omega\|_\infty$, Ω contains an open ball with radius $\|d_\Omega\|_\infty$ and centre p_Ω . By the monotonicity of the Dirichlet eigenvalues

$$\lambda_1(\Omega) \leq \lambda_1(B(p_\Omega; \|d_\Omega\|_\infty)) = j_{(m-2)/2}^2 \|d_\Omega\|_\infty^{-2}. \quad (22)$$

By the first inequality in (1), and (22),

$$\|v_\Omega\|_\infty^{-1} \leq j_{(m-2)/2}^2 \|d_\Omega\|_\infty^{-2}. \quad (23)$$

The upper bound in (13) follows from (23) and the upper bound in (12).

Proof of Theorem 1(ii). By (6), and suppressing n dependence,

$$\begin{aligned} \int_\Omega v_\Omega &= - \int_\Omega v_\Omega \Delta v_\Omega = \int_\Omega |\nabla v_\Omega|^2 \\ &\geq \frac{1}{c_\Omega} \int_\Omega \frac{v_\Omega^2}{d_\Omega^2} \geq \frac{1}{c_\Omega} \int_{\{d_\Omega < \eta\}} \frac{v_\Omega^2}{d_\Omega^2} \\ &\geq \frac{1}{c_\Omega \eta^2} \int_{\{d_\Omega < \eta\}} v_\Omega^2. \end{aligned} \quad (24)$$

On the other hand by Cauchy-Schwarz,

$$\left(\int_{\{d_\Omega < \eta\}} v_\Omega \right)^2 \leq |\{d_\Omega < \eta\}| \int_{\{d_\Omega < \eta\}} v_\Omega^2. \quad (25)$$

Combining (24) and (25) yields,

$$\left(\int_{\{d_\Omega < \eta\}} v_\Omega \right)^2 \leq c_\Omega \eta^2 |\Omega| \left(\int_{\{d_\Omega < \eta\}} v_\Omega + \int_{\{d_\Omega \geq \eta\}} v_\Omega \right). \quad (26)$$

Solving the quadratic inequality (26) gives,

$$\begin{aligned} \int_{\{d_\Omega < \eta\}} v_\Omega &\leq \frac{c_\Omega \eta^2 |\Omega|}{2} + \left(\frac{c_\Omega^2 \eta^4 |\Omega|^2}{4} + c_\Omega \eta^2 |\Omega| \int_{\{d_\Omega \geq \eta\}} v_\Omega \right)^{1/2} \\ &\leq c_\Omega \eta^2 |\Omega| + \left(c_\Omega \eta^2 |\Omega| \int_{\{d_\Omega \geq \eta\}} v_\Omega \right)^{1/2}. \end{aligned} \quad (27)$$

By (10),

$$\int_{\{d_\Omega \geq \eta\}} v_\Omega \geq \frac{1}{2m} \int_{\{d_\Omega \geq \eta\}} d_\Omega^2. \quad (28)$$

By (27) and (28)

$$\frac{\int_{\{d_\Omega < \eta\}} v_\Omega}{\int_{\{d_\Omega \geq \eta\}} v_\Omega} \leq \frac{2m c_\Omega \eta^2 |\Omega|}{\int_{\{d_\Omega \geq \eta\}} d_\Omega^2} + \left(\frac{2m c_\Omega \eta^2 |\Omega|}{\int_{\{d_\Omega \geq \eta\}} d_\Omega^2} \right)^{1/2}. \quad (29)$$

To complete the proof we suppose that (Ω_n) and (η_n) are sequences satisfying (15) and (16) respectively, and that c_{Ω_n} satisfies (14). If

$$\lim_{n \rightarrow \infty} \frac{\eta_n^2 |\Omega_n|}{\int_{\{d_{\Omega_n} \geq \eta_n\}} d_{\Omega_n}^2} = 0,$$

then by (29)

$$\lim_{n \rightarrow \infty} \frac{\int_{\{d_{\Omega_n} < \eta_n\}} v_{\Omega_n}}{\int_{\{d_{\Omega_n} \geq \eta_n\}} v_{\Omega_n}} = 0,$$

and

$$\lim_{n \rightarrow \infty} \frac{\int_{\{d_{\Omega_n} < \eta_n\}} v_{\Omega_n}}{\int_{\Omega_n} v_{\Omega_n}} = 0.$$

Hence

$$\lim_{n \rightarrow \infty} \frac{\int_{\{d_{\Omega_n} \geq \eta_n\}} v_{\Omega_n}}{\int_{\Omega_n} v_{\Omega_n}} = 1. \quad (30)$$

Let $A_n = \{d_{\Omega_n} \geq \eta_n\}$. Then A_n , $n \in \mathbb{N}$, is relatively closed in Ω , and hence measurable. By hypothesis (15),

$$\lim_{n \rightarrow \infty} \frac{|A_n|}{|\Omega_n|} = 0,$$

and by (30) we have $\limsup_{n \rightarrow \infty} \frac{\int_{A_n} v_{\Omega_n}}{\|v_{\Omega_n}\|_1} = 1$. Hence (8) holds with $\kappa = 1$.

Proof of Theorem 1(iii). By Cauchy-Schwarz, and suppressing n -dependence,

$$\left(\int_A v_\Omega \right)^2 \leq \left(\int_A d_\Omega^2 \right) \left(\int_A \frac{v_\Omega^2}{d_\Omega^2} \right). \quad (31)$$

By (6), the first two equalities in (24), and by (31) we obtain

$$\begin{aligned} \int_{\Omega} v_{\Omega} &\geq \frac{1}{c_{\Omega}} \int_A \frac{v_{\Omega}^2}{d_{\Omega}^2} \\ &\geq \frac{1}{c_{\Omega}} \frac{(\int_A v_{\Omega})^2}{\int_A d_{\Omega}^2}. \end{aligned}$$

Hence

$$\int_A v_{\Omega} \leq c_{\Omega}^{1/2} \left(\int_{\Omega} v_{\Omega} \right)^{1/2} \left(\int_A d_{\Omega}^2 \right)^{1/2},$$

and together with (11)

$$\frac{\int_A v_{\Omega}}{\int_{\Omega} v_{\Omega}} \leq c_{\Omega}^{1/2} \frac{\left(\int_A d_{\Omega}^2 \right)^{1/2}}{\left(\int_{\Omega} v_{\Omega} \right)^{1/2}} \leq (2mc_{\Omega})^{1/2} \left(\frac{\int_A d_{\Omega}^2}{\int_{\Omega} v_{\Omega}} \right)^{1/2}.$$

This implies the assertion of Theorem 1(iii). \square

Proof of Theorem 2(i). We use Theorem 1(i). First note that for a rectangle $R_{a,b}$ with side lengths a and b , where $a \geq b$ one has

$$\|d_{R_{a,b}}\|_2^2 = \frac{ab^3}{12} - \frac{b^4}{24} \geq \frac{ab^3}{24}. \quad (32)$$

To obtain a lower bound for $\|d_{\tilde{\Omega}_{\varepsilon_n, n}}^2\|_1$ we add a closed line segment of length 1 to the boundary of $\Omega_{\varepsilon_n, n}$, and obtain that the resulting set is the union of $n+1$ disjoint rectangles, denoted by $\tilde{\Omega}_{\varepsilon_n, n}$. See Figure 2. Hence, by (32),

$$\|d_{\tilde{\Omega}_{\varepsilon_n, n}}^2\|_1 \geq \|d_{\Omega_{\varepsilon_n, n}}^2\|_1 \geq \frac{1-\varepsilon_n}{24n^2} + \frac{\varepsilon_n^3}{24} \geq \frac{1}{48}(\varepsilon_n^3 + n^{-2}), \quad (33)$$

where we have used that $\varepsilon_n \leq \frac{1}{2}$ for $n \geq N_{\alpha, c}$.

To obtain an upper bound for $\|d_{\tilde{\Omega}_{\varepsilon_n, n}}^2\|_1$, we have the following estimate.

$$d_{\tilde{\Omega}_{\varepsilon_n, n}}(x) \leq \begin{cases} \frac{1}{2n}, & \forall (x_1, x_2) \in \{\Omega_{\varepsilon_n, n} : x_2 \leq 1 - \varepsilon_n\}, \\ 1 - x_2, & \forall (x_1, x_2) \in \{\Omega_{\varepsilon_n, n} : x_2 \geq 1 - \varepsilon_n\}. \end{cases} \quad (34)$$

Since

$$|\{\Omega_{\varepsilon_n, n} : x_2 \leq 1 - \varepsilon_n\}| = 1 - \varepsilon_n \leq 1, \quad (35)$$

we have by (34) and (35),

$$\int_{\{\Omega_{\varepsilon_n, n} : x_2 \leq 1 - \varepsilon_n\}} d_{\tilde{\Omega}_{\varepsilon_n, n}}^2 \leq \frac{1}{4n^2} \quad (36)$$

and

$$\int_{\{\Omega_{\varepsilon_n, n} : x_2 \geq 1 - \varepsilon_n\}} d_{\tilde{\Omega}_{\varepsilon_n, n}}^2 \leq \int_{1-\varepsilon_n}^1 dx_2 (1-x_2)^2 = \frac{\varepsilon_n^3}{3}. \quad (37)$$

Hence we have

$$\|d_{\tilde{\Omega}_{\varepsilon_n, n}}^2\|_1 \leq \frac{1}{3}(\varepsilon_n^3 + n^{-2}). \quad (38)$$




Figure 2: $\tilde{\Omega}_{\frac{1}{16}, 16}$

Finally we observe that

$$\frac{1}{2}\varepsilon_n \leq \|d_{\Omega_{\varepsilon_n, n}}\|_{\infty} \leq \varepsilon_n, \quad n \geq N_{\alpha, c}, \quad (39)$$

and (17) follows from (33), (38), (39), Theorem 1(i), $|\Omega_{\varepsilon_n, n}| = 1$, and that for a proper simply connected subset Ω in \mathbb{R}^2 , $c_{\Omega} = 16$, [1]. This proves the assertion of Theorem 2(i).

Proof of Theorem 2(ii). We use Theorem 1(ii), and choose $\eta_n = \frac{1}{2n}$. Then

$$|\{d_{\Omega_{\varepsilon_n, n}} \geq \frac{1}{2n}\}| \leq \varepsilon_n,$$

and (15) holds. Since for $n \geq N_{\alpha, c}$, $\{d_{\Omega_{\varepsilon_n, n}} \geq \frac{1}{2n}\}$ contains a rectangle with side lengths $1 - n^{-1}$ and $cn^{-\alpha} - \frac{1}{n}$ respectively, we have by (32)

$$\begin{aligned} \int_{\{d_{\Omega_{\varepsilon_n, n}} \geq \frac{1}{2n}\}} d_{\Omega_{\varepsilon_n, n}}^2 &\geq \frac{1}{24}(1 - n^{-1})(cn^{-\alpha} - n^{-1})^3 \\ &\geq \frac{c^3}{192}(1 - n^{-1})n^{-3\alpha} \\ &\geq \frac{c^3}{256}n^{-3\alpha}, \end{aligned}$$

where we have used that $cn^{1-\alpha} \geq 2$ for $n \geq N_{\alpha, c} \geq 4$. Hence

$$\frac{\eta_n^2 |\Omega_{\varepsilon_n, n}|}{\int_{\{d_{\Omega_{\varepsilon_n, n}} \geq \eta_n\}} d_{\Omega_{\varepsilon_n, n}}^2} \leq \frac{64n^{3\alpha-2}}{c^3}, \quad n \geq N_{\alpha, c},$$

and (16) holds for $0 < \alpha < \frac{2}{3}$. This proves the assertion of Theorem 2(ii) by Theorem 1(ii).

Proof of Theorem 2(iii). We use Theorem 1(iii), and let for $n \in \mathbb{N}$, A_n be an arbitrary measurable subset of $\Omega_{\varepsilon_n, n}$ such that $\lim_{n \rightarrow \infty} |A_n| = 0$. By (33),

$$\|d_{\Omega_{\varepsilon_n, n}}^2\|_1 \geq \frac{1}{48n^2}. \quad (40)$$

By (34),

$$\int_{A_n \cap \{x_2 \leq 1 - \varepsilon_n\}} d_{\Omega_{\varepsilon_n, n}}^2 \leq \frac{|A_n|}{4n^2}. \quad (41)$$

By (37), (40), (41),

$$\frac{\int_{A_n} d_{\Omega_{\varepsilon_n, n}}^2}{\int_{\Omega_{\varepsilon_n, n}} d_{\Omega_{\varepsilon_n, n}}^2} \leq c^3 n^{2-3\alpha} + 12|A_n|.$$

This implies

$$\lim_{n \rightarrow \infty} \frac{\int_{A_n} d_{\Omega_{\varepsilon_n, n}}^2}{\int_{\Omega_{\varepsilon_n, n}} d_{\Omega_{\varepsilon_n, n}}^2} = 0,$$

since $\frac{2}{3} < \alpha < 1$, and the hypothesis on A_n . This proves the assertion of Theorem 2(iii) by Theorem 1(iii). \square

3 Proof of Theorem 2(iv)

Let $(B(s), s \geq 0; \mathbb{P}_x, x \in \mathbb{R}^d)$ be Brownian motion associated to the Laplacian in \mathbb{R}^d . Here $(B(s), s \geq 0)$ takes values in \mathbb{R}^d , \mathbb{P}_x is Wiener measure with $\mathbb{P}_x(B(0) = x) = 1$, and for every Borel set $A \in \mathbb{R}^d$,

$$\mathbb{P}_x(B(s) \in A) = (4\pi s)^{-d/2} \int_A dy e^{-|x-y|^2/(4s)}, \quad s > 0. \quad (42)$$

If $\Omega \subset \mathbb{R}^d$ is open and $x \in \Omega$ then we denote the first exit time (or life time) of Ω by

$$T_\Omega = \inf\{s \geq 0 : B(s) \in \mathbb{R}^d \setminus \Omega\}.$$

It is convenient to denote the first hitting time of a closed set C by

$$\tau_C = \inf\{s \geq 0 : B(s) \in C\}.$$

It is well-known that the unique weak solution u_Ω of

$$\Delta u = \frac{\partial u}{\partial t}, \quad \text{in } \Omega \times (0, \infty),$$

with $u|_{\partial\Omega \times (0, \infty)} = 0$, and with initial datum

$$u(x; 0) = 1, \quad x \in \Omega,$$

is given by

$$u_\Omega(x; t) = \mathbb{P}_x(T_\Omega > t). \quad (43)$$

Since

$$v_\Omega(x) = \int_0^\infty dt u_\Omega(x; t) \quad (44)$$

we obtain by (43) and (44) that

$$v_\Omega(x) = \mathbb{E}_x(T_\Omega),$$

and

$$\|v_\Omega\|_1 = \int_\Omega dx \int_0^\infty dt u_\Omega(x; t).$$

For the facts above we refer to [16]. Now let $d = 2$, and let $B = (B_1, B_2)$, where B_1 and B_2 are independent one-dimensional Brownian motions with probability measures $\mathbb{P}_{x_1}^{(1)}$, and $\mathbb{P}_{x_2}^{(1)}$ respectively, and $\mathbb{P}_x = \mathbb{P}_{x_1}^{(1)} \otimes \mathbb{P}_{x_2}^{(2)}$. We have that $\inf\{s \geq 0 : B_2(s) = a\} = T_{(-\infty, a)}$. Since

$$\mathbb{P}_{x_2}^{(2)}[T_{(-\infty, a)} > t] = \frac{2}{\pi^{1/2}} \int_0^{(a-x_2)/(4t)^{1/2}} d\theta e^{-\theta^2}, \quad \forall x_2 < a,$$

we have that the density $\rho(a, \tau)$ of the random variable $\inf\{s \geq 0 : B_2(s) = a\}$ with $x_2 = 0$, is given by

$$\rho(a, \tau) = \frac{a}{2\pi^{1/2}\tau^{3/2}} e^{-a^2/(4\tau)} \mathbf{1}_{\mathbb{R}^+}(\tau). \quad (45)$$

The following lemma will be used in the proof of Theorem 2(iv).

Lemma 3. *Let Ω be an open, bounded and connected set in \mathbb{R}^2 which contains an open rectangle $R_{a,b} \equiv R_{a,b}(p)$, $a \geq b$, with sides $K_1 = [(p, 0) - (p+b, 0)]$, $K_2 = [(p, 0) - (p, a)]$, $K_3 = [(p+b, 0) - (p+b, a)]$, $K_4 = [(p+b, a) - (p, a)]$. Let*

$$\Omega_{a,b} \equiv \Omega_{a,b}(p) = \Omega \setminus (\cup_{j=1}^3 K_j).$$

If $x \in R_{a,b}$, then

$$v_{\Omega_{a,b}}(x) \leq \frac{1}{2}(x_1 - p)(p + b - x_1) + 2^{9/2} e^{-\pi(a-x_2)/(2b)} \lambda_1(\Omega_{a,b})^{-1}. \quad (46)$$

Proof. By (43) and (44), for any $x \in R_{a,b}$,

$$\begin{aligned} v_{\Omega_{a,b}}(x) &= \int_0^\infty dt \mathbb{P}_x[T_{\Omega_{a,b}} > t] \\ &= \int_0^\infty dt \mathbb{P}_x[T_{R_{a,b}} > t] + \int_0^\infty dt \mathbb{P}_x[T_{R_{a,b}} \leq t < T_{\Omega_{a,b}}] \\ &\leq v_{R_{a,b}}(x_1, x_2) + \int_0^\infty dt \mathbb{P}_x[T_{R_{a,b}} \leq t < T_{\Omega_{a,b}}] \\ &\leq \frac{1}{2}(x_1 - p)(p + b - x_1) + \int_0^\infty dt \mathbb{P}_x[T_{R_{a,b}} \leq t < T_{\Omega_{a,b}}], \end{aligned} \quad (47)$$

where we have used that $a \mapsto v_{R_{a,b}}(x_1, x_2)$ is monotone increasing and bounded from above by $\frac{1}{2}(x_1 - p)(p + b - x_1)$. See Figure 3. The latter is the torsion




Figure 3: $\Omega_{a,b} = \Omega \setminus (\cup_{j=1}^3 K_j)$: Rectangle $R_{a,b}$ with sides K_2, K_3 of length a , and K_1, K_4 of length b .

function for the interval $(p, p+b)$. To bound the second term in the right-hand side of (47) we use the inclusion,

$$\{T_{R_{a,b}} \leq t < T_{\Omega_{a,b}}\} \subset \{B(T_{R_{a,b}}) \in K_4\} \cap \{t < T_{\Omega_{a,b}}\}.$$

This inclusion states that the event of a Brownian path starting for example at x in $R_{a,b}$ exiting the rectangle $R_{a,b}$ before t but not exiting $\Omega_{a,b}$ before t has to exit the rectangle at K_4 , while staying in $\Omega_{a,b}$ until t . By Cauchy-Schwarz,

$$\begin{aligned} \mathbb{P}_x[T_{R_{a,b}} \leq t < T_{\Omega_{a,b}}] &\leq (\mathbb{P}_x[B(T_{R_{a,b}}) \in K_4])^{1/2} (\mathbb{P}_x[t < T_{\Omega_{a,b}}])^{1/2} \\ &\leq 2^{3/4} (\mathbb{P}_x[B(T_{R_{a,b}}) \in K_4])^{1/2} e^{-t\lambda_1(\Omega_{a,b})/8}, \end{aligned} \quad (48)$$

where we have used that for open sets $\Omega \subset \mathbb{R}^2$, or for open subsets $\Omega \subset \mathbb{R}^1$,

$$\mathbb{P}_x[T_\Omega > t] \leq 2^{3/2} e^{-t\lambda_1(\Omega)/4}. \quad (49)$$

See pp.10, 11 in [6]. Let $S_{a,b} = (p, p+b) \times (-\infty, a)$ be the half-strip of width b below the line segment K_4 . Note that $R_{a,b} \subset S_{a,b}$ and that $|S_{a,b}| = \infty$. By

(48), by applying (49) to the open set $\Omega = (0, b)$, and using (45),

$$\begin{aligned}
\mathbb{P}_x[B(T_{R_{a,b}}) \in K_4] &\leq \mathbb{P}_x[B(T_{S_{a,b}}) \in K_4] \\
&= \int_0^\infty d\tau \rho(a - x_2, \tau) \mathbb{P}_{x_1}^{(1)}[T_{(0,b)} > \tau] \\
&\leq 2^{3/2} \int_0^\infty d\tau \frac{a - x_2}{2\pi^{1/2}\tau^{3/2}} e^{-(a-x_2)^2/(4\tau) - \pi^2\tau/(4b^2)} \\
&= 2^{3/2} e^{-\pi(a-x_2)/(2b)}, \tag{50}
\end{aligned}$$

and where we have used formula 3.472.3 in [18] for the last equality. By (48) and (50)

$$\mathbb{P}_x[T_{R_{a,b}} \leq t < T_{\Omega_{a,b}}] \leq 2^{3/2} e^{-\pi(a-x_2)/(4b) - t\lambda_1(\Omega_{a,b})/8}. \tag{51}$$

Integrating both sides of (51) with respect to t , and using (47) completes the proof. \square

Proof of Theorem 2(iv). To prove (19) we first obtain a lower bound on the right-hand side of (8) by making a particular choice for (A_n) :

$$A_n = \{x \in \Omega_{\varepsilon_n, n} : x_2 > 1 - cn^{-2/3}\}. \tag{52}$$

We let

$$B_n = \{x \in \Omega_{\varepsilon_n, n} : 1 - cn^{-2/3} \geq x_2 > 1 - cn^{-2/3} - n^{-3/4}\}, \tag{53}$$

and

$$C_n = \{x \in \Omega_{\varepsilon_n, n} : 1 - cn^{-2/3} - n^{-3/4} \geq x_2\}, \tag{54}$$

so that $\Omega_{\varepsilon_n, n} = A_n \cup B_n \cup C_n$. By (8), and the monotonicity of the torsion function,

$$\begin{aligned}
\kappa &\geq \limsup_{n \rightarrow \infty} \frac{\int_{A_n} v_{\Omega_{\varepsilon_n, n}}}{\int_{A_n} v_{\Omega_{\varepsilon_n, n}} + \int_{B_n} v_{\Omega_{\varepsilon_n, n}} + \int_{C_n} v_{\Omega_{\varepsilon_n, n}}} \\
&\geq \limsup_{n \rightarrow \infty} \frac{\int_{A_n} v_{A_n}}{\int_{A_n} v_{A_n} + \int_{B_n} v_{\Omega_{\varepsilon_n, n}} + \int_{C_n} v_{\Omega_{\varepsilon_n, n}}}. \tag{55}
\end{aligned}$$

Since A_n is a rectangle with side lengths 1 and $cn^{-2/3}$ respectively, we have by (32)

$$\frac{c^3}{12n^2} \geq \int_{A_n} v_{A_n} \geq \frac{c^3}{12n^2} - \frac{c^4}{24n^{8/3}}. \tag{56}$$

By (55) and (56),

$$\begin{aligned}
\kappa &\geq \limsup_{n \rightarrow \infty} \frac{c^3 - \frac{1}{2}c^4n^{-2/3}}{c^3 + 12n^2 \int_{B_n} v_{\Omega_{\varepsilon_n, n}} + 12n^2 \int_{C_n} v_{\Omega_{\varepsilon_n, n}}} \\
&= \limsup_{n \rightarrow \infty} \frac{c^3}{c^3 + 12n^2 \int_{B_n} v_{\Omega_{\varepsilon_n, n}} + 12n^2 \int_{C_n} v_{\Omega_{\varepsilon_n, n}}} \tag{57}
\end{aligned}$$

By (21) and (39),

$$\|v_{\Omega_{\varepsilon_n, n}}\|_\infty \leq 16\mathfrak{c}_2 \|d_{\Omega_{\varepsilon_n, n}}\|_\infty^2 \leq 16\mathfrak{c}_2 c^2 n^{-4/3}, \quad n \geq N_{2/3, c}. \tag{58}$$

Since $|B_n| = n^{-3/4}$ we have by (58),

$$\begin{aligned} 12n^2 \int_{B_n} v_{\Omega_{\varepsilon_n, n}} &\leq 12n^2 |B_n| \|v_{\Omega_{\varepsilon_n, n}}\|_\infty \\ &\leq 192c_2 c^2 n^{-1/12}. \end{aligned}$$

This, together with (57), yields

$$\kappa \geq \limsup_{n \rightarrow \infty} \frac{c^3}{c^3 + 12n^2 \int_{C_n} v_{\Omega_{\varepsilon_n, n}}}. \quad (59)$$

In order to bound the integral in the right-hand side from above we use Lemma 3 for each of the n rectangles $R_{a,b}(p)$ in $\Omega_{\varepsilon_n, n}$, with $p = \frac{k}{n}$, $0 \leq k \leq n-1$, with $b = \frac{1}{n}$, and $a = 1 - cn^{-2/3}$. Note that for any $p = \frac{k}{n}$ the set $\Omega_{a,b}(p)$ introduced in Lemma 3 coincides with $\Omega_{\varepsilon_n, n}$, and that each point in C_n satisfies $a - x_2 \geq n^{-3/4}$. By (20) and (39),

$$\lambda_1(\Omega_{a,b}) = \lambda_1(\Omega_{\varepsilon_n, n}) \geq \frac{1}{16 \|d_{\Omega_{\varepsilon_n, n}}\|_\infty^2} \geq \frac{1}{16c^2 n^{-4/3}}.$$

Hence the second term the right-hand side of (46) is bounded from above by $2^{17/2} c^2 n^{-4/3} e^{-\pi n^{1/4}/2}$ uniformly for all points $x \in C_n$. Since $|C_n| \leq 1$ we have

$$\int_{C_n} 2^{9/2} e^{-\pi(a-x_2)/(2b)} \lambda_1(\Omega_{a,b})^{-1} \leq 2^{17/2} c^2 n^{-4/3} e^{-\pi n^{1/4}/2}. \quad (60)$$

Integrating the first term in the right-hand side of (46) over one rectangle in C_n , and adjusting the coordinate frame appropriately, gives a contribution

$$\int_0^{1-cn^{-2/3}-n^{-3/4}} dx_2 \int_0^{1/n} dx_1 \frac{x_1}{2} (n^{-1} - x_1) = \frac{1}{12n^3} (1 - cn^{-2/3} - n^{-3/4}).$$

Summing over all n rectangles in C_n gives, together with (60) and Lemma 3,

$$12n^2 \int_{C_n} v_{\Omega_{\varepsilon_n, n}} \leq 1 - cn^{-2/3} - n^{-3/4} + 3 \cdot 2^{21/2} c^2 n^{2/3} e^{-\pi n^{1/4}/2}. \quad (61)$$

By (59) and (61) we conclude

$$\kappa \geq \frac{c^3}{1 + c^3}.$$

We now prove the converse inequality by obtaining an upper bound for the right-hand side of (8). We first observe, by the monotonicity, that the torsion for $\Omega_{\varepsilon_n, n}$ is bounded from below by the torsion for the set in Figure 2. The latter is the union of n disjoint rectangles $R_{a,b}$ with $b = \frac{1}{n}$ and $a = 1 - cn^{-2/3}$, together with one disjoint rectangle with $a = 1$ and $b = cn^{-2/3}$. By (32) we find

$$\int_{\Omega_{\varepsilon_n, n}} v_{\Omega_{\varepsilon_n, n}} \geq \frac{1 - cn^{-2/3}}{12n^2} - \frac{1}{24n^3} + \frac{c^3}{12n^2} - \frac{c^4}{24n^{8/3}}. \quad (62)$$

In order to avoid abuse of notation we keep (52), (53) and (54), and denote by (D_n) an arbitrary sequence of in $\mathfrak{A}((\Omega_{\varepsilon_n, n}))$. That is $D_n \subset \Omega_{\varepsilon_n, n}$, measurable, $n \in \mathbb{N}$, with $\lim_{n \rightarrow \infty} |D_n| = 0$. We have

$$\begin{aligned} \int_{D_n} v_{\Omega_{\varepsilon_n, n}} &= \int_{D_n \cap (A_n \cup B_n)} v_{\Omega_{\varepsilon_n, n}} + \int_{D_n \setminus (A_n \cup B_n)} v_{\Omega_{\varepsilon_n, n}} \\ &\leq \int_{A_n} v_{\Omega_{\varepsilon_n, n}} + \int_{B_n} v_{\Omega_{\varepsilon_n, n}} + \int_{D_n \cap C_n} v_{\Omega_{\varepsilon_n, n}}. \end{aligned} \quad (63)$$

By (58),

$$\int_{B_n} v_{\Omega_{\varepsilon_n, n}} \leq 16\mathbf{c}_2 c^2 n^{-25/12}. \quad (64)$$

We use Lemma 3 to bound the third term in the right-hand side of (63) from above. We have, as before, that for each point $x \in C_n$, both $a - x_2 \geq n^{-3/4}$, and $\frac{x_1}{2}(\frac{1}{n} - x_1) \leq \frac{1}{8n^2}$, where the x_1 coordinate is adjusted to the rectangle under consideration. This gives,

$$v_{\Omega_{\varepsilon_n, n}}(x) \leq \frac{1}{8n^2} + 2^{17/2} c^2 n^{-4/3} e^{-\pi n^{1/4}/2}, \quad \forall x \in C_n.$$

Hence

$$\begin{aligned} \int_{D_n \cap C_n} v_{\Omega_{\varepsilon_n, n}} &\leq |D_n \cap C_n| \left(\frac{1}{8n^2} + 2^{17/2} c^2 n^{-4/3} e^{-\pi n^{1/4}/2} \right) \\ &\leq |D_n| \left(\frac{1}{8n^2} + 2^{17/2} c^2 n^{-4/3} e^{-\pi n^{1/4}/2} \right). \end{aligned} \quad (65)$$

By (62) and (64),

$$\limsup_{n \rightarrow \infty} \frac{\int_{B_n} v_{\Omega_{\varepsilon_n, n}}}{\int_{\Omega_{\varepsilon_n, n}} v_{\Omega_{\varepsilon_n, n}}} = 0.$$

Moreover, by (62) and (65),

$$\limsup_{n \rightarrow \infty} \frac{\int_{D_n \cap C_n} v_{\Omega_{\varepsilon_n, n}}}{\int_{\Omega_{\varepsilon_n, n}} v_{\Omega_{\varepsilon_n, n}}} \leq \limsup_{n \rightarrow \infty} \frac{4|D_n|}{3(1+c^3)} = 0.$$

It remains to find an upper bound for $\int_{A_n} v_{\Omega_{\varepsilon_n, n}}$. Let $\tilde{\Omega}_n$ be the connected open set in \mathbb{R}^2 with boundary consisting of the horizontal line $\{x_2 = 1\}$ and the vertical half-lines $\{x_2 \leq 1 - cn^{-2/3}, x_1 = \frac{k}{n}, k \in \mathbb{Z}\}$. See Figure 4.

We first show that the bottom of the spectrum of the Dirichlet Laplacian on $\tilde{\Omega}_n$ is bounded away from 0. By inserting Neumann boundary conditions on the line $x_2 = 1 - cn^{-2/3}$ the bottom of the spectrum decreases and decouples into the union of the spectra of the horizontal strip at width $cn^{-2/3}$ with Dirichlet boundary conditions on $x_2 = 1$, and Neumann boundary condition on $x_2 = 1 - cn^{-2/3}$, and vertical half-strips of width n^{-1} with Neumann boundary condition at the top end and Dirichlet conditions on the half lines. By taking the double of these half-strips we obtain that the bottom of the spectrum here is the same as the bottom of the infinite strip of width n^{-1} . Combining these inequalities gives for $n \geq N_{\alpha, c}$,

$$\begin{aligned} \lambda_1(\tilde{\Omega}_n) &\geq \min\left\{ \frac{\pi^2 n^{4/3}}{4c^2}, \pi^2 n^2 \right\} \\ &= \frac{\pi^2 n^{4/3}}{c^2}. \end{aligned} \quad (66)$$




Figure 4: $\tilde{\Omega}_n$: The vertical half lines, periodically extended, are at distance $\frac{1}{n}$, and have distance $cn^{-2/3}$ to the horizontal line.

Hence the torsion function for $\tilde{\Omega}_n$ exists (see (5) and the preceding paragraph), and by the monotonicity property (4)

$$v_{\Omega_{\varepsilon_n, n}}(x) \leq v_{\tilde{\Omega}_n}(x), \quad \forall x \in \Omega_{\varepsilon_n, n}.$$

Note that

$$v_{\tilde{\Omega}_n}(x_1, x_2) = v_{\tilde{\Omega}_n}(x_1 + \frac{1}{n}, x_2), \quad \forall x \in \tilde{\Omega}_n.$$

Let

$$E_n = \{x \in \mathbb{R}^2 : 1 > x_2 > 1 - cn^{-2/3} - n^{-3/4}\}.$$

We have

$$\begin{aligned} \mathbb{P}_x[T_{\Omega_{\varepsilon_n, n}} > t] &\leq \mathbb{P}_x[T_{\tilde{\Omega}_n} > t] \\ &= \mathbb{P}_x[T_{\tilde{\Omega}_n} > t, T_{E_n} > t] + \mathbb{P}_x[T_{E_n} \leq t < T_{\tilde{\Omega}_n}] \\ &\leq \mathbb{P}_x[T_{E_n} > t] + \mathbb{P}_x[T_{E_n} \leq t < T_{\tilde{\Omega}_n}]. \end{aligned} \quad (67)$$

Integrating both sides of (67) with respect to t over \mathbb{R}^+ gives

$$\begin{aligned} v_{\Omega_{\varepsilon_n, n}}(x) &\leq v_{E_n}(x) + \int_0^\infty dt \mathbb{P}_x[T_{E_n} \leq t < T_{\tilde{\Omega}_n}] \\ &= \frac{1}{2}(1 - x_2)(x_2 - 1 + d_n) + \int_0^\infty dt \mathbb{P}_x[T_{E_n} \leq t < T_{\tilde{\Omega}_n}], \end{aligned}$$

where

$$d_n = cn^{-2/3} + n^{-3/4}. \quad (68)$$

Hence

$$\begin{aligned} \int_{A_n} v_{\Omega_{\varepsilon_n, n}} &\leq \int_{A_n} dx \frac{1}{2}(1-x_2)(x_2-1+d_n) + \int_{A_n} dx \int_0^\infty dt \mathbb{P}_x[T_{E_n} \leq t < T_{\tilde{\Omega}_n}] \\ &\leq \frac{1}{12}d_n^3 + \int_{A_n} dx \int_0^\infty dt \mathbb{P}_x[T_{E_n} \leq t < T_{\tilde{\Omega}_n}]. \end{aligned}$$

By (62), (68) we have

$$\limsup_{n \rightarrow \infty} \frac{\frac{1}{12}d_n^3}{\int_{\Omega_{\varepsilon_n, n}} v_{\Omega_{\varepsilon_n, n}}} \leq \frac{c^3}{1+c^3}.$$

To complete the proof it therefore suffices to show that

$$\limsup_{n \rightarrow \infty} n^2 \int_{A_n} dx \int_0^\infty dt \mathbb{P}_x[T_{E_n} \leq t < T_{\tilde{\Omega}_n}] = 0. \quad (69)$$

To prove (69) we let $L_n = \{(x_1, 1-d_n) : x_1 \in \mathbb{R}\}$. We have by the strong Markov property,

$$\begin{aligned} \mathbb{P}_x[T_{E_n} \leq t < T_{\tilde{\Omega}_n}] &\leq \mathbb{E}_x \left(\int_0^t \mathbf{1}_{\tau_{L_n} \in d\tau} \mathbb{P}_{B(\tau_{L_n})}[T_{\tilde{\Omega}_n} > t - \tau] \right) \\ &\leq \mathbb{E}_x \left(\int_0^t \mathbf{1}_{\tau_{L_n} \in d\tau} \sup_{z \in L_n} \mathbb{P}_z[T_{\tilde{\Omega}_n} > t - \tau] \right) \\ &= \int_0^t d\tau \frac{\partial}{\partial \tau} (\mathbb{P}_x[\tau_{L_n} < \tau]) \sup_{z \in K_n} \mathbb{P}_z[T_{\tilde{\Omega}_n} > t - \tau] \\ &= \int_0^t d\tau \frac{\partial}{\partial \tau} (\mathbb{P}_x[\tau_{L_n} < \tau]) \mathbb{P}_{((2n)^{-1}, 1-d_n)}[T_{\tilde{\Omega}_n} > t - \tau], \quad (70) \end{aligned}$$

where $\mathbf{1}_{\tau_{L_n} \in d\tau}$ is the indicator function on the set of Brownian paths which hit L_n in the infinitesimal interval $d\tau$. We have used in the final equality in (70) that $x_1 \mapsto \mathbb{P}_{(x_1, 1-d_n)}[T_{\tilde{\Omega}_n} > t - \tau]$, $x_1 \in \mathbb{R}$ is periodic in x_1 with period n^{-1} , and which has equal maxima at $\{((2n)^{-1} + kn^{-1}, 1-d_n) : k \in \mathbb{Z}\}$. Hence the supremum in the third line in (70) is a maximum for $z = ((2n)^{-1}, 1-d_n)$. Integrating the convolution in the right-hand side of (70) with respect to t over \mathbb{R}^+ gives

$$\begin{aligned} \int_0^\infty dt \mathbb{P}_x[T_{E_n} \leq t < T_{\tilde{\Omega}_n}] &\leq \left(\int_0^\infty d\tau \frac{\partial}{\partial \tau} (\mathbb{P}_x[\tau_{L_n} < \tau]) \right) \int_0^\infty d\tau \mathbb{P}_{((2n)^{-1}, 1-d_n)}[T_{\tilde{\Omega}_n} > \tau] \\ &= v_{\tilde{\Omega}_n}((2n)^{-1}, 1-d_n). \end{aligned}$$

One verifies that Lemma 3 also holds if Ω contains the open rectangle $(p, p+b) \times (-\infty, a)$. Indeed the first term in the right-hand side of (46) is the torsion function for the infinite strip of width b . The second term takes into account that the strip is cut-off at a . With this we arrive at

$$\int_0^t dt \mathbb{P}_x[T_{E_n} \leq t < T_{\tilde{\Omega}_n}] \leq \frac{1}{8n^2} + 2^{9/2} e^{-\pi n^{1/4}/2} \lambda_1(\tilde{\Omega}_n)^{-1}. \quad (71)$$

Since $|A_n| = cn^{-2/3}$, we find by (71), and (66),

$$n^2 \int_{A_n} dx \int_0^t dt \mathbb{P}_x[T_{E_n} \leq t < T_{\Omega_n}] \leq \frac{c}{8n^{2/3}} + 2^{9/2} \pi^{-2} c^3 e^{-\pi n^{1/4}/2}.$$

This implies (69). □

Acknowledgments. Both authors acknowledge support by the Leverhulme Trust through Emeritus Fellowship EM-2018-011-9, and the Swiss National Science Foundation respectively.

References

- [1] A. Ancona, On strong barriers and an inequality of Hardy for domains in \mathbb{R}^n . *J. London Math. Soc.* **34** (1986), 274–290.
- [2] D. N. Arnold, G. David, M. Filoche, D. Jerison, S. Mayboroda, Computing spectra without solving eigenvalue problems. *SIAM J. Sci. Comput.* **41** (2019), B69–B92.
- [3] D. N. Arnold, G. David, M. Filoche, D. Jerison, S. Mayboroda, Localization of eigenfunctions via an effective potential. *Comm. Partial Differential Equations* **44** (2019), 1186–1216.
- [4] C. Bandle, *Isoperimetric Inequalities and Applications*, Monographs and Studies in Mathematics, Pitman, London, 1980.
- [5] M. van den Berg, Estimates for the torsion function and Sobolev constants. *Potential Analysis* **36** (2012), 607–616.
- [6] M. van den Berg, D. Bucur, T. Kappeler, On efficiency and localisation for the torsion function. *Potential Analysis*, to appear, arXiv:2005.06366.
- [7] M. van den Berg, T. Carroll, Hardy inequality and L^p estimates for the torsion function. *Bull. Lond. Math. Soc.* **41** (2009), 980–986.
- [8] M. van den Berg, F. Della Pietra, G. di Blasio, N. Gavitone, Efficiency and localisation for the first Dirichlet eigenfunction. *Journal of Spectral Theory*, to appear, arXiv:1905.06591.
- [9] L. Brasco, On torsional rigidity and principal frequencies: an invitation to the Kohler–Jobin rearrangement technique. *ESAIM: Control, Optimisation and Calculus of Variations* **20** (2014), 315–338.
- [10] D. Bucur, G. Buttazzo, *Variational Methods in Shape Optimization Problems*, Progress in Nonlinear Differential Equations and their Applications 65, Birkhäuser Boston, Inc., Boston, MA, 2005.
- [11] E. B. Davies *Heat Kernels and spectral theory*, Cambridge Tracts in Mathematics, **92**. Cambridge University Press, Cambridge, 1989.
- [12] E. B. Davies, The Hardy constant. *Quart. J. Math. Oxford Ser. (2)* **46** (1995), 417–431.

- [13] E. B. Davies, A review of Hardy inequalities. *Operator Theory Adv. Appl.* **110** (1999), 55–67.
- [14] E. B. Davies, Sharp boundary estimates for elliptic operators. *Math. Proc. Cambridge Philos. Soc.* **129** (2000), 165–178.
- [15] F. Della Pietra, N. Gavitone, S. Guarino Lo Bianco, On functionals involving the torsional rigidity related to some classes of nonlinear operators. *J. Differential Equations* **265** (2018), 6424–6442.
- [16] R. Durrett, *Brownian motion and martingales in analysis*, Wadsworth Mathematics Series, Belmont, CA, 1984.
- [17] T. Giorgi, R. G. Smits, Principal eigenvalue estimates via the supremum of torsion. *Indiana Univ. Math. J.* **59** (2010), 987–1011.
- [18] I. S. Gradshteym, I. M. Ryzhik, *Table of Integrals, Series, and Products*, 7th Edition, Academic Press, New York, 2007.
- [19] D. S. Grebenkov, B.-T. Nguyen, Geometrical structure of Laplacian eigenfunctions. *SIAM Rev.* **55** (2013), 601–667.
- [20] S. Markvorsen, V. Palmer, Torsional rigidity of minimal submanifolds. *Proc. London Math. Soc.* **93** (2006), 253–272.
- [21] L. E. Payne, I. Stakgold, On the mean value of the fundamental mode in the fixed membrane problem. Collection of articles dedicated to Alexander Weinstein on the occasion of his 75th birthday. *Applicable Anal.* **3** (1973), 295–306.
- [22] G. Pólya, G. Szegő, *Isoperimetric Inequalities in Mathematical Physics*, Ann. of Math. Stud. 27, Princeton University Press, Princeton, 1951.
- [23] R. Sperb, *Maximum principles and applications*, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1981.
- [24] H. Vogt, L_∞ - estimates for the torsion function and L_∞ - growth of semi-groups satisfying Gaussian bounds. *Potential Anal.* **51** (2019), 37–47.