

Snowmass2021 - Letter of Interest

New opportunities at the photon energy frontier

Coordinators of this LoI: Spencer Klein (LBNL)¹ and Daniel Tapia Takaki (U. Kansas)²

Authors: Jaroslav Adam⁹, Christine Aidala⁴⁰, Aaron Angerami³, Benjamin Audurier⁴⁷, Carlos Bertulani¹⁷, Christian Bierlich²⁴, Boris Blok³⁵, James Daniel Brandenburg⁹, Stanley Brodsky³⁴, Aleksandr Bylinkin², Veronica Canoa Roman⁴², Francesco Giovanni Celiberto⁵², Jan Cepila⁰, Grigorios Chachamis⁴⁶, Brian Cole²², Guillermo Contreras⁰, David d'Enterria¹⁴, Adrian Dumitru²⁸, Arturo Fernández Téllez²⁰, Leonid Frankfurt^{10,50}, Maria Beatriz Gay Ducati¹⁹, Frank Geurts²³, Gustavo Gil da Silveira¹¹, Francesco Giuli²⁶, Victor P. Goncalves¹⁶, Iwona Grabowska-Bold⁵, Vadim Guzey¹², Lucian Harland-Lang³², Martin Hentschinski²⁹, T. J. Hobbs²⁵, Jamal Jalilian-Marian²⁸, Valery A. Khoze¹⁵, Yongsun Kim³⁶, Spencer R. Klein¹, Simon Knapen²¹, Mariola Klusek-Gawenda⁴⁸, Michal Krelina⁰, Evgeny Kryshen¹², Tuomas Lappi³⁸, Constantin Loizides⁷, Agnieszka Luszczak⁴⁴, Magno Machado³⁹, Heikki Mäntysaari³⁸, Daniel Martins⁷, Ronan McNulty⁴⁵, Michael Murray², Jan Nemchik⁰, Jacquelyn Noronha-Hostler³³, Joakim Nystrand⁶, Alessandro Papa⁵¹, Bernard Pire³⁷, Mateusz Ploskon¹, Marius Przybycien⁵, John P. Ralston², Patricia Rebello Teles¹⁸, Christophe Royon², Björn Schenke⁹, William Schmidke⁹, Janet Seger⁸, Anna Stasto¹⁰, Peter Steinberg⁹, Mark Strikman¹⁰, Antoni Szczurek⁴⁸, Lech Szymanowski³¹, Daniel Tapia Takaki², Ralf Ulrich⁴⁹, Orlando Villalobos Baillie⁴¹, Ramona Vogt^{3,4}, Samuel Wallon³⁰, Michael Winn⁴³, Keping Xie²⁷, Zhangbu Xu⁹, Shuai Yang²³, Mikhail Zhilov¹², and Jian Zhou¹³

(0) Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic;

(1) Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley CA;

(2) Department of Physics and Astronomy, The University of Kansas, Lawrence KS;

(3) Lawrence Livermore National Laboratory, Livermore CA;

(4) Department of Physics, University of California, Davis CA;

(5) AGH University of Science and Technology, Cracow, Poland;

(6) Dept. of Physics and Technology, University of Bergen, Bergen, Norway;

(7) Oak Ridge National Laboratory, Oak Ridge, TN;

(8) Physics Department, Creighton University, Omaha, NE;

(9) Brookhaven National Laboratory, Upton, NY;

(10) Dept. of Physics, Pennsylvania State

University, University Park, PA;

(11) UFRGS and UERJ, Brazil;

(12) Petersburg Nuclear Physics Institute of National Research Centre "Kurchatov Institute;

(13) Shandong University(Qingdao), China;

(14) CERN, Geneva, Switzerland;

(15) IPPP, Durham University, Durham, UK;

(16) Universidade Federal de Pelotas, Pelotas, Brazil;

(17) Department of Physics and Astronomy, Texas A&M University-Commerce, Commerce, TX;

(18) Universidade Federal do Rio de Janeiro, UFRJ, Rio de Janeiro, Brazil;

(19) UFRGS (Universidade Federal do Rio Grande do Sul), Brazil

(20) Benemérita Universidad Autónoma de Puebla, Puebla, México;

¹srklein@lbl.gov

²Daniel.Tapia.Takaki@cern.ch

- (21) Physics Division, Lawrence Berkeley National Laboratory, Berkeley CA;
- (22) Dept. of Physics, Columbia University, New York, NY, 10027;
- (23) Department of Physics and Astronomy, Rice University, Houston, TX, 77005;
- (24) Department of Astronomy and Theoretical Physics, Lund University, Lund, Sweden;
- (25) Department of Physics, Southern Methodist University and JLab EIC Center;
- (26) University and INFN Rome 2, Via della Ricerca Scientifica, 1 – 00133 Roma;
- (27) PITT PACC, University of Pittsburgh, Pittsburgh, PA 15260;
- (28) Baruch College, CUNY, New York, NY 10010;
- (29) Departamento de Actuaría, Física y Matemáticas, Universidad de las Américas Puebla, Ex-Hacienda Santa Catarina Martir S/N, San Andrés Cholula 72820 Puebla, Mexico;
- (30) Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France;
- (31) National Centre for Nuclear Research, Warsaw, Poland;
- (32) Department of Physics, University of Oxford, Oxford, UK;
- (33) Illinois Center for Advanced Studies of the Universe, Department of Physics, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA;
- (34) SLAC National Accelerator Laboratory, Stanford University, Stanford, CA;
- (35) Physics Department, Technion, Haifa, Israel;
- (36) Sejong University, Seoul, South Korea;
- (37) CPHT, CNRS, Ecole polytechnique, I.P. Paris, 91128 Palaiseau, France;
- (38) Department of Physics, University of Jyväskylä, P.O. Box 35, 40014 University of Jyväskylä, Finland and Helsinki Institute of Physics, P.O. Box 64, 00014 University of Helsinki, Finland;
- (39) Institute of Physics - UFRGS, Brazil;
- (40) Physics Department, University of Michigan, Ann Arbor, MI 48109;
- (41) School of Physics and Astronomy, The University of Birmingham, Edgbaston, Birmingham, UK. B15 2TT;
- (42) Department of Physics and Astronomy, Stony Brook University, Stony Brook, NY, 11794;
- (43) Université Paris-Saclay Centre d'Etudes de Saclay (CEA), IRFU, Département de Physique Nucléaire(DPhN), Saclay, France;
- (44) Cracow University of Technology, Cracow, Poland;
- (45) University College Dublin, Dublin, Ireland;
- (46) LIP, Av. Prof. Gama Pinto 2, P-1649-003 Lisboa, Portugal;
- (47) Laboratoire Leprince-Ringuet, Palaiseau, France;
- (48) Institute of Nuclear Physics Polish Academy of Sciences, PL-31342 Krakow, Poland;
- (49) KIT, Karlsruhe Institute of Technology, 76131 Karlsruhe, Germany;
- (50) Sackler School of Exact Sciences, Tel Aviv University, Tel Aviv, 69978, Israel
- (51) Università della Calabria and INFN-Cosenza, Italy
- (52) Università degli Studi di Pavia and INFN, Sezione di Pavia, Italy

Abstract: Ultra-peripheral collisions (UPCs) involving heavy ions and protons are the energy frontier for photon-mediated interactions. UPC photons can be used for many purposes, including probing low- x gluons via photoproduction of dijets and vector mesons, probes of beyond-standard-model processes, such as those enabled by light-by-light scattering, and studies of two-photon production of the Higgs.

UPCs as the energy frontier Since the closure of the HERA ep collider, there have been no dedicated high-energy photon facilities. Instead, the photon energy frontier has been studied in UPCs at CERN’s Large Hadron Collider^{1–7}. The protons and heavy ions accelerated there carry electromagnetic fields which may be treated as a flux of nearly-real (virtuality $Q^2 < (\hbar c/R_A)^2$, where R_A is the hadron radius) photons. The photon spectra extend up to energies of $\gamma\hbar c/R_A$, where γ is the Lorentz boost of the ion. At the LHC, these photons lead to γp collisions at center of mass energies up to 5 TeV, γA collisions at center of mass energies up to 700 GeV/nucleon, and two-photon collisions up to $\sqrt{s_{\gamma\gamma}} = 4.2$ TeV. The γp energies are higher than were accessible at HERA, and the γA energies are many orders of magnitude higher than are accessible at fixed-target experiments. These photons have been used to study a wide variety of physics processes: measurements of low- x gluon densities in protons and studies of shadowing of parton densities in heavier nuclei, studies of higher order terms in dilepton production, and light-by-light scattering.

Photoproduction and parton distributions Parton distributions have been probed in γp and γA collisions. Photoproduction of dijets (and, still to come, open charm⁸) is, from the theoretical point of view, a fairly direct probe of the gluon distribution. So far, preliminary results from ATLAS on dijet production⁹ and from CMS on exclusive dijet photoproduction¹⁰ have been released. This can be used to measure the diffractive structure functions and study “elliptic gluon” dynamics^{11–17}. UPCs at the LHC can probe to Bjorken- x values of at least a few 10^{-6} ; this could reach even lower x with far-forward detectors like the proposed ALICE FoCal¹⁸.

Exclusive vector mesons are produced when an incident photon fluctuates to a virtual quark-antiquark pair, which then scatters elastically (or quasi-elastically) from a proton or nuclear target. Since elastic scattering requires two-gluon exchange for color neutrality, the cross section scales as the square of the gluon density. One limitation is that the color neutrality requirement introduces some systematic uncertainty^{19;20}. ALICE²¹ and LHCb have studied J/ψ production on proton targets, finding, for the most part, that the power-law behavior seen at HERA extends to higher energies. ALICE, LHCb and CMS have also studied $\psi(2S)$ ^{22;23} and Υ production^{24–26}, finding good agreement with NLO-inspired cross section calculations. Some calculations indicate that the saturation scale has been reached²⁷.

The nuclear-target cross sections are sensitive to gluon shadowing, and probe phenomena like gluon saturation of the color-glass condensate. ALICE^{28;29} and CMS³⁰ have studied J/ψ production on lead, and found moderate suppression, consistent with leading-twist calculations^{31–33}. There have also been studies on ρ photoproduction^{34–36}. The ρ cross section is about 40% smaller than predicted by the Glauber model, pointing to the importance of high-mass internal states, as expected in the Glauber-Gribov model³⁷.

Looking ahead, the LHC and HL-LHC will collect much more data, which should lead to precise measurements of shadowing (the anticipated error bars are shown in Fig. 1) and enable new types of measurements. In the Good-Walker paradigm³⁸, coherent and incoherent photoproduction allow access to qualitatively new studies^{32;39}. The coherent cross section $d\sigma/dt$ encodes the transverse spatial distribution of the targets - the nuclear equivalent of a Generalized Parton Distribution - while the incoherent $d\sigma/dt$ provides information on event-by-event fluctuations in the nuclear configuration, due to varying nucleon positions and gluonic hot spots^{39–42}. STAR has used $d\sigma/dt$ for ρ and direct $\pi^+\pi^-$ photoproduction to measure the spatial distribution of target scatters in gold⁴³. HL-LHC can do this measurement with heavier quarkonia, where pQCD is clearly applicable. CMS data on ρ ³⁴ shows sensitivity to the onset of gluon saturation in lead using differential studies in both momentum transfer (t) and γp center-of-mass energy⁴⁴. Other opportunities include study of perturbative Pomeron dynamics⁴⁵, color fluctuations in the photon^{46;47}, the gluonic Sivers function⁴⁸, search for the Odderon⁴⁹, among other studies. Finally, next-to-leading (NLO) order calculations for these processes are one of the future directions of the theoretical program^{50–52}.

Light-by-light scattering, W pair and dilepton production. Two-photon interactions at the LHCb probe the energy frontier. Photons couple to all electrically charged particles and some neutrals scalars (like




Figure 1: (left) Pseudodata projections for the nuclear suppression factors of the nuclear gluon density as a function of Bjorken- x for photoproduction of three vector mesons in PbPb UPCs. The points are EPS09-projections using the method described in³¹. From Ref.⁵³. (right) Exclusion limits on ALP-photon coupling ($1/\Lambda_a$) vs. ALP mass, from light-by-light scattering and other processes. From Ref.⁵⁴.

axions and the Higgs), so two-photon reactions are sensitive to many beyond-standard-model processes.

One process that has already been used to probe BSM physics is light-by-light scattering, $AA \rightarrow AA\gamma\gamma$ ^{55;56}. The subprocess $\gamma\gamma \rightarrow \gamma\gamma$ proceeds only via a charged-particle box diagram. The cross section is sensitive to all charged particles, including BSM particles^{57;58} such as vector fermions, GeV- mass axion-like particles (ALPs)^{59;60} and magnetic monopoles. The reaction also probes non-linear (BSM) corrections to electromagnetism^{61;62}. ATLAS^{54;63;64} and CMS⁶⁵ have both observed this process at a cross section consistent with the standard model. They then set limits on ALP production, as shown in Fig. 1.

In pp collisions, ATLAS⁶⁶ and CMS⁶⁷ have also studied $\gamma\gamma \rightarrow W^+W^-$ and CMS/TOTEM $\gamma\gamma \rightarrow \gamma\gamma$ ⁶⁸, thereby placing stringent limits on anomalous quartic gauge couplings. Production of electron and muon pairs via $\gamma\gamma \rightarrow l^+l^-$, has been studied by many collaborations. Exclusive $e - \mu$ events, as expected from tau pairs⁶⁹ have been seen. Tau pairs are of particular interest because the τ - γ coupling is sensitive to BSM physics, including the τ dipole moment, lepton compositeness or supersymmetry.

Looking ahead, $\gamma\gamma \rightarrow \gamma\gamma$ is a rare process and future HL-LHC running should lead to much larger data samples for this final states⁵³. ALICE and LHCb may be able to probe this reaction at lower diphoton masses than is currently possible⁷⁰. High statistics studies of dilepton production may also be sensitive to BSM processes, particularly at high masses. This is especially true for the τ , where the cross section⁷¹ and kinematic distributions⁷² are sensitive to new physics. It may also be possible to study two-photon production of heavy quark states, $\gamma\gamma \rightarrow p\bar{p}$ ⁷³, and search for pentaquarks⁷⁴, tetraquarks⁷⁵ and other exotica.

Strong fields, Quantum correlations and quantum tomography UPCs also offer opportunities to use the very strong fields to explore reactions involving multiple photon exchange, such as production of $\rho^0\rho^0$, where each ρ^0 is produced by a single, independent photon⁷⁶⁻⁷⁸. These pairs should exhibit quantum correlations which will allow for a more detailed study of decay dynamics, including with polarized photons. EPR (Einstein-Podolsky-Rosen)-type experiments can also be carried out to test quantum mechanics, and quantum tomography techniques can probe quantum correlations and entanglement^{79;80}.

UPCs at the FCC and synergies with future colliders Photon exchange at the future FCC, or at the proposed LHeC⁸¹, will allow probes at even higher energies, allowing for more extensive BSM physics and also the study of top photoproduction^{82;83} and $\gamma\gamma$ production of the Higgs⁸⁴. The future electron-ion collider⁸⁵, planned to be built at BNL around 2030, will make precise measurements of photon-mediated reactions over a wide Q^2 range, but at lower energies.

References

- [1] G. Baur, K. Hencken, D. Trautmann, S. Sadovsky, and Y. Kharlov, “Coherent gamma gamma and gamma-A interactions in very peripheral collisions at relativistic ion colliders,” *Phys. Rept.* **364** (2002) 359–450, [arXiv:hep-ph/0112211 \[hep-ph\]](#).
- [2] M. Greiner, M. Vidovic, and G. Soff, “Electromagnetic production of Higgs bosons, SUSY particles, glueballs and mesons in ultrarelativistic heavy ion collisions,” *Phys. Rev.* **C47** (1993) 2288–2298.
- [3] C. A. Bertulani, S. R. Klein, and J. Nystrand, “Physics of ultra-peripheral nuclear collisions,” *Ann. Rev. Nucl. Part. Sci.* **55** (2005) 271–310, [arXiv:nucl-ex/0502005 \[nucl-ex\]](#).
- [4] A. Baltz, “The Physics of Ultraperipheral Collisions at the LHC,” *Phys. Rept.* **458** (2008) 1–171, [arXiv:0706.3356 \[nucl-ex\]](#).
- [5] J. G. Contreras and J. D. Tapia Takaki, “Ultra-peripheral heavy-ion collisions at the LHC,” *Int. J. Mod. Phys.* **A30** (2015) 1542012.
- [6] S. Klein and J. Nystrand, “Ultraperipheral nuclear collisions,” *Phys. Today* **70** no. 10, (2017) 40–47.
- [7] S. Klein and P. Steinberg, “Photonuclear and Two-photon Interactions at High-Energy Nuclear Colliders,” [arXiv:2005.01872 \[nucl-ex\]](#).
- [8] S. R. Klein, J. Nystrand, and R. Vogt, “Heavy quark photoproduction in ultraperipheral heavy ion collisions,” *Phys. Rev. C* **66** (2002) 044906, [arXiv:hep-ph/0206220](#).
- [9] ATLAS Collaboration, ATLAS Collaboration, “Photo-nuclear dijet production in ultra-peripheral Pb+Pb collisions.” ATLAS-CONF-2017-011, 2017.
- [10] CMS Collaboration, “Angular correlations in exclusive dijet photoproduction in ultra-peripheral PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV CMS-PAS-HIN-18-011,”.
- [11] V. Guzey and M. Klasen, “Diffractive dijet photoproduction in ultraperipheral collisions at the LHC in next-to-leading order QCD,” *JHEP* **04** (2016) 158, [arXiv:1603.06055 \[hep-ph\]](#).
- [12] P. Kotko, K. Kutak, S. Sapeta, A. Stasto, and M. Strikman, “Estimating nonlinear effects in forward dijet production in ultra-peripheral heavy ion collisions at the LHC,” *Eur. Phys. J. C* **77** no. 5, (2017) 353, [arXiv:1702.03063 \[hep-ph\]](#).
- [13] T. Altinluk, N. Armesto, G. Beuf, and A. H. Rezaeian, “Diffractive Dijet Production in Deep Inelastic Scattering and Photon-Hadron Collisions in the Color Glass Condensate,” *Phys. Lett. B* **758** (2016) 373–383, [arXiv:1511.07452 \[hep-ph\]](#).
- [14] Y. Hagiwara, Y. Hatta, R. Pasechnik, M. Tasevsky, and O. Teryaev, “Accessing the gluon Wigner distribution in ultraperipheral pA collisions,” *Phys. Rev. D* **96** no. 3, (2017) 034009, [arXiv:1706.01765 \[hep-ph\]](#).
- [15] H. Mäntysaari, N. Mueller, and B. Schenke, “Diffractive Dijet Production and Wigner Distributions from the Color Glass Condensate,” *Phys. Rev. D* **99** no. 7, (2019) 074004, [arXiv:1902.05087 \[hep-ph\]](#).
- [16] Y. Hatta, B.-W. Xiao, and F. Yuan, “Probing the Small- x Gluon Tomography in Correlated Hard Diffractive Dijet Production in Deep Inelastic Scattering,” *Phys. Rev. Lett.* **116** no. 20, (2016) 202301, [arXiv:1601.01585 \[hep-ph\]](#).

- [17] J. Zhou, “Elliptic gluon generalized transverse-momentum-dependent distribution inside a large nucleus,” *Phys. Rev. D* **94** no. 11, (2016) 114017, [arXiv:1611.02397 \[hep-ph\]](#).
- [18] **ALICE FoCal** Collaboration, N. van der Kolk, “FoCal: A highly granular digital calorimeter,” *Nucl. Instrum. Meth. A* **958** (2020) 162059.
- [19] C. A. Flett, S. P. Jones, A. D. Martin, M. G. Ryskin, and T. Teubner, “Exclusive production of heavy quarkonia as a probe of the low x and low scale gluon PDF,” [arXiv:1912.09128 \[hep-ph\]](#).
- [20] C. A. Flett, S. P. Jones, A. D. Martin, M. G. Ryskin, and T. Teubner, “How to include exclusive J/ψ production data in global PDF analyses,” [arXiv:1908.08398 \[hep-ph\]](#).
- [21] **ALICE** Collaboration, S. Acharya *et al.*, “Energy dependence of exclusive J/ψ photoproduction off protons in ultra-peripheral p–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Eur. Phys. J.* **C79** no. 5, (2019) 402, [arXiv:1809.03235 \[nucl-ex\]](#).
- [22] **ALICE** Collaboration, “Coherent $\psi(2S)$ photo-production in ultra-peripheral Pb Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett.* **B751** (2015) 358–370, [arXiv:1508.05076 \[nucl-ex\]](#).
- [23] **LHCb** Collaboration, LHCb Collaboration, “Central exclusive production of J/ψ and $\psi(2S)$ mesons in pp collisions at $\sqrt{s} = 13$ TeV.” LHCb-CONF-2016-007, 2016.
- [24] **LHCb** Collaboration, R. Aaij *et al.*, “Measurement of the exclusive Y production cross-section in pp collisions at $\sqrt{s} = 7$ TeV and 8 TeV,” *JHEP* **09** (2015) 084, [arXiv:1505.08139 \[hep-ex\]](#).
- [25] R. McNulty, “Central Exclusive Production at LHCb,” *PoS DIS2016* (2016) 181, [arXiv:1608.08103 \[hep-ex\]](#).
- [26] **CMS** Collaboration, A. M. Sirunyan *et al.*, “Measurement of exclusive Υ photoproduction from protons in pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Eur. Phys. J.* **C79** no. 3, (2019) 277, [arXiv:1809.11080 \[hep-ex\]](#).
- [27] A. Arroyo Garcia, M. Hentschinski, and K. Kutak, “QCD evolution based evidence for the onset of gluon saturation in exclusive photo-production of vector mesons,” *Phys. Lett. B* **795** (2019) 569–575, [arXiv:1904.04394 \[hep-ph\]](#).
- [28] **ALICE** Collaboration, E. Abbas *et al.*, “Charmonium and e^+e^- pair photoproduction at mid-rapidity in ultra-peripheral Pb-Pb collisions at $\sqrt{s_{NN}}=2.76$ TeV,” *Eur. Phys. J.* **C73** no. 11, (2013) 2617, [arXiv:1305.1467 \[nucl-ex\]](#).
- [29] **ALICE** Collaboration, S. Acharya *et al.*, “Coherent J/ψ photoproduction at forward rapidity in ultra-peripheral Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett.* **B798** (2019) 134926, [arXiv:1904.06272 \[nucl-ex\]](#).
- [30] **CMS** Collaboration, V. Khachatryan *et al.*, “Coherent J/ψ photoproduction in ultra-peripheral PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the CMS experiment,” *Phys. Lett.* **B772** (2017) 489–511, [arXiv:1605.06966 \[nucl-ex\]](#).
- [31] V. Guzey, E. Kryshen, M. Strikman, and M. Zhalov, “Evidence for nuclear gluon shadowing from the ALICE measurements of PbPb ultraperipheral exclusive J/ψ production,” *Phys. Lett.* **B726** (2013) 290–295, [arXiv:1305.1724 \[hep-ph\]](#).
- [32] S. R. Klein and H. Mäntysaari, “Imaging the nucleus with high-energy photons,” *Nature Rev. Phys.* **1** no. 11, (2019) 662–674, [arXiv:1910.10858 \[hep-ex\]](#).

- [33] V. Guzey, E. Kryshen, M. Strikman, and M. Zhalov, “Nuclear suppression from coherent J/ψ photoproduction at the Large Hadron Collider,” [arXiv:2008.10891 \[hep-ph\]](#).
- [34] **CMS** Collaboration, A. M. Sirunyan *et al.*, “Measurement of exclusive $\rho(770)^0$ photoproduction in ultraperipheral pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Eur. Phys. J. C* **79** no. 8, (2019) 702, [arXiv:1902.01339 \[hep-ex\]](#).
- [35] **STAR** Collaboration, B. I. Abelev *et al.*, “ ρ^0 photoproduction in ultraperipheral relativistic heavy ion collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **77** (2008) 034910, [arXiv:0712.3320 \[nucl-ex\]](#).
- [36] **ALICE** Collaboration, J. Adam *et al.*, “Coherent ρ^0 photoproduction in ultra-peripheral Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **09** (2015) 095, [arXiv:1503.09177 \[nucl-ex\]](#).
- [37] L. Frankfurt, V. Guzey, M. Strikman, and M. Zhalov, “Nuclear shadowing in photoproduction of ρ^0 mesons in ultraperipheral nucleus collisions at RHIC and the LHC,” *Phys. Lett. B* **752** (2016) 51–58, [arXiv:1506.07150 \[hep-ph\]](#).
- [38] M. L. Good and W. D. Walker, “Diffraction dissociation of beam particles,” *Phys. Rev.* **120** (1960) 1857–1860.
- [39] H. Mantysaari and B. Schenke, “Revealing proton shape fluctuations with incoherent diffraction at high energy,” *Phys. Rev. D* **94** no. 3, (2016) 034042, [arXiv:1607.01711 \[hep-ph\]](#).
- [40] V. Guzey, M. Strikman, and M. Zhalov, “Accessing transverse nucleon and gluon distributions in heavy nuclei using coherent vector meson photoproduction at high energies in ion ultraperipheral collisions,” *Phys. Rev. C* **95** no. 2, (2017) 025204, [arXiv:1611.05471 \[hep-ph\]](#).
- [41] J. Cepila, J. Contreras, and J. D. Tapia Takaki, “Energy dependence of dissociative J/ψ photoproduction as a signature of gluon saturation at the LHC,” *Phys. Lett. B* **766** (2017) 186–191, [arXiv:1608.07559 \[hep-ph\]](#).
- [42] J. Cepila, J. Contreras, M. Krelina, and J. Tapia Takaki, “Mass dependence of vector meson photoproduction off protons and nuclei within the energy-dependent hot-spot model,” *Nucl. Phys. B* **934** (2018) 330–340, [arXiv:1804.05508 \[hep-ph\]](#).
- [43] **STAR** Collaboration, L. Adamczyk *et al.*, “Coherent diffractive photoproduction of ρ^0 mesons on gold nuclei at 200 GeV/nucleon-pair at the Relativistic Heavy Ion Collider,” *Phys. Rev. C* **96** no. 5, (2017) 054904, [arXiv:1702.07705 \[nucl-ex\]](#).
- [44] V. Gonçalves, F. Navarra, and D. Spiering, “Exclusive ρ and J/Ψ photoproduction in ultraperipheral pA collisions: Predictions of the gluon saturation models for the momentum transfer distributions,” *Phys. Lett. B* **791** (2019) 299–304, [arXiv:1811.09124 \[hep-ph\]](#).
- [45] B. Blok, L. Frankfurt, and M. Strikman, “DGLAP versus perturbative Pomeron in large momentum transfer hard diffractive processes at HERA and LHC,” *Phys. Lett. B* **690** (2010) 159–163, [arXiv:1002.3048 \[hep-ph\]](#).
- [46] M. Alvioli, L. Frankfurt, V. Guzey, M. Strikman, and M. Zhalov, “Mapping color fluctuations in the photon in ultraperipheral heavy ion collisions at the Large Hadron Collider,” *Phys. Lett. B* **767** (2017) 450–457, [arXiv:1605.06606 \[hep-ph\]](#).

- [47] L. Frankfurt, M. Strikman, and M. Zhalov, “Tracking fast small color dipoles through strong gluon fields at the LHC,” *Phys. Rev. Lett.* **102** (2009) 232001, [arXiv:0811.0368](#) [hep-ph].
- [48] R. Boussarie, Y. Hatta, L. Szymanowski, and S. Wallon, “Probing the Gluon Sivers Function with an Unpolarized Target: GTMD Distributions and the Odderons,” *Phys. Rev. Lett.* **124** no. 17, (2020) 172501, [arXiv:1912.08182](#) [hep-ph].
- [49] L. Harland-Lang, V. Khoze, A. Martin, and M. Ryskin, “Searching for the Odderon in Ultrapерipheral Proton–Ion Collisions at the LHC,” *Phys. Rev. D* **99** no. 3, (2019) 034011, [arXiv:1811.12705](#) [hep-ph].
- [50] R. Boussarie, A. Grabovsky, L. Szymanowski, and S. Wallon, “Impact factor for high-energy two and three jets diffractive production,” *JHEP* **09** (2014) 026, [arXiv:1405.7676](#) [hep-ph].
- [51] R. Boussarie, A. Grabovsky, L. Szymanowski, and S. Wallon, “On the one loop $\gamma^{(*)} \rightarrow q\bar{q}$ impact factor and the exclusive diffractive cross sections for the production of two or three jets,” *JHEP* **11** (2016) 149, [arXiv:1606.00419](#) [hep-ph].
- [52] R. Boussarie, A. Grabovsky, D. Y. Ivanov, L. Szymanowski, and S. Wallon, “Next-to-Leading Order Computation of Exclusive Diffractive Light Vector Meson Production in a Saturation Framework,” *Phys. Rev. Lett.* **119** no. 7, (2017) 072002, [arXiv:1612.08026](#) [hep-ph].
- [53] Z. Citron *et al.*, “Report from Working Group 5,” [arXiv:1812.06772](#) [hep-ph].
- [54] ATLAS Collaboration, G. Aad *et al.*, “Measurement of light-by-light scattering and search for axion-like particles with 2.2 nb⁻¹ of Pb+Pb data with the ATLAS detector,” [arXiv:2008.05355](#) [hep-ex].
- [55] D. d’Enterria and G. G. da Silveira, “Observing light-by-light scattering at the Large Hadron Collider,” *Phys. Rev. Lett.* **111** (2013) 080405, [arXiv:1305.7142](#) [hep-ph]. [Erratum: *Phys. Rev. Lett.* 116, 129901 (2016)].
- [56] R. Coelho, V. Gonçalves, D. Martins, and M. Rangel, “Exclusive and diffractive $\gamma\gamma$ production in *PbPb* collisions at the LHC, HE - LHC and FCC,” *Eur. Phys. J. C* **80** no. 5, (2020) 488, [arXiv:2002.03902](#) [hep-ph].
- [57] S. Fichet, G. von Gersdorff, B. Lenzi, C. Royon, and M. Saimpert, “Light-by-light scattering with intact protons at the LHC: from Standard Model to New Physics,” *JHEP* **02** (2015) 165, [arXiv:1411.6629](#) [hep-ph].
- [58] M. Bauer, M. Neubert, and A. Thamm, “Collider Probes of Axion-Like Particles,” *JHEP* **12** (2017) 044, [arXiv:1708.00443](#) [hep-ph].
- [59] S. Knapen, T. Lin, H. K. Lou, and T. Melia, “Searching for Axionlike Particles with Ultrapерipheral Heavy-Ion Collisions,” *Phys. Rev. Lett.* **118** no. 17, (2017) 171801, [arXiv:1607.06083](#) [hep-ph].
- [60] R. Coelho, V. Goncalves, D. Martins, and M. Rangel, “Production of axionlike particles in *PbPb* collisions at the LHC, HE–LHC and FCC: A phenomenological analysis,” *Phys. Lett. B* **806** (2020) 135512, [arXiv:2002.06027](#) [hep-ph].
- [61] J. Ellis, N. E. Mavromatos, and T. You, “Light-by-Light Scattering Constraint on Born-Infeld Theory,” *Phys. Rev. Lett.* **118** no. 26, (2017) 261802, [arXiv:1703.08450](#) [hep-ph].

- [62] P. Niau Akmansoy and L. G. Medeiros, “Constraining nonlinear corrections to Maxwell electrodynamics using $\gamma\gamma$ scattering,” *Phys. Rev.* **D99** no. 11, (2019) 115005, [arXiv:1809.01296 \[hep-ph\]](#).
- [63] ATLAS Collaboration, M. Aaboud *et al.*, “Evidence for light-by-light scattering in heavy-ion collisions with the ATLAS detector at the LHC,” *Nature Phys.* **13** no. 9, (2017) 852–858, [arXiv:1702.01625 \[hep-ex\]](#).
- [64] ATLAS Collaboration, G. Aad *et al.*, “Observation of light-by-light scattering in ultraperipheral Pb+Pb collisions with the ATLAS detector,” *Phys. Rev. Lett.* **123** no. 5, (2019) 052001, [arXiv:1904.03536 \[hep-ex\]](#).
- [65] CMS Collaboration, A. M. Sirunyan *et al.*, “Evidence for light-by-light scattering and searches for axion-like particles in ultraperipheral PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett.* **B797** (2019) 134826, [arXiv:1810.04602 \[hep-ex\]](#).
- [66] ATLAS Collaboration, “Observation of photon-induced W^+W^- production in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector,”.
- [67] CMS Collaboration, S. Chatrchyan *et al.*, “Study of Exclusive Two-Photon Production of W^+W^- in pp Collisions at $\sqrt{s} = 7$ TeV and Constraints on Anomalous Quartic Gauge Couplings,” *JHEP* **07** (2013) 116, [arXiv:1305.5596 \[hep-ex\]](#).
- [68] CMS, TOTEM Collaboration, “First search for exclusive diphoton production at high mass with intact protons in proton-proton collisions at $\sqrt{s} = 13$ TeV at the LHC CMS-PAS-EXO-18-014, TOTEM-NOTE-2020-003,”.
- [69] ATLAS Collaboration, “2018 heavy ion collision event (365914, 562492194): back-to-back electron-muon pair in an ultra-peripheral collision recorded with the ATLAS detector during the 2018 Pb+Pb data-taking period..” ATLAS-EVENTDISPLAY-2018-009, Nov, 2018.
- [70] M. Kłusek Gawenda, R. McNulty, R. Schicker, and A. Szczurek, “Light-by-light scattering in ultraperipheral heavy-ion collisions at low diphoton masses,” *Phys. Rev. D* **99** no. 9, (2019) 093013, [arXiv:1904.01243 \[hep-ph\]](#).
- [71] L. Beresford and J. Liu, “New physics and tau $g - 2$ using LHC heavy ion collisions,” [arXiv:1908.05180 \[hep-ph\]](#).
- [72] M. Dyndal, M. Klusek-Gawenda, M. Schott, and A. Szczurek, “Anomalous electromagnetic moments of τ lepton in $\gamma\gamma \rightarrow \tau^+\tau^-$ reaction in Pb+Pb collisions at the LHC,” *Phys. Lett. B* **809** (2020) 135682, [arXiv:2002.05503 \[hep-ph\]](#).
- [73] M. Kłusek Gawenda, P. Lebiedowicz, O. Nachtmann, and A. Szczurek, “From the $\gamma\gamma \rightarrow p\bar{p}$ reaction to the production of $p\bar{p}$ pairs in ultraperipheral ultrarelativistic heavy-ion collisions at the LHC,” *Phys. Rev. D* **96** no. 9, (2017) 094029, [arXiv:1708.09836 \[hep-ph\]](#).
- [74] V. Gonçalves and M. Jaime, “Photoproduction of pentaquark states at the LHC,” *Phys. Lett. B* **805** (2020) 135447, [arXiv:1911.10886 \[hep-ph\]](#).
- [75] S. R. Klein and Y.-P. Xie, “Photoproduction of charged final states in ultraperipheral collisions and electroproduction at an electron-ion collider,” *Phys. Rev. C* **100** no. 2, (2019) 024620, [arXiv:1903.02680 \[nucl-th\]](#).

- [76] S. Klein and J. Nystrand, “Exclusive vector meson production in relativistic heavy ion collisions,” *Phys. Rev.* **C60** (1999) 014903, [arXiv:hep-ph/9902259](#) [hep-ph].
- [77] M. Kłusek Gawenda and A. Szczurek, “Double-scattering mechanism in the exclusive $AA \rightarrow AA\rho^0\rho^0$ reaction in ultrarelativistic collisions,” *Phys. Rev. C* **89** no. 2, (2014) 024912, [arXiv:1309.2463](#) [nucl-th].
- [78] V. Goncalves, B. Moreira, and F. Navarra, “Double vector meson production in photon–hadron interactions at hadronic colliders,” *Eur. Phys. J. C* **76** no. 7, (2016) 388, [arXiv:1605.05840](#) [hep-ph].
- [79] S. R. Klein and J. Nystrand, “Does particle decay cause wave function collapse: An Experimental test,” *Phys. Lett. A* **308** (2003) 323–328, [arXiv:quant-ph/0206060](#).
- [80] J. C. Martens, J. P. Ralston, and J. D. T. Takaki, “Quantum tomography for collider physics: Illustrations with lepton pair production,” *Eur. Phys. J. C* **78** no. 1, (2018) 5, [arXiv:1707.01638](#) [hep-ph].
- [81] **LHeC, FCC-he Study Group** Collaboration, P. Agostini *et al.*, “The Large Hadron-Electron Collider at the HL-LHC,” [arXiv:2007.14491](#) [hep-ex].
- [82] S. R. Klein, J. Nystrand, and R. Vogt, “Photoproduction of top in peripheral heavy ion collisions,” *Eur. Phys. J. C* **21** (2001) 563–566, [arXiv:hep-ph/0005157](#) [hep-ph].
- [83] V. Gonçaves, “Photoproduction of top quarks in coherent hadron-hadron interactions,” *Phys. Rev. D* **88** no. 5, (2013) 054025.
- [84] D. d’Enterria, D. E. Martins, and P. Rebello Teles, “Higgs boson production in photon-photon interactions with proton, light-ion, and heavy-ion beams at current and future colliders,” *Phys. Rev. D* **101** no. 3, (2020) 033009, [arXiv:1904.11936](#) [hep-ph].
- [85] A. Accardi *et al.*, “Electron Ion Collider: The Next QCD Frontier: Understanding the glue that binds us all,” *Eur. Phys. J. A* **52** no. 9, (2016) 268, [arXiv:1212.1701](#) [nucl-ex].