Distributionally Robust Parametric Maximum Likelihood Estimation

Viet Anh Nguyen Xuhui Zhang José Blanchet
Stanford University, United States
{viet-anh.nguyen, xuhui.zhang, jose.blanchet}@stanford.edu

Angelos Georghiou

University of Cyprus, Cyprus georghiou.angelos@ucy.ac.cy

Abstract

We consider the parameter estimation problem of a probabilistic generative model prescribed using a natural exponential family of distributions. For this problem, the typical maximum likelihood estimator usually overfits under limited training sample size, is sensitive to noise and may perform poorly on downstream predictive tasks. To mitigate these issues, we propose a distributionally robust maximum likelihood estimator that minimizes the worst-case expected log-loss uniformly over a parametric Kullback-Leibler ball around a parametric nominal distribution. Leveraging the analytical expression of the Kullback-Leibler divergence between two distributions in the same natural exponential family, we show that the minmax estimation problem is tractable in a broad setting, including the robust training of generalized linear models. Our novel robust estimator also enjoys statistical consistency and delivers promising empirical results in both regression and classification tasks.

1 Introduction

We are interested in the relationship between a response variable Y and a covariate X governed by the generative model

$$Y|X = x \sim f(\cdot | \lambda(w_0, x)), \tag{1}$$

where λ is a pre-determined function that maps the weight w_0 and the covariate X to the parameter of the conditional distribution of Y given X. The weight w_0 is unknown and is the main quantity of interest to be estimated. Throughout this paper, we assume that the distribution f belongs to the exponential family of distributions. Given a ground measure ν on \mathcal{Y} , the exponential family is characterized by the density function

$$f(y|\theta) = h(y) \exp(\langle \theta, T(y) \rangle - \Psi(\theta))$$

with respect to ν , where $\langle \cdot, \cdot \rangle$ denotes the inner product, θ is the natural parameters, Ψ is the log-partition function and T is the sufficient statistics. The space of natural parameters is denoted by $\Theta = \left\{\theta: \int h(y) \exp(\langle \theta, T(y) \rangle) < \infty \right\} \subseteq \mathbb{R}^p$. We assume that the exponential family of distributions is regular, hence Θ is an open set, and $T_1(y), \ldots, T_p(y)$ are affinely independent [5, Chapter 8].

The generative setting (1) encapsulates numerous models which are suitable for regression and classification [17]. It ranges from logistic regression for classification [26], Poisson counting regression [25], log-linear models [15] to numerous other generalized linear models [17].

Given data $\{(\widehat{x}_i,\widehat{y}_i)\}_{i=1,\dots,N}$ which are assumed to be independently and identically distributed (i.i.d.) following the generative model (1), we want to estimate the true value of w_0 that dictates (1). If we use $\widehat{\mathbb{P}}^{\text{emp}} = N^{-1} \sum_{i=1}^{N} \delta_{(\widehat{x}_i, \widehat{y}_i)}$ to denote the empirical distribution supported on the training data, and define ℓ_{λ} as the log-loss function with the parameter mapping λ

$$\ell_{\lambda}(x, y, w) = \Psi(\lambda(w, x)) - \langle T(y), \lambda(w, x) \rangle, \tag{2}$$

then the maximum likelihood estimation (MLE) produces an estimate w_{MLE} by solving the following two equivalent optimization problems

nt optimization problems
$$w_{MLE} = \arg\min_{w \in \mathcal{W}} \sum_{i=1}^{N} \frac{1}{N} \left(\Psi(\lambda(w, \widehat{x}_i)) - \left\langle T(\widehat{y}_i), \lambda(w, \widehat{x}_i) \right\rangle \right)$$

$$= \arg\min_{w \in \mathcal{W}} \mathbb{E}_{\widehat{\mathbb{P}}^{emp}} [\ell_{\lambda}(X, Y, w)].$$
(3b)

$$= \arg\min_{w \in \mathcal{W}} \mathbb{E}_{\widehat{\mathbb{P}}^{emp}}[\ell_{\lambda}(X, Y, w)]. \tag{3b}$$

The popularity of MLE can be attributed to its consistency, asymptotic normality and efficiency [45, Section 5]. Unfortunately, this estimator exhibits several drawbacks in the finite sample regime, or when the data carry high noise and may be corrupted. For example, the ML estimator for the Gaussian model recovers the sample mean, which is notoriously susceptible to outliers [38]. The MLE for multinomial logistic regression yields over-fitted models for small and medium sized data [16].

Various strategies can be utilized to counter these adverse effects of the MLE in the limited data regime. The most common approach is to add a convex penalty term such as a 1-norm or 2-norm of w into the objective function of problem (3a) to obtain different regularization effects, see [35, 30] for regularized logistic regression. However, this approach relies on strong prior assumptions, such as the sparsity of w_0 for the 1-norm regularization, which may rarely hold in reality. Recently, dropout training has been used to prevent overfit and improve the generalization of the MLE [43, 46, 47]. Specific instances of dropout have been shown to be equivalent to a 2-norm regularization upon a suitable transformation of the inputs [46, Section 4]. Another popular strategy to regularize problem (3a) is by reweighting the samples instead of using a constant weight 1/N when calculating the loss. This approach is most popular in the name of weighted least-squares, which is a special instance of MLE problem under the Gaussian assumption with heteroscedastic noises.

Distributionally robust optimization (DRO) is an emerging scheme aiming to improve the out-ofsample performance of the statistical estimator, whereby the objective function of problem (3b) is minimized with respect to the most adverse distribution Q in some ambiguity set. The DRO framework has produced many interesting regularization effects. If the ambiguity set is defined using the Kullback-Leibler (KL) divergence, then we can recover an adversarial reweighting scheme [31, 10], a variance regularization [33, 20], and adaptive gradient boosting [12]. DRO models using KL divergence is also gaining recent attraction in many machine learning learning tasks [21, 41]. Another popular choice is the Wasserstein distance function which has been shown to have strong connections to regularization [40, 29], and has been used in training robust logistic regression classifiers [39, 11]. Alternatively, the robust statistics literature also consider the robustification of the MLE problem, for example, to estimate a robust location parameter [28]

Existing efforts using DRO typically ignore, or have serious difficulties in exploiting, the available information regarding the generative model (1). While existing approaches using the Kullback-Leibler ball around the empirical distribution completely ignore the possibility of perturbing the conditional distribution, the Wasserstein approach faces the challenge of elicitating a sensible ground metric on the response variables. For a concrete example, if we consider the Poisson regression application, then Y admits values in the space of natural numbers \mathbb{N} , and deriving a global metric on N that carries meaningful local information is nearly impossible because one unit of perturbation of an observation with $\hat{y}_i = 1$ does not carry the same amount of information as perturbing $\hat{y}_i = 1000$. The drawbacks of the existing methods behoove us to investigate a novel DRO approach that can incorporate the available information on the generative model in a systematic way.

Contributions. We propose the following distributionally robust MLE problem

$$\min_{w \in \mathcal{W}} \max_{\mathbb{Q} \in \mathbb{B}(\widehat{\mathbb{P}})} \mathbb{E}_{\mathbb{Q}} \left[\ell_{\lambda}(X, Y, w) \right], \tag{4}$$

which is a robustification of the MLE problem (3b) for generative models governed by an exponential family of distributions. The novelty in our approach can be summarized as follows.

· We advocate a new nominal distribution which is calibrated to reflect the available parametric information, and introduce a Kullback-Leibler ambiguity set that allows perturbations on both the marginal distribution of the covariate and the conditional distributions of the response.

- We show that the min-max estimation problem (4) can be reformulated as a single finite-dimensional minimization problem. Moreover, this reformulation is a convex optimization problem in broadly applicable settings, including the training of many generalized linear models.
- We demonstrate that our approach can recover the adversarial reweighting scheme as a special case, and it is connected to the variance regularization surrogate. Further, we prove that our estimator is consistent and provide insights on the practical tuning of the parameters of the ambiguity set. We also shed light on the most adverse distribution in the ambiguity set that incurs the extremal loss for any estimate of the statistician.

Technical notations. The variables (X,Y) admit values in $\mathcal{X} \times \mathcal{Y} \subseteq \mathbb{R}^n \times \mathbb{R}^m$, and \mathcal{W} is a finite-dimensional set. The mapping $\lambda: \mathcal{W} \times \mathcal{X} \to \Theta \subseteq \mathbb{R}^p$ is jointly continuous, and $\langle \cdot, \cdot \rangle$ denotes the inner product in \mathbb{R}^p . For any set \mathcal{S} , $\mathcal{M}(\mathcal{S})$ is the space of all probability measures with support on \mathcal{S} . We use \xrightarrow{p} to denote convergence in probability, and \xrightarrow{d} to denote convergence in distribution. All proofs are relegated to the appendix.

2 Distributionally Robust Estimation with a Parametric Ambiguity Set

We delineate in this section the ingredients of our distributionally robust MLE using parametric ambiguity set. Since the log-loss function is pre-determined, we focus solely on eliciting a nominal probability measure and the neighborhood surrounding it, which will serve as the ambiguity set.

While the typical empirical measure $\widehat{\mathbb{P}}^{\text{emp}}$ may appear at first as an attractive option for the nominal measure, $\widehat{\mathbb{P}}^{\text{emp}}$ does not reflect the parametric nature of the conditional measure of Y given X. Consequently, to robustify the MLE model, we need a novel construction of the nominal distribution $\widehat{\mathbb{P}}$.

Before proceeding, we assume w.l.o.g. that the dataset $\{(\widehat{x}_i,\widehat{y}_i)\}_{i=1,\dots,N}$ consists of $C \leq N$ distinct observations of X, each value is denoted by \widehat{x}_c for $c=1,\dots,C$, and the number of observations with the same covariate value \widehat{x}_c is denoted by N_c . This regrouping of the data by \widehat{x}_c typically enhances the statistical power of estimating the distribution conditional on the event $X=\widehat{x}_c$.

We posit the following parametric nominal distribution $\widehat{\mathbb{P}} \in \mathcal{M}(\mathcal{X} \times \mathcal{Y})$. This distribution is fully characterized by (p+1)C parameters: a probability vector $\widehat{p} \in \mathbb{R}_+^C$ whose elements sum up to 1 and a vector of nominal natural parameters $\widehat{\theta} \in \Theta^C \subseteq (\mathbb{R}^p)^C$. Mathematically, $\widehat{\mathbb{P}}$ satisfies

$$\begin{cases} \widehat{\mathbb{P}}(\{\widehat{x}_c\} \times A) = \widehat{\mathbb{P}}_X(\{\widehat{x}_c\})\widehat{\mathbb{P}}_{Y|\widehat{x}_c}(A) & \forall \widehat{x}_c, \forall A \subseteq \mathcal{Y} \text{ measurable} \\ \widehat{\mathbb{P}}_X = \sum_{c=1}^C \widehat{p}_c \delta_{\widehat{x}_c}, & \widehat{\mathbb{P}}_{Y|\widehat{x}_c} \sim f(\cdot|\widehat{\theta}_c) \ \forall c. \end{cases}$$
 (5)

The first equation indicates that the nominal measure $\widehat{\mathbb{P}}$ can be decomposed into a marginal distribution of the covariates X and a collection of conditional measures of Y given X using the definition of the conditional probability measure [44, Theorem 9.2.2]. The second line stipulates that the nominal marginal distribution $\widehat{\mathbb{P}}_X$ of the covariates is a discrete distribution supported on \widehat{x}_c , $c=1,\ldots,C$. Moreover, for each c, the nominal conditional distribution of Y given $X=\widehat{x}_c$ is a distribution in the exponential family with parameter $\widehat{\theta}_c$. Notice that the form of $\widehat{\mathbb{P}}$ in (5) is chosen to facilitate the injection of parametric information $\widehat{\theta}_c$ into the nominal distribution, and it is also necessary to tie $\widehat{\mathbb{P}}$ to the MLE problem using the following notion of MLE-compatibility.

Definition 2.1 (MLE-compatible nominal distribution). A nominal distribution $\widehat{\mathbb{P}}$ of the form (5) is MLE-compatible with respect to the log-loss function ℓ_{λ} if the optimal solution $\widehat{w} = \arg\min_{w \in \mathcal{W}} \mathbb{E}_{\widehat{\mathbb{P}}}[\ell_{\lambda}(X,Y,w)]$ coincides with the estimator w_{MLE} that solves (3a).

Definition 2.1 indicates that $\widehat{\mathbb{P}}$ is compatible for the MLE problem if the MLE solution w_{MLE} is recovered by solving problem (3b) where the expectation is now taken under $\widehat{\mathbb{P}}$. Therefore, MLE-compatibility implies that $\widehat{\mathbb{P}}$ and $\widehat{\mathbb{P}}^{\text{emp}}$ are equivalent in the MLE problem.

The next examples suggest two possible ways of calibrating an MLE-compatible $\widehat{\mathbb{P}}$ of the form (5).

Example 2.2 (Compatible nominal distribution I). If $\widehat{\mathbb{P}}$ is chosen of the form (5) with $\widehat{p}_c = N_c/N$ and $\widehat{\theta}_c = (\nabla \Psi)^{-1} \left((N_c)^{-1} \sum_{\widehat{x}_i = \widehat{x}_c} T(\widehat{y}_i) \right) \in \Theta$ for all c, then $\widehat{\mathbb{P}}$ is MLE-compatible.

Example 2.3 (Compatible nominal distribution II). If $\widehat{\mathbb{P}}$ is chosen of the form (5) with $\widehat{p}_c = N_c/N$ and $\widehat{\theta}_c = \lambda(w_{MLE}, \widehat{x}_c)$ for all c, where w_{MLE} solves (3a), then $\widehat{\mathbb{P}}$ is MLE-compatible.

We now detail the choice of the dissimilarity measure which is used to construct the neighborhood surrounding the nominal measure $\widehat{\mathbb{P}}$. For this, we will use the Kullback-Leiber divergence.

Definition 2.4 (Kullback-Leibler divergence). Suppose that \mathbb{P}_1 is absolutely continuous with respect to \mathbb{P}_2 , the Kullback-Leibler (KL) divergence from \mathbb{P}_1 to \mathbb{P}_2 is defined as $\mathrm{KL}(\mathbb{P}_1 \parallel \mathbb{P}_2) \triangleq \mathbb{E}_{\mathbb{P}_1} [\log(d\mathbb{P}_1/d\mathbb{P}_2)]$, where $d\mathbb{P}_1/d\mathbb{P}_2$ is the Radon-Nikodym derivative of \mathbb{P}_1 with respect to \mathbb{P}_2 .

The KL divergence is an ideal choice in our setting for numerous reasons. Previously, DRO problems with a KL ambiguity set often result in tractable finite-dimensional reformulations [7, 27, 10]. More importantly, the manifold of exponential family of distributions equipped with the KL divergence inherits a natural geometry endowed by a dually flat and invariant Riemannian structure [2, Chapter 2]. Furthermore, the KL divergence between two distributions in the same exponential family admits a closed form expression [4, 2].

Lemma 2.5 (KL divergence between distributions from exponential family). The KL divergence from $\mathbb{Q}_1 \sim f(\cdot | \theta_1)$ to $\mathbb{Q}_2 \sim f(\cdot | \theta_2)$ amounts to $\mathrm{KL}(\mathbb{Q}_1 \parallel \mathbb{Q}_2) = \langle \theta_1 - \theta_2, \nabla \Psi(\theta_1) \rangle - \Psi(\theta_1) + \Psi(\theta_2)$.

Using the above components, we are now ready to introduce our ambiguity set $\mathbb{B}(\widehat{\mathbb{P}})$ as

$$\mathbb{B}(\widehat{\mathbb{P}}) \triangleq \left\{ \mathbb{Q} \in \mathcal{M}(\mathcal{X}), \exists \theta_{c} \in \Theta \text{ such that } \mathbb{Q}_{Y \mid \widehat{x}_{c}} \sim f(\cdot \mid \theta_{c}) \quad \forall c \\ \mathbb{Q}(\{\widehat{x}_{c}\} \times A) = \mathbb{Q}_{X}(\{\widehat{x}_{c}\}) \mathbb{Q}_{Y \mid \widehat{x}_{c}}(A) \quad \forall c, \forall A \subseteq \mathcal{Y} \text{ measurable} \\ \mathrm{KL}(\mathbb{Q}_{Y \mid \widehat{x}_{c}} \parallel \widehat{\mathbb{P}}_{Y \mid \widehat{x}_{c}}) \leq \rho_{c} \quad \forall c \\ \mathrm{KL}(\mathbb{Q}_{X} \parallel \widehat{\mathbb{P}}_{X}) + \mathbb{E}_{\mathbb{Q}_{X}}[\sum_{c=1}^{C} \rho_{c} \mathbb{1}_{\widehat{x}_{c}}(X)] \leq \varepsilon \right\}$$
(6)

parametrized by a marginal radius ε and a collection of the conditional radii ρ_c . Any distribution $\mathbb{Q} \in \mathbb{B}(\widehat{\mathbb{P}})$ can be decomposed into a marginal distribution \mathbb{Q}_X of the covariate and an ensemble of parametric conditional distributions $\mathbb{Q}_{Y|\widehat{x}_c} \sim f(\cdot|\theta_c)$ at every event $X = \widehat{x}_c$. The first inequality in (6) restricts the parametric conditional distribution $\mathbb{Q}_{Y|\widehat{x}_c}$ to be in the ρ_c -neighborhood from the nominal $\widehat{\mathbb{P}}_{Y|\widehat{x}_c}$ prescribed using the KL divergence, while the second inequality imposes a similar restriction for the marginal distribution \mathbb{Q}_X . One can show that for any conditional radii $\rho \in \mathbb{R}_+^C$ satisfying $\sum_{c=1}^C \widehat{p}_c \rho_c \leq \varepsilon$, $\mathbb{B}(\widehat{\mathbb{P}})$ is non-empty with $\widehat{\mathbb{P}} \in \mathbb{B}(\widehat{\mathbb{P}})$. Moreover, if all ρ and ε are zero, then $\mathbb{B}(\widehat{\mathbb{P}})$ becomes the singleton set $\{\widehat{\mathbb{P}}\}$ that contains only the nominal distribution.

The set $\mathbb{B}(\widehat{\mathbb{P}})$ is a parametric ambiguity set: all conditional distributions $\mathbb{Q}_{Y|\widehat{x}_c}$ belong to the same parametric exponential family, and at the same time, the marginal distribution \mathbb{Q}_X is absolutely continuous with respect to a discrete distribution $\widehat{\mathbb{P}}_X$ and hence \mathbb{Q}_X can be parametrized using a C-dimensional probability vector.

At first glance, the ambiguity set $\mathbb{B}(\widehat{\mathbb{P}})$ looks intricate and one may wonder whether the complexity of $\mathbb{B}(\widehat{\mathbb{P}})$ is necessary. In fact, it is appealing to consider the ambiguity set

$$\mathcal{B}(\widehat{\mathbb{P}}) \triangleq \left\{ \mathbb{Q} \in \mathcal{M}(\mathcal{X} \times \mathcal{Y}) : \mathbb{Q}(\{\widehat{x}_c\} \times A) = \mathbb{Q}_X(\{\widehat{x}_c\}) \mathbb{Q}_{Y|\widehat{x}_c}(A) \ \forall c, \forall A \subseteq \mathcal{Y} \text{ measurable } \right\}$$
(7)
$$\text{KL}(\mathbb{Q} \parallel \widehat{\mathbb{P}}) \leq \varepsilon$$

which still preserves the parametric conditional structure and entails only one KL divergence constraint on the *joint* distribution space. Unfortunately, the ambiguity set $\mathcal{B}(\widehat{\mathbb{P}})$ may be overly conservative as pointed out in the following result.

Proposition 2.6. Denote momentarily the ambiguity sets (6) and (7) by $\mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}})$ and $\mathcal{B}_{\varepsilon}(\widehat{\mathbb{P}})$ to make the dependence on the radii explicit. For any nominal distribution $\widehat{\mathbb{P}}$ of the form (5) and any radius $\varepsilon \in \mathbb{R}_+$, we have

$$\mathcal{B}_{\varepsilon}(\widehat{\mathbb{P}}) = \bigcup_{\rho \in \mathbb{R}_{+}^{C}} \mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}}).$$

Proposition 2.6 suggests that the ambiguity set $\mathcal{B}(\widehat{\mathbb{P}})$ can be significantly bigger than $\mathbb{B}(\widehat{\mathbb{P}})$, and that the solution of the distributionally robust MLE problem (4) with $\mathbb{B}(\widehat{\mathbb{P}})$ being replaced by $\mathcal{B}(\widehat{\mathbb{P}})$ is potentially too conservative and may lead to undesirable or uninformative results.

The ambiguity set $\mathbb{B}(\widehat{\mathbb{P}})$ requires 1+C parameters, including one marginal radius ε and C conditional radii ρ_c , $c=1,\ldots,C$, which may be cumbersome to tune in the implementation. Fortunately, by the asymptotic result in Lemma 4.4, the set of radii ρ_c can be tuned simultaneously using the same scaling rate, which will significantly reduce the computational efforts for parameter tuning.

3 Tractable Reformulation

We devote this section to study the solution method for the min-max problem (4) by transforming it into a finite dimensional minimization problem. To facilitate the exposition, we denote the ambiguity set for the conditional distribution of Y given $X = \widehat{x}_c$ as

$$\mathbb{B}_{Y|\widehat{x}_c} \triangleq \left\{ \mathbb{Q}_{Y|\widehat{x}_c} \in \mathcal{M}(\mathcal{Y}) : \exists \theta \in \Theta, \ \mathbb{Q}_{Y|\widehat{x}_c}(\cdot) \sim f(\cdot|\theta), \ \mathrm{KL}(\mathbb{Q}_{Y|\widehat{x}_c} \parallel \widehat{\mathbb{P}}_{Y|\widehat{x}_c}) \leq \rho_c \right\}. \tag{8}$$

As a starting point, we first show the following decomposition of the worst-case expected loss under the ambiguity set $\mathbb{B}(\widehat{\mathbb{P}})$ for any measurable loss function.

Proposition 3.1 (Worst-case expected loss). Suppose that $\mathbb{B}(\widehat{\mathbb{P}})$ is defined as in (6) for some $\varepsilon \in \mathbb{R}_+$ and $\rho \in \mathbb{R}_+^C$ such that $\sum_{c=1}^C \widehat{p}_c \rho_c \leq \varepsilon$. For any function $L: \mathcal{X} \times \mathcal{Y} \to \mathbb{R}$ measurable, we have

$$\sup_{\mathbb{Q}\in\mathbb{B}(\widehat{\mathbb{P}})}\mathbb{E}_{\mathbb{Q}}\left[L(X,Y)\right] = \left\{ \begin{array}{ll} \inf & \alpha + \beta\varepsilon + \beta\sum_{c=1}^{C}\widehat{p}_{c}\exp\left(\beta^{-1}(t_{c} - \alpha) - \rho_{c} - 1\right) \\ \text{s. t.} & t\in\mathbb{R}^{C}, \ \alpha\in\mathbb{R}, \ \beta\in\mathbb{R}_{++} \\ & \sup_{\mathbb{Q}_{Y\mid\widehat{x}_{c}}\in\mathbb{B}_{Y\mid\widehat{x}_{c}}}\mathbb{E}_{\mathbb{Q}_{Y\mid\widehat{x}_{c}}}\left[L(\widehat{x}_{c},Y)\right] \leq t_{c} \quad \forall c=1,\ldots,C. \end{array} \right.$$

Proposition 3.1 leverages the decomposition structure of the ambiguity set $\mathbb{B}(\widehat{\mathbb{P}})$ to reformulate the worst-case expected loss into an infimum problem that involves C constraints, where each constraint is a hypergraph reformulation of a worst-case conditional expected loss under the ambiguity set $\mathbb{B}_{Y|\widehat{x}_c}$. Proposition 3.1 suggests that to reformulate the min-max estimation problem (4), it suffices now to reformulate the worst-case conditional expected log-loss

$$\sup_{\mathbb{Q}_{Y|\widehat{x}_c} \in \mathbb{B}_{Y|\widehat{x}_c}} \mathbb{E}_{\mathbb{Q}_{Y|\widehat{x}_c}} \left[\ell_{\lambda}(\widehat{x}_c, Y, w) \right] \tag{9}$$

for each value of \hat{x}_c into a dual infimum problem. Using Lemma 2.5, one can rewrite $\mathbb{B}_{Y|\hat{x}_c}$ in (8) using the natural parameter representation as

$$\mathbb{B}_{Y|\widehat{x}_c} = \left\{ \mathbb{Q}_{Y|\widehat{x}_c} \in \mathcal{M}(\mathcal{Y}) : \exists \theta \in \Theta, \mathbb{Q}_{Y|\widehat{x}_c}(\cdot) \sim f(\cdot|\theta), \left\langle \theta - \widehat{\theta}_c, \nabla \Psi(\theta) \right\rangle - \Psi(\theta) + \Psi(\widehat{\theta}_c) \leq \rho_c \right\}.$$

Since Ψ is convex [4, Lemma 1], it is possible that $\mathbb{B}_{Y|\widehat{x}_c}$ is represented by a non-convex set of natural parameters and hence reformulating (9) is non-trivial. Surprisingly, the next proposition asserts that problem (9) always admits a convex reformulation.

Proposition 3.2 (Worst-case conditional expected log-loss). For any $\widehat{x}_c \in \mathcal{X}$ and $w \in \mathcal{W}$, the worst-case conditional expected log-loss (9) is equivalent to the univariate convex optimization problem

$$\inf_{\gamma_c \in \mathbb{R}_{++}} \gamma_c \left(\rho_c - \Psi(\widehat{\theta}_c) \right) + \gamma_c \Psi(\widehat{\theta}_c - \gamma_c^{-1} \lambda(w, \widehat{x}_c)) + \Psi(\lambda(w, \widehat{x}_c)). \tag{10}$$

A reformulation for the worst-case conditional expected log-loss was proposed in [27]. Nevertheless, the results in [27, Section 5.3] requires that the sufficient statistics T(y) is a linear function of y. The reformulation (10), on the other hand, is applicable when T is a nonlinear function of y. Examples of exponential family of distributions with nonlinear T are (multivariate) Gaussian, Gamma and Beta distributions. The results from Propositions 3.1 and 3.2 lead to the reformulation of the distributionally robust estimation problem (4), which is the main result of this section.

Theorem 3.3 (Distributionally robust MLE reformulation). The distributionally robust MLE problem (4) is tantamount to the following finite dimensional optimization problem

inf
$$\alpha + \beta \varepsilon + \beta \sum_{c=1}^{C} \widehat{p}_c \exp(\beta^{-1}(t_c - \alpha) - \rho_c - 1)$$

s.t. $w \in \mathcal{W}, \ \alpha \in \mathbb{R}, \ \beta \in \mathbb{R}_{++}, \ \gamma \in \mathbb{R}_{++}^{C}, \ t \in \mathbb{R}^{C}$
 $\gamma_c(\rho_c - \Psi(\widehat{\theta}_c)) + \gamma_c \Psi(\widehat{\theta}_c - \gamma_c^{-1}\lambda(w, \widehat{x}_c)) + \Psi(\lambda(w, \widehat{x}_c)) \leq t_c \quad \forall c = 1, \dots, C.$

In generalized linear models with $\lambda:(w,x)\mapsto w^{\top}x$ and \mathcal{W} being convex, problem (11) is convex.

Below we show how the Poisson and logistic regression models fit within this framework.

Example 3.4 (Poisson counting model). The Poisson counting model with the ground measure ν being a counting measure on $\mathcal{Y} = \mathbb{N}$, the sufficient statistic T(y) = y, the natural parameter space $\Theta = \mathbb{R}$ and the log-partition function $\Psi(\theta) = \exp(\theta)$. If $\lambda(w, x) = w^{\top}x$, we have

$$Y|X = x \sim \text{Poisson}(w_0^{\top}x), \qquad \mathbb{P}(Y = k|X = x) = (k!)^{-1} \exp(kw_0^{\top}x - e^{w_0^{\top}x}).$$

The distributionally robust MLE is equivalent to the following convex optimization problem

inf
$$\alpha + \beta \varepsilon + \beta \sum_{c=1}^{C} \widehat{p}_c \exp\left(\beta^{-1}(t_c - \alpha) - \rho_c - 1\right)$$

s.t. $w \in \mathcal{W}, \ \alpha \in \mathbb{R}, \ \beta \in \mathbb{R}_{++}, \ \gamma \in \mathbb{R}_{++}^{C}, \ t \in \mathbb{R}^{C}$
 $\gamma_c(\rho_c - \exp(\widehat{\theta}_c)) + \gamma_c \exp\left(\widehat{\theta}_c - w^{\top}\widehat{x}_c/\gamma_c\right) + \exp\left(w^{\top}\widehat{x}_c\right) \le t_c \quad \forall c = 1, \dots, C.$

Example 3.5 (Logistic regression). The logistic regression model is specified with ν being a counting measure on $\mathcal{Y} = \{0,1\}$, the sufficient statistic T(y) = y, the natural parameter space $\Theta = \mathbb{R}$ and the log-partition function $\Psi(\theta) = \log \left(1 + \exp(\theta)\right)$. If $\lambda(w, x) = w^{\top} x$, we have

$$Y|X=x\sim \mathrm{Bernoulli}\big((1+\exp(-w_0^\top x))^{-1}\big), \qquad \mathbb{P}(Y=1|X=x)=(1+\exp(-w_0^\top x))^{-1}.$$

The distributionally robust MLE is equivalent to the following convex optimization problem

inf
$$\alpha + \beta \varepsilon + \beta \sum_{c=1}^{C} \widehat{p}_c \exp\left(\beta^{-1}(t_c - \alpha) - \rho_c - 1\right)$$

s.t. $w \in \mathcal{W}, \ \alpha \in \mathbb{R}, \ \beta \in \mathbb{R}_{++}, \ \gamma \in \mathbb{R}_{++}^{C}, \ t \in \mathbb{R}^{C}$
 $\gamma_c \left(\rho_c - \log(1 + \exp(\widehat{\theta}_c))\right) + \gamma_c \log\left(1 + \exp(\widehat{\theta}_c - w^{\top}\widehat{x}_c/\gamma_c)\right) + \log\left(1 + \exp(w^{\top}\widehat{x}_c)\right) \le t_c \ \forall c.$
(13)

Problems (13) and (12) can be solved by exponential conic solvers such as ECOS [18] and MOSEK [32].

4 Theoretical Analysis

In this section, we provide an in-depth theoretical analysis of our estimator. We first show that our proposed estimator is tightly connected to several existing regularization schemes.

Proposition 4.1 (Connection to the adversarial reweighting scheme). Suppose that \widehat{x}_i are distinct and $\rho_c=0$ for any $c=1,\ldots,N$. If $\widehat{\mathbb{P}}$ is of the form (5) and chosen according to Example 2.2, then the distributionally robust estimation problem (4) is equivalent to

$$\min_{w \in \mathcal{W}} \sup_{\mathbb{Q}: \mathrm{KL}(\mathbb{Q} \| \widehat{\mathbb{P}}^{\mathrm{emp}}) \leq \varepsilon} \mathbb{E}_{\mathbb{Q}}[\ell_{\lambda}(X, Y, w)].$$

Proposition 4.1 asserts that by setting the conditional radii to zero, we can recover the robust estimation problem where the ambiguity set is a KL ball around the empirical distribution $\widehat{\mathbb{P}}^{emp}$, which has been shown to produce the adversarial reweighting effects [31, 10]. Recently, it has been shown that distributionally robust optimization using f-divergences is statistically related to the variance regularization of the empirical risk minimization problem [34]. Our proposed estimator also admits a variance regularization surrogate, as asserted by the following proposition.

Proposition 4.2 (Variance regularization surrogate). Suppose that Ψ has locally Lipschitz continuous gradients. For any fixed $\widehat{\theta}_c \in \Theta$, $c=1,\ldots,C$, there exists a constant m>0 that depends only on Ψ and $\widehat{\theta}_c$, $c=1,\ldots,C$, such that for any $w\in \mathcal{W}$ and $\varepsilon\geq\sum_{c=1}^C\widehat{p}_c\rho_c$, we have

$$\sup_{\mathbb{Q}\in\mathbb{B}(\widehat{\mathbb{P}})} \mathbb{E}_{\mathbb{Q}}[\ell_{\lambda}(X,Y,w)] \leq \mathbb{E}_{\widehat{\mathbb{P}}}[\ell_{\lambda}(X,Y,w)] + \kappa_{1}\sqrt{\operatorname{Var}_{\widehat{\mathbb{P}}}(\ell_{\lambda}(X,Y,w))} + \kappa_{2}\|\lambda(w,\widehat{x}_{c})\|_{2},$$

where
$$\kappa_1 = \sqrt{2\varepsilon}/(\min_c \sqrt{\widehat{p}_c})$$
 and $\kappa_2 = \sqrt{2 \max_c \rho_c/m}$.

One can further show that for sufficiently small ρ_c , the value of m is proportional to the inverse of the local Lipschitz constant of $\nabla \Psi$ at $\hat{\theta}_c$, in which case κ_2 admits an explicit expression (see Appendix D). Next, we show that our robust estimator is also consistent, which is a highly desirable statistical property.

Theorem 4.3 (Consistency). Assume that w_0 is the unique solution of the problem $\min_{w \in \mathcal{W}} \mathbb{E}_{\mathbb{P}} \left[\ell_{\lambda}(X,Y,w) \right]$, where \mathbb{P} denotes the true distribution. Assume that \mathcal{X} has finite cardinality, $\Theta = \mathbb{R}^p$, Ψ has locally Lipschitz continuous gradients, and $\ell_{\lambda}(x,y,w)$ is convex in w for each x and y. If $\widehat{\theta}_c \xrightarrow{p} \lambda(w_0,\widehat{x}_c)$ for each $c, \varepsilon \to 0, \rho_c \to 0$ and $\varepsilon \ge \sum_{c=1}^C \widehat{p}_c \rho_c$ with probability going to 1, then the distributionally robust estimator w^* that solves (4) exists with probability going to 1, and $w^* \xrightarrow{p} w_0$.

One can verify that choosing $\widehat{\theta}_c$ using Examples 2.2 and 2.3 will satisfy the condition $\widehat{\theta}_c \stackrel{p.}{\longrightarrow} \lambda(w_0, \widehat{x}_c)$, and as a direct consequence, choosing $\widehat{\mathbb{P}}$ following these two examples will result in a consistent estimator under the conditions of Theorem 4.3.

We now consider the asymptotic scaling rate of ρ_c as the number N_c of samples with the same covariate \widehat{x}_c tends to infinity. Lemma 4.4 below asserts that ρ_c should scale at the rate N_c^{-1} . Based on this result, we can set $\rho_c = aN_c^{-1}$ for all c, where a>0 is a tuning parameter. This reduces significantly the burden of tuning ρ_c down to tuning a single parameter a.

Lemma 4.4 (Joint asymptotic convergence). Suppose that $|\mathcal{X}| = C$ with $\mathbb{P}(X = \widehat{x}_c) > 0$. Let $\theta_c = \lambda(w_0, \widehat{x}_c)$ and $\widehat{\mathbb{P}}$ be defined as in Example 2.2. Let $V_c = D_c \mathrm{Cov}_{f(\cdot|\theta_c)}(T(Y))D_c^{\top}$, where $D_c = J(\nabla \Psi)^{-1}(\mathbb{E}_{f(\cdot|\theta_c)}[T(Y)])$ and J denotes the Jacobian operator. Then the following joint convergence holds

Assuming w_{MLE} that solves (3a) is asymptotically normal with square-root convergence rate, we remark that the asymptotic joint convergence (14) also holds for $\widehat{\mathbb{P}}$ in Example 2.3, though in this case the limiting distribution Z takes a more complex form that can be obtained by the delta method.

Finally, we study the structure of the worst-case distribution $\mathbb{Q}^{\star} = \arg \max_{\mathbb{Q} \in \mathbb{B}(\widehat{\mathbb{P}})} \mathbb{E}_{\mathbb{Q}} [\ell_{\lambda}(X, Y, w)]$ for any value of input w. This result explicitly quantifies how the adversary will generate the adversarial distribution adapted to any estimate w provided by the statistician.

Theorem 4.5 (Worst-case joint distribution). Given $\rho \in \mathbb{R}_+^C$ and $\varepsilon \in \mathbb{R}_+$ such that $\sum_{c=1}^C \widehat{p}_c \rho_c \leq \varepsilon$. For any w and $c=1,\ldots,C$, let $\mathbb{Q}_{Y|\widehat{x}_c}^{\star} \sim f(\cdot|\theta_c^{\star})$ with $\theta_c^{\star} = \widehat{\theta}_c - \lambda(w,\widehat{x}_c)/\gamma_c^{\star}$, where $\gamma_c^{\star} > 0$ is the solution of the nonlinear equation

$$\Psi(\widehat{\theta}_c - \gamma^{-1}\lambda(w, \widehat{x}_c)) + \gamma^{-1}\langle\nabla\Psi(\widehat{\theta}_c - \gamma^{-1}\lambda(w, \widehat{x}_c)), \lambda(w, \widehat{x}_c)\rangle = \Psi(\widehat{\theta}_c) - \rho_c,$$

and let $t_c^\star = \Psi(\lambda(w, \widehat{x}_c)) - \langle \nabla \Psi(\theta_c^\star), \lambda(w, \widehat{x}_c) \rangle$. Let $\alpha^\star \in \mathbb{R}$ and $\beta^\star \in \mathbb{R}_{++}$ be the solution of the following system of nonlinear equations

$$\sum_{c=1}^{C} \widehat{p}_c \exp\left(\beta^{-1}(t_c^{\star} - \alpha) - \rho_c - 1\right) - 1 = 0$$
$$\sum_{c=1}^{C} \widehat{p}_c(t_c^{\star} - \alpha) \exp\left(\beta^{-1}(t_c^{\star} - \alpha) - \rho_c - 1\right) - (\varepsilon + 1)\beta = 0,$$

then the worst-case distribution is $\mathbb{Q}^{\star} = \sum_{c=1}^{C} \widehat{p}_{c} \exp\left((\beta^{\star})^{-1} (t_{c}^{\star} - \alpha^{\star}) - \rho_{c} - 1\right) \delta_{\widehat{x}_{c}} \otimes \mathbb{Q}_{Y|\widehat{x}_{c}}^{\star}$.

Notice that \mathbb{Q}^* is decomposed into a worst-case marginal distribution of X supported on \widehat{x}_c and a collection of worst-case conditional distributions $\mathbb{Q}_{Y|\widehat{x}_c}^*$.

5 Numerical Experiments

We now showcase the abilities of the proposed framework in the distributionally robust Poisson and logistic regression settings using a combination of simulated and empirical experiments. All optimization problems are modeled in MATLAB using CVX [23] and solved by the exponential conic solver MOSEK [32] on an Intel i7 CPU (1.90GHz) computer. Optimization problems (12) and (13) are solved in under 3 seconds for all instances both in the simulated and empirical experiments. The MATLAB code is available at https://github.com/angelosgeorghiou/DR-Parametric-MLE.




Figure 1: Median (solid blue line) and the 10th-90th percentile region (shaded) of out-of-sample divergence loss collected from 100 independent runs.

		N = 50	N = 100	N = 500
CI _{95%}	100(DRO-MLE)/MLE	$-69.84 \pm 3.33\%$	$-52.66 \pm 3.98\%$	$-22.25 \pm 4.11\%$
	$100(DRO - L_1)/L_1$	$-22.59 \pm 4.02\%$	$-19.90 \pm 4.16\%$	$-13.20 \pm 3.38\%$
	$100(DRO - L_2)/L_2$	$-14.98 \pm 4.08\%$	$-9.98 \pm 4.14\%$	$-5.62 \pm 2.83\%$
CVaR _{5%}	MLE	0.4906	0.1651	0.0246
	L_1	0.0967	0.0742	0.0195
	L_2	0.0894	0.0692	0.0176
	DRO	0.0547	0.0518	0.0172

Table 1: Comparison between the DRO estimator with the other methods. Lower values are better.

5.1 Poisson Regression

We will use simulated experiments to demonstrate the behavior of the tuning parameters and to compare the performance of our estimator with regard to other established methods. We assume that the true distribution \mathbb{P} is discrete, the 10-dimensional covariate X is supported on K=100 points and their locations \widehat{x}_k are generated i.i.d. using a standard normal distribution. We then generate a K-dimensional vector whose components are i.i.d. uniform over $M_k \in [0,10000]$, then normalize it to get the probability vector $p_k = M_k/M$ of the true marginal distribution of X. The value w_0 that determines the true conditional distribution $\mathbb{P}_{Y|X}$ via the generative model (1) is assigned to $w_0 = \widehat{w}/\|\widehat{w}\|_1$, where \widehat{w} is drawn randomly from a 10-dimensional standard normal distribution.

Our experiment comprises 100 simulation runs. In each run we generate $N \in \{50, 100, 500\}$ training samples i.i.d. from $\mathbb P$ and use the MLE-compatible nominal distribution $\widehat{\mathbb P}$ of the form (5) as in Example 2.3. We calibrate the regression model (12) by tuning $\rho_c = aN_c^{-1}$ with $a \in [10^{-4}, 1]$ and $\varepsilon \in [\sum_{c=1}^C \widehat{p}_c \rho_c, 1]$, both using a logarithmic scale with 20 discrete points. The quality of an estimate w^* with respect to the true distribution $\mathbb P$ is evaluated by the out-of-sample divergence loss

$$\mathbb{E}_{\mathbb{P}_X}[\mathrm{KL}(\mathbb{P}_{Y|X} \parallel \mathbb{Q}_{w^\star,Y|X})] = \sum_{k=1}^K p_k \left(\exp(w_0^\top \widehat{x}_k) \left((w_0 - w^\star)^\top \widehat{x}_k - 1 \right) + \exp(\widehat{x}_k^\top w^\star) \right).$$

In the first numerical experiment, we fix the marginal radius $\varepsilon=1$ and examine how tuning the conditional radii ρ_c can improve the quality of the estimator. Figure 1 shows the 10th, 50th and 90th percentile of the out-of-sample divergence for different samples sizes. If the constant a is chosen judiciously, incorporating the uncertainty in the conditional distribution can reduce the out-of-sample divergence loss by 17.65%, 10.55% and 1.82% for N=50,100 and 500, respectively.

Next, we compare the performance of our proposed estimator to the w_{MLE} that solves (3a) and the 1-norm (L_1) and 2-norm (L_2) MLE regularization, where the regularization weight takes values in $[10^{-4},1]$ on the logarithmic scale with 20 discrete points. In each run, we choose the optimal parameters that give the lowest of out-of-sample divergence for each method, and construct the empirical distribution of the out-of-sample divergence collected from 100 runs. Table 1 reports the 95% confidence intervals of $100(\mathrm{DRO-MLE})/\mathrm{MLE}$, $100(\mathrm{DRO-}L_1)/L_1$ and $100(\mathrm{DRO-}L_2)/L_2$, as well as the 5% Conditional Value-at-Risk (CVaR). Our approach delivers lower out-of-sample divergence loss compared to the other methods, and additionally ensures a lower value of CVaR for all sample sizes. This improvement is particularly evident in small sample sizes.

	AUC				CCR					
Dataset	DRO	KL	L_1	L_2	MLE	DRO	KL	L_1	L_2	MLE
australian ($N = 690, n = 14$)	92.74	92.62	92.73	92.71	92.61	85.75	85.72	85.52	85.60	85.72
banknote ($N = 1372, n = 4$)	98.46	98.46	98.43	98.45	98.45	94.31	94.32	94.16	94.35	94.32
climate $(N = 540, n = 18)$	94.30	82.77	94.85	94.13	82.76	95.04	93.89	94.85	94.83	93.89
german $(N = 1000, n = 19)$	75.75	75.68	75.74	75.74	75.67	73.86	74.05	73.82	73.70	74.05
haberman ($N = 306, n = 3$)	66.86	67.21	69.19	68.17	67.20	73.83	73.80	73.20	73.18	73.80
housing $(N = 506, n = 13)$	76.24	75.73	75.37	75.57	75.73	91.65	91.70	92.68	92.65	91.70
ILPD $(N = 583, n = 10)$	74.01	73.66	73.56	73.77	73.66	71.11	71.07	71.68	71.79	71.07
mammo. $(N = 830 n = 5)$	87.73	87.72	87.70	87.68	87.71	81.00	81.20	80.99	80.94	81.20

Table 2: Average area under the curve (AUC) and correct classification rates (CCR) on UCI datasets (m = 1).

5.2 Logistic Regression

We now study the performance of our proposed estimation in a classification setting using data sets from the UCI repository [19]. We compare four different models: our proposed DRO estimator (13), the w_{MLE} that solves (3a), the 1-norm (L_1) and 2-norm (L_2) MLE regularization. In each independent trial, we randomly split the data into train-validation-test set with proportion 50%-25%-25%. For our estimator, we calibrate the regression model (13) by tuning $\rho_c = aN_c^{-1}$ with $a \in [10^{-4}, 10]$ using a logarithmic scale with 10 discrete points and setting $\varepsilon = 2\sum_{c=1}^C \widehat{p}_c \rho_c$. Similarly, for the L_1 and L_2 regularization, we calibrate the regularization weight from $[10^{-4}, 1]$ on the logarithmic scale with 10 discrete points. Leveraging Proposition 4.1, we also compare our approach versus the DRO nonparametric Kullback-Leibler (KL) MLE by setting $\rho_c = 0$ and tune only with $\varepsilon \in [10^{-4}, 10]$ with 10 logarithmic scale points. The performance of the methods was evaluated on the testing data using two popular metrics: the correct classification rate (CCR) with a threshold level of 0.5, and the area under the receiver operating characteristics curve (AUC). Table 2 reports the performance of each method averaged over 100 runs. One can observe that our estimator performs reasonably well compared to other regularization techniques in both performance metrics.

Remark 5.1 (Uncertainty in \widehat{x}_c). The absolute continuity condition of the KL divergence implies that our proposed model cannot hedge against the error in the covariate \widehat{x}_c . It is natural to ask which model can effectively cover this covariate error. Unfortunately, answering this question needs to overcome to technical difficulties: first, the log-partition function Ψ is convex; second, the there are multiplicative terms between X and Y in the objective function. Maximizing over the X space to find the worst-case covariate is thus difficult. Alternatively, one can think of perturbing each \widehat{x}_c in a finite set but this approach will lead to trivial modifications of the constraints of problem (11).

Acknowledgments. Material in this paper is based upon work supported by the Air Force Office of Scientific Research under award number FA9550-20-1-0397. Additional support is gratefully acknowledged from NSF grants 1915967, 1820942, 1838676 and from the China Merchant Bank.

This appendix is organized as follows. Section A-C provide the detailed proofs for all the technical results in the main paper. Section D provides further discussion on the variance regularization surrogate result in Proposition 4.2.

A Proofs of Section 2

Proof of Example 2.2. We note that

$$\min_{w \in \mathcal{W}} \mathbb{E}_{\widehat{\mathbb{P}}}[\ell_{\lambda}(X, Y, w)] = \min_{w \in \mathcal{W}} \sum_{c=1}^{C} \widehat{p}_{c} \left(\Psi(\lambda(w, \widehat{x}_{c})) - \left\langle \mathbb{E}_{\widehat{\mathbb{P}}_{Y|\widehat{x}_{c}}}[T(Y)], \lambda(w, \widehat{x}_{c}) \right\rangle \right)$$

$$= \min_{w \in \mathcal{W}} \sum_{c=1}^{C} \widehat{p}_{c} \left(\Psi(\lambda(w, \widehat{x}_{c})) - \left\langle \nabla \Psi(\widehat{\theta}_{c}), \lambda(w, \widehat{x}_{c}) \right\rangle \right).$$

If $\widehat{\theta}_c = (\nabla \Psi)^{-1} ((N_c)^{-1} \sum_{\widehat{x}_i = \widehat{x}_o} T(\widehat{y}_i))$, then we have

$$\min_{w \in \mathcal{W}} \mathbb{E}_{\widehat{\mathbb{P}}}[\ell_{\lambda}(X, Y, w)] = \min_{w \in \mathcal{W}} \sum_{c=1}^{C} \widehat{p}_{c} \left(\Psi(\lambda(w, \widehat{x}_{c})) - \left\langle \frac{\sum_{\widehat{x}_{i} = \widehat{x}_{c}} T(\widehat{y}_{i})}{N_{c}}, \lambda(w, \widehat{x}_{c}) \right) \right)$$

$$= \min_{w \in \mathcal{W}} \frac{1}{N} \sum_{i=1}^{N} \left(\Psi(\lambda(w, \widehat{x}_{i})) - \left\langle T(\widehat{y}_{i}), \lambda(w, \widehat{x}_{i}) \right\rangle \right),$$

where we used $\widehat{p}_c = N_c/N$. Therefore w_{MLE} solves $\min_{w \in \mathcal{W}} \mathbb{E}_{\widehat{\mathbb{p}}}[\ell_{\lambda}(X,Y,w)]$.

Proof of Example 2.3. We find

$$\min_{w \in \mathcal{W}} \mathbb{E}_{\widehat{\mathbb{P}}}[\ell_{\lambda}(X, Y, w)] = \min_{w \in \mathcal{W}} \sum_{c=1}^{C} \widehat{p}_{c} \left(\Psi(\lambda(w, \widehat{x}_{c})) - \left\langle \mathbb{E}_{\widehat{\mathbb{P}}_{Y | \widehat{x}_{c}}}[T(Y)], \lambda(w, \widehat{x}_{c}) \right\rangle \right) \\
\geq \sum_{c=1}^{C} \widehat{p}_{c} \min_{w_{c} \in \mathcal{W}} \left(\Psi(\lambda(w_{c}, \widehat{x}_{c})) - \left\langle \mathbb{E}_{\widehat{\mathbb{P}}_{Y | \widehat{x}_{c}}}[T(Y)], \lambda(w_{c}, \widehat{x}_{c}) \right\rangle \right) \\
= \sum_{c=1}^{C} \widehat{p}_{c} \left(\Psi(\lambda(w_{MLE}, \widehat{x}_{c})) - \left\langle \mathbb{E}_{\widehat{\mathbb{P}}_{Y | \widehat{x}_{c}}}[T(Y)], \lambda(w_{MLE}, \widehat{x}_{c}) \right\rangle \right),$$

where the first equality follows from the definition of the log-loss function ℓ_{λ} , the inequality follows because $\widehat{p}_c > 0$, and the last equality follows because of the convex conjugate relationship that implies the optimal solution w_c^{\star} should satisfy

$$\nabla \Psi \big(\lambda(w_c^\star, \widehat{x}_c) \big) = \mathbb{E}_{\widehat{\mathbb{P}}_{Y \mid \widehat{x}_c}}[T(Y)] = \nabla \Psi \big(\lambda(w_{MLE}, \widehat{x}_c) \big) \implies w_c^\star = w_{MLE}.$$

This implies that w_{MLE} solves $\min_{w \in \mathcal{W}} \mathbb{E}_{\widehat{\mathbb{P}}}[\ell_{\lambda}(X,Y,w)]$ and completes the proof.

Proof of Proposition 2.6. Fix any set of conditional radii $\rho \in \mathbb{R}_+^C$. If $\mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}})$ is empty then it is trivial that $\mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}}) \subset \mathcal{B}_{\varepsilon}(\widehat{\mathbb{P}})$. Suppose that $\mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}})$ is non-empty and pick any $\mathbb{Q} \in \mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}})$. By definition of the set $\mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}})$, \mathbb{Q} can be decomposed into a marginal \mathbb{Q}_X and a collection of conditional measures $\mathbb{Q}_{Y|\widehat{x}_c}$. Furthermore, because ε is finite, the marginal \mathbb{Q}_X should be absolutely continuous with respect to $\widehat{\mathbb{P}}_X$. We have

$$\begin{split} \operatorname{KL}(\mathbb{Q} \parallel \widehat{\mathbb{P}}) &= \operatorname{KL}(\mathbb{Q}_X \parallel \widehat{\mathbb{P}}_X) + \mathbb{E}_{\mathbb{Q}_X} [\operatorname{KL}(\mathbb{Q}_{Y|X} \parallel \widehat{\mathbb{P}}_{Y|X})] \\ &\leq \operatorname{KL}(\mathbb{Q}_X \parallel \widehat{\mathbb{P}}_X) + \mathbb{E}_{\mathbb{Q}_X} [\sum_{c=1}^C \rho_c \mathbb{1}_{\widehat{x}_c}(X)] \leq \varepsilon, \end{split}$$

where the equality is from the chain rule of the conditional relative entropy [24, Lemma 7.9]. The first inequality follows from the fact that $\mathrm{KL}(\mathbb{Q}_{Y|\widehat{x}_c} \parallel \widehat{\mathbb{P}}_{Y|\widehat{x}_c}) \leq \rho_c$ for every c. The second inequality follows from the last constraint defining the set $\mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}})$. This implies that $\mathbb{Q} \in \mathcal{B}_{\varepsilon}(\widehat{\mathbb{P}})$, and because \mathbb{Q} was chosen arbitrarily, we have $\mathbb{B}_{\varepsilon,\rho} \subseteq \mathcal{B}_{\varepsilon}(\widehat{\mathbb{P}})$. As a consequence, $\bigcup_{\rho \in \mathbb{R}^c} \mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}}) \subseteq \mathcal{B}_{\varepsilon}(\widehat{\mathbb{P}})$.

Regarding the reverse relation, pick an arbitrary $\mathbb{Q} \in \mathcal{B}_{\varepsilon}(\widehat{\mathbb{P}})$ which admits the decomposition into a marginal \mathbb{Q}_X and conditional measures $\mathbb{Q}_{Y|\widehat{x}_c}$. By setting the conditional radii $\rho \in \mathbb{R}_+^C$ with $\rho_c = \mathrm{KL}(\mathbb{Q}_{Y|\widehat{x}_c} \parallel \widehat{\mathbb{P}}_{Y|\widehat{x}_c})$ for every c, one can verify using the chain rule of the conditional relative entropy that $\mathbb{Q} \in \mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}})$. This implies that $\mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}}) \subseteq \bigcup_{\rho \in \mathbb{R}_+^C} \mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}})$.

Concerning the last statement, notice that the condition $\sum_{c=1}^{C} \widehat{p}_c \rho_c \leq \varepsilon$ implies that $\widehat{\mathbb{P}} \in \mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}})$ and thus $\mathbb{B}_{\varepsilon,\rho}(\widehat{\mathbb{P}})$ is non-empty. The proof is complete.

B Proofs of Section 3

The proof of Proposition 3.1 relies on the following preliminary result.

Lemma B.1. Let $\widehat{p} \in \mathbb{R}_{++}^C$ be a probability vector summing up to one. For any $\varepsilon \in \mathbb{R}_+$ and $\rho \in \mathbb{R}_+^C$ satisfying $\sum_{c=1}^C \widehat{p}_c \rho_c \leq \varepsilon$, the finite dimensional set

$$Q \triangleq \left\{ q \in \mathbb{R}_+^C : \sum_{c=1}^C q_c = 1, \ \sum_{c=1}^C q_c (\log q_c - \log \widehat{p}_c + \rho_c) \le \varepsilon \right\}$$
 (16)

is compact and convex. Moreover, the support function $h_\mathcal{Q}$ of \mathcal{Q} satisfies

$$\forall t \in \mathbb{R}^C : \quad h_{\mathcal{Q}}(t) \triangleq \sup_{q \in \mathcal{Q}} q^\top t = \inf_{\alpha \in \mathbb{R}, \ \beta \in \mathbb{R}_{++}} \left\{ \alpha + \beta \varepsilon + \beta \sum_{c=1}^C \widehat{p}_c \exp\left(\frac{t_c - \alpha}{\beta} - \rho_c - 1\right) \right\}.$$

Proof of Lemma B.1. The function $\mathbb{R}_+^C \ni q \mapsto \sum_{c=1}^C q_c (\log q_c - \log \widehat{p}_c + \rho_c) \in \mathbb{R}_+$ is continuous and convex, hence, the set $\{q \in \mathbb{R}_+^C : \sum_{c=1}^C q_c (\log q_c - \log \widehat{p}_c + \rho_c) \le \varepsilon\}$ is closed and convex. Consequentially, \mathcal{Q} can be written as the intersection between a simplex (thus compact and convex) and a closed, convex set, so \mathcal{Q} is compact and convex.

The proof of the support function of $\mathcal Q$ proceeds in 2 steps. First, we prove the support function for the ϵ -inflated set

$$Q_{\epsilon} = \left\{ q \in \mathbb{R}_{+}^{C} : \sum_{c=1}^{C} q_{c} = 1, \sum_{c=1}^{C} q_{c} (\log q_{c} - \log \widehat{p}_{c} + \rho_{c}) \leq \varepsilon + \epsilon \right\}$$

with the right-hand side of the last constraint being inflated with $\epsilon \in \mathbb{R}_{++}$. In the second step, we use a limit argument to show that the support function of \mathcal{Q} is attained as the limit of the support function of \mathcal{Q}_{ϵ} as ϵ tends to 0.

Reminding that Δ is the C-dimensional simplex. For any $t \in \mathbb{R}^C$ and any $\epsilon \in \mathbb{R}_{++}$, by the definition of the support function, we have for every $t \in \mathbb{R}^C$

$$h_{\mathcal{Q}_{\epsilon}}(t) = \begin{cases} \sup q^{\top} t \\ \text{s. t.} \quad q \in \Delta, \quad \sum_{c=1}^{C} q_{c}(\log q_{c} - \log \widehat{p}_{c} + \rho_{c}) \leq \varepsilon + \epsilon \end{cases}$$

$$= \sup_{q \in \Delta} \inf_{\beta \in \mathbb{R}_{+}} q^{\top} t + \beta(\varepsilon + \epsilon - \sum_{c=1}^{C} q_{c}(\log q_{c} - \log \widehat{p}_{c} + \rho_{c})$$

$$= \inf_{\beta \in \mathbb{R}_{+}} \sup_{q \in \Delta} q^{\top} t + \beta(\varepsilon + \epsilon - \sum_{c=1}^{C} q_{c}(\log q_{c} - \log \widehat{p}_{c} + \rho_{c}),$$

$$(17a)$$

where the interchange of the sup-inf operators in (17b) is justified by strong duality [9, Proposition 5.3.1] because \hat{p} constitutes a Slater point of the set \mathcal{Q}_{ϵ} . By Berge's maximum theorem [8], the optimal value of the inner supremum problem is a continuous function in β because the simplex Δ is compact and the objective function is continuous in the decision variable q. As a consequence, we can restrict $\beta \in \mathbb{R}_{++}$ without any loss of optimality. Because Δ is prescribed using linear

constraints, strong duality implies that

$$h_{\mathcal{Q}_{\epsilon}}(t) = \inf_{\alpha \in \mathbb{R}, \ \beta \in \mathbb{R}_{++}} \left\{ \alpha + \beta(\varepsilon + \epsilon) + \sup_{q \in \mathbb{R}_{+}^{C}} \sum_{c=1}^{C} q_{c}(t_{c} - \alpha + \beta \log \widehat{p}_{c} - \beta \rho_{c} - \beta \log q_{c}) \right\}$$
$$= \inf_{\alpha \in \mathbb{R}, \ \beta \in \mathbb{R}_{++}} \left\{ \alpha + \beta(\varepsilon + \epsilon) + \sum_{c=1}^{C} \sup_{q_{c} \in \mathbb{R}_{+}} q_{c}(t_{c} - \alpha + \beta \log \widehat{p}_{c} - \beta \rho_{c} - \beta \log q_{c}) \right\},$$

where the last equality holds because the supremum problem is separable in each decision variable q_c . It now follows from the first-order optimality condition that the maximizer q_c^{\star} is

$$q_c^{\star} = \exp\left(\frac{t_c - \alpha + \beta \log \widehat{p}_c - \beta \rho_c - \beta}{\beta}\right) > 0,$$

and by substituting this maximizer into the objective function, the value of the support function $h_{\mathcal{Q}_{\epsilon}}(t)$ is then equal to the optimal value of the below optimization problem

$$\inf_{\alpha \in \mathbb{R}, \ \beta \in \mathbb{R}_{++}} \alpha + \beta(\varepsilon + \epsilon) + \beta \sum_{c=1}^{C} \widehat{p}_c \exp\left(\frac{t_c - \alpha}{\beta} - \rho_c - 1\right).$$

We now proceed to the second step. Denote temporarily the objective function of the above problem as $G(\epsilon, \gamma)$, where $\gamma = [\alpha; \beta]$ combines both dual variables α and β . Define the function

$$g(\epsilon) = \inf_{\gamma \in \Gamma} \ G(\epsilon, \gamma), \qquad \text{with } \Gamma \triangleq \mathbb{R} \times \mathbb{R}_{++}.$$

Because G is continuous, [36, Lemma 2.7] implies that g is upper-semicontinuous at 0. Furthermore, G is calm from below at $\epsilon=0$ because $G(\epsilon,\gamma)-G(0,\gamma)=\beta\epsilon\geq 0$, thus [36, Lemma 2.7] implies that g is lower-semicontinuous at 0. These two facts lead to the continuity of g at 0. From the first part of the proof, we have $g(\epsilon)=h_{\mathcal{Q}_{\epsilon}}(t)$ for any $\epsilon\in\mathbb{R}_+$. Moreover, by applying Berge's maximum theorem [8] to (17a), $h_{\mathcal{Q}_{\epsilon}}(t)$ is a continuous function of ϵ over \mathbb{R}_+ . Thus we find

$$h_{\mathcal{Q}}(t) = h_{\mathcal{Q}_0}(t) = \lim_{\epsilon \downarrow 0} h_{\mathcal{Q}_{\epsilon}}(t) = \lim_{\epsilon \downarrow 0} g(\epsilon) = g(0),$$

where the chain of equalities follows from the definition of \mathcal{Q}_{ϵ} , the continuity of $h_{\mathcal{Q}_{\epsilon}}(t)$ in ϵ , the fact that $g(\epsilon) = h_{\mathcal{Q}_{\epsilon}}(t)$ for $\epsilon > 0$, and the continuity of g at 0 established previously. The proof is now completed.

Proof of Proposition 3.1. To facilitate the proof, we define the following ambiguity set over the marginal distribution of the covariate X as

$$\mathbb{B}_X \triangleq \left\{ \mathbb{Q}_X \in \mathcal{M}(\mathcal{X}) : \mathrm{KL}(\mathbb{Q}_X \parallel \widehat{\mathbb{P}}_X) + \mathbb{E}_{\mathbb{Q}_X} [\sum_{c=1}^C \rho_c \mathbb{1}_{\widehat{x}_c}(X)] \le \varepsilon \right\}.$$

Given a nominal marginal distribution $\widehat{\mathbb{P}}_X$ supported on a finite set $\{\widehat{x}_c\}_{c\in\mathcal{C}}$, the absolute continuity requirement suggests that $\mathrm{KL}(\mathbb{Q}_X\parallel\widehat{\mathbb{P}}_X)$ is finite if and only if \mathbb{Q}_X is absolutely continuous with respect to $\widehat{\mathbb{P}}_X$. Thus, any \mathbb{Q}_X of interest should be supported on the same set $\{\widehat{x}_c\}_{c=1,\ldots,C}$, and \mathbb{Q}_X and be finitely parametrized by a C-dimensional vector $\{q_c\}_{c=1,\ldots,C}$. Let \mathcal{Q} denote the convex compact feasible set in \mathbb{R}^C , that is,

$$Q \triangleq \left\{ q \in \mathbb{R}_+^C : \sum_{c=1}^C q_c = 1, \sum_{c=1}^C q_c (\log q_c - \log \widehat{p}_c + \rho_c) \le \varepsilon \right\},\,$$

and the ambiguity set \mathbb{B}_X can now be finitely parametrized as

$$\mathbb{B}_X = \left\{ \mathbb{Q}_X \in \mathcal{M}(\mathcal{X}) : \exists q \in \mathcal{Q}, \ \mathbb{Q}_X = \sum_{i=1}^C q_c \delta_{\widehat{x}_c} \right\}.$$

By coupling \mathbb{B}_X with the conditional ambiguity sets $\mathbb{B}_{Y|\widehat{x}_c}$, $\mathbb{B}(\widehat{\mathbb{P}})$ can be re-written as

$$\mathbb{B}(\widehat{\mathbb{P}}) = \left\{ \mathbb{Q} \in \mathcal{M}(\mathcal{X} \times \mathcal{Y}) : \begin{array}{l} \exists \mathbb{Q}_X \in \mathbb{B}_X, \ \mathbb{Q}_{Y \mid \widehat{x}_c} \in \mathbb{B}_{Y \mid \widehat{x}_c} \ \forall c = 1, \dots, C \\ \mathbb{Q}(\{\widehat{x}_c\} \times A) = \mathbb{Q}_X(\{\widehat{x}_c\}) \mathbb{Q}_{Y \mid \widehat{x}_c}(A) \ \forall A \in \mathcal{F}(\mathcal{Y}) \quad \forall c = 1, \dots, C \end{array} \right\}$$

The worst-case expected loss becomes

$$\begin{split} \sup_{\mathbb{Q} \in \mathbb{B}(\widehat{\mathbb{P}})} \; \mathbb{E}_{\mathbb{Q}}[L(X,Y)] &= \sup_{\mathbb{Q}_X \in \mathbb{B}_X} \mathbb{E}_{\mathbb{Q}_X} \left[\sup_{\mathbb{Q}_{Y|X} \in \mathbb{B}_{Y|X}} \mathbb{E}_{\mathbb{Q}_{Y|X}} \left[L(X,Y) \right] \right] \\ &= \sup_{q \in \mathcal{Q}} \; \sum_{c=1}^C q_c \sup_{\mathbb{Q}_{Y|\hat{x}_c} \in \mathbb{B}_{Y|\hat{x}_c}} \mathbb{E}_{\mathbb{Q}_{Y|\hat{x}_c}} \left[L(\hat{x}_c,Y) \right], \end{split}$$

where the first equality follows from the law of total expectation, and the second equality follows from the finite reparametrization of \mathbb{B}_X . If we denote by \mathcal{T} the epigraph reformulation of the worst-case conditional expectations

$$\mathcal{T} \triangleq \left\{ t \in \mathbb{R}^C : \sup_{\mathbb{Q}_{Y \mid \widehat{x}_c} \in \mathbb{B}_{Y \mid \widehat{x}_c}} \mathbb{E}_{\mathbb{Q}_{Y \mid \widehat{x}_c}} \left[L(\widehat{x}_c, Y) \right] \leq t_c \quad \forall c = 1, \dots, C \right\},$$

then the worst-case expected loss can be further re-expressed as

$$\sup_{\mathbb{Q}\in\mathbb{B}(\widehat{\mathbb{P}})} \mathbb{E}_{\mathbb{Q}}[L(X,Y)] = \sup_{q\in\mathcal{Q}} \inf_{t\in\mathcal{T}} q^{\top}t$$
(18a)

$$= \inf_{t \in \mathcal{T}} \sup_{q \in \mathcal{O}} q^{\top} t \tag{18b}$$

$$= \begin{cases} & \text{inf} \quad \alpha + \beta \varepsilon + \beta \sum_{c=1}^{C} \widehat{p}_c \exp\left(\frac{t_c - \alpha}{\beta} - \rho_c - 1\right) \\ & \text{s. t.} \quad t \in \mathcal{T}, \ \alpha \in \mathbb{R}, \ \beta \in \mathbb{R}_{++}, \end{cases}$$
(18c)

where the sup-inf formulation (18a) is justified because q is non-negative and we can resort to the epigraph formulations of the worst-case conditional expected loss. In (18b) we applied Sion's minimax theorem [42], which is valid because the sup-inf program (18a) is a concave-convex saddle problem, and Q is convex and compact and T is convex. In (18c) we have used Lemma B.1 to reformulate the supremum over q. The claim then follows.

Instead of solving the problem in the natural parameters θ coupled with its log-partition function Ψ , we will use the reparametrization to the mean parameters using the conjugate function of Ψ . More specifically, let ϕ be the convex conjugate of Ψ , that is,

$$\phi: \mu \mapsto \sup_{\theta \in \Theta} \left\{ \left\langle \mu, \theta \right\rangle - \Psi(\theta) \right\}$$

Before proceeding to the technical proofs, the below lemma collects from the existing literature the necessary background knowledge about the log-partition function Ψ and its conjugate ϕ , along with the relationship between the natural parameter θ and its corresponding expectation parameter μ .

Lemma B.2 (Relevant facts). The following assertions hold for regular exponential family.

- (i) The function ϕ is closed, convex and proper on \mathbb{R}^p .
- (ii) (Θ, Ψ) and $(\operatorname{int}(\operatorname{dom}(\phi)), \phi)$ are convex functions of Legendre type, and they are Legendre duals of each other.
- (iii) The gradient function $\nabla \Psi$ is a one-to-one function from the open convex set Θ onto the open convex set $\operatorname{int}(\operatorname{dom}(\phi))$.
- (iv) The gradient functions $\nabla \Psi$ and $\nabla \phi$ are continuous, and $\nabla \phi = (\nabla \Psi)^{-1}$.
- (v) The function ϕ is essentially smooth over $int(dom(\phi))$.

Proof of Lemma B.2. Assertion (i) holds since $\langle \mu, \theta \rangle - \Psi(\theta)$ is convex and closed for each θ , thus taking supremum, ϕ is convex and closed. ϕ is proper since $dom(\phi)$ is non-empty. Assertions (ii) to (iv) follows from [4, Lemma 1] and [4, Theorem 2]. Assertion (v) follows from [4, Lemma 1] and [37, Theorem 26.3], and the fact that Ψ and ϕ is a convex conjugate pair.

From Assertion (ii), we have the mappings between the dual spaces $\operatorname{int}(\operatorname{dom}(\phi))$ and Θ are given by the Legendre transformation

$$\mu(\theta) = \nabla \Psi(\theta) \quad \text{and} \quad \theta(\mu) = \nabla \phi(\mu).$$

For any $\mu \in \operatorname{int}(\operatorname{dom}(\phi))$, the conjugate function ϕ can be expressed as

$$\phi(\mu) = \langle \mu, \theta(\mu) \rangle - \Psi(\theta(\mu)).$$

Lemma B.3 (KL divergence between distributions from exponential family). Suppose that \mathbb{Q}_1 and \mathbb{Q}_2 belong to the exponential family of distributions with the same log-partition function Ψ and with natural parameters θ_1 and θ_2 respectively. The KL divergence from \mathbb{Q}_1 to \mathbb{Q}_2 amounts to

$$\mathrm{KL}(\mathbb{Q}_1 \parallel \mathbb{Q}_2) = \langle \theta_1 - \theta_2, \mu_1 \rangle - \Psi(\theta_1) + \Psi(\theta_2) = \phi(\mu_1) - \phi(\mu_2) - \langle \mu_1 - \mu_2, \theta_2 \rangle,$$

where ϕ is the convex conjugate of Ψ , and $\mu_j = \nabla \Psi(\theta_j)$ for any $j \in \{1, 2\}$.

The result of Lemma B.3 can be found in [4, Appendix A], but the explicit proof is included here for completeness.

Proof of Lemma B.3. One finds

$$KL(\mathbb{Q}_1 \parallel \mathbb{Q}_2) = \mathbb{E}_{\mathbb{Q}_1}[\log(d\mathbb{Q}_1/d\mathbb{Q}_2)]$$

$$= \mathbb{E}_{\mathbb{Q}_1}[\langle T(Y), \theta_1 - \theta_2 \rangle - \Psi(\theta_1) + \Psi(\theta_2)]$$

$$= \langle \mu_1, \theta_1 - \theta_2 \rangle - \Psi(\theta_1) + \Psi(\theta_2),$$
(19a)
$$(19b)$$

where equality (19a) follows by calculating the logarithm of the Radon-Nikodym derivatives between two distributions, and equality (19b) follows by noting that $\mu_1 = \mathbb{E}_{\mathbb{Q}_1}[T(Y)]$.

By [4, Theorem 4], one can also rewrite the density using the mean parameter $\mu = \mu(\theta)$ as

$$f(y|\mu) = h(y) \exp (\langle \theta, T(y) \rangle - \Psi(\theta))$$

= $h(y) \exp (\phi(\mu) + \langle T(y) - \mu, \nabla \phi(\mu) \rangle)$

The KL divergence from \mathbb{Q}_1 to \mathbb{Q}_2 amounts to

$$KL(\mathbb{Q}_1 \parallel \mathbb{Q}_2) = \mathbb{E}_{\mathbb{Q}_1}[\log(d\mathbb{Q}_1/d\mathbb{Q}_2)]$$

$$= \mathbb{E}_{\mathbb{Q}_1}[\phi(\mu_1) - \phi(\mu_2) + \langle T(Y), \nabla \phi(\mu_1) - \nabla \phi(\mu_2) \rangle - \langle \mu_1, \nabla \phi(\mu_1) \rangle + \langle \mu_2, \nabla \phi(\mu_2) \rangle]$$

$$= \langle \mu_2 - \mu_1, \theta_2 \rangle + \phi(\mu_1) - \phi(\mu_2).$$
(20b)

From Assertion (iv) in Lemma B.2, we notice that $\theta_2 = \nabla \phi(\mu_2)$, which completes the proof. \square

Recall that the conditional ambiguity set defined in (8) is

$$\mathbb{B}_{Y|\widehat{x}_c} \triangleq \left\{ \mathbb{Q}_{Y|\widehat{x}_c} \in \mathcal{M}(\mathcal{Y}) : \exists \theta \in \Theta, \ \mathbb{Q}_{Y|\widehat{x}_c}(\cdot) \sim f(\cdot|\theta), \ \mathrm{KL}(\mathbb{Q}_{Y|\widehat{x}_c} \parallel \widehat{\mathbb{P}}_{Y|\widehat{x}_c}) \leq \rho_c \right\}$$

for a parametric, nominal conditional measure $\widehat{\mathbb{P}}_{Y|\widehat{x}_c} \sim f(\cdot|\widehat{\theta}_c), \widehat{\theta}_c \in \Theta$ and a radius $\rho_c \in \mathbb{R}_+$. The uncertainty set \mathcal{S}_c of expectation parameters induced by the ambiguity set $\mathbb{B}_{Y|\widehat{x}_c}$ is defined as

$$\mathcal{S}_c \triangleq \left\{ \mu \in \text{dom}(\phi) : \exists \mathbb{Q}_{Y \mid \widehat{x}_c} \in \mathbb{B}_{Y \mid \widehat{x}_c}, \ \mu = \mathbb{E}_{\mathbb{Q}_{Y \mid \widehat{x}_c}}[T(Y)] \right\}.$$

Lemma B.4 (Compactness of expectation parameter uncertainty set). The set S_c is compact, and it has an interior point whenever $\rho_c > 0$.

Proof of Lemma B.4. By Lemma B.3 and the definition of the set S_c , we can write S_c as

$$S_c = \left\{ \mu \in \text{dom}(\phi) : \phi(\mu) - \phi(\widehat{\mu}_c) - \left\langle \mu - \widehat{\mu}_c, \widehat{\theta}_c \right\rangle \le \rho_c \right\}.$$

Because ϕ is closed, convex, proper, and that $\widehat{\theta}_c \in \operatorname{int}(\Theta) = \Theta$, the function $\phi(\cdot) - \langle \cdot, \widehat{\theta}_c \rangle$ is coercive by [37, Corollary 14.2.2] and [6, Fact 2.11]. As a consequence, \mathcal{S}_c is bounded.

Because Ψ is essentially strictly convex on Θ , ϕ is essentially smooth on $\operatorname{int}(\operatorname{dom}(\phi))$ by [37, Theorem 26.3]. [6, Theorem 3.8] now implies that if μ' is a boundary point of $\operatorname{int}(\operatorname{dom}(\phi))$ then as $\operatorname{int}(\operatorname{dom}(\phi)) \ni \mu_k \xrightarrow{k \to \infty} \mu'$ then $\phi(\mu_k) - \langle \mu_k, \widehat{\theta}_c \rangle \xrightarrow{k \to \infty} + \infty$. Moreover, because ϕ is continuous over $\operatorname{int}(\operatorname{dom}(\phi))$, the set \mathcal{S}_c is closed. This implies that \mathcal{S}_c , being a closed and bounded set of finite dimension, is compact.

The continuity of ϕ leads a straightforward manner to the non-empty interior of S_c when $\rho_c > 0$. This observation completes the proof.

Proof of Proposition 3.2. Because λ is a mapping onto the space Θ of natural parameters, we use the shorthand $\lambda_c = \lambda(w, \widehat{x}_c) \in \Theta$. Moreover, let $\widehat{\mu}_c = \nabla \Psi(\widehat{\theta}_c)$. The worst-case conditional expectation of the log-loss function becomes

$$\begin{split} \sup_{\mathbb{Q}_{Y|\hat{x}_c} \in \mathbb{B}_{Y|\hat{x}_c}} \mathbb{E}_{\mathbb{Q}_{Y|\hat{x}_c}} \left[\ell_{\lambda}(\hat{x}_c, Y, w) \right] &= \sup_{\mathbb{Q}_{Y|\hat{x}_c} \in \mathbb{B}_{Y|\hat{x}_c}} \mathbb{E}_{\mathbb{Q}_{Y|\hat{x}_c}} \left[\Psi(\lambda(w, \hat{x}_c)) - \left\langle T(Y), \lambda(w, \hat{x}_c) \right\rangle \right] \\ &= \sup_{\mathbb{Q}_{Y|\hat{x}_c} \in \mathbb{B}_{Y|\hat{x}_c}} \Psi(\lambda(w, \hat{x}_c)) - \left\langle \mathbb{E}_{\mathbb{Q}_{Y|\hat{x}_c}} [T(Y)], \lambda(w, \hat{x}_c) \right\rangle \\ &= \left\{ \begin{array}{ll} \sup & \Psi(\lambda_c) - \left\langle \mu, \lambda_c \right\rangle \\ \mathrm{s. t.} & \phi(\mu) - \phi(\hat{\mu}_c) - \left\langle \mu - \hat{\mu}_c, \hat{\theta}_c \right\rangle \leq \rho_c, \end{array} \right. \end{split}$$

where the first equality is from the definition of ℓ_λ and the second equality follows from the linearity of the expectation operator. The last equality follows from the definition of the ambiguity set $\mathbb{B}_{Y|\widehat{x}_c}$ using the ϕ function by Lemma B.3. Because the term $\Psi(\lambda_c)$ does not involve the decision variable μ , it suffices now to consider the optimization problem

$$\sup \left\{ \left\langle -\lambda_c, \mu \right\rangle : \phi(\mu) - \left\langle \mu, \widehat{\theta}_c \right\rangle \le \rho_c + \phi(\widehat{\mu}_c) - \left\langle \widehat{\mu}_c, \widehat{\theta}_c \right\rangle \right\}. \tag{22}$$

Suppose at this moment that $\lambda_c \neq 0$ and $\rho_c > 0$. When $\rho_c > 0$, the feasible set of (22) satisfies the Slater condition because ϕ is a continuous function. Hence, by a strong duality argument, the convex optimization problem (22) is equivalent to

$$\sup_{\mu} \inf_{\gamma \geq 0} \left\langle -\lambda_c, \mu \right\rangle + \gamma(\bar{\rho}_c - \phi(\mu) + \left\langle \mu, \widehat{\theta}_c \right\rangle) = \inf_{\gamma \geq 0} \left\{ \gamma \bar{\rho}_c + \sup_{\mu} \left\langle \mu, \gamma \widehat{\theta}_c - \lambda_c \right\rangle - \gamma \phi(\mu) \right\},$$

where $\bar{\rho}_c \triangleq \rho_c + \phi(\hat{\mu}_c) - \left\langle \hat{\mu}_c, \hat{\theta}_c \right\rangle \in \mathbb{R}$ and the interchange of the supremum and the infimum operators is justified thanks to [9, Proposition 5.3.1]. Consider now the infimum problem on the right hand side of the above equation. If $\gamma=0$, then the inner supremum subproblem on the right hand side is unbounded because $\lambda_c \neq 0$, thus $\gamma=0$ is never an optimal solution to the infimum problem. By utilizing the definition of the conjugate function, one thus deduce that problem (22) is equivalent to

$$\inf_{\gamma>0} \gamma \bar{\rho}_c + (\gamma \phi)^* (\gamma \hat{\theta}_c - \lambda_c) = \inf_{\gamma>0} \gamma \bar{\rho}_c + \gamma \phi^* \left(\hat{\theta}_c - \frac{\lambda_c}{\gamma} \right), \tag{23}$$

where the equality exploits the fact that $(\gamma \phi)^*(\theta) = \gamma \phi^*(\theta/\gamma)$ for any $\gamma > 0$ [13, Table 3.2].

We now show that the reformulation problem (23) is valid when $\rho_c = 0$. Indeed, when $\rho_c = 0$, problem (22) has a unique feasible solution $\widehat{\mu}_c$, thus its optimal value is $\langle -\lambda_c, \widehat{\mu}_c \rangle$. Moreover, in this case, problem (23) becomes

$$\inf_{\gamma>0} \gamma \left[\phi(\widehat{\mu}_c) - \langle \widehat{\mu}_c, \widehat{\theta}_c \rangle + \phi^* \left(\widehat{\theta}_c - \frac{\lambda_c}{\gamma} \right) \right] \\
= \langle -\lambda_c, \widehat{\mu}_c \rangle + \inf_{\gamma>0} \gamma \left[\phi(\widehat{\mu}_c) - \langle \widehat{\mu}_c, \widehat{\theta}_c - \frac{\lambda_c}{\gamma} \rangle + \phi^* \left(\widehat{\theta}_c - \frac{\lambda_c}{\gamma} \right) \right].$$

Notice that the term in the square bracket of the optimization problem on the right hand side is non-negative by the definition of the conjugate function. Thus, the infimum problem over γ admits the optimal value of 0 as γ tends to $+\infty$. As a consequence, when $\rho_c=0$, both problem (22) and (23) have the same optimal value and they are equivalent.

Consider now the situation where $\lambda_c = 0$. In this case, problem (23) becomes

$$\inf_{\gamma>0} \gamma \rho_c + \gamma \left(\phi(\widehat{\mu}_c) - \left\langle \widehat{\mu}_c, \widehat{\theta}_c \right\rangle + \phi^*(\widehat{\theta}_c) \right).$$

By definition of the conjugate function, we have $\phi^*(\widehat{\theta}_c) \geq \langle \widehat{\mu}_c, \widehat{\theta}_c \rangle - \phi(\widehat{\mu}_c)$, and thus, by combining with the fact that $\rho_c \geq 0$, this infimum problem will admit the optimal value of 0. Notice that when $\lambda_c = 0$, the optimal value of problem (22) is also 0. This shows that (23) is equivalent to (22) for any possible value of λ_c . Replacing ϕ^* in (23) by its equivalence Ψ and substituting $\langle \widehat{\mu}_c, \widehat{\theta}_c \rangle - \phi(\widehat{\mu}_c)$ by its equivalence $\Psi(\widehat{\theta}_c)$ complete the reformulation (10).

Proof of Theorem 3.3. By applying Proposition 3.1, the distributionally robust MLE problem (4) can be reformulated as

$$\min_{w \in \mathcal{W}} \max_{\mathbb{Q} \in \mathbb{B}(\widehat{\mathbb{P}})} \mathbb{E}_{\mathbb{Q}} \Big[\ell_{\lambda}(X, Y, w) \Big] = \left\{ \begin{array}{ll} \inf & \alpha + \beta \varepsilon + \beta \sum_{c=1}^{C} \widehat{p}_{c} \exp \left(\frac{t_{c} - \alpha}{\beta} - \rho_{c} - 1 \right) \\ \text{s.t.} & w \in \mathcal{W}, \ t \in \mathbb{R}^{C}, \ \alpha \in \mathbb{R}, \ \beta \in \mathbb{R}_{++} \\ & \sup_{\mathbb{Q}_{Y \mid \widehat{x}_{c}} \in \mathbb{B}_{Y \mid \widehat{x}_{c}}} \mathbb{E}_{\mathbb{Q}_{Y \mid \widehat{x}_{c}}} \left[\ell_{\lambda}(\widehat{x}_{c}, Y, w) \right] \leq t_{c} \ \forall c = 1, \dots, C. \end{array} \right.$$

Using Proposition 3.2 to reformulate each constraint of the above optimization problem leads to the desired result.

C Proofs of Section 4

Proof of Proposition 4.1. Let $\mathbbm{1}$ denote the N dimensional vector of all 1's. Let $\mathrm{KL}(q \parallel p) = \sum_{i=1}^{N} q_i \log(q_i/p_i)$, we have

$$\sup_{\mathbb{Q}: \mathrm{KL}(\mathbb{Q}\|\widehat{\mathbb{P}}^{\mathrm{emp}}) \leq \varepsilon} \mathbb{E}_{\mathbb{Q}}[\ell_{\lambda}(X, Y, w)] = \sup_{q: \mathrm{KL}(q\|\frac{1}{N}\mathbb{1}) \leq \varepsilon} \sum_{i=1}^{N} q_{i} \ell_{\lambda}(\widehat{x}_{i}, \widehat{y}_{i}, w)$$

$$= \sup_{q: \mathrm{KL}(q\|\frac{1}{N}\mathbb{1}) \leq \varepsilon} \sum_{i=1}^{N} q_{i} \left(\Psi(\lambda(w, \widehat{x}_{i})) - \left\langle T(\widehat{y}_{i}), \lambda(w, \widehat{x}_{i}) \right\rangle \right).$$

On the other hand, we note

$$\begin{split} \sup_{\mathbb{Q} \in \mathbb{B}(\widehat{\mathbb{P}})} \mathbb{E}_{\mathbb{Q}} \Big[\ell_{\lambda}(X,Y,w) \Big] &= \sup_{q: \mathrm{KL}(q \parallel \frac{1}{N} \mathbb{1}) \leq \varepsilon} \sum_{i=1}^{N} q_{i} \mathbb{E}_{\mathbb{Q}_{Y \mid \widehat{x}_{i}}} \left[\ell_{\lambda}(\widehat{x}_{i},Y,w) \right] \\ &= \sup_{q: \mathrm{KL}(q \parallel \frac{1}{N} \mathbb{1}) \leq \varepsilon} \sum_{i=1}^{N} q_{i} \left(\Psi(\lambda(w,\widehat{x}_{i})) - \left\langle \nabla \Psi(\widehat{\theta}_{i}), \lambda(w,\widehat{x}_{i}) \right\rangle \right) \\ &= \sup_{q: \mathrm{KL}(q \parallel \frac{1}{N} \mathbb{1}) \leq \varepsilon} \sum_{i=1}^{N} q_{i} \left(\Psi(\lambda(w,\widehat{x}_{i})) - \left\langle T(\widehat{y}_{i}), \lambda(w,\widehat{x}_{i}) \right\rangle \right). \end{split}$$

Therefore the objective functions are the same and the two problems are equivalent.

The proof of Proposition 4.2 relies on the following result.

Lemma C.1. Let $\Delta \subset \mathbb{R}^C$ be a simplex and $\widehat{p} \in \operatorname{int}(\Delta)$ be a probability vector. For any two vectors $\widehat{t}, t^\star \in \mathbb{R}^C$, any vector $\rho \in \mathbb{R}^C_+$ and any scalar $\varepsilon \geq \widehat{p}^\top \rho$, we have

$$\sup \left\{ q^{\top} t^{\star} - \widehat{p}^{\top} \widehat{t} : q \in \Delta, \sum_{c=1}^{C} q_{c} (\log q_{c} - \log \widehat{p}_{c} + \rho_{c}) \leq \varepsilon \right\} \\
\leq \|t^{\star} - \widehat{t}\|_{\infty} + \frac{\sqrt{2\varepsilon}}{\min_{c} \sqrt{\widehat{p}_{c}}} \sqrt{\sum_{c=1}^{C} \widehat{p}_{c} (\widehat{t}_{c} - \overline{t})^{2}},$$

where $\bar{t} = \hat{p}^{\top} \hat{t}$.

Proof of Lemma C.1. Let $\mathbb{1}$ denote the C dimensional vector of 1's, we have

$$\begin{cases} \sup & q^{\top}t^{\star} - \widehat{p}^{\top}\widehat{t} \\ \text{s. t.} & q \in \Delta, \sum_{c=1}^{C} q_{c}(\log q_{c} - \log \widehat{p}_{c} + \rho_{c}) \leq \varepsilon \end{cases}$$

$$= \begin{cases} \sup & q^{\top}(t^{\star} - \widehat{t}) + (q - \widehat{p})^{\top}\widehat{t} \\ \text{s. t.} & q \in \Delta, \sum_{c=1}^{C} q_{c}(\log q_{c} - \log \widehat{p}_{c} + \rho_{c}) \leq \varepsilon \end{cases}$$

$$\leq \begin{cases} \sup & q^{\top}(t^{\star} - \widehat{t}) + (q - \widehat{p})^{\top}\widehat{t} \\ \text{s. t.} & q \in \Delta, \sum_{c=1}^{C} (q_{c} - \widehat{p}_{c})^{2} \leq 2\varepsilon \end{cases}$$

$$\leq \sup_{\|q\|_{1}=1} q^{\top}(t^{\star} - \widehat{t}) + \sup_{c=1}^{C} \{(q - \widehat{p})^{\top}(\widehat{t} - \widehat{t}\mathbb{1}) : \|q - \widehat{p}\|_{2}^{2} \leq 2\varepsilon \}$$

$$\leq \sup_{\|q\|_{1}=1} q^{\top}(t^{\star} - \widehat{t}) + \sup_{c=1}^{C} \sum_{c=1}^{C} \frac{q_{c} - \widehat{p}_{c}}{\sqrt{\widehat{p}_{c}}} \sqrt{\widehat{p}_{c}}(\widehat{t}_{c} - \widehat{t}) : \|q - \widehat{p}\|_{2}^{2} \leq 2\varepsilon \}$$

$$\leq \sup_{\|q\|_{1}=1} q^{\top}(t^{\star} - \widehat{t}) + \sup_{c=1}^{C} \sum_{c=1}^{C} \frac{q_{c} - \widehat{p}_{c}}{\sqrt{\widehat{p}_{c}}} \sqrt{\widehat{p}_{c}}(\widehat{t}_{c} - \widehat{t}) : \|q - \widehat{p}\|_{2}^{2} \leq 2\varepsilon \}$$

$$\leq \sup_{\|q\|_{1}=1} q^{\top}(t^{\star} - \widehat{t}) + \frac{\sqrt{2\varepsilon}}{\min_{c} \sqrt{\widehat{p}_{c}}} \sqrt{\widehat{p}_{c}}(\widehat{t}_{c} - \widehat{t})^{2},$$

where the first inequality follows from Pinsker's inequality [14, Theorem 4.19] and the fact that $\|q-\widehat{p}\|_2^2 \leq \|q-\widehat{p}\|_1^2 = 4\|q-\widehat{p}\|_{TV}^2$, the second inequality follows from the fact that $(q-\widehat{p})^\top \mathbb{1} = 0$ and dropping the constraint $q \in \Delta$, and the last inequality is from Cauchy-Schwarz.

In the last step, we have

$$\sup_{\|q\|_1 = 1} q^{\top} (t^* - \hat{t}) = \|t^* - \hat{t}\|_{\infty},$$

which completes the proof.

We now ready to prove Proposition 4.2.

Proof of Proposition 4.2. Let t^* and \hat{t} be two C-dimensional vectors whose elements are defined as

$$t_c^{\star} = \sup_{\mathbb{Q}_{Y \mid \widehat{x}_c} \in \mathbb{B}_{Y \mid \widehat{x}_c}} \mathbb{E}_{\mathbb{Q}_{Y \mid \widehat{x}_c}} \left[\ell_{\lambda}(\widehat{x}_c, Y, w) \right], \quad \widehat{t}_c = \mathbb{E}_{\widehat{\mathbb{P}}_{Y \mid \widehat{x}_c}} \left[\ell_{\lambda}(\widehat{x}_c, Y, w) \right] \qquad \forall c$$

By Lemma C.1, we find

$$\sup_{\mathbb{Q}\in\mathbb{B}(\widehat{\mathbb{P}})} \mathbb{E}_{\mathbb{Q}}[\ell_{\lambda}(X,Y,w)] - \mathbb{E}_{\widehat{\mathbb{P}}}[\ell_{\lambda}(X,Y,w)] = \begin{cases} \sup & q^{\top}t^{\star} - \widehat{p}^{\top}\widehat{t} \\ \text{s.t.} & q \in \Delta, \sum_{c=1}^{C} q_{c}(\log q_{c} - \log \widehat{p}_{c} + \rho_{c}) \leq \varepsilon \end{cases}$$

$$\leq \|t^{\star} - \widehat{t}\|_{\infty} + \frac{\sqrt{2\varepsilon}}{\min_{c} \sqrt{\widehat{p}_{c}}} \sqrt{\sum_{c=1}^{C} \widehat{p}_{c}(\widehat{t}_{c} - \overline{t})^{2}},$$

where $\bar{t} = \hat{p}^{T} \hat{t}$. In the last step, notice that

$$\sum_{c=1}^{C} \widehat{p}_c(\widehat{t}_c - \overline{t})^2 = \operatorname{Var}_{\widehat{\mathbb{P}}_X} \left(\mathbb{E}_{\widehat{\mathbb{P}}_{Y|X}} \left[\ell_{\lambda}(X, Y, w) \right] \right) \leq \operatorname{Var}_{\widehat{\mathbb{P}}} \left(\ell_{\lambda}(X, Y, w) \right).$$

It now remains to provide the bounds for $||t^* - \hat{t}||_{\infty}$. For any c, let $\lambda_c = \lambda(w, \hat{x}_c)$, we have

$$t_c^{\star} - \widehat{t}_c = \left\{ \begin{array}{ll} \sup & \left\langle \mu - \widehat{\mu}_c, \lambda_c \right\rangle \\ \mathrm{s.\,t.} & \phi(\mu) - \phi(\widehat{\mu}_c) - \left\langle \mu - \widehat{\mu}_c, \widehat{\theta}_c \right\rangle \leq \rho_c. \end{array} \right.$$

Because Ψ has locally Lipschitz continuous gradients, ϕ is locally strongly convex [22, Theorem 4.1]. Moreover, the feasible set S_c of the above problem is compact by Lemma B.4, hence there exists a constant $0 < m_c$ such that

$$\frac{m_c}{2} \|\mu - \widehat{\mu}_c\|_2^2 \le \phi(\mu) - \phi(\widehat{\mu}_c) - \left\langle \mu - \widehat{\mu}_c, \widehat{\theta}_c \right\rangle \quad \forall \mu \in \mathcal{S}_c.$$

Notice that the constants m_c depends only on Ψ and $\widehat{\theta}_c$. Thus, we find

$$t_c^{\star} - \hat{t}_c \leq \sup\left\{\left\langle \mu - \hat{\mu}_c, \lambda_c \right\rangle : m_c \|\mu - \hat{\mu}_c\|_2^2 \leq 2\rho_c\right\} = \sqrt{2\rho_c/m_c} \|\lambda(w, \hat{x}_c)\|_2.$$

By setting $m = \min_c m_c$, we have

$$||t^{\star} - \hat{t}||_{\infty} \le \sqrt{\frac{2 \max_{c} \rho_{c}}{m}} ||\lambda(w, \hat{x}_{c})||_{2}.$$

Combining terms leads to the postulated results.

For any $\widehat{\theta}_c \in \Theta$, $\rho_c \in \mathbb{R}_+$, let $\mathcal{R}_{\widehat{\theta}_c,\rho_c}(w)$ denote the value of the worst-case expected log-loss

$$\mathcal{R}_{\widehat{\theta}_c,\rho_c}(w) = \sup_{\mathbb{Q}_{Y|\widehat{x}_c} \in \mathbb{B}_{Y|\widehat{x}_c}} \mathbb{E}_{\mathbb{Q}_{Y|\widehat{x}_c}} \left[\ell_{\lambda}(\widehat{x}_c,Y,w) \right].$$

Lemma C.2. Suppose that the log-partition function Ψ has locally Lipschitz continuous gradients, that $\Theta = \mathbb{R}^p$ and that $\Theta_c \subset \Theta$ is a compact set. For any fixed $\overline{\rho}_c \in \mathbb{R}_{++}$, there exist constants $0 < m < M < +\infty$ that depend only on Ψ , Θ_c and $\overline{\rho}_c$ such that for any value $\lambda(w, \widehat{x}_c) \in \mathbb{R}^p$ and any radius $\overline{\rho}_c \geq \rho_c \geq 0$

$$\sqrt{2\rho_c/M}\|\lambda(w,\widehat{x}_c)\|_2 \leq \mathcal{R}_{\widehat{\theta}_c,\rho_c}(w) - \mathcal{R}_{\widehat{\theta}_c,0}(w) \leq \sqrt{2\rho_c/m}\|\lambda(w,\widehat{x}_c)\|_2 \qquad \forall \widehat{\theta}_c \in \Theta_c.$$

Proof of Lemma C.2. Consider the set

$$\mathcal{D} \triangleq \{ \widehat{\mu}_c : \exists \widehat{\theta}_c \in \Theta_c \text{ such that } \widehat{\mu}_c = \nabla \Psi(\widehat{\theta}_c) \}$$

and its $\overline{\rho}_c$ -inflated set

$$\mathcal{D}_{\overline{\rho}_c} \triangleq \{\mu: \exists \widehat{\mu}_c \in \mathcal{D} \text{ such that } \phi(\mu) - \phi(\widehat{\mu}_c) - \left\langle \mu - \widehat{\mu}_c, \widehat{\theta}_c \right\rangle \leq \overline{\rho}_c \}.$$

Because Θ_c is compact and $\nabla \Psi$ is a continuous function, \mathcal{D} is compact [1, Theorem 2.34]. Note that we can rewrite $\mathcal{D}_{\overline{\rho}_c}$ as

$$\mathcal{D}_{\overline{\rho}_c} = \{\mu: \exists \widehat{\mu}_c \in \mathcal{D} \text{ such that } \phi(\mu) + \left\langle \mu, -\widehat{\theta}_c \right\rangle \leq \overline{\rho}_c + \phi(\widehat{\mu}_c) - \left\langle \widehat{\mu}_c, \widehat{\theta}_c \right\rangle \}.$$

Let S be temporarily the set

$$S = \left\{ \mu : \phi(\mu) + \inf_{\widehat{\theta}_c \in \Theta_c} \left\langle \mu, -\widehat{\theta}_c \right\rangle \leq \overline{\rho}_c + \sup_{\widehat{\theta}_c \in \Theta_c} \phi(\widehat{\mu}_c) - \left\langle \widehat{\mu}_c, \widehat{\theta}_c \right\rangle < \infty \right\}.$$

We have that $\mathcal{D}_{\overline{\rho}_c} \subseteq S$. Recall the definition of ϕ :

$$\phi: \mu \mapsto \sup_{\theta \in \Theta} \left\{ \left\langle \mu, \theta \right\rangle - \Psi(\theta) \right\}.$$

Therefore $\phi(\cdot)$ is closed, convex and proper. Therefore by [6, Proposition 2.16], $\Theta = \mathbb{R}^p$ implies that $\phi(\cdot)$ is super-coercive, i.e., $\lim_{\|\mu\|_2 \to \infty} \phi(\mu) / \|\mu\|_2 \to \infty$. Thus

$$\lim_{\|\mu\|_2 \to \infty} \phi(\mu) + \inf_{\widehat{\theta}_c \in \Theta_c} \left\langle \mu, -\widehat{\theta}_c \right\rangle \to \infty.$$

Therefore S is bounded, which implies that $\mathcal{D}_{\overline{\rho}_{c}}$ is also bounded.

Since Θ_c is compact, there exists a subsequence $\{\widehat{\theta}_c^{k_n}\}_{n\geq 1}$ such that $\widehat{\theta}_c^{k_n}\to\widehat{\theta}_c^\infty\in\Theta_c$ as $n\to\infty$. Since \mathcal{D}_{ρ_c} is bounded, it suffices to show that \mathcal{D}_{ρ_c} is closed. Choose any sequence $\{\mu^k\}_{k\geq 1}\in\mathcal{D}_{\rho_c}$ such that $\mu^k\to\mu^\infty$ as $k\to\infty$, we want to show that $\mu^\infty\in\mathcal{D}_{\rho_c}$. For each k, since $\mu^k\in\mathcal{D}_{\rho_c}$, there exists $\widehat{\mu}_c^k\in\mathcal{D}$ and $\widehat{\theta}_c^k\in\Theta_c$ such that $\phi(\mu^k)-\phi(\widehat{\mu}_c^k)-\left\langle\mu^k-\widehat{\mu}_c^k,\widehat{\theta}_c^k\right\rangle\leq\rho_c$. Since \mathcal{D} and

 Θ_c are compact, there exists a subsequence $\{k_n\}_{n\geq 1}$ such that $\widehat{\mu}_c^{k_n} \to \widehat{\mu}_c^{\infty}$ and $\widehat{\theta}_c^{k_n} \to \widehat{\theta}_c^{\infty}$ for some $\widehat{\mu}_c^{\infty} \in \mathcal{D}$ and $\widehat{\theta}_c^{\infty} \in \Theta_c$. Since $\widehat{\mu}_c^{k_n} = \nabla \Psi(\widehat{\theta}_c^{k_n})$, by continuity we have $\widehat{\mu}_c^{\infty} = \nabla \Psi(\widehat{\theta}_c^{\infty})$. Note that

$$\phi(\mu^{k_n}) - \phi(\widehat{\mu}_c^{k_n}) - \langle \mu^{k_n} - \widehat{\mu}_c^{k_n}, \widehat{\theta}_c^{k_n} \rangle \le \rho_c,$$

by continuity of ϕ , we have

$$\phi(\mu^{\infty}) - \phi(\widehat{\mu}_c^{\infty}) - \langle \mu^{\infty} - \widehat{\mu}_c^{\infty}, \widehat{\theta}_c^{\infty} \rangle \le \rho_c.$$

Therefore $\mu^{\infty} \in \mathcal{D}_{\rho_c}$ and hence \mathcal{D}_{ρ_c} is closed.

The finite dimensional set $\mathcal{D}_{\overline{\rho}_c}$ is closed and bounded, thus it is compact, and moreover $\mathcal{D} \subseteq \mathcal{D}_{\rho_c}$. The convex hull $\overline{\mathcal{D}}_{\overline{\rho}_c}$ of $\mathcal{D}_{\overline{\rho}_c}$ is also compact [1, Corollary 5.33]. Because Ψ has locally Lipschitz continuous gradients, ϕ is locally strongly convex [22, Theorem 4.1]. Moreover, ϕ is also essentially smooth by Lemma B.2(v). Thus over the set $\overline{\mathcal{D}}_{\overline{\rho}_c}$, there exist constants $0 < m \le M < +\infty$ such that

$$\frac{m}{2}\|\mu-\mu'\|_2^2 \leq \phi(\mu) - \phi(\mu') - \left\langle \mu-\mu',\theta' \right\rangle \leq \frac{M}{2}\|\mu-\mu'\|_2^2 \quad \forall \mu,\mu' \in \overline{\mathcal{D}}_{\overline{\rho}_c}, \mu' = \nabla \Psi(\theta').$$

Notice that the constants m and M depend only on ϕ , and thus on Ψ , $\overline{\rho}_c$ and Θ_c

Denote temporarily the shorthand $\lambda_c = \lambda(w, \widehat{x}_c)$. We have $\mathcal{R}_{\widehat{\theta}_c,0}(w) = \Psi(\lambda_c) - \langle \widehat{\mu}_c, \lambda_c \rangle$, and so

$$\mathcal{R}_{\widehat{\theta}_c, \rho_c}(w) - \mathcal{R}_{\widehat{\theta}_c, 0}(w) = \begin{cases} \sup & \langle \mu - \widehat{\mu}_c, \lambda_c \rangle \\ \text{s. t.} & \phi(\mu) - \phi(\widehat{\mu}_c) - \langle \mu - \widehat{\mu}_c, \widehat{\theta}_c \rangle \leq \rho_c. \end{cases}$$

Because μ and $\widehat{\mu}_c$ are both in $\overline{D}_{\overline{\rho}_c}$, we have

$$\frac{m}{2}\|\mu - \widehat{\mu}_c\|_2^2 \le \phi(\mu) - \phi(\widehat{\mu}_c) - \left\langle \mu - \widehat{\mu}_c, \widehat{\theta}_c \right\rangle \le \frac{M}{2}\|\mu - \widehat{\mu}_c\|_2^2.$$

We now have

$$\mathcal{R}_{\widehat{\theta}_c,\rho_c}(w) - \mathcal{R}_{\widehat{\theta}_c,0}(w) \le \sup\left\{\left\langle \mu - \widehat{\mu}_c, \lambda_c \right\rangle : \|\mu - \widehat{\mu}_c\|_2^2 \le 2\rho_c/m\right\} = \sqrt{2\rho_c/m} \|\lambda_c\|_2.$$

A similar argument leads to the lower bound. This observation completes the proof.

Proof of Theorem 4.3. Without loss of generality consider $W \subseteq \mathbb{R}^q$. For notational simplicity, denote

$$R_{\widehat{\theta},\varepsilon,\rho}(w) = \sup_{\mathbb{Q} \in \mathbb{B}(\widehat{\mathbb{P}})} \mathbb{E}_{\mathbb{Q}} \left[\ell_{\lambda}(X,Y,w) \right].$$

Since $\varepsilon \geq \sum_{c=1}^{C} \widehat{p}_c \rho_c$ with probability going to 1, following the same argument as in the proof of Proposition 4.2, we have that with probability going to 1, for any $w \in \mathcal{W}$,

$$R_{\widehat{\theta},\varepsilon,\varrho}(w) - R_{\widehat{\theta},0,0}(w) \le ||t^{\star} - \widehat{t}||_1 + \sqrt{2\varepsilon}||\widehat{t}||_1,$$

where

$$\|\widehat{t}\|_1 = \sum_{c=1}^C |\mathbb{E}_{\widehat{\mathbb{P}}_Y|\widehat{x}_c}\left[\ell_\lambda(\widehat{x}_c, Y, w)\right]| \quad \text{and} \quad \|t^\star - \widehat{t}\|_1 = \sum_{c=1}^C |\mathcal{R}_{\widehat{\theta}_c, \rho_c}(w) - \mathcal{R}_{\widehat{\theta}_c, 0}(w)|.$$

For each w, since $\widehat{\theta}_c \to \lambda(w_0, \widehat{x}_c)$ in probability, we have $\mathbb{P}(\|\widehat{\theta}_c - \lambda(w_0, \widehat{x}_c)\|_2 > 1) \to 0$. Therefore there exists compact set Θ_c for each c such that $\widehat{\theta}_c$ is contained in Θ_c with probability going to 1. Choose $\overline{\rho}_c = 1$, since $\rho_c \to 0$, we have $\overline{\rho}_c \ge \rho_c$ eventually. Therefore, by Lemma C.2, for each c with probability going to 1

$$|\mathcal{R}_{\widehat{\theta}_c,\rho_c}(w) - \mathcal{R}_{\widehat{\theta}_c,0}(w)| \le \sqrt{2\rho_c/m} \|\lambda(w,\widehat{x}_c)\|_2,$$

where the above constant m can be chosen independent of c due to the finite cardinality assumption of \mathcal{X} . Since the function $\lambda(w, \widehat{x}_c)$ is continuous in w for any \widehat{x}_c , we have $\|\lambda(w, \widehat{x}_c)\|_2$ is bounded for all w ranging over a compact set $W \subset \mathcal{W}$. Thus for each c with probability going to 1, we have

$$\sup_{w \in W} |\mathcal{R}_{\widehat{\theta}_c, \rho_c}(w) - \mathcal{R}_{\widehat{\theta}_c, 0}(w)| \leq \sqrt{2\rho_c/m} \sup_{w \in W} \|\lambda(w, \widehat{x}_c)\|_2.$$

Since $\rho_c \to 0$, we have for each c

$$\sup_{w \in W} |\mathcal{R}_{\widehat{\theta}_c, \rho_c}(w) - \mathcal{R}_{\widehat{\theta}_c, 0}(w)| = o_{\mathbb{P}}(1).$$

Thus $\sup_{w\in W} \|t^* - \widehat{t}\|_1 = o_{\mathbb{P}}(1)$. On the other hand, since $\sup_{w\in W} \mathcal{R}_{\widehat{\theta}_c,0}(w)$ is $O_{\mathbb{P}}(1)$, we have $\sup_{w\in W} \|\widehat{t}\|_1 = O_{\mathbb{P}}(1)$. Therefore as $\varepsilon \to 0$, $\rho_c \to 0$,

$$\sup_{w \in W} |R_{\widehat{\theta}, \varepsilon, \rho}(w) - R_{\widehat{\theta}, 0, 0}(w)| = o_{\mathbb{P}}(1)$$

for any compact set W. Next, since $\widehat{\theta}_c \to \lambda(w_0, \widehat{x}_c)$ in probability, we have by continuous mapping theorem

$$\nabla \Psi(\widehat{\theta}_c) \to \nabla \Psi(\lambda(w_0, \widehat{x}_c))$$
 in probability.

Besides, by the strong law of large number,

$$\widehat{p}_c \to \mathbb{P}(X = \widehat{x}_c)$$
 almost surely.

Recall that

$$\begin{split} R_{\widehat{\theta},0,0}(w) &= \mathbb{E}_{\widehat{\mathbb{P}}}[\ell_{\lambda}(X,Y,w)] = \sum_{c=1}^{C} \widehat{p}_{c} \mathbb{E}_{\widehat{\mathbb{P}}_{Y|\widehat{x}_{c}}}[\ell_{\lambda}(\widehat{x}_{c},Y,w)] \\ &= \sum_{c=1}^{C} \widehat{p}_{c} \left(\Psi(\lambda(w,\widehat{x}_{c})) - \langle \nabla \Psi(\widehat{\theta}_{c}), \lambda(w,\widehat{x}_{c}) \rangle \right). \end{split}$$

Therefore, for each w, we have

$$R_{\widehat{\theta},0,0}(w) \to R(w)$$
 in probability,

where

$$R(w) = \mathbb{E}_{\mathbb{P}}[\ell_{\lambda}(X, Y, w)] = \sum_{c=1}^{C} \mathbb{P}(X = \widehat{x}_{c}) \left(\Psi(\lambda(w, \widehat{x}_{c})) - \left\langle \nabla \Psi(\lambda(w_{0}, \widehat{x}_{c})), \lambda(w, \widehat{x}_{c}) \right\rangle \right).$$

Since for each c,

$$w_0 = \min_{w \in \mathcal{W}} \Psi(\lambda(w, \widehat{x}_c)) - \left\langle \nabla \Psi(\lambda(w_0, \widehat{x}_c)), \lambda(w, \widehat{x}_c) \right\rangle$$

Therefore w_0 solves $\min_{w \in \mathcal{W}} R(w)$. If R(w) admits an unique solution, then clearly w_0 is such a solution. Since $R_{\widehat{\theta}_{0,0}}(\cdot)$ is convex, by [3, Theorem II.1],

$$\sup_{w \in W} |R_{\widehat{\theta},0,0}(w) - R(w)| = o_{\mathbb{P}}(1)$$

for any compact set W. Thus by triangle inequality

$$\sup_{w \in W} |R_{\widehat{\theta}, \varepsilon, \rho}(w) - R(w)| = o_{\mathbb{P}}(1)$$

for any compact set W. Let B denote the unit closed ball in \mathbb{R}^q , then $w_0 + \eta B$ is compact for any $\eta > 0$. Thus $R_{\widehat{\theta},\varepsilon,\rho}(w) - R(w) = o_{\mathbb{P}}(1)$ uniformly over $w_0 + \eta B$. Since R(w) is convex and w_0 is its unique optimal solution, we have

$$\inf_{w \in w_0 + \eta B \setminus \frac{\eta}{2}B} R(w) > R(w_0).$$

Therefore, with probability going to 1,

$$\inf_{w \in w_0 + \frac{\eta}{2}B} R_{\widehat{\theta}, \varepsilon, \rho}(w) < \inf_{w \in w_0 + \eta B \setminus \frac{\eta}{2}B} R_{\widehat{\theta}, \varepsilon, \rho}(w).$$

Thus by convexity of $R_{\widehat{\theta},\varepsilon,\rho}$, also

$$\inf_{w \in w_0 + \frac{\eta}{2}B} R_{\widehat{\theta}, \varepsilon, \rho}(w) < \inf_{w \notin w_0 + \eta B} R_{\widehat{\theta}, \varepsilon, \rho}(w).$$

Thus the solution w^* that solves $\inf_{w \in \mathcal{W}} R_{\widehat{\theta}, \varepsilon, o}(w)$ satisfies

$$\mathbb{P}(\|w^* - w_0\|_2 \le \frac{\eta}{2}) \to 1.$$

Since η is chosen arbitrarily, we conclude that $w^* \to w_0$ in probability.

Proof of Lemma 4.4. Denote

$$W_c = \sqrt{N_c} \left(\frac{\sum_{\widehat{x}_i = \widehat{x}_c} T(\widehat{y}_i)}{N_c} - \mathbb{E}_{f(\cdot \mid \theta_c)} [T(Y)] \right).$$

W.l.o.g. we can assume that $\mathbb{E}_{f(\cdot \mid \theta_c)}[T(Y)] = 0$. We first show the joint convergence

$$(W_1^{\top}, \dots, W_C^{\top})^{\top} \xrightarrow{d} \mathcal{N}(0, G)$$
 as $N \to \infty$,

where G is a block-diagonal matrix with diagonal blocks given by $G_c = \text{Cov}_{f(\cdot \mid \theta_c)}(T(Y)), c = 1, \ldots, C$. Note that

$$N_c/N \to \mathbb{P}(X = \hat{x}_c) > 0$$
 a.s. for each c .

For convenience denote $r_c = \mathbb{P}(X = \hat{x}_c)$. We let

$$\tilde{W_c} = \sqrt{\lfloor r_c N \rfloor} \cdot \frac{\sum_{\widehat{x}_i = \widehat{x}_c} T(\widehat{y}_i)}{\lfloor r_c N \rfloor} = \frac{\sum_{\widehat{x}_i = \widehat{x}_c} T(\widehat{y}_i)}{\sqrt{\lceil r_c N \rceil}}.$$

Let $\left[\sum_{\widehat{x}_i=\widehat{x}_c} T(\widehat{y}_i)\right]_{\lfloor r_c N \rfloor}$ be the sum of the first $\lfloor r_c N \rfloor$ samples of $T(\widehat{y}_i)$ such that $\widehat{x}_i=\widehat{x}_c$. If $N_c < \lfloor r_c N \rfloor$, we add additional $\lfloor r_c N \rfloor - N_c$ independent copies of T(Y) where $Y \sim f(\cdot | \theta_c)$ to the sum $\sum_{\widehat{x}_i=\widehat{x}_c} T(\widehat{y}_i)$, and denote it by $\left[\sum_{\widehat{x}_i=\widehat{x}_c} T(\widehat{y}_i)\right]_{|r_c N \rfloor}$ as well. Denote

$$\bar{W_c} = \frac{\left[\sum_{\widehat{x}_i = \widehat{x}_c} T(\widehat{y}_i)\right]_{\lfloor r_c N \rfloor}}{\sqrt{|r_c N|}}.$$

Note that $\bar{W}_1, \dots, \bar{W}_C$ are independent, by i.i.d central limit theorem

$$(\bar{W}_1^\top, \dots, \bar{W}_C^\top)^\top \xrightarrow{d} \mathcal{N}(0, G)$$
 as $N \to \infty$,

where G is a block-diagonal matrix with $G_c = \operatorname{Cov}_{f(\cdot \mid \theta_c)}(T(Y))$. We next show that

$$\tilde{W}_c - \bar{W}_c = o_{\mathbb{P}}(1).$$

Note that

$$\tilde{W}_c - \bar{W}_c = \frac{\left[\sum_{\widehat{x}_i = \widehat{x}_c} T(\widehat{y}_i)\right]_{\lfloor r_c N \rfloor} - \sum_{\widehat{x}_i = \widehat{x}_c} T(\widehat{y}_i)}{\sqrt{|r_c N|}}.$$

By Chebyshev inequality

$$\begin{split} \mathbb{P}(\|\tilde{W}_c - \bar{W}_c\|_2 > \epsilon) &\leq \frac{\mathbb{E}\left[\left\|\left[\sum_{\widehat{x}_i = \widehat{x}_c} T(\widehat{y}_i)\right]_{\lfloor r_c N \rfloor} - \sum_{\widehat{x}_i = \widehat{x}_c} T(\widehat{y}_i)\right\|_2^2\right]}{\epsilon^2 \lfloor r_c N \rfloor} \\ &= \frac{\mathbb{E}\left[\mathbb{E}\left[\left\|\left[\sum_{\widehat{x}_i = \widehat{x}_c} T(\widehat{y}_i)\right]_{\lfloor r_c N \rfloor} - \sum_{\widehat{x}_i = \widehat{x}_c} T(\widehat{y}_i)\right\|_2^2\right] \left|N_c\right|}{\epsilon^2 \lfloor r_c N \rfloor} \\ &= \frac{\mathbb{E}[\|T(\widehat{y}_i)\|_2^2]}{\epsilon^2} \frac{\mathbb{E}[|\lfloor r_c N \rfloor - N_c|]}{\lfloor r_c N \rfloor}. \end{split}$$

Since $N_c/\lfloor r_c N \rfloor \to 1$ almost surely, by dominated convergence theorem

$$\frac{\mathbb{E}[|\lfloor r_c N \rfloor - N_c|]}{\lfloor r_c N \rfloor} \to 0.$$

Thus

$$\mathbb{P}(\|\tilde{W}_c - \bar{W}_c\|_2 > \epsilon) \to 0$$
 as $N \to \infty$,

which means that $\tilde{W}_c - \bar{W}_c = o_{\mathbb{P}}(1)$. Thus by Slutsky's lemma

$$\left(\tilde{W}_1^{\top}, \dots, \tilde{W}_C^{\top}\right)^{\top} \xrightarrow{d} \mathcal{N}(0, G)$$
 as $N \to \infty$.

Finally, since $W_c = (1 + o_{\mathbb{P}}(1))\tilde{W}_c$, by Slutsky's lemma,

$$(W_1^{\top}, \dots, W_C^{\top})^{\top} \xrightarrow{d} \mathcal{N}(0, G)$$
 as $N \to \infty$.

Now note that

$$\widehat{\theta}_c = (\nabla \Psi)^{-1} \left((N_c)^{-1} \sum_{\widehat{x}_i = \widehat{x}_c} T(\widehat{y}_i) \right)$$

and

$$\theta_c = (\nabla \Psi)^{-1} \left(\mathbb{E}_{f(\cdot \mid \theta_c)} [T(Y)] \right).$$

Also note that the vector-valued function $(\nabla \Psi)^{-1}(\cdot)$ is continuously differentiable at $\mathbb{E}_{f(\cdot \mid \theta_c)}[T(Y)]$, therefore, by the delta method

$$\left(\sqrt{N_1}(\widehat{\theta}_1 - \theta_1)^\top, \dots, \sqrt{N_C}(\widehat{\theta}_C - \theta_C)^\top\right)^\top \xrightarrow{d} D \cdot \mathcal{N}(0, G),$$

where D is a block-diagonal matrix with diagonal elements given by

$$D_c = J(\nabla \Psi)^{-1}(\mathbb{E}_{f(\cdot \mid \theta_c)}[T(Y)])$$

the Jacobian matrix of $(\nabla \Psi)^{-1}$ evaluated at $\mathbb{E}_{f(\cdot \mid \theta_c)}[T(Y)]$. Thus

$$V_c = D_c \operatorname{Cov}_{f(\cdot \mid \theta_c)}(T(Y)) D_c^{\top}.$$

Note that by Lemma B.3, we find

$$\mathrm{KL}(f(\cdot|\theta_c) \parallel f(\cdot|\widehat{\theta}_c)) = \langle \theta_c - \widehat{\theta}_c, \mu_c \rangle + \Psi(\widehat{\theta}_c) - \Psi(\theta_c).$$

Note that Ψ is infinitely-many differentiable, we have the follow Taylor expansion

$$\Psi(\widehat{\theta}_c) - \Psi(\theta_c) = \langle \widehat{\theta}_c - \theta_c, \mu_c \rangle + \frac{1}{2} \langle \widehat{\theta}_c - \theta_c, \nabla^2 \Psi(\theta_c + \eta(\widehat{\theta}_c - \theta_c)) (\widehat{\theta}_c - \theta_c) \rangle,$$

where η is a random variable with values between 0 and 1. Therefore

$$\mathrm{KL}(f(\cdot|\theta_c) \parallel f(\cdot|\widehat{\theta}_c)) = \frac{1}{2} \langle \widehat{\theta}_c - \theta_c, \nabla^2 \Psi (\theta_c + \eta(\widehat{\theta}_c - \theta_c)) (\widehat{\theta}_c - \theta_c) \rangle.$$

Because $\sqrt{N_c}(\widehat{\theta}_c-\theta_c) \xrightarrow{d.} \mathcal{N}(0,V_c)$, and $\nabla^2 \Psi(\,\cdot\,)$ is continuous, we have

$$\nabla^2 \Psi (\theta_c + \eta(\widehat{\theta}_c - \theta_c)) = \nabla^2 \Psi (\theta_c) + o_{\mathbb{P}}(1).$$

Moreover, since we have the joint convergence

$$\left(\sqrt{N_1}(\widehat{\theta}_1 - \theta_1)^\top, \dots, \sqrt{N_C}(\widehat{\theta}_C - \theta_C)^\top\right)^\top \xrightarrow{d} \mathcal{N}(0, V),$$

by continuous mapping theorem

$$\left(N_1 \times \mathrm{KL}(f(\cdot|\theta_1) \parallel f(\cdot|\widehat{\theta}_1)), \dots, N_C \times \mathrm{KL}(f(\cdot|\theta_C) \parallel f(\cdot|\widehat{\theta}_C))\right)^{\top} \xrightarrow{d.} Z \quad \text{as} \quad N \to \infty,$$

where
$$Z = (Z_1, \dots, Z_C)^{\top}$$
 with $Z_c = \frac{1}{2} R_c^{\top} \nabla^2 \Psi(\theta_c) R_c$, $R_c \sim \mathcal{N}(0, V_c)$ and are independent for $c = 1, \dots, C$.

Before proving the result on the worst-case distribution in Theorem 4.5, we first prove the worst-case conditional measure that maximize problem (9).

Proposition C.3 (Worst-case conditional distribution). For any $w \in \mathcal{W}$ and $\rho_c \in \mathbb{R}_{++}$, then the supremum problem (9) is attained by $\mathbb{Q}_{Y|\widehat{x}_c}^{\star} \sim f(\cdot|\theta_c^{\star})$ with $\theta_c^{\star} = \widehat{\theta}_c - \lambda(w,\widehat{x}_c)/\gamma_c^{\star}$, where $\gamma_c^{\star} > 0$ is the solution of the nonlinear algebraic equation

$$\Psi(\widehat{\theta}_c - \gamma^{-1}\lambda(w,\widehat{x}_c)) + \gamma^{-1}\langle\nabla\Psi(\widehat{\theta}_c - \gamma^{-1}\lambda(w,\widehat{x}_c)),\lambda(w,\widehat{x}_c)\rangle = \Psi(\widehat{\theta}_c) - \rho_c.$$
 (24)

Proof of Proposition C.3. Reminding that problem (9) is written as

$$\sup_{\mathbb{Q}_{Y|\widehat{x}_c} \in \mathbb{B}_{Y|\widehat{x}_c}} \mathbb{E}_{\mathbb{Q}_{Y|\widehat{x}_c}} \left[\ell_{\lambda}(\widehat{x}_c, Y, w) \right].$$

In the first step, we show that $\mathbb{Q}_{Y|\widehat{x}_c}^{\star}$ is feasible in problem (9), which means that $\mathbb{Q}_{Y|\widehat{x}_c}^{\star} \in \mathbb{B}_{Y|\widehat{x}_c}$. Indeed, we find that

$$\mathrm{KL}(\mathbb{Q}_{Y|\widehat{x}_c}^{\star} \parallel \widehat{\mathbb{P}}_{Y|\widehat{x}_c}) = -\Psi\left(\widehat{\theta}_c - \frac{\lambda(w, \widehat{x}_c)}{\gamma_c^{\star}}\right) - \frac{1}{\gamma_c^{\star}} \left\langle \nabla \Psi\left(\widehat{\theta}_c - \frac{\lambda(w, \widehat{x}_c)}{\gamma_c^{\star}}\right), \lambda(w, \widehat{x}_c) \right\rangle + \Psi(\widehat{\theta}_c) = \rho_c,$$

where the first equality exploits the expression of the KL divergence between two distributions from the same family in Lemma B.3, and the second equality follows from the fact that γ_c^{\star} solves (24).

Proposition 3.2 asserts that the worst-case conditional expected log-loss problem (9) is equivalent to the convex program (10). Noticing that (24) is the first-order optimality condition of problem (10), thus, by definition, γ_c^{\star} is the minimizer of (10). The objective value of $\mathbb{Q}_{Y|\widehat{x}_c}^{\star}$ in (9) amounts to

$$\begin{split} \mathbb{E}_{\mathbb{Q}_{Y|\hat{x}_{c}}^{\star}}\left[\ell_{\lambda}(\hat{x}_{c},Y,w)\right] &= \Psi(\lambda(w,\hat{x}_{c})) - \left\langle \mathbb{E}_{\mathbb{Q}_{Y|\hat{x}_{c}}^{\star}}[T(Y)],\lambda(w,\hat{x}_{c})\right\rangle \\ &= \Psi(\lambda(w,\hat{x}_{c})) - \left\langle \nabla\Psi\left(\widehat{\theta}_{c} - \frac{\lambda(w,\hat{x}_{c})}{\gamma_{c}^{\star}}\right),\lambda(w,\hat{x}_{c})\right\rangle \\ &= \gamma_{c}^{\star}\left(\rho_{c} - \Psi(\widehat{\theta}_{c})\right) + \gamma_{c}^{\star}\Psi\left(\widehat{\theta}_{c} - \frac{\lambda(w,\hat{x}_{c})}{\gamma_{c}^{\star}}\right) + \Psi(\lambda(w,\hat{x}_{c})), \end{split}$$

where the first equality follows by substituting the expression of ℓ_{λ} and the linearity of the expectation operator, the second equality follows from the convex conjugate relationship between the expectation parameters and the log-partition function Ψ , and the last equality follows from the fact that γ_c^{\star} solves (24). Notice that the last expression coincide with the objective value of (10) evaluated at the optimal solution γ_c^{\star} . This observation implies that $\mathbb{Q}_{Y|\widehat{x}_c}^{\star}$ attains the optimal value in (9). \square

Next, we establish the following result on the optimal solution of the support function h_Q of the set Q defined as in Lemma B.1.

Lemma C.4 (Support point of Q). Let Q be defined as in (16). For any $t \in \mathbb{R}^C$, if there exist $\alpha^* \in \mathbb{R}$ and $\beta^* \in \mathbb{R}_{++}$ that solve the following system of nonlinear algebraic equation

$$\sum_{c=1}^{C} \widehat{p}_c \exp\left(\frac{t_c - \alpha}{\beta} - \rho_c - 1\right) - 1 = 0$$
 (25a)

$$\sum_{c=1}^{C} \widehat{p}_c(t_c - \alpha) \exp\left(\frac{t_c - \alpha}{\beta} - \rho_c - 1\right) - (\varepsilon + 1)\beta = 0$$
 (25b)

then the optimal solution $q^* \in \mathcal{Q}$ that attains $t^\top q^* = h_{\mathcal{Q}}(t)$ is

$$q_c^* = \widehat{p}_c \exp\left(\frac{t_c - \alpha^*}{\beta^*} - \rho_c - 1\right) \quad \forall c = 1, \dots, C.$$
 (25c)

Proof of Lemma C.4. By definition of q^* in (25c), one can verify that $q^* \ge 0$ and that $\sum_{c=}^{C} q_c^* = 1$, where the equality follows from (25a). Moreover,

$$\sum_{c=1}^{C} q_c^{\star} (\log q_c^{\star} - \log \widehat{p}_c + \rho_c) = \sum_{c=1}^{C} \widehat{p}_c \left(\frac{t_c - \alpha^{\star}}{\beta^{\star}} - 1 \right) \exp \left(\frac{t_c - \alpha^{\star}}{\beta^{\star}} - \rho_c - 1 \right)$$
$$= \sum_{c=1}^{C} \widehat{p}_c \left(\frac{t_c - \alpha^{\star}}{\beta^{\star}} \right) \exp \left(\frac{t_c - \alpha^{\star}}{\beta^{\star}} - \rho_c - 1 \right) - 1 = \varepsilon,$$

where the equalities follow from the definition of q^* in (25c), and the equations (25a) and (25b), respectively. This implies that $q^* \in \mathcal{Q}$.

It now remains to show that $t^{\top}q^{\star}=h_{\mathcal{Q}}(t).$ By Lemma B.1, we have

$$h_{\mathcal{Q}}(t) = \begin{cases} & \text{inf} \quad \alpha + \varepsilon \beta + \beta \sum_{c=1}^{C} \widehat{p}_{c} \exp\left(\frac{t_{c} - \alpha}{\beta} - \rho_{c} - 1\right) \\ & \text{s. t.} \quad \alpha \in \mathbb{R}, \ \beta \in \mathbb{R}_{++}. \end{cases}$$

If $(\alpha^*, \beta^*) \in \mathbb{R} \times \mathbb{R}_{++}$ is the solution of (25a)-(25b), then (α^*, β^*) satisfy the Karush-Kuhn-Tucker condition of the above infimum optimization problem, and thus we have

$$h_{\mathcal{Q}}(t) = \alpha^* + \varepsilon \beta^* + \beta^* \sum_{c=1}^{C} \widehat{p}_c \exp\left(\frac{t_c - \alpha^*}{\beta^*} - \rho_c - 1\right).$$

Moreover, we find

$$\sum_{c=1}^{C} t_c q_c^* = \sum_{c=1}^{C} t_c \widehat{p}_c \exp\left(\frac{t_c - \alpha^*}{\beta^*} - \rho_c - 1\right)$$

$$= (\varepsilon + 1)\beta^* + \alpha^* \sum_{c=1}^{C} \widehat{p}_c \exp\left(\frac{t_c - \alpha^*}{\beta^*} - \rho_c - 1\right)$$

$$= \alpha^* + \varepsilon \beta^* + \beta^* \sum_{c=1}^{C} \widehat{p}_c \exp\left(\frac{t_c - \alpha^*}{\beta^*} - \rho_c - 1\right) = h_{\mathcal{Q}}(t),$$

where the first equality follows from the definition of q^* , the second equality follows from (25b) and the third equality follows from (25a). This observation completes the proof.

Proof of Theorem 4.5. It is easy to verify that \mathbb{Q}^{\star} is a probability measure because each $\delta_{\widehat{x}_c}$ and $\mathbb{Q}^{\star}_{Y|\widehat{x}_c}$ is a probability measure, and $\sum_{c=1}^{C}\widehat{p}_c\exp\left((t_c^{\star}-\alpha^{\star})/\beta^{\star}-\rho_c-1\right)=1$ since $\alpha^{\star},\beta^{\star}$ solves

$$\sum_{c=1}^{C} \widehat{p}_c \exp\left(\beta^{-1} (t_c^* - \alpha) - \rho_c - 1\right) - 1 = 0$$
 (26)

$$\sum_{c=1}^{C} \widehat{p}_c(t_c^{\star} - \alpha) \exp\left(\beta^{-1}(t_c^{\star} - \alpha) - \rho_c - 1\right) - (\varepsilon + 1)\beta = 0, \tag{27}$$

If we set $\mathbb{Q}_X^{\star} = \sum_{c=1}^{C} \widehat{p}_c \exp\left((t_c^{\star} - \alpha^{\star})/\beta^{\star} - \rho_c - 1\right) \delta_{\widehat{x}_c}$, then we have

$$\mathbb{Q}^{\star}(\{\widehat{x}_c\} \times A) = \mathbb{Q}_X^{\star}(\{\widehat{x}_c\}) \mathbb{Q}_{Y|\widehat{x}_c}^{\star}(A) \quad \forall A \in \mathcal{F}(\mathcal{Y}), \ \forall c.$$

Moreover, because $\mathbb{Q}_{Y|\widehat{x}_c}^{\star}$ is constructed using Proposition C.3, we have $\mathrm{KL}(\mathbb{Q}_{Y|\widehat{x}_c} \parallel \widehat{\mathbb{P}}_{Y|\widehat{x}_c}) \leq \rho_c$ for all c. Furthermore, we also have

$$\begin{aligned} \operatorname{KL}(\mathbb{Q}_{X}^{\star} \parallel \widehat{\mathbb{P}}_{X}) + \mathbb{E}_{\mathbb{Q}_{X}^{\star}} [\sum_{c=1}^{C} \rho_{c} \mathbb{1}_{\widehat{x}_{c}}(X)] &= \sum_{c=1}^{C} \widehat{p}_{c} \left(\frac{t_{c}^{\star} - \alpha^{\star}}{\beta^{\star}} - 1 \right) \exp \left(\frac{t_{c}^{\star} - \alpha^{\star}}{\beta^{\star}} - \rho_{c} - 1 \right) \\ &= \sum_{c=1}^{C} \widehat{p}_{c} \left(\frac{t_{c}^{\star} - \alpha^{\star}}{\beta^{\star}} \right) \exp \left(\frac{t_{c}^{\star} - \alpha^{\star}}{\beta^{\star}} - \rho_{c} - 1 \right) - 1 = \varepsilon, \end{aligned}$$

where the equalities follow from the construction of \mathbb{Q}_X^* and the equations (26) and (27), respectively. This implies that $\mathbb{Q}^* \in \mathbb{B}(\widehat{\mathbb{P}})$.

It now remains to show that \mathbb{Q}^* is optimal. For any weight w, by the definition of t_c^* , we have

$$t_c^\star = \mathbb{E}_{\mathbb{Q}_{Y|\widehat{x}_c}^\star}\left[\ell_\lambda(\widehat{x}_c,Y,w)\right] = \sup_{\mathbb{Q}_{Y|\widehat{x}}} \ \mathbb{E}_{\mathbb{Q}_{Y|\widehat{x}_c}}\left[\ell_\lambda(\widehat{x}_c,Y,w)\right]$$

We thus find

$$\max_{\mathbb{Q}\in\mathbb{B}(\widehat{\mathbb{P}})} \mathbb{E}_{\mathbb{Q}} \left[\ell_{\lambda}(X,Y,w) \right] = \sup_{\mathbb{Q}_{X}\in\mathbb{B}_{X}} \mathbb{E}_{\mathbb{Q}_{X}} \left[\sup_{\mathbb{Q}_{Y|X}\in\mathbb{B}_{Y|X}} \mathbb{E}_{\mathbb{Q}_{Y|X}} \left[\ell_{\lambda}(X,Y,w) \right] \right] \\
= \sup_{\mathbb{Q}_{X}\in\mathbb{B}_{X}} \mathbb{E}_{\mathbb{Q}_{X}} \left[\sum_{c=1}^{C} t_{c}^{\star} \mathbb{1}_{\widehat{x}_{c}}(X) \right] \\
= \sup_{q\in\mathcal{Q}} q^{\top} t^{\star} \qquad (28) \\
= \sum_{c=1}^{C} \widehat{p}_{c} t_{c}^{\star} \exp\left(\frac{t_{c}^{\star} - \alpha^{\star}}{\beta^{\star}} - \rho_{c} - 1 \right) \qquad (29) \\
= \mathbb{E}_{\mathbb{Q}_{X}^{\star}} \left[\sum_{c=1}^{C} t_{c}^{\star} \mathbb{1}_{\widehat{x}_{c}}(X) \right] \\
= \mathbb{E}_{\mathbb{Q}_{X}^{\star}} \left[\mathbb{E}_{\mathbb{Q}_{Y|X}^{\star}} \left[\ell_{\lambda}(X,Y,w) \right] \right] = \mathbb{E}_{\mathbb{Q}^{\star}} \left[\ell_{\lambda}(X,Y,w) \right].$$

where the set \mathcal{Q} in (28) is defined as in (16). Equality (29) follows from Lemma C.4 and from the definition of α^* and β^* that solve (26)-(27). Equality (30) follows from the definition of \mathbb{Q}_X^* . The proof is completed.

D Auxiliary Results

Lemma D.1 (Locally strongly convex parameter). If Ψ is locally strongly smooth, and at $\widehat{\theta}$, the smoothness parameter is σ , then ϕ is locally strongly convex at $\widehat{\mu} = \nabla \Psi(\widehat{\theta})$ with strongly convex parameter $1/\sigma$ in a sufficiently small neighbourhood of $\widehat{\mu}$.

Proof of Lemma D.1. The proof follows directly from the proof of [22, Theorem 4.1]. By the definition of locally strongly smooth, for some $\Theta' \subseteq \Theta$ neighborhood of $\widehat{\theta}$, we have for $\theta \in \Theta'$

$$\Psi(\theta) \leq \Psi(\widehat{\theta}) + \left\langle \nabla \Psi(\widehat{\theta}), \theta - \widehat{\theta} \right\rangle + \frac{\sigma}{2} \|\theta - \widehat{\theta}\|_2^2.$$

Since $\widehat{\mu} = \nabla \Psi(\widehat{\theta})$ and $\phi(\widehat{\mu}) = \langle \widehat{\mu}, \widehat{\theta} \rangle - \Psi(\widehat{\theta})$, we have

$$\begin{split} \phi(\mu) &= \sup_{\theta \in \Theta} \left(\left\langle \mu, \theta \right\rangle - \Psi(\theta) \right) \\ &\geq \sup_{\theta \in \Theta'} \left(\left\langle \mu, \theta \right\rangle - \Psi(\widehat{\theta}) - \left\langle \widehat{\mu}, \theta - \widehat{\theta} \right\rangle - \frac{\sigma}{2} \|\theta - \widehat{\theta}\|_2^2 \right) \\ &= \left\langle \widehat{\mu}, \widehat{\theta} \right\rangle - \Psi(\widehat{\theta}) + \sup_{\theta \in \Theta'} \left(\left\langle \mu, \theta \right\rangle - \left\langle \widehat{\mu}, \theta \right\rangle - \frac{\sigma}{2} \|\theta - \widehat{\theta}\|_2^2 \right) \\ &= \phi(\widehat{\mu}) + \left\langle \widehat{\theta}, \mu - \widehat{\mu} \right\rangle + \sup_{\theta \in \Theta'} \left(\left\langle \mu - \widehat{\mu}, \theta - \widehat{\theta} \right\rangle - \frac{\sigma}{2} \|\theta - \widehat{\theta}\|_2^2 \right). \end{split}$$

In the last step, note that $\widehat{\theta} = \nabla \phi(\widehat{\mu})$. Taking $\theta - \widehat{\theta} = \alpha(\mu - \widehat{\mu})$ where $\alpha = 1/\sigma$. $\theta \in \Theta'$ if $\mu - \widehat{\mu}$ is sufficiently small. We have

$$\sup_{\theta \in \Theta'} \left(\left\langle \mu - \widehat{\mu}, \theta - \widehat{\theta} \right\rangle - \frac{\sigma}{2} \|\theta - \widehat{\theta}\|_2^2 \right) \ge (\alpha - \frac{\sigma}{2} \alpha^2) \|\mu - \widehat{\mu}\|_2^2 = \frac{1}{2\sigma} \|\mu - \widehat{\mu}\|_2^2.$$

Therefore ϕ is locally strongly convex at $\widehat{\mu}$ with strongly convex parameter $1/\sigma$.

In Proposition 4.2, since Ψ is locally Lipschitz continuous, we have that Ψ is locally strongly smooth with smoothness parameter σ_c at $\hat{\theta}_c$, where σ_c can be chosen as the local Lipschitz constant for a neighborhood around $\hat{\theta}_c$. By Lemma D.1 and the proof of Proposition 4.2, for sufficiently small ρ_c , $c = 1, \ldots, C$, we can choose m explicitly as $m = \min_c 1/\sigma_c$, thus $\kappa_2 = \sqrt{2 \max_c \rho_c \cdot \max_c \sigma_c}$.

References

- [1] C. D. ALIPRANTIS AND K. C. BORDER, *Infinite Dimensional Analysis: A Hitchhiker's Guide*, Springer, 2006.
- [2] S. AMARI, Information Geometry and Its Applications, Springer, 2016.
- [3] P. K. Andersen and R. D. Gill, Cox's regression model for counting processes: A large sample study, Annals of Statistics, 10 (1982), pp. 1100–1120.
- [4] A. BANERJEE, S. MERUGU, I. DHILLON, AND J. GHOSH, *Clustering with Bregman divergences*, Journal of Machine Learning Research, 6 (2005), pp. 1705–1749.
- [5] O. BARNDORFF-NIELSEN, Information and Exponential Families, John Wiley & Sons, 2014.
- [6] H. BAUSCHKE AND J. BORWEIN, Legendre functions and the method or random Bregman projections, Journal of Convex Analysis, 4 (1997), pp. 27–67.
- [7] A. BEN-TAL, D. DEN HERTOG, A. DE WAEGENAERE, B. MELENBERG, AND G. RENNEN, *Robust solutions of optimization problems affected by uncertain probabilities*, Management Science, 59 (2013), pp. 341–357.
- [8] C. BERGE, Topological Spaces: Including a Treatment of Multi-Valued Functions, Vector Spaces, and Convexity, Courier Corporation, 1963.
- [9] D. BERTSEKAS, Convex Optimization Theory, Athena Scientific, 2009.
- [10] D. BERTSIMAS, V. GUPTA, AND N. KALLUS, *Data-driven robust optimization*, Mathematical Programming, 167 (2018), p. 235–292.
- [11] J. BLANCHET, K. MURTHY, AND F. ZHANG, Optimal transport based distributionally robust optimization: Structural properties and iterative schemes, arXiv preprint arXiv:1810.02403, (2018).
- [12] J. BLANCHET, F. ZHANG, Y. KANG, AND Z. HU, A distributionally robust boosting algorithm, in 2019 Winter Simulation Conference, 2019, pp. 3728–3739.
- [13] J. M. BORWEIN AND A. S. LEWIS, Convex Analysis and Nonlinear Optimization, Springer, 2006.
- [14] S. BOUCHERON, G. LUGOSI, AND P. MASSART, Concentration Inequalities: A Nonasymptotic Theory of Independence, OUP Oxford, 2013.
- [15] R. CHRISTENSEN, Log-Linear Models, Springer, 1990.
- [16] V. M. T. DE JONG, M. J. C. EIJKEMANS, B. VAN CALSTER, D. TIMMERMAN, K. G. M. MOONS, E. W. STEYERBERG, AND M. VAN SMEDEN, Sample size considerations and predictive performance of multinomial logistic prediction models, Statistics in Medicine, 38 (2019), pp. 1601–1619.
- [17] A. J. DOBSON AND A. G. BARNETT, An Introduction To Generalized Linear Models, CRC press, 2018.
- [18] A. DOMAHIDI, E. CHU, AND S. BOYD, *ECOS: An SOCP solver for embedded systems*, in 2013 European Control Conference (ECC), IEEE, 2013, pp. 3071–3076.
- [19] D. DUA AND C. GRAFF, UCI machine learning repository, 2017.
- [20] J. C. DUCHI, P. W. GLYNN, AND H. NAMKOONG, Statistics of robust optimization: A generalized empirical likelihood approach, arXiv preprint arXiv:1610.03425, (2016).
- [21] L. FAURY, U. TANIELIAN, F. VASILE, E. SMIRNOVA, AND E. DOHMATOB, *Distributionally robust counterfactual risk minimization*, in AAAI Conference on Artificial Intelligence, 2020.
- [22] R. GOEBEL AND R. T. ROCKAFELLAR, Local strong convexity and local Lipschitz continuity of the gradient of convex functions, Journal of Convex Analysis, 15 (2008), p. 263–270.
- [23] M. GRANT AND S. BOYD, CVX: Matlab software for disciplined convex programming, version 2.1, Mar. 2014.
- [24] R. GRAY, Entropy and Information Theory, Springer, 2011.
- [25] J. M. HILBE, Modeling Count Data, Cambridge University Press, 2014.
- [26] D. W. HOSMER JR, S. LEMESHOW, AND R. X. STURDIVANT, *Applied Logistic Regression*, John Wiley & Sons, 2013.

- [27] Z. HU AND J. HONG, Kullback-Leibler divergence constrained distributionally robust optimization, Available on Optimization Online, (2013).
- [28] P. J. Huber, *Robust estimation of a location parameter*, Annals of Mathematical Statistics, 35 (1964), pp. 73–101.
- [29] D. KUHN, P. M. ESFAHANI, V. A. NGUYEN, AND S. SHAFIEEZADEH-ABADEH, Wasserstein distributionally robust optimization: Theory and applications in machine learning, in Operations Research & Management Science in the Age of Analytics, INFORMS, 2019, pp. 130– 166.
- [30] S. LEE, H. LEE, P. ABBEEL, AND A. Y. NG, *Efficient L1 regularized logistic regression*, in Proceedings of the Twenty-First National Conference on Artificial Intelligence and the Eighteenth Innovative Applications of Artificial Intelligence Conference, 2006, pp. 401–408.
- [31] M. LI AND D. B. DUNSON, Comparing and weighting imperfect models using D-probabilities, Journal of the American Statistical Association, (2019), pp. 1–26.
- [32] MOSEK APS, The MOSEK optimization toolbox. Version 9.2., 2019.
- [33] H. NAMKOONG AND J. C. DUCHI, Stochastic gradient methods for distributionally robust optimization with f-divergences, in Advances in Neural Information Processing Systems 29, 2016, pp. 2208–2216.
- [34] ——, *Variance-based regularization with convex objectives*, in Advances in Neural Information Processing Systems 30, 2017, pp. 2971–2980.
- [35] A. Y. NG, Feature selection, L1 vs. L2 regularization, and rotational invariance, in Proceedings of the 21st International Conference on Machine Learning, 2004, pp. 78–85.
- [36] V. NGUYEN, D. KUHN, AND P. MOHAJERIN ESFAHANI, Distributionally robust inverse covariance estimation: The Wasserstein shrinkage estimator, arXiv preprint arXiv:1805.07194, (2018).
- [37] R. T. ROCKAFELLAR, Convex Analysis, Princeton University Press, 1970.
- [38] P. J. ROUSSEEUW AND M. HUBERT, *Robust statistics for outlier detection*, WIREs Data Mining and Knowledge Discovery, 1 (2011), pp. 73–79.
- [39] S. SHAFIEEZADEH-ABADEH, P. M. ESFAHANI, AND D. KUHN, Distributionally robust logistic regression, in Advances in Neural Information Processing Systems 28, 2015, pp. 1576– 1584.
- [40] S. SHAFIEEZADEH-ABADEH, D. KUHN, AND P. M. ESFAHANI, *Regularization via mass transportation*, Journal of Machine Learning Research, 20 (2019), pp. 1–68.
- [41] N. SI, F. ZHANG, Z. ZHOU, AND J. BLANCHET, *Distributionally robust policy evaluation and learning in offline contextual bandits*, in Proceedings of the 37th International Conference on Machine Learning, 2020.
- [42] M. SION, On general minimax theorems, Pacific Journal of Mathematics, 8 (1958), pp. 171–176.
- [43] N. SRIVASTAVA, G. HINTON, A. KRIZHEVSKY, I. SUTSKEVER, AND R. SALAKHUTDI-NOV, *Dropout: A simple way to prevent neural networks from overfitting*, Journal of Machine Learning Research, 15 (2014), pp. 1929–1958.
- [44] D. STROOCK, Probability Theory: An Analytic View, Cambridge University Press, 2011.
- [45] A. W. VAN DER VAART, Asymptotic Statistics, Cambridge University Press, 2000.
- [46] S. WAGER, S. WANG, AND P. S. LIANG, *Dropout training as adaptive regularization*, in Advances in Neural Information Processing Systems 26, 2013, pp. 351–359.
- [47] S. WANG AND C. MANNING, *Fast dropout training*, in Proceedings of the 30th International Conference on Machine Learning, 2013, pp. 118–126.