DOI: xxx/xxxx

ARTICLE TYPE

Gravitational memory effects in Brans-Dicke theory

Shaoqi Hou*

¹School of Physics and Technology, Wuhan University, Hubei, China

Correspondence

* Email: hou.shaoqi@whu.edu.cn

Present Address

School of Physics and Technology, Wuhan University, Wuhan, Hubei 430072, China

Funding Information

National Natural Science Foundation of China, 11633001 and 11920101003. Strategic Priority Research Program of the Chinese Academy of Sciences, XDB23000000. China Postdoctoral Science Foundation, 2020M672400. There exist gravitational memory effects in Brans-Dicke theory. They are closely related to the Bondi-Metzner-Sachs symmetries living on the null infinity in an isolated system. By studying the asymptotically flat spacetime in Brans-Dicke theory and the asymptotic symmetries, one discovers that the displacement memory effect in the tensor sector is due to the vacuum transition caused by the null energy fluxes penetrating the null infinity, while in the scalar sector, the vacuum transition is due to the angular momentum fluxes. Together with the spin and the center-of-mass memory effects, the displacement memories are constrained by various flux-balance laws.

KEYWORDS:

gravitational memory effect, Brans-Dicke theory, Bondi-Metzner-Sachs symmetry, conservation laws

1 | INTRODUCTION

Gravitational memory effects are fascinating phenomena that occur in general relativity (GR) (Braginsky & Grishchuk (1985);Christodoulou (1991); Thorne (1992);Zel'dovich & Polnarev (1974)). There are several types of memories. The displacement memory refers to the permanent change in the relative distance between two test particles after the passage of gravitational waves (GWs). The spin memory manifests itself in the different propagating times it takes for two test particles, orbiting in a circle in opposite directions, to return to their original positions, when GWs pass by (Pasterski, Strominger, & Zhiboedov (2016)). And the center-of-mass (CM) memory is the lasting shift of the CM of an isolated system due to GWs (Nichols (2018)). There are other memories, e.g., velocity memory effect (Zhang, Duval, Gibbons, & Horvathy (2017a, 2018, 2017b)), none of which will be discussed in this work. Both displacement and spin memories might be detected by interferometers and pulsar timing arrays (Boersma, Nichols, & Schmidt (2020);Hübner, Talbot, Lasky, & Thrane (2020);Johnson, Kapadia, Osborne, Hixon, & Kennefick (2019);Lasky, Thrane, Levin, Blackman, & Chen (2016); Madison (2020); McNeill, Thrane, & Lasky (2017); Seto (2009);

Wang et al. (2015)), but it is difficult to observe the CM memory experimentally (Nichols (2018)).

Memory effects might also exist in modified theories of gravity (Du & Nishizawa (2016); Kilicarslan (2018, 2019); Kilicarslan & Tekin (2019); Lang (2014, 2015)). Brans-Dicke (BD) theory is the simplest modified gravity (Brans & Dicke (1961)), whose action is

$$S = \frac{1}{16\pi G_0} \int d^4x \sqrt{-g} \left(\varphi R - \frac{\omega}{\varphi} \nabla_a \varphi \nabla^a \varphi \right), \quad (1)$$

where ω is a constant, G_0 is the bare gravitational constant. The memory effect in this theory would be the topic of this work. It is found out that there are not only the same memories in BD as those in GR, but also a new memory effect due to the BD scalar field, named S memory in Du & Nishizawa (2016). These memories also have something to do with the asymptotic symmetries of an isolated system, the so-called Bondi-Metzner-Sachs (BMS) group, which is a semi-direct product of the supertranslation group and the Lorentz group as in GR. On the one hand, the supertranslation causes the vacuum transition in the tensor sector, and is responsible for the displacement memory, while the Lorentz transformation switches vacua in the scalar sector, so the S memory takes place. On the other hand, the BMS symmetries imply the existence of supermomenta, angular momenta and Lorentz boost charges at the null infinity *I*. When there are GWs, these

⁰Abbreviations: GR, general relativity; BD, Brans-Dicke theory; GW, gravitational wave; BMS, Bondi-Metzner-Sachs

quantities vary over time, and their variations equal the corresponding fluxes escaping from the source of gravity. The flux-balance laws provide constraints on the memory effects. For more details, please refer to Hou & Zhu (2020a, 2020b); Tahura, Nichols, Saffer, Stein, & Yagi (2020).

This work mainly discusses the displacement memories in the tensor and the scalar sectors. Their relations with the BMS symmetries will be deciphered. Finally, we will compute the constraints on various memory effects using the flux-balance laws. We hope this work will set up the basis for using memory effects to probe the nature of gravity (Koyama (2020)). Recently, the properties of gravity has been studied by the scattering of Dirac spinors and the scalar field Fu, Xing, & Na (2020); Gao, Song, Li, Shan, & Wang (2019). So this work is organized in the following way. Section 2 discusses the asymptotically flat spacetime in BD. Based on that, the BMS symmetry is defined and computed in Sec. 2.1, and then, the "conserved charges" and fluxes are determined in Sec. 2.2. Section 3 focuses on memory effects. There, the displacement memories are analyzed in Sec. 3.1, and Sec. 3.2 is devoted to the spin and the CM memories. In the end, there is a short summary in Sec. 4. In this work, the abstract index notation is used (Wald (1984)), and the speed of light in vacuum is c = 1.

2 | ASYMPTOTICALLY FLAT SPACETIMES

Memory effects also happen in cosmological background (Bonga & Prabhu (2020); Donnay & Giribet (2019)), but here, we will focus on those occuring in an isolated system. At the distances very far away from the source of gravity, the spacetime is nearly Minkowskian, so such kind of spacetime is said to be asymptotically flat. An asymptotically flat spacetime at null infinity in GR can be defined in a coordinate independent manner; see Wald (1984). In BD, one can also propose a similar definition as presented in Hou & Zhu (2020a). But here, we will simply define such a spacetime using the generalized Bondi-Sachs coordinates { u, r, θ, ϕ } ($x^A = \theta, \phi$), in which the metric is

$$ds^2 = g_{uu}du^2 + 2g_{ur}dudr + 2g_{uA}dudx^A + h_{AB}dx^Adx^B.$$
 (2)

One requires that the metric components satisfy the following boundary conditions,

$$g_{uu} = -1 + \mathcal{O}(r^{-1}), \quad g_{ur} = -1 + \mathcal{O}(r^{-1}), \quad (3a)$$

$$g_{uA} = \mathcal{O}(1), \quad h_{AB} = r^2 \gamma_{AB} + \mathcal{O}(r),$$
 (3b)

where γ_{AB} is the round metric on a unit 2-sphere,

$$\gamma_{AB} \mathrm{d}x^A \mathrm{d}x^B = \mathrm{d}\theta^2 + \sin^2\theta \mathrm{d}\phi^2. \tag{4}$$

The scalar field behaves like $\varphi = \varphi_0 + \mathcal{O}(r^{-1})$ with φ_0 a constant. One also imposes the determinant condition,

$$\det(h_{AB}) = r^4 \left(\frac{\varphi_0}{\varphi}\right)^2 \sin^2 \theta, \tag{5}$$

so that *r* becomes the luminosity distance as it approaches the null infinity.

With these conditions, one can obtain the series solutions to the equations of motion, i.e.,

$$\varphi = \varphi_0 + \frac{\varphi_1}{r} + \frac{\varphi_2}{r^2} + \mathcal{O}\left(\frac{1}{r^3}\right), \tag{6a}$$

$$g_{uu} = -1 + \frac{2m + \varphi_1/\varphi_0}{r} + \mathcal{O}\left(\frac{1}{r^2}\right),$$
 (6b)

$$g_{ur} = -1 + \frac{\varphi_1}{\varphi_0 r} + \frac{1}{r^2} \left[\frac{1}{16} \hat{c}_A^B \hat{c}_B^A + \frac{2\omega - 5}{8} \left(\frac{\varphi_1}{\varphi_0} \right)^2 + \frac{\varphi_2}{\varphi_0} \right] \\ + \mathcal{O}\left(\frac{1}{r^3} \right), \tag{6c}$$

$$g_{uA} = \frac{\mathscr{D}_B \hat{c}_A^B}{2} + \frac{2}{3r} \left[N_A + \frac{1}{4} \hat{c}_{AB} \mathscr{D}_C \hat{c}^{BC} - \frac{\varphi_1}{12\varphi_0} \mathscr{D}_B \hat{c}_A^B \right] \\ + \mathcal{O}\left(\frac{1}{r^2}\right), \tag{6d}$$

$$g_{AB} = r^{2} \gamma_{AB} + r \left(\hat{c}_{AB} - \gamma_{AB} \frac{\varphi_{1}}{\varphi_{0}} \right) + \hat{d}_{AB} + \gamma_{AB} \left(\frac{1}{4} \hat{c}_{C}^{D} \hat{c}_{D}^{C} + \frac{\varphi_{1}^{2}}{\varphi_{0}^{2}} - \frac{\varphi_{2}}{\varphi_{0}} \right) + \mathcal{O}\left(\frac{1}{r} \right).$$
(6e)

Here, φ_1 , φ_2 , \hat{c}_{AB} , and \hat{d}_{AB} are functions of u and x^A with $\gamma^{AB}\hat{c}_{AB} = \gamma^{AB}\hat{d}_{AB} = 0$, and \mathcal{D}_A is the covariant derivative for γ_{AB} . $m = m(u, x^A)$ and $N_A = N_A(u, x^B)$ are called the Bondi mass and angular momentum aspects, respectively. Their evolution equations are

$$\dot{m} = -\frac{\mathcal{D}_A \mathcal{D}_B N^{AB}}{4} - \frac{N_{AB} N^{AB}}{8} - \frac{2\omega + 3}{4} \left(\frac{N}{\varphi_0}\right)^2 ,(7a)$$
$$\dot{N}_A = \mathcal{D}_A m + \frac{1}{4} (\mathcal{D}_B \mathcal{D}_A \mathcal{D}_C \hat{c}^{BC} - \mathcal{D}_B \mathcal{D}^B \mathcal{D}_C \hat{c}^C_A)$$
$$- \frac{1}{16} \mathcal{D}_A (N_C^B \hat{c}_B^C) + \frac{1}{4} N_C^B \mathcal{D}_A \hat{c}_B^C$$
$$+ \frac{1}{4} \mathcal{D}_B (N_A^C \hat{c}_C^B - \hat{c}_A^C N_C^B)$$
$$+ \frac{2\omega + 3}{8\varphi_0^2} (\varphi_1 \mathcal{D}_A N - 3N \mathcal{D}_A \varphi_1),$$
(7b)

where $N_{AB} = -\partial \hat{c}_{AB}/\partial u$ is the news tensor, and $N = \partial \varphi_1/\partial u$. When there are no GWs, both N_{AB} and N vanish. So they are called the radiative degrees of freedom. In particular, N_{AB} corresponds to the tensor GW, while N to the scalar GW (Hou, Gong, & Liu (2018)).

At the null infinity, although the spacetime is very similar to the flat one, the symmetry group living there is not simply the Poincaré group. It is a much larger group, instead. In the next subsection, this group will be analyzed. After that, the "conserved charges" and fluxes will be obtained.

2.1 | Bondi-Metzner-Sachs symmetries

The asymptotic symmetries, or BMS symmetries, are diffeomorphisms that preserve the boundary conditions defined in the previous section. An infinitesimal asymptotic symmetry ξ^a is the easiest to be determined by solving the following conditions (Hou & Zhu (2020b))

$$\mathscr{L}_{\xi}g_{rr} = \mathscr{L}_{\xi}g_{rA} = 0, \tag{8a}$$

$$g^{AB}\mathscr{L}_{\xi}g_{AB} = -\frac{2\mathscr{L}_{\xi}\varphi}{\varphi},\tag{8b}$$

$$\mathscr{L}_{\xi}g_{ur} = \mathcal{O}\left(r^{-1}\right), \ \mathscr{L}_{\xi}g_{uA} = \mathcal{O}\left(1\right), \ \mathscr{L}_{\xi}g_{AB} = \mathcal{O}\left(r\right), \quad (8c)$$

$$\mathscr{L}_{\xi}g_{uu} = \mathcal{O}\left(r^{-1}\right),\tag{8d}$$

$$\mathscr{L}_{\xi}\varphi = \mathcal{O}\left(r^{-1}\right). \tag{8e}$$

The result shows that ξ^a is parameterized by two functions $\alpha(x^A)$ and $Y^A(x^B)$, and explicitly given by

ξ

$$f' = f, (9a)$$

$$\xi^{A} = Y^{A} - \frac{\mathscr{D}^{A}f}{r} + \mathcal{O}\left(\frac{1}{r^{2}}\right), \tag{9b}$$

$$\xi^{r} = -\frac{r}{2}\psi + \frac{1}{2}\mathscr{D}^{2}f + \mathcal{O}\left(\frac{1}{r}\right).$$
(9c)

with $f = \alpha + u\psi/2$ and $\psi = \mathcal{D}_A Y^A$. The symmetries parameterized by α are called supertranslations. If $\alpha = \alpha_0 + \alpha_1 \sin \theta \cos \phi + \alpha_2 \sin \theta \sin \phi + \alpha_3 \cos \theta$ with α_i (i = 0, 1, 2, 3)constant, one obtains the usual translation. A generic supertranslation is described by an arbitrary linear combination of spherical harmonics Y_{lm} , so the group of supertranslations, the supertranslation group, is infinitely dimensional. The (global) conformal Killing vector field Y^A , satisfying $\mathcal{L}_Y \gamma_{AB} = \psi \gamma_{AB}$, determines an infinitesimal Lorentz transformation. The BMS group is thus the semi-direct product of the supertranslation group and the Lorentz group as in GR. In particular, the supertranslation group is abelian and the normal subgroup of the BMS group.

The BMS group is much larger than the Poincaré group, as the former is infinitely dimensional, while the later has the dimension of ten. In fact, the BMS group contains an infinite copies of the Poincaré group, which are related to each other through the supertranslation.

2.2 | "Conserved charges" and fluxes

By Noether's theorem, one may define conserved quantities for the BMS symmetries. However, when there are GWs, none of these quantities is exactly a constant of time (u). Their changes are given by the corresponding fluxes.

With the Hamiltonian formalism designed by Wald & Zoupas (2000), the charges and fluxes can be calculated (Hou & Zhu (2020a)). For a supertranslation, one finds the supermomentum,

$$\mathcal{P}_{\alpha}[\mathscr{C}] = \frac{\varphi_0}{4\pi G_0} \oint_{\mathscr{C}} \alpha m \mathrm{d}^2 \mathbf{\Omega}, \qquad (10)$$

where \mathscr{C} represents a cross section on \mathscr{I} , and $d^2 \Omega = \sin \theta d\theta d\phi$. If $\alpha = 1$, one has the Bondi mass, and if α is a linear combination of Y_{lm} with l = 1, one has the spatial momentum. The flux is

$$F_{\alpha}[\mathscr{B}] = \frac{\varphi_0}{16\pi G_0} \int_{\mathscr{B}} \alpha \left[\mathscr{D}_A \mathscr{D}_B N^{AB} + \frac{1}{2} N^B_A N^A_B + (2\omega + 3) \left(\frac{\varphi_1}{\varphi_0}\right)^2 \right] du d^2 \Omega, \quad (11)$$

where \mathscr{B} is a subset of \mathscr{I} . The first term in the square brackets represents the so-called soft part, and the remaining are the hard part. If \mathscr{C} and \mathscr{C}' are the past and the future boundaries of \mathscr{B} , the flux-balance law is

$$F_{\alpha}[\mathscr{B}] = -(\mathcal{P}_{\alpha}[\mathscr{C}'] - \mathcal{P}_{\alpha}[\mathscr{C}]).$$
(12)

This will be useful for constraining the displacement memory in the tensor sector.

For an infinitesimal Lorentz transformation, one writes $Y^A = \mathcal{D}^A \mu + \epsilon^{AB} \mathcal{D}_B v$, then the angular momentum is

$$\mathcal{J}_{v}[\mathscr{C}] = -\frac{\varphi_{0}}{8\pi G_{0}} \oint_{\mathscr{C}} v \epsilon^{AB} \mathscr{D}_{A} N_{B} \mathrm{d}^{2} \mathbf{\Omega}, \qquad (13)$$

and the Lorentz boost charge

$$\begin{aligned} \mathcal{K}_{\mu}[\mathscr{C}] &= -\frac{\varphi_0}{8\pi G_0} \oint_{\mathscr{C}} \quad \mu \left(\mathscr{D}^A N_A + 2um \right. \\ &\left. -\frac{\hat{c}_A^B \hat{c}_B^A}{16} - \frac{2\omega + 3}{8} \frac{\varphi_1^2}{\varphi_0^2} \right) \mathrm{d}^2 \mathbf{\Omega}. \ (14) \end{aligned}$$

In the above relations, μ and v are linear combinations of Y_{lm} with l = 1. The corresponding flux is

$$F_{Y}[\mathscr{B}] = F_{\alpha'}[\mathscr{B}] + \frac{\varphi_{0}}{32\pi G_{0}} \int_{\mathscr{B}} Y^{A} \left[\frac{1}{2} (\hat{c}_{B}^{C} \mathscr{D}_{A} N_{C}^{B} - N_{B}^{C} \mathscr{D}_{A} \hat{c}_{C}^{B}) \right. \\ \left. + \mathscr{D}^{B} (N_{B}^{C} \hat{c}_{AC} - \hat{c}_{B}^{C} N_{AC}) \right. \\ \left. + \frac{2\omega + 3}{\varphi_{0}^{2}} (N \mathscr{D}_{A} \varphi_{1} - \varphi_{1} \mathscr{D}_{A} N) \right] du d^{2} \mathbf{\Omega}, \qquad (15)$$

where $\alpha' = u\psi/2$. The flux-balance law in this case is

$$F_{Y}[\mathscr{B}] = -(\mathcal{Q}_{Y}[\mathscr{C}'] - \mathcal{Q}_{Y}[\mathscr{C}]), \qquad (16)$$

with Q_Y either \mathcal{J}_v or \mathcal{K}_{μ} .

The flux-balance laws are very useful to constrain various memories, which will be discussed in the next section.

3 | MEMORIES

Finally, the memory effect is discussed. Let us first use the obtained metric for an isolated system to compute the geodesic deviation equation, since the ground-based interferometer uses it to detect GWs. Let T^a be the 4-velocity of the test particles, and S^a be the deviation vector. Then one has (Wald (1984))

$$T^{c}\nabla_{c}(T^{b}\nabla_{b}S^{a}) = -R_{cbd}{}^{a}T^{c}S^{b}T^{d}.$$
 (17)

Assume the test particles are placed at a fixed radius r_0 and the fixed directions x_0^A , so these particles are called BMS detectors (Strominger & Zhiboedov (2016)). In general, these particles will be accelerated, but as long as r_0 is very large, they approximately freely fall. One then has $T^a \approx (\partial_u)^a$ at a far distance r_0 . Define orthogonal spatial vectors $(e_{\hat{\theta}})^a = r^{-1}(\partial/\partial\theta)^a$, and $(e_{\hat{\phi}})^a = (r \sin \theta)^{-1}(\partial/\partial\phi)^a$. Then, let $S^a = S^{\hat{A}}(e_{\hat{A}})^a$, and the above equation becomes

$$\ddot{S}^{\hat{A}} \approx -R_{u\hat{B}u}{}^{\hat{A}}S^{\hat{B}},\tag{18}$$

where the electric part of the Riemann tensor $R_{abc}^{\ \ d}$ is

$$R_{u\hat{A}u\hat{B}} = -\frac{1}{2r} \left(\partial_u^2 \hat{c}_{\hat{A}\hat{B}} - \delta_{\hat{A}\hat{B}} \frac{\partial_u^2 \varphi_1}{\varphi_0} \right) + \mathcal{O}\left(\frac{1}{r^2}\right).$$
(19)

Now, integrating eq. (18) twice results in (Hou & Zhu (2020b); Tahura et al. (2020))

$$\Delta S_{\hat{A}} \approx \frac{1}{2r} \left(\Delta \hat{c}_{\hat{A}\hat{B}} - \delta_{\hat{A}\hat{B}} \frac{\Delta \varphi_1}{\varphi_0} \right) S_0^{\hat{B}} + \mathcal{O}\left(\frac{1}{r^2}\right), \qquad (20)$$

in which S_0^B is the initial deviation vector at the retarded time u_0 when there were no GWs, i.e., $N_{AB}(u_0, x^C) = 0$ and $N(u_0, x^A) = 0$. A radiating isolated system will eventually settle down to a state in which no GWs can be emitted, and so N_{AB} and N vanish again. But $S^{\hat{A}}$ may not return to its original value, that is,

$$\Delta S_{\hat{A}} \neq 0. \tag{21}$$

If so, there exists a permanent change in the relative distances between test particles. This is the displacement memory effect. By Eq. (20), there are two contributions to the total displacement memory, one of which is from the tensor part $\Delta \hat{c}_{AB}$, and the other from the scalar part $\Delta \varphi_1$. So we will discuss the two contributions separately in Sec. 3.1. Section 3.2 concentrates on the spin and CM memories, mainly the constraints.

For the following discussion, one writes

$$\hat{\varepsilon}_{AB} = \left(\mathscr{D}_A \mathscr{D}_B - \frac{1}{2} \gamma_{AB} \mathscr{D}^2\right) \Phi + \epsilon_{C(A} \mathscr{D}_{B)} \mathscr{D}^C \Upsilon, \qquad (22)$$

where Φ is the electric part and Υ the magnetic part.

3.1 | Displacement memories

The displacement memory in the tensor sector is very similar to the one in GR. It is related to the vacuum transition caused by the supertranslation as in GR (Strominger & Zhiboedov (2016)). As in GR, a vacuum state in the tensor sector is the spacetime with $\hat{c}_{AB} = (\mathscr{D}_A \mathscr{D}_B - \frac{\gamma_{AB}}{2} \mathscr{D}^2) \Phi$ for some function $\Phi(x^A)$, so $N_{AB} = 0$ (Hou & Žhu (2020b)). This kind of state can be transformed by a supertranslation α to a state with $\hat{c}'_{AB} = (\mathcal{D}_A \mathcal{D}_B - \frac{\gamma_{AB}}{2} \mathcal{D}^2) \Phi'$ and $\Phi' = \Phi - 2\alpha$. Therefore, the new state is also a vacuum, but different from the original one. This observation suggests that there are infinitely many vacua in the tensor sector, and the transition between any pair of them is induced by a supertranslation. Usually, one chooses one of the vacua as the physical one, e.g., $\hat{c}_{AB} = 0$. An infinitesimal Lorentz transformation results in $\delta_Y \hat{c}_{AB}$ = $\mathscr{L}_{Y}\hat{c}_{AB} - \frac{\psi}{2}\hat{c}_{AB}$ which is not a vacuum state, but it preserves the physical vacuum, so the S-matrix and the soft theorem are still Lorentz covariant. The above observation suggests a similar explanation for the displacement memory to the one in GR. A process in question starts with an initial state which is a vacuum described by \hat{c}_{AB} , then the tensor GW penetrates \mathcal{I} and eventually disappears. So the final state is again a vacuum, which is likely a new one \hat{c}'_{AB} . The difference

$$\Delta \hat{c}_{AB} = \hat{c}'_{AB} - \hat{c}_{AB} = \left(\mathscr{D}_A \mathscr{D}_B - \frac{\gamma_{AB}}{2} \mathscr{D}^2 \right) \Delta \Phi, \qquad (23)$$

or equivalently, $\Delta \Phi = \Phi' - \Phi$, measures the memory effect. This is nothing but

$$\Delta \Phi = -2\alpha, \tag{24}$$

i.e., the displacement memory is induced by a certain supertranslation.

The displacement memory in the scalar sector, or S memory, is new. Let the vacuum in the scalar sector be described by $\varphi_1 = \varphi_1(x^A)$ and N = 0. Then one finds out that a supertranslation does not transform it, i.e., $\delta_{\alpha}\varphi_1 = 0$, but a Lorentz generator Y^A changes it according to $\delta_Y\varphi_1 = \mathscr{L}_Y\varphi_1 + \frac{\psi}{2}\varphi_1$. It is interesting to find out that the new state $\varphi'_1 = \varphi_1 + \delta_Y\varphi_1$ is also a vacuum state $(\partial_u \varphi'_1 = 0)$. Therefore, like the displacement memory in the tensor sector, the displacement memory $\Delta\varphi_1$ in the scalar sector is also the vacuum transition due to the Lorentz transformation.

The flux-balance law Eq. (12) can be used to constrain the displacement memory in the tensor sector. In terms of the variation in Φ , one has

$$\oint_{\mathscr{C}} \alpha \mathscr{D}^2 (\mathscr{D}^2 + 2) \Delta \Phi d^2 \Omega = \frac{32\pi G_0}{\varphi_0} (\mathscr{E}_{\alpha} + \Delta \mathcal{P}_{\alpha}).$$
(25)

Here, \mathscr{C}_{α} is Eq. (11) without the first term in the square brackets, that is, it is the null energy fluxes of the tensor and the scalar GWs. The displacement memory in the scalar sector can be constrained by the evolution equation Eq. (7b), i.e.,

$$\Delta \varphi_1^2 = \frac{16\varphi_0^2}{2\omega + 3} \left\{ \frac{1}{32} \Delta (\hat{c}_A^B \hat{c}_B^A) + \mathcal{D}^{-2} \mathcal{D}^A \Delta N_A - \int_{u_i}^{u_f} \mathrm{d}u \left[m + \frac{1}{2} \mathcal{D}^{-2} \mathcal{D}^A J_A \right] \right\}, (26)$$

where

$$J_A = \frac{1}{2} N_C^B \mathcal{D}_A \hat{c}_B^C - \frac{2\omega + 3}{\varphi_0^2} N \mathcal{D}_A \varphi_1, \qquad (27)$$

and \mathcal{D}^{-2} is the inverse operator of \mathcal{D}^2 and is explicitly given in Hou & Zhu (2020b).

From the above discussion, one understands that indeed, the displacement memory in the tensor sector is very similar to the one in GR. For example, the both effects are due to the vacuum transition caused by the supertranslation in the tensor sector, i.e., Eq. (24) (Hou & Zhu (2020b); Strominger & Zhiboedov (2016)). At the same time, Eq. (25) is also similar to the constraint in GR. For instance, Eq. (3.11a) in Compère, Oliveri, & Seraj (2020) is the equation for constraining the displacement memory in GR based on the flux-balance law. However, this equation is a surface integral equation. To obtain the similar form to Eq. (25), one simply integrates it over u. Note that in that expression, Δ corresponds to \mathcal{D}^2 here, T is α , and \mathscr{P}_T is \mathscr{P}_{α} . Finally, I do not consider matter fields, so one may set $\hat{T}_{\mu\nu}$ in Eq. (3.11a) to zero, for the purpose of comparison. There are also two main differences between the displacement memories in BD and that in GR. First, the displacement memory effect in the tensor sector has a new contribution, i.e., the null energy flux of the scalar GW contained in \mathscr{C}_{α} (the last term in Eq. (11)), which is absent in GR. Second, the S memory Eq. (26) never appears in GR, because there does not exist a scalar degree of freedom in GR.

3.2 | Spin and center-of-mass memories

As mentioned in the Introduction, the spin memory can be detected by observing two counter-orbiting particles in a circle. If there are GWs, they will return to their starting points at different times, given that they were released at the same time. This effect is related to the leading term in g_{uA} component (Pasterski et al. (2016)), which does not depend on the scalar field. So the spin memory exists only in the tensor sector, and is very similar to the one in GR.

As in GR, the spin memory should be constrained by the flux-balance law associated with Y^A . However, for this purpose, one allows Y^A be a local conformal Killing vector field for γ_{AB} . Then Y^A has a finite number of singular points on the unit 2-sphere, and the flux-balance law obtained in Sec. 2.2 should be modified, for example, keeping the charges Eq. (13) and (14) while adding to Eq. (15) the following term,

$$\Delta F_{Y}[\mathscr{B}] = \frac{\varphi_{0}}{32\pi G_{0}} \int_{\mathscr{B}} Y^{A} \mathscr{D}^{B}(\mathscr{D}_{A} \mathscr{D}_{C} \hat{c}_{B}^{C} - \mathscr{D}_{B} \mathscr{D}_{C} \hat{c}_{A}^{C}) dud^{2} \Omega$$
$$= \frac{\varphi_{0}}{64\pi G_{0}} \int_{\mathscr{B}} \epsilon_{AB} Y^{A} \mathscr{D}^{B} \mathscr{D}^{2}(\mathscr{D}^{2} + 2) \Upsilon dud^{2} \Omega.$$
(28)

5

Then, the constraint on the spin memory, measured by $\Delta \mathcal{R} = \int du \Upsilon$ (Flanagan & Nichols (2017)), reads,

$$\oint_{\mathscr{C}} v \mathscr{D}^2 \mathscr{D}^2 (\mathscr{D}^2 + 2) \Delta \mathcal{R} d^2 \mathbf{\Omega} = -\frac{32\pi G_0}{\varphi_0} (\Delta \mathcal{J}_v + \mathscr{Q}_v + \mathscr{J}_v), \quad (29)$$

where one defines

$$\mathcal{Q}_{v} = -\frac{\varphi_{0}}{16\pi G_{0}} \int_{\mathscr{B}} v \epsilon_{AB} N^{AC} \hat{c}_{C}^{B} \mathrm{d} u \mathrm{d}^{2} \mathbf{\Omega}, \qquad (30a)$$

$$\mathcal{J}_{v} = \frac{\varphi_{0}}{16\pi G_{0}} \int_{\mathscr{B}} v \epsilon^{AB} \mathscr{D}_{A} J_{B} \mathrm{d} u \mathrm{d}^{2} \mathbf{\Omega}.$$
(30b)

Note that here, v is generally not a linear combination of l = 1 spherical harmonics.

Finally, consider the constraint on the CM memory. One may split $\Phi = \Phi_n + \Phi_o$ such that

$$\oint_{\mathcal{S}} \alpha \mathscr{D}^2 (\mathscr{D}^2 + 2) \Delta \Phi_n \mathrm{d}^2 \mathbf{\Omega} = \frac{32\pi G_0}{\varphi_0} \mathscr{E}_{\alpha}, \qquad (31a)$$

$$\oint_{\mathcal{K}} \alpha \mathscr{D}^2 (\mathscr{D}^2 + 2) \Delta \Phi_o d^2 \mathbf{\Omega} = \frac{32\pi G_0}{\varphi_0} \Delta \mathcal{P}_{\alpha}.$$
 (31b)

Then the CM memory effect is measured by (Nichols (2018); Tahura et al. (2020))

$$\Delta \mathscr{K} = \int_{u_i}^{u_f} u \partial_u \Phi_o \mathrm{d}u, \qquad (32)$$

which is contained in $F_{\alpha'}[\mathscr{B}]$ in Eq. (15), i.e.,

$$F_{\alpha'}[\mathscr{B}] = -\frac{\varphi_0}{64\pi G_0} \oint_{\mathscr{C}} \mu \mathscr{D}^2 \mathscr{D}^2 (\mathscr{D}^2 + 2) \Delta \mathscr{R} d^2 \mathbf{\Omega}.$$
(33)

So the flux-balance law Eq. (16) can be used to obtain,

$$\oint_{\mathscr{C}} \mu \mathscr{D}^2 \mathscr{D}^2 (\mathscr{D}^2 + 2) \Delta \mathscr{K} d^2 \mathbf{\Omega} = \frac{64\pi G_0}{\varphi_0} (\mathscr{J}_{\mu} - \Delta \mathcal{K}'_{\mu}), \quad (34)$$

where one defines

$$\Delta \mathcal{K}'_{\mu} = -\frac{\varphi_0}{8\pi G_0} \oint_{\mathscr{C}} \mu \Delta(\mathscr{D}^A N_A + 2um) \mathrm{d}^2 \mathbf{\Omega}.$$
(35)

Therefore, the CM memory is constrained by Eq. (34), as long as μ is not simply a linear combination of l = 1 spherical harmonics. Since CM memory is related to Φ_o , which is a part of \hat{c}_{AB} , then there does not exist an analogous effect in the scalar sector.

4 | CONCLUSION

This work shows that the memory effect of an isolated system in BD shares some similarities with that in GR, and at the same time, has its own unique features. Both theories share the displacement memory effect for the tensor GW, which is induced by the passage of the null energy fluxes through \mathcal{I} , because the supertranslations transform the degenerate vacua among each other in the tensor sector. However, in BD, there exists the scalar GW. It not only contributes to the tensor displacement memory effect by providing a new energy flux, but also has its own memory effect. The S memory effect is due to the angular momentum fluxes penetrating \mathscr{I} , and degenerate vacua in the scalar sector are transformed to each other by Lorentz transformations. Using flux-balance laws, one obtains the constraints on the displacement memories, the spin memory and the CM memory.

Since the displacement memories in the tensor and the scalar sectors cause the permanent changes in the configuration of the interferometer's arms, the basic ideas for detecting them should be similar to those for detecting GR's displacement memory effect (Boersma et al. (2020); Hübner et al. (2020); Johnson et al. (2019); Lasky et al. (2016); McNeill et al. (2017)). It is also possible to use pulsar timing arrays to detect them as in GR, since the memory waveform generated by a supermassive binary black hole system can be modeled as a step function of time (Seto (2009); Wang et al. (2015)). As in GR, the spin memory effect in BD should also be detected by LISA (Amaro-Seoane et al. (2017); Nichols (2017)), and likewise, Tiangin (Luo et al. (2016)) or Taiji (Hu & Wu (2017)). But the CM memory would be difficult to be detected by the current and even the planed detectors (Nichols (2018)). The smoking gun would be the S memory effect. The constraints from the flux-balance laws are very useful to predict how strong memory effects are, and estimate whether they can be detected by the interferometer or pulsar timing arrays. Memory effect might provide a new method to test the nature of gravity.

ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China under grant Nos.11633001 and 11920101003, and the Strategic Priority Research Program of the Chinese Academy of Sciences, grant No. XDB23000000. This was also a project funded by China Postdoctoral Science Foundation (No. 2020M672400).

REFERENCES

- Amaro-Seoane, P., et al. 2017, 2,
- Boersma, O. M., Nichols, D. A., & Schmidt, P. 2020, *Phys. Rev. D*, *101*(8), 083026. doi:
- Bonga, B., & Prabhu, K. 2020, 9,
- Braginsky, V., & Grishchuk, L. 1985, Sov. Phys. JETP, 62, 427-430.
- Brans, C., & Dicke, R. 1961, Phys. Rev., 124, 925-935. doi:
- Christodoulou,
 D.
 1991,
 Sep,
 Phys.
 Rev.

 Lett.,
 67,
 1486–1489.
 Retrieved
 from

SHAOQI HOU

- https://link.aps.org/doi/10.1103/PhysRevLett.67.1486 doi:
- Compère, G., Oliveri, R., & Seraj, A. 2020, JHEP, 20, 116. doi:
- Donnay, L., & Giribet, G. 2019, Class. Quant. Grav., 36(16), 165005. doi:
- Du, S. M., & Nishizawa, A. 2016, Phys. Rev. D, 94(10), 104063. doi:
- Flanagan, E. E., & Nichols, D. A. 2017, *Phys. Rev. D*, *95*(4), 044002. doi:
- Fu, G. Z., Xing, C. C., & Na, W. 2020, *Eur. Phys. J. C*, 80(6), 582. doi:
- Gao, Z., Song, D., Li, X., Shan, H., & Wang, N. 2019, *Astron. Nachr.*, 340(1-3), 241–246. doi:
- Hou, S., Gong, Y., & Liu, Y. 2018, Eur. Phys. J. C, 78, 378. doi:
- Hou, S., & Zhu, Z.-H. 2020a, 8,
- Hou, S., & Zhu, Z.-H. 2020b, 5,
- Hu, W.-R., & Wu, Y.-L. 2017, *National Science Review*, 4(5), 685-686. Retrieved from http://dx.doi.org/10.1093/nsr/nwx116 doi:
- Hübner, M., Talbot, C., Lasky, P. D., & Thrane, E. 2020, January, *Phys. Rev. D*, 101(2), 023011. doi:
- Johnson, A. D., Kapadia, S. J., Osborne, A., Hixon, A., & Kennefick, D. 2019, *Phys. Rev. D*, 99(4), 044045. doi:
- Kilicarslan, E. 2018, Phys. Rev. D, 98(6), 064048. doi:
- Kilicarslan, E. 2019, Turk. J. Phys., 43(1), 126-134. doi:
- Kilicarslan, E., & Tekin, B. 2019, Eur. Phys. J. C, 79(2), 114. doi:
- Koyama, K. 2020, Phys. Rev. D, 102(2), 021502. doi:
- Lang, R. N. 2014, Phys. Rev. D, 89(8), 084014. doi:
- Lang, R. N. 2015, Phys. Rev. D, 91(8), 084027. doi:
- Lasky, P. D., Thrane, E., Levin, Y., Blackman, J., & Chen, Y. 2016, *Phys. Rev. Lett.*, *117*(6), 061102. doi:
- Luo, J., et al. 2016, Class. Quant. Grav., 33(3), 035010. doi:
- Madison, D. R. 2020, Phys. Rev. Lett., 125(4), 041101. doi:
- McNeill, L. O., Thrane, E., & Lasky, P. D. 2017, *Phys. Rev. Lett.*, 118(18), 181103. doi:
- Nichols, D. A. 2017, Phys. Rev. D, 95(8), 084048. doi:
- Nichols, D. A. 2018, Phys. Rev. D, 98(6), 064032. doi:
- Pasterski, S., Strominger, A., & Zhiboedov, A. 2016, *JHEP*, *12*, 053. doi:
- Seto, N. 2009, Mon. Not. Roy. Astron. Soc., 400, L38. doi:
- Strominger, A., & Zhiboedov, A. 2016, JHEP, 01, 086. doi:
- Tahura, S., Nichols, D. A., Saffer, A., Stein, L. C., & Yagi, K. 2020, 7,
- Thorne, K. S. 1992, Phys. Rev. D, 45(2), 520-524. doi:
- Wald, R. M. 1984, General Relativity. Chicago, IL: University of Chicago Press. doi:
- Wald, R. M., & Zoupas, A. 2000, Phys. Rev. D, 61, 084027. doi:
- Wang, J. B., Hobbs, G., Coles, W. et al. 2015, January, Mon. Not. Roy. Astron. Soc., 446(2), 1657-1671. doi:
- Zel'dovich, Y. B., & Polnarev, A. G. 1974, Sov. Astron., 18, 17.
- Zhang, P.-M., Duval, C., Gibbons, G., & Horvathy, P. 2017a, *Phys. Lett. B*, 772, 743–746. doi:
- Zhang, P.-M., Duval, C., Gibbons, G., & Horvathy, P. 2018, *JCAP*, 05, 030. doi:
- Zhang, P.-M., Duval, C., Gibbons, G. W., & Horvathy, P. A. 2017b, *Phys. Rev. D*, 96(6), 064013. doi: