

An exact $\sin\Theta$ formula for matrix perturbation analysis and its applications

He Lyu, Rongrong Wang

Michigan State University

October 10, 2023

Abstract

In this paper, we establish a useful set of formulae for the $\sin\Theta$ distance between the original and the perturbed singular subspaces. These formulae explicitly show that how the perturbation of the original matrix propagates into singular vectors and singular subspaces, thus providing a direct way of analyzing them. Following this, we derive a collection of new results on SVD perturbation related problems, including a tighter bound on the $\ell_{2,\infty}$ -norm of the singular vector perturbation errors under Gaussian noise, a new stability analysis of the Principal Component Analysis and an error bound on the hard singular value thresholding operator. For the latter two, we consider the most general rectangular matrices with full matrix rank.

1 Introduction

Singular value decomposition (SVD) is a fundamental tool in computational mathematics. Many widely used algorithms in numerical analysis and statistics (e.g., principal component analysis [34, 8, 2], matrix completion [12, 11, 26], matrix denoising [20, 25], community detection [44, 18, 1], graph inference [36, 5], etc.) involve the SVD computation. Since the singular vectors and singular subspaces can be sensitive to noise, a careful study of the stability of SVD is necessary.

For a given matrix A , let $A = U\Sigma V^T$ be its SVD, and $\tilde{A} = \tilde{U}\tilde{\Sigma}\tilde{V}^T$ be the SVD of the noisy version $\tilde{A} = A + \Delta A$. In perturbation analysis, our goal is to characterize the robustness of the left or right singular subspaces under an arbitrary perturbation ΔA .

Classical subspace perturbation results, including Davis-Kahan's theorem [19], Wedin's $\sin\Theta$ theorem [42] and many others (e.g., [35, 21, 22]), provide bounds on the $\sin\Theta$ angles between original and perturbed subspaces. For general symmetric matrices with no statistical assumption on the noise, Davis-Kahan's $\sin\Theta$ bound is already tight and easy to use. However, when the perturbation matrix ΔA is random, Davis-Kahan's bound becomes a random quantity. To solve this problem, deterministic variants of the Davis-Kahan's $\sin\Theta$ theorem were introduced in [43, 10] that are particularly useful for statistical applications. Additionally, various asymptotic bounds on eigenvector perturbations have also been derived in [13, 36, 24, 4].

Recently, it was noticed that in addition to norms of $\sin\Theta$ angles, one may derive perturbation bound under other metrics, which can bring extra benefits. One such metric is the $\ell_{2,\infty}$ -norm of the difference between

the original and perturbed singular vectors. In applications such as clustering, classification, and dimension reduction, the $\ell_{2,\infty}$ metric is more accurate in the sense that it provides finer entry-wise error bounds of the embedded data. In addition, bounds on the $\ell_{2,\infty}$ -norm are possible to be much smaller than those on the $\sin\Theta$ angle [14], which is another benefit of using it. Recently, many interesting results have been derived along this direction, including [3, 15, 17, 14, 2, 8], but the problem is still open.

Besides the perturbation bounds on individual factors (i.e., U, Σ, V) in the SVD, bounds on combinations of the factors are also desired by many applications. For example, in PCA, the target quantity is the PC scores $U\Sigma$. Several works related to perturbation of this quantity exist. In [7], the perturbation error of eigenspaces of kernel PCA is studied. In [2], an ℓ_p stability analysis of PC scores is developed for the hollowed version of PCA, which is the normal PCA with the diagonal entries of the Gram matrix removed before carrying out the SVD. To the best of our knowledge, a tight bound for the PC scores of the vanilla PCA is still missing. Another example that requires perturbation analysis on combinations of factors is singular value truncation [37, 20, 9], where the target quantity whose perturbation we care about is the best rank- r approximation, A_r , of A . Previously, sharp perturbation bounds on A_r only exist for low-rank matrices [31], and that for general full-rank matrices is still missing.

In this paper, we present several new perturbation results including an improved $\ell_{2,\infty}$ -norm bound on the singular vector perturbation under Gaussian noise, a perturbation bound for the hard singular value thresholding operator applied to full-rank matrices, and a useful error bound for the perturbation of the PC scores. These new results are either derived or motivated by the new set of $\sin\Theta$ expressions we shall present in Section 3.

2 Collection of new perturbation results on SVD and its derivatives

Before stating our main mathematical tool in Section 3, we first present the three aforementioned implications, as they might be of independent interests.

2.1 A new bound on $\ell_{2,\infty}$ -norm of the singular vector perturbation

Deriving tight $\ell_{2,\infty}$ -norm bounds for the singular vector perturbation is an active research area in statistics [3, 15, 17, 14]. For many machine learning tasks (e.g., spectral clustering and Principal Component Analysis), the $\ell_{2,\infty}$ -norm provides a better characterization of the embedding quality than the $\sin\Theta$ angle as it is a point-wise metric reflecting the error of individual embedding. Previous literature has presented various bounds on the $\ell_{2,\infty}$ -norm of singular vector perturbation. To list a few, [3] established an $\ell_{2,\infty}$ -norm bound for eigenspaces of symmetric random matrices whose expectations are of low-rank. The result is shown to be useful for analyzing spectral methods on the stochastic block model. [2] developed an $\ell_{2,p}$ analysis for a hollowed PCA, for any $1 \leq p \leq \infty$. Hollowed PCA in the presence of missing data has also been investigated in [8], with a special focus in the case where the number of features is significantly larger than the number of samples. Our problem setting and approach are more closely related to [14], which established an entry-wise singular subspace perturbation bound for low-rank matrices through a Procrustean matrix decomposition.

In this section, we are interested in the case where each entry in the perturbation matrix follows i.i.d. Gaussian distribution. Statistical applications that fall into this regime include Gaussian Mixture Model with isotropic noise covariance matrix [30].

To facilitate the illustration, we introduce some notation. For rectangular matrices A , $\Delta A \in \mathbb{R}^{n \times m}$, we can

write the conformal SVD of the original matrix A and its perturbed version $\tilde{A} = A + \Delta A$ as

$$A = U\Sigma V^T = \begin{pmatrix} U_1 & U_2 \end{pmatrix} \begin{pmatrix} \Sigma_1 & \\ & \Sigma_2 \end{pmatrix} \begin{pmatrix} V_1^T \\ V_2^T \end{pmatrix}, \quad \tilde{A} = \tilde{U}\tilde{\Sigma}\tilde{V}^T = \begin{pmatrix} \tilde{U}_1 & \tilde{U}_2 \end{pmatrix} \begin{pmatrix} \tilde{\Sigma}_1 & \\ & \tilde{\Sigma}_2 \end{pmatrix} \begin{pmatrix} \tilde{V}_1^T \\ \tilde{V}_2^T \end{pmatrix}. \quad (2.1)$$

Here $U_1 \in \mathbb{R}^{n \times r}$, $U_2 \in \mathbb{R}^{n \times (n-r)}$, $V_1 \in \mathbb{R}^{m \times r}$, $V_2 \in \mathbb{R}^{m \times (m-r)}$, $[U_1, U_2] \in \mathbb{R}^{n \times n}$, $[V_1, V_2] \in \mathbb{R}^{m \times m}$ are orthogonal matrices, $\Sigma_1 = \text{diag}\{\sigma_1, \sigma_2, \dots, \sigma_r\} \in \mathbb{R}^{r \times r}$, $\Sigma_2 = \text{diag}\{\sigma_{r+1}, \sigma_{r+2}, \dots, \sigma_{\min\{m, n\}}\} \in \mathbb{R}^{(n-r) \times (m-r)}$, and the singular values are indexed in non-increasing order, i.e., $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r \geq \sigma_{r+1} \geq \dots \geq \sigma_{\min\{m, n\}}$. When $n \neq m$, Σ_2 is rectangular, and the extra columns/rows are padded with 0s. The decomposition of \tilde{A} has a similar structure with non-increasing singular values $\tilde{\Sigma}_1 = \text{diag}\{\tilde{\sigma}_1, \tilde{\sigma}_2, \dots, \tilde{\sigma}_r\} \in \mathbb{R}^{r \times r}$, $\tilde{\Sigma}_2 = \text{diag}\{\tilde{\sigma}_{r+1}, \tilde{\sigma}_{r+2}, \dots, \tilde{\sigma}_{\min\{m, n\}}\} \in \mathbb{R}^{(n-r) \times (m-r)}$.

For a matrix A , the $\ell_{2, \infty}$ -norm is defined as

$$\|A\|_{2, \infty} := \sup_{\|x\|_2=1} \|Ax\|_\infty.$$

One can show that $\|A\|_{2, \infty} = \max_i \|a_i\|_2$, where a_i is the i th row of A .

In dimensionality reduction, to characterize the difference between \tilde{U}_1 and U_1 , a desirable measure is $\min_{Q \in \mathbb{O}_r} \|\tilde{U}_1 - U_1 Q\|_{2, \infty}$, where Q is a rotation matrix and \mathbb{O}_r is the orthogonal matrix group in dimension r . In other words, we consider the difference between \tilde{U}_1 and U_1 after they are maximally aligned by a proper rotation Q .

A well known property about $\min_{Q \in \mathbb{O}_r} \|\tilde{U}_1 - U_1 Q\|_{2, \infty}$ is that it is smaller than a constant multiple of the $\sin\Theta$ angle [10]

$$\min_{Q \in \mathbb{O}_r} \|\tilde{U}_1 - U_1 Q\|_{2, \infty} \leq \min_{Q \in \mathbb{O}_r} \|\tilde{U}_1 - U_1 Q\| \leq \sqrt{2} \|\sin\Theta(U_1, \tilde{U}_1)\|. \quad (2.2)$$

This provides a trivial bound on the $\ell_{2, \infty}$ -norm error, but can be very pessimistic. To see why, we need the following definition of incoherent matrices.

Definition 2.1 (Incoherent). A matrix $U \in \mathbb{R}^{n \times r}$ with orthonormal columns ($n \geq r$) is said to be μ -incoherent ($\mu \geq 1$) if $\|U\|_{2, \infty} \leq \mu \sqrt{\frac{r}{n}}$.

Suppose the perturbation matrix ΔA has i.i.d. Gaussian entries $N(0, \sigma^2)$, and the matrix U_1 of leading singular vector is μ -incoherent. Then the bound (2.2) is pessimistic in that it only gives a bound of $O(1)$, while the bound we are about to provide is $O\left(\frac{r}{\sqrt{n}}\right)$. Explicitly, (2.2) combined with Wedin's $\sin\Theta$ bound yields that, provided

$$c_1 \sigma \sqrt{\max\{n, m\}} \leq \sigma_r(A) - \sigma_{r+1}(A), \quad (\text{gap condition})$$

it holds with high probability that,

$$\min_{Q \in \mathbb{O}_r} \|\tilde{U}_1 - U_1 Q\|_{2, \infty} \leq \sqrt{2} \min \left\{ 1, \frac{c_2 \sqrt{\max\{n, m\}} \sigma}{\sigma_r - \sigma_{r+1}} \right\} \sim O(1). \quad (2.3)$$

Here c_1, c_2 are absolute constants, and the big O notation is with respect to the size variables n and m . The gap condition implies that the noise level σ can be as large as $O(1/\sqrt{\max\{n, m\}})$, so the order of $\sigma \sqrt{\max\{n, m\}}$ is $O(1)$. Hence the bound in (2.3) is $O(1)$.

We show that (2.3) is pessimistic by deriving a bound that is of order $O\left(\frac{r}{\sqrt{n}}\right)$.

Theorem 2.1. Suppose $\tilde{A} = A + \Delta A \in \mathbb{R}^{n \times m}$, $\bar{n} := \max\{n, m\}$, ΔA has i.i.d. $N(0, \sigma^2)$ entries, and assume $\sigma_r(A) - \sigma_{r+1}(A) > 21\sigma\sqrt{\bar{n}}$. Then with probability at least $1 - \frac{c}{n^2}$,

$$\min_{Q \in \mathbb{O}_r} \|\tilde{U}_1 - U_1 Q\|_{2, \infty} \leq c_1 \|U_1\|_{2, \infty} \frac{\sigma^2 \bar{n}}{(\sigma_r(A) - \sigma_{r+1}(A))^2} + c_2 \sigma \frac{R(r, n)}{\sigma_r(A) - \sigma_{r+1}(A)}, \quad (2.4)$$

where

$$R(r, n) = \begin{cases} \sqrt{r} + \sqrt{\log n}, & \text{if } A \text{ is rank } r; \\ r + \sqrt{r \log n}, & \text{else,} \end{cases}$$

and c, c_1, c_2, σ are absolute constants independent of n and m .

The following corollary of Theorem 2.1 may be easier to digest.

Corollary 2.2. Under the same assumption as in Theorem 2.1, if we additionally assume that the matrix U_1 holding the leading r left singular vectors of A is μ_1 -incoherent with some constant μ_1 , then with probability at least $1 - \frac{c}{n^2}$,

$$\min_{Q \in \mathbb{O}_r} \|\tilde{U}_1 - U_1 Q\|_{2, \infty} \leq C \cdot \frac{\mu_1 \sqrt{r} + \sqrt{r \log n} + r}{\sqrt{n}} \sim O\left(\frac{r}{\sqrt{n}}\right),$$

where c and C are absolute constants, and the logarithmic factors are omitted in the big O expression. A similar result holds for the right singular subspace.

For low-rank matrices, Corollary 2.2 can be further improved.

Corollary 2.3. When A is of rank- r , (2.4) reduces to

$$\min_{Q \in \mathbb{O}_r} \|\tilde{U}_1 - U_1 Q\|_{2, \infty} \leq c_1 \|U_1\|_{2, \infty} \frac{\sigma^2 \bar{n}}{\sigma_r^2(A)} + c_2 \sigma \frac{\sqrt{r} + \sqrt{\log n}}{\sigma_r(A)}. \quad (2.5)$$

Under the same assumption as in Corollary 2.2, with probability at least $1 - \frac{c}{n^2}$,

$$\min_{Q \in \mathbb{O}_r} \|\tilde{U}_1 - U_1 Q\|_{2, \infty} \leq C \cdot \frac{\mu_1 \sqrt{r} + \sqrt{r \log n} + \sqrt{r}}{\sqrt{n}} \sim O\left(\sqrt{\frac{r}{n}}\right),$$

where c and C are absolute constants. A similar result holds for the right singular subspace.

Remark 2.4. To get a sense of the tightness of this corollary, we mention the following minimax lower bound derived in Theorem 2.5 [8]. The original result considered the case of missing entries with probability $1 - p$, we plug in $p = 1$ to reduce it to the fully observed case. (2.6) below indicates that when A is of rank- r , $n \asymp m$, and $\sigma_r \asymp \sigma\sqrt{n}$, the result in Corollary 2.3 matches the minimax lower bound $O\left(\frac{1}{\sqrt{n}}\right)$ up to a factor of \sqrt{r} .

Theorem 2.5 (Theorem 3.3 in [8]). Suppose $1 \leq r \leq n/2$, and $(\Delta A)_{ij} \stackrel{i.i.d.}{\sim} N(0, \sigma^2)$. Define

$$\mathcal{M} := \{B \in \mathbb{R}^{n \times m} \mid \text{rank}(B) = r, \sigma_r(B) \in [0.9\sigma_r^*, 1.1\sigma_r^*]\}.$$

Denote by $U(B) \in \mathbb{R}^{n \times r}$ the matrix containing the r left singular vectors of B . Then there exists some universal constant $c_{ub} > 0$ such that

$$\inf_{\hat{U}} \sup_{A \in \mathcal{M}} \mathbb{E} \left[\min_{Q \in \mathbb{O}_r} \|\hat{U} Q - U(A)\|_{2, \infty} \right] \geq c_{ub} \min \left\{ \frac{\sigma^2}{\sigma_r^{*2}} \sqrt{nm} + \frac{\sigma}{\sigma_r^*} \sqrt{n}, 1 \right\} \frac{1}{\sqrt{n}}, \quad (2.6)$$

where the infimum is taken over all estimators for $U(A)$ based on the noisy observation $A + \Delta A$.

	Achieve $O(\sqrt{r/n})$ for rank- r matrices	Do not require $\kappa = O(1)$	Do not require addition assumptions	Achieve $O(r/\sqrt{n})$ for general matrices
[14]	✗	✓	✓	✗
[28]	✓	✓	✗	✗
[3]	✓	✗	✓	✗
[8]	✓	✗	✗	✗
This paper	✓	✓	✓	✓

Table 1: Comparison of result in this paper with existing works about the $\ell_{2,\infty}$ bound. Here we compare these results under the assumptions that ΔA has i.i.d. $N(0, \sigma^2)$ entries, $\|U_1\|_{2,\infty} \leq c\sqrt{r/n}$, where σ, c are constants, and $n \asymp m$. Here $\kappa = \frac{\sigma_1(A)}{\sigma_r(A)}$ is the condition number of A . For simplicity, we ignore $\log n$ in the big O notation.

We take a moment here to make a comparison between several existing works with the results derived in this paper (Theorem 2.1 and its corollaries). Table 1 summarizes the $\ell_{2,\infty}$ -norm bound and requirements in each work. The purpose of the comparison is to show the effectiveness of the derived results under the setting where the perturbation matrix ΔA has i.i.d. Gaussian entries and the matrix A is nearly square ($n \asymp m$). To be fair, we would like to mention that some of the existing results might be better suited for other settings (such as when $m \gg n$).

From the table, we can see that the result in this paper achieves the $O(\sqrt{r/n})$ order upper bound for low-rank matrices and $O(r/\sqrt{n})$ for full-rank matrices. In comparison, previous results in [14] do not achieve the $O(\sqrt{r/n})$ order for the low-rank case under the assumptions as in Corollary 2.3. The result in [28] achieves the same order of accuracy $O(\sqrt{r/n})$ but only for rank- r matrices and under a more restrictive gap condition (below is a simplified version)

$$\sigma_r - \sigma_{r+1} \geq c \left(\min\left\{\frac{\sigma_1}{\sigma_r}, 2r\right\} \sigma \sqrt{n} + \|A\|_{2,\infty} \right),$$

where c is a constant. [3, 15, 17, 23] consider the perturbation of one eigen-vector instead of a set of eigen-vectors. [3] obtained an $O(\sqrt{r/n})$ perturbation bound for low-rank matrices, but for full rank matrices, their bound is $O(1)$. Besides, the bound contains in it the condition number $\tilde{\kappa}(A) = \frac{\sigma_1(A)}{\sigma_r(A) - \sigma_{r+1}(A)}$, which

potentially makes it very large. Likewise, the result in [8] also contains a condition number $\kappa(A) = \frac{\sigma_1(A)}{\sigma_r(A)}$.

In contrast, the condition number does not show up in Theorem 2.1, hence it is more suitable for matrices with large condition numbers. In addition, all previous analyses except for the one in [14] are based on techniques developed for eigen-decomposition. When they are applied to rectangular matrices $A \in \mathbb{R}^{n \times m}$, the matrix needs to be symmetrized, causing the resulting upper bounds potentially depend on both the left and the right singular vectors. In contrast, our bound (2.4) is one-sided, in that the perturbation of U_1 only depends on $\|U_1\|_{2,\infty}$ but not $\|V_1\|_{2,\infty}$.

We make a more detailed comparison between the present Theorem 2.1 and Theorem 4.3 in [14]. For a fair comparison, we restrict ourselves to the setting where both [14] and our results hold, that is, when the data matrix $A \in \mathbb{R}^{n \times m}$ is of rank r and the perturbation matrix ΔA has i.i.d. Gaussian $N(0, \sigma^2)$ entries with some constant σ . Theorem 4.3 in [14] reads if $n \geq m$, $\sigma_r(A) \gtrsim \sigma(n/\sqrt{m})$, then

$$\min_{Q \in \mathbb{O}_r} \|\tilde{U}_1 - U_1 Q\|_{2,\infty} \leq C_r \left(\frac{\log n}{\sigma_r(A)} \right) \left(1 + \frac{m}{\sigma_r(A)} + \frac{\sqrt{m}}{\log n} \|U_1\|_{2,\infty} \right), \quad (2.7)$$

where $C(r) \sim O(\sqrt{r})$. Comparing (2.7) and (2.5), we notice that under the allowable gap condition $\sigma_r(A) \gtrsim \sigma\sqrt{\max\{n, m\}}$ and the incoherence condition $\|U_1\|_{2,\infty} \leq c_r/\sqrt{n}$, the second term in (2.7) can be as large as $O(1)$, which is much larger than our bound $O(\sqrt{r/n})$.

Admittedly, our current result only holds for Gaussian perturbation due to the proof techniques we use. We leave the study of other perturbation types as future work.

2.2 Stability of principal component analysis (PCA)

As one of the arguably most popular tools for data visualization and exploration, PCA is used to extract the main features from a dataset or to reduce the dimensionality of the data [33]. There is a vast literature on the analysis of PCA. Most previous works focused on the consistency of Principal Component directions or eigenvalues [16, 8, 39, 32], while the stability of PC scores (i.e., the projection of data matrix A onto its PC directions, using the notations in this paper, PC scores are given by $U_1\Sigma_1$) is less explored, despite its importance in the analysis of various spectral methods.

There are several relevant works investigating the stability of PC scores, but their analyses were under different settings. For completeness, we include a brief review here. [2] developed an ℓ_p analysis for a hollowed version of PCA, where SVD is conducted on the hollowed gram matrix $G = \mathcal{H}(AA^T)$. Here, A is the data matrix and the operator $\mathcal{H}(\cdot)$ zeros out all diagonal entries of a square matrix. The PC scores are given by $U\Lambda^{\frac{1}{2}}$, where U and Λ are the eigenvector matrix and corresponding eigenvalues of G . Perturbation bounds on PC scores in $\ell_{2,p}$ norm were derived in [2] to characterize entrywise behaviour of PCA. Another line of research studied adjacency spectral embedding (ASE) for random dot product graphs (RDPG), which is closely related to PCA in that they both return a weighted singular vector matrix. Central limit theorems for rows of ASE have been provided in [36, 5].

Different from these previous studies, in this section, we focus on the stability of PC scores of the original PCA algorithm, which does not use the hollowed gram matrix. Given a centered data matrix A and its conformal SVD, PCA returns $U_1\Sigma_1$ (or $V_1\Sigma_1$) as the low dimensional projection into \mathbb{R}^r . Due to the possible similarity among singular values within Σ_1 , the PCA embedding may be subject to rotations. Hence when computing the error, we mode out this rotation and aim to bound $\min_{Q \in \mathbb{O}_r} \|U_1\Sigma_1 - \tilde{U}_1\tilde{\Sigma}_1Q\|$ or $\min_{Q \in \mathbb{O}_r} \|U_1\Sigma_1 - \tilde{U}_1\tilde{\Sigma}_1Q\|_F$, where $\|\cdot\|$ is the spectral norm.

The main difference between these quantities and the $\sin\Theta$ angle between singular subspaces is that U_1 is now multiplied by the corresponding singular values, and it is the perturbation of this product that we want to analyze. Naively, one may expect that the perturbation of $U_1\Sigma_1$ is approximately equal to the perturbation of U_1 times $\|\Sigma_1\|$ plus the perturbation of Σ times $\|U_1\|$, and the perturbation of U_1 can in turn be controlled by the $\sin\Theta$ theorem. This argument leads to

$$\min_{Q \in \mathbb{O}_r} \|U_1\Sigma_1 - \tilde{U}_1\tilde{\Sigma}_1Q\| \leq c \cdot \frac{\sigma_1(A)\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, \quad (2.8)$$

where c is some absolute constant. However, this bound is quite large due to the existence of $\sigma_1(A)$ in the numerator. Noticing that $\sigma_1(A)$ appears in (2.8) because we consider $U_1\Sigma_1$ as a whole, in the following theorem, we show that the perturbed singular vectors corresponding to different singular values actually have different levels of stability, which in turn enables a tighter bound on the PC scores. More specifically, the next theorem shows that the singular vectors associated with larger singular values are more stable.

Theorem 2.6. *For $j = 1, \dots, r$, let $\sin\Theta(\tilde{u}_j, U_1)$ be the $\sin\Theta$ angle between the j th left perturbed singular vector \tilde{u}_j and the leading r -dimensional singular subspace $\text{span}(U_1)$ of A . Then provided that $3\|\Delta A\| \leq$*

$\sigma_r - \sigma_{r+1}$, we have

$$\|\sin\Theta(\tilde{u}_j, U_1)\| \leq \frac{C\|\Delta A\|}{\sigma_j - \sigma_{r+1}}.$$

where C is some universal constant and by definition $\|\sin\Theta(\tilde{u}_j, U_1)\| \equiv \|\tilde{u}_j^T U_2\|$, U_2 is the orthogonal complement of U_1 .

The different levels of stability of singular vectors observed in Theorem 2.6 will help us get rid of the $\sigma_1(A)$ and establish a tighter bound on the PC scores.

Theorem 2.7. $\tilde{A} = A + \Delta A$, $U_1\Sigma_1$ is the PCA embedding of A and $\tilde{U}_1\tilde{\Sigma}_1$ is that of \tilde{A} , we have

$$\begin{aligned} \min_{Q \in \mathbb{O}_r} \|U_1\Sigma_1 - \tilde{U}_1\tilde{\Sigma}_1Q\| &\leq 3\|\Delta A\| + 3\sigma_{r+1} \min\left\{\frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1\right\}, \\ \min_{Q \in \mathbb{O}_r} \|U_1\Sigma_1 - \tilde{U}_1\tilde{\Sigma}_1Q\|_F &\leq \left(2\|(\Delta A)_r\|_F^2 + 3\left(\|(\Delta A)_r\|_F + \|(\Sigma_2)_r\|_F \min\left\{\frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1\right\}\right)^2\right)^{1/2} \\ &\quad + \|(\Delta A)_r\|_F + \|(\Sigma_2)_r\|_F \min\left\{\frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1\right\}. \end{aligned}$$

Here $(\Delta A)_r$ is the best rank- r approximation of ΔA .

The upper bound is tighter than (2.8) and can be used to facilitate the error analysis of PCA-related methods (e.g., [29]).

Remark 2.8. When A is rank- r , the above result reduces to

$$\begin{aligned} \min_{Q \in \mathbb{O}_r} \|U_1\Sigma_1 - \tilde{U}_1\tilde{\Sigma}_1Q\| &\leq 3\|\Delta A\|, \\ \min_{Q \in \mathbb{O}_r} \|U_1\Sigma_1 - \tilde{U}_1\tilde{\Sigma}_1Q\|_F &\leq (\sqrt{5} + 1)\|(\Delta A)_r\|_F. \end{aligned}$$

Remark 2.9. A tight $\ell_{2,p}$ norm perturbation bound of PC scores for hollowed PCA was developed in [2]. However, unlike the previous theorem for vanilla PCA, it seems not possible to eliminate $\sigma_1(A)$ from the bound in hollowed PCA, due to the fact that the hollowed PCA conducts the decomposition on the gram matrix instead of the original data matrix A . In the noisy setting, the noise on the Gram matrix contains the term $A^T\Delta A$, whose norm may reach $O(\sigma_1(A)\|\Delta A\|)$ with σ_1 included in the expression.

2.3 A new stability result on singular value truncation

In addition to studying the perturbations of U_1 and $U_1\Sigma_1$, we also investigate the stability of the hard singular value thresholding operator, which provides the best rank- r approximation of A , i.e., $A_r = U_1\Sigma_1V_1^T$. This operator, also known as singular value truncation, is widely used in matrix completion and matrix denoising for promoting low-rankness or reducing the noise [37, 20, 9, 25]. Let $\tilde{A} = A + \Delta A$ be the noisy matrix, and let \tilde{A}_r denote its best rank- r approximation. We characterize the stability of the hard singular value thresholding operator through a bound on $\|A_r - \tilde{A}_r\|$. Previous works have investigated the stability of truncated SVD [31, 41], and tight error bounds for low-rank matrices have been derived. However, a tight bound for general matrices is still missing in the literature.

For rank- r matrix A with $r < \min\{m, n\}$, the following perturbation result was obtained in [31],

$$\|A - \tilde{A}_r\| \leq 2\|\Delta A\|. \quad (2.9)$$

Since in practice, A may not be exactly rank- r , we hope to establish upper bounds for general full-rank matrices.

We comment that although we can easily derive an upper bound of full-rank matrices from that of the low-rank ones, the resulting bound is not tight. Explicitly, for a full-rank matrix A , A_r is of low rank, so we can apply (2.9) on A_r to get

$$\|A_r - \tilde{A}_r\| = \|A_r - (A + \Delta A)_r\| = \|A_r - (A_r + \tilde{E})_r\| \leq 2\|\tilde{E}\| \leq 2\|\Delta A\| + 2\|A - A_r\| = 2\|\Delta A\| + 2\sigma_{r+1},$$

where $\tilde{E} = \Delta A + A - A_r$ and the first inequality used (2.9).

Apparently, this bound is not optimal as it does not shrink to 0 when $\Delta A \rightarrow 0$. This then motivates us to establish the following tighter bound.

Theorem 2.10 (Perturbation result on singular value truncation). *Let $A \in \mathbb{R}^{n \times m}$ be any $n \times m$ matrix and $\tilde{A} = A + \Delta A$ be its noisy version. Denote by A_r and \tilde{A}_r their rank- r thresholding with all but the first r singular values set to 0. Let σ_i be the i th largest singular value of A and Σ_2 be the diagonal matrix containing $\sigma_{r+1}, \dots, \sigma_{\min}$ (the $(r+1)$ 'th to the last singular values of A) on the diagonal. Then*

$$\|A_r - \tilde{A}_r\| \leq 2\|\Delta A\| + 2\sigma_{r+1} \min \left\{ \frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1 \right\}, \quad (2.10)$$

$$\|A_r - \tilde{A}_r\|_F \leq \left(2\|(\Delta A)_r\|_F^2 + 3 \left(\|(\Delta A)_r\|_F + \|(\Sigma_2)_r\|_F \min \left\{ \frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1 \right\} \right)^2 \right)^{1/2}. \quad (2.11)$$

This error bound has exactly the same form as the PCA perturbation bound established in the previous section, except that here A_r and \tilde{A}_r do not differ by a rotation. Intuitively, this indicates that the noise-induced rotation on \tilde{U}_1 and that on \tilde{V}_1^T can essentially cancel with each other.

Remark 2.11. When A is a rank- r matrix, the bound in Theorem 2.10 reduces to the result in [31]:

$$\begin{cases} \|A_r - \tilde{A}_r\| \leq 2\|\Delta A\|, \\ \|A_r - \tilde{A}_r\|_F \leq \sqrt{5}\|(\Delta A)_r\|_F. \end{cases} \quad (2.12)$$

3 Closed-form expression of $\sin\Theta$ distance between two singular spaces

The several new results presented in the previous section are derived either directly or indirectly from a set of $\sin\Theta$ formulae we shall establish in this section. In other words, these $\sin\Theta$ formulae serve as useful tools to analyze SVD based perturbation problems.

3.1 First order equivalent expressions of the $\sin\Theta$ distance

Following the same notation as in Section 2.1, our goal is to compute the exact expressions of perturbation angles of the leading left singular subspace U_1 under noise ΔA .

For two matrices $U_1, \tilde{U}_1 \in \mathbb{R}^{n \times r}$ with orthonormal columns, let the singular values of $U_1^T \tilde{U}_1$ be $\gamma_1 \geq \gamma_2 \geq \dots \geq \gamma_r \geq 0$, then $\cos^{-1} \gamma_i$, $i = 1, \dots, r$ are the principal angles, and the $\sin\Theta$ matrix is the following diagonal matrix

$$\sin\Theta(U_1, \tilde{U}_1) = \text{diag}\{\sin \cos^{-1}(\gamma_1), \sin \cos^{-1}(\gamma_2), \dots, \sin \cos^{-1}(\gamma_r)\}.$$

The angles are usually measured under either the spectral norm $\|\sin\Theta(U_1, \tilde{U}_1)\|$ or the Frobenius norm $\|\sin\Theta(U_1, \tilde{U}_1)\|_F$. It is well known that (e.g., [10, 27])

$$\|\sin\Theta(U_1, \tilde{U}_1)\| = \|U_2^T \tilde{U}_1\| = \|\tilde{U}_2^T U_1\|, \quad (3.1)$$

$$\|\sin\Theta(U_1, \tilde{U}_1)\|_F = \|U_2^T \tilde{U}_1\|_F = \|\tilde{U}_2^T U_1\|_F, \quad (3.2)$$

where U_2 is the orthogonal complement of U_1 as defined in (2.1).

(3.1) and (3.2) indicate that the matrices $U_2^T \tilde{U}_1$ and $\tilde{U}_2^T U_1$ are key intermediate quantities to bound the $\sin\Theta$ angles. In the following theorem, we provide useful expressions of these key quantities.

Theorem 3.1 (Angular perturbation formula). *Let $A, \tilde{A} = A + \Delta A$ be two $n \times m$ matrices and their conformal SVDs are defined as (2.1). The rank of A is at least r . Assume there is a gap between the r th and the $(r+1)$ th singular values, i.e., $\sigma_r - \tilde{\sigma}_{r+1} > 0$ and $\tilde{\sigma}_r - \sigma_{r+1} > 0$. Then the following expressions hold:*

$$\begin{aligned} U_1^T \tilde{U}_2 &= F_U^{12} \circ (U_1^T (\Delta A) \tilde{V}_2 \tilde{\Sigma}_2^T + \Sigma_1 V_1^T (\Delta A)^T \tilde{U}_2), \\ U_2^T \tilde{U}_1 &= F_U^{21} \circ (U_2^T (\Delta A) \tilde{V}_1 \tilde{\Sigma}_1^T + \Sigma_2 V_2^T (\Delta A)^T \tilde{U}_1), \\ V_1^T \tilde{V}_2 &= F_V^{12} \circ (\Sigma_1^T U_1^T (\Delta A) \tilde{V}_2 + V_1^T (\Delta A)^T \tilde{U}_2 \tilde{\Sigma}_2), \\ V_2^T \tilde{V}_1 &= F_V^{21} \circ (\Sigma_2^T U_2^T (\Delta A) \tilde{V}_1 + V_2^T (\Delta A)^T \tilde{U}_1 \tilde{\Sigma}_1). \end{aligned} \quad (3.3)$$

More specifically, the assumption $\sigma_r - \tilde{\sigma}_{r+1} > 0$ is required for the first and the third expressions of (3.3) to hold, and $\tilde{\sigma}_r - \sigma_{r+1} > 0$ is required for the second and the last expressions to hold. Here \circ means the Hadamard product, or element-wise product between two matrices. $F_U^{12} \in \mathbb{R}^{r \times (n-r)}$ has entries $(F_U^{12})_{i,j} = \frac{1}{\tilde{\sigma}_{j+r}^2 - \sigma_i^2}$, $1 \leq i \leq r$, $1 \leq j \leq n-r$; $F_U^{21} \in \mathbb{R}^{(n-r) \times r}$ has entries $(F_U^{21})_{i,j} = \frac{1}{\tilde{\sigma}_j^2 - \sigma_{i+r}^2}$, $1 \leq i \leq n-r$, $1 \leq j \leq r$. Similarly, $F_V^{12} \in \mathbb{R}^{r \times (m-r)}$ has entries $(F_V^{12})_{i,j} = \frac{1}{\tilde{\sigma}_{j+r}^2 - \sigma_i^2}$, $1 \leq i \leq r$, $1 \leq j \leq m-r$; and $F_V^{21} \in \mathbb{R}^{(m-r) \times r}$ with entries $(F_V^{21})_{i,j} = \frac{1}{\tilde{\sigma}_j^2 - \sigma_{i+r}^2}$, $1 \leq i \leq m-r$, $1 \leq j \leq r$. Here if $i > \min\{n, m\}$, we enforce σ_i and $\tilde{\sigma}_i$ to be 0.

Taking the spectral norm on both hand sides of (3.3) gives us the following new expression of the $\sin\Theta$ distance.

Corollary 3.2. *If the condition in Theorem 3.1 is satisfied, then the $\sin\Theta$ distances between the r leading singular spaces of the original and the perturbed matrices satisfy*

$$\begin{aligned} \|\sin\Theta(U_1, \tilde{U}_1)\| &= \|F_U^{12} \circ (U_1^T (\Delta A) \tilde{V}_2 \tilde{\Sigma}_2^T + \Sigma_1 V_1^T (\Delta A)^T \tilde{U}_2)\| \\ &= \|F_U^{21} \circ (U_2^T (\Delta A) \tilde{V}_1 \tilde{\Sigma}_1^T + \Sigma_2 V_2^T (\Delta A)^T \tilde{U}_1)\|, \\ \|\sin\Theta(V_1, \tilde{V}_1)\| &= \|F_V^{12} \circ (\Sigma_1^T U_1^T (\Delta A) \tilde{V}_2 + V_1^T (\Delta A)^T \tilde{U}_2 \tilde{\Sigma}_2)\| \\ &= \|F_V^{21} \circ (\Sigma_2^T U_2^T (\Delta A) \tilde{V}_1 + V_2^T (\Delta A)^T \tilde{U}_1 \tilde{\Sigma}_1)\|. \end{aligned}$$

Remark 3.3. In the expressions of corollary 3.2, the singular value gaps are contained in the terms F_U^{21} , F_U^{12} , F_V^{21} , F_V^{12} as denominators. In this sense, (3.3) conveys the same insight as Wedin's $\sin\Theta$ theorem.

Everything else in the right hand sides of Corollary 3.2 is straightforward to bound except perhaps for the Hadamard products. The following lemma shows that the Hadamard product is also relatively easy to treat.

Lemma 3.4. *Assume $\sigma_r - \tilde{\sigma}_{r+1} > 0$, $\tilde{\sigma}_r - \sigma_{r+1} > 0$, let F_U^{12} , F_U^{21} , Σ_1 , $\tilde{\Sigma}_1$, Σ_2 , $\tilde{\Sigma}_2$, be the same as in Theorem 3.1 and let $H_1 \in \mathbb{R}^{(n-r) \times r}$, $H_2 \in \mathbb{R}^{(m-r) \times r}$, $H_3 \in \mathbb{R}^{r \times (m-r)}$, $H_4 \in \mathbb{R}^{r \times (n-r)}$ be some arbitrary matrices. Then*

$$\|F_U^{21} \circ (H_1 \tilde{\Sigma}_1)\| \leq \frac{\tilde{\sigma}_r}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} \|H_1\|, \quad \|F_U^{21} \circ (\Sigma_2 H_2)\| \leq \frac{\sigma_{r+1}}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} \|H_2\|, \quad (3.4)$$

$$\|F_U^{12} \circ (H_3 \tilde{\Sigma}_2^T)\| \leq \frac{\tilde{\sigma}_{r+1}}{\sigma_r^2 - \tilde{\sigma}_{r+1}^2} \|H_3\|, \quad \|F_U^{12} \circ (\Sigma_1 H_4)\| \leq \frac{\sigma_r}{\sigma_r^2 - \tilde{\sigma}_{r+1}^2} \|H_4\|, \quad (3.5)$$

where $\|\cdot\|$ can be either the spectral or the Frobenius norm. Similar results also hold for F_V^{12} and F_V^{21} .

3.2 Examples in using Theorem 3.1

We demonstrate how to use Theorem 3.1 to simplify proofs of some existing perturbation bounds in the literature. The theorem we use to derive all the new results in this paper is in the next section (Theorem 3.7). Curious readers may safely jump to the next section from here.

Example 1: The angular perturbation formulae in Theorem 3.1 naturally yield the one-sided $\sin\Theta$ bounds first discovered in [10]. Theorem 3.1 now introduces a very straightforward derivation of these bounds.

Theorem 3.5 (One-sided $\sin\Theta$ theorem). *Using the same notation and quantities as in Theorem 3.1, if $\sigma_r - \tilde{\sigma}_{r+1} > 0$, $\tilde{\sigma}_r - \sigma_{r+1} > 0$, then*

$$\|\sin\Theta(U_1, \tilde{U}_1)\| \leq \min \left\{ \frac{\tilde{\sigma}_r \|(\Delta A) \tilde{V}_1\|}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} + \frac{\sigma_{r+1} \|(\Delta A)^T \tilde{U}_1\|}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2}, \frac{\sigma_r \|(\Delta A) V_1\|}{\sigma_r^2 - \tilde{\sigma}_{r+1}^2} + \frac{\tilde{\sigma}_{r+1} \|U_1^T (\Delta A)\|}{\sigma_r^2 - \tilde{\sigma}_{r+1}^2} \right\}, \quad (3.6)$$

$$\|\sin\Theta(V_1, \tilde{V}_1)\| \leq \min \left\{ \frac{\tilde{\sigma}_r \|\tilde{U}_1^T (\Delta A)\|}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} + \frac{\sigma_{r+1} \|(\Delta A) \tilde{V}_1\|}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2}, \frac{\sigma_r \|U_1^T (\Delta A)\|}{\sigma_r^2 - \tilde{\sigma}_{r+1}^2} + \frac{\tilde{\sigma}_{r+1} \|(\Delta A) V_1\|}{\sigma_r^2 - \tilde{\sigma}_{r+1}^2} \right\}. \quad (3.7)$$

Moreover,

$$\max\{\|\sin\Theta(U_1, \tilde{U}_1)\|, \|\sin\Theta(V_1, \tilde{V}_1)\|\} \leq \min \left\{ \frac{1}{\sigma_r - \tilde{\sigma}_{r+1}}, \frac{1}{\tilde{\sigma}_r - \sigma_{r+1}} \right\} \|\Delta A\|. \quad (3.8)$$

(3.6) and (3.7) are individual bounds on $\|\sin\Theta(U_1, \tilde{U}_1)\|$ and $\|\sin\Theta(V_1, \tilde{V}_1)\|$, while the classical Wedin's $\sin\Theta$ theorem is a uniform bound on both $\|\sin\Theta(U_1, \tilde{U}_1)\|$ and $\|\sin\Theta(V_1, \tilde{V}_1)\|$. The benefit of obtaining the individual bounds was clearly pointed out in [10] by an example. When $A \in \mathbb{R}^{n \times m}$ is a fixed rank- r matrix with $r < n \ll m$, and $\Delta A \in \mathbb{R}^{n \times m}$ is a small random matrix with i.i.d. standard normal entries. The Wedin's theorem implies

$$\max\{\|\sin\Theta(U_1, \tilde{U}_1)\|, \|\sin\Theta(V_1, \tilde{V}_1)\|\} \leq \frac{C \max\{\sqrt{n}, \sqrt{m}\}}{\sigma_r}, \quad (3.9)$$

while the one-sided bounds approximately give,

$$\|\sin\Theta(U_1, \tilde{U}_1)\| \leq \frac{C\sqrt{n}}{\sigma_r}, \quad \|\sin\Theta(V_1, \tilde{V}_1)\| \leq \frac{C\sqrt{m}}{\sigma_r}. \quad (3.10)$$

Since we assumed $n \ll m$, only the one-sided bound successfully indicated that U_1 is more stable than V_1 .

The proof of Theorem 3.5 is a simple application of Theorem 3.1.

Proof. From Theorem 3.1 we have

$$U_2^T \tilde{U}_1 = F_U^{21} \circ (U_2^T (\Delta A) \tilde{V}_1 \tilde{\Sigma}_1^T + \Sigma_2 V_2^T (\Delta A)^T \tilde{U}_1),$$

$$U_1^T \tilde{U}_2 = F_U^{12} \circ (U_1^T (\Delta A) \tilde{V}_2 \tilde{\Sigma}_2^T + \Sigma_1 V_1^T (\Delta A)^T \tilde{U}_2).$$

By (3.4) in Lemma 3.4,

$$\begin{aligned} \|U_2^T \tilde{U}_1\| &\leq \frac{\tilde{\sigma}_r}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} \|U_2^T (\Delta A) \tilde{V}_1\| + \frac{\sigma_{r+1}}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} \|V_2^T (\Delta A)^T \tilde{U}_1\| \\ &\leq \frac{\tilde{\sigma}_r}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} \|(\Delta A) \tilde{V}_1\| + \frac{\sigma_{r+1}}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} \|(\Delta A)^T \tilde{U}_1\| \end{aligned} \quad (3.11)$$

$$\leq \frac{\|\Delta A\|}{\tilde{\sigma}_r - \sigma_{r+1}}. \quad (3.12)$$

Similarly,

$$\|U_1^T \tilde{U}_2\| \leq \frac{\tilde{\sigma}_{r+1}}{\sigma_r^2 - \tilde{\sigma}_{r+1}^2} \|U_1^T \Delta A\| + \frac{\sigma_r}{\sigma_r^2 - \tilde{\sigma}_{r+1}^2} \|(\Delta A) V_1\| \quad (3.13)$$

$$\leq \frac{\|\Delta A\|}{\sigma_r - \tilde{\sigma}_{r+1}}. \quad (3.14)$$

Inserting (3.11) and (3.13) into $\|\sin\Theta(U_1, \tilde{U}_1)\| = \min\{\|U_1^T \tilde{U}_2\|, \|U_2^T \tilde{U}_1\|\}$, we obtain (3.6). Similarly, (3.7) also holds. (3.8) is obtained by using (3.12) and (3.14). \square

Example 2: In this example, we show that one may obtain some interesting results when applying Theorem 3.1 to some less usual choices of ΔA .

Explicitly, we use Theorem 3.1 to re-derive a useful result in [10] but with a more straightforward proof. The result, copied in Proposition 3.6, is about the $\sin\Theta$ distance between the leading singular subspace of a matrix A and an arbitrary subspace.

Proposition 3.6 (Proposition 1 in [10]). *Suppose $A \in \mathbb{R}^{n \times m}$. The orthonormal matrix $V = [V_1, V_2] \in \mathbb{R}^{m \times m}$ is the matrix of right singular vectors of A , i.e., $V_1 \in \mathbb{R}^{m \times r}$, $V_2 \in \mathbb{R}^{m \times (m-r)}$ correspond to the first r and last $m-r$ singular vectors respectively. $[W_1, W_2] \in \mathbb{R}^{m \times m}$ is any orthonormal matrix with $W_1 \in \mathbb{R}^{m \times r}$, $W_2 \in \mathbb{R}^{m \times (m-r)}$. Given that $\sigma_r(AW_1) > \sigma_{r+1}(A)$, we have*

$$\|\sin\Theta(V_1, W_1)\| \leq \min \left\{ \frac{\sigma_r(AW_1) \|\mathbb{P}_{(AW_1)} AW_2\|}{\sigma_r^2(AW_1) - \sigma_{r+1}^2(A)}, 1 \right\}. \quad (3.15)$$

$$\|\sin\Theta(V_1, W_1)\|_F \leq \min \left\{ \frac{\sigma_r(AW_1) \|\mathbb{P}_{(AW_1)} AW_2\|_F}{\sigma_r^2(AW_1) - \sigma_{r+1}^2(A)}, \sqrt{r} \right\}. \quad (3.16)$$

In order to use Theorem 3.1 to prove Proposition 3.6, we recognize that Proposition 3.6 is actually a $\sin\Theta$ bound under a special perturbation. Specifically, if we set $\Delta A = AW_1 W_1^T - A$, then the quantity $\|\sin\Theta(V_1, W_1)\|$ bounded in Proposition 3.6 is exactly the $\sin\Theta$ angle between A and $\tilde{A} = A + \Delta A$. In addition, this particular choice of ΔA has small magnitude of norm therefore leading to a small perturbation bound.

Proof. Apply Theorem 3.1 to A and $\tilde{A} = AW_1 W_1^T$, which means $\Delta A = \tilde{A} - A = AW_1 W_1^T - A = -AW_2 W_2^T$. Assume $U_i, V_i, \Sigma_i, \tilde{U}_i, \tilde{V}_i, \tilde{\Sigma}_i$, $i = 1, 2$ are from the conformal SVDs (2.1) of this A and \tilde{A} . Then using the

notation in Theorem 3.1, we have $(F_V^{21})_{i,j} = \frac{1}{\sigma_j^2(AW_1) - \sigma_{i+r}^2(A)}$, $\tilde{V}_1 = W_1$, $\Sigma_2^T U_2^T (\Delta A) \tilde{V}_1 = 0$. Theorem 3.1 in this case gives

$$V_2^T W_1 = F_V^{21} \circ (V_2^T (\Delta A)^T \tilde{U}_1 \tilde{\Sigma}_1).$$

By Lemma 3.4, this implies

$$\|V_2^T W_1\| \leq \frac{\sigma_r(AW_1) \|\tilde{U}_1^T A W_2 W_2^T V_2\|}{\sigma_r^2(AW_1) - \sigma_{r+1}^2(A)} \leq \frac{\sigma_r(AW_1) \|P_{AW_1} A W_2\|}{\sigma_r^2(AW_1) - \sigma_{r+1}^2(A)},$$

where $\|\cdot\|$ can be either the spectral or Frobenius norm. Also, we directly have $\|V_2^T W_1\| \leq 1$ and $\|V_2^T W_1\|_F \leq \sqrt{r}$, thus (3.15) and (3.16) hold. \square

3.3 High order $\sin\Theta$ distance formulae using series expansions

Although the formulae in Theorem 3.1 are already quite useful, they are still only first-order formulae in the following sense. Looking at the first formula in (3.3) of Theorem 3.1, a closer examination shows that the unknown left hand side $U_1^T \tilde{U}_2$ also appears implicitly in the right hand side, albeit as high order terms. Since we consider upper bounds in the non-asymptotic regime, high order errors may sometimes affect the tightness of the bound, so we hope to get rid of them.

To be more specific about the implicit appearances of the high order terms, we denote the left hand sides of the four formulae in Theorem 3.1 as X, Y, W, Z

$$X := U_1^T \tilde{U}_2, \quad Y := U_2^T \tilde{U}_1, \quad W := V_1^T \tilde{V}_2, \quad Z := V_2^T \tilde{V}_1.$$

First focus on the expression of Y in Theorem 3.1

$$\begin{aligned} Y &\equiv U_2^T \tilde{U}_1 = F_U^{21} \circ (U_2^T (\Delta A) \tilde{V}_1 \tilde{\Sigma}_1^T + \Sigma_2 V_2^T (\Delta A)^T \tilde{U}_1) \\ &= F_U^{21} \circ (\Sigma_2 V_2^T (\Delta A)^T U_1 U_1^T \tilde{U}_1 + \Sigma_2 V_2^T (\Delta A)^T U_2 U_2^T \tilde{U}_1 + U_2^T (\Delta A) V_1 V_1^T \tilde{V}_1 \tilde{\Sigma}_1^T \\ &\quad + U_2^T (\Delta A) V_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^T) \\ &= \underbrace{F_U^{21} \circ (\Sigma_2 \alpha_{12}^T U_1^T \tilde{U}_1 + \alpha_{21} V_1^T \tilde{V}_1 \tilde{\Sigma}_1^T)}_{:=C_1} + F_U^{21} \circ (\Sigma_2 \alpha_{22}^T Y) + F_U^{21} \circ (\alpha_{22} Z \tilde{\Sigma}_1^T), \end{aligned} \quad (3.17)$$

where the second line used $U_1 U_1^T + U_2 U_2^T = I$ and $V_1 V_1^T + V_2 V_2^T = I$, the third line is a re-grouping of terms, and $\alpha_{ij} := U_i^T \Delta A V_j$, $1 \leq i, j \leq 2$. We can get the same expression for Z

$$\begin{aligned} Z &\equiv V_2^T \tilde{V}_1 = F_V^{21} \circ (\Sigma_2^T U_2^T (\Delta A) \tilde{V}_1 + V_2^T (\Delta A)^T \tilde{U}_1 \tilde{\Sigma}_1) \\ &= F_V^{21} \circ (V_2^T (\Delta A)^T U_1 U_1^T \tilde{U}_1 \tilde{\Sigma}_1 + V_2^T (\Delta A)^T U_2 U_2^T \tilde{U}_1 \tilde{\Sigma}_1 + \Sigma_2^T U_2^T (\Delta A) V_1 V_1^T \tilde{V}_1 \\ &\quad + \Sigma_2^T U_2^T (\Delta A) V_2 V_2^T \tilde{V}_1) \\ &= \underbrace{F_V^{21} \circ (\alpha_{12}^T U_1^T \tilde{U}_1 \tilde{\Sigma}_1 + \Sigma_2^T \alpha_{21} V_1^T \tilde{V}_1)}_{:=C_2} + F_V^{21} \circ (\alpha_{22}^T Y \tilde{\Sigma}_1) + F_V^{21} \circ (\Sigma_2^T \alpha_{22} Z). \end{aligned} \quad (3.18)$$

Looking at the last right hand sides of (3.17) and (3.18), we see that Y and Z are contained in the second and third terms, respectively, so they appear on both hand sides.

To highlight this structure, we shorten the notation by letting \mathcal{F} be the linear operator defined as

$$\mathcal{F} \left(\begin{pmatrix} Y \\ Z \end{pmatrix} \right) = \begin{pmatrix} F_U^{21} \circ (\Sigma_2 \alpha_{22}^T Y) + F_U^{21} \circ (\alpha_{22} Z \tilde{\Sigma}_1^T) \\ F_V^{21} \circ (\alpha_{22}^T Y \tilde{\Sigma}_1) + F_V^{21} \circ (\Sigma_2^T \alpha_{22} Z) \end{pmatrix}.$$

Then (3.17) and (3.18) become,

$$\begin{bmatrix} Y \\ Z \end{bmatrix} = \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} + \mathcal{F} \left(\begin{bmatrix} Y \\ Z \end{bmatrix} \right).$$

Clearly, this is an implicit equation system of Y and Z .

Provided $\|\mathcal{F}\| < 1$, we can move \mathcal{F} to the left and take the inverse

$$\begin{pmatrix} U_2^T \tilde{U}_1 \\ V_2^T \tilde{V}_1 \end{pmatrix} \equiv \begin{bmatrix} Y \\ Z \end{bmatrix} = (1 - \mathcal{F})^{-1} \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \sum_{k=0}^{\infty} \mathcal{F}^k \begin{pmatrix} C_1 \\ C_2 \end{pmatrix}.$$

This gives us a series expression of the quantities $U_2^T \tilde{U}_1$ and $V_2^T \tilde{V}_1$, which allows us to derive Theorem 2.1 and Theorem 2.6 presented in Section 2. We summarize this result in the following theorem.

Theorem 3.7 (Angular perturbation formula using series expansion). *Using the same notation and quantities as in Theorem 3.1, we have*

$$\begin{pmatrix} U_2^T \tilde{U}_1 \\ V_2^T \tilde{V}_1 \end{pmatrix} = \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} + \mathcal{F} \left(\begin{bmatrix} U_2^T \tilde{U}_1 \\ V_2^T \tilde{V}_1 \end{bmatrix} \right), \quad \begin{pmatrix} U_1^T \tilde{U}_2 \\ V_1^T \tilde{V}_2 \end{pmatrix} = \begin{pmatrix} C_3 \\ C_4 \end{pmatrix} + \mathcal{G} \left(\begin{bmatrix} U_1^T \tilde{U}_2 \\ V_1^T \tilde{V}_2 \end{bmatrix} \right). \quad (3.19)$$

In addition, provided that $\|\mathcal{F}\| < 1$ and $\|\mathcal{G}\| < 1$, we have

$$\begin{pmatrix} U_2^T \tilde{U}_1 \\ V_2^T \tilde{V}_1 \end{pmatrix} = \sum_{k=0}^{\infty} \mathcal{F}^k \begin{pmatrix} C_1 \\ C_2 \end{pmatrix}, \quad \begin{pmatrix} U_1^T \tilde{U}_2 \\ V_1^T \tilde{V}_2 \end{pmatrix} = \sum_{k=0}^{\infty} \mathcal{G}^k \begin{pmatrix} C_3 \\ C_4 \end{pmatrix}. \quad (3.20)$$

Here

$$\begin{aligned} \mathcal{F} \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} &= \begin{pmatrix} F_U^{21} \circ (\Sigma_2 \alpha_{22}^T C_1) + F_U^{21} \circ (\alpha_{22} C_2 \tilde{\Sigma}_1^T) \\ F_V^{21} \circ (\alpha_{22}^T C_1 \tilde{\Sigma}_1) + F_V^{21} \circ (\Sigma_2^T \alpha_{22} C_2) \end{pmatrix}, \\ \mathcal{G} \begin{pmatrix} C_3 \\ C_4 \end{pmatrix} &= \begin{pmatrix} F_U^{12} \circ (\alpha_{11} C_4 \tilde{\Sigma}_2^T) + F_U^{12} \circ (\Sigma_1 \alpha_{11}^T C_3) \\ F_V^{12} \circ (\Sigma_1^T \alpha_{11} C_4) + F_V^{12} \circ (\alpha_{11}^T C_3 \tilde{\Sigma}_2) \end{pmatrix}. \end{aligned}$$

$$\begin{aligned} C_1 &= F_U^{21} \circ (\Sigma_2 \alpha_{12}^T U_1^T \tilde{U}_1 + \alpha_{21} V_1^T \tilde{V}_1 \tilde{\Sigma}_1^T), & C_2 &= F_V^{21} \circ (\alpha_{12}^T U_1^T \tilde{U}_1 \tilde{\Sigma}_1 + \Sigma_2^T \alpha_{21} V_1^T \tilde{V}_1), \\ C_3 &= F_U^{12} \circ (\alpha_{12} V_2^T \tilde{V}_2 \tilde{\Sigma}_2^T + \Sigma_1 \alpha_{21}^T U_2^T \tilde{U}_2), & C_4 &= F_V^{12} \circ (\Sigma_1^T \alpha_{12} V_2^T \tilde{V}_2 + \alpha_{21}^T U_2^T \tilde{U}_2 \tilde{\Sigma}_2). \end{aligned}$$

and $\alpha_{ij} := U_i^T \Delta A V_j$.

Remark 3.8. Careful readers may observe that, although we removed all cross terms $U_1^T \tilde{U}_2$, $U_2^T \tilde{U}_1$, $V_1^T \tilde{V}_2$, $V_2^T \tilde{V}_1$ from the right hand sides of the expressions (3.20), there are still terms like $U_1^T \tilde{U}_1$ and $V_1^T \tilde{V}_1$ appearing on the right hand side. In fact, these terms are of order $O(1)$ thus will not degrade the tightness of the upper bounds by any order of magnitudes and only possibly affect the constants.

When A has rank r , Theorem 3.7 reduces to the following simpler formulae.

Corollary 3.9. *Using the definitions above, when A has rank r and $\|\Delta A\| < \sigma_r(\tilde{A})$,*

$$\begin{aligned} U_2^T \tilde{U}_1 &= \sum_{k=0}^{+\infty} (\alpha_{22} \alpha_{22}^T)^k (\alpha_{21} V_1^T \tilde{V}_1 + \alpha_{22} \alpha_{12}^T U_1^T \tilde{U}_1 \tilde{\Sigma}_1^{-1}) \tilde{\Sigma}_1^{-(2k+1)}, \\ V_2^T \tilde{V}_1 &= \sum_{k=0}^{+\infty} (\alpha_{22}^T \alpha_{22})^k (\alpha_{12}^T U_1^T \tilde{U}_1 + \alpha_{22}^T \alpha_{21} V_1^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}) \tilde{\Sigma}_1^{-(2k+1)}. \end{aligned} \quad (3.21)$$

Remark 3.10. When A has rank- r and α_{22} is full rank, Corollary 3.9 can also be derived using series expansion for Sylvester-type equations. Denote matrix $M = \begin{pmatrix} & \alpha_{22} \\ \alpha_{22}^T & \end{pmatrix}^{-1}$, $X = \begin{pmatrix} & \alpha_{22} \\ \alpha_{22}^T & \end{pmatrix} \begin{pmatrix} U_2^T \tilde{U}_1 \\ V_2^T \tilde{V}_1 \end{pmatrix}$, $B = \tilde{\Sigma}_1^{-1}$, and $Y = \begin{pmatrix} \alpha_{21} V_1^T \tilde{V}_1 \tilde{\Sigma}_1^{-1} \\ \alpha_{12}^T U_1^T \tilde{U}_1 \tilde{\Sigma}_1^{-1} \end{pmatrix}$. Direct calculation gives $MX - XB = Y$. By the assumption in Corollary 3.9, $\|\alpha_{22}\| \leq \|\Delta A\| < \sigma_r(\tilde{A})$, we can see for any eigenvalue λ of matrix M , it holds that $|\lambda| > \frac{1}{\sigma_r(\tilde{A})}$. Classical series expansion for Sylvester-type equations (Theorem VII.2.2 in [6]) also leads to equation (3.21).

3.4 Examples of Using Theorem 3.7

Theorem 3.7 is used to derive the refined $\ell_{2,\infty}$ bound (Theorem 2.1) and the $\sin\Theta$ bound between singular vectors and their resided singular subspace (Theorem 2.6), which provided the main intuition behind our PCA and singular value truncation results (Theorem 2.7 and Theorem 2.10) in Section 2. Here, we only present the proof of Theorem 2.6, and defer the proof of the rest to the appendix and the supplementary material since they are more involved.

Proof. Again we denote $Y := U_2^T \tilde{U}_1$ and $Z := V_2^T \tilde{V}_1$. Restricting (3.19) in Theorem 3.7 to the j th columns ($1 \leq j \leq r$), we have

$$\begin{aligned} Y_j &= (C_1)_j + \tilde{\sigma}_j (F_U^{21})_j \circ (\alpha_{22} Z_j) + (F_U^{21})_j \circ (\Sigma_2 \alpha_{22}^T Y_j), \\ Z_j &= (C_2)_j + \tilde{\sigma}_j (F_V^{21})_j \circ (\alpha_{22}^T Y_j) + (F_V^{21})_j \circ (\Sigma_2^T \alpha_{22} Z_j). \end{aligned}$$

It is easy to verify that $\|(C_1)_j\| \leq \frac{\tilde{\sigma}_j}{\tilde{\sigma}_j^2 - \sigma_{r+1}^2} \|\alpha_{21}\| + \frac{\sigma_{r+1}}{\tilde{\sigma}_j^2 - \sigma_{r+1}^2} \|\alpha_{12}\|$, $\|(C_2)_j\| \leq \frac{\sigma_{r+1}}{\tilde{\sigma}_j^2 - \sigma_{r+1}^2} \|\alpha_{21}\| + \frac{\tilde{\sigma}_j}{\tilde{\sigma}_j^2 - \sigma_{r+1}^2} \|\alpha_{12}\|$, then

$$\|Y_j\| \leq \frac{1}{\tilde{\sigma}_j^2 - \sigma_{r+1}^2} (\tilde{\sigma}_j \|\alpha_{21}\| + \sigma_{r+1} \|\alpha_{12}\| + \tilde{\sigma}_j \|\alpha_{22}\| \|Z_j\| + \sigma_{r+1} \|\alpha_{22}\| \|Y_j\|),$$

$$\|Z_j\| \leq \frac{1}{\tilde{\sigma}_j^2 - \sigma_{r+1}^2} (\sigma_{r+1} \|\alpha_{21}\| + \tilde{\sigma}_j \|\alpha_{12}\| + \tilde{\sigma}_j \|\alpha_{22}\| \|Y_j\| + \sigma_{r+1} \|\alpha_{22}\| \|Z_j\|).$$

Summing up the first inequality multiplied by $\tilde{\sigma}_j^2 - \sigma_{r+1}^2 - \sigma_{r+1} \|\alpha_{22}\|$ and the second inequality multiplied by $\tilde{\sigma}_j \|\alpha_{22}\|$, after some simplification we get

$$\begin{aligned} \|Y_j\| &= \|U_2^T \tilde{u}_j\| \leq \frac{\tilde{\sigma}_j \|\alpha_{21}\| + \sigma_{r+1} \|\alpha_{12}\| + \|\alpha_{22}\| \|\alpha_{12}\|}{\tilde{\sigma}_j^2 - (\sigma_{r+1} + \|\alpha_{22}\|)^2} \\ &\leq \frac{(\sigma_j - \|\Delta A\|) \|\Delta A\| + \sigma_{r+1} \|\Delta A\| + \|\Delta A\|^2}{(\sigma_j - \|\Delta A\|)^2 - (\sigma_{r+1} + \|\Delta A\|)^2} \\ &\leq \frac{\|\Delta A\|}{\sigma_j - \sigma_{r+1} - 2\|\Delta A\|} \\ &\leq \frac{3\|\Delta A\|}{\sigma_j - \sigma_{r+1}}, \end{aligned}$$

provided that $3\|\Delta A\| \leq \sigma_r - \sigma_{r+1}$. Here the second inequality is because the upper bound on the right hand side is decreasing with respect to $\tilde{\sigma}_j$ and increasing with respect to $\|\alpha_{22}\|$. Similarly, we also have

$$\|Z_j\| = \|V_2^T \tilde{v}_j\| \leq \frac{\tilde{\sigma}_j \|\alpha_{12}\| + \sigma_{r+1} \|\alpha_{21}\| + \|\alpha_{22}\| \|\alpha_{21}\|}{\tilde{\sigma}_j^2 - (\sigma_{r+1} + \|\alpha_{22}\|)^2} \leq \frac{3\|\Delta A\|}{\sigma_j - \sigma_{r+1}}.$$

□

4 Proof of the main results

In Section 4.1, we derive the proof of Theorem 3.1 and Lemma 3.4. After that, we present the proof of Theorem 2.1 in Section 4.2. Since the proof of one key lemma (Lemma 4.3) is long and involved, we divide it into low-rank case and full-rank case. We prove the low-rank case in Section 4.3, and the proof of full-rank case is deferred to appendix. In Section 4.4 we provide the proof of Theorem 2.10, while the proof of Theorem 2.7 can be found in Section 4.5.

4.1 Proof of Theorem 3.1 and Lemma 3.4

Proof of Theorem 3.1. First, decompose the perturbation ΔA in the following two ways :

$$\begin{aligned} \Delta A &= \tilde{A} - A = \tilde{U}\tilde{\Sigma}\tilde{V}^T - U\Sigma V^T \\ &= (U + \Delta U)\tilde{\Sigma}\tilde{V}^T - U\Sigma(\tilde{V} - \Delta V)^T \\ &= U\tilde{\Sigma}\tilde{V}^T + (\Delta U)\tilde{\Sigma}\tilde{V}^T - U\Sigma\tilde{V}^T + U\Sigma(\Delta V)^T \\ &= U(\Delta\Sigma)\tilde{V}^T + (\Delta U)\tilde{\Sigma}\tilde{V}^T + U\Sigma(\Delta V)^T, \end{aligned} \tag{4.1}$$

and

$$\begin{aligned} \Delta A &= \tilde{A} - A = \tilde{U}\tilde{\Sigma}\tilde{V}^T - U\Sigma V^T \\ &= \tilde{U}\tilde{\Sigma}(V + \Delta V)^T - (\tilde{U} - \Delta U)\Sigma V^T \\ &= \tilde{U}\tilde{\Sigma}V^T + \tilde{U}\tilde{\Sigma}(\Delta V)^T - \tilde{U}\Sigma V^T + (\Delta U)\Sigma V^T \\ &= \tilde{U}(\Delta\Sigma)V^T + \tilde{U}\tilde{\Sigma}(\Delta V)^T + (\Delta U)\Sigma V^T. \end{aligned} \tag{4.2}$$

Multiplying (4.1) with U^T on the left and \tilde{V} on the right leads to

$$U^T(\Delta A)\tilde{V} = \Delta\Sigma + U^T(\Delta U)\tilde{\Sigma} + \Sigma(\Delta V)^T\tilde{V}. \tag{4.3}$$

Similarly, multiplying (4.2) with \tilde{U}^T on the left and V on the right we obtain

$$\tilde{U}^T(\Delta A)V = \Delta\Sigma + \tilde{\Sigma}(\Delta V)^T V + \tilde{U}^T(\Delta U)\Sigma. \tag{4.4}$$

Denote $dP = U^T(\Delta A)\tilde{V}$, $d\bar{P} = \tilde{U}^T(\Delta A)V$, $\Delta\Omega_U = U^T(\Delta U)$, $\Delta\Omega_V = V^T(\Delta V)$. Notice that $I = \tilde{U}^T\tilde{U} = U^T U$ gives $(U + \Delta U)^T\tilde{U} = U^T(\tilde{U} - \Delta U)$, hence $U^T\Delta U = -\Delta U^T\tilde{U}$. Similarly, we also have $V^T\Delta V = -\Delta V^T\tilde{V}$. Plugging these into (4.3) and (4.4), we have

$$\begin{cases} dP = U^T\Delta A\tilde{V} = \Delta\Sigma + \Delta\Omega_U\tilde{\Sigma} - \Sigma\Delta\Omega_V, \\ d\bar{P} = \tilde{U}^T\Delta A V = \Delta\Sigma + \tilde{\Sigma}\Delta\Omega_V^T - \Delta\Omega_U^T\Sigma. \end{cases} \tag{4.5}$$

Next, from (4.5) we can cancel $\Delta\Omega_V$ by

$$\begin{aligned} G_U &:= dP\tilde{\Sigma}^T + \Sigma d\bar{P}^T \\ &= \Delta\Sigma\tilde{\Sigma}^T + \Sigma(\Delta\Sigma)^T + \Delta\Omega_U\tilde{\Sigma}\tilde{\Sigma}^T - \Sigma\Sigma^T\Delta\Omega_U \\ &= \tilde{\Sigma}\tilde{\Sigma}^T - \Sigma\Sigma^T + \Delta\Omega_U\tilde{\Sigma}\tilde{\Sigma}^T - \Sigma\Sigma^T\Delta\Omega_U. \end{aligned}$$

Let $\Delta\Omega_U = \{w_{ij}\}_{i,j=1}^n$, then for all $1 \leq i, j \leq n$, the following equations hold

$$(G_U)_{ij} = \begin{cases} (\tilde{\sigma}_j^2 - \sigma_i^2)w_{ij}, & i \neq j, \\ (\tilde{\sigma}_j^2 - \sigma_i^2)(w_{ij} + 1), & i = j. \end{cases} \quad (4.6)$$

Here if $i > \min\{n, m\}$, we define σ_i or $\tilde{\sigma}_i$ to be 0. Also, define F_U^{12} , F_U^{21} , F_V^{12} , F_V^{21} as in the statement of Theorem 3.1. By assumption, $\tilde{\sigma}_r - \sigma_{r+1} > 0$, $\sigma_r - \tilde{\sigma}_{r+1} > 0$, we can directly check that the denominators in these four matrices only have nonzero entries, thus are well defined. Consider the upper right part in $\Delta\Omega_U = U^T(\Delta U)$, that is, $1 \leq i \leq r$, $r+1 \leq j \leq n$, from (4.6) we have

$$w_{ij} = \frac{1}{\tilde{\sigma}_j^2 - \sigma_i^2}(G_U)_{ij}, \quad 1 \leq i \leq r, \quad r+1 \leq j \leq n.$$

Therefore,

$$U_1^T \tilde{U}_2 = U_1^T(\Delta U_2) = F_U^{12} \circ (G_U^{12}) = F_U^{12} \circ (U_1^T(\Delta A)\tilde{V}_2\tilde{\Sigma}_2^T + \Sigma_1 V_1^T(\Delta A)^T \tilde{U}_2).$$

Following the same reasoning, we also obtain

$$\begin{aligned} U_2^T \tilde{U}_1 &= U_2^T(\Delta U_1) = F_U^{21} \circ (U_2^T(\Delta A)\tilde{V}_1\tilde{\Sigma}_1^T + \Sigma_2 V_2^T(\Delta A)^T \tilde{U}_1), \\ V_1^T \tilde{V}_2 &= V_1^T(\Delta V_2) = F_V^{12} \circ (\Sigma_1^T U_1^T(\Delta A)\tilde{V}_2 + V_1^T(\Delta A)^T \tilde{U}_2 \tilde{\Sigma}_2), \\ V_2^T \tilde{V}_1 &= V_2^T(\Delta V_2) = F_V^{21} \circ (\Sigma_2^T U_2^T(\Delta A)\tilde{V}_1 + V_2^T(\Delta A)^T \tilde{U}_1 \tilde{\Sigma}_1). \end{aligned}$$

□

Proof of Lemma 3.4. Here we only prove the first inequality in (3.4), i.e., $\|F_U^{21} \circ (H_1 \tilde{\Sigma}_1)\| \leq \frac{\tilde{\sigma}_r}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} \|H_1\|$, the other three inequalities can be proved similarly. Recall the definition of F_U^{21} is $(F_U^{21})_{i-r,j} = \frac{1}{\tilde{\sigma}_j^2 - \sigma_i^2}$, $r+1 \leq i \leq n$, $1 \leq j \leq r$. We directly have

$$F_U^{21} \circ (H_1 \tilde{\Sigma}_1) = \bar{F}_U^{21} \circ H_1,$$

where

$$(\bar{F}_U^{21})_{i-r,j} = \frac{\tilde{\sigma}_j}{\tilde{\sigma}_j^2 - \sigma_i^2}, \quad r+1 \leq i \leq n, \quad 1 \leq j \leq r.$$

Let $B_1 = F_U^{21} \circ (H_1 \tilde{\Sigma}_1)$, then $H_1 = \tilde{F}_U^{21} \circ B_1$, where

$$(\tilde{F}_U^{21})_{i-r,j} = \frac{\tilde{\sigma}_j^2 - \sigma_i^2}{\tilde{\sigma}_j} = \tilde{\sigma}_j - \frac{\sigma_i^2}{\tilde{\sigma}_j}, \quad r+1 \leq i \leq n, \quad 1 \leq j \leq r.$$

Inserting the above expression of \tilde{F} into $H_1 = \tilde{F}_U^{21} \circ B_1$, we have

$$H_1 = B_1 \begin{pmatrix} \tilde{\sigma}_1 & & & \\ & \tilde{\sigma}_2 & & \\ & & \ddots & \\ & & & \tilde{\sigma}_r \end{pmatrix} - \begin{pmatrix} \sigma_{r+1}^2 & & & \\ & \sigma_{r+2}^2 & & \\ & & \ddots & \\ & & & \sigma_n^2 \end{pmatrix} B_1 \begin{pmatrix} \frac{1}{\tilde{\sigma}_1} & & & \\ & \frac{1}{\tilde{\sigma}_2} & & \\ & & \ddots & \\ & & & \frac{1}{\tilde{\sigma}_r} \end{pmatrix}.$$

Take norm on both sides, we obtain

$$\|H_1\| \geq \tilde{\sigma}_r \|B_1\| - \frac{\sigma_{r+1}^2}{\tilde{\sigma}_r} \|B_1\| = \frac{\tilde{\sigma}_r^2 - \sigma_{r+1}^2}{\tilde{\sigma}_r} \|B_1\|,$$

which further gives $\|B_1\| \leq \frac{\tilde{\sigma}_r}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} \|H_1\|$. \square

Remark 4.1. When $\tilde{\sigma}_r > \sigma_{r+1}$, $\sigma_r > \tilde{\sigma}_{r+1}$, the bounds in Lemma 3.4 are tight. That is, in this case, there exists H_i , $1 \leq i \leq 4$, such that the equalities in (3.4) and (3.5) hold. Specifically, let

$$H_1 = \begin{pmatrix} 0 & \cdots & 0 & \epsilon \\ 0 & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & \cdots & \cdots & 0 \end{pmatrix} \in \mathbb{R}^{(n-r) \times r}, \quad H_3 = \begin{pmatrix} 0 & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & \vdots & \vdots & \vdots \\ \epsilon & 0 & \cdots & 0 \end{pmatrix} \in \mathbb{R}^{r \times (m-r)},$$

$$H_2 = \begin{pmatrix} 0 & \cdots & 0 & \epsilon \\ 0 & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & \cdots & \cdots & 0 \end{pmatrix} \in \mathbb{R}^{(m-r) \times r}, \quad H_4 = \begin{pmatrix} 0 & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & \vdots & \vdots & \vdots \\ \epsilon & 0 & \cdots & 0 \end{pmatrix} \in \mathbb{R}^{r \times (n-r)},$$

then we can directly check that the equalities in (3.4) and (3.5) hold.

4.2 Proof of Theorem 2.1

To prove Theorem 2.1, we need to decompose $\tilde{U}_1 - U_1 Q$ into a sum of several components and bound them separately. For convenience, we put the decomposition in the following lemma, which is similar in nature to Theorem 3.1 in [14].

Proposition 4.2. *Set the rotation Q to be $Q = Q_1 Q_2^T$, where Q_1 and Q_2 are the left and right singular vectors from the SVD: $U_1^T \tilde{U}_1 = Q_1 S Q_2^T$, then*

$$\tilde{U}_1 - U_1 Q = U_2 U_2^T \Delta A V_1 V_1^T \tilde{V}_1 \tilde{\Sigma}_1^{-1} + U_2 U_2^T \Delta A V_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1} + U_2 \Sigma_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1} + U_1 Q_1 (S - I) Q_2^T, \quad (4.7)$$

and

$$\|S - I\| \leq \|\sin\Theta(U_1, \tilde{U}_1)\|^2. \quad (4.8)$$

Proof. By direct calculation, we have

$$\begin{aligned} \tilde{U}_1 - U_1 Q &= \tilde{U}_1 - U_1 Q_1 Q_2^T \\ &= \tilde{U}_1 - U_1 Q_1 S Q_2^T + U_1 Q_1 (S - I) Q_2^T \\ &= \tilde{U}_1 - U_1 U_1^T \tilde{U}_1 + U_1 Q_1 (S - I) Q_2^T \\ &= U_2 U_2^T \tilde{U}_1 + U_1 Q_1 (S - I) Q_2^T \\ &= U_2 U_2^T \Delta A \tilde{V}_1 \tilde{\Sigma}_1^{-1} + U_2 \Sigma_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1} + U_1 Q_1 (S - I) Q_2^T \\ &= U_2 U_2^T \Delta A V_1 V_1^T \tilde{V}_1 \tilde{\Sigma}_1^{-1} + U_2 U_2^T \Delta A V_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1} + U_2 \Sigma_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1} + U_1 Q_1 (S - I) Q_2^T. \end{aligned} \quad (4.9)$$

In addition, since $\|S\| = \|U_1^T \tilde{U}_1\| \leq 1$,

$$\|S - I\| = 1 - \min_i S_i \leq 1 - \min_i S_i^2 = \|\sin\Theta(U_1, \tilde{U}_1)\|^2,$$

where S_i is the i th diagonal entry of S . Hence

$$\|U_1 Q_1 (S - I) Q_2^T\|_{2,\infty} \leq \|U_1\|_{2,\infty} \|S - I\| \leq \|U_1\|_{2,\infty} \|\sin\Theta(U_1, \tilde{U}_1)\|^2.$$

□

The first and the last terms in the expansion (4.7) are easy to bound, the following lemma is devoted to bounding the middle terms, which requires invoking the angular perturbation formula Theorem 3.7.

Lemma 4.3. *Under the assumption of Theorem 2.1, it holds that*

$$\max\{\|U_2 U_2^T \Delta A V_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty}, \|U_2 \Sigma_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty}\} \leq C \frac{\sigma R(r, n)}{\sigma_r(A) - \sigma_{r+1}(A)},$$

where C is some constant and

$$R(r, n) = \begin{cases} \sqrt{r} + \sqrt{\log n}, & \text{if } A \text{ is of rank } r; \\ r + \sqrt{r \log n}, & \text{else.} \end{cases}$$

Before proving this lemma, let us first see how to use it to prove Theorem 2.1.

Proof of Theorem 2.1. Due to (4.7), we have

$$\begin{aligned} \min_{\tilde{Q} \in \mathcal{O}_r} \|\tilde{U}_1 - U_1 \tilde{Q}\|_{2,\infty} &\leq \underbrace{\|U_2 U_2^T \Delta A V_1 V_1^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty}}_{\text{(I)}} + \|U_2 U_2^T \Delta A V_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty} \\ &\quad + \|U_2 \Sigma_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty} + \underbrace{\|U_1\|_{2,\infty} \|\sin\Theta(U_1, \tilde{U}_1)\|^2}_{\text{(II)}}. \end{aligned}$$

The two middle terms are bounded in Lemma 4.3. We are left to bound the first and the last terms. For the last term, we have

$$\text{(II)} \leq \|U_1\|_{2,\infty} \left(\frac{2\|\Delta A\|}{\sigma_r(A) - \sigma_{r+1}(A)} \right)^2 \leq \|U_1\|_{2,\infty} \frac{36\sigma^2 \bar{n}}{(\sigma_r(A) - \sigma_{r+1}(A))^2}.$$

Here the first inequality used (4.14) in Lemma 4.7, the second one used Corollary 7.3.3 of [40] which bounds the spectral norm of i.i.d. Gaussian matrices: with probability at least $1 - e^{-c\bar{n}}$ for some absolute constant c , $\|\Delta A\| \leq 3\sigma\sqrt{\bar{n}}$.

Next we bound (I).

$$\begin{aligned} \text{(I)} &= \|U_2 U_2^T \Delta A V_1 V_1^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty} \leq \|U_2 U_2^T \Delta A V_1\|_{2,\infty} \|V_1^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\| \\ &\leq \frac{1}{\sigma_r(\tilde{A})} \|U_2 U_2^T \Delta A V_1\|_{2,\infty} \leq \frac{7}{6\sigma_r(A)} \|U_2 U_2^T \Delta A V_1\|_{2,\infty}. \end{aligned} \tag{4.10}$$

Here the last inequality is by Weyl's bound and the assumption $\sigma_r(A) > 21\sigma\sqrt{\bar{n}}$. (4.10) implies that it suffices to bound the row norms of $U_2 U_2^T \Delta A V_1$. Since ΔA is i.i.d. $N(0, \sigma^2)$, U_2 and V_1 are independent of ΔA and that $\|U_2\| = \|V_1\| = 1$, then each row of $U_2 U_2^T \Delta A V_1$ is a Gaussian vector having independent

Gaussian entries with mean 0 and variance at most σ^2 . By exactly the same proof as Theorem 3.1.1 in [40], there exists a constant c such that for all $t > 0$,

$$\mathbb{P}(\|u_i^T U_2^T \Delta A V_1\| - \sigma \|u_i^T U_2^T\| \sqrt{r} \geq t) < 2e^{-\frac{ct^2}{\sigma^2 \|u_i^T U_2^T\|^2}},$$

where u_i^T is the i th row of U_2 .

Setting in the above $t = \sigma \|u_i^T U_2^T\| \sqrt{3 \log n / c}$, then with probability at least $1 - \frac{2}{n^3}$,

$$\|u_i^T U_2^T \Delta A V_1\| \leq c_1 \sigma (\sqrt{r} + \sqrt{\log n}) \|u_i^T U_2^T\| \leq c_1 \sigma (\sqrt{r} + \sqrt{\log n}),$$

with some constant c_1 . By the union bound, the probability of failure for all the rows is at most $\frac{2}{n^2}$. Hence with probability at least $1 - \frac{2}{n^2}$, it holds

$$\|U_2 U_2^T \Delta A V_1\|_{2,\infty} \leq c_1 \sigma (\sqrt{r} + \sqrt{\log n}).$$

Plugging this into (4.10), we obtain

$$(I) \leq \frac{c_1 \sigma (\sqrt{r} + \sqrt{\log n})}{\sigma_r(A)}.$$

Combining the bounds on I, II and Lemma 4.3 completes the proof. \square

4.3 Proof of Lemma 4.3

Here we first provide the proof for the low-rank case to give the reader some intuition. The full-rank case follows a similar idea but is quite notationally heavy, we defer the proof of Lemma 4.3 for full-rank case to appendix.

Proof of Lemma 4.3- the low-rank case. When A is of rank r , the second quantity to be bounded in Lemma 4.3 is 0, hence we focus on the first quantity $\|U_2 U_2^T \Delta A V_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty}$.

Let u_i^T be the i th row of U_2 , then by Corollary 3.9, the i th row of $U_2 U_2^T \Delta A V_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}$ can be expressed as

$$\begin{aligned} u_i^T U_2^T \Delta A V_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1} &= u_i^T \alpha_{22} \left(\sum_{k=0}^{\infty} (\alpha_{22}^T \alpha_{22})^k (\alpha_{12}^T U_1^T \tilde{U}_1 \tilde{\Sigma}_1^{-1} + \alpha_{22}^T \alpha_{21} V_1^T \tilde{V}_1 \tilde{\Sigma}_1^{-2}) (\tilde{\Sigma}_1^{-2})^k \right) \tilde{\Sigma}_1^{-1} \\ &= u_i^T \left(\sum_{k=0}^{\infty} (\alpha_{22} \alpha_{22}^T)^k (\alpha_{22} \alpha_{12}^T U_1^T \tilde{U}_1 \tilde{\Sigma}_1^{-1} + \alpha_{22} \alpha_{22}^T \alpha_{21} V_1^T \tilde{V}_1 \tilde{\Sigma}_1^{-2}) (\tilde{\Sigma}_1^{-2})^k \right) \tilde{\Sigma}_1^{-1}, \end{aligned} \quad (4.11)$$

where $\alpha_{ij} = U_i^T \Delta A V_j$. Due to the orthogonality of U and V , the entries in each α_{ij} follow i.i.d. $N(0, \sigma^2)$ distribution, and α_{22} is independent of α_{12} . This further implies that $u_i^T (\alpha_{22} \alpha_{22}^T)^k \alpha_{22}$ and $u_i^T (\alpha_{22} \alpha_{22}^T)^k \alpha_{22} \alpha_{22}^T$ are independent of α_{12} and α_{21} , respectively. Conditional on α_{22} , $u_i^T (\alpha_{22} \alpha_{22}^T)^k \alpha_{22} \alpha_{12}^T$ varies with α_{12} , and it follows normal distribution. Again by Theorem 3.1.1 in [40], for fixed $k = 0, \dots$, there exists a constant c such that

$$\mathbb{P}(\|u_i^T (\alpha_{22} \alpha_{22}^T)^k \alpha_{22} \alpha_{12}^T\| - \sigma \sqrt{r} \|u_i^T (\alpha_{22} \alpha_{22}^T)^k \alpha_{22}\| > t) \leq 2 \exp\left(-\frac{ct^2}{\sigma^2 \|u_i^T (\alpha_{22} \alpha_{22}^T)^k \alpha_{22}\|^2}\right).$$

Setting in the above $t = \sigma \|u_i^T (\alpha_{22} \alpha_{22}^T)^k \alpha_{22}\| \sqrt{\log(n^3 \cdot 2^k)}/c$, we get with probability at least $1 - \frac{2}{2^k n^3}$,

$$\begin{aligned} \|u_i^T (\alpha_{22} \alpha_{22}^T)^k \alpha_{22} \alpha_{12}^T\| &\leq \sigma \sqrt{r} \|u_i^T (\alpha_{22} \alpha_{22}^T)^k \alpha_{22}\| + t \\ &\leq c_2 \sigma \|u_i^T (\alpha_{22} \alpha_{22}^T)^k \alpha_{22}\| (\sqrt{r} + \sqrt{\log n} + \sqrt{k}), \end{aligned} \quad (4.12)$$

where c_2 is some absolute constant. Then

$$\|u_i^T (\alpha_{22} \alpha_{22}^T)^k \alpha_{22} \alpha_{12}^T U_1^T \tilde{U}_1 \tilde{\Sigma}_1^{-(2k+2)}\| \leq c_2 \frac{\sigma}{\tilde{\sigma}_r} \left(\frac{\|\alpha_{22}\|}{\tilde{\sigma}_r} \right)^{2k+1} (\sqrt{r} + \sqrt{\log n} + \sqrt{k}).$$

Let $\lambda = \frac{\|\alpha_{22}\|}{\tilde{\sigma}_r}$. We next argue that $\lambda < 1/2$. By Corollary 7.3.3 of [40], $\|\Delta A\| \leq 3\sigma\sqrt{n}$ with probability at least $1 - e^{-cn}$. On this event, by Weyl's bound,

$$\tilde{\sigma}_r \geq \sigma_r - \|\Delta A\| \geq \sigma_r - 3\sigma\sqrt{n} \geq 18\sigma\sqrt{n} \geq 6\|\Delta A\| \geq 6\|\alpha_{22}\|,$$

which implies $\lambda < 1/2$. The third inequality above is due to the assumption $21\sigma\sqrt{n} < \sigma_r$. By union bound on the probability of failure of (4.12) over all $k = 0, \dots$, we have with probability at least $1 - \frac{4}{n^3}$,

$$\begin{aligned} \sum_{k=0}^{\infty} \|u_i^T (\alpha_{22} \alpha_{22}^T)^k \alpha_{22} \alpha_{12}^T U_1^T \tilde{U}_1 \tilde{\Sigma}_1^{-(2k+2)}\|_2 &\leq c_3 \frac{\sigma}{\tilde{\sigma}_r} \left(\sum_{k=0}^{\infty} \sqrt{k} \lambda^{2k+1} + (\sqrt{\log n} + \sqrt{r}) \sum_{k=0}^{\infty} \lambda^{2k+1} \right) \\ &\leq c_4 \frac{\sigma}{\tilde{\sigma}_r} (\sqrt{r} + \sqrt{\log n}) \\ &\leq c_5 \sigma \frac{\sqrt{r} + \sqrt{\log n}}{\sigma_r(A)}, \end{aligned}$$

with $c_3 - c_5$ being absolute constants, where the last inequality used Weyl's bound and the assumption $\sigma_r(A) > 21\sigma\sqrt{n}$, and the second inequality used the fact that for any $0 < \lambda < 1/2$, we have

$$\sum_{k=0}^{\infty} \sqrt{k} \lambda^{2k+1} \leq \sum_{k=0}^{\infty} \sqrt{k} \lambda^k \leq \sum_{k=1}^{\infty} (k+1) \lambda^k = \frac{d}{d\lambda} \left(\frac{1}{1-\lambda} \right) - 1 \leq \frac{2\lambda}{(1-\lambda)^2} < 4. \quad (4.13)$$

Following the same reasoning, with probability at least $1 - \frac{4}{n^3}$,

$$\sum_{k=0}^{\infty} \|u_i^T (\alpha_{22} \alpha_{22}^T)^{(k+1)} \alpha_{21} V_1^T \tilde{V}_1 \tilde{\Sigma}_1^{-(2k+3)}\|_2 \leq c_6 \sigma \frac{\sqrt{r} + \sqrt{\log n}}{\sigma_r(A)},$$

for some constant c_6 . Using these in (4.11), by the union bound, we obtain that with probability at least $1 - \frac{8}{n^2}$,

$$\|U_2 U_2^T \Delta A V_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty} \leq c_7 \sigma \frac{\sqrt{r} + \sqrt{\log n}}{\sigma_r(A)},$$

where c_7 is some constant. □

4.4 Proof of Theorem 2.10

Although Theorem 2.10 is motivated and could be proved by Theorem 2.6, we provide an alternative proof that is more straightforward. For this purpose, we will need the following lemmas.

Lemma 4.4. *Under the same assumption as Theorem 2.10, we have*

$$A_r - \tilde{A}_r = U \begin{pmatrix} -U_1^T \Delta A \tilde{V}_1 & -U_1^T \Delta A \tilde{V}_2 \\ -U_2^T A V_2 V_2^T \tilde{V}_1 & 0 \end{pmatrix} \tilde{V}^T + U \begin{pmatrix} 0 & U_1^T \tilde{U}_2 \tilde{U}_2^T \tilde{A} \tilde{V}_2 \\ -U_2^T \Delta A \tilde{V}_1 & 0 \end{pmatrix} \tilde{V}^T.$$

Proof. The lemma can be straightforwardly verified by using the relation $A_r = U_1 \Sigma_1 V_1^T$ and $\tilde{A}_r = \tilde{U}_1 \tilde{\Sigma}_1 \tilde{V}_1^T$. \square

Lemma 4.5 (Lemma 2 in [31]). *Suppose $x_1 \geq x_2 \geq \dots \geq x_k \geq 0$ and $y_1 \geq y_2 \geq \dots \geq y_k \geq 0$. For any $1 \leq j \leq k$, $\sum_{i=1}^j x_i \leq \sum_{i=1}^j y_i$. Then for any $p \geq 1$,*

$$\sum_{i=1}^k x_i^p \leq \sum_{i=1}^k y_i^p.$$

The equality holds if and only if $(x_1, x_2, \dots, x_k) = (y_1, y_2, \dots, y_k)$.

Lemma 4.6 (Theorem 1 in [38]). *Assume $A, B, C = A + B$ are (not necessarily square) matrices of the same size, with singular values*

$$\alpha_1 \geq \alpha_2 \geq \dots, \beta_1 \geq \beta_2 \geq \dots, \gamma_1 \geq \gamma_2 \geq \dots,$$

respectively. Let $i_1 < i_2 < \dots < i_m$ and $j_1 < j_2 < \dots < j_m$ be positive integers, and set

$$k_t = i_t + j_t - t, \quad t = 1, 2, \dots, m.$$

Then the singular values of A, B, C satisfy

$$\sum_{t=1}^m \gamma_{k_t} \leq \sum_{t=1}^m \alpha_{i_t} + \sum_{t=1}^m \beta_{j_t}.$$

Lemma 4.7. *We have the following uniform error bound on $\sin\Theta$ distance*

$$\max\{\|\sin\Theta(U_1, \tilde{U}_1)\|, \|\sin\Theta(V_1, \tilde{V}_1)\|\} \leq \min\left\{\frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1\right\}. \quad (4.14)$$

Proof of Lemma 4.7. If $\sigma_r = \sigma_{r+1}$, (4.14) holds trivially, here we consider the case $\sigma_r > \sigma_{r+1}$. Consider the two possibilities $\sigma_r - \sigma_{r+1} > 2\|\Delta A\|$ and $\sigma_r - \sigma_{r+1} \leq 2\|\Delta A\|$. When $\sigma_r - \sigma_{r+1} > 2\|\Delta A\|$, this and the Weyl's bound

$$|\tilde{\sigma}_r - \sigma_r| \leq \|\Delta A\|,$$

together give

$$\tilde{\sigma}_r - \sigma_{r+1} > \sigma_r - \sigma_{r+1} - \|\Delta A\| > \frac{1}{2}(\sigma_r - \sigma_{r+1}) > 0,$$

which ensures the assumption in Theorem 3.5 to hold, and then (3.8) in Theorem 3.5 implies

$$\|\sin\Theta(U_1, \tilde{U}_1)\| = \|U_2^T \tilde{U}_1\| \leq \frac{\|\Delta A\|}{\sigma_r - \|\Delta A\| - \sigma_{r+1}} \leq \frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}.$$

When $\sigma_r - \sigma_{r+1} \leq 2\|\Delta A\|$, we directly have

$$\|U_2^T \tilde{U}_1\| \leq 1 \leq \frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}.$$

Putting the two cases together, we have

$$\|\sin\Theta(U_1, \tilde{U}_1)\| \leq \min\left\{\frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1\right\}.$$

Following the same reasoning, we also have $\|\sin\Theta(V_1, \tilde{V}_1)\| \leq \min\left\{\frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1\right\}$, thus (4.14) holds. \square

Proof of Theorem 2.10. By Lemma 4.4,

$$\begin{aligned} \|A_r - \tilde{A}_r\| &\leq \left\| \begin{pmatrix} -U_1^T \Delta A \tilde{V}_1 & -U_1^T \Delta A \tilde{V}_2 \\ -U_2^T A V_2 V_2^T \tilde{V}_1 & 0 \end{pmatrix} \right\| + \left\| \begin{pmatrix} 0 & U_1^T \tilde{U}_2 \tilde{U}_2^T \tilde{A} \tilde{V}_2 \\ -U_2^T \Delta A \tilde{V}_1 & 0 \end{pmatrix} \right\| \\ &\leq \sqrt{\|U_1^T \Delta A\|^2 + \|U_2^T A V_2 V_2^T \tilde{V}_1\|^2} + \max\{\|U_2^T \Delta A \tilde{V}_1\|, \|U_1^T \tilde{U}_2 \tilde{U}_2^T \tilde{A} \tilde{V}_2\|\} \\ &= \sqrt{\|U_1^T \Delta A\|^2 + \tau^2} + \max\{\|U_2^T \Delta A \tilde{V}_1\|, \nu\}, \end{aligned} \quad (4.15)$$

where we have let $\nu = \|U_1^T \tilde{U}_2 \tilde{U}_2^T \tilde{A} \tilde{V}_2\|$ and $\tau = \|U_2^T A V_2 V_2^T \tilde{V}_1\|$, we next bound τ and ν .

$$\nu = \|U_1^T \tilde{U}_2 \tilde{U}_2^T \tilde{A} \tilde{V}_2\| = \|U_1^T \tilde{U}_2 \tilde{\Sigma}_2\| \leq \tilde{\sigma}_{r+1} \|U_1^T \tilde{U}_2\|. \quad (4.16)$$

Due to the Weyl's bound, we also have

$$|\tilde{\sigma}_{r+1} - \sigma_{r+1}| < \|\Delta A\|. \quad (4.17)$$

By (4.14),

$$\nu \leq (\sigma_{r+1} + \|\Delta A\|) \min\left\{\frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1\right\} \leq \|\Delta A\| + \sigma_{r+1} \min\left\{\frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1\right\}.$$

Similarly, we can also derive

$$\tau = \|\Sigma_2 V_2^T \tilde{V}_1\| \leq \sigma_{r+1} \min\left\{\frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1\right\}.$$

Inserting the upper bounds of τ and ν back to (4.15) completes the proof of (2.10). For Frobenius norm:

$$\begin{aligned} \|A_r - \tilde{A}_r\|_F^2 &= \left\| \begin{pmatrix} -U_1^T \Delta A \tilde{V}_1 & -U_1^T \Delta A \tilde{V}_2 + U_1^T \tilde{U}_2 \tilde{U}_2^T \tilde{A} \tilde{V}_2 \\ -U_2^T \Delta A \tilde{V}_1 - U_2^T A V_2 V_2^T \tilde{V}_1 & 0 \end{pmatrix} \right\|_F^2 \\ &= \underbrace{\left\| \begin{pmatrix} -U_1^T \Delta A \tilde{V}_1 \\ -U_2^T \Delta A \tilde{V}_1 - U_2^T A V_2 V_2^T \tilde{V}_1 \end{pmatrix} \right\|_F^2}_{:=R_1} + \underbrace{\left\| \begin{pmatrix} -U_1^T \Delta A \tilde{V}_2 + U_1^T \tilde{U}_2 \tilde{U}_2^T \tilde{A} \tilde{V}_2 \\ 0 \end{pmatrix} \right\|_F^2}_{:=R_2}. \end{aligned}$$

Next we bound R_1 and R_2 separately. First consider R_2 , let $M\Lambda W^T$ be the singular value decomposition of $U_1^T \tilde{U}_2$, where $M \in \mathbb{R}^{r \times r}$, $\Lambda \in \mathbb{R}^{r \times r}$, $W \in \mathbb{R}^{(n-r) \times r}$. Then

$$\begin{aligned} R_2 &= \|-U_1 \Delta A \tilde{V}_2 + U_1 \tilde{U}_2 \tilde{\Sigma}_2\|_F^2 \\ &\leq 2\|U_1^T \Delta A \tilde{V}_2\|_F^2 + 2\|U_1^T \tilde{U}_2 \tilde{\Sigma}_2\|_F^2 \\ &\leq 2\|(\Delta A)_r\|_F^2 + 2\|U_1^T \tilde{U}_2 W W^T \tilde{\Sigma}_2\|_F^2 \\ &\leq 2\|(\Delta A)_r\|_F^2 + 2\left(\min\left\{\frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1\right\}\right)^2 \|W^T \tilde{\Sigma}_2\|_F^2 \\ &\leq 2\|(\Delta A)_r\|_F^2 + 2\left(\min\left\{\frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1\right\}\right)^2 \sum_{k=1}^r \sigma_{r+k}^2(\tilde{A}). \end{aligned}$$

In the second to last inequality, we used the fact that $\|AB\|_F \leq \|A\|\|B\|_F$ and in the last inequality, we used $\|P_\Omega A\|_F \leq \|A_r\|_F$ for any r -dimensional subspace Ω . By Lemma 4.6,

$$\sum_{i=1}^k \sigma_{r+i}(\tilde{A}) = \sum_{i=1}^k \sigma_{r+i}(A + \Delta A) \leq \sum_{i=1}^k \sigma_i(\Delta A) + \sum_{i=1}^k \sigma_{r+i}(A) = \sum_{i=1}^k (\sigma_i(\Delta A) + \sigma_{r+i}(A)), 1 \leq k \leq r.$$

From Lemma 4.5, we have

$$\sum_{k=1}^r \sigma_{r+k}^2(\tilde{A}) \leq \sum_{k=1}^r (\sigma_k(\Delta A) + \sigma_{r+k}(A))^2 \leq (\|(\Delta A)_r\|_F + \|(\Sigma_2)_r\|_F)^2.$$

Hence

$$R_2 \leq 2\|(\Delta A)_r\|_F^2 + 2 \left(\min \left\{ \frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1 \right\} \right)^2 (\|(\Delta A)_r\|_F + \|(\Sigma_2)_r\|_F)^2. \quad (4.18)$$

Next we consider R_1 . Notice that

$$R_1 = \left\| \begin{pmatrix} -U_1^T \Delta A \tilde{V}_1 \\ -U_2^T \Delta A \tilde{V}_1 \end{pmatrix} + \begin{pmatrix} 0 \\ -U_2^T A V_2 V_2^T \tilde{V}_1 \end{pmatrix} \right\|_F^2 \leq (\|(\Delta A)_r\|_F + \|\Sigma_2 V_2^T \tilde{V}_1\|_F)^2.$$

Following the same reasoning as in bounding R_2 , we have

$$R_1 \leq \left(\|(\Delta A)_r\|_F + \|(\Sigma_2)_r\|_F \min \left\{ \frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1 \right\} \right)^2. \quad (4.19)$$

Combining (4.18) and (4.19), we obtain

$$\|A_r - \tilde{A}_r\|_F^2 \leq 2\|(\Delta A)_r\|_F^2 + 3 \left(\|(\Delta A)_r\|_F + \|(\Sigma_2)_r\|_F \min \left\{ \frac{2\|\Delta A\|}{\sigma_r - \sigma_{r+1}}, 1 \right\} \right)^2.$$

□

4.5 Proof of Theorem 2.7

Proof of Theorem 2.7. Let $\tilde{V}_1^T V_1 = Q_1 S Q_2^T$ be the SVD of $\tilde{V}_1^T V_1$. Define a special rotation $\hat{Q} = Q_1 Q_2^T$, and we bound $\|U_1 \Sigma_1 - \tilde{U}_1 \tilde{\Sigma}_1 \hat{Q}\|$, where $\|\cdot\|$ can be either the spectral or the Frobenius norm. This yields an upper bound on $\min_{Q \in \mathcal{O}_r} \|U_1 \Sigma_1 - \tilde{U}_1 \tilde{\Sigma}_1 Q\|$. By a direct calculation,

$$\begin{aligned} \|U_1 \Sigma_1 - \tilde{U}_1 \tilde{\Sigma}_1 \hat{Q}\| &\leq \|U_1 \Sigma_1 - \tilde{U}_1 \tilde{\Sigma}_1 \tilde{V}_1^T V_1 + \tilde{U}_1 \tilde{\Sigma}_1 \tilde{V}_1^T V_1 - \tilde{U}_1 \tilde{\Sigma}_1 \hat{Q}\| \\ &\leq \|U_1 \Sigma_1 - \tilde{U}_1 \tilde{\Sigma}_1 \tilde{V}_1^T V_1\| + \|\tilde{U}_1 \tilde{\Sigma}_1 \tilde{V}_1^T V_1 - \tilde{U}_1 \tilde{\Sigma}_1 \hat{Q}\| \\ &= \|(U_1 \Sigma_1 V_1^T - \tilde{U}_1 \tilde{\Sigma}_1 \tilde{V}_1^T) V_1\| + \|\tilde{U}_1 \tilde{\Sigma}_1 (\tilde{V}_1^T V_1 - \hat{Q})\| \\ &\leq \|U_1 \Sigma_1 V_1^T - \tilde{U}_1 \tilde{\Sigma}_1 \tilde{V}_1^T\| + \|\tilde{U}_1 \tilde{\Sigma}_1 (\tilde{V}_1^T V_1 - \hat{Q})\| \\ &= \|A_r - \tilde{A}_r\| + \|\tilde{U}_1 \tilde{\Sigma}_1 (\tilde{V}_1^T V_1 - \hat{Q})\|. \end{aligned} \quad (4.20)$$

The first term of (4.20) can be bounded by Theorem 2.10. Let us focus on the second term. Let $Z = V_2^T \tilde{V}_1$. Observe $Z^T Z + \tilde{V}_1^T V_1 (\tilde{V}_1^T V_1)^T = I_r$, this implies $\tilde{V}_1^T V_1 = \sqrt{I_r - Z^T Z} \hat{Q}$ (Lemma 4.8), where the square root of a positive semi-definite matrix B is defined to be the positive semi-definite matrix \tilde{B} such that $\tilde{B} \tilde{B} = B$.

Using this observation on the quantity inside the norm of the second term on the right hand side of (4.20), we have

$$\tilde{U}_1 \tilde{\Sigma}_1 (\tilde{V}_1^T V_1 - \hat{Q}) = \tilde{U}_1 \tilde{\Sigma}_1 (\sqrt{I - Z^T Z} - I) \hat{Q} = -\tilde{U}_1 \tilde{\Sigma}_1 Z^T Z (\sqrt{I - Z^T Z} + I)^{-1} \hat{Q},$$

where the last equality used the fact that $(\sqrt{I - Z^T Z} - I)(\sqrt{I - Z^T Z} + I) = -Z^T Z$. Hence

$$\| \tilde{U}_1 \tilde{\Sigma}_1 (\tilde{V}_1^T V_1 - \hat{Q}) \| \leq \| \tilde{U}_1 \tilde{\Sigma}_1 Z^T Z \| \cdot \| (\sqrt{I - Z^T Z} + I)^{-1} \hat{Q} \| \leq \| \tilde{\Sigma}_1 Z^T \|. \quad (4.21)$$

The last inequality used $\|Z\| \leq 1$, and $\|(\sqrt{I - Z^T Z} + I)^{-1}\| \leq 1$. Notice that

$$(\tilde{\Sigma}_1 Z^T)^T = Z \tilde{\Sigma}_1 = V_2^T \tilde{V}_1 \tilde{\Sigma}_1 = V_2^T \tilde{A}^T \tilde{U}_1 = V_2^T \Delta A^T \tilde{U}_1 + V_2^T A^T \tilde{U}_1 = V_2^T \Delta A^T \tilde{U}_1 + \Sigma_2^T U_2^T \tilde{U}_1.$$

Then for the spectral norm of $\tilde{\Sigma}_1 Z^T$, we have

$$\| \tilde{\Sigma}_1 Z^T \| \leq \| \Delta A \| + \sigma_{r+1} \| U_2^T \tilde{U}_1 \| \leq \| \Delta A \| + \sigma_{r+1} \min \left\{ \frac{2 \| \Delta A \|}{\sigma_r - \sigma_{r+1}}, 1 \right\}.$$

For the Frobenius norm, we have

$$\| \tilde{\Sigma}_1 Z^T \|_F \leq \| V_2^T \Delta A^T \tilde{U}_1 \|_F + \| \Sigma_2^T U_2^T \tilde{U}_1 \|_F \leq \| (\Delta A)_r \|_F + \| (\Sigma_2)_r \|_F \min \left\{ \frac{2 \| \Delta A \|}{\sigma_r - \sigma_{r+1}}, 1 \right\}.$$

Combining this with (4.20) and (4.21) completes the proof. \square

Lemma 4.8. *Let $A \in \mathbb{R}^{r \times r}$ be a semi-definite matrix with eigenvalues no greater than 1, $B \in \mathbb{R}^{r \times r}$ has SVD $B = U_B S_B V_B^T$. In addition, A and B satisfy $A + BB^T = I$, then*

$$B = \sqrt{I - A} U_B V_B^T.$$

Proof. Since $BB^T = U_B S_B^2 U_B^T$, then $\sqrt{BB^T} = U_B S_B U_B^T$, and therefore $B = \sqrt{BB^T} U_B V_B^T$. By assumption, $BB^T = I - A$, then the result of the lemma follows. \square

References

- [1] E. Abbe. Community detection and stochastic block models: recent developments. *The Journal of Machine Learning Research*, 18(1):6446–6531, 2017.
- [2] E. Abbe, J. Fan, and K. Wang. An ℓ_p theory of pca and spectral clustering. *The Annals of Statistics*, 50(4):2359–2385, 2022.
- [3] E. Abbe, J. Fan, K. Wang, and Y. Zhong. Entrywise eigenvector analysis of random matrices with low expected rank. *The Annals of Statistics*, 48(3):1452–1474, 2020.
- [4] J. Agterberg, Z. Lubberts, and C. E. Priebe. Entrywise estimation of singular vectors of low-rank matrices with heteroskedasticity and dependence. *IEEE Transactions on Information Theory*, 68(7):4618–4650, 2022.
- [5] A. Athreya, M. Tang, Y. Park, and C. E. Priebe. On estimation and inference in latent structure random graphs. *Statistical Science*, 36(1):68–88, 2021.
- [6] R. Bhatia. *Matrix analysis*, volume 169. Springer Science & Business Media, 2013.

-
- [7] G. Blanchard, O. Bousquet, and L. Zwald. Statistical properties of kernel principal component analysis. *Machine Learning*, 66:259–294, 2007.
- [8] C. Cai, G. Li, Y. Chi, H. V. Poor, and Y. Chen. Subspace estimation from unbalanced and incomplete data matrices: $\ell_{2,\infty}$ statistical guarantees. *The Annals of Statistics*, 49(2):944–967, 2021.
- [9] J. F. Cai, E. J. Candès, and Z. Shen. A singular value thresholding algorithm for matrix completion. *SIAM Journal on optimization*, 20(4):1956–1982, 2010.
- [10] T. T. Cai and A. Zhang. Rate-optimal perturbation bounds for singular subspaces with applications to high-dimensional statistics. *The Annals of Statistics*, 46(1):60–89, 2018.
- [11] E. J. Candès and Y. Plan. Matrix completion with noise. *Proceedings of the IEEE*, 98(6):925–936, 2010.
- [12] E. J. Candès and B. Recht. Exact matrix completion via convex optimization. *Communications of the ACM*, 55(6):111–119, 2012.
- [13] J. Cape, M. Tang, and C. E. Priebe. Signal-plus-noise matrix models: eigenvector deviations and fluctuations. *Biometrika*, 106(1):243–250, 2019.
- [14] J. Cape, M. Tang, and C. E. Priebe. The two-to-infinity norm and singular subspace geometry with applications to high-dimensional statistics. *The Annals of Statistics*, 47(5):2405–2439, 2019.
- [15] Y. Chen, C. Cheng, and J. Fan. Asymmetry helps: Eigenvalue and eigenvector analyses of asymmetrically perturbed low-rank matrices. *The Annals of statistics*, 49(1):435, 2021.
- [16] Y. Chen, Y. Chi, J. Fan, C. Ma, et al. Spectral methods for data science: A statistical perspective. *Foundations and Trends® in Machine Learning*, 14(5):566–806, 2021.
- [17] C. Cheng, Y. Wei, and Y. Chen. Inference for linear forms of eigenvectors under minimal eigenvalue separation: Asymmetry and heteroscedasticity. *arXiv preprint arXiv:2001.04620*, 2020.
- [18] P. Chin, A. Rao, and V. Vu. Stochastic block model and community detection in sparse graphs: A spectral algorithm with optimal rate of recovery. In *Conference on Learning Theory*, pages 391–423. PMLR, 2015.
- [19] C. Davis and W. M. Kahan. The rotation of eigenvectors by a perturbation. iii. *SIAM Journal on Numerical Analysis*, 7(1):1–46, 1970.
- [20] D. Donoho and M. Gavish. Minimax risk of matrix denoising by singular value thresholding. *The Annals of Statistics*, 42(6):2413–2440, 2014.
- [21] F. M. Dopico. A note on $\sin \theta$ theorems for singular subspace variations. *BIT Numerical Mathematics*, 40(2):395–403, 2000.
- [22] F. M. Dopico and J. Moro. Perturbation theory for simultaneous bases of singular subspaces. *BIT Numerical Mathematics*, 42(1):84–109, 2002.
- [23] J. Eldridge, M. Belkin, and Y. Wang. Unperturbed: spectral analysis beyond davis-kahan. In *Algorithmic Learning Theory*, pages 321–358. PMLR, 2018.
- [24] J. Fan, Y. Fan, X. Han, and J. Lv. Asymptotic theory of eigenvectors for random matrices with diverging spikes. *Journal of the American Statistical Association*, 117(538):996–1009, 2022.
- [25] M. Gavish and D. L. Donoho. The optimal hard threshold for singular values is $4/\sqrt{3}$. *IEEE Transactions on Information Theory*, 60(8):5040–5053, 2014.

-
- [26] R. Keshavan, A. Montanari, and S. Oh. Matrix completion from noisy entries. *Advances in neural information processing systems*, 22, 2009.
- [27] A. V. Knyazev and M. E. Argentati. Principal angles between subspaces in an a -based scalar product: algorithms and perturbation estimates. *SIAM Journal on Scientific Computing*, 23(6):2008–2040, 2002.
- [28] L. Lei. Unified $l_2 \rightarrow \infty$ eigenspace perturbation theory for symmetric random matrices. *arXiv preprint arXiv:1909.04798*, 2019.
- [29] A. Little, Y. Xie, and Q. Sun. An analysis of classical multidimensional scaling. *arXiv preprint arXiv:1812.11954*, 2018.
- [30] M. Löffler, A. Y. Zhang, and H. H. Zhou. Optimality of spectral clustering in the gaussian mixture model. *The Annals of Statistics*, 49(5):2506–2530, 2021.
- [31] Y. Luo, R. Han, and A. Zhang. A Schatten- q low-rank matrix perturbation analysis via perturbation projection error bound. *Linear Algebra and its Applications*, 630:225–240, 2021.
- [32] P. Narayanamurthy and N. Vaswani. Fast robust subspace tracking via pca in sparse data-dependent noise. *IEEE Journal on Selected Areas in Information Theory*, 1(3):723–744, 2020.
- [33] L. H. Nguyen and S. Holmes. Ten quick tips for effective dimensionality reduction. *PLoS computational biology*, 15(6):e1006907, 2019.
- [34] K. Pearson. Liii. on lines and planes of closest fit to systems of points in space. *The London, Edinburgh, and Dublin philosophical magazine and journal of science*, 2(11):559–572, 1901.
- [35] M. Stewart. Perturbation of the svd in the presence of small singular values. *Linear algebra and its applications*, 419(1):53–77, 2006.
- [36] M. Tang and C. E. Preibe. Limit theorems for eigenvectors of the normalized laplacian for random graphs. *The Annals of Statistics*, 46(5):2360–2415, 2018.
- [37] J. Tanner and K. Wei. Normalized iterative hard thresholding for matrix completion. *SIAM Journal on Scientific Computing*, 35(5):S104–S125, 2013.
- [38] R. C. Thompson. Singular value inequalities for matrix sums and minors. *Linear Algebra and its Applications*, 11(3):251–269, 1975.
- [39] N. Vaswani and P. Narayanamurthy. Finite sample guarantees for pca in non-isotropic and data-dependent noise. In *2017 55th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 783–789. IEEE, 2017.
- [40] R. Vershynin. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge university press, 2018.
- [41] T. Vu, E. Chumikhina, and R. Raich. Perturbation expansions and error bounds for the truncated singular value decomposition. *Linear Algebra and its Applications*, 627:94–139, 2021.
- [42] H. Weyl. Das asymptotische verteilungsgesetz der eigenwerte linearer partieller differentialgleichungen (mit einer anwendung auf die theorie der hohlraumstrahlung). *Mathematische Annalen*, 71(4):441–479, 1912.
- [43] Y. Yu, T. Wang, and R. J. Samworth. A useful variant of the davis–kahan theorem for statisticians. *Biometrika*, 102(2):315–323, 2015.
- [44] S-Y Yun and A. Proutiere. Accurate community detection in the stochastic block model via spectral algorithms. *arXiv preprint arXiv:1412.7335*, 2014.

5 Appendix

Proof of Lemma 4.3- the full-rank case. We first bound $\|U_2 U_2^T \Delta A V_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty}$. To do so, we need Theorem 3.7 to obtain the expansion of $V_2^T \tilde{V}_1$. Let us first check that in the setting of this lemma (i.e., $21\sigma\sqrt{\bar{n}} < \sigma_r(A) - \sigma_{r+1}(A)$), the condition in Theorem 3.7 is satisfied with high probability, that is, $\|\mathcal{F}\| < 1$, where

$$\mathcal{F} \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} F_U^{21} \circ (\Sigma_2 \alpha_{22}^T C_1) + F_U^{21} \circ (\alpha_{22} C_2 \tilde{\Sigma}_1^T) \\ F_V^{21} \circ (\alpha_{22}^T C_1 \tilde{\Sigma}_1) + F_V^{21} \circ (\Sigma_2^T \alpha_{22} C_2) \end{pmatrix}.$$

As discussed in the proof of Theorem 2.1, $\|\Delta A\| \leq 3\sigma\sqrt{\bar{n}}$ with probability at least $1 - e^{-c\bar{n}}$ with some constant c . By the assumption $21\sigma\sqrt{\bar{n}} < \sigma_r(A) - \sigma_{r+1}(A)$, we have with probability at least $1 - e^{-c\bar{n}}$,

$$\begin{aligned} & \left\| \mathcal{F} \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} \right\| \\ & \leq \|F_U^{21} \circ (\Sigma_2 \alpha_{22}^T C_1)\| + \|F_U^{21} \circ (\alpha_{22} C_2 \tilde{\Sigma}_1^T)\| + \|F_V^{21} \circ (\alpha_{22}^T C_1 \tilde{\Sigma}_1)\| + \|F_V^{21} \circ (\Sigma_2^T \alpha_{22} C_2)\| \\ & \leq \frac{\sigma_{r+1}}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} \|\alpha_{22}\| \|C_1\| + \frac{\tilde{\sigma}_r}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} \|\alpha_{22}\| \|C_2\| + \frac{\tilde{\sigma}_r}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} \|\alpha_{22}\| \|C_1\| + \frac{\sigma_{r+1}}{\tilde{\sigma}_r^2 - \sigma_{r+1}^2} \|\alpha_{22}\| \|C_2\| \\ & = \frac{\|\alpha_{22}\|}{\tilde{\sigma}_r - \sigma_{r+1}} (\|C_1\| + \|C_2\|) \\ & \leq \frac{\|\Delta A\|}{\sigma_r - \sigma_{r+1} - \|\Delta A\|} \cdot \sqrt{2(\|C_1\|^2 + \|C_2\|^2)} \\ & \leq \frac{\sqrt{2}}{6} \sqrt{\|C_1\|^2 + \|C_2\|^2}. \end{aligned}$$

Here the second inequality is due to Lemma 3.4. Now we have $\|\mathcal{F}\| < 1$ with high probability, which enables us to use Theorem 3.7 to decompose $U_2 U_2^T \Delta A V_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}$. Denote

$$a_1 = F_U^{21} \circ (\Sigma_2 \alpha_{12}^T U_1^T \tilde{U}_1), \quad a_2 = F_U^{21} \circ (\alpha_{21} V_1^T \tilde{V}_1 \tilde{\Sigma}_1^T), \quad a_3 = F_V^{21} \circ (\alpha_{12}^T U_1^T \tilde{U}_1 \tilde{\Sigma}_1), \quad a_4 = F_V^{21} \circ (\Sigma_2^T \alpha_{21} V_1^T \tilde{V}_1),$$

$$f_1(X) = F_U^{21} \circ (\Sigma_2 \alpha_{22}^T X), \quad f_2(X) = F_U^{21} \circ (\alpha_{22} X \tilde{\Sigma}_1^T), \quad f_3(X) = F_V^{21} \circ (\alpha_{22}^T X \tilde{\Sigma}_1), \quad f_4(X) = F_V^{21} \circ (\Sigma_2^T \alpha_{22} X).$$

By Theorem 3.7, each term in the expansion of $V_2^T \tilde{V}_1$ is of the form

$$f_{i_1}(f_{i_2}(\dots(f_{i_k}(a_{i_0}))))), \quad 1 \leq i_0, i_1, \dots, i_k \leq 4, \quad k = 0, 1, 2, \dots$$

Now assume i_1, \dots, i_k and k are fixed. Let w be the i th row in $U_2 \alpha_{22} f_{i_1}(f_{i_2}(\dots(f_{i_k}(a_{i_0})))) \tilde{\Sigma}_1^{-1}$, and let $b^T = u_i^T \alpha_{22}$, where u_i^T is the i th row of U_2 . Then $w = b^T f_{i_1}(f_{i_2}(\dots(f_{i_k}(a_{i_0})))) \tilde{\Sigma}_1^{-1}$. Notice that a_{i_0} and each f_{i_s} , $1 \leq s \leq k$, either contains Σ_2 or $\tilde{\Sigma}_1$, let $h_{i_s} = 1$ if f_{i_s} contains Σ_2 , and $h_{i_s} = 0$ if f_{i_s} contains $\tilde{\Sigma}_1$. Let $h_{i_0} = 1$ if a_{i_0} contains Σ_2 and $h_{i_0} = 0$ if a_{i_0} contains $\tilde{\Sigma}_1$. Also, let m be the total number of times that $\tilde{\Sigma}_1$ appears in f_{i_s} and a_{i_0} . Then

$$h_{i_0} + h_{i_1} + \dots + h_{i_k} + m = k + 1. \quad (5.1)$$

Likewise, each f_{i_s} , $1 \leq s \leq k$ either contains α_{22} or α_{22}^T . Let $d_{i_s} = \alpha_{22}$ if f_{i_s} contains α_{22} and $d_{i_s} = \alpha_{22}^T$ if it contains α_{22}^T . Also, let $\gamma = \alpha_{12}^T$ if a_{i_0} contains α_{12}^T and $\gamma = \alpha_{21}$ if a_{i_0} contains α_{21} . Last, denote $\beta = V_1^T \tilde{V}_1$ if a_{i_0} contains $V_1^T \tilde{V}_1$ and $\beta = U_1^T \tilde{U}_1$, if a_{i_0} contains $U_1^T \tilde{U}_1$. For the γ and β defined above, let γ_l^T be the l th row of γ , i.e., $\gamma = [\gamma_1^T; \gamma_2^T; \dots; \gamma_{n-r}^T]$ and β_i be the i th column of β , i.e., $\beta = [\beta_1, \beta_2, \dots, \beta_r]$. Then for

$1 \leq j \leq r$, the j th entry in w is

$$\begin{aligned} w_j &= \sum_{l_1} \frac{b_{l_1} \sigma_{r+l_1}^{h_{i_1}}}{\tilde{\sigma}_j^2 - \sigma_{r+l_1}^2} \sum_{l_2} \frac{(d_{i_1})_{l_1 l_2} \sigma_{r+l_2}^{h_{i_2}}}{\tilde{\sigma}_j^2 - \sigma_{r+l_2}^2} \cdots \sum_{l_k} \frac{(d_{i_{k-1}})_{l_{k-1} l_k} \sigma_{r+l_k}^{h_{i_k}}}{\tilde{\sigma}_j^2 - \sigma_{r+l_k}^2} \sum_{l_0} \frac{(d_{i_k})_{l_k l_0} \sigma_{r+l_0}^{h_{i_0}}}{\tilde{\sigma}_j^2 - \sigma_{r+l_0}^2} \gamma_{l_0}^T (\tilde{\sigma}_j^{m-1} \beta_j) \\ &= \underbrace{\left\langle \sum_{l_1} \frac{b_{l_1} \sigma_{r+l_1}^{h_{i_1}}}{\tilde{\sigma}_j^2 - \sigma_{r+l_1}^2} \sum_{l_2} \frac{(d_{i_1})_{l_1 l_2} \sigma_{r+l_2}^{h_{i_2}}}{\tilde{\sigma}_j^2 - \sigma_{r+l_2}^2} \cdots \sum_{l_k} \frac{(d_{i_{k-1}})_{l_{k-1} l_k} \sigma_{r+l_k}^{h_{i_k}}}{\tilde{\sigma}_j^2 - \sigma_{r+l_k}^2} \sum_{l_0} \frac{(d_{i_k})_{l_k l_0} \sigma_{r+l_0}^{h_{i_0}}}{\tilde{\sigma}_j^2 - \sigma_{r+l_0}^2} \gamma_{l_0}, \tilde{\sigma}_j^{m-1} \beta_j \right\rangle}_{\equiv M_j}. \end{aligned}$$

In the above, let

$$M_j = \sum_{l_1} \frac{b_{l_1} \sigma_{r+l_1}^{h_{i_1}}}{\tilde{\sigma}_j^2 - \sigma_{r+l_1}^2} \sum_{l_2} \frac{(d_{i_1})_{l_1 l_2} \sigma_{r+l_2}^{h_{i_2}}}{\tilde{\sigma}_j^2 - \sigma_{r+l_2}^2} \cdots \sum_{l_k} \frac{(d_{i_{k-1}})_{l_{k-1} l_k} \sigma_{r+l_k}^{h_{i_k}}}{\tilde{\sigma}_j^2 - \sigma_{r+l_k}^2} \sum_{l_0} \frac{(d_{i_k})_{l_k l_0} \sigma_{r+l_0}^{h_{i_0}}}{\tilde{\sigma}_j^2 - \sigma_{r+l_0}^2} \gamma_{l_0}.$$

We first bound $\|M_j\|$. Denote $\eta_j = \sigma_j^2 - \tilde{\sigma}_j^2$, and $\Delta_{jl} = \sigma_j^2 - \sigma_{r+l}^2$. Notice that

$$\begin{aligned} \frac{1}{\tilde{\sigma}_j^2 - \sigma_{r+l}^2} &= \frac{1}{\sigma_j^2 - \sigma_{r+l}^2 + (\tilde{\sigma}_j^2 - \sigma_j^2)} = \frac{1}{(\sigma_j^2 - \sigma_{r+l}^2)(1 + \frac{\tilde{\sigma}_j^2 - \sigma_j^2}{\sigma_j^2 - \sigma_{r+l}^2})} \\ &= \frac{1}{\Delta_{jl}(1 - \frac{\eta_j}{\Delta_{jl}})} = \frac{1}{\Delta_{jl}} \left(1 + \frac{\eta_j}{\Delta_{jl}} + \left(\frac{\eta_j}{\Delta_{jl}}\right)^2 + \dots\right). \end{aligned}$$

Hence

$$\begin{aligned} \|M_j\| &= \left\| \sum_{l_1, \dots, l_k, l_0} \frac{b_{l_1} \prod_{s=1}^k (d_{i_s})_{l_s l_{s+1}} \prod_{s=0}^k \sigma_{r+l_s}^{h_{i_s}}}{\prod_{s=0}^k \Delta_{j l_s}} \prod_{s=0}^k \left(1 + \frac{\eta_j}{\Delta_{j l_s}} + \left(\frac{\eta_j}{\Delta_{j l_s}}\right)^2 + \dots\right) \gamma_{l_0} \right\| \\ &= \left\| \sum_{q_0, q_1, \dots, q_k=0}^{\infty} \sum_{l_1, \dots, l_k, l_0} \frac{b_{l_1} \prod_{s=1}^k (d_{i_s})_{l_s l_{s+1}} \prod_{s=0}^k \sigma_{r+l_s}^{h_{i_s}}}{(\prod_{s=0}^k \Delta_{j l_s}) \prod_{s=0}^k \Delta_{j l_s}^{q_s}} \gamma_{l_0} \cdot \eta_j^{\sum_{s=0}^k q_s} \right\| \\ &\leq \sum_{q_0, q_1, \dots, q_k=0}^{\infty} \underbrace{\left\| \sum_{l_1, \dots, l_k, l_0} \frac{b_{l_1} \prod_{s=1}^k (d_{i_s})_{l_s l_{s+1}} \prod_{s=0}^k \sigma_{r+l_s}^{h_{i_s}}}{\prod_{s=0}^k \Delta_{j l_s}^{1+q_s}} \gamma_{l_0} \right\|}_{T(j, q_0, \dots, q_k)} \cdot \left| \eta_j^{\sum_{s=0}^k q_s} \right|. \end{aligned}$$

Here we let $l_{k+1} = l_0$. In the above, denote

$$T(j, q_0, \dots, q_k) \equiv \sum_{l_1, \dots, l_k, l_0} \frac{b_{l_1} \prod_{s=1}^k (d_{i_s})_{l_s l_{s+1}} \prod_{s=0}^k \sigma_{r+l_s}^{h_{i_s}}}{\prod_{s=0}^k \Delta_{j l_s}^{1+q_s}} \gamma_{l_0} \in \mathbb{R}^{r \times 1}.$$

Next, we bound the ℓ_2 norm of $T(j, q_0, \dots, q_k)$. Notice that we can rewrite $T(j, q_0, \dots, q_k)$ in the following way

$$T(j, q_0, \dots, q_k) = \left(b^T \hat{f}_{i_1}^{q_1} \hat{f}_{i_2}^{q_2} \cdots \hat{f}_{i_k}^{q_k} \hat{a}_{i_0}^{q_0} \right)^T, \quad 1 \leq i_0, i_2, \dots, i_k \leq 4, \quad k \geq 0.$$

Here matrices \hat{f}_{i_s} are modified from the functions f_{i_s} , and matrix \hat{a}_{i_0} is modified from the function a_{i_0} . Explicitly,

$$\begin{aligned} \hat{a}_1^q &= \hat{F}_U^{21, q} \Sigma_2 \alpha_{12}^T, \quad \hat{a}_2^q = \hat{F}_U^{21, q} \alpha_{21}, \quad \hat{a}_3^q = \hat{F}_V^{12, q} \alpha_{12}^T, \quad \hat{a}_4^q = \hat{F}_V^{21, q} \Sigma_2^T \alpha_{21}, \\ \hat{f}_1^q &= \hat{F}_U^{21, q} \Sigma_2 \alpha_{22}^T, \quad \hat{f}_2^q = \hat{F}_U^{21, q} \alpha_{22}, \quad \hat{f}_3^q = \hat{F}_V^{21, q} \alpha_{22}^T, \quad \hat{f}_4^q = \hat{F}_V^{21, q} \Sigma_2^T \alpha_{22}, \end{aligned}$$

where $\hat{F}_U^{21,q}$ and $\hat{F}_V^{21,q}$ are diagonal matrices with diagonal entries

$$(\hat{F}_U^{21,q})_{i'-r, i'-r} = \frac{1}{(\sigma_j^2 - \sigma_{i'}^2)^{1+q}}, \quad r+1 \leq i' \leq n,$$

$$(\hat{F}_V^{21,q})_{i'-r, i'-r} = \frac{1}{(\sigma_j^2 - \sigma_{i'}^2)^{1+q}}, \quad r+1 \leq i' \leq m.$$

Similar as in Theorem 3.1, if $i' > \min\{n, m\}$, we define $\sigma_{i'}$ to be 0.

As before, In the above expression of $T(j, q_0, \dots, q_k)$, $\hat{a}_{i_0}^{q_0}$ either contains α_{21} or α_{12}^T . If it contains the former, let γ be the former, and it contains the latter, let γ be the latter. It is easy to check that this γ coincides with the γ defined in the paragraph under (5.1).

Conditional on α_{22} , the only random variable in $T(j, q_0, \dots, q_k)$ is α_{21} or α_{12}^T , that is γ . Therefore, if we write $T(j, q_0, \dots, q_k) = G(\gamma)^T$, then the linear operator G is independent of γ , and it is straightforward to check that

$$\|G\| \leq \underbrace{\|b\| \frac{\|\alpha_{22}\|^k \sigma_{r+1}^{\sum_{s=0}^k h_{i_s}}}{(\sigma_j^2 - \sigma_{r+1}^2)^{k+1 + \sum_{s=0}^k q_s}}}_{\equiv K}, \quad 1 \leq j \leq r.$$

Again, conditional on α_{22} , since for each $1 \leq p \leq r$, the p th entry of $T(j, q_0, \dots, q_k)$ only depends on the p th column of α_{21} or α_{12}^T , then different entries of $T(j, q_0, \dots, q_k)$ are independent of each other, each following a Gaussian distribution $\mathcal{N}(0, \sigma^2 \xi_p^2)$, and $\xi_p \leq K$, $1 \leq p \leq r$, where K denotes the above bound. By Theorem 3.1.1 in [40], there exists some constant c such that

$$\mathbb{P}(\|T(j, q_0, \dots, q_k)\| - K\sigma\sqrt{r} \geq t) \leq 2e^{-ct^2/\sigma^2 K^2}.$$

Let $t = K\sigma\sqrt{\log(n^3 \cdot 2^{\sum_{s=0}^k q_s} \cdot 8^{kr})}/c$, then with probability at least $1 - \frac{2}{n^3 \cdot 2^{\sum_{s=0}^k q_s} \cdot 8^{kr}}$,

$$\|T(j, q_0, \dots, q_k)\| \leq c_1 \sigma \|b\| \frac{\sigma_{r+1}^{\sum_{s=0}^k h_{i_s}} \|\alpha_{22}\|^k}{(\sigma_j^2 - \sigma_{r+1}^2)^{k+1 + \sum_{s=0}^k q_s}} (\sqrt{\log n} + \sum_{s=0}^k \sqrt{q_s} + \sqrt{k} + \sqrt{r}),$$

where c_1 is some constant. Hence

$$\begin{aligned} \|M_j\| &\leq \sum_{q_0, q_1, \dots, q_k=0}^{\infty} \|T(j, q_0, \dots, q_k)\| \cdot \left| \eta_j^{\sum_{s=0}^k q_s} \right| \\ &\leq c_1 \sigma \frac{\|b\| \sigma_{r+1}^{\sum_{s=0}^k h_{i_s}} \|\alpha_{22}\|^k}{(\sigma_j^2 - \sigma_{r+1}^2)^{k+1}} \sum_{q_0, q_1, \dots, q_k=0}^{\infty} (\sqrt{\log n} + \sum_{s=0}^k \sqrt{q_s} + \sqrt{k} + \sqrt{r}) \cdot \left| \frac{\eta_j}{\sigma_j^2 - \sigma_{r+1}^2} \right|^{\sum_{s=0}^k q_s} \\ &\leq c_1 \sigma \frac{\|b\| \sigma_{r+1}^{\sum_{s=0}^k h_{i_s}} \|\alpha_{22}\|^k}{(\sigma_j^2 - \sigma_{r+1}^2)^{k+1}} \left((\sqrt{\log n} + \sqrt{k} + \sqrt{r}) \frac{1}{(1 - \frac{|\eta_j|}{\sigma_j^2 - \sigma_{r+1}^2})^{k+1}} + \frac{2(k+1)}{(1 - \frac{|\eta_j|}{\sigma_j^2 - \sigma_{r+1}^2})^{k+1}} \right) \\ &= c_2 \sigma \frac{\|b\| \sigma_{r+1}^{\sum_{s=0}^k h_{i_s}} \|\alpha_{22}\|^k}{(\sigma_j^2 - \sigma_{r+1}^2 - |\eta_j|)^{k+1}} (\sqrt{\log n} + \sqrt{r} + k), \end{aligned}$$

where c_2 is a constant. In the second inequality above, we used the fact that

$$\begin{aligned} \sum_{q_0, q_1, \dots, q_k=0}^{\infty} \left| \frac{\eta_j}{\sigma_j^2 - \sigma_{r+1}^2} \right|^{\sum_{s=0}^k q_s} &= \sum_{q_0, q_1, \dots, q_{k-1}=0}^{\infty} \left| \frac{\eta_j}{\sigma_j^2 - \sigma_{r+1}^2} \right|^{\sum_{s=0}^{k-1} q_s} \sum_{q_k=0}^{\infty} \left| \frac{\eta_j}{\sigma_j^2 - \sigma_{r+1}^2} \right|^{q_k} \\ &= \frac{1}{\left(1 - \frac{|\eta_j|}{\sigma_j^2 - \sigma_{r+1}^2}\right)} \sum_{q_0, q_1, \dots, q_{k-1}=0}^{\infty} \left| \frac{\eta_j}{\sigma_j^2 - \sigma_{r+1}^2} \right|^{\sum_{s=0}^{k-1} q_s} \\ &= \dots = \frac{1}{\left(1 - \frac{|\eta_j|}{\sigma_j^2 - \sigma_{r+1}^2}\right)^{k+1}}. \end{aligned}$$

and that

$$\begin{aligned} &\sum_{q_0, q_1, \dots, q_k=0}^{\infty} \left(\sum_{s=0}^k \sqrt{q_s} \right) \left| \frac{\eta_j}{\sigma_j^2 - \sigma_{r+1}^2} \right|^{\sum_{s=0}^k q_s} \\ &= \sum_{s=0}^k \sum_{q_0, \dots, q_{s-1}, q_{s+1}, \dots, q_k=0}^{\infty} \left| \frac{\eta_j}{\sigma_j^2 - \sigma_{r+1}^2} \right|^{\sum_{l \neq s} q_l} \sum_{q_s=0}^{\infty} \sqrt{q_s} \left| \frac{\eta_j}{\sigma_j^2 - \sigma_{r+1}^2} \right|^{q_s} \\ &\leq \sum_{s=0}^k \frac{2}{\left(1 - \frac{|\eta_j|}{\sigma_j^2 - \sigma_{r+1}^2}\right)} \sum_{q_0, \dots, q_{s-1}, q_{s+1}, \dots, q_k=0}^{\infty} \left| \frac{\eta_j}{\sigma_j^2 - \sigma_{r+1}^2} \right|^{\sum_{l \neq s} q_l} \\ &= \frac{2(k+1)}{\left(1 - \frac{|\eta_j|}{\sigma_j^2 - \sigma_{r+1}^2}\right)^{k+1}}. \end{aligned}$$

Here the inequality is due to (4.13), which holds under the condition

$$\begin{aligned} \frac{|\eta_j|}{\sigma_j^2 - \sigma_{r+1}^2} &= \frac{|\tilde{\sigma}_j^2 - \sigma_j^2|}{\sigma_j^2 - \sigma_{r+1}^2} \leq \frac{(\sigma_j + \|\Delta A\|)^2 - \sigma_j^2}{\sigma_j^2 - \sigma_{r+1}^2} \\ &\leq \frac{2\sigma_j \|\Delta A\| + \|\Delta A\|^2}{(\sigma_j - \sigma_{r+1})(\sigma_j + \sigma_{r+1})} \leq \frac{(2\sigma_j + \frac{1}{7}\sigma_j)\|\Delta A\|}{7\|\Delta A\|\sigma_j} < \frac{1}{2}. \end{aligned}$$

Therefore with probability at least $1 - \frac{4}{n^3 \cdot 4^k r}$,

$$\begin{aligned} \|w_j\| &\leq \|M_j\| \cdot \tilde{\sigma}_j^{m-1} \|\beta_j\| \\ &\leq c_2 \sigma \frac{\|b\| \tilde{\sigma}_j^{m-1} \sigma_{r+1}^{\sum_{s=0}^k h_{i_s}} \|\alpha_{22}\|^k}{(\sigma_j^2 - \sigma_{r+1}^2 - |\eta_j|)(\sigma_j^2 - \sigma_{r+1}^2 - |\eta_j|)^k} (\sqrt{\log n} + \sqrt{r} + k) \\ &\leq c_2 \sigma \frac{\|b\| (\frac{8}{7}\sigma_j)^k \|\Delta A\|^k}{\frac{2}{3}(\sigma_j^2 - \sigma_{r+1}^2) (\frac{34}{7}\sigma_j \|\Delta A\|)^k} (\sqrt{\log n} + \sqrt{r} + k) \\ &\leq \frac{3}{2} c_2 \sigma \frac{\|b\|}{\sigma_r^2 - \sigma_{r+1}^2} \left(\frac{4}{17}\right)^k (\sqrt{\log n} + \sqrt{r} + k). \end{aligned}$$

Here the third inequality is due to $\sum_{s=0}^k h_{i_s} + m - 1 = k$ and

$$\begin{aligned} \sigma_j^2 - \sigma_{r+1}^2 - |\tilde{\sigma}_j^2 - \sigma_j^2| &\geq \sigma_j(\sigma_j - \sigma_{r+1}) - ((\sigma_j + \|\Delta A\|)^2 - \sigma_j^2) \\ &\geq 7\sigma_j \|\Delta A\| - 2\sigma_j \|\Delta A\| - \|\Delta A\|^2 \geq \frac{34}{7}\sigma_j \|\Delta A\|, \end{aligned}$$

as well as

$$\begin{aligned}
\sigma_j^2 - \sigma_{r+1}^2 - |\tilde{\sigma}_j^2 - \sigma_j^2| &\geq \sigma_j^2 - \sigma_{r+1}^2 - 2\sigma_j \|\Delta A\| - \|\Delta A\|^2 \\
&\geq \sigma_j^2 - \sigma_{r+1}^2 - \|\Delta A\| (2\sigma_j + \|\Delta A\|) \\
&\geq \sigma_j^2 - \sigma_{r+1}^2 - \frac{1}{7}(\sigma_j - \sigma_{r+1}) \cdot \frac{15}{7}(\sigma_j + \sigma_{r+1}) \\
&\geq \frac{2}{3}(\sigma_j^2 - \sigma_{r+1}^2).
\end{aligned}$$

With probability at least $1 - \frac{4}{4^k n^3}$,

$$\|w\| \leq \frac{3}{2} c_2 \sigma \frac{\sqrt{r} \|u_i^T \alpha_{22}\|}{\sigma_r^2 - \sigma_{r+1}^2} \left(\frac{4}{17}\right)^k (\sqrt{\log n} + \sqrt{r} + k).$$

By Theorem 3.7, we can see that there are 2^{k+1} terms in the expansion of $V_2^T \tilde{V}_1$ that has order k , hence by (4.13), with probability at least $1 - \frac{16}{n^3}$,

$$\|u_i^T \alpha_{22} V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\| \leq \sum_{k=0}^{\infty} 3c_2 \sigma \frac{\sqrt{r} \|u_i^T \alpha_{22}\|}{\sigma_r^2 - \sigma_{r+1}^2} \left(\frac{8}{17}\right)^k (\sqrt{\log n} + \sqrt{r} + k) \leq C\sigma \frac{(\sqrt{r \log n} + r) \|u_i^T \alpha_{22}\|}{\sigma_r^2 - \sigma_{r+1}^2}.$$

By the union bound, with probability at least $1 - \frac{16}{n^2}$,

$$\begin{aligned}
\|U_2 U_2^T \Delta A V_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty} &\leq C\sigma \frac{(\sqrt{r \log n} + r) \|\alpha_{22}\|}{\sigma_r^2 - \sigma_{r+1}^2} \\
&= C\sigma \frac{(\sqrt{r \log n} + r) \|\Delta A\|}{(\sigma_r - \sigma_{r+1})(\sigma_r + \sigma_{r+1})} \leq C\sigma \frac{\sqrt{r \log n} + r}{\sigma_r + \sigma_{r+1}}.
\end{aligned}$$

Next, we consider $\|U_2 \Sigma_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty}$. Following the same reasoning, we have with probability at least $1 - \frac{16}{n^2}$,

$$\|U_2 \Sigma_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty} \leq C\sigma \frac{(\sqrt{r \log n} + r) \|\Sigma_2\|}{\sigma_r^2 - \sigma_{r+1}^2} = C\sigma \frac{(\sqrt{r \log n} + r) \sigma_{r+1}}{\sigma_r^2 - \sigma_{r+1}^2}.$$

Combining the above two bounds,

$$\begin{aligned}
\max\{\|U_2 \Sigma_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty}, \|U_2 U_2^T \Delta A V_2 V_2^T \tilde{V}_1 \tilde{\Sigma}_1^{-1}\|_{2,\infty}\} &\leq C\sigma \left(\frac{(\sqrt{r \log n} + r) \sigma_{r+1}}{\sigma_r^2 - \sigma_{r+1}^2} + \frac{\sqrt{r \log n} + r}{\sigma_r + \sigma_{r+1}} \right) \\
&\leq C\sigma \frac{\sqrt{r \log n} + r}{\sigma_r - \sigma_{r+1}}.
\end{aligned}$$

□